1 **Hepatic loss of AATF attenuates MASH by suppressing AKT–mTORC1**
2 **signaling and reprogramming lipid metabolism**
3

Akshatha N. Srinivas[1], Diwakar Suresh[1], Prajna Anirvan[2], Manju Moorthy[3],

Gopalkrishna Ramaswamy[3], Suma M. Nataraj[1], Prasanna Kumar Santhekadur[1],

Deepak Suvarna[4], Shivaram P. Singh[2], and Divya P. Kumar[1]*

[1]Department of Biochemistry, CEMR, JSS Medical College, JSS Academy of Higher

Education and Research, Mysuru, Karnataka, India

[2] Department of Translational Research, Kalinga Gastroenterology Foundation,

Cuttack, Odisha, India

[3]TheraCUES Innovations Pvt. Ltd, Bangalore, India

[4]Department of Gastroenterology, JSS Medical College and Hospital, JSS Academy

of Higher Education and Research, Mysuru- 570015, Karnataka, India

*Correspondence:

Dr. Divya P. Kumar

Associate Professor

Department of Biochemistry

JSS Medical College, JSS AHER

SS Nagar, Mysuru- 570015

Karnataka, India

Email: divyapk243@gmail.com/

divyapkumar@jssuni.edu.in

Abstract

Background & aims

Metabolic dysfunction-associated steatohepatitis (MASH) is a multifactorial disease

driven by complex molecular mechanisms. Identifying key regulators is critical for

developing targeted therapies. Here, we demonstrate the impact of the loss of

apoptosis antagonizing transcription factor (AATF) on hepatic lipid metabolism and

MASH progression.

Methods

A preclinical mouse model recapitulating human MASH was established by feeding

C57Bl/6 mice either a chow diet (CD) or a western diet with sugar water (WD).

Hepatic AATF silencing was achieved by tail vein injection of siAATF delivered by

adeno-associated virus 8 (AAV8) with the liver-specific thyroxine-binding globulin

(TBG) promoter. In addition to histological, biochemical, and molecular biology

evaluations, the mechanistic insights were derived from whole transcriptomics and

untargeted metabolomics analyses.

Results

AAV8-mediated specific knockdown of AATF in hepatocytes significantly reduced

body weight, liver weight, and insulin resistance in mice fed with western diet (WD).

However, no such effects were observed in mice fed with a chow diet (CD). Further

analyses showed reduced liver injury, steatosis, and steatohepatitis in WDsiAATF

mice. Transcriptomic analysis demonstrated that loss of AATF alleviated cellular

stress, inflammation, and fibrosis in WD-fed mice. Moreover, AATF silencing led to

alterations in lipid metabolism, notably by decreasing hepatic lipogenesis in the WD

mice. Interestingly, untargeted metabolomics revealed an increase in the

biosynthesis of glycerophospholipids and beta-oxidation of fatty acids in WDsiAATF

mice.

Conclusion

Our findings reveal a previously unrecognized role of AATF as a central regulator of hepatic lipid metabolism in MASH, acting through the AKT–mTORC1 signaling pathway, and establish its inhibition as a promising therapeutic strategy for metabolic liver disease.

Keywords

Apoptosis antagonizing transcription factor, metabolic dysfunction-associated steatotic liver disease, fatty acid β-oxidation, therapeutic target, transcriptomics, and metabolomics

Background

Metabolic dysfunction-associated steatotic liver disease (MASLD), formerly termed non-alcoholic fatty liver disease (NAFLD), stands as the prevailing chronic liver ailment impacting more than 30% of the adult population worldwide (1,2). It notably afflicts individuals with obesity and type 2 diabetes, emerging as a primary contributor to advanced liver disease and hepatocellular carcinoma (HCC) (3,4). MASLD encompasses a spectrum of advancing steatotic liver conditions, spanning from benign metabolic dysfunction-associated steatotic liver (MASL) to steatohepatitis (MASH), presenting with varying degrees of liver fibrosis that could advance to cirrhosis (5). MASLD not only heightens the likelihood of liver-related complications such as cirrhosis, end-stage liver disease, and hepatocellular carcinoma (HCC) but also escalates the risk of multiple extraneous manifestations, including cardiovascular disease (CVD), chronic kidney disease (CKD), and specific forms of extrahepatic cancers (6). As MASLD progresses to the end stage of liver disease, treatment options become increasingly limited, often leading to consideration for liver transplantation (7). Early detection and intervention are, therefore, paramount. While lifestyle modifications remain fundamental to managing the condition, pharmacological treatments can complement these interventions (8,9). Despite numerous clinical studies, the pathogenesis of MASH remains elusive, and the mechanisms underlying its occurrence and progression are not yet fully understood (10). Therefore, elucidating the molecular pathways involved in MASH pathogenesis is crucial for identifying potential therapeutic targets.

MASH manifests with the accumulation of lipids in the liver, stemming from disturbances in both lipid uptake (from circulating sources or de novo synthesis) and lipid disposal (via fatty acid oxidation or secretion of triglyceride-rich lipoproteins) (11). This disruption leads to the accumulation of lipotoxic compounds, cellular stress, injury, and fibrosis (11,12). The regulation of lipid metabolism is complex and multifaceted, involving various factors like membrane transporters, metabolic

enzymes, and transcriptional regulators (13). Glycerophospholipids, a subclass of

phospholipids that are the major constituents of cell membranes, play a crucial role in

lipid metabolism. Alterations in the composition of glycerophospholipids have been

reported in individuals with MASH (14,15). Phosphatidylcholine (PC),

phosphatidylethanolamine (PE), phosphatidylserine (PS), phosphatidylinositol (PI),

and cardiolipin (CL) are a diverse array of glycerophospholipids that are well-studied

for their roles in cell signaling and lipid metabolism (15). Interestingly,

glycerophospholipids are involved in inflammatory, oxidative stress, and bile acid

metabolism pathways, which are key drivers of disease pathogenesis in MASH (16).

Notably, glycerophospholipids are crucial for maintaining the integrity of the

mitochondrial membrane, which is essential for various mitochondrial functions,

including the beta-oxidation of fatty acids (17). It is observed that, in MASH subjects,

diminished β-oxidation of fatty acids in the liver fails to adequately mitigate the

lipotoxic effects of harmful lipids (18). Disruption of β-oxidation pathways leads to

lipid accumulation in the liver, increased oxidative stress, and inflammation, all of

which contribute to the progression of the disease (19). Thus, glycerophospholipid

levels and efficient β-oxidation of fatty acids in the liver contribute to the protection

and alleviation of MASLD.

Apoptosis antagonizing transcription factor (AATF) is a multifaceted transcription

factor recognized for its pivotal involvement in essential cellular functions (20). It

serves as a dynamic regulator of fundamental cellular processes, including

transcriptional control, regulation of the cell cycle and apoptosis, ribosome

biogenesis, and response to cellular stress (20). AATF protects cells from oxidative

stress, ER stress, DNA damage, and hypoxia by modulating the cell cycle,

autophagy, and apoptosis (21). The widely studied role of AATF is the regulation of

apoptosis, exerted by the activation of antiapoptotic pathways and the inhibition of

proapoptotic pathways (22). Pathologically, AATF has been implicated in numerous

cancer types, including leukemia, breast cancer, neuroblastoma, lung cancer, and

colon cancer (23-25). Our previous investigations have revealed elevated levels of

hepatic AATF in HCC, where it plays a significant role in promoting tumor

angiogenesis (26). Studies have also demonstrated the inhibitory effects of curcumin

on MASH-HCC by downregulating AATF via the KLF4-SP1 pathway (27). We

recently investigated the role of the SIRT1-TIMP3-TACE axis in AATF upregulation,

with subsequent TACE inhibition resulting in AATF downregulation and the

attenuation of MASH-associated liver tumorigenesis (28). However, the mechanistic

role of AATF in the pathophysiology of MASH remains poorly understood.

Despite growing evidence implicating metabolic and inflammatory pathways in

MASH, the transcriptional regulators that coordinate these complex responses

remain poorly defined. Here, we demonstrate for the first time that hepatocyte-

specific loss of AATF ameliorates MASH through metabolic reprogramming, including

enhanced   β-oxidation,   suppressed   lipogenesis,   and   restoration   of

glycerophospholipid homeostasis. By integrating transcriptomic and untargeted

metabolomic profiling with in vivo AAV8-mediated gene silencing, our study provides

a novel mechanistic framework linking AATF to MASH pathogenesis and offers a

previously unexplored therapeutic target for this widespread liver disease.

Methods

Animal study

Male C57Bl/6NCrl mice aged 6 weeks procured from Hylasco Biotechnology Pvt. Ltd.

(Charles River Laboratories, Hyderabad, Telangana, India) were used for the study.

All the mice were accommodated in the central animal housing facility at the Center

for Experimental Pharmacology and Toxicology, JSS AHER, following a 12-hour

light-dark cycle, consistent with previously established protocols (59). Following one

week of acclimatization, study protocols were commenced. All animal-related

procedures adhered to the guidelines sanctioned by the Committee for the Purpose

of Control and Supervision of Experiments on Animals (CPCSEA), Govt of India

(JSSAHER/CPT/IAEC/118/2022).

<u>Diet-induced MASH model:</u>

After a week of acclimatization, the mice were fed with either a chow diet normal

water (CD) or a western diet (21% fat, 41% sucrose, and 1.25% cholesterol by

weight, Research Diet, Inc. USA) along with sugar water (WD) containing 18.9 g/L d-

glucose and 23.1 g/L d-fructose (HiMedia Laboratories) ad libitum for 24 weeks. Mice

were continuously monitored for food intake, water intake, and body weight. After 24

weeks, mice were euthanized, and blood and tissues were collected for further

histological, biochemical, and molecular analysis.

AAV expression system

Liver gene expression was modulated using AAV-mediated gene transfer technology,

as previously defined (60). The siRNA sequences targeting AATF were incorporated

into an adeno-associated virus (AAV) vector under the regulation of a liver-specific

thyroxine-binding globulin (TBG) promoter. AAV8-TBG-siAATF and AAV8-TBG-

siControl constructs were generated by packaging into serotype 8, titrated by Applied

Biological Materials Inc. (ABM). The siRNA construct design mimics the behavior of

shRNA due to its vector-based expression system. The siRNA sequence is cloned

into a plasmid vector under the control of a dual convergent promoter system, U6

and H1, which drives the transcription of both strands of DNA to form a shRNA-like

structure within the cell. This technology enhances the stability of siRNA, enabling

prolonged gene silencing. Henceforth, the one-time insertion of siRNA in the study

provided a stable and sustained expression.

Hepatocyte-specific silencing of AATF in experimental MASH

The mice were fed either CD or WD. After 12 weeks of dietary intervention, mice

were injected via the tail vein with 1×10^{11} genome copies of AAV8 serotype particles

and randomized into four groups: (i) CD with AAV8-TBG-siControl, (ii) CD with AAV8-

TBG-siAATF, (iii) WD with AAV8-TBG-siControl, and (iv) WD with AAV8-TBG-

siAATF. The same dietary regimen was followed until the end of 24 weeks.

Throughout the study, mice were continuously monitored for food intake, water

intake, and body weight. The GTT and ITT were performed during the study period.

After 24 weeks, mice were euthanized under fasting conditions, followed by blood

and tissue collection for further analysis.

Isolation of primary mouse hepatocytes

Primary hepatocytes were isolated from the livers of mice injected with AAV8-TBG-

Null and AAV8-TBG-siAATF following previously established protocols (60). Briefly,

mice were anesthetized, and the inferior vena cava was cannulated. Liver perfusion

was carried out using pre-warmed perfusion buffer, followed by digestion buffer

containing 25 µg/ml Liberase (Sigma-Aldrich). The isolated hepatocytes were then

filtered through a 70-micron cell strainer (HiMedia) and purified via centrifugation in

DMEM low-glucose medium (HiMedia Laboratories) at 50g for 5 minutes at room

temperature. Percoll (Sigma Aldrich) gradient separation was employed to isolate

hepatocytes from nonparenchymal cells. The isolated cells were suspended in

DMEM supplemented with 10% FBS and 100 U penicillin-streptomycin. Collagen-

coated plates were utilized for cell culture in William's E medium (HiMedia

Laboratories).

Glucose and Insulin Tolerance Tests

Mice were fasted overnight, or 5-6 hours for GTT and ITT, respectively. The basal

level of blood glucose was measured using a glucometer (Roche). Furthermore, mice

were administered glucose orally (1 g/kg body weight) and insulin intraperitoneally

(0.75 U/kg body weight). Venous tail blood samples were taken at various intervals

between 0 and 120 minutes to measure blood glucose levels.

Biochemical analysis

Blood collected from the fasting mice was centrifuged at 1500 g for 15 min at 4°C.

The serum levels of AST, ALT, total cholesterol (TC), total triglycerides (TG), and

LDL-cholesterol were measured using commercial kits (Monlab tests) according to

the manufacturer's protocol.

Histological analysis

Liver tissues were subjected to formalin (Sigma Aldrich) fixation, and paraffin blocks

were prepared. 4µm tissue sections were used for hematoxylin and eosin staining to

assess the liver morphology. For pathological grading, stained liver sections were

scored by pathologists according to the MASLD activity score, which comprises the

sum of steatosis (0-3), lobular inflammation (0-3), and hepatocyte ballooning (0-2). A

cumulative activity score of 5 or greater was indicative of MASH (61,62).

Oil Red O staining

Oil Red O staining was performed using frozen liver tissues. Cryosections, 12 µm

thick, were fixed with 10% formaldehyde (Sigma Aldrich) for 5 minutes and then

washed with distilled water. Further, the slides were incubated with 60% isopropanol

(Sisco Research Laboratories) and Oil Red O solution (Sigma Aldrich) for one hour.

After rinsing off excess stain with distilled water, nuclei were counterstained with

hematoxylin (HiMedia Laboratories). The slides were then mounted, and images

were captured using the Leica DM570 microscope (Leica Microsystems). For

quantitative analysis, the areas of lipid droplets were quantified using ImageJ

software.

Picrosirius red staining

The formalin-fixed paraffin-embedded (FFPE) liver tissue sections were subjected to

deparaffinization and rehydration using xylene (HiMedia), followed by decreasing

percentages of ethanol. The slides were first stained with hematoxylin for 2 minutes,

followed by incubation with the picrosirius red solution (Sigma Aldrich) for 1 hour.

Slides were washed in 0.5% acidified water and mounted. The images were captured

using the Leica DM570 microscope (Leica Microsystems) and quantified using

ImageJ software.

ELISA

Serum insulin and TNFα levels were quantified using mouse ELISA kits (Krishgen

Biosystems) according to the manufacturer's instructions.

Immunohistochemistry

Immunohistochemistry was performed according to standard protocol. Briefly, after

deparaffinization and rehydration, FFPE liver tissues were subjected to antigen

retrieval with citrate buffer (pH 6) at 94°C for 15 minutes. Following this, the tissue

sections were blocked with 3% hydrogen peroxide (PathnSitu Biotechnologies) and

normal goat serum (Abcam) for 2 hours. Primary antibodies (AATF; Sigma Aldrich,

1:500, F4/80; Cell Signaling Technologies, 1:100, and Desmin; Cell Signaling

Technologies, 1:500) were used to stain the respective tissue sections overnight at

4°C in a humidified chamber. The detection of signals was accomplished using the

Polyexcel HRP/DAB detection system (PathnSitu, Biotechnologies). Images were

captured using the Leica DM570 microscope (Leica Microsystems), and quantitative

analysis was conducted using ImageJ software.

Immunoblot analysis

Liver tissues and cell pellets were dissolved in RIPA buffer (Sigma Aldrich)

containing protease/phosphatase inhibitor cocktail (Thermo Fischer Scientific). The

lysates were obtained by centrifugation at 12000 rpm at 4^0C for 20 minutes. Protein

concentration was estimated by Bradford's (Bio-Rad) protein estimation method.

Equivalent amounts of protein were resolved in SDS-PAGE, followed by transfer into

the nitrocellulose membrane (Bio-Rad). Blots were blocked in 5% nonfat dry milk for

1 hour and probed with primary antibodies of interest (anti-AATF-Sigma Aldrich

1:1000, anti-pAKT, anti-AKT, anti-pmTOR, anti-mTOR, anti-Raptor, and anti-β-actin -

Cell Signaling Technologies 1:1000, anti-pJNK, anti-JNK, anti-pERK1/2, and anti-

ERK1/2 – Santa Cruz Biotechnologies 1:100) overnight at 4°C and secondary

antibodies (Cell Signaling Technologies) for 2 hours at room temperature. Blots were

visualized using Western Bright ECL HRP substrate (Advansta) in the UVitec

Alliance Q9 chemiluminescence imaging system and analyzed for densitometric

measurements in ImageJ software. The band intensity was normalized with the

respective endogenous control.

RNA isolation, cDNA synthesis, and Quantitative real-time PCR

Total RNA was extracted from frozen live tissues or cell pellets using TRIzol reagent

(Sigma-Aldrich) according to the manufacturer's instructions. RNA was quantified

using a Nanodrop spectrophotometer and reverse-transcribed using the Verso cDNA

synthesis kit, following the manufacturer's protocol (Thermo Fisher Scientific).

Relative gene expressions were assessed using quantitative real-time PCR (qRT-

PCR) using the Rotor-Gene Q 5plex HRM system. qRT-PCR was performed in

triplicate per gene using the Dynamo Colorflash SYBR Green kit (Thermo Fisher

Scientific) with 0.5 µM primers (IDT), and 50 ng of cDNA in a 20 µL reaction volume

was used to set up the reaction. The mRNA levels of selected genes were calculated

after normalization to β-actin by using the $2^{(-\Delta\Delta C(T))}$ method. Primer sequences are

provided in Table S1.

RNA sequencing and transcriptomics analysis

RNA extraction was performed using the All-prep DNA/RNA kit (Qiagen). RNA was

quantified using a Qubit fluorometer, and RIN was determined with a TapeStation

4150 using HS RNA screen tape. The library preparation was carried out using the

SMART-Seq Library Prep Kit. The sequencing was performed using the Illumina

NovaSeq 6000 platform with a read length of 150 bp × 2. The reads were mapped to

STAR-indexed Mus musculus (GRCm39: GCF_000001635.27) using the STAR

aligner version 2.7.9a. The alignment file (sorted BAM) from individual samples was

quantified using Feature Counts version 2.0.1, based on the filtered GTF file, to

obtain gene counts. These gene counts for Mus musculus were used as inputs to

DESeq2 for estimating differential expression.

Differential gene expression analysis

Differential expression analysis was performed using the Bioconductor package

DESeq2 (R, version [latest version]). Genes with read counts of less than 10 across

the samples were removed before proceeding with downstream analysis. Raw read

counts were normalized by DESeq2. The magnitude (log 2 transformed fold change)

and significance (P-value) of differential expression between groups were calculated,

and genes with P-values <0.05 were considered significant. We obtained DEGs with

the following criteria: P value <0.05 and |log 2-fold change|≥2. PLS-DA analysis was

carried out in R using the plsda function in the "mixOmics" package.

Functional Gene Set Enrichment Analysis

Functional gene set enrichment analysis (GSEA) was performed using the R

package ClusterProfiler for Gene Ontology Biological Process (GO BP), Molecular

Function (GO MF), and Kyoto Encyclopedia of Genes and Genomes (KEGG)

pathways. For multiple test corrections, P-values were adjusted by the Benjamini-

Hochberg false discovery rate (BH-FDR). Gene sets with a p-value < 0.05 were

considered significantly enriched. Gene sets were downloaded from the MSigDB

database (https://www.gsea-msigdb.org/gsea/msigdb).

Gene signature heatmaps

Normalized expression of leading genes in the enriched cellular stress, inflammation,

and fibrosis pathways identified by GSEA was used to generate heatmaps using the

pheatmap R package. Glycerophospholipid metabolism, lipid metabolism, and fatty

acid degradation pathway gene lists were obtained from the KEGG database.

AATF PPI network

The Search Tool for the Retrieval of Interacting Genes (STRING version 12.0)

(http://string-db.org) was used to systematically screen the top 100 mouse protein

interactors of AATF, set with a high confidence threshold of > 0.7. Interactors of

AATF were also retrieved from the BioGRID database. Furthermore, the protein-

protein interaction network was created using Cytoscape software, where the color of

the node represents the log2 fold change of the gene within the comparison of

WDsiAATF vs WDC. Interactions between AATF and genes involved in biological

processes related to lipid metabolism, as defined by the Gene Ontology, were

screened separately within the STRING database. The gene ontology biological

process and molecular functions, as well as KEGG pathways enriched by the

interactors of the AATF gene, were analyzed and visualized using the ClueGO app

within Cytoscape. For all the data related to transcriptomics, P<0.05 was considered

to indicate a statistically significant difference. All statistical analysis and

visualizations were performed in R v.3.2 unless indicated otherwise.

Untargeted metabolomics- Metabolite extraction and LC-MS analysis

Metabolites were extracted following previously established protocols (63). Briefly,

liver tissues were lyophilized in a vacuum freeze dryer for 24 hours, and 50 mg of

samples were homogenized with 50 mM ammonium bicarbonate (3:1 w/v). Samples

were extracted in cold methanol by centrifugation at 18,000× g for 15 minutes at 4°C.

Supernatants are dried and reconstituted in a 3:1 (v/v) mixture of MeOH and water

for LC/MS analysis. The hepatic metabolome was analyzed using a Vanquish

UHPLC system (Thermo Fisher) coupled with an Orbitrap Eclipse Tribrid mass

spectrometer (Thermo Fisher). Samples were separated on an RP column, 2.1 x 150

mm, 1.8 µm particle size, and 0.3 mL/min flow rate with two eluents for positive and

negative polarity modes.

Metabolomics data analysis

The data processing was done using the Compound Discoverer 3.3 software

(Thermo Fisher Scientific). A pre-existing workflow template titled 'Untargeted

Metabolomics with Statistics Detect Unknowns with ID using Online Databases and

mzLogic' was used to identify the differences between samples. For identification, the

mass tolerance was 5 ppm, and the retention time tolerance was 0.2 min. The

minimum peak intensity was 10,000, and the chromatographic S/N threshold was

1.5. The solvent blanks were used to create exclusion lists, identify background

components, and filter these components from the results table. A pooled quality

control (QC) sample was used for compound identification only.

To visualize the similarity between the metabolomic profiles of the samples from

different groups, principal component analysis (PCA) and heatmap plots were

generated. All the studies were conducted based on the compound's peak areas.

Differential metabolite analysis was performed using a student's t-test, considering p

value<0.05 as significant and log2 foldchange ≥ |1| as either an upregulated or

downregulated metabolite. The log2 fold change per metabolite was calculated as

the ratio between group areas, transformed to a log2 scale, with the group area being

calculated as the median of the compound area per sample within the group. The up-

or down-regulated metabolites were visualized using volcano plots. P-values were

adjusted for multiple testing using the Benjamini-Hochberg correction for false

discovery rate.

The GSEA algorithm was applied to the KEGG Mus musculus (House Mouse)

pathway library, specifically the lipid superclass and its subclasses. Significant

pathways (p < 0.05) identified by the GSEA method were further analyzed. Pathway

enrichment against the Small Molecule Pathway Database (SMPDB) was performed

separately for upregulated and downregulated metabolites using MetaboAnalyst 6.0.

For overlaying the regulation of metabolites involved in the glycerophospholipid

metabolism pathway, the compound list and pathway were accessed from the KEGG

database and visualized using the Pathview R package. To visualize the metabolic

network, MetScape (http://metscape.ncibi.org/) was used, and the Cytoscape plugin

was employed to identify the interactions between the metabolites involved in the

glycerophospholipid metabolism pathway.

Quantification and statistical analysis

Statistical analyses were performed using GraphPad Prism 6.0 software (GraphPad,

Inc., La Jolla, CA, USA). Data were represented as the mean ± standard error of the

mean. Statistical significance was analyzed using Student's t-test for two groups or

analysis of variance (ANOVA) with post hoc Bonferroni correction for multiple

comparisons, and p values < 0.05 (*/#) or < 0.001 (**/##) were considered significant.

Results

Loss of hepatic AATF ameliorates western diet-induced insulin resistance, liver

injury, steatosis, and steatohepatitis

To assess the role of AATF in MASH, we targeted AATF expression in mouse

hepatocytes using a liver-directed adeno-associated virus serotype 8 (AAV8) vector

driven by the thyroxine-binding globulin (TBG) promoter. Mice were fed a chow diet

with normal water (CD) or a western diet with sugar water (WD). After 12 weeks, they

were administered either AAV8-TBG-siControl or AAV8-TBG-siAATF (The stable

gene silencing effect is consistent with the engineered siRNA construct's shRNA-like

behavior, driven by a dual-promoter expression system- *Detailed in the Methods*

section) via tail vein injection and monitored for an additional 12 weeks. At the end of

the 24-week study, serum and tissues from several organs, including the liver,

adipose tissue, heart, kidneys, pancreas, and spleen, were collected. Liver tissue

was used for histological, biochemical, and molecular analyses, as well as

transcriptomic and metabolomic profiling (**Fig. 1A**)**.** AATF expression was assessed

across the collected tissues, confirming that AATF was silenced exclusively in the

livers of AAV8-TBG-siAATF mice, compared to AAV8-TBG-siControl mice (**Fig. S1A**

and **Fig. S1B**). Furthermore, AATF silencing was specific to hepatocytes, with no

impact on non-parenchymal cells (**Fig. S1C**).

Silencing of hepatic AATF significantly reduced body weight, liver weight, liver-to-

body ratio, and body fat mass in WD-fed mice, but did not affect CD mice (**Fig. 1B-**

1E, Fig. S2A). Importantly, AAV8 administration had no adverse effects on the mice,

as evidenced by stable bodyweight, normal behavior, and no mortality or

abnormalities in major organs. Food intake was monitored throughout the study, and

the cumulative calorie intake remained identical for all groups of mice **(Fig. S2B and**

S2C). AATF silencing also improved glucose tolerance and insulin sensitivity in WD

mice, as demonstrated by the glucose tolerance test (GTT) and the insulin tolerance

test (ITT) (**Fig. 1F and 1G**). Notably, WD siAATF mice exhibited lower blood glucose

and fasting insulin levels, along with a reduced homeostatic model assessment for

insulin resistance (HOMA-IR) compared to WD control mice. These findings indicate

AATF silencing ameliorates Western diet-induced impairments in glucose

metabolism and insulin sensitivity (**Fig. 1H-1J**).

Furthermore, we also examined the effect of hepatic AATF silencing on liver injury by

measuring serum levels of liver enzymes, including ALT, AST, total cholesterol,

triglycerides, and LDL cholesterol. Consistently, WD siAATF mice showed reduced

levels of liver enzymes compared to WD controls (**Fig. 2A-2E**). The macroscopic

examination revealed that WD siAATF mice had a healthier liver morphology

compared to WD controls (**Fig. 2F**). Histological analysis of liver sections showed

decreased steatosis and substantial improvement in steatohepatitis upon hepatic

AATF silencing compared to controls in WD mice (**Fig. 2G**). Similar results were

obtained with Oil Red O staining of liver sections, where there was reduced

intracellular lipid droplet accumulation in WD siAATF mice compared to WD controls

(**Fig. 2H**). Consistent with these findings, histological scoring for steatosis, lobular

inflammation, and hepatocyte ballooning was reduced in hepatic AATF-silenced WD

mice, with an improvement in MASLD activity score compared to WD controls (**Fig.**

2I), indicating that hepatic AATF inhibition improves metabolic health by ameliorating

liver injury and dysregulated lipid metabolism.

Effect of AATF silencing on hepatic transcriptomic and metabolomic profiles in

MASH mice

To explore the liver-specific molecular pathways regulated by AATF in MASH, we

performed RNA sequencing and untargeted metabolomic profiling of the liver tissues

from all four groups. Comparisons were made between WDC (western diet-fed mice

administered with AAV8-TBG-siControl) and CDC (chow diet-fed mice administered

with AAV8-TBG-siControl), WDsiAATF (western diet-fed mice administered with

AAV8-TBG-siAATF) and WDC (western diet-fed mice administered with AAV8-TBG-

siControl), and CDsiAATF (chow diet-fed mice administered with AAV8-TBG-siAATF)

and CDC (chow diet-fed mice administered with AAV8-TBG-siControl). The samples

in transcriptomics analysis formed clear clusters between the WDsiAATF, WDC, and

CD groups with good intra-group sample correlation (**Fig. 3A and 3B)**. Volcano plots

showed a distribution of differentially altered genes in WDsiAATF vs. WDC (**Fig. 3C)**.

Differential expression analysis showed 224 up and 144 downregulated genes in the

WDsiAATF vs. WDC group (P value <0.05 and log 2-fold change≥2). In WDsiAATF

mice, genes related to the anti-inflammatory response, such as neutrophilic granule

protein (NGP), lactoferrin (LTF), and major urinary protein 1 (MUP1), which regulates

gluconeogenesis and lipogenesis in the liver, were found to be upregulated. In

contrast, genes like EPH receptor B2 (EPHB2) and suppression of tumorigenicity 18

(ST18), which play roles in liver fibrosis and inflammation, were downregulated.

Additionally, the early growth response gene-2 (EGR2), which is crucial for the

differentiation of monocytes into hepatic lipid-associated macrophages, and tubulin

beta 2B class IIb (TUBB2B), involved in cholesterol biosynthesis, were also

downregulated. GSEA analysis of the KEGG and GO databases showed suppressed

extracellular matrix and collagen binding, immune response, activation of primary bile

acid synthesis, and nitrogen metabolism (**Fig. 3D)**. The PLS-DA plot for hepatic

metabolite profiling showed the differences between experimental groups and

identified a significant separation of liver metabolites between the WDsiAATF and

WDC groups, as well as the WDC and CDC groups (**Fig. 3E).** Univariate analysis

identified 487 metabolites altered in the WD siAATF, of which 412 were upregulated,

in contrast to 197 upregulated metabolites in WDC. Similarly, 867 metabolites were

downregulated in WDC mice. In contrast, only 75 metabolites were downregulated in

WDsiAATF mice, clearly indicating marked differences in metabolome profiling upon

hepatic AATF silencing (Fig. 3F). Consistently, the volcano plots also showed a

similar distribution of differentially altered metabolites in the WDsiAATF vs. WDC

groups **(Fig. 3G).** No significant changes were observed in the hepatic transcriptomic

and metabolomic profiles between CDsiAATF and CDC (**Fig. S3**).

Hepatic AATF loss alleviates cellular stress, inflammation, and fibrosis in

experimental MASH

Cellular stress plays a prime role in the pathogenesis of MASH by promoting hepatic

inflammation and fibrosis. Several factors contribute to cellular stress, stemming from

lipid accumulation in hepatocytes, mitochondrial dysfunction, oxidative stress, and

ER stress (30). In our investigation, we explored the impact of hepatic AATF

silencing on cellular stress responses. Transcriptomics data showed downregulation

of the ERK1/2 cascade, MAPK pathway, reactive oxygen species, PKA pathways,

and apoptotic pathways in the WDsiAATF group (**Fig. 4A and 4B).** We subsequently

validated the expression of key ER stress markers using qRT-PCR. Remarkably, in

WDsiAATF mice, we found significant downregulation in the mRNA levels of CHOP,

GRP78, PERK, and ATF4 compared to WDC (**Fig. 4C).**

Furthermore, to elucidate the intricate crosstalk between cellular stress and

inflammation, we hypothesized that hepatocellular stress triggers inflammatory

signaling pathways, notably stress kinases, leading to the release of pro-

inflammatory cytokines in WDC mice. Consequently, DEGs in WDsiAATF are

attributed to the suppression of TNF-α superfamily cytokine production, macrophage

migration, leukocyte proliferation, and differentiation (**Fig. 4D and 4E**). Silencing of

hepatic AATF in WD mice remarkably reduced inflammation, as evidenced by the

reduced expression of pro-inflammatory cytokines IL-6, IL-1β, and TNF-α (**Fig. 4F**).

Concurrently, phosphorylation of JNK and ERK1/2 was noticeably diminished in

WDsiAATF mice compared to WDC (**Fig. 4G and 4H**). The immunostaining for the

mouse macrophage marker F4/80 corroborated these findings, showing a marked

reduction in response to AATF silencing in WD mice (**Fig. 4I**). These findings indicate

that AATF is an important mediator of cellular stress-induced hepatic inflammation.

More interestingly, transcriptomic analysis of liver tissues from MASH mice revealed

key genes associated with fibrosis, as evidenced by negative regulation of

extracellular matrix organization, collagen metabolism, the TGF-β receptor signaling

pathway, and fibroblast proliferation in WDsiAATF mice (**Fig. 5A and 5B**). The

further validation also confirmed reduced expression of α-SMA, Col1A1, Col3A1, and

TGF-β, the key markers for liver fibrosis in the WDsiAATF group (**Fig. 5C**).

Additionally, picrosirius red staining and hepatic stellate cell activation marked by

desmin were attenuated in WDsiAATF mice, suggesting that AATF silencing reduces

fibrogenic activation in WDC (**Fig. 5D and 5E**). Taken together, these results indicate

the impact of hepatic silencing of AATF on key MASH signatures, including cellular

stress, inflammation, and fibrosis.

Hepatic loss of AATF regulates lipid metabolism and ameliorates metabolic

dysfunction-associated steatohepatitis

Dysregulated lipid metabolism is the primary culprit in MASH pathogenesis, as it

contributes to lipid accumulation, exacerbates insulin resistance, and triggers

inflammation (31). This lipid accumulation results from excessive lipid influx, reduced

fatty acid oxidation, and promotes de novo lipogenesis in the liver, ultimately leading

to steatohepatitis (32). Differentially expressed genes (DEGs) from RNA sequencing

in WDsiAATF were enriched in pathways such as the downregulation of ceramide

and sphingolipids, the fatty acyl-CoA metabolic process, regulation of lipid

localization, and transport (**Fig. 6A and 6B**). Based on this observation, we further

validated the genes involved in fatty acid synthesis, such as FASN, ACC1, ACC2,

and SCD1. We found that lipogenic genes were significantly downregulated in

WDsiAATF mice compared to WDC (**Fig. 6C**).

To comprehend the significance of the dysregulated metabolites for specific

biological processes, a metabolic pathway analysis was conducted using

MetaboAnalyst 6.0. The KEGG pathway enrichment analysis revealed that 74

pathways were enriched by the differentially expressed metabolites (DEMs) in the

WD siAATF vs. WD control groups. Among these pathways, glycerophospholipid

metabolism, ether lipid metabolism, nicotinamide metabolism, linoleic acid

metabolism, riboflavin metabolism, and pyruvate metabolism were significantly

upregulated upon hepatic silencing of AATF in MASH mice. In contrast, branched-

chain amino acid biosynthesis, sulfur metabolism, and glycerolipid metabolism were

downregulated **(Fig. 6D).** With the focus on only lipid-related pathways, DEMs were

further enriched against the lipid superclass within the KEGG database, which

unveiled the upregulation of glycerophospholipids (glycerophosphocholines,

glycerophosphoserines, and glycerophosphoethanolamines), fatty esters, and

eicosanoids in WDsiAATF mice **(Fig. 6E).** Delving deeper into lipid subclass analysis,

PC, PS, PE, lysoPC, lysoPE, lysoPS, and vitamin D3 were elevated in WDsiAATF

mice **(Fig. 6F).**

Glycerophospholipids are key structural elements of cell membranes and are

involved in several cell signaling pathways (16). Notably, disruption in

glycerophospholipid metabolism is known to compromise membrane stability and

thereby contribute to the development of MASLD (17). Along the same lines, a

network analysis of putative interacting proteins of the DEMs identified them as being

enriched in the glycerophospholipid metabolism pathway in the KEGG database **(Fig.**

7A). We further measured the relative abundance of different glycerophospholipids.

We found that the levels of phosphocholine, PC, lysoPE, and lysoPC were

significantly increased in WD siAATF mice compared to WD controls (**Fig. 7B**). This

was further supported by the transcriptomic analysis, where the genes involved in

glycerophospholipid metabolism were upregulated in WDsiAATF mice. (**Fig. 7C**). To

gain deeper insights into the biological functions of the differentiated metabolites, a

functional relationship network was constructed utilizing MetScape. The resulting

network revealed that most of the metabolites were primarily involved in the

metabolism of glycerophospholipids, linoleate, and phosphatidyl inositol (**Fig. S4**).

Furthermore, we investigated the relationships between clinical variables related to

MASH (anthropometry, liver enzymes, lipid profile, blood glucose, and insulin levels)

and metabolite signatures using a debiases sparse partial correlation (DSPC)

network based on MetaboAnalyst software (**Fig. 7D**). The correlation networks

suggested that the majority of the metabolites belonging to glycerophospholipid

metabolism showed strong positive interactions with each other and negative

interactions with clinical variables, indicating a crucial role of glycerophospholipid

metabolism in MASH pathogenesis. Collectively, these findings underscore the

profound impact of AATF silencing on liver metabolites, particularly in the

upregulation of glycerophospholipid biosynthesis.

The process of beta-oxidation of fatty acids plays a crucial role in the development of

MASH. When beta-oxidation pathways are disrupted, it leads to the accumulation of

lipids in the liver, increased oxidative stress, and inflammation, all of which contribute

to disease progression (31). In contrast, alpha-linolenic acid (ALA), an essential

omega-3 (n-3) polyunsaturated fatty acid (PUFA), provides substantial antioxidant

benefits, as well as anti-inflammatory and lipid-lowering effects (33). Along similar

lines, our studies revealed the upregulation of metabolites involved in the

mitochondrial beta-oxidation pathway and alpha-linolenic acid metabolism alongside

the biosynthesis of glycerophospholipids in WDsiAATF through pathway enrichment

analysis using the SMPB database (**Fig. 8A**). Consistently, genes involved in fatty

acid oxidation were upregulated in WDsiAATF mice (**Fig. 8B)**. Based on this, we

hypothesized whether silencing AATF enhances beta-oxidation of fatty acids, thereby

improving MASH. Supporting this, our results showed a significant upregulation of

key genes involved in fatty acid oxidation, including carnitine palmitoyltransferase-1

alpha (CPT1α), acyl-CoA oxidase 1 (ACOX1), peroxisome proliferator-activated

receptor gamma coactivator-1 alpha (PGC1α), and acyl-CoA synthetase long-chain

family member 1 (ACSL1) in WD siAATF mice compared to the controls (**Fig. 8C**).

Furthermore, to understand the possible interacting partners of AATF in regulating

MASH progression, we performed an integrative analysis of AATF interacting

proteins with the transcriptomic signature of the WDsiAATF group using the STRING

and BioGRID databases (**Fig. S5)**. The protein-protein interaction network revealed

that AATF is involved in the positive regulation of fatty acid metabolic processes

through CD74 and MAPK9; however, further validation is necessary to elucidate the

signaling pathway.

Molecular mechanisms underlying hepatic AATF-driven MASH progression

mTORC1 (mammalian target of rapamycin complex 1) is a central regulator of cell

growth and metabolism that contributes to the pathogenesis of MASH through its

effects on lipid metabolism (34-36). In MASH, the activation of the AKT (protein

kinase B) and mTORC1 signaling pathways is often dysregulated, contributing to the

disease progression (37). We hypothesized that AATF silencing inactivates the AKT-

mTORC1 signaling pathway, thereby mobilizing hepatic lipid accumulation.

Accordingly, we found that phosphorylation of AKT at Thr308 was inhibited in

WDsiAATF mice compared to the WDC mice. Following this, activation of mTORC1,

represented by the phosphorylation of mTOR and Raptor, was downregulated in

WDsiAATF mice (**Fig. 8D)**. Further, SREBP1 levels were downregulated in

WDsiAATF mice, supporting the mTORC1-mediated lipogenesis (**Fig. 8E**). Previous

studies have shown that mTORC1 inhibits lipolysis and β-oxidation of fatty acids

(38,39). In our study, we found that inactivation of mTORC1 via AATF activates long-

chain acyl-CoA dehydrogenase (LCAD), an enzyme crucial for the initiation of β-

oxidation (**Fig. 8F**). Interestingly, PNPLA2 (Patatin-Like Phospholipase Domain-

Containing Protein 2), an enzyme involved in lipolysis, particularly breakdown of

triglycerides, was upregulated in WDsiAATF mice compared to WDC (**Fig. 8G**).

PNPLA2 is responsible for the conversion of triglycerides to diacylglycerol and free

fatty acids, which further undergo glycerophospholipid synthesis and β-oxidation,

respectively (40). Collectively, hepatic silencing of AATF ameliorates MASH by

enhancing glycerophospholipid biosynthesis and β-oxidation of fatty acids via

inhibition of AKT-mTORC1 signaling pathways (**Fig. 8H**).

Discussion

MASLD is a multifactorial condition influenced by various molecular drivers (5).

Dysregulation of lipid metabolism is a hallmark of MASLD. The increased influx of

free fatty acids (FFAs) from adipose tissue and dietary sources, combined with

impaired fatty acid oxidation and the export of triglycerides from the liver, contributes

to hepatic steatosis (11). Insulin resistance in hepatocytes leads to impaired

suppression of gluconeogenesis and increased lipogenesis, promoting hepatic lipid

accumulation (31). Several factors associated with metabolic dysfunction, including

lipid accumulation, oxidative stress, and insulin resistance, can trigger ER stress in

hepatocytes (30). Inflammatory pathways play a crucial role in the progression of

MASLD from simple steatosis to MASH. Activation of inflammatory signaling

pathways leads to the production of pro-inflammatory cytokines and chemokines,

contributing to hepatocyte injury and fibrosis (41). Thus, identifying and

understanding the molecular drivers involved in MASH pathogenesis is essential for

developing targeted therapeutic strategies aimed at preventing disease progression

and improving clinical outcomes for patients with MASLD. The evolving

understanding of the disease underscores the urgent need for new therapeutic

targets. The complex pathophysiology of MASH involves multiple metabolic,

inflammatory, and fibrotic pathways, making it essential to identify novel targets that

can address these different aspects. Advanced molecular profiling techniques,

including transcriptomics, proteomics, metabolomics, lipidomics, and epigenomics,

can provide valuable insights into the underlying mechanisms driving MASH and

guide the development of personalized treatment approaches.

AATF has been implicated in the pathogenesis of several cancers due to its role in

modulating apoptotic pathways (20). We, for the first time, elucidated how TNF-α

upregulates the expression of AATF via SREBP-1, promoting hepatocarcinogenesis

(42). Further studies revealed the role of AATF in impeding HCC angiogenesis by

downregulating PEDF (26). Recently, we investigated the role of AATF in MASH-

HCC and showed that TACE inhibition by Marimastat prevents AATF-mediated

inflammation, fibrosis, and oncogenesis (28). Building on these experimental

findings, we sought to understand the role of AATF in MASH. In the current study, we

explored for the first time how AATF regulates lipid metabolism and its implications in

the development and progression of MASH. We employed a preclinical murine model

of MASH comprising C57BL/6 mice that were fed a standard chow diet, normal

water, or western diet sugar water. This model effectively mimics the features of

metabolic syndrome and MASH, including the progression of fibrosis. The mice

develop fatty liver within 12 weeks of being fed with a western diet, sugar water,

progressing to MASH and fibrosis by 24 weeks. This spectrum mirrors the dietary

pattern, systemic milieu, and metabolic and histological characteristics of MASH,

demonstrating the activation of key cellular pathways associated with human

disease. This approach overcomes the constraint of the previously employed

DIAMOND mouse model, which was an isogenic cross between C57BL/6J and

129S1/SvImJ mouse strains (29). Although the DIAMOND model closely mimics the

human disease characteristics, the genetic instability and associated inconsistency in

phenotypic outcomes render it less accepted. In the current study, we assessed the

impact of liver-specific silencing of AATF on hepatic lipid accumulation in

experimental MASH. Of note, while siRNA was chosen for gene silencing, the

approach employed in this study closely resembles the characteristics of shRNA,

particularly in terms of its stability. The technology developed by Applied Biological

Materials Inc. (ABM) utilizes a convergent promoter vector system, which allows

siRNA to be expressed in both directions— a feature typical of shRNA. Additionally,

the AAV8 vector used for gene delivery offers tissue-specific targeting, providing

stability and sustained gene expression. We examined how this silencing affected the

genes, proteins, and metabolites associated with the biological processes involved in

the development and progression of MASH. The silencing of hepatic AATF showed a

significant reduction in body weight and liver weight in WDSW mice and did not affect

CDNW mice. Insulin resistance was notably improved, and we also observed

substantial improvement in both histological and biochemical parameters in

WDsiAATF mice compared to WDC. In line with these findings, liver transcriptomics

analysis and on-bench validations demonstrated that the loss of AATF reduced

cellular stress responses, leading to decreased production of inflammatory cytokines

and fibrogenesis in WDSW mice. These results underscore the potential role of

AATF in the progression of MASLD and set the groundwork for more detailed

molecular investigations.

This is the first study to conduct both hepatic transcriptomic and metabolomic

profiling in mice with AATF silenced. Since the precise mechanism(s) by which the

loss of AATF alleviates MASH are unclear, we concentrated on detecting changes in

liver metabolites in the experimental MASH group with AATF silencing compared to

the control group using untargeted metabolomics, which offers a comprehensive

overview of the entire metabolome. Based on transcriptomic analysis and our key

interest in elucidating the role of AATF in lipid metabolism, we found it intriguing that

the leading unregulated pathways among the 74 pathways enriched by DEMs in the

WD siAATF compared to the WD control group included glycerophospholipid

metabolism, ether lipid metabolism, nicotinamide metabolism, linoleic acid

metabolism, riboflavin metabolism, and pyruvate metabolism. Notably, there was a

marked increase in glycerophospholipids, particularly PC, PS, and PE, in the WD

siAATF mice. These subclasses of lipids are implicated in MASLD progression,

where dysregulation of these glycerophospholipids is linked with insulin resistance,

inflammation, and fibrosis, which are the hallmarks of MASLD (43). Additionally, their

breakdown products, like LysoPC, LysoPS, and LysoPE, were also increased. The

metabolite-metabolite interaction network highlighted the potential functional

relationships of glycerophospholipid with linoleate metabolism. It is noteworthy that

ALA, an essential omega-3 fatty acid, is primarily attributed to its anti-inflammatory,

antioxidant, and lipid-lowering properties in MASLD (44). ALA can influence lipid

metabolism by modulating the expression of genes involved in fatty acid oxidation

and lipid synthesis. It also promotes the activation of peroxisome proliferator-

activated receptors (PPARs), which play a crucial role in lipid metabolism (45).

Previous studies have also reported alterations in phospholipid content in the liver

biopsies of MASH patients, suggesting that impaired phospholipid synthesis is

associated with disease progression (14,15). The low hepatic PC levels and the

altered hepatic PC to PE ratio appear to play significant roles in the development of

MASLD (46). However, the pathophysiology of these lipid-induced processes

remains not fully understood. Moreover, low PC levels have previously been shown

to activate SREBP1 (47). Essential phospholipids (EPLs) rich in PC are a widely

used treatment option for fatty liver disease. There is substantial and consistent

clinical evidence indicating that treatment with EPLs leads to the regression of

steatosis (48). Studies show that phosphatidylcholine metabolism intersects with

various metabolic pathways, and optimizing PC levels could potentially help regulate

lipid metabolism and insulin sensitivity, thereby mitigating the progression of MASLD

(49). Additionally, PC may possess antioxidant properties, which could counteract

oxidative damage in the liver and protect against disease progression (50). In

addition to various lipid classes, the liver metabolite profile in WDsiAATF mice

showed decreased branched-chain amino acid metabolism and sulfur metabolism

involved in the MASH pathogenesis, in contrast to increased pyruvate metabolism,

NAD, and FAD metabolism, as well as one-carbon metabolic pathways. The effect of

AATF silencing on non-lipid metabolic pathways reveals new dimensions of AATF's

functional significance in MASLD (data not shown; manuscript under preparation).

Another intriguing observation in the current study using pathway enrichment

analysis is the upregulation of mitochondrial beta-oxidation of the fatty acid pathway

in WDsiAATF mice. Fatty acid oxidation (FAO) helps to efficiently metabolize fatty

acids, and thereby enhanced beta-oxidation may alleviate inflammation and oxidative

stress in the liver, which are key drivers of MASLD progression (51). Several genes,

such as CPT1α, PPARs, and ACOX1, encode enzymes involved in FAO, and

henceforth, upregulated expression of these genes would enhance fatty acid

oxidation capacity, potentially improving metabolic health (52,53). Consistent with

this, our study also showed the upregulation of FAO genes in the liver tissue of

WDsiAATF compared to WDC. The development of hepatic steatosis is primarily

associated with the excessive accumulation of triglycerides in hepatocytes. Our

research suggests that loss of hepatic AATF may redirect the liver's metabolism from

triglyceride synthesis to the production of diacylglycerol (DAG) and, subsequently,

glycerophospholipids, helping to manage or decrease fat accumulation in the liver.

Furthermore, increased fatty acid oxidation could mobilize the stored triglycerides,

improving metabolic balance, reducing oxidative stress, and potentially alleviating

MASH. An in-depth, targeted lipidomics approach coupled with triglyceride profiling

will yield definitive insights, which is the central focus of our forthcoming research.

Interestingly, the protein-protein interaction network of AATF-interacting proteins, in

conjunction with the transcriptomic profile of WDsiAATF, revealed CD74 as a

possible link between AATF and lipid metabolism. Supporting this, studies by Li et al.

have shown that CD74 levels were positively correlated with triglyceride levels in

human atherosclerotic lesions (54).

Studies by Ishigaki et al. have demonstrated that AATF activates AKT in response to

cellular stress, thereby promoting cell survival (55). AKT phosphorylated at Thr308

activates mTORC1, which particularly plays a dominant role in promoting lipogenesis

and inhibiting lipid breakdown (56, 57). Thus, we evaluated the effect of the AATF-

AKT-mTOR signaling pathway on hepatic lipid metabolism. AATF silencing

effectively inactivated mTORC1 activation by AKT, further affecting the lipogenic

genes such as SREBP1, FASN, and ACC. The inactivated mTORC1 complex led to

the upregulation of LCAD, a key enzyme required for initiating the β-oxidation of fatty

acids. Moreover, mTORC1 suppresses PNPLA2, a key lipase involved in the lipolysis

of triglyceride to diacylglycerol and free fatty acids (58). The stored fat is mobilized in

the form of diacylglycerol, a precursor for glycerophospholipids and free fatty acids,

which undergo β-oxidation for energy production. In our study, AATF silencing

increased PNPLA2 levels, as reflected by decreased triglyceride levels and

upregulated glycerophospholipid synthesis and β-oxidation in WDsiAATF mice.

Despite extensive studies on MASLD, understanding the molecular drivers of lipid

metabolism, insulin resistance, oxidative stress, and inflammation is essential for

addressing its pathogenesis. Although knockout models remain the gold standard for

functional validation, we utilized AAV8-mediated, hepatocyte-specific siRNA delivery

to achieve temporal and liver-restricted silencing of AATF. This approach avoids

developmental compensations inherent to germline knockouts and offers

translational relevance, mimicking therapeutic gene silencing strategies. The use of

the TBG promoter ensured hepatocyte specificity, minimizing off-target effects in

non-parenchymal cells. Given AATF's pleiotropic functions, a complete knockout may

result in systemic effects not confined to liver pathology. Importantly, AAV8 vectors

are widely used in translational gene therapy platforms, providing a clinically relevant

proof of concept. While we acknowledge that future studies using hepatocyte-specific

AATF knockout mice will be essential to validate the long-term impact and specificity

of our findings fully, our results provide compelling evidence for the functional role of

AATF in MASH and establish a foundation for therapeutic targeting. Another

promising alternative is the development of small-molecule inhibitors targeting AATF,

offering a more practical and sustained therapeutic approach. Identifying and

optimizing these inhibitors will be critical for advancing AATF-targeted therapies for

MASH and other metabolic diseases.

In conclusion, this study identifies AATF as a novel transcriptional regulator of lipid

metabolism in the context of MASH. It elucidates a previously unknown AKT–

mTORC1 axis, which is modulated by AATF, in hepatocytes. Our use of liver-specific

AAV8-mediated siRNA delivery enabled sustained and targeted gene silencing,

representing a translationally relevant model for future therapeutic development.

Furthermore, the application of integrated transcriptomics and metabolomics

provides a systems-level understanding of how AATF silencing reprograms lipid

metabolism, particularly through the upregulation of glycerophospholipid biosynthesis

and fatty acid oxidation. These findings open new avenues for precision-targeted

treatments and highlight AATF as a promising candidate for pharmacological

inhibition in MASH.

Declarations

Ethics approval and consent to participate: Not applicable

Consent for publication: Not applicable

Availability of data and materials: Data presented herein are available from the corresponding author upon request. RNA sequencing data have been uploaded to the GEO database (Accession number: GSE280320)

Competing interests: The authors declare that there are no conflicts of interest.

Funding: This study was supported in whole or in part by the SERB-POWER grant (SPG/2021/002524) and Ramalingaswami Re-entry Fellowship from the Department of Biotechnology (BT/RLF/Reentry/58/2017) to DPK, and Senior Research Fellowship (SRF) from ICMR to ANS.

Authors' contributions: Conceptualization: ANS, DPK; Methodology: ANS, DS, MM, PA, DPK; Investigation: ANS, PKS, DS, SPS, SMN, DPK; Formal Analysis: ANS, DS, MM, GR, PKS, DPK; Visualization: ANS, DPK; Resources: ANS, MM, GR, SPS, SMN, PKS, DPK; Supervision: ANS, DPK; Funding Acquisition: DPK; Writing- original draft: ANS, DPK; Writing- Review and Editing: All authors.

Acknowledgements: RNA sequencing was carried out at Molsys Pvt. Ltd. (Molsys Scientific). We thank the DBT-SAHAJ National Facility at Rajiv Gandhi Centre for Biotechnology (RGCB) for their work on metabolomics. We also thank TheraCUES Innovations Pvt Ltd. for assisting with bioinformatic analysis of the transcriptomics and untargeted metabolomics assay data.

The authors also acknowledge funding support from the Department of Biotechnology- Boost to University Interdisciplinary Life Science Departments for Education and Research program [DBT-BUILDER: BT/INF/22/SP43045/2021] and the Department of Science and Technology–Promotion of University Research and Scientific Excellence [DST-PURSE: SR/PURSE/2021/81(c)].

References

1. Stefan N, Yki-Järvinen H, Neuschwander-Tetri BA. Metabolic dysfunction-associated steatotic liver disease: heterogeneous pathomechanisms and effectiveness of metabolism-based treatment. *Lancet Diabetes Endocrinol*. 2025;13(2):134-148. doi:10.1016/S2213-8587(24)00318-8

2. Younossi ZM, Golabi P, Paik JM, Henry A, Van Dongen C, Henry L. The global epidemiology of nonalcoholic fatty liver disease (NAFLD) and nonalcoholic steatohepatitis (NASH): a systematic review. Hepatology. 2023;77(4) 1335-1347 doi:10.1097/HEP.0000000000000004.

3. Targher G, Corey KE, Byrne CD, Roden M. The complex link between NAFLD and type 2 diabetes mellitus - mechanisms and treatments. Nat Rev Gastroenterol Hepatol. 2021;18(9):599-612. doi:10.1038/s41575-021-00448-y

4. Suresh D, Srinivas AN, Kumar DP. Etiology of Hepatocellular Carcinoma: Special Focus on Fatty Liver Disease. Front Oncol. 2020;30;10:601710. doi: 10.3389/fonc.2020.601710.

5. Chan WK, Chuah KH, Rajaram RB, Lim LL, Ratnasingam J, Vethakkan SR. Metabolic Dysfunction-Associated Steatotic Liver Disease (MASLD): A State-of-the-Art Review. J Obes Metab Syndr. 2023;32(3):197-213. doi: 10.7570/jomes23052.

6. Byrne CD, Targher G. NAFLD: a multisystem disease. J Hepatol. 2015;62(1 Suppl):S47-64. doi: 10.1016/j.jhep.2014.12.012.

7. Ratziu V, Francque S, Sanyal A. Breakthroughs in therapies for NASH and remaining challenges. J Hepatol. 2022;76(6):1263-1278. doi:10.1016/j.jhep.2022.04.002

8. European Association for the Study of the Liver (EASL); European Association for the Study of Diabetes (EASD); European Association for the Study of Obesity

(EASO). EASL-EASD-EASO Clinical Practice Guidelines on the management of

metabolic dysfunction-associated steatotic liver disease (MASLD). J Hepatol.

2024;81(3):492-542. doi:10.1016/j.jhep.2024.04.031

9. Rinella, M. E., Neuschwander-Tetri, B. A., Siddiqui, M. S., Abdelmalek, M. F.,

Caldwell, S., Barb, D., et al. AASLD Practice Guidance on the clinical

assessment and management of nonalcoholic fatty liver disease. Hepatology.

2023;77(5):1797-1835. doi:10.1097/HEP.0000000000000323

10. Li Y, Yang P, Ye J, Xu Q, Wu J, Wang Y. Updated mechanisms of MASLD

pathogenesis. Lipids Health Dis. 2024;23(1):117. doi: 10.1186/s12944-024-

02108-x.

11. Rao G, Peng X, Li X, An K, He H, Fu X, et al. Unmasking the enigma of lipid

metabolism in metabolic dysfunction-associated steatotic liver disease: from

mechanism to the clinic. Front Med (Lausanne). 2023;10:1294267. doi:

10.3389/fmed.2023.1294267

12. Venkatesan N, Doskey LC, Malhi H. The Role of Endoplasmic Reticulum in

Lipotoxicity during Metabolic Dysfunction-Associated Steatotic Liver Disease

(MASLD) Pathogenesis. Am J Pathol. 2023;193(12):1887-1899. doi:

10.1016/j.ajpath.2023.08.007.

13. Carli F, Della Pepa G, Sabatini S, Vidal Puig A, Gastaldelli A. Lipid metabolism

in MASLD and MASH: From mechanism to the clinic. JHEP Rep.

2024;6(12):101185. doi:10.1016/j.jhepr.2024.101185

14. Puri P, Baillie RA, Wiest MM, Mirshahi F, Choudhury J, Cheung O, et al. A

lipidomic analysis of nonalcoholic fatty liver disease. Hepatology 2007; 46: 1081-

1090 DOI: 10.1002/hep.21763

15. Chiappini F, Coilly A, Kadar H, Gual P, Tran A, Desterke C, et al. Metabolism

dysregulation induces a specific lipid signature of nonalcoholic steatohepatitis in

patients. Sci Rep 2017; 7: 46658. doi:10.1038/srep46658

16. Marc Prentki, S. R. Murthy Madiraju. Glycerolipid Metabolism and Signaling in

Health   and   Disease.   Endocrine   Reviews,   2008;29(6)   647–676,

https://doi.org/10.1210/er.2008-0007

17. Acoba MG, Senoo N, Claypool SM. Phospholipid ebb and flow makes

mitochondria   go.   J   Cell   Biol.   2020;219(8):e202003131.

doi:10.1083/jcb.202003131

18. Koliaki C, Szendroedi J, Kaul K, Jelenik T, Nowotny P, Jankowiak F, et al.

Adaptation of hepatic mitochondrial function in humans with non-alcoholic fatty

liver   is   lost   in   steatohepatitis.   Cell   Metab.   2015;21(5):739-746.

doi:10.1016/j.cmet.2015.04.004

19. Selen ES, Choi J, Wolfgang MJ. Discordant hepatic fatty acid oxidation and

triglyceride hydrolysis leads to liver disease. JCI Insight. 2021;6(2):e135626.

Published 2021 Jan 25. doi:10.1172/jci.insight.135626

20. Srinivas AN, Suresh D, Mirshahi F, Santhekadur PK, Sanyal AJ, Kumar DP.

Emerging roles of AATF: Checkpoint signaling and beyond. J Cell Physiol.

2021;236(5):3383-3395. doi: 10.1002/jcp.30141.

21. Iezzi S, Fanciulli M. Discovering Che-1/AATF: a new attractive target for cancer

therapy. Front Genet. 2015;6:141. doi: 10.3389/fgene.2015.00141.

22. Jackson JG, Lozano G. Che-ating death: CHE1/AATF protects from p53-

mediated   apoptosis.   EMBO   J.   2012;31(20):3951-3.   doi:

10.1038/emboj.2012.258.

23. Welcker D, Jain M, Khurshid S, Jokić M, Höhne M, Schmitt A, et al. AATF

suppresses apoptosis, promotes proliferation and is critical for Kras-driven lung

cancer. Oncogene. 2018;37(11):1503-1518. doi: 10.1038/s41388-017-0054-6.

24. Haanpää M, Reiman M, Nikkilä J, Erkko H, Pylkäs K, Winqvist R. Mutation

analysis of the AATF gene in breast cancer families. BMC Cancer. 2009;9:457.

doi: 10.1186/1471-2407-9-457.

25. Kaul D, Mehrotra A. Functional characterization of AATF transcriptome in human

leukemic cells. Mol Cell Biochem. 2007;297(1-2):215-20. doi: 10.1007/s11010-

006-9317-1.

26. Suresh D, Srinivas AN, Prashant A, Satish S, Vishwanath P, Nataraj SM, et al.

AATF inhibition exerts antiangiogenic effects against human hepatocellular

carcinoma. Front Oncol. 2023;13:1130380. doi: 10.3389/fonc.2023.1130380.

27. Srinivas AN, Suresh D, Chidambaram SB, Santhekadur PK, Kumar DP.

Apoptosis antagonizing transcription factor-mediated liver damage and

inflammation to cancer: Therapeutic intervention by curcumin in experimental

metabolic dysfunction associated steatohepatitis-hepatocellular carcinoma. J

Cell Physiol. 2024;239(1):135-151. doi: 10.1002/jcp.31151.

28. Srinivas AN, Suresh D, Vishwanath PM, Satish S, Santhekadur PK, Koka S, et

al. TACE inhibition: a promising therapeutic intervention against AATF-mediated

steatohepatitis to hepatocarcinogenesis. Mol Oncol. 2024; doi: 10.1002/1878-

0261.13646.

29. Asgharpour A, Cazanave SC, Pacana T, Seneshaw M, Vincent R, Banini BA, et

al. A diet-induced animal model of non-alcoholic fatty liver disease and

hepatocellular cancer. J Hepatol. 2016;65(3):579-88. doi:

10.1016/j.jhep.2016.05.005.

30. Han J, Kaufman RJ. The role of ER stress in lipid metabolism and lipotoxicity. J
Lipid Res. 2016;57(8):1329-38. doi: 10.1194/jlr.R067595.

31. Perry RJ, Samuel VT, Petersen KF, Shulman GI. The role of hepatic lipids in
hepatic insulin resistance and type 2 diabetes. Nature. 2014;510(7503):84-91.
doi: 10.1038/nature13478.

32. Dempsey JL, Ioannou GN, Carr RM. Mechanisms of Lipid Droplet Accumulation
in Steatotic Liver Diseases. Semin Liver Dis. 2023;43(4):367-382. doi:
10.1055/a-2186-3557.

33. Calder PC. n-3 polyunsaturated fatty acids, inflammation, and inflammatory
diseases. Am J Clin Nutr. 2006;83(6 Suppl):1505S-1519S. doi:
10.1093/ajcn/83.6.1505S.

34. Georgila K, Gounis M, Havaki S, Gorgoulis VG, Eliopoulos AG. mTORC1-
dependent protein synthesis and autophagy uncouple in the regulation of
Apolipoprotein A-I expression. Metabolism. 2020;105:154186.
doi:10.1016/j.metabol.2020.154186

35. Gosis BS, Wada S, Thorsheim C, Li K, Jung S, Rhoades JH, et al. Inhibition of
nonalcoholic fatty liver disease in mice by selective inhibition of mTORC1.
Science. 2022;376(6590):eabf8271. doi: 10.1126/science.abf8271.

36. He A, Chen X, Tan M, Chen Y, Lu D, Zhang X, et al. Acetyl-CoA Derived from
Hepatic Peroxisomal β-Oxidation Inhibits Autophagy and Promotes Steatosis via
mTORC1 Activation. Mol Cell. 2020;79(1):30-42.e4. doi:
10.1016/j.molcel.2020.05.007.

37. Liu DD, Han CC, Wan HF, He F, Xu HY, Wei SH, et al. Effects of inhibiting PI3K-
Akt-mTOR pathway on lipid metabolism homeostasis in goose primary
hepatocytes. Animal. 2016;10(8):1319-1327. doi:10.1017/S1751731116000380

38. Chakrabarti P, English T, Shi J, Smas CM, Kandror KV. Mammalian target of

rapamycin complex 1 suppresses lipolysis, stimulates lipogenesis, and promotes

fat storage. Diabetes. 2010;59(4):775-781. doi:10.2337/db09-1602

39. Soliman GA, Acosta-Jaquez HA, Fingar DC. mTORC1 inhibition via rapamycin

promotes triacylglycerol lipolysis and release of free fatty acids in 3T3-L1

adipocytes. Lipids. 2010;45(12):1089-1100. doi:10.1007/s11745-010-3488-y

40. Fischer J, Lefèvre C, Morava E, Mussini JM, Laforêt P, Negre-Salvayre A, et al.

The gene encoding adipose triglyceride lipase (PNPLA2) is mutated in neutral

lipid storage disease with myopathy. *Nat Genet.* 2007;39(1):28-30.

doi:10.1038/ng1951

41. Duan Y, Pan X, Luo J, Xiao X, Li J, Bestman PL, et al. Association of

Inflammatory Cytokines With Non-Alcoholic Fatty Liver Disease. Front Immunol.

2022;13:880298. doi: 10.3389/fimmu.2022.880298.

42. Kumar DP, Santhekadur PK, Seneshaw M, Mirshahi F, Uram-Tuculescu C,

Sanyal AJ. A Regulatory Role of Apoptosis Antagonizing Transcription Factor in

the Pathogenesis of Nonalcoholic Fatty Liver Disease and Hepatocellular

Carcinoma. Hepatology. 2019;69(4):1520-1534. doi: 10.1002/hep.30346.

43. Cohn, J. S., Wat, E., Kamili, A., and Tandy, S. Dietary Phospholipids, Hepatic

Lipid Metabolism and Cardiovascular Disease. Curr. Opin. Lipidol. 2008;19,

257–262. doi:10.1097/MOL.0b013e3282ffaf96

44. Jordão Candido C, Silva Figueiredo P, Del Ciampo Silva R, Candeloro Portugal

L, Augusto Dos Santos Jaques J, Alves de Almeida J, et al. Protective Effect of

α-Linolenic Acid on Non-Alcoholic Hepatic Steatosis and Interleukin-6 and -10 in

Wistar Rats. Nutrients. 2019;12(1):9. doi:10.3390/nu12010009

45. Yang L, Yuan J, Liu L, Shi C, Wang L, Tian F, et al. α-linolenic acid inhibits

human renal cell carcinoma cell proliferation through PPAR-γ activation and

COX-2 inhibition. Oncol Lett. 2013;6(1):197-202. doi: 10.3892/ol.2013.1336.

46. Arendt BM, Ma DW, Simons B, Noureldin SA, Therapondos G, Guindi M, et al.

Nonalcoholic fatty liver disease is associated with lower hepatic and erythrocyte

ratios of phosphatidylcholine to phosphatidylethanolamine. Appl Physiol Nutr

Metab. 2013;38(3):334-40. doi: 10.1139/apnm-2012-0261.

47. Walker AK, Jacobs RL, Watts JL, Rottiers V, Jiang K, Finnegan DM, et al. A

conserved SREBP-1/phosphatidylcholine feedback circuit regulates lipogenesis

in metazoans. Cell. 2011;147(4):840-52. doi: 10.1016/j.cell.2011.09.045.

48. Dajani AI, Popovic B. Essential phospholipids for nonalcoholic fatty liver disease

associated with metabolic syndrome: A systematic review and network meta-

analysis. World J Clin Cases. 2020;8(21):5235-5249.

doi:10.12998/wjcc.v8.i21.5235

49. Osipova D, Kokoreva K, Lazebnik L, Golovanova E, Pavlov C, Dukhanin A, et al.

Regression of Liver Steatosis Following Phosphatidylcholine Administration: A

Review of Molecular and Metabolic Pathways Involved. Front Pharmacol.

2022;13:797923. doi: 10.3389/fphar.2022.797923.

50. Maev IV, Samsonov AA, Palgova LK, Pavlov CS, Vovk EI, Shirokova EN,

Starostin KM. Effectiveness of phosphatidylcholine in alleviating steatosis in

patients with non-alcoholic fatty liver disease and cardiometabolic comorbidities

(MANPOWER study). BMJ Open Gastroenterol. 2020;7(1):e000341. Published

2020 Jan 13. doi:10.1136/bmjgast-2019-000341

51. Naguib G, Morris N, Yang S, Fryzek N, Haynes-Williams V, Huang WA, et al.

Dietary fatty acid oxidation is decreased in non-alcoholic fatty liver disease: A

palmitate breath test study. Liver Int. 2020;40(3):590-597. doi:10.1111/liv.14309

52. Fondevila MF, Fernandez U, Heras V, Parracho T, Gonzalez-Rellan MJ, Novoa

E, et al. Inhibition of carnitine palmitoyltransferase 1A in hepatic stellate cells

protects against fibrosis. J Hepatol. 2022;77(1):15-28. doi:

10.1016/j.jhep.2022.02.003.

53. Régnier M, Polizzi A, Smati S, Lukowicz C, Fougerat A, Lippi Y, et al.

Hepatocyte-specific deletion of Pparα promotes NAFLD in the context of obesity.

Sci Rep. 2020;10(1):6489. doi: 10.1038/s41598-020-63579-3.

54. Li W, Sultana N, Yuan L, Forssell C, Yuan XM. CD74 in Apoptotic Macrophages

Is Associated with Inflammation, Plaque Progression and Clinical Manifestations

in Human Atherosclerotic Lesions. Metabolites. 2022;12(1):54. doi:

10.3390/metabo12010054.

55. Ishigaki S, Fonseca SG, Oslowski CM, Jurczyk A, Shearstone JR, Zhu LJ, et al.

AATF mediates an antiapoptotic effect of the unfolded protein response through

transcriptional regulation of AKT1. Cell Death Differ. 2010;17(5):774-786.

doi:10.1038/cdd.2009.175

56. Memmott RM, Dennis PA. Akt-dependent and -independent mechanisms of

mTOR regulation in cancer. Cell Signal. 2009;21(5):656-664.

doi:10.1016/j.cellsig.2009.01.004

57. Cui J, Zhang F, Wang Y, Liu J, Ming X, Hou J, et al. Macrophage migration

inhibitory factor promotes cardiac stem cell proliferation and endothelial

differentiation through the activation of the PI3K/Akt/mTOR and AMPK

pathways. Int J Mol Med. 2016;37(5):1299-309. doi: 10.3892/ijmm.2016.2542.

58. Grabner GF, Xie H, Schweiger M, Zechner R. Lipolysis: cellular mechanisms for

lipid mobilization from fat stores. Nat Metab 2021;3, 1445–1465.

doi.org/10.1038/s42255-021-00493-6.

59. Srinivas AN, Suresh D, Suvarna D, Pathak P, Giri S, Suman, et al. Unraveling

the Potential Role of Tecomella undulata in Experimental NASH. Int J Mol Sci.

2023;24(4):3244. doi: 10.3390/ijms24043244.

60. Banini BA, Kumar DP, Cazanave S, Seneshaw M, Mirshahi F, Santhekadur PK,

et al. Identification of a Metabolic, Transcriptomic, and Molecular Signature of

Patatin-Like Phospholipase Domain Containing 3-Mediated Acceleration of

Steatohepatitis. Hepatology. 2021; doi: 10.1002/hep.31609.

61. Poynard T, Munteanu M, Charlotte F, et al. Impact of steatosis and inflammation

definitions on the performance of NASH tests. Eur J Gastroenterol Hepatol.

2018;30(4):384-391. doi:10.1097/MEG.0000000000001033

62. Brunt EM, Kleiner DE, Wilson LA, Sanyal AJ, Neuschwander-Tetri BA;

Nonalcoholic Steatohepatitis Clinical Research Network. Improvements in

Histologic Features and Diagnosis Associated With Improvement in Fibrosis in

Nonalcoholic Steatohepatitis: Results From the Nonalcoholic Steatohepatitis

Clinical Research Network Treatment Trials. Hepatology. 2019;70(2):522-531.

doi:10.1002/hep.30418

63. Liu Z, Wang P, Liu Z, Wei C, Li Y, Liu L. Evaluation of liver tissue extraction

protocol for untargeted metabolomics analysis by ultra-high-performance liquid

chromatography/tandem mass spectrometry. J Sep Sci. 2021;44(18):3450-3461.

doi: 10.1002/jssc.202100051

Figure Legends

Figure 1. Hepatic silencing of AATF improves metabolic health in experimental

MASH

(A) Study design- C57Bl/6 mice were fed chow diet normal water (CD) or western

diet sugar water (WD) for 12 weeks. Liver-specific AAV8-TBG-siControl (Control) or

AAV8-TBG-siAATF (siAATF) were injected via the tail vein into mice and monitored

for an additional 12 weeks. (B) Body weight, (C) liver weight, (D) liver-to-body weight

ratio, and (E) adipose tissue weight were determined. Mean±SEM for 6-8 mice per

group, ##p<0.001 compared to CD control; **p<0.001 compared to WD control,

unpaired t-test. (F) Glucose tolerance test (G) Insulin tolerance test were performed.

Mouse serum was collected after 24 weeks. (H) Fasting blood glucose, (I) fasting

insulin, and (J) HOMA-IR were determined. Mean±SEM for 6-8 mice per group,

##p<0.001 compared to CD control; **p<0.001 compared to WD control, unpaired t-

test. AAV; adeno-associated virus, TBG; tyrosine binding globulin, AATF; apoptosis

antagonizing transcription factor, CD; chow diet, WD; western diet.

1051

Figure 2. Hepatic loss of AATF improves liver injury and impaired lipid

metabolism in MASH mice

Mice were euthanized after 24 weeks, and blood was collected. Serum levels of (A)

AST, (B) ALT, (C) total cholesterol, (D) triglycerides, and (E) LDL-cholesterol were

measured. Mean±SEM for 6-8 mice per group, ##p<0.001 compared to CD control;

**p<0.001 compared to WD control, unpaired t-test. Representative (F) gross liver

images, (G) H&E staining, and (H) Oil Red O staining of the mouse liver tissues. n=6

mice per group. Scale bar 50 μm. (I) Histology scores were calculated for the liver

sections of mice. Representative images and scoring of steatosis, lobular

inflammation, hepatocyte ballooning, and MASLD activity score. Mean ±SEM for 6-8

mice per group. Scale bar 50 μm, unpaired t-test. **p<0.001 compared to WD

control. HOMA-IR; homeostatic model assessment for insulin resistance, ALT,

alanine transaminase; AST, aspartate transaminase; LDL; low-density lipoprotein,

MASLD; metabolic dysfunction associated steatotic liver disease, CD; chow diet,

WD; western diet.

Figure 3: Effect of AATF silencing on hepatic transcriptomic and metabolic

profiles in experimental MASH

C57Bl/6 mice were fed chow diet normal water (CD) or western diet sugar water

(WD) for 12 weeks. Liver-specific AAV8-TBG-siControl (Control) or AAV8-TBG-

siAATF (siAATF) were injected via the tail vein into mice and monitored for an

additional 12 weeks. After mice were euthanized, liver tissues were subjected to

transcriptomic and untargeted metabolomic analyses. In transcriptomics, the PLS-DA

plot (A) among the four groups shows a clear separation between the groups. (B)

sample-sample correlation within the four groups. (C) Volcano plot for differentially

expressed genes in WDsiAATF vs. WDC groups. The x-axis denotes fold change in

expression (log2 scale), and the y-axis denotes adjusted p-value (negative log10

scale) for the analyzed genes in the dataset (each dot represents a single gene). Red

and green dots represent genes significantly upregulated and downregulated,

respectively. Blue dots (p<0.05) and grey dots represent nonsignificant genes.

(DESeq2; cut-offs used: adjusted p-value < 0.05 and log fold change ≥ 2). (D) Top 10

up- and downregulated KEGG pathways, GO-BP, and GO-MF in WDsiAATF vs.

WDC groups. In metabolomics, (E) PLS-DA plot among the four groups shows a

clear group separation between the groups. (F) Venn diagram showing the up- and

downregulated metabolites between the WDC vs. CDC and WDsiAATF vs. WDC.

(G) Volcano plot for differential metabolites in WDsiAATF vs. WDC. Red and green

dots represent metabolites significantly upregulated and downregulated, respectively.

Blue dots (p<0.05) and grey dots represent nonsignificant metabolites. n=3. CD;

chow diet, WD; western diet, AAV; adeno-associated virus, TBG; tyrosine binding

globulin, AATF; apoptosis antagonizing transcription factor, PLS-DA; partial least

squares discriminant analysis, WDC; western diet control, KEGG; Kyoto

encyclopedia of genes and genomes, GO; gene ontology, BP; biological processes,

MF; molecular function.

Figure 4. Effect of silencing hepatic AATF on cellular stress and inflammation

in experimental MASH

Transcriptomic analysis of liver tissues from mice fed with chow diet normal water
(CD) or western diet sugar water (WD), administered with AAV8-TBG-siControl or
AAV8-TBG-siAATF, n=3 (A) p-value plot of cellular stress-related pathways. (B)
Heatmap of cellular stress-related genes. (C) hepatic mRNA expression of ER stress
markers- CHOP, Grp78, PERK, and ATF4. Mean±SEM, n=4, unpaired t-test. (D) p-
value plot of inflammation-related pathways. (E) Heatmap of inflammation-related
genes. (F) hepatic mRNA expression of inflammatory markers- IL-1β, IL-6, and TNF-
α, and serum TNF-α by ELISA. Liver tissue lysates were used for immunoblotting of
inflammatory markers- (G) pJNK and (H) pERK1/2. (I) F4/80 staining, Scale bar 50
μm. Mean ±SEM, n=4, unpaired t-test. $^{##}$p<0.001 or $^{#}$p<0.05 compared to CD control;
**p<0.001 or *p<0.05 compared to WD control. CD; chow diet, WD; western diet,
AAV; adeno-associated virus, TBG; tyrosine binding globulin, AATF; apoptosis
antagonizing transcription factor, CHOP; C/EBP homologous protein, GRP78;
glucose-regulated protein 78, PERK; protein kinase R (PKR)-like endoplasmic
reticulum kinase, ATF4; activating transcription factor 4, IL; interleukins, TNF-α;
tumor necrosis factor-alpha, ELISA; enzyme-linked immunosorbent assay.

Figure 5. Silencing of hepatic AATF ameliorates liver fibrosis

Transcriptomic analysis of liver tissues (n=3) showing (A) p-value plot of fibrosis-
related pathways. (B) Heatmap of fibrosis-related genes. (C) Hepatic mRNA
expression of Col1A1, Col3A1, α-SMA, and TGF-β quantified. Mean ±SEM, n=4,
unpaired t-test. (D) Representative images of picrosirius red staining with fibrosis
scoring. (E) Desmin immunostaining with scoring. Scale bar 50 μm. Mean ±SEM for
6-8 mice, unpaired t-test. $^{##}$p<0.001 or $^{#}$p<0.05 compared to CD control; **p<0.001
compared to WD control. Col; collagen, α-SMA; alpha-smooth muscle actin, TGF-β;
transforming growth factor-beta. CD; chow diet, WD; western diet

Figure 6. Effect of silencing hepatic AATF on lipid metabolism in MASH mice

Transcriptomic analysis of liver tissues (n=3) showing (A) p-value plot of lipid
metabolism-related pathways. (B) Heatmap of lipid metabolism-related genes. (C)

Hepatic mRNA expression of FASN, ACC1, ACC2, and SCD1 was evaluated. Mean

±SEM, n=4, unpaired t-test. Untargeted metabolomics was carried out in mouse liver

tissues (n = 3). (D) Bubble plots showing the top 20 KEGG pathways enrichment, (E)

lipid superclass, and (F) lipid subclass using the mummichog GSEA algorithm in

WDC vs. WDsiAATF. P<0.05 was considered significant. CD; chow diet, WD;

western diet, AAV; adeno-associated virus, TBG; tyrosine binding globulin, AATF;

apoptosis antagonizing transcription factor, FASN; fatty acid synthase, ACC; acetyl-

CoA carboxylase, SCD; stearoyl-coa 9-desaturase, KEGG; Kyoto encyclopedia of

genes and genomes, GSEA; gene set enrichment analysis.

Figure 7. Metabolic networking of glycerophospholipid metabolism

Transcriptomics and untargeted metabolomics analyses were performed on mouse

liver tissues (n = 3). Metabolites of glycerophospholipid metabolism were analyzed,

and a functional correlation network analysis was performed. (A) The

glycerophospholipid metabolism pathway, visualized in the Pathview network,

highlights the dysregulated compounds (up- or downregulated) in WDsiAATF

compared to WDC. (B) Relative abundance of phosphocholine, PC

(phosphatidylcholine), LPE (lysophophatidylethanolamine), and LPC

(lysophosphatidylcholine). **p<0.001 or *p<0.05 compared to WDC. (C) Heatmap of

genes involved in glycerophospholipid metabolism. (D) Debiased sparse partial

correlation network plot visualizing the major relationships of glycerophospholipid

metabolic alterations with the clinical variables. Statistical significance was analyzed

by a student's t-test. P<0.05 was considered significant. CD; chow diet, WD; western

diet, AATF; apoptosis antagonizing transcription factor.

Figure 8. Hepatic loss of AATF alters lipid metabolism via AKT-mTORC1

signaling pathways in MASH mice

Transcriptomics and untargeted metabolomics analyses were performed on mouse

liver tissues (n = 3). Pathway analysis and enrichment analysis were performed on

both datasets. An enrichment analysis based on the Small Molecule Pathway

Database (SMPDB) was performed on metabolites. (A) upregulated and downregulated pathways. (B) Heatmap of genes involved in fatty acid oxidation. (C) hepatic mRNA expression of CPT-1α, ACOX1, PGC-1α, and ACSL1. (D) Liver tissue lysates were used for immunoblotting of pAKT (Thr308), pmTOR, and Raptor. Hepatic mRNA expression of (E) SREBP1, (F) LCAD, and (G) PNPLA2. Mean±SEM, n=4, unpaired t-test, **P < 0.001 or *P < 0.05 compared to WDC. (H) Schematic representation of the effect of hepatic AATF silencing in experimental MASH. ACOX1; Acyl-CoA Oxidase 1, PGC1; Peroxisome proliferator-activated receptor gamma coactivator-1alpha, ACSL1; acyl-CoA synthetase long-chain family member 1, AKT; protein kinase B, mTOR; mammalian target of rapamycin, SREBP1; sterol regulatory element-binding protein 1, LCAD; long-chain acyl-CoA dehydrogenase, PNPLA2; patatin-like phospholipase domain-containing protein 2.

Supplemental Figure 1: (A) Tissue distribution of Control and siAATF in the experimental mice. (B) liver-specific, and (C) hepatocyte-specific AATF silencing as determined by qRT-PCR and western blot, respectively. Mean±SEM for n=4 mice per group, **p<0.001 compared to AAV8 control, unpaired t-test.

Supplemental Figure 2: (A) Weekly measurement of body weight from week 12 to week 24, followed by the administration of AAV8-TBG-siControl (Control) and AAV8-TBG-siAATF (siAATF) at week 12 to the chow diet normal water (CD) and western diet sugar water (WD) fed mice. (B) Food intake and (C) Calorie intake in mice. Mean±SEM for 6-8 mice per group, one-way ANOVA.

Supplemental Figure 3: (A) Volcano plot presentation of differentially expressed genes between liver samples of mice from WDC and CDC. (B) Volcano plot presentation of differentially expressed genes between liver samples of mice from CDsiAATF and CDC. (C) Volcano plot presentation of differential metabolites from liver samples of mice from WDC and CDC. (D) Volcano plot presentation of differential metabolites from liver samples of mice from CDsiAATF and CDC. X-axis denotes fold change in expression (log2 scale), y-axis denotes adjusted p-value

(negative log10 scale) for the analyzed genes or metabolites in the data set (each dot

represents a single gene or metabolite). Red and green dots represent genes or

metabolites significantly upregulated and downregulated, respectively. Blue dots

(p<0.05) and grey dots represent nonsignificant genes or metabolites. (DESeq2; cut-

offs used: adjusted p-value < 0.05 and log fold change ≥ 2).

Supplemental Figure 4: Metabolite functional interaction network of

glycerophospholipid metabolism pathway in WDsiAATF and WDC.

Supplemental Figure 5: Protein-protein interaction of AATF interacting proteins with

the lipid metabolism-related proteins was determined using BioGRID and STRING

databases and visualized in Cytoscape.

Supplemental Table 1: List of primer sequences used in qRT-PCR in the study.

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8