Tonotopy is not preserved in a descending stage of auditory cortex

Miaoqing Gu[1,2,10], Shanshan Liang[3,10], Jiahui Zhu[2,3], Ruijie Li[3], Ke Liu[3], Xuanyue Wang[4],

Frank Ohl[5,6,7], Yun Zhang[4], Xiang Liao[3,4], Chunqing Zhang[3], Hongbo Jia[1,5,9], Yi Zhou[2,8,*],

Jianxiong Zhang[3,*], Xiaowei Chen[3,4,*]

[1]School of Physical Science and Technology, Guangxi University, Nanning 530004,

China.

[2]Guangxi Key Laboratory of Special Biomedicine and Advanced Institute for Brain and

Intelligence, School of Medicine, Guangxi University, Nanning, 530004, China

[3]Brain Research Center and State Key Laboratory of Trauma and Chemical Poisoning,

Third Military Medical University, Chongqing 400038, China.

[4]Chongqing Institute for Brain and Intelligence, Guangyang Bay Laboratory,

Chongqing 400064, China.

[5]Leibniz Institute for Neurobiology (LIN), Magdeburg, Germany.

[6]Insitute of Biology (IBIO), Otto-von-Guericke University, Magdeburg, Germany.

[7]Center for Behavioral Brain Sciences (CBBS), Magdeburg, Germany.

[8]Department of Neurobiology, Chongqing Key Laboratory of Neurobiology, School of

Basic Medicine, Third Military Medical University, Chongqing 400038, China.

[9]Brain Research Instrument Innovation Center, Suzhou Institute of Biomedical

Engineering and Technology, Chinese Academy of Sciences, Suzhou 215163, China

[10]These authors contributed equally to this work.

*Correspondence: yzhou@tmmu.edu.cn (Y.Z.), jianxiong_zhang1988@tmmu.edu.cn

(J.Z.), xiaowei_chen@tmmu.edu.cn (X.C.)

Abstract

Previous studies based on layer specificity suggest that ascending signals from the

thalamus to sensory neocortex preserve spatially organized information, but it

remains unknown whether sensory information descending from sensory neocortex

to thalamus also maintains such spatial organization pattern. By focusing on

projection specificity, we mapped tone response properties of two groups of cortical

neurons in the primary auditory cortex (A1), based on the relationship between their

specific connections to other regions and their function in ascending

(thalamocortical recipient, TR neurons) or descending (corticothalamic, CT neurons)

auditory information. A clear tonotopic gradient was observed among TR, but not CT

neurons. Additionally, CT neurons exhibited markedly higher heterogeneity in their

frequency tuning and had broader bandwidth than TR neurons. These results reveal

that the information flow descending from A1 to thalamus via CT neurons does not

arrange tonotopically, suggesting that the descending information flow possibly

contributes to higher-order feedback processing of diverse auditory inputs.

Introduction

In the mammalian auditory system, one of the most prominent features is the

tonotopic organization — a spatially ordered gradient of neuronal frequency

preference (Bizley et al., 2005; Kajikawa et al., 2005; Merzenich et al., 1976; Morel et

al., 1993; Nelken et al., 2004; Reale and Imig, 1980; Stiebler et al., 1997; Tani et al.,

2018). Clear tonotopic maps have been found in the auditory cortex (AuC) of many

mammalian species including humans (Clopton et al., 1974; Humphries et al., 2010),

macaque monkeys (Hackett et al., 1998), cats (Lee et al., 2004), ferrets (Bizley et al.,

2005), Mongolian gerbils (Budinger et al., 2000; Ohl et al., 2000; Thomas et al., 1993),

rats (Polley et al., 2007), and mice (Guo et al., 2012; King et al., 2018). Tonotopy

originates in the cochlea and is relayed and preserved throughout all the ascending

stages including the medial geniculate body (MGB) and the AuC (Jasmin et al., 2019;

Simon et al., 2009; Smith and Wever, 1949). As one of the most explored pathways in

the auditory system, the connections between MGB and AuC play an essential role in

the perception of auditory information (Lee, 2013; Ohga et al., 2017; Pardi et al.,

2020).

Core fields of the AuC receive their predominant thalamic input from ventral

division of the MGB (MGBv), which confers well-defined frequency tuning arranged

into smoothly varying tonotopic gradients (Andersen et al., 1980; Merzenich and

Brugge, 1973; Winer et al., 2005). Non-core AuC fields are innervated by non-primary

divisions of the MGB and from intracortical inputs originating outside of the AuC

(Jones, 2001; Lee and Winer, 2005; Reale and Imig, 1980), which show weak

tonotopy and selectivity for processing conspecific communication sounds

(Norman-Haignere et al., 2015; Schneider and Woolley, 2013). Feedforward auditory

information is conveyed from the MGBv to the AuC, primarily received in layer 4 (L4)

but also extending to other layers (Constantinople and Bruno, 2013; Petreanu et al.,

2009). Subsequently, it is projected corticofugally to downstream targets from either

layer 5 (L5) or layer 6 (L6) of the AuC (Shepherd and Yamawaki, 2021), including

feedback information to MGB (Happel et al., 2014; Happel et al., 2010; Homma and

Bajo, 2021; Kirchgessner et al., 2020). Several studies have revealed the sophisticated

inputs and the ability in the perception of complex sounds of CT neurons in A1

(Clayton et al., 2021; Homma et al., 2017), suggesting the descending information

flow in the core fields of the AuC possibly contributes to higher-order feedback

processing of diverse auditory inputs.

It is also known that the tonotopic gradient is well preserved across all layers

within the A1 (Guo et al., 2012; Montes-Lourido et al., 2021; Tischbirek et al., 2019;

Winkowski and Kanold, 2013). However, all of these previous discoveries in tonotopic

mapping are based on layer specificity. While the laminar structure is closely

associated with functional projections in the AuC, they are not precisely identical.

Neurons within the same layer can exhibit diversity in their molecular, morphological,

physiological, and connectional features (Triarhou, 2021; Yarmohammadi et al.,

2014), and therefore, categorizing cortical neurons based on their projection

specificity can provide a deeper fundamental understanding of their functional

organization. The A1-MGB projecting neurons play an essential role in the

feedforward-feedback loop of the thalamus-neocortex interplay for integrated sound

information processing flow (Happel et al., 2014; Happel et al., 2010) and the

perception of complex sounds (Clayton et al., 2021; Homma et al., 2017). However,

due to technical difficulties, there has been limited understanding of the functional

properties of these specific corticothalamic projecting neurons (Antunes and

Malmierca, 2021; Harris et al., 2019; Sherman and Guillery, 1998; Winer et al., 2001)

because they are located in deeper layers of the cortex and more difficult to be

accessed by means of commonly-used, single-cell-resolving recording techniques

such as conventional two-photon (2P) microscopy (Kobat et al., 2009; Oheim et al.,

2001; Takasaki et al., 2020; Tischbirek et al., 2015).

Recent work has established a rabies-based retrograde labeling method with

pathway specificity (Sun et al., 2019; Zhu et al., 2020) to circumvent the neuropil

contamination problem for conventional 2P imaging for CT neurons. Here, using viral

tracing and 2P Ca^{2+} imaging of awake head-fixed mice, we systematically revisited the

functional organization in the A1, specifically targeting two types of neurons with

projection specificity: TR (Zingg et al., 2017) and CT neurons (Gu et al., 2023), akin to

the input and output of the AuC. The projection specificity-based approach provides

detailed insight into the differences in population activity and frequency

response-related topographic organization between neurons with different

connectivity specificity.

Results

Pathway-specific labeling of TR and CT neurons

To label TR neurons, we used an anterograde trans-synaptic tracing strategy by

injecting a Cre-recombinase-expressing AAV in the MGB, then injecting a second AAV

carrying Cre-dependent GCaMP6s (AAV2/9-CaMKⅡ-DIO-GCaMP6s) in the AuC

(Figure 1A and B) (Zingg et al., 2017). Neurons labeled with AAV-GCaMP6s were

observed across all layers in AuC (Figure 1C and D). To label CT neurons, we

performed retrograde tracing with a rabies virus CVS-N2c-derived vector

(CVS-GCaMP6s) by injecting it into MGB (Figure 1E). Neurons labeled with

CVS-GCaMP6s were restricted to AuC L6 (Figure 1F and G) and then were imaged by

2P Ca^{2+} imaging, in line with our recent work (Gu et al., 2023). Whole-cell

patch-clamp recordings and in vivo 2P imaging showed no change in

electrophysiological, morphological or functional characteristics of CVS-labeled CT

neurons (Gu et al., 2023). To locate A1, we conducted wide-field imaging in awake

GCaMP6s-labeled mice, and defined that low-frequency tone (4 kHz) elicited spatially

restricted responses in the regions as "low-frequency (LF) area" and high-frequency

tone (32 kHz) elicited responses in the regions as "high-frequency (HF) area" (Figure

1H). In AAV-GCaMP6s labeled mice, we identified the locations of A1, anterior

auditory field (AAF), and secondary auditory field (A2) based on the known tonotopy

in mice (Figure 1H, left panel) (Issa et al., 2014; Liu et al., 2019).

It is worth noting that, in CVS-GCaMP6s labeled (i.e., CT neurons) mice, pure

tone stimulation elicited wide-field signals in only one auditory region that exhibited

a tonotopic gradient (Figure 1H, right panel). To identify this region in AuC, we

injected a retrograde tracing indicator cholera toxin subunit B conjugated with Alexa

555 (CTB-555) into this region, which resulted in clearly visible CTB-555 labeled cell

bodies in the MGBv (Figure 1I), suggesting that CVS-GCaMP6s labeled neurons were

located in A1(Rothschild et al., 2010). When performing 2P imaging, the average

power delivered to the brain was in the range of 30 - 120 mW, depending on the

depth of imaging. To determine whether neurons labeled with the CVS retained

normal response properties after deep 2P imaging, with higher laser power than

superficial layers imaging, we observed the broadband noise-evoked (BBN-evoked)

responses of these neurons over time (days) (Figure 1- figure supplementary 1A). At

7 and 13 days after injection, we found that CVS-labeled neurons had completely

normal response properties (Figure 1- figure supplementary 1B and C).

Next, we performed Ca^{2+} imaging of TR or CT neurons from A1 identified by

wide-field imaging (Figure 1J). High-titer AAV1-Cre virus has been reported to

anterogradely label postsynaptic neurons and also retrogradely label presynaptic

neurons (Zingg et al., 2017). To minimize this labeling ambiguity, we restricted the 2P

imaging depth of TR neurons to 200 - 400 µm, without imaging the neurons in L5 and

L6, which were reciprocally connected with MGB (Harris and Mrsic-Flogel, 2013).

These results suggest the feasibility of our approach to label and record population

Ca^{2+} signals with pathway specificity at single-cell resolution in awake mice.

CT neurons exhibit no tonotopic gradient

To study tonal response profiles, we delivered 330 pure tones at each focal plane (11

frequencies ranging from 2 - 40 kHz, 6 attenuation levels, 5 repetitions). We

determined the best frequency (BF) (Guo et al., 2012) of all tuned TR (Figure 2A) or

CT neurons (Figure 2B) at each 2P imaging focal plane. In total, we imaged 1041 TR

neurons in 5 animals. 46% were "tuned" neurons, 41% neurons were "irregular"

neurons and 13% were "silent" neurons (Figure 2- figure supplementary 2A, the

definitions of "irregular", "tuned" and "silent" see METHOD DETAILS). For CT neurons,

we imaged a total of 2721 neurons from 10 animals. 18% were "tuned" neurons, 57%

neurons were "irregular" neurons and 25% were "silent" neurons (Figure 2- figure

supplementary 2B).

Based on the BFs, we determined the median BF of all tuned neurons in each 2P

focal plane (e.g., Figure 2C and D) and related them to the position within the

wide-field imaging areas (Figure 2E and F), which were then used for computing the

tonotopic gradient. For TR neurons, we observed a clear tonotopic axis in the

organization of median BF (R = 0.77, P < 0.001; 2.13 octaves/mm; Figure 2G).

However, no such tonotopic gradient was observed in CT neurons (R = 0.20, P = 0.21;

0.21 octaves/mm; Figure 2H). Plotting the relative positions of all individual tuned TR

or CT neurons in each mouse revealed a significant low-to-high frequency gradient

from caudal to rostral for TR neurons (R = 0.52, P < 0.001; 1.76 octaves/mm; Figure 2I

and J) but an absence of tonotopy in CT neurons (R = 0.079, P = 0.084; 0.37

octaves/mm; Figure 2K and L). These results at single-neuron resolution confirmed

that TR neuronal population does possess tonotopy, as previous reports have shown

on a coarse scale (Kalatsky et al., 2005; Wu et al., 2006).

The fraction of different BF responsive neurons among TR and CT neurons

showed that both TR and CT neurons displayed no obvious pattern throughout A1

(Figure 2M). When categorized into LF or HF areas, higher proportions of TR neurons

responded to low-frequency sounds in the LF areas or high-frequency sounds in the

HF areas, whereas CT neurons showed no frequency preference in either area

(Figure 2N and O). Note the obvious discrepancy that the 2P imaging data here

shows no tonotopic gradient of L6 CT neurons, but widefield epifluorescence imaging

data shows a presence of tonotopic gradient from the same mice (Figure 1H), which

could be largely due to differences in the imaging methods and will be discussed

later on.

CT neurons exhibit high heterogeneity in tuning preference

Next, we performed analysis of BF spatial distribution within each focal plane.

Examination of BF distribution among TR or CT neurons revealed that TR neurons

exhibited similar frequency preference in HF or LF areas (Figure 3A), whereas CT

neurons displayed high heterogeneity in their frequency preferences, regardless of LF

or HF areas (Figure 3B).

To quantify the degree of BF heterogeneity, we examined BF spatial distribution

at the focal-plane, local, and nearest-neighboring scales. To this end, the

inter-quartile range of BF (IQR_{BF}, in octaves) was calculated for all tuned neurons in

each focal plane (Figure 3C and D). Quantitative analysis showed that CT neurons had

higher IQR_{BF} than TR neurons across all focal planes (Figure 3C). Similarly, evaluation

of LF or HF areas indicated that CT neurons had significantly higher IQR_{BF} than that of

TR neurons in LF areas, but showed no significant difference from TR neurons in HF

areas at the focal-plane level (Figure 3D).

To assess tone response heterogeneity at the local scale, we then calculated

IQR_{BF} for all tuned neurons within a 25 µm radius (Winkowski and Kanold, 2013; Zeng

et al., 2019) around each randomly selected CT or TR neuron (Figure 3E-G). The

results revealed that TR neurons also exhibited lower median IQR_{BF} than CT neurons

at the local scale (Figure 3F). Consistent with the focal plane scale, CT neurons had

significantly higher local IQR_{BF} than TR neurons in LF areas (Figure 3G). Unexpectedly,

the local IQR$_{BF}$ of TR neurons in HF areas was significantly higher than that of TR

neurons in LF areas (Figure 3G). Further analysis across local regions revealed that

the local IQR$_{BF}$ of TR neurons followed a low-to-high gradient along the

caudal-to-rostral axis (R = 0.2655, *P* = 0.0032; Figure 3H).

The analysis of Δfrequency (BF variability) (Bandyopadhyay et al., 2010; Zeng et

al., 2019) between nearest-neighboring TR or CT neurons was consistent with results

of local IQR$_{BF}$ analysis (Figure 3I). That is, Δfrequency followed a clear gradient in the

organization of TR neurons along the caudal-to-rostral axis (R = 0.2463, *P* < 0.001;

Figure 3J). Moreover, TR neurons, but not CT neurons, displayed high IQR$_{BF}$ in HF

areas compared to LF areas (Figure 3G and I). These results suggest the relative

heterogeneity of frequency preference among CT neurons but homogeneity among

TR neurons.

Receptive field properties of TR versus CT neurons

Given the above differences in the functional organization of frequency preference

between TR and CT neurons, we next focused our analysis on the frequency

responsive areas (FRAs) of individual neurons, as the FRAs directly reflect neuronal

auditory selectivity essential for sound processing (Sadagopan and Wang, 2008). For

this analysis, all tuned neurons were categorized into either a V-shaped (decreasing

frequency selectivity with increasing intensity), I-shaped (narrow, level-tolerant

tuning), or O-shaped (non-monotonic) FRA (Figure 4A and B). We found that a higher

proportion of CT neurons was associated with V-shaped FRAs than TR neurons,

whereas the proportion of CT neurons associated with I-shaped FRAs was lower than

that of TR neurons (Figure 4C). By contrast, no difference was detected in the

proportions of CT and TR neurons associated with O-shaped FRAs (Figure 4C).

A key feature of neuronal tuning curves is their sharpness. Common measures

of sharpness in V shape are half-peak bandwidth and the "quality factor" (Q), which

was obtained by dividing the BF of the neuron by a measure of tuning (Micheyl et al.,

2013), the width of the tuning curve at half-peak in this study. This analysis showed

that the half-peak bandwidth of CT neurons was significantly wider than that of TR

neurons (Figure 4D), while the Q value of CT neurons was lower than that of TR

neurons, especially in the HF areas (Figure 4E and F). Since higher Q values indicate

sharper tuning of neurons, which suggests higher frequency discrimination (Micheyl

et al., 2013), these results suggest that TR neurons might be responsible for finer

tone discrimination.

As the bandwidth of neurons' FRA can also reflect the selectivity of overall tonal

responses (Rodrigues-Dagaeff et al., 1989; Schreiner and Sutter, 1992), we examined

the maximum FRA width of TR or CT neurons (Figure 4G-J). BW_{max} was defined as the

maximum FRA width at any sound level. The quantitative analysis showed that the

BW_{max} of TR neurons was significantly narrower than that of CT neurons, irrespective

of the FRA shape (Figure 4G). In addition, the distribution of TR neurons generally

peaked at a BW_{max} of 1 - 2 octaves, while the distribution of CT neurons peaked at a

BW_{max} of 3 - 4 octaves (Figure 4H). The statistical analysis showed that CT neurons

had significantly higher BW_{max} than TR neurons (Figure 4I). Within LF or HF areas, TR

neurons had a significantly narrower BW_{max} range than CT neurons in either area.

Surprisingly, TR neurons in HF areas had a significantly lower BW_{max} than TR neurons

in LF areas (Figure 4J), suggesting that the BW_{max} distribution among TR neurons

might change along the caudal-to-rostral axis.

We then investigated the organization for intensity tuning, another crucial

feature in auditory perception, based on the monotonicity index (MI, see Methods

for details), wherein 1 indicated monotonic response increase with intensity and 0

indicated strong nonmonotonic tuning (de la Rocha et al., 2008; Sutter and Schreiner,

1995; Watkins and Barbour, 2011). The MI distribution across both TR and CT

neuronal populations peaked at 1 (Figure 4K). Subsequent categorization as either

monotonic or non-monotonic (using a criterion of MI = 0.5 (de la Rocha et al., 2008;

Moore and Wehr, 2013), neurons with an index < 0.5 considered intensity-tuned),

showed that the fraction of neurons exhibiting non-monotonic intensity response did

not significantly differ between TR and CT neurons (Figure 4L). However, examination

of MI values within LF or HF areas showed that a higher proportion of

non-monotonic CT neurons than TR were located in LF areas, while a larger

proportion of non-monotonic TR neurons were located in HF areas than in LF areas

(Figure 4M). In addition, among non-monotonic neurons, TR neurons had higher MI

than CT neurons, whereas monotonic TR neurons had a lower MI than monotonic CT

neurons (Figure 4N). Furthermore, we note that non-monotonic CT neurons had a

higher BF than monotonic CT neurons, whereas the BF was similar between

monotonic and non-monotonic TR neurons (Figure 4O). Finally, BW_{max} did not

significantly differ between monotonic and non-monotonic neurons in either the TR

or CT populations (Figure 4P).

Discussion

A key finding in this study is that the information flow descending from A1 to

thalamus via CT neurons does not preserve tonotopy, which contrasted with the

observation that the information flow ascending from the thalamus to A1 via TR

neurons exhibits clear tonotopy.

Layer specific versus projection specific functional organization in AuC

Previous studies have reported the existence of tonotopic gradients across all layers

of A1 including L6 (Guo et al., 2012; Tischbirek et al., 2019). CT neurons labeled with

CVS-GCaMP6s were restricted to AuC L6 but did not preserve tonotopy. This is

because only 30 - 50% of the pyramidal cells in L6 are CT neurons (Thomson, 2010),

and the neurons we image are restricted to MGB-projecting CT neurons, excluding CT

neurons projecting to other downstream nuclei (Clayton et al., 2021). Using

CVS-based labeling approach and 2P single-cell-resolved imaging, we successfully

dissected a small fraction of CT neurons from all L6 neurons and studied the

functional organization of this group of neurons.

290   The studies examining functional organization in AuC have mostly focused on

layer specificity (Bandyopadhyay et al., 2010; Montes-Lourido et al., 2021; Rothschild

et al., 2010; Tischbirek et al., 2019; Winkowski and Kanold, 2013), and these studies

have provided us with a basic understanding of the tonotopy and heterogeneity of

AuC. However, layer-specific representation is a relatively coarse approach.

Projection specificity-based approach provides detailed insight into differences in

population activity and frequency response-related topographic organization

between neurons with different connectivity specificity. For example, anatomical

studies of AuC in several species have identified the topographical organization of L5

corticocollicular projections (Bajo and Moore, 2005; Saldana et al., 1996; Stebbings

et al., 2014). By contrast, a recent study using a pathway-specific labeling method

showed that L5 corticocollicular neurons displayed a relatively weak topological

organization than the non-corticocollicular neurons in the same layer (Schmitt et al.,

2023). Those studies and our data consistently support the notion that different

anatomically organized projections do not necessarily transmit information with the

same topological alignment. Therefore, combining pathway-specific labeling and

in-vivo single-cell-resolved functional imaging could reveal unexpected fine-scale

details, and settle discrepancies arising from results obtained by individual methods

alone.

CVS virus provides an effective means of mapping neuronal function in L6

The CVSN2c-ΔG rabies virus strain, a recently engineered self-inactivating ΔG rabies

virus lacking the polymerase gene, exhibits strong neurotropism and reduced

cytotoxicity (Reardon et al., 2016). Previous studies have revealed that reliable neural

activity using the CVS-GCaMP6s virus can be maintained for at least 21 days, as

recorded by fiber photometry (Lin et al., 2023). In acute forebrain slices,

CVS-N2c-ΔG-hChR2-YFP reliably elicits action potentials in transfected cortical

neurons for at least 28 days after infection (Reardon et al., 2016). Our recent studies

also confirm that CVS-labeled CT neurons exhibit electrophysiological properties that

are indistinguishable from those of normal neurons (Gu et al., 2023). Here, chronic

2P calcium imaging shows that CT neurons retain stable sound response properties

for at least 13 days (Figure 1- figure supplementary 1). All of these studies suggest

that the CVS virus-labeling method provides an effective means of mapping neuronal

circuitry and manipulating neuronal activity *in vivo* in the mammalian central

nervous system.

Tonotopy of CT neurons can be observed in wide-field but not 2P imaging

Our 2P imaging results are not contradictory with our wide-field epifluorescence imaging data (Figure 1) showing a tonotopy in A1 from the same mice that were later used for 2P imaging experiments. First, the source of the fluorescence signal differs between wide-field imaging and 2P imaging. Wide-field epifluorescence imaging reports all fluorescent proteins from cell bodies and neuropils across layers, whereas 2P imaging only examines fluorescent signals from the cell bodies (Scott et al., 2018). Furthermore, BF tuning from individual neurons results in a significantly closer match to wide-field mapping before removal of the neuropil contribution, thus confirming that the neuropil produces greater local homogeneity in BFs along with a clearer global tonotopic organization (Romero et al., 2020). For L6 CT neurons, the contribution of the neuropil (dendrites) is important due to the specificity of their morphological structure: the cell bodies are located in L6, while their corresponding apical dendrites terminate in L4 (32%) or extend to L1 (68%)(Olsen et al., 2012).

Profiling of auditory features of "in-and-out" neurons in A1

At the single-neuron level, the tonotopic organization or functional distribution of adjacent neuron clusters in the AuC can be highly heterogeneous in mice (Issa et al., 2014; Tao et al., 2017; Winkowski and Kanold, 2013). Our findings in this study reveal that TR neurons exhibit more homogeneous functional and spatial distribution (Figure 3), which contrasts with some other single-cell-resolved studies in the mouse AuC (Li et al., 2017; Rothschild et al., 2010; Winkowski and Kanold, 2013). This discrepancy can be also explained by the pathway specificity, i.e., only a small fraction of neurons in the upper layers of AuC are thalamocortical recipients. It is expected that TR neurons possess high-gradient tonotopy and sharp tuning because feedforward thalamocortical projections are known to possess the same features (Guo et al., 2012; Kanold et al., 2014; Winkowski and Kanold, 2013). What is unexpected is that the same predictions based on the classical literature of functional mapping of AuC (Guo et al., 2012; Imig et al., 1977; Recanzone et al., 1999;

Rothschild et al., 2010) do not apply to L6 CT neurons. This fits perfectly with a

previous study, which found a transition from precise, homogenous frequency

organization in L4 to coarse, diffuse organization in L2/3, suggesting that information

flow exhibits diverse selectivity after cortical processing (King et al., 2018; Winkowski

and Kanold, 2013).

What could be the functional meaning of this change in, or elimination of,

tonotopic organization? We suggest that descending information flow of A1-MGB

projection could potentially support higher cortical functions such as higher-level

feedback processing of complex (Homma et al., 2017; Malmierca et al., 2015) and

behaviorally meaningful sound (Jeschke et al., 2021; Ohl et al., 2001; Wang et al.,

2020; Wang et al., 2022). In addition to frequency preference, this study also

provides evidence illustrating the effects of cortical processing on response

properties. For example, BW_{max}, which reflects the degree of integration of synaptic

inputs from presynaptic neurons (Kratz and Manis, 2015; Li et al., 2019; Schreiner

and Sutter, 1992), is 3.0 octaves in CT neurons, but only 1.3 octaves in TR neurons

(Figure 4I), indicative of an increase in synaptic integration during cortical processing

(Li et al., 2019). The previous study shows that L6 CT neurons are more narrowly

tuned than L5 corticofugal neurons (Williamson and Polley, 2019), which is opposite

to what is reported here because the neurons we image are MGB-projecting CT

neurons. The difference in BW_{max} between TR and CT neurons is consistent with a

scenario in which CT neurons non-selectively "pool" inputs for higher-order feedback

processing of sophisticated auditory inputs that affect behavior (Wang et al., 2020;

Wang et al., 2022).

Conclusion

In summary, our finding fills in a knowledge gap in the auditory physiology: whereas

the corticothalamic feedback projection is known to contribute to higher cognitive

processing of behaviorally-relevant complex sounds, the basic pure-tone map that

underlies and facilitates these advanced processing remained unclear. Our results

reveal that the sensory information flow descending from the AuC to thalamus via L6

CT neurons does not arrange tonotopically. The new shift, categorizing cortical

neurons based on their projection specificity, represents an advance in the

conceptual framework for functional organization.

 Figures

Figure 1. Imaging TR or CT neurons in A1 of awake mice. (**A**) Cartoon illustration of the

auditory thalamocortical and corticothalamic circuits. (**B**) Schematic diagram of the injection

site of AAV2/1-Cre in the MGB and AAV2/9-CaMKⅡ-DIO -GCaMP6s in AuC. (**C**) Coronal slice

showing AAV-GCaMP6s expression in the AuC on day 21. (**D**) Fluorescence intensity of TR

neurons in AuC with distance from pia (0%) to the L6/WM border (100%). (**E**) Schematic

diagram of the injection site of CVS-GCaMP6s in the MGB. (**F**) Coronal slice showing the

CVS-GCaMP6s retrogradely labeled neurons in the AuC on day 7. (**G**) Fluorescence intensity

of CT neurons in AuC with distance from pia (0%) to the L6/WM border (100%). (**H**) Left:

wide-field imaging in AAV-GCaMP6s-expressing mouse (TR neurons), fluorescence response

to pure-tone stimulation with 4 kHz (blue area) and 32 kHz (red area). White dotted lines

outline the A1, A2, and AAF boundaries. Right: wide-field imaging in CVS-GCaMP6s-expressing mouse (CT neurons). (**I**) Left: a fluorescent micrograph of a coronal slice after CTB-555 loading guided by 2P imaging into A1 in a CVS-GCaMP6s-expressing mouse. Right: Micrograph of a coronal slice of the MGB from the same mouse. CTB-555 retrogradely labeled neurons were mainly concentrated in the MGBv. (**J**) Left: wide-field imaging (in a GCaMP6s-expressing mouse) in A1, fluorescence response to pure-tone stimulation with 4 kHz (low-frequency (LF) area) and 32 kHz (high-frequency (HF) area), same abbreviation for all subsequent figures. Inset panels outlined by dashed boxes show the magnified views of 2P imaging of TR or CT neurons. Right: examples of 2P images of TR and CT neurons *in vivo*, respectively.

Figure 1- figure supplementary 1. Chronic *in vivo* 2P imaging of CT neurons. (**A**) Calcium imaging from L6 CT neurons in a CVS-expressing mouse, again with repeated imaging of the same focal planes on day 7 and 13 after virus injection. The dashed circle indicates the example neurons, showing fluorescence traces of neurons in the lower panel. (**B**) The comparison of the fraction of active CT neurons on day 7 and 13. N = 14 focal planes from 5 mice. "Day 7": 79.14\59.26–86.11%, "Day 13": 75.33\62.07–86.11%; P = 0.96, two-sided Wilcoxon signed-rank test, *P < 0.05, **P < 0.01, and ***P < 0.001, same statistics for boxplots. (**C**) The comparison of fraction of reliable CT neurons on day 7 and 13. "Day 7": 12.70\9.09–26.67%, "Day 13": 13.19\7.69–20.00%; P = 0.53. The definitions of "active" and "reliable" neurons see METHOD DETAILS.

Figure 2. Tonotopic gradients of TR versus CT neurons. (**A**) Two-photon image of the focal plane used to image TR neurons in A1. The dashed circle indicates the tuned neurons. Fluorescence traces of neurons in the right panel are indicated by dashed circle ordered according to sound frequency and level. The five traces associated with the five repeats of each stimulus are plotted in gray. The average calcium signals are plotted in black. The red outline marks the BF response of the neuron. (**B**) Same as panel **A**) but for CT neurons in A1. (**C**) BF map of TR neurons. (**D**) BF map of CT neurons. (**E**) Schematic illustrating the recording locations of individual fields of view of TR neurons in A1. Outlines are color-coded according to the median BF response in the respective fields of view, with their BFs (kHz) color-coded

by the scheme below. N = 23 focal planes, from 5 mice. (**F**) Same as panel **E**) but for CT

neurons. N = 40 focal planes, from 10 mice. (**G**) BF$_{median}$ plotted against distance along the

tonotopic axis for TR neurons. Scatter plots showing the correlation between the cellular

BF$_{median}$ values measured in the fields of view with 2P imaging and the corresponding

extrapolated brain surface frequencies determined by wide-field imaging of TR neurons. (**H**)

Same as panel **G**) but for CT neurons. (**I**) Reconstruction of the relative spatial locations of

tuned TR neurons that covered the whole A1 area and color-coded according to each

neuron's BF. N = 481 neurons from 5 mice. (**J**) Plots of TR neurons' BF and their relative

distances along the caudal-to-rostral axis. (**K**) Same as panel **i** but for CT neurons. N = 491

neurons from 10 mice. (**L**) Same as panel **J**) but for CT neurons. (**M**) Distribution histogram of

BF from all neurons in the dataset (TR: N = 481 neurons; CT: N = 491 neurons). (**N**)

Distribution histogram of BF neurons that were identified in the LF area by wide-field

imaging (TR: N = 116 neurons; CT: N = 98 neurons). (**O**) Distribution histogram of BF neurons

that were identified in HF area by wide-field imaging (TR: N = 91 neurons; CT: N = 90

neurons).

Figure 2- figure supplementary 2. Pie charts showing the percentage of responsive neurons

to pure-tone stimulation of TR and CT neurons. N = 5 animals for TR neurons, N = 10 animals

for CT neurons. The definitions of ''silent'' and ''irregular'' neurons see METHOD DETAILS.

Figure 3. Heterogeneity of frequency preference in TR versus CT neurons. (**A**) Spatial

distribution of all tone-selective neurons in an example imaging plane of TR neurons from LF

(left) and HF (right) imaging area. (**B**) Same as panel **A**) but for CT neurons. BFs (kHz) color

code is on panel **A**). (**C**) The comparison of field (200 μm) IQR of TR and CT. N = 23 focal

planes from 5 mice for TR; N = 40 focal planes from 10 mice for CT. "TR": 1.29\0.91–1.88

octaves (median\25–75% percentiles, same notation for all subsequent data), "CT":

1.94\1.29–2.58 octaves. P = 0.016, two-sided Wilcoxon rank-sum test, *P < 0.05, **P < 0.01,

and ***P < 0.001, same statistics for boxplots. (**D**) The comparison of the field IQR of TR and

CT neurons in LF and HF areas. LF area: N = 4 focal planes from 3 mice for TR; N = 9 focal

planes from 5 mice for CT. HF area: N = 5 focal planes from 5 mice for TR; N = 11 focal planes

from 10 mice for CT. "TR: LF area": 0.43\0.22–0.86 octaves, "TR: HF area": 1.29\1.02–1.32

octaves, "CT: LF area": 1.94\1.45–2.42 octaves, "CT: HF area": 1.83\0.94–2.58 octaves. P (TR:

LF area, HF area) = 0.11, P (CT: LF area, HF area) = 0.82, P (LF area: TR, CT) = 0.0084, P (HF

area: TR, CT) = 0.25. (**E**) Cartoon showing how the local and minimum percentages were

computed. (**F**) Cumulative percentage plot displaying local (25 μm radius) IQR around each

neuron of TR and CT. Median local IQR$_{BF}$ of TR neurons = 0.86 octaves, N = 120 local planes

from 5 mice; Median local IQR$_{BF}$ of CT neurons = 1.29 octaves; N = 105 local planes from 10

mice. (**G**) The comparison of the local (25 µm radius) IQR of TR and CT neurons in LF and HF

areas. LF area: N = 34 local planes for TR; N = 20 local planes for CT. HF area: N = 23 local

planes for TR; N = 16 local planes for CT. "TR: LF area": 0.65\0.32–0.97 octaves, "TR: HF area":

1.29\0.54–2.58 octaves, "CT: LF area": 1.29\0.75–2.04 octaves, "CT: HF area": 0.91\0.86–1.61

octaves. P (TR: LF area, HF area) = 0.0021, P (CT: LF area, HF area) = 0.69, P (LF area: TR, CT) =

0.0057, P (HF area: TR, CT) = 0.34. (**H**) Local IQR$_{BF}$ plotted against distance along the

tonotopic axis for TR neurons. (**I**) The comparison of the minimum distance Δ Frequency of

TR and CT neurons in LF and HF area. LF area: N = 116 paired neurons for TR; N = 87 paired

neurons for CT. HF area: N = 91 paired neurons for TR; N = 78 paired neurons for CT. "TR: LF

area": 0.43\0.00–0.86 octaves, "TR: HF area": 0.86\0.00–1.72 octaves, "CT: LF area":

0.43\0.00–1.29 octaves, "CT: HF area": 0.43 \0.00–1.29 octaves. P (TR: LF area, HF area) =

4.25e-04, P (CT: LF area, HF area) = 0.79, P (LF area: TR, CT) = 0.18, P (HF area: TR, CT) = 0.025.

(**J**) Δ Frequency plotted against distance along the tonotopic axis for TR neurons.

Figure 4. Receptive field properties of TR versus CT neurons. (**A**) One representative FRA from a specific type of TR neuron. Types are V shape, I shape, and O shape. See text for further details. (**B**) One representative FRA from a specific type of CT neuron. (**C**) Comparison of fraction of neurons from V, I, and O shaped neurons in TR and CT. (**D**) Comparison of half-peak bandwidth of V-shaped TR and CT neurons. N = 190 neurons from 5 mice for TR; N = 249 neurons from 10 mice for CT. "TR": 1.08\0.86–1.29 octaves, "CT": 1.94\1.29–2.37 octaves. P = 4.86e-29. (**E**) Comparison of Q value (bandwidth of BF/half bandwidth) of V-shaped TR and CT neurons. N = 190 neurons from 5 mice for TR; N = 249 neurons from 10 mice for CT. "TR": 2.00\1.50–2.29, "CT": 1.78\ 1.50–2.00. P = 0.035. (**F**) Comparison of the Q value of V-shaped TR and CT neurons in LF and HF areas. LF area: N = 56 neurons for TR; N = 49 neurons for CT. HF area: N = 37 neurons for TR; N = 41 neurons for CT. "TR: LF area": 1.78\1.60–2.00, "TR: HF area": 2.00\1.58–2.43, "CT: LF area": 1.78\1.50–2.22, "CT: HF area": 1.60\1.33–1.88. P (TR: LF area, HF area) = 0.20, P (CT: LF area, HF area) = 0.032, P (LF area: TR, CT) = 0.88, P (HF area: TR, CT) = 0.0032. (**G**) Comparison of maximum bandwidth (BW$_{max}$)

from V, I, and O shaped neurons in TR and CT. V shape: N = 190 neurons for TR, N = 249

neurons for CT; "TR": 2.15\1.72–2.58 octaves, "CT": 3.44\2.58–3.87 octaves; P = 4.18e-29. I

shape: N = 174 neurons for TR, N = 112 neurons for CT; "TR": 0.86\0.43–1.29 octaves, "CT":

1.29\0.43–3.44 octaves; P = 4.66e-05. O shape: N = 117 neurons for TR, N = 130 neurons for

CT; "TR": 1.29\0.86–2.15 octaves, "CT": 3.01\2.15–3.44 octaves; P = 4.30e-19. (**H**)

Distribution histogram of BW_{max} from all neurons in the dataset. (**I**) Comparison of the BW_{max}

of TR and CT neurons. N = 481 neurons from 5 mice for TR, N = 491 neurons from 10 mice for

CT; "TR": 1.29\0.86–2.15 octaves, "CT": 3.01\2.15–3.87 octaves; P = 1.89e-51. (**J**)

Comparison of the BW_{max} of TR and CT neurons in LF and HF areas. LF area: N = 116 neurons

for TR; N = 98 neurons for CT. HF area: N = 91 neurons for TR; N = 90 neurons for CT. "TR: LF

area": 1.72\1.29–2.15, "TR: HF area": 1.29\ 0.86–2.47, "CT: LF area": 2.58\ 1.72–3.44, "CT:

HF area": 3.01\ 1.29–3.87. P (TR: LF area, HF area) = 0.034, P (CT: LF area, HF area) = 0.92, P

(LF area: TR, CT) = 2.44e-08, P (HF area: TR, CT) = 2.28e-06. (**K**) Distribution of the

monotonicity index (MI) of TR and CT neurons. (**L**) Comparison of the fraction of

non-monotonic neurons of TR and CT. N = 23 focal planes from 5 mice for TR, N = 40 focal

planes from 10 mice for CT; "TR": 20.00%\12.92–31.49%, "CT": 25.00%\14.58–40.00%; P =

0.33. (**M**) Comparison of the fraction of non-monotonic neurons of TR and CT in LF and HF

areas. LF area: N = 4 focal planes from 3 mice for TR; N = 9 focal planes from 5 mice for CT.

HF area: N = 5 focal planes from 5 mice for TR; N = 11 focal planes from 10 mice for CT. "TR:

LF area": 9.55%\6.08–13.50%, "TR: HF area": 15.38%\14.66–29.38%; "CT: LF area":

22.22%\17.59–31.67%, "CT: HF area": 38.18%\16.67–60.00%; P (TR: LF area, HF area) = 0.032,

P (CT: LF area, HF area) = 0.19, P (LF area: TR, CT) = 0.016, P (HF area: TR, CT) = 0.17. (**N**)

Comparison of the MI of non-monotonic and monotonic neurons in TR and CT. Non-mon: N =

107 neurons for TR, N = 128 neurons for CT; "TR": 0.42\0.33–0.47, "CT": 0.35\0.26–0.44; P =

7.03e-04. Mon: N = 374 neurons for TR, N = 363 neurons for CT; "TR": 0.78\0.63–1.00, "CT":

1.00\0.73–1.00; P = 6.19e-11. (**O**) Comparison of the BF between monotonic and

non-monotonic neurons in TR and CT. TR: N = 107 neurons for non-mon, N = 374 neurons for

mon; "non-mon": 6.60\2.93–12.10 kHz, "mon": 8.90\3.60–16.30 kHz; P = 0.69. CT: N = 128

neurons for non-mon, N = 363 neurons for mon; "non-mon": 8.90\3.60–19.15 kHz, "mon":

6.60\2.70–12.10 kHz; P = 1.64e-04. (**P**) Comparison of the BW_{max} between monotonic and

non-monotonic neurons in TR and CT. TR: N = 107 neurons for non-mon, N = 374 neurons for

mon; "non-mon": 1.72\0.86–2.15 octaves, "mon": 1.29\0.86–2.15 octaves; P = 0.92. CT: N =

128 neurons for non-mon, N = 363 neurons for mon; "non-mon": 3.44\2.15–3.87 octaves,

"mon": 3.01\2.15–3.87 octaves; P = 0.052.

536 Materials and methods

537 Key resources table

Reagent type (species) or resource	Designation	Source or reference	Identifiers	Additional information
Genetic reagent (Mus musculus)	Mouse: C57BL/6J	Beijing HFK	RRID:IMSR_JAX:000664	https://www.jax.org/
Recombinant DNA reagent	AAV2/1-hSyn-Cre-WPRE-hGH polyA	BrainVTA	PT-0136	
Recombinant DNA reagent	AAV2/9-CaMK Ⅱ -DIO-GCaMP6s-WPRE-hGH pA	BrainVTA	PT-0071	
Recombinant RNA reagent	CVS-N2c-ΔG-GCaM6s	BrainCase Co., Ltd.	BC-RV-CVS715	
Peptide, recombinant protein	Alexa Fluor 555-conjugated Cholera Toxin Subunit B	Invitrogen	C34776	
Chemical compound	DAPI	Sigma-Aldrich	Cat# D9542	
Software, algorithm	Matlab	Mathworks	N/A	https://www.mathworks.com/
Software, algorithm	LabVIEW	National Instruments	N/A	https://www.ni.com/zh-cn.html
Software, algorithm	Igor Pro	Wavemetrics	N/A	https://www.wavemetrics.com/
Software, algorithm	Prism	GraphPad	N/A	https://ww

			w.graphpad. com/
Other	UHU Super Glue	DETAI	N/A
Other	Sun Medical Super Bond C & B Kit Bonding Kit	Super-Bond	N/A

Animals

C57BL/6J male mice (2–3 months old) were provided by the Laboratory Animal

Center at the Third Military Medical University. The mice were housed in a

temperature- and humidity-controlled room on a cycle of 12-h light/dark (lights off at

19:00) with free access to food and water. All animal procedures were approved by

the Animal Care Committee of the Third Military Medical University and were

performed in accordance with the principles outlined in the National Institutes of

Health Guide for the Care and Use of Laboratory Animals.

Virus injections and confocal imaging

Mice were anesthetized with 1% to 2% isoflurane in pure oxygen and placed in a

stereotactic frame (Beijing Zhongshi Dichuang Technology Development Co., Ltd.). A

warm heating pad was used to keep the animals at a proper body temperature

(36.5°C to 37.5°C). To achieve TR neurons labeling, a Cre-expressing AAV

(AAV2/1-hSyn-Cre-WPRE-hGH polyA, titer ≥ 1E+13vg/ml, PT-0136, BrainVTA) was

injected into the MGB (AP -3.1 mm, ML 2.0 mm, DV -2.8 mm from dura) and

DIO-expressing AAV (AAV2/9-CaMKⅡ-DIO-GCaMp6s-WPRE-hGH pA, PT-0071,

BrainVTA) was injected into the AuC (AP -3.1 mm, ML 3.8 mm, DV -1.4 mm from dura

with manipulator tilted 25° from the z-axis). For CT neuron labeling, the CVS virus

(CVS-N2c-ΔG-GCaM6s), packaged by BrainCase Co., Ltd., Shenzhen, China, was

injected into the MGB. In all the injections, approximately 100 nL of virus was

performed. After injection, the pipette was held in place for 10 min before retraction.

Tissue glue (3M Animal Care Products, Vetbond) was used for bonding the scalp

incision.

Mice were killed for histology 4 weeks after AAV injection or 7 days after CVS

injection. For slice preparation, mice were perfused transcardially with 4%

paraformaldehyde in phosphate-buffered saline. Brains were postfixed in 4%

paraformaldehyde overnight at 4 °C and cut into 50 μm sections on a cryostat

microtome (Thermo Fisher, NX50, Waltham, MA). Mounted sections were imaged on

a scanning confocal microscope (TCS SP5, Leica).

Auditory stimulation

Sound stimuli were delivered by an ED1 electrostatic speaker driver and a free-field

ES1 speaker (both from Tucker Davis Technologies). During experiments, the speaker

was put at a distance of ~6 cm to the left ear of the animal. The sound stimulus was

produced by a custom-written, LabVIEW-based program (LabVIEW 2012, National

Instruments) and transformed to analogue voltage through a PCI6731 card (National

Instruments). All the data were obtained at 1 MHz via a data acquisition device

(USB-6361, National Instruments) and examined by our custom-made LabVIEW

program. For BBN (BBN, bandwidth 0–50 kHz), the sound level was ~65 dB sound

pressure level (SPL). We generated a waveform segment of BBN and used the same

waveform segment for all experiments involving BBN.

For functional mapping of TR and CT neurons, sequences of randomly ordered

pure tones with 11 frequencies (2–40 kHz) at 6 attenuation levels (30–80 dB SPL)

were used. Each frequency-attenuation combination was presented 5–8 times.

Background noise generated by the recording hardware, especially the resonance

scanner of the 2P microscope, was measured to be below < 30 dB SPL for the

relevant frequency range. As described in our previous reports, low frequencies (<1

kHz) were major components of background noise. With a spectral density of ~33

dB/sqrt (Hz), the peak of background noise was below 1 kHz. Neither visible light nor

other sensory stimuli were present. The duration of a sound stimulus (tone or BBN)

was 100 ms.

In vivo **wide-field epifluorescence imaging**

A homemade binocular microscope (BM01, SIBET, CAS) with a 4X, 0.2 NA objective

(Olympus) was used to record wide-field fluorescence images in the mouse cortex for

establishing the reference cortical map of A1. A light-emitting diode (470 nm,

M470L4, Thorlabs) was used for blue illumination. Green fluorescence passed

through a filter cube was measured at 10 Hz with a sCMOS camera (Zyla 4.2, Andor

Technology). Mice injected with AAV-GCaM6s or CVS-GCaM6s were used to

functionally identify the region of A1. The mouse was anesthetized by isoflurane and

kept on a warm plate (37.5°C). A piece of bone (~5 mm × 5 mm) was removed and

replaced by a coverslip (3 mm in diameter). To localize A1, two pure tones (4 and 32

kHz) were repeatedly presented 20 times at an interval of 6 s.

We observed that pure tone stimulation evoked wide-field signals in only one

auditory region in CVS-GCaM6s mice. To confirm the region, we used the criterion

that the ventral part of the medial geniculate body (MGBv) is connecting with A1.

Injections were performed under visual guidance using 2P excitation. In the

experiment, we inserted the electrode into the cortical region at a depth of ~500 μm

below the surface. We used Alexa Fluor 555-conjugated cholera toxin subunit B (CTB)

as the neural tracer and injected the fluorescent CTB solution with 0.5% in phosphate

buffer by pressure (700 mbar) for 15min. Seven days after the fluorescent CTB

injection, the mice were anesthetized with pentobarbital (1.0 g/kg ip). The brain was

first dissected out and then it was immersed with 4% paraformaldehyde overnight.

To visualize fluorescent tracers, a consecutive series of coronal or horizontal sections

(50 μm thick) were prepared using a sliding cryotome, and then all sections were

mounted onto glass slides and imaged with a stereoscope (Olympus).

Two-photon Ca^{2+} imaging in A1

For 2P imaging in head-fixed awake mice, we removed the skin and muscles over the

right A1 after local lidocaine injection under isoflurane anesthesia (1–2%). A

custom-made plastic chamber (head post) designed for head-fixed mouse

experiments was then glued to the skull with cyanoacrylic glue (UHU). After recovery

from surgery for 3 days, the mouse underwent head-fixation training for 3–5 days

(from 1 to 4 h per day). After head-fixation training, mice gradually adapted to this

posture and were able to sit comfortably for 4 h. On the recording day, a small

craniotomy was performed under local anesthesia. Then the field of interest was

determined by comparing the wide-field map with the blood vessel patterns to

ensure A1 was imaged.

Two-photon imaging was performed with a custom-built 2P microscope system

based on a 12.0 kHz resonant scanner (model "LotosScan 1.0", Suzhou Institute of

Biomedical Engineering and Technology). Two-photon excitation light was delivered

by a mode-locked Ti: Sa laser (model "Mai-Tai DeepSee", Spectra Physics), and a

40×/0.8 numerical aperture (NA) water immersion objective (Nikon) was used for

imaging. For Ca^{2+} imaging experiments, the excitation wavelength was set to 920 nm.

The typical size of the field of view (FOV) was ~200 μm × 200 μm. The average power

delivered to the brain was in the range of 30–120 mW, depending on the depth of

imaging.

Chronic 2P Ca^{2+} imaging in L6

For chronic 2P imaging, A circular cranial window (2.5 mm diameter coverslip) was

implanted over the right AuC after head-post implantation. To this end, the muscle

and skull fascia above the temporal skull were removed after local lidocaine (2%)

injection. Afterward, a custom-made plastic chamber was fixed to the skull with

dental cement (Superbond, Sun Medical Co., Ltd.) and a small craniotomy (~2.7 mm

in diameter, the center point: AP: -3.0 mm, ML: -4.5 mm) was performed. The dura

was removed, and the craniotomy was polished to match the size of the coverslip. A

coverslip was carefully placed on top of the cortex with mild compression by

tweezers. The coverslip was sealed with UV-curing dental cement (Tetric N-flow,

Ivoclar Co., Ltd.). Antibiotics (Cefazolin, 500 mg/kg, North China Pharmaceutical

Group Corporation) were administered before surgery, as well as until 3 days after

surgery. 3 days of post-surgery recovery were needed before head-fixation training.

Continuous 2P Ca^{2+} imaging was performed on day 7 and 13. To minimize the

bleaching of cells, we used low laser illumination in the deep imaging, based on

high-quality imaging of CT neurons with CVS labeling strategy. The repeated imaging

FOVs were identified on consecutive days based on superficial blood vessels and

nearby blood vessels, then further refined by visually matching reference images

acquired from precedent days.

QUANTIFICATION AND STATISTICAL ANALYSIS

Data were analyzed using custom-written software in LabVIEW 2012 (National

Instruments), Igor Pro 5.0 (Wavemetrics), Image 1.51 (NIH), Prism 8.4 (GraphPad)

and MATLAB 2014a (MathWorks).

Wide-field imaging data analysis

In each mouse, the recorded cortical images were first down sampled from the

original 750 × 1200 pixels to 75 × 120 pixels. After that, the frames recorded with

sound stimuli were averaged across 20 trials. To enhance the signal-to-noise ratio,

spatial averaging was conducted over 5 × 5 pixels by a matrix filter, and temporal

averaging was conducted with three consecutive images. The pre-processed images

were then temporally normalized to obtain the relative changes in fluorescence (f)

pixel-by-pixel. With the baseline fluorescence (f0) obtained by averaging the images

of 800 ms before sound stimulation, the relative fluorescence changes of each pixel

were calculated as $\Delta f/f = (f − f0)/f0$. The normalized images are shown on a

color-coded scale to visualize the relative fluorescence changes ($\Delta f/f$) in the cortex.

Two-photon imaging data analysis

To correct motion-related artifacts in imaging data, a frame-by-frame alignment

algorithm was used to minimize the sum of squared intensity differences between

each frame image and a template, which was the average of the selected image

frames. To extract fluorescence signals, neurons were visually identified, and drawing

regions of interest (ROIs) based on fluorescence intensity was performed.

Fluorescence changes (f) were calculated by averaging the corresponding pixel values

for each ROI. Relative fluorescence changes Δf/f = (f−f0)/f0 were calculated as Ca^{2+}

signals, where the baseline fluorescence f0 was estimated as the 25th percentile of

the entire fluorescence recording. To calculate the amplitude of sound-evoked Ca^{2+}

transients, we performed automatic Ca^{2+} transient detection based on threshold

criteria regarding peak amplitude and rising rate. The noise level was set to be 3

times the standard deviation of the baseline (window length: 1 s). The peak

amplitude and the rate of rising of the Ca^{2+} signals were calculated to determine

whether it was a true transient. The trace of the detected Ca^{2+} transient was first

extracted by exponential infinite impulse response (IIR) filtering (window length: 200

ms) and then subtracted from the original signal. The residual fluorescence trace was

used as the baseline for the next transient detection, similar to previously published

peeling approaches.

Sound-evoked Ca^{2+} responsiveness

BBN-evoked Ca^{2+} responsiveness

For data from chronic Ca^{2+} imaging, we tracked the same FOV based on the last

training day. We removed the outer ~10% of the image from each ROI to account for

edge effects or imaging deviation. The success rate was defined as the fraction of

sound-evoked responses during 10 consecutive BBN stimuli. Note that "Active"

neurons showed clear Ca^{2+} transients during the entire recording duration, including

spontaneous and BBN-evoked responses. The neurons with a success rate ≥ 50%

were defined as "Reliable" neurons among active neurons.

Pure tone-evoked Ca^{2+} responsiveness

The frequency response area (FRA) of each ROI was constructed from the 55 average

responses to all of the unique frequency-attenuation combinations. The frequency

tuning curves were constructed by plotting the average values (and the s.e.m.) of the

calcium signal amplitudes from single trials for each frequency tested. If more than

one area of contiguous frequency-level combinations remained, the largest one was

defined as the FRA. The amplitude of a Ca^{2+} signal was determined as the average

value for a period of 200 ms around the peak of the calcium transient. The baseline

value was calculated for a period of 100 ms before the onset of the auditory stimulus.

Pure tone frequencies that induced response amplitudes higher than half of the

maximal response were defined as effective frequencies.

"Irregular" neurons exhibited spontaneous activity with highly variable

responses to sound stimulation. "Tuned" neurons were responsive neurons that

demonstrated significant selectivity for certain stimuli. "Silent" neurons were defined

as those that remained completely inactive during our recording period (> 30 min).

For tuned neurons, the best frequency (BF) was defined as the sound frequency

associated with the highest response averaged across all sound levels. FRAs showing

decreasing frequency selectivity with increasing intensity were classified as being V

shape, constant frequency selectivity with increasing intensity was classified as being

I shape, responsiveness to a small range of intensities and frequencies was classified

as being O shape if the peak response was not to the highest intensity. Maximum

bandwidth (BW_{max}) was defined as the maximum FRA width at any level.

To compare the BF heterogeneity, we analyzed the BF distribution in fixed-size

analysis windows. The BF heterogeneity was quantified by the interquartile range

(IQR) of the distribution. For filed IQR, if less than 5 neurons were within a 100 μm

radius (including the center neuron), no IQR was calculated(Schmitt et al., 2023). For

Large-scale analysis, imaging regions in separate mice were aligned with each other

according to the caudal-to-rostral gradient that was identifiable in the wide-field

images.

The monotonicity index (MI) characterizes the strength of a neuron's intensity

tuning. We defined MI as the neuron's response at maximum intensity (80 dB)

divided by its maximum response(Moore and Wehr, 2013; Sutter and Schreiner,

1995). An MI of 1 indicates no intensity tuning; an MI near zero indicates very strong

intensity tuning.

Statistics

To compare data between groups, we used the nonparametric Wilcoxon rank sum test (unpaired), and Wilcoxon signed-rank test (paired) to determine statistical significance (P < 0.05) between them. In the text, summarized data are presented as the median\25th–75th percentiles. In the figures, the data presented in the box-and-whisker plot indicate the median (center line), 25th and 75th percentiles (Q1 and Q3), i.e., IQR (box), Q1-1.5 × IQR and Q3 + 1.5 × IQR (whiskers), and all other data with error bars are presented as the mean ± s.e.m.

Acknowledgments

The authors would like to thank Jia Lou for the cartoon art and figure layout. This work was supported by grants from the National Key R&D Program of China (2021YFA0805000), the National Natural Science Foundation of China (31925018, 32127801, 32300937, T2241002), the Jiangsu Provincial Big Science Facility Initiative (BM2022010), and Guangxi Science and Technology Base & Talents Fund (GUIKE AD22035948). X.C. is a member of the CAS Center for Excellence in Brain Science and Intelligence Technology.

Additional information

Funding

Funder	Grant reference number	Author
National Key R&D Program of China	2021YFA0805000	Xiaowei Chen
National Natural Science Foundation of China	31925018	Xiaowei Chen
National Natural Science Foundation of China	32127801	Xiaowei Chen
National Natural Science Foundation of China	32300937	Jianxiong Zhang
National Natural Science Foundation of China	T2241002	Jianxiong Zhang
Jiangsu Provincial Big Science Facility Initiative	BM2022010	Hongbo Jia

Guangxi Science and Technology Base & GUIKE AD22035948   Xiaowei Chen
Talents Fund

Author contributions

X.C. and H.J. conceived the project. X.C., JX.Z. and Y.Z. designed the experiments. M.G. performed the experiments with the help of R.L., K.L., X.W. and JH.Z.; X.C., H.J., Y.Z., S.L., JX.Z., and M.G. devised the data analysis methods; M.G., S.L., Yun.Z., X.L., C.Z. and JX.Z. performed the data analysis; JX.Z., H.J., Y.Z. and X.C. inspected the data and evaluated the findings; M.G., JX.Z., Y.Z., F.O., X.C. and H.J. wrote the manuscript with the help from all authors.

Author ORCIDs

Xiaowei Chen: https://orcid.org/0000-0003-0906-6666

Jianxiong Zhang: https://orcid.org/0000-0003-2245-8752

Yi Zhou: https://orcid.org/0000-0002-2623-8960

Ethics

All experiments were approved by the Animal Care Committee of the Third Military Medical University (Approval number: AMUWEC20230061).

Additional files

Data availability

Wide-field imaging and two-photon imaging are available upon request from the corresponding author. Source data underlying Figs. 2–4 is available as a Source data file. No datasets that require mandatory deposition into a public database were generated during the current study. Source data are provided in this paper.

Code availability

The codes supporting the current study have not been deposited in a public

787   repository, but are available from the corresponding author upon request.

References

Andersen RA, Knight PL, Merzenich MM. 1980. The thalamocortical and corticothalamic connections
of AI, AII, and the anterior auditory field (AAF) in the cat: evidence for two largely segregated systems
of connections. *Journal of Comparative Neurology* 194: 663-701. DIO:
https://doi.org/10.1002/cne.901940312, PMID: 7451688

Antunes FM, Malmierca MS. 2021. Corticothalamic Pathways in Auditory Processing: Recent Advances
and Insights From Other Sensory Systems. *Front Neural Circuits* 15: 721186. DIO:
https://doi.org/10.3389/fncir.2021.721186, PMID: 34489648

Bajo VM, Moore DR. 2005. Descending projections from the auditory cortex to the inferior colliculus in
the gerbil, Meriones unguiculatus. *Journal of Comparative Neurology* 486: 101-16. DIO:
https://doi.org/10.1002/cne.20542, PMID: 15844210

Bandyopadhyay S, Shamma SA, Kanold PO. 2010. Dichotomy of functional organization in the mouse
auditory cortex. *Nat Neurosci* 13: 361-8. DIO: https://doi.org/10.1038/nn.2490, PMID: 20118924

Bizley JK, Nodal FR, Nelken I, King AJ. 2005. Functional organization of ferret auditory cortex. *Cerebral
Cortex* 15: 1637-53. DIO: https://doi.org/10.1093/cercor/bhi042, PMID: 15703254

Budinger E, Heil P, Scheich H. 2000. Functional organization of auditory cortex in the Mongolian gerbil
(Meriones unguiculatus). IV. Connections with anatomically characterized subcortical structures.
European Journal of Neuroscience 12: 2452-74. DIO:
https://doi.org/10.1046/j.1460-9568.2000.00143.x, PMID: 10947822

Clayton KK, Williamson RS, Hancock KE, Tasaka GI, Mizrahi A, Hackett TA, Polley DB. 2021. Auditory
Corticothalamic Neurons Are Recruited by Motor Preparatory Inputs. *Current Biology* 31: 310-321 e5.
DIO: https://doi.org/10.1016/j.cub.2020.10.027, PMID: 33157020

Clopton BM, Winfield JA, Flammino FJ. 1974. Tonotopic organization: review and analysis. *Brain
Research* 76: 1-20. DIO: https://doi.org/10.1016/0006-8993(74)90509-5, PMID: 4367399

Constantinople CM, Bruno RM. 2013. Deep cortical layers are activated directly by thalamus. *Science*
340: 1591-4. DIO: https://doi.org/10.1126/science.1236425, PMID: 23812718

de la Rocha J, Marchetti C, Schiff M, Reyes AD. 2008. Linking the response properties of cells in
auditory cortex with network architecture: cotuning versus lateral inhibition. *Journal of Neuroscience*
28: 9151-63. DIO: https://doi.org/10.1523/JNEUROSCI.1789-08.2008, PMID: 18784296

Gu M, Li X, Liang S, Zhu J, Sun P, He Y, Yu H, Li R, Zhou Z, Lyu J, Li SC, Budinger E, Zhou Y, Jia H, Zhang J,
Chen X. 2023. Rabies virus-based labeling of layer 6 corticothalamic neurons for two-photon imaging
in vivo. *iScience* 26: 106625. DIO: https://doi.org/10.1016/j.isci.2023.106625, PMID: 37250327

Guo W, Chambers AR, Darrow KN, Hancock KE, Shinn-Cunningham BG, Polley DB. 2012. Robustness of
cortical topography across fields, laminae, anesthetic states, and neurophysiological signal types.
Journal of Neuroscience 32: 9159-72. DIO: https://doi.org/10.1523/JNEUROSCI.0065-12.2012, PMID:
22764225

Hackett TA, Stepniewska I, Kaas JH. 1998. Subdivisions of auditory cortex and ipsilateral cortical
connections of the parabelt auditory cortex in macaque monkeys. *Journal of Comparative Neurology*
394: 475-95. DIO: https://doi.org/10.1002/(sici)1096-9861(19980518)394:4<475::aid-cne6>3.0.co;2-z,
PMID: 9590556

Happel MF, Deliano M, Handschuh J, Ohl FW. 2014. Dopamine-modulated recurrent corticoefferent

feedback in primary sensory cortex promotes detection of behaviorally relevant stimuli. *Journal of Neuroscience* 34: 1234-47. DIO: https://doi.org/10.1523/JNEUROSCI.1990-13.2014, PMID: 24453315

Happel MF, Jeschke M, Ohl FW. 2010. Spectral integration in primary auditory cortex attributable to temporally precise convergence of thalamocortical and intracortical input. *Journal of Neuroscience* 30: 11114-27. DIO: https://doi.org/10.1523/JNEUROSCI.0689-10.2010, PMID: 20720119

Harris JA, Mihalas S, Hirokawa KE, Whitesell JD, Choi H, Bernard A, Bohn P, Caldejon S, Casal L, Cho A, Feiner A, Feng D, Gaudreault N, Gerfen CR, Graddis N, Groblewski PA, Henry AM, Ho A, Howard R, Knox JE, Kuan L, Kuang X, Lecoq J, Lesnar P, Li Y, Luviano J, McConoughey S, Mortrud MT, Naeemi M, Ng L, Oh SW, Ouellette B, Shen E, Sorensen SA, Wakeman W, Wang Q, Wang Y, Williford A, Phillips JW, Jones AR, Koch C, Zeng H. 2019. Hierarchical organization of cortical and thalamic connectivity. *Nature* 575: 195-202. DIO: https://doi.org/10.1038/s41586-019-1716-z, PMID: 31666704

Harris KD, Mrsic-Flogel TD. 2013. Cortical connectivity and sensory coding. *Nature* 503: 51-8. DIO: https://doi.org/10.1038/nature12654, PMID: 24201278

Homma NY, Bajo VM. 2021. Lemniscal Corticothalamic Feedback in Auditory Scene Analysis. *Frontiers in Neuroscience* 15: 723893. DIO: https://doi.org/10.3389/fnins.2021.723893, PMID: 34489635

Homma NY, Happel MFK, Nodal FR, Ohl FW, King AJ, Bajo VM. 2017. A Role for Auditory Corticothalamic Feedback in the Perception of Complex Sounds. *Journal of Neuroscience* 37: 6149-6161. DIO: https://doi.org/10.1523/JNEUROSCI.0397-17.2017, PMID: 28559384

Humphries C, Liebenthal E, Binder JR. 2010. Tonotopic organization of human auditory cortex. *Neuroimage* 50: 1202-11. DIO: https://doi.org/10.1016/j.neuroimage.2010.01.046, PMID: 20096790

Imig TJ, Ruggero MA, Kitzes LM, Javel E, Brugge JF. 1977. Organization of auditory cortex in the owl monkey (Aotus trivirgatus). *Journal of Comparative Neurology* 171: 111-28. DIO: https://doi.org/10.1002/cne.901710108, PMID: 401509

Issa JB, Haeffele BD, Agarwal A, Bergles DE, Young ED, Yue DT. 2014. Multiscale optical Ca2+ imaging of tonal organization in mouse auditory cortex. *Neuron* 83: 944-59. DIO: https://doi.org/10.1016/j.neuron.2014.07.009, PMID: 25088366

Jasmin K, Lima CF, Scott SK. 2019. Understanding rostral-caudal auditory cortex contributions to auditory perception. *Nature Reviews Neuroscience* 20: 425-434. DIO: https://doi.org/10.1038/s41583-019-0160-2, PMID: 30918365

Jeschke M, Ohl FW, Wang X. 2021. Effects of Cortical Cooling on Sound Processing in Auditory Cortex and Thalamus of Awake Marmosets. *Front Neural Circuits* 15: 786740. DIO: https://doi.org/10.3389/fncir.2021.786740, PMID: 35069125

Jones EG. 2001. The thalamic matrix and thalamocortical synchrony. *Trends in Neurosciences* 24: 595-601. DIO: https://doi.org/10.1016/s0166-2236(00)01922-6, PMID: 11576674

Kajikawa Y, de La Mothe L, Blumell S, Hackett TA. 2005. A comparison of neuron response properties in areas A1 and CM of the marmoset monkey auditory cortex: tones and broadband noise. *Journal of Neurophysiology* 93: 22-34. DIO: https://doi.org/10.1152/jn.00248.2004, PMID: 15342713

Kalatsky VA, Polley DB, Merzenich MM, Schreiner CE, Stryker MP. 2005. Fine functional organization of auditory cortex revealed by Fourier optical imaging. *Proceedings of the National Academy of Sciences of the United States of America* 102: 13325-30. DIO: https://doi.org/10.1073/pnas.0505592102, PMID: 16141342

Kanold PO, Nelken I, Polley DB. 2014. Local versus global scales of organization in auditory cortex. *Trends in Neurosciences* 37: 502-10. DIO: https://doi.org/10.1016/j.tins.2014.06.003, PMID: 25002236

King AJ, Teki S, Willmore BDB. 2018. Recent advances in understanding the auditory cortex. *F1000Res*

7 DIO: https://doi.org/10.12688/f1000research.15580.1, PMID: 30345008

Kirchgessner MA, Franklin AD, Callaway EM. 2020. Context-dependent and dynamic functional
influence of corticothalamic pathways to first- and higher-order visual thalamus. *Proceedings of the*
National Academy of Sciences of the United States of America 117: 13066-13077. DIO:
https://doi.org/10.1073/pnas.2002080117, PMID: 32461374

Kobat D, Durst ME, Nishimura N, Wong AW, Schaffer CB, Xu C. 2009. Deep tissue multiphoton
microscopy using longer wavelength excitation. *Optics Express* 17: 13354-13364. DIO:
https://doi.org/10.1364/Oe.17.013354, PMID: WOS:000268843700009

Kratz MB, Manis PB. 2015. Spatial organization of excitatory synaptic inputs to layer 4 neurons in
mouse primary auditory cortex. *Front Neural Circuits* 9: 17. DIO:
https://doi.org/10.3389/fncir.2015.00017, PMID: 25972787

Lee CC. 2013. Thalamic and cortical pathways supporting auditory processing. *Brain and Language*
126: 22-8. DIO: https://doi.org/10.1016/j.bandl.2012.05.004, PMID: 22728130

Lee CC, Schreiner CE, Imaizumi K, Winer JA. 2004. Tonotopic and heterotopic projection systems in
physiologically defined auditory cortex. *Neuroscience* 128: 871-87. DIO:
https://doi.org/10.1016/j.neuroscience.2004.06.062, PMID: 15464293

Lee CC, Winer JA. 2005. Principles governing auditory cortex connections. *Cerebral Cortex* 15: 1804-14.
DIO: https://doi.org/10.1093/cercor/bhi057, PMID: 15800026

Li H, Liang F, Zhong W, Yan L, Mesik L, Xiao Z, Tao HW, Zhang LI. 2019. Synaptic Mechanisms for
Bandwidth Tuning in Awake Mouse Primary Auditory Cortex. *Cerebral Cortex* 29: 2998-3009. DIO:
https://doi.org/10.1093/cercor/bhy165, PMID: 30010857

Li J, Zhang J, Wang M, Pan J, Chen X, Liao X. 2017. Functional imaging of neuronal activity of auditory
cortex by using Cal-520 in anesthetized and awake mice. *Biomed Opt Express* 8: 2599-2610. DIO:
https://doi.org/10.1364/boe.8.002599, PMID: 28663893

Lin KZ, Li L, Ma WY, Yang X, Han ZP, Luo NS, Wang J, Xu FQ. 2023. A rabies virus-based toolkit for
efficient retrograde labeling and monosynaptic tracing. *Neural Regen Res* 18: 1827-1833. DIO:
https://doi.org/10.4103/1673-5374.358618, PMID: 36751812

Liu J, Whiteway MR, Sheikhattar A, Butts DA, Babadi B, Kanold PO. 2019. Parallel Processing of Sound
Dynamics across Mouse Auditory Cortex via Spatially Patterned Thalamic Inputs and Distinct Areal
Intracortical Circuits. *Cell Rep* 27: 872-885 e7. DIO: https://doi.org/10.1016/j.celrep.2019.03.069,
PMID: 30995483

Malmierca MS, Anderson LA, Antunes FM. 2015. The cortical modulation of stimulus-specific
adaptation in the auditory midbrain and thalamus: a potential neuronal correlate for predictive coding.
Front Syst Neurosci 9: 19. DIO: https://doi.org/10.3389/fnsys.2015.00019, PMID: 25805974

Merzenich MM, Brugge JF. 1973. Representation of the cochlear partition of the superior temporal
plane of the macaque monkey. *Brain Research* 50: 275-96. DIO:
https://doi.org/10.1016/0006-8993(73)90731-2, PMID: 4196192

Merzenich MM, Kaas JH, Roth GL. 1976. Auditory cortex in the grey squirrel: tonotopic organization
and architectonic fields. *Journal of Comparative Neurology* 166: 387-401. DIO:
https://doi.org/10.1002/cne.901660402, PMID: 1270613

Micheyl C, Schrater PR, Oxenham AJ. 2013. Auditory frequency and intensity discrimination explained
using a cortical population rate code. *PLoS Computational Biology* 9: e1003336. DIO:
https://doi.org/10.1371/journal.pcbi.1003336, PMID: 24244142

Montes-Lourido P, Kar M, David SV, Sadagopan S. 2021. Neuronal selectivity to complex vocalization

features emerges in the superficial layers of primary auditory cortex. *PLoS Biology* 19: e3001299. DIO:
https://doi.org/10.1371/journal.pbio.3001299, PMID: 34133413
Moore AK, Wehr M. 2013. Parvalbumin-expressing inhibitory interneurons in auditory cortex are
well-tuned for frequency. *Journal of Neuroscience* 33: 13713-23. DIO:
https://doi.org/10.1523/JNEUROSCI.0663-13.2013, PMID: 23966693
Morel A, Garraghty PE, Kaas JH. 1993. Tonotopic organization, architectonic fields, and connections of
auditory cortex in macaque monkeys. *Journal of Comparative Neurology* 335: 437-59. DIO:
https://doi.org/10.1002/cne.903350312, PMID: 7693772
Nelken I, Bizley JK, Nodal FR, Ahmed B, Schnupp JW, King AJ. 2004. Large-scale organization of ferret
auditory cortex revealed using continuous acquisition of intrinsic optical signals. *Journal of
Neurophysiology* 92: 2574-88. DIO: https://doi.org/10.1152/jn.00276.2004, PMID: 15152018
Norman-Haignere S, Kanwisher NG, McDermott JH. 2015. Distinct Cortical Pathways for Music and
Speech Revealed by Hypothesis-Free Voxel Decomposition. *Neuron* 88: 1281-1296. DIO:
https://doi.org/10.1016/j.neuron.2015.11.035, PMID: 26687225
Oheim M, Beaurepaire E, Chaigneau E, Mertz J, Charpak S. 2001. Two-photon microscopy in brain
tissue: parameters influencing the imaging depth. *Journal of Neuroscience Methods* 111: 29-37. DIO:
https://doi.org/10.1016/s0165-0270(01)00438-1, PMID: 11574117
Ohga S, Tsukano H, Horie M, Terashima H, Nishio N, Kubota Y, Takahashi K, Hishida R, Takebayashi H,
Shibuki K. 2017. DIO: https://doi.org/10.1101/226100,
Ohl FW, Scheich H, Freeman WJ. 2000. Topographic analysis of epidural pure-tone-evoked potentials in
gerbil auditory cortex. *Journal of Neurophysiology* 83: 3123-32. DIO:
https://doi.org/10.1152/jn.2000.83.5.3123, PMID: 10805706
Ohl FW, Scheich H, Freeman WJ. 2001. Change in pattern of ongoing cortical activity with auditory
category learning. *Nature* 412: 733-6. DIO: https://doi.org/10.1038/35089076, PMID: 11507640
Olsen SR, Bortone DS, Adesnik H, Scanziani M. 2012. Gain control by layer six in cortical circuits of
vision. *Nature* 483: 47-52. DIO: https://doi.org/10.1038/nature10835, PMID: 22367547
Pardi MB, Vogenstahl J, Dalmay T, Spano T, Pu DL, Naumann LB, Kretschmer F, Sprekeler H, Letzkus JJ.
2020. A thalamocortical top-down circuit for associative memory. *Science* 370: 844-848. DIO:
https://doi.org/10.1126/science.abc2399, PMID: 33184213
Petreanu L, Mao T, Sternson SM, Svoboda K. 2009. The subcellular organization of neocortical
excitatory connections. *Nature* 457: 1142-5. DIO: https://doi.org/10.1038/nature07709, PMID:
19151697
Polley DB, Read HL, Storace DA, Merzenich MM. 2007. Multiparametric auditory receptive field
organization across five cortical fields in the albino rat. *Journal of Neurophysiology* 97: 3621-38. DIO:
https://doi.org/10.1152/jn.01298.2006, PMID: 17376842
Reale RA, Imig TJ. 1980. Tonotopic organization in auditory cortex of the cat. *Journal of Comparative
Neurology* 192: 265-91. DIO: https://doi.org/10.1002/cne.901920207, PMID: 7400399
Reardon TR, Murray AJ, Turi GF, Wirblich C, Croce KR, Schnell MJ, Jessell TM, Losonczy A. 2016. Rabies
Virus CVS-N2c(DeltaG) Strain Enhances Retrograde Synaptic Transfer and Neuronal Viability. *Neuron*
89: 711-24. DIO: https://doi.org/10.1016/j.neuron.2016.01.004, PMID: 26804990
Recanzone GH, Schreiner CE, Sutter ML, Beitel RE, Merzenich MM. 1999. Functional organization of
spectral receptive fields in the primary auditory cortex of the owl monkey. *Journal of Comparative
Neurology* 415: 460-81. DIO:
https://doi.org/10.1002/(sici)1096-9861(19991227)415:4<460::aid-cne4>3.0.co;2-f, PMID: 10570456

Rodrigues-Dagaeff C, Simm G, De Ribaupierre Y, Villa A, De Ribaupierre F, Rouiller EM. 1989. Functional organization of the ventral division of the medial geniculate body of the cat: evidence for a rostro-caudal gradient of response properties and cortical projections. *Hearing Research* 39: 103-25. DIO: https://doi.org/10.1016/0378-5955(89)90085-3, PMID: 2737959

Romero S, Hight AE, Clayton KK, Resnik J, Williamson RS, Hancock KE, Polley DB. 2020. Cellular and Widefield Imaging of Sound Frequency Organization in Primary and Higher Order Fields of the Mouse Auditory Cortex. *Cerebral Cortex* 30: 1603-1622. DIO: https://doi.org/10.1093/cercor/bhz190, PMID: 31667491

Rothschild G, Nelken I, Mizrahi A. 2010. Functional organization and population dynamics in the mouse primary auditory cortex. *Nat Neurosci* 13: 353-60. DIO: https://doi.org/10.1038/nn.2484, PMID: 20118927

Sadagopan S, Wang X. 2008. Level invariant representation of sounds by populations of neurons in primary auditory cortex. *Journal of Neuroscience* 28: 3415-26. DIO: https://doi.org/10.1523/JNEUROSCI.2743-07.2008, PMID: 18367608

Saldana E, Feliciano M, Mugnaini E. 1996. Distribution of descending projections from primary auditory neocortex to inferior colliculus mimics the topography of intracollicular projections. *Journal of Comparative Neurology* 371: 15-40. DIO: https://doi.org/10.1002/(SICI)1096-9861(19960715)371:1<15::AID-CNE2>3.0.CO;2-O, PMID: 8835717

Schmitt TTX, Andrea KMA, Wadle SL, Hirtz JJ. 2023. Distinct topographic organization and network activity patterns of corticocollicular neurons within layer 5 auditory cortex. *Front Neural Circuits* 17: 1210057. DIO: https://doi.org/10.3389/fncir.2023.1210057, PMID: 37521334

Schneider DM, Woolley SM. 2013. Sparse and background-invariant coding of vocalizations in auditory scenes. *Neuron* 79: 141-52. DIO: https://doi.org/10.1016/j.neuron.2013.04.038, PMID: 23849201

Schreiner CE, Sutter ML. 1992. Topography of excitatory bandwidth in cat primary auditory cortex: single-neuron versus multiple-neuron recordings. *Journal of Neurophysiology* 68: 1487-502. DIO: https://doi.org/10.1152/jn.1992.68.5.1487, PMID: 1479426

Scott BB, Thiberge SY, Guo C, Tervo DGR, Brody CD, Karpova AY, Tank DW. 2018. Imaging Cortical Dynamics in GCaMP Transgenic Rats with a Head-Mounted Widefield Macroscope. *Neuron* 100: 1045-1058 e5. DIO: https://doi.org/10.1016/j.neuron.2018.09.050, PMID: 30482694

Shepherd GMG, Yamawaki N. 2021. Untangling the cortico-thalamo-cortical loop: cellular pieces of a knotty circuit puzzle. *Nature Reviews Neuroscience* 22: 389-406. DIO: https://doi.org/10.1038/s41583-021-00459-3, PMID: 33958775

Sherman SM, Guillery RW. 1998. On the actions that one nerve cell can have on another: distinguishing "drivers" from "modulators". *Proceedings of the National Academy of Sciences of the United States of America* 95: 7121-6. DIO: https://doi.org/10.1073/pnas.95.12.7121, PMID: 9618549

Simon E, Perrot X, Mertens P. 2009. [Functional anatomy of the cochlear nerve and the central auditory system]. *Neuro-Chirurgie* 55: 120-6. DIO: https://doi.org/10.1016/j.neuchi.2009.01.017, PMID: 19304300

Smith KR, Wever EG. 1949. The problem of stimulation deafness; the functional and histological effects of a high-frequency stimulus. *Journal of Experimental Psychology* 39: 238-41. DIO: https://doi.org/10.1037/h0061044, PMID: 18125722

Stebbings KA, Lesicko AM, Llano DA. 2014. The auditory corticocollicular system: molecular and circuit-level considerations. *Hearing Research* 314: 51-9. DIO: https://doi.org/10.1016/j.heares.2014.05.004, PMID: 24911237

Stiebler I, Neulist R, Fichtel I, Ehret G. 1997. The auditory cortex of the house mouse: left-right
differences, tonotopic organization and quantitative analysis of frequency representation. *J Comp*
Physiol A 181: 559-71. DIO: https://doi.org/10.1007/s003590050140, PMID: 9449817

Sun L, Tang Y, Yan K, Yu J, Zou Y, Xu W, Xiao K, Zhang Z, Li W, Wu B, Hu Z, Chen K, Fu ZF, Dai J, Cao G.
2019. Differences in neurotropism and neurotoxicity among retrograde viral tracers. *Molecular*
Neurodegeneration 14: 8. DIO: https://doi.org/10.1186/s13024-019-0308-6, PMID: 30736827

Sutter ML, Schreiner CE. 1995. Topography of intensity tuning in cat primary auditory cortex:
single-neuron versus multiple-neuron recordings. *Journal of Neurophysiology* 73: 190-204. DIO:
https://doi.org/10.1152/jn.1995.73.1.190, PMID: 7714564

Takasaki K, Abbasi-Asl R, Waters J. 2020. Superficial Bound of the Depth Limit of Two-Photon Imaging
in Mouse Brain. *eNeuro* 7 DIO: https://doi.org/10.1523/ENEURO.0255-19.2019, PMID: 31907211

Tani T, Abe H, Hayami T, Banno T, Miyakawa N, Kitamura N, Mashiko H, Ichinohe N, Suzuki W. 2018.
Sound Frequency Representation in the Auditory Cortex of the Common Marmoset Visualized Using
Optical Intrinsic Signal Imaging. *eNeuro* 5 DIO: https://doi.org/10.1523/ENEURO.0078-18.2018, PMID:
29736410

Tao C, Zhang G, Zhou C, Wang L, Yan S, Tao HW, Zhang LI, Zhou Y, Xiong Y. 2017. Diversity in
Excitation-Inhibition Mismatch Underlies Local Functional Heterogeneity in the Rat Auditory Cortex.
Cell Rep 19: 521-531. DIO: https://doi.org/10.1016/j.celrep.2017.03.061, PMID: 28423316

Thomas H, Tillein J, Heil P, Scheich H. 1993. Functional organization of auditory cortex in the
mongolian gerbil (Meriones unguiculatus). I. Electrophysiological mapping of frequency
representation and distinction of fields. *European Journal of Neuroscience* 5: 882-97. DIO:
https://doi.org/10.1111/j.1460-9568.1993.tb00940.x, PMID: 8281300

Thomson AM. 2010. Neocortical layer 6, a review. *Front Neuroanat* 4: 13. DIO:
https://doi.org/10.3389/fnana.2010.00013, PMID: 20556241

Tischbirek C, Birkner A, Jia H, Sakmann B, Konnerth A. 2015. Deep two-photon brain imaging with a
red-shifted fluorometric Ca2+ indicator. *Proceedings of the National Academy of Sciences of the United*
States of America 112: 11377-82. DIO: https://doi.org/10.1073/pnas.1514209112, PMID: 26305966

Tischbirek CH, Noda T, Tohmi M, Birkner A, Nelken I, Konnerth A. 2019. In Vivo Functional Mapping of
a Cortical Column at Single-Neuron Resolution. *Cell Rep* 27: 1319-1326 e5. DIO:
https://doi.org/10.1016/j.celrep.2019.04.007, PMID: 31042460

Triarhou LC. 2021. Pre-Brodmann pioneers of cortical cytoarchitectonics I: Theodor Meynert, Vladimir
Betz and William Bevan-Lewis. *Brain Structure and Function* 226: 49-67. DIO:
https://doi.org/10.1007/s00429-020-02168-6, PMID: 33165657

Wang M, Liao X, Li R, Liang S, Ding R, Li J, Zhang J, He W, Liu K, Pan J, Zhao Z, Li T, Zhang K, Li X, Lyu J,
Zhou Z, Varga Z, Mi Y, Zhou Y, Yan J, Zeng S, Liu JK, Konnerth A, Nelken I, Jia H, Chen X. 2020.
Single-neuron representation of learned complex sounds in the auditory cortex. *Nature*
Communications 11: 4361. DIO: https://doi.org/10.1038/s41467-020-18142-z, PMID: 32868773

Wang M, Liu K, Pan J, Li J, Sun P, Zhang Y, Li L, Guo W, Xin Q, Zhao Z, Liu Y, Zhou Z, Lyu J, Zheng T, Han Y,
Zhang C, Liao X, Zeng S, Jia H, Chen X. 2022. Brain-wide projection reconstruction of single functionally
defined neurons. *Nature Communications* 13: 1531. DIO:
https://doi.org/10.1038/s41467-022-29229-0, PMID: 35318336

Watkins PV, Barbour DL. 2011. Rate-level responses in awake marmoset auditory cortex. *Hearing*
Research 275: 30-42. DIO: https://doi.org/10.1016/j.heares.2010.11.011, PMID: 21145961

Williamson RS, Polley DB. 2019. Parallel pathways for sound processing and functional connectivity

among layer 5 and 6 auditory corticofugal neurons. *Elife* 8 DIO: https://doi.org/10.7554/eLife.42974, PMID: 30735128

Winer JA, Diehl JJ, Larue DT. 2001. Projections of auditory cortex to the medial geniculate body of the cat. *Journal of Comparative Neurology* 430: 27-55. DIO, PMID: 11135244

Winer JA, Miller LM, Lee CC, Schreiner CE. 2005. Auditory thalamocortical transformation: structure and function. *Trends in Neurosciences* 28: 255-63. DIO: https://doi.org/10.1016/j.tins.2005.03.009, PMID: 15866200

Winkowski DE, Kanold PO. 2013. Laminar transformation of frequency organization in auditory cortex. *J Neurosci* 33: 1498-508. DIO: https://doi.org/10.1523/jneurosci.3101-12.2013, PMID: 23345224

Wu GK, Li P, Tao HW, Zhang LI. 2006. Nonmonotonic synaptic excitation and imbalanced inhibition underlying cortical intensity tuning. *Neuron* 52: 705-15. DIO: https://doi.org/10.1016/j.neuron.2006.10.009, PMID: 17114053

Yarmohammadi H, Mahmoudi Nezhad GS, Dalfardi B, Ghanizadeh A. 2014. Theodor Meynert (1833-1892). *Journal of Neurology* 261: 1647-8. DIO: https://doi.org/10.1007/s00415-013-7077-x, PMID: 23963472

Zeng HH, Huang JF, Chen M, Wen YQ, Shen ZM, Poo MM. 2019. Local homogeneity of tonotopic organization in the primary auditory cortex of marmosets. *Proceedings of the National Academy of Sciences of the United States of America* 116: 3239-3244. DIO: https://doi.org/10.1073/pnas.1816653116, PMID: 30718428

Zhu X, Lin K, Liu Q, Yue X, Mi H, Huang X, He X, Wu R, Zheng D, Wei D, Jia L, Wang W, Manyande A, Wang J, Zhang Z, Xu F. 2020. Rabies Virus Pseudotyped with CVS-N2C Glycoprotein as a Powerful Tool for Retrograde Neuronal Network Tracing. *Neurosci Bull* 36: 202-216. DIO: https://doi.org/10.1007/s12264-019-00423-3, PMID: 31444652

Zingg B, Chou XL, Zhang ZG, Mesik L, Liang F, Tao HW, Zhang LI. 2017. AAV-Mediated Anterograde Transsynaptic Tagging: Mapping Corticocollicular Input-Defined Neural Pathways for Defense Behaviors. *Neuron* 93: 33-47. DIO: https://doi.org/10.1016/j.neuron.2016.11.045, PMID: 27989459