Impaired hippocampal circuit function underlying memory encoding and consolidation

precede robust Aβ deposition in a mouse model of Alzheimer's disease

Hanyan Li [1,2,#], Zhuoyang Zhao [1,2,#], Aline Fassini [1,2], Han K. Lee [1,2], Reese J. Green [1,2], and

Stephen N. Gomperts [1,2,*]

[1] Department of Neurology, Massachusetts General Hospital, Boston, Massachusetts, USA.

[2] Harvard Medical School, Boston, Massachusetts, USA.

# Contributed equally to this work

* Correspondence: Gomperts.stephen@mgh.harvard.edu

Keywords: Alzheimer's disease, amyloid-beta, hippocampus, place cell, electrophysiology, sleep

Abstract

Current therapeutic strategies for Alzheimer's disease (AD) target amyloid-beta (Aβ) fibrils and

high molecular weight protofibrils associated with plaques, but molecular cascades associated

with AD may drive neural systems failure before Aβ plaque deposition in AD. Employing

hippocampal electrophysiological recordings and dynamic calcium imaging across the sleep-

wake cycle in a freely behaving mouse model of AD before Aβ plaques accumulated, we

detected marked impairments of hippocampal systems function: In a spatial behavioral task,

phase-amplitude coupling (PAC) of the hippocampal theta and gamma oscillations was impaired

and place cell calcium fluctuations were hyper-synchronized with the theta oscillation. These

changes were not observed in REM sleep. In subsequent slow wave sleep (SWS), place cell

reactivation was reduced. These degraded neural functions underlying memory encoding and

consolidation support targeting pathological processes of the pre-plaque phase of AD to treat and

prevent hippocampal impairments.

Introduction

In Alzheimer's disease (AD), deposition of amyloid-beta42 (Aβ) in amyloid plaques and aggregation of intracellular tau in neurofibrillary tangles are associated with neuronal loss and progressive memory impairment on the basis of hippocampal systems failure [1]. The advent of amyloid-targeted immunotherapies such as Lecanemab (targeting high molecular weight Aβ protofibrils as well as fibrils in plaques) [2], Donanemab (targeting the pyroglutamate modification seen in Aβ fibrils in plaques) [3], and Aducanumab (targeting Aβ fibrils in plaques) [4] for the treatment of AD heralds the beginning of a new era, with evidence to date demonstrating marked reductions of brain amyloid. Even so, these treatments as yet have only modestly improved cognitive outcomes of AD patients [2,3,5]. Several plausible explanations may explain this discrepancy, including the possibility of problems with the timing or molecular targets of these state-of-the-art treatments against neurodegeneration in AD.

Hippocampal functions are diffusely disrupted in Aβ plaque-laden mice such as the APPswe/PS1dE9 (APP/PS1) transgenic model that harbors genetic mutations that drive production of the Aβ42 peptide and cause autosomal dominant AD [6,7]. During online memory encoding processes, hippocampal place cells form place fields that are activated sequentially as animals explore an environment in coordination with the hippocampal theta oscillation [8,9]. In Aβ plaque-laden mice, place representations [10,11] and the coupling of the amplitude of gamma oscillations to the phase of theta [12–19] are impaired, and neuronal calcium fluctuations hyper-synchronize with the theta oscillation [20]. Aβ plaque-laden mice also show impaired off-line hippocampal memory consolidation processes: In SWS, reduced reactivation of hippocampal place cell pairs [21] and reduced coordination of hippocampal sharp wave ripples (SWRs) with

cortical spindles and slow oscillations (SO) [20,22]; in REM sleep, the same oscillatory impairments

seen in wakefulness, namely, reduced theta-gamma coupling [20,23,24], and hyper-synchronization

of neuronal calcium fluctuations with theta [20].

Some molecular species arising from aberrant degradation of the amyloid precursor protein (APP)

have been found to be cytotoxic [25], including soluble Aβ oligomers (AβO). AβO impair synapse

function, plasticity, and density [26–36] and have been found to drive hippocampal neuronal hyper-

excitability in anesthetized AD models [37–39]. As AβO arise before plaques form, we hypothesized

that impairments of hippocampal systems functions underlying memory encoding and

consolidation would be evident in advance of plaque deposition. To test this hypothesis, here we

acquired hippocampal dynamic calcium imaging and hippocampal and cortical

electrophysiological recordings across behavioral states in freely-behaving APP/PS1 transgenic

mice in the window before robust plaque deposition [6,40]. We found marked impairments of

hippocampal functions underlying memory encoding and consolidation processes despite

minimal Aβ plaque burden.

Results

To evaluate hippocampal function before plaque deposition, we acquired hippocampal CA1

dynamic calcium imaging with GCaMP6f using the Inscopix miniscope[TM] together with

simultaneous cortical and hippocampal local field potential (LFP) recordings of young APP/PS1

mice (5.2 ± 0.6 months, n=8, male) and their littermate controls (4.7 ± 0.5 months, n=7, male;

age, p=0.4, T=0.8, df=13, t-test), as animals explored a linear track (RUN), and subsequently in

quiet wakefulness (QW), slow wave sleep (SWS) and REM sleep in their home cage (Fig. 1a).

Recording lengths of each state were comparable between genotypes (Fig. S1).

Minimal burden of Aβ plaques and abundant soluble Aβ oligomers in young APP/PS1 mice

In the APP/PS1 model, Aβ plaques have been reported to first deposit around 6 months, with

dense accumulation by 12 months [6,40]. In line with these reports, we observed minimal deposition

of Aβ plaques in hippocampus and cortex in young APP/PS1 mice, in contrast to 12-month-old

mice (hippocampus: repeated measures ANOVA $F_{(4,15)}=42.1$, $p<0.001$, post hoc comparison

with positive control $p<0.001$; cortex: repeated measures ANOVA $F_{(4,15)}=105.4$, $p<0.001$, post

hoc comparison with positive control $p<0.001$, Fig. 1b, c). Soluble AβOs were detected in both

hippocampus and cortex and increased with age (hippocampus $R=0.5$, $p=0.03$; cortex $R=0.6$,

$p=0.006$, Pearson correlations, Fig. 1d, consistent with prior results [41,42]).

*Figure 1. Young APP/PS1 mice expressing minimal Aβ plaque burden and abundant soluble AβO showed preserved place cell function and no calcium hyperactivity across the sleep-wake cycle. a) Schematic of hippocampal calcium imaging and simultaneous ipsilateral LFP recording across the sleep-wake cycle. Insert: examples of LFP spectrogram (color bar indicates power) and one CA1 neuron's calcium trace across behavioral states; and example dataset of place cells sorted by their place fields (color bar indicates probability) in inbound (LR) and outbound (RL) directions. Yellow shows the place field_of each cell. Exploratory behavior (RUN); quiet wakefulness (QW); slow wave sleep (SWS); rapid eye movement (REM) sleep. b) Immunohistochemistry against Aβ in hippocampus of a wild-type control mouse (6 months) and APP/PS1 mice at 6 and 12 months. Green: Aβ plaques (D54D2). c) Aβ plaque counts in hippocampus and cortex were low in 4-6 month-old APP/PS1 mice and increased markedly at 12 months. d) Levels of soluble AβO in hippocampus and cortex increased with age in APP/PS1 mice. e) Spatial information (SI) and f) place field width were comparable across genotypes. g) Calcium event rates of place cells and non place cells in each behavioral state were similar to those of littermate control mice, and calcium event rates of place cells were higher in RUN than SWS in both genotypes. h) In RUN, APP/PS1 had fewer low rate and high rate cells; In QW and SWS, APP/PS1 had more low rate cells and fewer high rate cells. In REM, APP/PS1 had fewer high rate cells. * p<0.05, ** p<0.01, ***p<0.001; c, one-way ANOVA with post-hoc comparisons; d, Pearson correlation; e, f, t-test; g, two-way ANOVA with post-hoc comparisons; h, mixed effects model. b-d: each group n=4 mice; e-h: Control n=5 mice, 395 place cells, 2169 total cells; APP/PS1 n=6 mice, 237 place cells, 2133 total cells. Data are represented as mean □±□S.E.M.*

Intact place representations but reduced dynamic calcium activity in quiet wakefulness and slow wave sleep before plaque deposition

Behavior was comparable across young APP/PS1 and littermate control groups, with similar run velocity (Control: 4.1 ± 0.7 cm/s; APP/PS1: 3.5 ± 0.6 cm/s, p=0.2, T=1.3, df=13, t-test) and laps on the track (Control: 34.2 ± 3.2 laps; APP/PS1: 39.8 ± 2.5 laps, p=0.2, T=1.3, df=13, t-test), consistent with prior reports [43].

To investigate place cell representations across the genotypes, which are often directional [44], we considered inbound (LR) and outbound (RL) trajectories separately (Fig. 1a insert). The proportion of place cells was similar in young APP/PS1 and control mice (Control: 16.8% ± 1.9%; APP/PS1: 11.0% ± 2.1%. p=0.07, T=1.9, df=9, t-test). Spatial information (p=0.5, T=0.5,

df=9, t-test, Fig. 1e) and place field width (p=0.3, T=0.9, df=9; t-test, Fig. 1f) were also similar

across the genotypes. These results are consistent with prior work [45].

Previous studies in plaque-laden AD models have detected aberrant neuronal activity [20,37,38], and

studies in young anesthetized AD mice [37,39] have reported hyperactivity of hippocampal neurons

before amyloid plaque formation. We therefore extended analyses of neuronal activity of young

APP/PS1 mice across the sleep-wake cycle, free from the complex effects of anesthesia [39].

Compared to littermate control mice, young APP/PS1 mice had similar place cell and nonplace

cell mean calcium event rates in each behavioral state (Control place cell vs APP/PS1 place cell:

RUN p = 0.07, QW p = 0.1, SWS p = 0.1, REM p = 0.1; Control non place cell vs APP/PS1 non

place cell: RUN p = 0.1, QW p = 0.1, SWS p = 0.1, REM p = 0.1; two-way repeated-measures

ANOVA with post-hoc comparisons, Fig. 1g, Supplementary Table 1, 2). We did not observe

hyperactivity (defined as calcium event rates > 2SD above the control group mean [20]) in young

APP/PS1 mice in any behavioral state (high rate cell proportions, Control vs APP/PS1, RUN: 4.2%

± 0.4% vs 0.7% ± 0.6%, p<0.001; QW: 5.1% ± 0.4% vs 2.8% ± 1.9%, p=0.001; SWS: 4.9% ±

0.3% vs 1.8% ± 1.5%, p<0.001; REM sleep: 4.4% ± 0.5% vs 1.0% ± 0.8%, p<0.001. Mixed

effects model, Fig 1h). Instead, we observed an increased proportion of hypoactive cells (< 0.25

spikes/min [20,37]) in APP/PS1 mice in QW and SWS (low rate cell proportions, Control vs

APP/PS1, RUN: 19.5% ± 8.2% vs 11.7% ± 1.5%, p<0.001; QW: 31.2% ± 4.7% vs 39.8% ± 5.8%,

p=0.009; SWS: 30.6% ± 3.5% vs 42.5% ± 11.5%, p<0.001; REM sleep: 21.0% ± 4.0% vs 21.5%

± 3.5%, ns. Mixed effects model, Fig 1h), regardless of place cell identity. This finding persisted

when high rate cells were defined using a calcium event rate threshold of 3 Z scores: APP/PS1

mice showed no hyperactivity in any behavioral state (RUN p<0.001, QW p=0.001. SWS

p<0.001; REM p<0.001, Mixed effects model).

Consistent with prior results in plaque-laden APP/PS1 mice [20] (but see [39]), calcium event rates of

place cells were higher in RUN than in SWS in both APP/PS1 and control mice (Control: RUN

vs SWS p = 0.009, RUN vs REM p = 0.01; APP/PS1: RUN vs SWS p = 0.02; two-way repeated-

measures ANOVA with post-hoc comparisons, Fig. 1g, Supplementary Table 1, 2). In both

APP/PS1 and control mice, the calcium event rates of non-place cells were similar across

behavioral states.

Reduced place cell pairwise reactivation despite intact slow wave sleep oscillations before

robust plaque deposition

Since place cell properties of young APP/PS1 mice were essentially intact at the single cell level,

we next evaluated place cell coordination in offline behavioral states. Place cells with

overlapping place fields during spatial behavioral tasks tend to fire together during subsequent

rest and sleep in the service of memory consolidation [46–48], and prior work has found that

reactivation of place cell pairs in SWS is reduced in Aβ plaque-bearing mice [21]. We therefore set

out to evaluate whether place cell reactivation is perturbed prior to robust Aβ plaque deposition.

SWR frequency and duration in SWS were comparable in APP/PS1 and control mice (SWR rate:

Control 0.061 ± 0.014 event/s, APP/PS1 0.089 ± 0.020 event/s, p=0.2, T=1, df=13; duration:

Control 0.024 ± 0.001 s, APP/PS1 0.025 ± 0.001 s, p=0.2, T=1.2, df=13, t-tests). We calculated

the spike time cross-correlation of all place cell pairs during exploratory behavior and in

subsequent offline states, and identified reactivation when a place cell pair was coactive in both

online and offline states [48] (Fig. 2a, see Methods and Fig. S2). Pairwise reactivation was detected

in each genotype by comparing to the distribution of 500 cross-correlations of shuffled spike

times (Supplementary Fig. S3a). The percentage of place cell pairs that reactivated in SWS was

significantly reduced in APP/PS1 compared to control mice ($p=0.005$, two-way repeated-

measure ANOVA with post-hoc comparison, Fig. 2a, b, Supplementary Table 3).

Place cell activity and numbers were comparable in APP/PS1 and control mice (see above), and

this difference persisted after matching place cell event rate and place cell numbers in control

mice and APP/PS1 mice (Supplementary Fig. S3b-d). In contrast to SWS, pairwise reactivation

of APP/PS1 and control mice was comparable in QW (as well as in REM sleep) (QW $p=0.9$,

REM $p=0.2$, two-way repeated-measure ANOVA with post-hoc comparison, Fig 2a, b). Thus,

hippocampal reactivation of place cell pairs in SWS was reduced in APP/PS1 mice prior to

robust Aβ plaque deposition.

Figure 2. Young APP/PS1 mice showed reduced place cell pairwise reactivation but intact oscillations of slow wave sleep. a) Spike time cross-correlations of 30 randomly selected place cells from one control and one APP/PS1 mouse, sorted by place field location on the track (colorbar). Grey line: coactivation. b) The percentage of place cell pairs that were temporally reactivated in slow wave sleep (SWS) was reduced in APP/PS1 compared to control mice, but not in quiet wakefulness (QW) or REM sleep (REMs). c) In SWS, power in the cortical slow oscillation, delta, spindle (left panel) and hippocampal ripple band (right panel) was comparable between genotypes. Inset: cortical slow oscillation and delta power. d) Cortical spindle-triggered ripple power was comparable between genotypes. **p□<□0.01, two-way ANOVA with post-hoc comparison. Control n=5 mice, 395 place cells, 2169 total cells; APP/PS1 n=6 mice, 237 place cells, 2133 total cells. Data are represented as mean□±□S.E.M.

To investigate possible mechanisms underlying reduced APP/PS1 place cell reactivation, we next

evaluate the coordination of SWS hippocampal and cortical oscillations. Hippocampal SWRs,

cortical spindles, cortical slow oscillations and their nested coordination are fundamental features

of SWS that contribute to memory consolidation [49 50]. Whereas these SWS oscillations and their

coordination are degraded in Aβ plaque-laden AD models [20,22], before plaques predominate we

found that they remained intact. Specifically, SWR rates and duration (see above), cortical SO

power, cortical spindle power, and hippocampal ripple power (SO, $p=0.3$, $T=1$, $df=13$; spindle

$p=0.1$, $T=1.4$, $df=13$; ripple: $p=0.3$, $T=1$, $df=13$, t-tests, Fig. 2c) were comparable between young

APP/PS1 and control mice, and the coordination of cortical spindles with hippocampal ripples

was similar across genotypes ($p=0.1$, $T=1.5$, $df=13$, t-test, Fig. 2d). Thus, in contrast to impaired

place cell pairwise reactivation at this early stage of Aβ amyloidosis, the characteristic

hippocampal and cortical oscillations of SWS remain intact.

Impairment of online theta-gamma coupling before plaque deposition

Since the physiological oscillations of SWS were intact in young APP/PS1 mice, we considered

the possibility that the reduction in place cell pairwise reactivation could be related to changes in

online oscillatory physiology linked to memory encoding. Intra-hippocampal and hippocampal-

cortical theta-gamma phase amplitude coupling (PAC) both contribute to memory encoding[13,24]

and degrade in Aβ plaque-bearing AD models [11,20,51–54]. We found that hippocampal theta and

gamma power were comparable across young APP/PS1 and control mice (theta $p=0.9$, $T=0.01$,

$df=13$; gamma $p=0.4$, $T=0.8$, $df=13$, t-tests, Fig. 3a), and theta power was not significantly

correlated with age in either group ($p>0.05$, Pearson's correlation, Fig. 3a). However, intra-

hippocampal theta (6-8 Hz) -gamma (40-80 Hz) PAC in APP/PS1 mice was impaired ($p=0.03$,

$T=2.3$, $df=13$, t-test, Fig. 3b). Consistent with the possibility of accumulating toxicity,

hippocampal theta-gamma PAC in APP/PS1 mice was negatively correlated with age (APP/PS1

R=-0.7, p=0.02; Control p=0.2; Pearson correlations, Fig. 3c). These results were unchanged

after controlling for the possible contribution of theta harmonics [55] (Supplementary Fig. S4 a-c).

Cortical gamma power was also comparable between genotypes (p=0.6, T=0.4, df=13, t-test, Fig.

3d), while theta sampled in cortex was reduced in APP/PS1 mice (p=0.005, T=3.3, df=13, t-test;

(Fig. 3d). Cortical theta power was not significantly correlated with age in APP/PS1 or control

mice (Fig. 3d). In this context, hippocampal theta (6-8 Hz) - cortical gamma (40-80 Hz) PAC

was also impaired during exploratory behavior in young APP/PS1 mice (p=0.02, T=2.5, df=13, t-

test, Fig. 3e). As with intrahippocampal theta-gamma PAC, in APP/PS1 but not control mice,

hippocampal-cortical PAC was negatively correlated with age (APP/PS1 R=-0.8, p=0.007;

Control p=0.9; Pearson correlation, Fig. 3f). This result persisted after controlling for the

possible contribution of theta harmonics [12] (Supplementary Fig. S4 d-f). Together, these results

suggest that accumulating APP degradation products impair theta-gamma PAC well before

plaques robustly deposit.

Interestingly, the magnitude of hippocampal theta-gamma PAC in RUN was strongly correlated

with the extent of place cell pairwise reactivation in subsequent SWS in APP/PS1 mice

(APP/PS1 R=0.8, p=0.02; Control p=0.3; Pearson correlations, Fig. 3g). In contrast, pairwise

reactivation was not correlated with age (APP/PS1 p=0.2; Control p=0.2; Pearson correlations,

Fig. 3h). These observations raise the possibility that the reduction of theta-gamma PAC during

online behavior may contribute to the impairment of place cell pairwise reactivation that arises in

subsequent SWS.

Aberrant synchrony of somatic calcium fluctuations with hippocampal theta before plaque deposition

As oscillations both coordinate and reflect the coordination of neuronal activity, and as calcium signaling links synaptic activity to synaptic plasticity and gene expression, we considered the possibility that breakdown in the coordination of neuronal activity reflected in impaired theta-gamma PAC might manifest at the cellular level in altered coordination of somatic calcium fluctuations with the LFP. In exploratory behavior, neuronal calcium levels can synchronize with theta oscillations, and this synchronization is aberrantly increased in Aβ plaque-laden mice [20]. We therefore measured the coherence of hippocampal neuronal calcium traces with the LFP in APP/PS1 mice before significant plaque accumulation. Compared to control mice, APP/PS1 mice showed hyper-synchronization of place cell calcium fluctuations with theta (p=0.01, two-way repeated-measured ANOVA post-hoc comparisons, Fig. 3i, Supplementary Table 4). Thus, at the pre-plaque stage preceding robust Aβ plaque deposition, dysfunction of memory encoding processes manifest at both the oscillatory and the cellular level.

Figure 3. In pre-plaque APP/PS1 mice, theta-gamma phase-amplitude coupling in exploratory behavior was impaired and dynamic calcium fluctuations were excessively synchronized with hippocampal theta. a) *In RUN, hippocampal theta and gamma power in APP/PS1 and control mice were comparable, and hippocampal theta power was not significantly correlated with age in either group.* ***b)*** *Intra-hippocampal theta (6-8Hz) - gamma (40-80Hz) phase - amplitude coupling (PAC) was reduced in APP/PS1 compared to control mice.* ***c)*** *Intrahippocampal theta-gamma PAC was negatively correlated with the age of APP/PS1 mice but not control mice.* ***d)*** *Cortical theta power was reduced in APP/PS1 mice while cortical gamma power was unchanged. Cortical theta power was not significantly correlated with age in either group.* ***e)*** *PAC of hippocampal theta (6-8Hz) - cortical gamma (40-80Hz) was reduced in APP/PS1 compared to control mice.* ***f)*** *The reduction of hippocampal theta- cortical gamma PAC was negatively correlated with age in APP/PS1 but not control mice.* ***g)*** *In APP/PS1 but not controls, the extent of place cell pairwise reactivation in SWS correlated with the magnitude of intra-hippocampal theta-gamma PAC in RUN.* ***h)*** *The proportion of reactivated place cell pairs in SWS was not correlated with age.* ***i)*** *Average theta-range LFP-cellular coherence of place cells was higher in APP/PS1 than control mice. *p□<□0.05; a, c, d, f, g, h Pearson correlation; a, d, b, e, 2 sided t-test; i, two-way ANOVA with post-hoc comparisons. a – f: Control n=7 mice, APP/PS1 n=8 mice; g – i: Control n=5 mice, 395 place cells, 1774 non-place cells, 2169 total cells; APP/PS1 n=6*

280 *mice, 237 place cells, 1896 non-place cells, 2133 total cells. Data are represented as*
281 *mean $\pm$ S.E.M. pc, place cells. Non-pc, nonplace cells.*
282

REM sleep theta-gamma coupling before plaque deposition

As intra-hippocampal and hippocampal-cortical theta-gamma PAC in REM sleep contribute to

memory consolidation [56,57] and are diminished in Aβ plaque-bearing AD mice [20,53], we evaluated

REM sleep theta-gamma PAC prior to robust Aβ plaque deposition. Hippocampal REM sleep

theta and gamma power (theta p=0.7, T=0.3, df=13; gamma p=0.5, T=0.6, df=13 t-tests, Fig. 4a),

and intra-hippocampal theta (6-8 Hz) -gamma (40-80 Hz) PAC were comparable across

genotypes (p=0.9, T=0.01, df=13, t-test, Fig. 4b), consistent with prior results [53]. In contrast to

RUN, hippocampal theta power in REM sleep of APP/PS1 mice was negatively correlated with

age (R=-0.8, p=0.008, Pearson correlation, Fig. 4a). This correlation was not observed in control

mice. Only a trend level correlation was observed between intra-hippocampal PAC magnitude in

REM sleep of APP/PS1 mice and age (R=-0.6, p=0.05, Pearson correlation, Fig. 4c), and this

result did not survive after controlling for the possible contribution of theta harmonics

(Supplementary Fig. S5 a-c).

REM sleep cortical gamma power (p=0.1, T=1.5, df=13, t-test, Fig. 4d), and hippocampal theta

(6-8 Hz) - cortical gamma (40-80 Hz) PAC were also comparable across genotypes (p=0.2, T=1,

df=13, t-test, Fig. 4e). Even so, in young APP/PS1 but not control mice, cortical theta power

(R=-0.7, p=0.04, Pearson correlation, Fig. 4d) and hippocampal-cortical PAC in REM sleep (R=-

0.8, p=0.005, Pearson correlation, Fig. 4f) were significantly and negatively correlated with age,

and this result persisted after controlling for the possible contribution of theta harmonics

(Supplementary Fig. S5 d-f). These results suggest that as in exploratory behavior, theta-gamma

PAC during REM sleep also falls with accumulation of toxic species before robust plaque

deposition.

Because theta-gamma PAC in APP/PS1 mice was differentially affected in RUN and REM sleep

but negatively correlated with age in both behavioral states, we compared theta-gamma PAC in

RUN versus REM sleep. In both APP/PS1 and control mice, the magnitude of hippocampal theta

- cortical gamma PAC in RUN and REM sleep were strongly correlated (Control R=0.9, p=0.001;

APP/PS1 R=0.6, p=0.04; Pearson correlations, Fig 4h), with only a trend level correlation

between intra-hippocampal theta-gamma PAC in RUN and REM sleep (Control R=0.6, p=0.05;

APP/PS1 R=0.5, p=0.06; Pearson correlations, Fig. 4g).

We next sought to determine whether the hyper-synchronization of somatic calcium levels with

the theta oscillation that we observed in exploratory behavior (Fig. 3i) extended to REM sleep. In

contrast to findings in exploratory behavior, neuronal calcium synchronization with theta in

REM sleep was similar across APP/PS1 and control mice (p=0.2, two-way repeated measures

ANOVA post-hoc comparison, Fig. 4i, Supplementary Table 5). These results suggest that

hippocampal function in REM sleep may be more resilient than in other brain states to the

accumulation of toxic APP cleavage products.

Figure 4. In young APP/PS1 mice, theta-gamma phase amplitude coupling in REM sleep was intact and correlated with theta-gamma phase-amplitude coupling during exploratory behavior. a) *In REM sleep, hippocampal theta and gamma power were comparable between groups. Hippocampal theta power of APP/PS1 but not control mice was negatively correlated with age.* **b)** *Intra-hippocampal theta (6-8Hz) - gamma (40-80Hz) phase-amplitude coupling (PAC) was comparable in APP/PS1 and control mice.* **c)** *Intrahippocampal theta-gamma PAC showed only a trending negative correlation with the age of APP/PS1 mice but not control mice.* **d)** *Cortical theta and gamma power were comparable between groups. Cortical theta power of APP/PS1 mice but not control mice was negatively correlated with age.* **e)** *PAC of hippocampal theta (6-8Hz) - cortical gamma (40-80Hz) was comparable across genotypes.* **f)** *Hippocampal theta- cortical gamma PAC was negatively correlated with age in APP/PS1 mice but not control mice.* **g)** *Intra-hippocampal theta-gamma PAC between RUN and REM sleep showed a trending correlation in both genotypes.* **h)** *Hippocampal theta - cortical gamma PAC in RUN and in REM sleep were significantly correlated in both AP/PS1 and control mice.* **i)** *Average theta-range LFP-cellular coherence in REM sleep was comparable across APP/PS1 and control mice. a, c, d, f, g, h, Pearson correlation; a, d, b, e, 2 sided t-test; i, two-way ANOVA with post-hoc comparisons. a – g: Control n=7 mice, APP/PS1 n=8 mice; i: Control n=5 mice, 395 place cells, 2169 total cells; APP/PS1 n=6 mice, 237 place cells, 2133 total cells. Data are represented as mean□±□S.E.M. pc, place cells. Non-pc, nonplace cells.*

Discussion

In the context of well-described impairments of hippocampal neurophysiology in Aβ plaque-

laden mice yet limited therapeutic benefits in manifest AD from targeting Aβ plaques, we

investigated hippocampal functions that contribute to memory encoding and consolidation in

young APP/PS1 mice harboring AD-associated gene mutations but lacking significant Aβ plaque

burden. We found reduced theta-gamma PAC and excessive theta synchronization of

hippocampal place cells with their surrounding LFP during exploratory behavior, and reduced

place cell pairwise reactivation in subsequent SWS. These diverse impairments of neural

functions underlying memory encoding and consolidation manifest before significant plaque

burden and implicate other toxic processes in AD.

Hippocampal place cells encode the environment during online behavior, and their network

reactivation during offline behaviors, including hippocampal replay during sharp wave ripples

[46,58], contributes to sleep-dependent learning [46,59] and memory consolidation [47,60,61]. Prior work

has reported reduced place cell pairwise reactivation in Aβ plaque bearing mice [21], and the

present results show that this result arises prior to significant Aβ plaque burden. The impairment

of pairwise reactivation did not result from differences in behavior, LFP properties such as ripple

rate and duration, or cellular properties such as calcium event rate between young APP/PS1 and

control mice. Likewise, although the coordination of cortical slow oscillations, corticothalamic

spindles, and hippocampal sharp wave ripples is a key feature of memory consolidation in SWS

[49,50], and this coordination degrades in Aβ plaque-bearing animals [20,22], these oscillations and

their coordination were preserved in young APP/PS1 mice lacking robust plaque deposits. As

such, the reduced place cell pairwise reactivation observed in these mice stands in contrast with

intact SWS oscillatory coordination.

Brain oscillations and their coupling play a crucial role in hippocampal-dependent memory

functions. Online intra-hippocampal and hippocampal-cortical theta-gamma PACs have been

linked to memory encoding [12–19], and, like the coordinated oscillations of SWS, are also impaired

in multiple AD models harboring Aβ plaques [11,20,51–54]. In contrast to the preserved oscillations of

SWS, online hippocampal and hippocampal-cortical theta-gamma PACs were reduced in young

APP/PS1 mice. As the extent of these reductions correlated with age, online theta-gamma PAC

appears to degrade as APP degradation products such as soluble AβO accumulate.

The intra-hippocampal and hippocampal-cortical theta-gamma PACs of REM sleep have also

been linked to memory consolidation [12,13]. While these deteriorate in plaque-laden AD model

mice [20], the present findings show that before significant plaque deposition, theta-gamma PACs

of REM sleep remain grossly preserved, consistent with prior results [22]. Even so, a subtle

vulnerability was evident, in that hippocampal-cortical theta-gamma PAC declined with

increasing age.

In plaque-laden APP/PS1 mice, CA1 neuronal calcium fluctuations are excessively synchronized

with hippocampal theta [20]. Consistent with both this result and the greater vulnerability of online

theta-gamma PAC compared to REM sleep in young APP/PS1 mice, hyper-synchronization of

neuronal calcium oscillations with the hippocampal LFP was detected in young APP/PS1 mice in

exploratory behavior alone. Although the mechanisms underlying impaired theta-gamma PAC

and aberrant synchronization of neuronal calcium levels with the theta oscillation remain to be

determined, brain oscillations and their coupling are known to reflect the coordinated activity of

large populations of excitatory and inhibitory neurons, and their imbalance or dyscoordination

may contribute. In this context, it is possible that Aβ-mediated dysfunction of inhibitory neurons

and their synapses may contribute to these findings [62].

Interestingly, despite the relatively small dataset, the degree of impairment of hippocampal theta-

gamma PAC in exploratory behavior was correlated with the reduction of place cell pairwise

reactivation in subsequent SWS. This observation raises the possibility of a primary deleterious

effect of AβO or other APP degradation product on memory encoding online that then impairs

memory consolidation in SWS. Linked impairments across behavioral states mediated by the

effects of impaired theta-gamma PAC and hyper-synchronized neuronal calcium fluctuations on

theta sequences that link the activity of adjacent place cells into trajectories within each theta

cycle is one potential mechanism for this observation [63–65]. This possibility will be worth

assessing in future studies.

Although we observed marked abnormalities of hippocampal function across the sleep-wake

cycle of young APP/PS1 mice, we did not detect aberrant neuronal hyperactivity, as has been

reported in studies of anesthetized APP/PS1 mice [37–39]. Instead, we observed a tendency towards

hypoactivity in non-theta offline states. Reports of changes in neuronal activity in APP/PS1 mice

have been heterogeneous [39], and prior work has shown that hyperactivity in young APP/PS1

mice is brought out by and depends on anesthesia and recorded cell types (based on the use of

the hsyn vs CAMKII promoter) [39]. In this context, the present results suggest that hippocampal

dynamic calcium hyperactivity in the behaving animal does not appear to be a prominent and

requisite finding of early Aβ pathology.

Interestingly, the hippocampal systems level impairments observed here in young APP/PS1 mice

precede many of the behavioral impairments that arise in this AD model on standard behavioral

assays, although deficits of contextual memory have been reported to start around 6 months [66,67].

Thus, sampling hippocampal systems level function appears to provide a sensitive readout of

brain dysfunction in AD models.

While tau fibrils and neuronal loss are strong predictors of cognitive impairment in AD [68,69], the

current findings support the early contribution of additional AD-associated processes linked to

APP processing. Prior to plaque deposition, APP cleavage products including toxic soluble

species of Aβ accumulate. Among these, AβO are well-known to be cytotoxic and have been

shown to induce cognitive impairment [31,36,70]. Among soluble Aβ species, small oligomers are

particularly toxic [36], and levels of small brain AβO but not monomer have been shown to

correlate with AD progression in APP/PS1 mice in the age range studied here [42].

Given the cellular and synaptic toxicity of AβO and their relevance to AD progression, as well as

the minimal levels of Aβ protofibrils detectable in young APP/PS1 mice [42], the accumulation of

AβO is well-positioned to contribute to the marked systems-level hippocampal dysfunctions

observed here. Even so, other non-Aβ mechanisms resulting from APP overexpression and the

PS1 mutation [25,71] may also contribute. Among these, non-Aβ APP cleavage products such as

sAPPɑ, APP intracellular domain (AICD) and P3 fragment are biologically active [72,73], and

soluble APP products have been associated with cognitive impairment, neuronal loss,

electrophysiological abnormalities and GABA reduction [74]. In addition, PS1 mutations through

their impact on γ-secretase function affect digestion of γ-secretase substrates including but not

limited to APP [75–77]. Like the PS1dE9 mutation present in the APP/PS1 mouse [78], PS1 mutations

typically reduce γ-secretase activity, leading to complex downstream effects including altered

calcium homeostasis [79], increased inflammatory responses in the brain [76], and contributing to

progressive brain atrophy, memory deficits, and synaptic impairments [76]. Future experiments to

dissect the contributions of each potential mechanism to hippocampal systems failure will be of

value.

Strengths and innovations of this study include the use of a well-characterized, freely behaving

AD model to evaluate hippocampal functions underlying memory encoding and consolidation

before Aβ plaques predominate, free from the complexities associated with anesthesia. The

acquisitions of simultaneous hippocampal dynamic calcium imaging and ipsilaterally placed

local electrophysiological recordings is a further strength, enabling us to relate brain oscillations

critical for normal hippocampal mnemonic functions to neuronal calcium fluctuations, place cell

activity, and their reactivation in SWS. Limitations of this study include the use of a single model

of amyloidosis, and additional work in other models will be informative. A further limitation is

the correlative nature of the work. In this regard, future, causal experiments will prove useful to

identify the factors underlying the hippocampal physiological impairments observed here, which

could derive from elevated levels of soluble Aβ42 oligomers or other fragments, and/or from

presenilin 1 dysfunction.

In summary, the results of this study in AD model mice prior to robust Aβ plaque deposition

reveal impairments in hippocampal cellular and oscillatory functions that contribute to both

memory encoding and consolidation. Targeting APP related cascades rather than Aβ plaques, per

se, may improve upon the significant but, to date, only modest effectiveness of anti-amyloid

antibody therapies for AD [1–4].

Materials and Methods

All procedures were approved by the Institutional Animal Care and Use Committees of the

Massachusetts General Hospital and the study was conducted in accordance with the ethical

guidelines of the US National Institutes of Health. This study is reported in accordance with

ARRIVE guidelines.

Surgical preparation

All procedures were similar to our previous studies [20,80]. The mouse strain used for this research

project, B6C3-Tg(APPswe,PSEN1dE9)85Dbo/Mmjax, RRID:MMRRC_034829-JAX, was

obtained from the Mutant Mouse Resource and Research Center (MMRRC) at The Jackson

Laboratory, an NIH-funded strain repository, and was donated to the MMRRC by David

Borchelt, Ph.D., McKnight Brain Institute, University of Florida [7]. Briefly, APP/PS1 mice and

their age/gender matched non-transgenic control litters were first injected with 1 μL AAV-hSyn-

GCaMP6f-GFP into the hippocampal CA1 region (AP -2.1 mm, ML -1.65 mm, DV -1.4 mm)

under anesthesia (isoflurane 1.5-2%). Two weeks later, a grin lens (1 mm diameter, AP -2.1 mm,

ML -1.65 mm, DV -1.4 mm) and a custom drivable probe holding 7 stereotrodes (AP -2.8 mm,

ML 2.6 mm, -36 $^{\circ}$ relative to the AP axis, -39 $^{\circ}$ relative to the inter-aural axis; DV -1.2 mm) were

implanted ipsilaterally, to sample the LFP in the vicinity of the calcium imaging site, with

cerebellar ground. The microscope baseplate was attached three weeks later. Carprofen (10

mg/kg, s.c) was used for postoperative analgesia. Animals were housed in individual cages with

a 12 hr light/12hr dark- standard light cycle. All mice were male to avoid sex differences in Aβ

pathogenesis in APP/PS1 mice [81].

Calcium imaging and electrophysiology recordings

The Inscopix^{TM} nVista3 mini-microscope and acquisition system were used to sample calcium

imaging at 20 Hz. Local field potential (LFP) (Neuralynx^{TM}) was sampled at 2 kHz followed by

0.5-900 Hz filtering. Animal position was recorded with overhead camera tracking of two LED

diodes mounted on the headstage. Electrophysiology and calcium imaging were acquired

simultaneously: Electrophysiology was recorded continuously, while calcium imaging was

recorded in 10 min acquisition blocks, with 4-5 minute intervals off-camera to avoid

photobleaching. These datasets were synchronized and aligned by TTL pulses offline.

Behavior and sleep stages

Data were acquired as animals first traversed a linear track and were then moved to their home

cage within the same recording room for immediate acquisition of subsequent Quiet wakefulness

(QW), slow-wave sleep (SWS), and REM sleep. Running behavior on the track was identified

when run speed exceeded 1 cm/sec; QW was defined as wakeful immobility (< 0.5 cm/sec, eyes

open) in the home cage; SWS was identified when SWRs were present and delta/theta LFP

power ratio exceeded 2, while the animal maintained a sleep posture; REM sleep was identified

when the theta/delta LFP power ratio exceeded 2 in sleep.

Data analysis. All analyses were performed using MATLAB (2019b) and were similar to our previous studies [20,80].

Calcium imaging: Inscopix Data Processing Software (IDPS, Inscopix, Palo Alto, California) was used for movie preprocessing, motion correction, normalization ΔF/F and constrained non-negative matrix factorization (CNMFE) cell identification[82]. The shape and signal-noise-ratio (SNR) of each cell soma was reviewed manually for quality control. Calcium events were then detected at 2 Z scores above the mean, with onset at 30% of peak amplitude. Low rate cells were defined as < 0.25 transients/min [20,83]; high rate cells were defined using a calcium event rate threshold of 2 Z scores above the mean of the control group for each behavioral state [20].

Place cell analyses: Spatial tuning curves (3 cm bins) were constructed for each cell for each running direction using all detected calcium events acquired during running. Spatial information was computed and compared to 1000 shuffled spatial information distributions to test significance (Monte Carlo p-value < 0.05) [84]. Place field width was defined at 50% of peak firing distribution over track location [85]. Spike time cross-correlations of all place cell pairs in both online (exploratory/RUN) and subsequent offline (QW, SWS) states were calculated. Reactivation was defined when the same pair was coactivated during both online and specific offline states [48]. Coactivation was defined based on the distinguished clusters (cluster 1 not coactivated: <0.001, and cluster 2 coactivated: >0.02) in the distribution of correlations of all place cell pairs in each state (Fig. S2). Results persisted across a broad range of thresholds (between >0.001 and <0.02), here we used >0.002 per previous work [48]. Proportions of reactivated pairs among all pairs of each animal were used to compare between genotypes.

Recordings with at least 10 place cells (5 out of 7 control mice and 6 out of 8 APP/PS1 mice)

were included in the pairwise reactivation analysis.

LFP analyses: LFPs were filtered to obtain slow oscillations (0.5-1 Hz), delta (1-4 Hz), theta (4-

10 Hz), gamma (30-150 Hz), and ripples (100-300 Hz). The spectrograms of each channel were

calculated using the Chronux toolbox [86] with a moving window of size 2 s and step size 1 s. A

stereotrode in s. radiatum showing high amplitude theta was used for assessments of RUN and

REM sleep; a stereotrode in s. pyramidale showing high amplitude SWRs was used for

assessments of QW and SWS; a stereotrode within cortex showing slow oscillations and sleep

spindles was also used. Power spectra were computed from 0 to 200 Hz using the Chronux

toolbox [86], and the sum of relative power of each frequency band was used for quantification.

Phase-amplitude-coupling (PAC) modulation index (MI) was calculated with 1-25 Hz for phase

frequency and 1-300 Hz for amplitude frequency using the MATLAB toolbox

https://data.mrc.ox.ac.uk/data-set/matlab-toolbox-estimating-phase-amplitude-coupling [87,88], and

the mean value of the high coupling region was used for quantification [22]. Spindle events were

detected at 1.25 Z score with minimum length of 200 ms; SWR events were detected at 3 Z score

with minimum length of 20 ms [49]. Ripple power triggered on spindle onsets was averaged $\pm$ 0.5 s

around lag = 0 s and compared between groups. LFPs were downsampled to the same sampling

rate as calcium imaging for the Chronux coherence analysis [86] for oscillatory-cellular

synchronization [20].

Biochemistry:

Euthanasia: Mice were euthanized by ketamine-xylazine overdose (240-300 mg/kg ketamine +

15-30 mg/kg xylazine IP).

Brain protein extraction: Mice were perfused with cold PBS containing protease inhibitor (cat #

A32957), and the brain was rapidly excised. One hemisphere was separated for

immunohistochemistry analyses. Cortex and hippocampus were dissected from the remaining

hemisphere and immediately frozen in liquid nitrogen, then stored at −80°C before use. Protein

extraction was performed as indicated by the ELISA kit. In summary, frozen brain samples were

homogenized in 10 vol (w/v) of cold TBS containing protease inhibitor using a Teflon-glass and

centrifuged at 16,000 x g for 30 min at 4°C. The pellets were re-suspended in TBS plus 1%

Triton X-100 containing protease inhibitor (TBS-T) and centrifuged at 16,000 x g for another 30

min at 4°C. The TBS-T fraction was used to measure oligomeric β-amyloid. The total protein

concentration in TBS-T fractions was measured using the BCA protein assay kit (cat# 23225).

ELISA: ELISA was used to quantify oligomeric β-amyloid in cortical and hippocampal samples

(Biosensis cat# BEK-2215-1P/2P). Oligomeric β-amyloid concentration was determined using a

standard curve, and values are reported as ng/mg of protein. Analyses were blinded to animal

age.

Immunohistochemistry: Hemibrains were drop-fixed in 4% paraformaldehyde for two days and

sectioned at 40 µm on a vibratome (Leica, Buffalo Grove, IL). Sections were collected in a 1 in 4

series and stored in PBS at 4°C. Sections were mounted, and antigen retrieval was performed

with citrate buffer at 95°C for 30 min. Next, sections were blocked and permeabilized in TBS-T

with 5% bovine serum albumin (BSA) and 5% normal goat serum (NGS) for 1h. Sections were

incubated overnight in Alexa Fluor 488 Conjugate primary β-Amyloid antibody (D54D2, cat#

51374S) at 4C. Following several washes in PBS, slides were cover slipped with anti-fade

mounting media, and images were acquired using an Olympus VS120 Slide Scanner.

Statistical analysis

All group values were represented as mean ± s.e.m. For analyses of dynamic calcium imaging hypo- and hyperactivity, mixed effects models were used. For the remaining analyses, two-sided *t*-test and two-way ANOVA with repeated measures followed by *post hoc* Tukey-Kramer tests were used for statistical analyses according to the experimental design (SPSS). The significance threshold was set at $p < 0.05$.

Acknowledgements

We thank Joseph J. Locascio for statistical consultation.

Funding disclosure

This work was supported by NIH grants to SNG (R01 AG054551; R01 AG077611).

Author contributions

Conceptualization, H.L., Z.Z. and S.N.G.; Methodology, H.L., Z.Z., A.F. and S.N.G.; Software, H.L. and S.N.G.; Validation, H.L., Z.Z., A.F., H.K.L. and R.J.G.; Formal Analysis, H.L.; Investigation, Z.Z., A.F., H.K.L. and R.J.G.; Resources, H.L., Z.Z., A.F., H.K.L. and R.J.G.; Data Curation, H.L. and R.J.G.; Writing – Original Draft, H.L. and S.N.G.; Writing – Review & Editing, H.L., Z.Z., A.F., H.K.L., R.J.G. and S.N.G.; Visualization, H.L.; Supervision, S.N.G.; Project Administration, S.N.G.; Funding Acquisition, S.N.G.

Competing interests

The authors declare no competing interests.

Data availability statement

598   The datasets generated and analyzed in the current study are available in the Dryad repository,

599   https://datadryad.org/stash/share/gN7A2L9rPMErDH74jxAWQUdjKm7A89v4ERaftNsJOiA

References

1. Bloom, G. S. Amyloid-β and Tau: The Trigger and Bullet in Alzheimer Disease Pathogenesis.

JAMA Neurology **71**, 505–508 (2014).

2. van Dyck, C. H. *et al.* Lecanemab in Early Alzheimer's Disease. *New England Journal of*

Medicine **388**, 9–21 (2023).

3. Mintun, M. A. *et al.* Donanemab in Early Alzheimer's Disease. *N Engl J Med* **384**, 1691–

1704 (2021).

4. Budd Haeberlein, S. *et al.* Two Randomized Phase 3 Studies of Aducanumab in Early

Alzheimer's Disease. *J Prev Alzheimers Dis* **9**, 197–210 (2022).

5. Honig, L. S. *et al.* Trial of Solanezumab for Mild Dementia Due to Alzheimer's Disease. *N*

Engl J Med **378**, 321–330 (2018).

6. Jankowsky, J. L. *et al.* Mutant presenilins specifically elevate the levels of the 42 residue β-

amyloid peptide in vivo: evidence for augmentation of a 42-specific γ secretase. *Human*

Molecular Genetics **13**, 159–170 (2004).

7. Jankowsky, J. L. *et al.* Co-expression of multiple transgenes in mouse CNS: a comparison of

strategies. *Biomol Eng* **17**, 157–165 (2001).

8. O'Keefe, J. Place units in the hippocampus of the freely moving rat. *Experimental Neurology*

51, 78–109 (1976).

9. O'Keefe, J., Burgess, N., Donnett, J. G., Jeffery, K. J. & Maguire, E. A. Place cells,

navigational accuracy, and the human hippocampus. *Philos Trans R Soc Lond B Biol Sci* **353**,

1333–1340 (1998).

10. Cacucci, F., Yi, M., Wills, T. J., Chapman, P. & O'Keefe, J. Place cell firing correlates with

memory deficits and amyloid plaque burden in Tg2576 Alzheimer mouse model.

Proceedings of the National Academy of Sciences **105**, 7863–7868 (2008).

11. Mably, A. J., Gereke, B. J., Jones, D. T. & Colgin, L. L. Impairments in spatial

representations and rhythmic coordination of place cells in the 3xTg mouse model of

Alzheimer's disease. *Hippocampus* **27**, 378–392 (2017).

12. Belluscio, M. A., Mizuseki, K., Schmidt, R., Kempter, R. & Buzsaki, G. Cross-Frequency

Phase-Phase Coupling between Theta and Gamma Oscillations in the Hippocampus. *Journal*

of Neuroscience **32**, 423–435 (2012).

13. Colgin, L. L. Rhythms of the hippocampal network. *Nat Rev Neurosci* **17**, 239–249 (2016).

14. Jacobson, T. K. *et al.* Hippocampal theta, gamma, and theta-gamma coupling: effects of

aging, environmental change, and cholinergic activation. *Journal of Neurophysiology* **109**,

1852–1865 (2013).

15. Kendrick, K. M. *et al.* Learning alters theta amplitude, theta-gamma coupling and neuronal

synchronization in inferotemporal cortex. *BMC Neuroscience* **12**, 55 (2011).

16. Lisman, J. E. & Jensen, O. The Theta-Gamma Neural Code. *Neuron* **77**, 1002–1016 (2013).

17. Tamura, M., Spellman, T. J., Rosen, A. M., Gogos, J. A. & Gordon, J. A. Hippocampal-

prefrontal theta-gamma coupling during performance of a spatial working memory task. *Nat*

Commun **8**, 2182 (2017).

18. Tort, A. B. L., Komorowski, R. W., Manns, J. R., Kopell, N. J. & Eichenbaum, H. Theta–

gamma coupling increases during the learning of item–context associations. *Proc. Natl. Acad.*

Sci. U.S.A. **106**, 20942–20947 (2009).

19. Vivekananda, U. *et al.* Theta power and theta-gamma coupling support long-term spatial

memory retrieval. *Hippocampus* **31**, 213–220 (2021).

20. Zhou, H. *et al.* Disruption of hippocampal neuronal circuit function depends upon behavioral

state in the APP/PS1 mouse model of Alzheimer's disease. *Sci Rep* **12**, 21022 (2022).

21. Prince, S. M. *et al.* Alzheimer's pathology causes impaired inhibitory connections and

reactivation of spatial codes during spatial navigation. *Cell Rep* **35**, 109008 (2021).

22. Zhurakovskaya, E. *et al.* Impaired hippocampal-cortical coupling but preserved local

synchrony during sleep in APP/PS1 mice modeling Alzheimer's disease. *Sci Rep* **9**, 5380

(2019).

23. Buzsáki, G. Theta Oscillations in the Hippocampus. *Neuron* **33**, 325–340 (2002).

24. Colgin, L. L. Theta–gamma coupling in the entorhinal–hippocampal system. *Current*

Opinion in Neurobiology **31**, 45–50 (2015).

25. O'Brien, R. J. & Wong, P. C. Amyloid precursor protein processing and Alzheimer's disease.

Annu Rev Neurosci **34**, 185–204 (2011).

26. Arbel-Ornath, M. *et al.* Soluble oligomeric amyloid-β induces calcium dyshomeostasis that

precedes synapse loss in the living mouse brain. *Molecular Neurodegeneration* **12**, 27 (2017).

27. Cleary, J. P. *et al.* Natural oligomers of the amyloid-β protein specifically disrupt cognitive

function. *Nat Neurosci* **8**, 79–84 (2005).

28. Cline, E. N., Bicca, M. A., Viola, K. L. & Klein, W. L. The Amyloid-β Oligomer Hypothesis:

Beginning of the Third Decade. *JAD* **64**, S567–S610 (2018).

29. Ding, Y. *et al.* Amyloid Beta Oligomers Target to Extracellular and Intracellular Neuronal

Synaptic Proteins in Alzheimer's Disease. *Frontiers in Neurology* **10**, (2019).

30. Faucher, P., Mons, N., Micheau, J., Louis, C. & Beracochea, D. J. Hippocampal Injections of

Oligomeric Amyloid β-peptide (1–42) Induce Selective Working Memory Deficits and Long-

lasting Alterations of ERK Signaling Pathway. *Frontiers in Aging Neuroscience* **7**, (2016).

31. Ferreira, S. T., Lourenco, M. V., Oliveira, M. M. & De Felice, F. G. Soluble amyloid-β

oligomers as synaptotoxins leading to cognitive impairment in Alzheimer's disease. *Front*

Cell Neurosci **9**, 191 (2015).

32. Nicole, O. *et al.* Soluble amyloid beta oligomers block the learning-induced increase in

hippocampal sharp wave-ripple rate and impair spatial memory formation. *Sci Rep* **6**, 22728

(2016).

33. Özcan, G. G., Lim, S., Leighton, P. L., Allison, W. T. & Rihel, J. Sleep is bi-directionally

modified by amyloid beta oligomers. *eLife* **9**, e53995 (2020).

34. Shankar, G. M. *et al.* Amyloid-β protein dimers isolated directly from Alzheimer's brains

impair synaptic plasticity and memory. *Nat Med* **14**, 837–842 (2008).

35. Malinow, R. & Malenka, R. C. AMPA receptor trafficking and synaptic plasticity. *Annu Rev*

Neurosci **25**, 103–126 (2002).

36. Yang, T., Li, S., Xu, H., Walsh, D. M. & Selkoe, D. J. Large Soluble Oligomers of Amyloid

β-Protein from Alzheimer Brain Are Far Less Neuroactive Than the Smaller Oligomers to

Which They Dissociate. *J Neurosci* **37**, 152–163 (2017).

37. Busche, M. A. *et al.* Critical role of soluble amyloid-β for early hippocampal hyperactivity in

a mouse model of Alzheimer's disease. *Proc. Natl. Acad. Sci. U.S.A.* **109**, 8740–8745 (2012).

38. Busche, M. A. & Konnerth, A. Neuronal hyperactivity – A key defect in Alzheimer's disease?

BioEssays **37**, 624–632 (2015).

39. Zarhin, D. *et al.* Disrupted neural correlates of anesthesia and sleep reveal early circuit

dysfunctions in Alzheimer models. *Cell Reports* **38**, 110268 (2022).

40. Garcia-Alloza, M. *et al.* Characterization of amyloid deposition in the APPswe/PS1dE9

mouse model of Alzheimer disease. *Neurobiol Dis* **24**, 516–524 (2006).

41. Takeda, S. *et al.* Brain interstitial oligomeric amyloid β increases with age and is resistant to

clearance from brain in a mouse model of Alzheimer's disease. *FASEB J* **27**, 3239–3248

(2013).

42. Zhang, L. *et al.* Dynamic Changes in the Levels of Amyloid-β42 Species in the Brain and

Periphery of APP/PS1 Mice and Their Significance for Alzheimer's Disease. *Front Mol*

Neurosci **14**, 723317 (2021).

43. Webster, S. J., Bachstetter, A. D. & Van Eldik, L. J. Comprehensive behavioral

characterization of an APP/PS-1 double knock-in mouse model of Alzheimer's disease.

Alzheimers Res Ther **5**, 28 (2013).

44. Knierim, J. J., Kudrimoti, H. S. & McNaughton, B. L. Place cells, head direction cells, and

the learning of landmark stability. *J Neurosci* **15**, 1648–1659 (1995).

45. Jun, H. *et al.* Disrupted Place Cell Remapping and Impaired Grid Cells in a Knockin Model

of Alzheimer's Disease. *Neuron* **107**, 1095-1112.e6 (2020).

46. Ego-Stengel, V. & Wilson, M. A. Disruption of ripple-associated hippocampal activity during

rest impairs spatial learning in the rat. *Hippocampus* **20**, 1–10 (2010).

47. Ólafsdóttir, H. F., Bush, D. & Barry, C. The Role of Hippocampal Replay in Memory

and Planning. *Curr Biol* **28**, R37–R50 (2018).

48. Wilson, M. A. & McNaughton, B. L. Reactivation of Hippocampal Ensemble Memories

During Sleep. *Science* **265**, 676–679 (1994).

49. Siapas, A. G. & Wilson, M. A. Coordinated Interactions between Hippocampal Ripples and

Cortical Spindles during Slow-Wave Sleep. *Neuron* **21**, 1123–1128 (1998).

50. Born, J., Rasch, B. & Gais, S. Sleep to Remember. *Neuroscientist* **12**, 410–424 (2006).

51. Bazzigaluppi, P. *et al.* Early-stage attenuation of phase-amplitude coupling in the

hippocampus and medial prefrontal cortex in a transgenic rat model of Alzheimer's disease.

Journal of Neurochemistry **144**, 669–679 (2018).

52. Etter, G. *et al.* Optogenetic gamma stimulation rescues memory impairments in an

Alzheimer's disease mouse model. *Nat Commun* **10**, 5322 (2019).

53. Gurevicius, K., Lipponen, A. & Tanila, H. Increased Cortical and Thalamic Excitability in

Freely Moving APPswe/PS1dE9 Mice Modeling Epileptic Activity Associated with

Alzheimer's Disease. *Cerebral Cortex* **23**, 1148–1158 (2013).

54. Stoiljkovic, M., Kelley, C., Stutz, B., Horvath, T. L. & Hajós, M. Altered Cortical and

Hippocampal Excitability in TgF344-AD Rats Modeling Alzheimer's Disease Pathology.

Cerebral Cortex **29**, 2716–2727 (2019).

55. Rodriguez-Larios, J. & Alaerts, K. Tracking Transient Changes in the Neural Frequency

Architecture: Harmonic Relationships between Theta and Alpha Peaks Facilitate Cognitive

Performance. *J Neurosci* **39**, 6291–6298 (2019).

56. Bandarabadi, M. *et al.* Dynamic modulation of theta-gamma coupling during rapid eye

movement sleep. *Sleep* **42**, zsz182 (2019).

57. Montgomery, S. M., Sirota, A. & Buzsáki, G. Theta and Gamma Coordination of

Hippocampal Networks during Waking and Rapid Eye Movement Sleep. *J. Neurosci.* **28**,

6731–6741 (2008).

58. Joo, H. R. & Frank, L. M. The hippocampal sharp wave-ripple in memory retrieval for

immediate use and consolidation. *Nat Rev Neurosci* **19**, 744–757 (2018).

59. Girardeau, G., Cei, A. & Zugaro, M. Learning-Induced Plasticity Regulates Hippocampal

Sharp Wave-Ripple Drive. *J. Neurosci.* **34**, 5176–5183 (2014).

60. Atherton, L. A., Dupret, D. & Mellor, J. R. Memory trace replay: the shaping of memory

consolidation by neuromodulation. *Trends Neurosci* **38**, 560–570 (2015).

61. Louie, K. & Wilson, M. A. Temporally Structured Replay of Awake Hippocampal Ensemble

Activity during Rapid Eye Movement Sleep. *Neuron* **29**, 145–156 (2001).

62. Kurucu, H. *et al.* Inhibitory synapse loss and accumulation of amyloid beta in inhibitory

presynaptic terminals in Alzheimer's disease. *European Journal of Neurology* **29**, 1311–1323

(2022).

63. Tang, W. & Jadhav, S. P. Multiple-Timescale Representations of Space: Linking Memory to

Navigation. *Annu Rev Neurosci* **45**, 1–21 (2022).

64. Lisman, J. The theta/gamma discrete phase code occuring during the hippocampal phase

precession may be a more general brain coding scheme. *Hippocampus* **15**, 913–922 (2005).

65. Skaggs, W. E., McNaughton, B. L., Wilson, M. A. & Barnes, C. A. Theta phase precession in

hippocampal neuronal populations and the compression of temporal sequences.

Hippocampus **6**, 149–172 (1996).

66. Lonnemann, N., Korte, M. & Hosseini, S. Repeated performance of spatial memory tasks

ameliorates cognitive decline in APP/PS1 mice. *Behavioural Brain Research* **438**, 114218

(2023).

67. Kilgore, M. *et al.* Inhibitors of Class 1 Histone Deacetylases Reverse Contextual Memory

Deficits in a Mouse Model of Alzheimer's Disease. *Neuropsychopharmacology* **35**, 870–880

(2010).

68. Medeiros, R., Baglietto-Vargas, D. & LaFerla, F. M. The role of tau in Alzheimer's disease

and related disorders. *CNS Neurosci Ther* **17**, 514–524 (2011).

69. Niikura, T., Tajima, H. & Kita, Y. Neuronal Cell Death in Alzheimer's Disease and a

Neuroprotective Factor, Humanin. *Curr Neuropharmacol* **4**, 139–147 (2006).

70. Lambert, M. P. *et al.* Diffusible, nonfibrillar ligands derived from Abeta1-42 are potent

central nervous system neurotoxins. *Proc Natl Acad Sci U S A* **95**, 6448–6453 (1998).

71. Hampel, H. *et al.* The Amyloid-β Pathway in Alzheimer's Disease. *Mol Psychiatry* **26**, 5481–

5503 (2021).

72. Chow, V. W., Mattson, M. P., Wong, P. C. & Gleichmann, M. An Overview of APP

Processing Enzymes and Products. *Neuromolecular Med* **12**, 1–12 (2010).

73. Bukhari, H. *et al.* Small things matter: Implications of APP intracellular domain AICD

nuclear signaling in the progression and pathogenesis of Alzheimer's disease. *Prog*

Neurobiol **156**, 189–213 (2017).

74. Kreis, A. *et al.* Overexpression of wild-type human amyloid precursor protein alters

GABAergic transmission. *Sci Rep* **11**, 17600 (2021).

75. Do, H. N., Devkota, S., Bhattarai, A., Wolfe, M. S. & Miao, Y. Effects of presenilin-1

familial Alzheimer's disease mutations on γ-secretase activation for cleavage of amyloid

precursor protein. *Commun Biol* **6**, 1–14 (2023).

76. Soto-Faguás, C. M., Sanchez-Molina, P. & Saura, C. A. Loss of presenilin function enhances

tau phosphorylation and aggregation in mice. *Acta Neuropathol Commun* **9**, 162 (2021).

77. Kelleher, R. J. & Shen, J. Presenilin-1 mutations and Alzheimer's disease. *Proc Natl Acad*

Sci U S A **114**, 629–631 (2017).

78. Woodruff, G. *et al.* The Presenilin-1 ΔE9 mutation results in reduced γ-secretase activity, but

not total loss of PS1 function, in isogenic human stem cells. *Cell Rep* **5**,

10.1016/j.celrep.2013.10.018 (2013).

79. Bagaria, J., Bagyinszky, E. & An, S. S. A. Genetics, Functions, and Clinical Impact of

Presenilin-1 (PSEN1) Gene. *Int J Mol Sci* **23**, 10970 (2022).

80. Zhou, H. *et al.* Cholinergic modulation of hippocampal calcium activity across the sleep-

wake cycle. *eLife* **8**, e39777 (2019).

81. Li, X. *et al.* Sex differences between APPswePS1dE9 mice in A-beta accumulation and

pancreatic islet function during the development of Alzheimer's disease. *Lab Anim* **50**, 275–

285 (2016).

82. Zhou, P. *et al.* Efficient and accurate extraction of in vivo calcium signals from

microendoscopic video data. *eLife* **7**, e28728 (2018).

83. Lerdkrai, C. *et al.* Intracellular Ca $^{2+}$ stores control in vivo neuronal hyperactivity in a mouse

model of Alzheimer's disease. *Proc. Natl. Acad. Sci. U.S.A.* **115**, (2018).

84. Markus, E. J., Barnes, C. A., McNaughton, B. L., Gladden, V. L. & Skaggs, W. E. Spatial

information content and reliability of hippocampal CA1 neurons: Effects of visual input.

Hippocampus **4**, 410–421 (1994).

85. Go, M. A. *et al.* Place Cells in Head-Fixed Mice Navigating a Floating Real-World

Environment. *Frontiers in Cellular Neuroscience* **15**, (2021).

86. Bokil, H., Andrews, P., Kulkarni, J. E., Mehta, S. & Mitra, P. P. Chronux: A platform for

analyzing neural signals. *Journal of Neuroscience Methods* **192**, 146–151 (2010).

87. Onslow, A. C. E., Bogacz, R. & Jones, M. W. Quantifying phase–amplitude coupling in

neuronal network oscillations. *Progress in Biophysics and Molecular Biology* **105**, 49–57

(2011).

88. Osipova, D., Hermes, D. & Jensen, O. Gamma Power Is Phase-Locked to Posterior Alpha

Activity. *PLoS ONE* **3**, e3990 (2008).