# INSTINCT: Multi-sample integration of spatial chromatin accessibility sequencing data via stochastic domain translation

Yuyao Liu[1, *], Zhen Li[1, *], Xiaoyang Chen[1], Xuejian Cui[1], Zijing Gao[1], and Rui Jiang[1, ✉]

[1] Ministry of Education Key Laboratory of Bioinformatics, Bioinformatics Division at the Beijing National Research Center for Information Science and Technology, Center for Synthetic and Systems Biology, Department of Automation, Tsinghua University, Beijing 100084, China

[*] These authors contributed equally: Yuyao Liu, Zhen Li

[✉] Corresponding author: ruijiang@tsinghua.edu.cn

## Abstract

Recent advances in spatial epigenomic techniques have given rise to spatial assay for transposase-accessible chromatin using sequencing (spATAC-seq) data, enabling the characterization of epigenomic heterogeneity and spatial information simultaneously. Integrative analysis of multiple spATAC-seq samples, for which no method has been developed, allows for effective identification and elimination of unwanted non-biological factors within the data, enabling comprehensive exploration of tissue structures and providing a holistic epigenomic landscape, thereby facilitating the discovery of biological implications and the study of regulatory processes. In this article, we present INSTINCT, a method for multi-sample INtegration of Spatial chromaTIN accessibility sequencing data via stochastiC domain Translation. INSTINCT can efficiently handle the high dimensionality of spATAC-seq data and eliminate the complex noise and batch effects of samples from different conditions through a stochastic domain translation procedure. We demonstrate the superiority and robustness of INSTINCT in integrating spATAC-seq data across multiple simulated scenarios and real datasets. Additionally, we highlight the advantages of INSTINCT in spatial domain identification, visualization, spot-type annotation, and various downstream analyses, including expression enrichment analysis and partitioned heritability analysis.

**Keywords:** Data integration; spATAC-seq data; multi-sample; batch correction; stochastic domain translation

## Introduction

Chromatin accessibility is a measurement of the degree to which trans-acting factors can physically bind with cis-regulatory elements within open chromatin regions during DNA replication or transcription[1,2]. With the advances of single-cell epigenomic techniques[3-7] in the recent years, abundant single-cell assay for transposase-accessible chromatin using sequencing (scATAC-seq) data have been generated, facilitating the study of heterogeneity among cells and contributing to the exploration of the mechanisms of biological regulatory networks[8]. However, the majority of intricate regulatory processes rely on the intercellular transmission of ligand-receptor pairs within the tissue microenvironment, a dependence largely contingent upon the physical proximity of cells[9-11], whereas scATAC-seq data offers only epigenetic profiles of individual cells without spatial information. This limitation impedes the attainment of a comprehensive understanding of tissues and organs through the analysis of scATAC-seq data.

With the innovation of spatial omics, emerging spatial epigenomic techniques, such as spatial ATAC[12], spatial-ATAC-seq[13], and SCA-seq[14], along with some spatial multi-omics techniques, such as MISAR-seq[15] and spatial ATAC-RNA-seq[16], have given rise to spatial assay for transposase-accessible chromatin using sequencing (spATAC-seq) data. Although such data presently do not attain single-cell resolution, wherein each barcoded spot on a sequencing array typically aggregates multiple cells, these techniques do provide coordinates for each spot based on its spatial location on the array. This capability empowers researchers

to amalgamate spatial and epigenomic data, thereby facilitating the dissection of the structure and function of tissues and organs at various hierarchical levels. As the number of spATAC-seq datasets continues rising, it becomes more and more urgent to conduct joint comparative analysis across multiple samples, for the purpose of fully leveraging information in each sample towards comprehensive understanding of the regulatory process within the tissue microenvironment.

Integrative analysis has long been a challenging and meaningful topic for both single-cell omics[17-21] and spatial omics[22,23]. Joint modeling of multiple samples can establish connections between different samples and enhance the statistical power for the analysis of shared biological variations, making samples comparable by effectively identifying and eliminating the unwanted non-biological factors within the data. Furthermore, joint modeling enables more detailed analyses and comparisons of temporal dynamics and spatial structures[24], shedding light on the fundamental processes of biology and disease[21]. By integrating multiple spATAC-seq samples, researchers can comprehensively characterize spatial tissue structures and obtain a holistic epigenomic landscape, thereby facilitating the discovery of biological implications and contributing to the study of regulatory logic.

Many computational methods have been developed for multi-batch integration of single-cell or spatially resolved transcriptomics (SRT) data, enabling the integration of data from different samples, tissues, donors, developmental stages and technologies[25-31]. SCALEX[25] is a method based on the variational autoencoder (VAE)[32] for modeling single-cell RNA-seq or ATAC-seq data. By providing batch information only to the decoder, the batch-free encoder is enabled to merely extract biological information. PeakVI[26] uses a Bernoulli distribution to model scATAC-seq data and employs a VAE combined with an auxiliary neural network to predict distribution parameters and perform batch correction. SEDR[27] fuses spatial

information with transcriptional profiles by leveraging a masked autoencoder and a variational graph convolutional autoencoder (GCN)[33], while combining Harmony[20] to eliminate batch effects. STAligner[28] employs a graph attention autoencoder (GAT)[34] for modeling SRT data, utilizing spot triplet loss for batch correction. STitch3D[31] first pre-aligns multiple SRT samples to establish connections between them, and achieves the removal of batch effects using a GAT-based batch-free encoder and a batch-specific decoder. However, no integration method specifically for spATAC-seq data has been developed.

Several fundamental challenges need to be addressed to accomplish the integrative analysis of spATAC-seq datasets. Firstly, spatial information should be incorporated into the analysis of spATAC-seq data effectively[9,35]. Methods for modeling scATAC-seq data mentioned above generally suffer from a common problem: they lack a design to utilize spatial information. Directly applying these methods to integrate spATAC-seq data may result in the wastage of spatial information, leading to unreliable biological analysis. For instance, after clustering, spots of the same cluster may be scattered in space, making it difficult to identify meaningful spatial domains with strong continuity. Secondly, spATAC-seq data generally exhibits high dimensionality, great sparsity, and suffers from a notable incidence of data missing due to a low capturing rate[3,36,37]. These characteristics result in different data patterns compared to SRT data, rendering some mechanisms for modeling SRT data unsuitable for analyzing spATAC-seq data, and therefore, may not yield satisfactory results. For example, STAligner[28] identifies nearest neighbors across various samples using Euclidean distance and employs spot triplet learning to bring these identified neighbors, likely belonging to the same category, closer in latent space, thereby mitigating batch effects. However, we found that utilizing cosine similarity is more appropriate for spATAC-seq data, yielding better results compared to using Euclidean distance (Supplementary Figs. 1 and 2).

Therefore, there is an urgent need for integration algorithms specifically designed for spATAC-seq data. Thirdly, spATAC-seq data presents complex patterns of noise and batch effects, posing challenges in distinguishing biological variations from non-biological factors. A procedure that can effectively remove noise and batch effects while preserving biological variations needs to be introduced, enabling comparability across multiple spATAC-seq samples for joint analysis[38]. Fourthly, the absence of methodologies capable of annotating spATAC-seq data complicates the exploration of biological significance relying on annotated domains.

To this end, we present INSTINCT, a method for multi-sample integration of spatial chromatin accessibility sequencing data via stochastic domain translation, to address these gaps. To the best of our knowledge, INSTINCT is the first method specifically designed for integrative analysis of multiple spATAC-seq samples. INSTINCT trains a variant of graph attention[34] autoencoder to effectively integrate spatial information and epigenetic profiles, implements a stochastic domain translation procedure[39-41] to facilitate batch correction, and finally obtains low-dimensional representations of spots in a shared latent space where the batch effects are eliminated and biological variations are preserved. The representations of spots can then be utilized for spatial domain identification, visualization, spot-type annotation and various downstream analyses, including expression enrichment analysis and partitioned heritability analysis. Additionally, based on the spot-type annotation results, INSTINCT can reveal the underlying biological implications, illustrating its potential to guide and contribute to the study of biological traits and pathology.

## Results

### Overview of INSTINCT

INSTINCT is designed based on the premise that, due to noise and batch effects resulting from varying experimental conditions and techniques, spots from different samples may belong to different domains in the feature space, even when some spots share similar biological attributes. With this consideration, INSTINCT aims to properly integrate these domains by eliminating unwanted non-biological bias while preserving biological variations among spots from different samples, thereby facilitating the integrative analysis of multi-sample spATAC-seq data.

As shown in Fig. 1a, INSTINCT adopts a two-stage strategy. In stage 1, the model comprises a graph attention (GAT) encoder[34], a multilayer perceptron (MLP) decoder[42], and a discriminator. Taking the preprocessed data matrix of multiple samples ($X$) and the concatenated neighbor graph as input, the GAT encoder fuses epigenetic profiles and spatial information of spots, thereby projecting the data from different slices into a shared latent space ($Z$). The MLP decoder reconstructs each spot back to the domain of its original slice. The discriminator is pretrained to simultaneously judge the authenticity of the input data ($X$) and the reconstructed data ($X_{rec}$), as well as the category of spots. In stage 2, the model further incorporates a noise generator component besides the GAT encoder, MLP decoder, and discriminator that are already trained in stage 1. A stochastic domain translation procedure[39-41] is implemented to facilitate batch correction (Fig. 1b), wherein different patterns of noise and batch effects are artificially added to the low-dimensional representation of each spot, enabling flexible projection to the domains of different samples (Methods).

Specifically, during each training epoch in stage 2, each slice is assigned at random a target domain that is different from its original one, and spots from the slice need to be

mapped to this new target domain. After the extraction of biological variations performed by the encoder, the noise generator captures the original pattern of batch effects of each spot in the latent space using the latent representation matrix and a one-hot matrix indicating the original slices of spots, and a pseudo-noise generation (PNG) process simulates a different pattern of batch effects for each spot based on the output of the noise generator and the target domain of that spot (Fig. 1a, lower). These two patterns of batch effects are then added separately to the low-dimensional representation of the corresponding spot, enabling either reconstruction to the domain of its original slice ($X_{rec}$) or generation to the randomly selected target domain ($Y$) through the decoder. The discriminator further performs the discrimination and classification process by simultaneously judging the authenticity of the input data and the generated data, as well as the category of spots, thereby ensuring a more accurate data generation and domain translation process through adversarial learning. This approach encourages the encoder to extract merely the biological variations and eliminate the batch effects.

After training, the latent representation matrix $Z$, which contains joint representations of spots without noise and batch effects, is used for spatial domain identification, visualization, spot-type annotation, and various downstream analyses such as expression enrichment analysis and partitioned heritability analysis (Fig. 1c, Methods).

**INSTINCT shows superiority and robustness in comprehensive simulated scenarios**

We first conducted comprehensive and realistic simulated experiments in six scenarios to validate the ability of INSTINCT in data integration.

Inspired by SpatialPCA[43], we adopted a similar approach to simulate spATAC-seq datasets (Fig. 2a, Methods). Initially, we utilized simCAS[44] to generate three scATAC-seq

datasets that exhibited varying degrees of batch effects, each comprising 10,000 cells distributed among five cell-types labeled as integers 0 to 4. Each of these three datasets was used to generate one spATAC-seq sample. We pre-defined a single-cell resolution slice of size $120 \times 150$ as a template, which was divided into five domains represented by domain 0 to 4, respectively. Each domain corresponded to a major cell-type with the same number as its index (Fig. 2a), which was assigned to locations within that domain with a probability of $r$, and other cell-types were each assigned with a probability of $\frac{1-r}{4}$. Subsequently, we defined each $3 \times 3$ cell grid as a spot to simulate the resolution of spATAC-seq techniques. The category of each spot was determined by the cell-type that predominantly occupied that spot. Evident batch effects between the three simulated slices can be observed in all scenarios (Supplementary Fig. 3).

For scenario 1, the probability $r$, which is used to assign major cell-types to domains, is set to 0.8. We integrated the three slices and compared INSTINCT with several baseline integration methods for both single-cell omics and spatial omics (Fig. 2b, Methods). Integration methods for single-cell omics include Scanorama[21], SCALEX[25], PeakVI[26], with the latter two methods being specifically developed for integrating scATAC-seq data. Meanwhile, methods for spatial omics encompass SEDR[27], STAligner[28] and GraphST[29], all designed for integrating SRT data, as no method has been specifically developed for spATAC-seq data. We compared the capabilities of these integration methods in three aspects: clustering performance, representation quality, and batch correction, using a total of 12 metrics (Methods), with the first two aspect assessing biological conservation. INSTINCT achieved the highest score in 9 of the 12 metrics and second in two. It also ranked first on the group-specific overall scores of clustering performance and representation quality, as well as the final overall score, and ranked second on the overall score of batch correction.

Visualizations in Supplementary Fig. 4 also demonstrate the superior data integration capability of INSTINCT. Most of the baseline methods only integrated the two slices where the batch effects between them were not very pronounced (slices 0 and 2), while failing to mix the data from the other slice with them. However, INSTINCT evenly mixed the data from three batches, with spots of identical type gathering together while different types remain separated. While STAligner also integrated data from different batches, it mistakenly mixed different spot-types together, making it difficult to distinguish between them, which resulted in significantly inferior clustering performance compared to INSTINCT.

We reduced the value of $r$ continuously across the first three scenarios, resulting in increasing levels of confusion among spot-types and progressively more complex batch effects (Methods, Supplementary Fig. 3). The same procedure was applied to evaluate the performance of all the methods in scenario 2 and 3. In the progressively challenging scenarios, INSTINCT consistently outperformed other methods (Fig. 2c, Supplementary Figs. 5 and 6). Specifically, in terms of clustering performance and representation quality, INSTINCT consistently performed the best across all three scenarios. While in batch correction, it ranked second in the first two scenarios, with slightly lower scores than STAligner. As the scenarios became more challenging, INSTINCT surpassed STAligner in batch correction in the third scenario, securing the first position. Besides, INSTINCT achieved the highest final overall score in all the three scenarios, demonstrating its superiority and robustness. SCALEX and STAligner performed relatively well compared to other baseline methods. However, as the scenarios became more complex, and different types of spots became increasingly mixed before integration, the confusion in the low-dimensional representations obtained by SCALEX for each spot also increased (Supplementary Figs. 4 and 7). Consequently, distinguishing between different spot-types became more difficult. The same issue

consistently occurred for STAligner across all the three scenarios (Supplementary Figs. 4 and 7). In contrast, INSTINCT not only evenly mixed data from three slices but also accurately distinguished between different spot-types in the latent space across scenarios 1 to 3 (Supplementary Figs. 4 and 7). This suggests that INSTINCT possesses the ability to not only effectively remove noise and batch effects but also accurately capture and highlight biological variations, contributing to subsequent analyses and downstream tasks.

In certain situations, vertically adjacent slices may not exhibit complete overlap, where different slices display varying proportions of spot-types, or a slice may lack some spot-types found in other slices. This issue can also arise in samples from different developmental stages. Therefore, based on scenario 1, we designed scenarios 4 and 5 to simulate two cases where the proportions of spot-types differ between slices (Methods). In scenario 4, we applied different cropping to the three slices from scenario 1 to generate three new slices (Supplementary Fig. 8). For slice 0, the proportion of spot-type 1 noticeably decreased, while for slice 2, the proportions of spot-types 2 and 3 decreased significantly. Despite the clear differences in the number and proportion of certain spot-types across different slices, INSTINCT still yielded excellent results when integrating the three slices in this scenario (Fig. 2d). In scenario 5, we retained slice 0 and slice 1 from scenario 1 and removed one domain from slice 2 (Supplementary Fig. 8). We applied INSTINCT to scenario 5 and observed that, although there was no spot identical to spot-type 3 in slice 2, INSTINCT did not incorrectly mix different spot-types in slice 2 together with spot-type 3 in other slices, as marked in Fig. 2d. These experiments demonstrate the flexibility and reliability of INSTINCT in various situations.

We further simulated scenario 6 based on scenario 1 to validate the ability of INSTINCT to preserve rare and specific spot-types (Methods). We excluded a certain domain from slices

1 and 2 of scenario 1, retaining only a $5 \times 5$ area of that domain in slice 0 and removing the rest, resulting in a rare spot-type with merely about 25 spots (Supplementary Fig. 8). After data integration performed by INSTINCT, the rare spot-type was perfectly preserved in latent space, as marked in Fig. 2e, indicating that INSTINCT can effectively identify unique biological signals in samples and prevent overfitting.

**INSTINCT integrates multiple samples across different developmental stages**

We applied INSTINCT to a mouse brain dataset[15] generated by MISAR-seq (MISAR-seq MB). The dataset comprises eight spATAC-seq data slices corresponding to developmental stages E11.0, E13.5, E15.5, and E18.5, with two vertically adjacent slices from each stage. We applied INSTINCT and baseline methods to integrate the four section 1 (S1) slices with annotations from each stage and evaluated the integration performance using various metrics. We used the provided annotations as ground truth spot-types (Fig. 3f). Since certain spot-types appear exclusively in specific slices, we introduced two additional metrics, average silhouette width across isolated labels (isolated label ASW) and isolated label F1-score[45], to the metric group of representation quality to evaluate the preservation of sample-specific spot-types (Methods). INSTINCT ranked first in 10 out of 14 evaluation metrics and second in one. In terms of overall scores, INSTINCT secured the top position in clustering performance, representation quality, as well as the final overall score, and came second in batch correction (Fig. 3a, b).

We used Uniform Manifold Approximation and Projection (UMAP)[46] (Methods) to visualize the data before and after integration (Fig. 3c, d, Supplementary Fig. 9). Prior to integration, noticeable batch effects were observed among the four slices. However, after integrating with INSTINCT, the data from different slices were effectively mixed, removing

the batch effects between them. Through visualization (Fig. 3e), we found that INSTINCT not only accurately distinguished major categories such as cartilage, mesenchyme, and cerebral regions (diencephalon and hindbrain etc.), but also preserved many rare spot-types, such as muscle, thalamus, ventricular zone of dorsal pallium (DPallv), a subtype of cartilage (cartilage_1), and the two subtypes of subpallium (subpallium_1 and subpallium_2). As marked in Fig. 3e, most of these rare spot-types only exist in one or two developmental stages. For instance, subpallium_1 only exists in slice of E18.5, while thalamus and subpallium_2 only exist in slices of E15.5 and E18.5. INSTINCT ensured that these types of spots retained their unique biological characteristics in the low-dimensional representation, enabling them to be distinguishable from other spot-types after integration. In contrast, most of the baseline methods could only roughly differentiate several major categories with significant biological differences and erroneously mixed many rare spot-types into these major categories (Fig. 3e, Supplementary Fig. 10), resulting in the failure to capture and preserve the biological variations between different spot-types. Among all the rare spot-types, SCALEX and PeakVI only distinguished subpallium_1 and subpallium_2 from other spot-types, Scanorama and STAligner barely moved subpallium_1 to the edge, but its boundary with other spot-types was not clear, and the remaining two methods failed to preserve any rare spot-types. It is worth noting that none of the methods distinguished muscle, thalamus, DPallv, and cartilage_1 like INSTINCT did. This demonstrates that INSTINCT can not only effectively remove batch effects but also preserve extensive biological variations, thereby obtaining meaningful results in practical applications.

Subsequently, we jointly clustered the low-dimensional representations for each spot learnt by INSTINCT and mapped the identified clusters to the annotated categories (Fig. 3g, Methods). INSTINCT accurately identified numerous spatial domains across slices that hold

practical significance. For instance, in the E11.0 slice, INSTINCT captured a relatively intact structure of the primary brain (cluster 11 and 12). Consistently across the other three slices, INSTINCT identified DPallv (cluster 6) and the mantle zone of dorsal pallium (DPallm) (cluster 5). Particularly, the structures of DPallm and DPallv identified in the E15.5 and E18.5 slices closely resembled the ground truth. Cartilage_1 (cluster 1) and subpallium_1 (cluster 13) were accurately identified in the E18.5 slice. Subpallium_2 primarily exists in the E15.5 slice and only remains in a small portion at the lower end of the subpallium in the E18.5 slice. We found that this domain (cluster 14) was accurately identified not only in the E15.5 slice but also in the E18.5 slice. Thalamus (cluster 15), present only in the slices of the E15.5 and E18.5, was also accurately identified. Compared to other methods, INSTINCT also exhibited more accurate and comprehensive recognition of large domains, such as mesenchyme (cluster 8), cartilage (cluster 1 to 4), and a domain representing diencephalon and hindbrain (cluster 7).

All baseline methods performed poorly in the spatial domain identification task (Fig. 3g, Supplementary Fig. 11). While the three integration methods for scATAC-seq data were able to cluster most spots within certain large spatial domains into a single category, the clustering results were not spatially coherent. For example, SCALEX, while relatively accurately depicting the boundary of a large domain representing the diencephalon and hindbrain (cluster 7), exhibited scattered spots of multiple different clusters within it. This not only rendered the identified spatial domains unreliable but also led to the failure to identify domains occupying smaller portions of the slices, such as thalamus, which none of the single-cell epigenomic integration methods recognized. This is because spATAC-seq data, representing the combined epigenetic profiles of multiple cells at each spot[16], is more prone to confusion and less distinguishable between different categories compared to scATAC-seq

data. Moreover, these methods did not consider the association of each spot with its surrounding environment by overlooking the spatial information provided by spATAC-seq data. While the three integration methods for SRT data could identify continuous spatial domains, their mechanisms were not suitable for spATAC-seq data, resulting in structures of the identified domains vastly differing from the actual tissue structures and lacking practical significance. For instance, although STAligner seemed to identify the thalamus region (cluster 15) in the E15.5 and E18.5 slices, its structure significantly differed from the ground truth and had a much larger volume than the actual thalamus (Fig. 3f, g). INSTINCT overcomes the limitations of both types of methods by modeling specifically for spATAC-seq data and incorporating spatial information with it. The low-dimensional representations of spots can be utilized to yield superior results in spatial domain identification by accurately identifying continuous domains with boundaries consistent with those of real tissues.

**INSTINCT acquires joint representations of samples across different scales**

We utilized INSTINCT to integrate three samples of mouse postnatal day 21/22 (P21/22) brains[16] generated by spatial ATAC-RNA-seq (spatial ATAC-RNA-seq MB), a multi-omics sequencing technique. It is worth noting that although these three slices were sequenced from similar developmental stages of the same organ, the first two slices (slice 0 and 1) have a size of $50 \times 50$ barcodes, while the third slice (slice 2) contains $100 \times 100$ barcodes. This resulted in the sequencing of brain tissues of different scales, with slice 2 essentially encompassing the entire hemisphere of the coronal brain, while slice 0 and 1 only contained approximately one-quarter of the size.

We used INSTINCT to integrate these three samples with different tissue scales and performed clustering (Methods). The spots exhibited more obvious multi-group gathering

patterns, and 13 clusters were identified, including cluster 11 and 12 unique to slice 2, and cluster 9 predominantly present in slice 2 (Fig. 4b). We colored spots according to their clusters and restored their original arrangements and found that these clusters exhibited strong continuity in space (Fig. 4c). According to the anatomical reference of the coronal brain of adult mouse from Allen Brain Atlas website[47] (Fig. 4a), we found that clustering restored the true tissue structure and identified spatial domains with practical significance. These include the caudate putamen (clusters 0 and 6), lateral ventricle (cluster 10), corpus callosum (cluster 3), lateral septal nucleus (cluster 4), and nucleus accumbens (cluster 9). Additionally, hierarchical structures were identified in the cortex (clusters 1, 2, 5, 8, and 11), and a small domain representing the island of Calleja (cluster 12), which only exists in slice 2, was also identified.

Since spatial ATAC-RNA-seq involves multi-omics sequencing, the dataset also includes paired SRT data for these three slices, which provided the transcriptional landscapes. Using the SRT data corresponding to each spot, we identified the differentially expressed genes (DEGs) for each region (Methods) and visualized the expression levels of some marker genes (Fig. 4d, Supplementary Fig. 12). For example, in the caudate putamen, marker genes such as $Rgs9$[48], $Pde10a$[49], $Gng7$[50], $Bcl11b$[51], and $Foxp1$[52] showed significantly higher expression levels. INSTINCT made the spots in this domain more clustered compared to before integration. $Sox4$[53], $Dlx1$[54], and $Zbtb20$[55] were predominantly expressed in the lateral ventricle. Before integration, these spots were not clearly distinguished from others, but after integration, they were separated and aggregated in a small area. Cluster 3, representing the corpus callosum, showed high expression of genes such as $Plp1$[56] and $Mbp$[57]. While spots of cluster 3 were mixed internally and difficult to distinguish from other categories before integration, INSTINCT made them separately gathered and identified as a distinct category

through clustering. Marker genes for the island of Calleja, such as *Isl1*[58] and *Rreb1*[59], were highly expressed only in cluster 12 unique to slice 2, indicating that INSTINCT, when integrating samples with different tissue scales, can preserve non-shared rare spot-types. Additionally, the expression of *Dgkg*[60] in the lateral septal nucleus and *Mef2c*[61] in the cortex further demonstrated that INSTINCT better distinguishes between spot-types that were previously difficult to differentiate (Supplementary Fig. 12), proving its effectiveness not only in integrating samples with different tissue scales but also in highlighting the biological variations between different spot-types.

**INSTINCT facilitates the integrated analysis of whole mouse organisms**

We further applied INSTINCT to a mouse embryo dataset[12] generated by spatial ATAC (spatial ATAC ME), which consists of six slices from developmental stages E12.5, E13.5, and E15.5, with two vertically adjacent slices from each stage (Fig. 5d, Supplementary Figs. 13 and 14). Initially, we utilized INSTINCT to integrate the S1 slices of the three stages and calculated the evaluation metrics based on their ground truth annotations. We did not utilize metrics for evaluating the preservation of sample-specific spot-types as each slice contained the same categories. INSTINCT ranked first in 8 out of 12 metrics (Supplementary Fig. 15), and for the overall scores, it ranked first in clustering performance, representation quality, as well as the final overall score (Fig. 5a).

However, the performance of INSTINCT in batch correction was not particularly high. We note that most baseline methods overfitted on this dataset, failing to accurately capture biological variations. As a result, these overfitted methods obtained low representation quality and clustering scores but relatively higher batch correction performance. Visualization results revealed that most methods improperly mixed data from the three slices,

resulting in a blend of various spot-types that made it challenging to accurately discern any spot-type (Fig. 5b, c, Supplementary Fig. 16). In contrast, INSTINCT accurately distinguished the liver from other spot-types. Furthermore, INSTINCT did not excessively mix all slices and accurately categorized these spots into four major groups: central nervous system (CNS), meningeal and peripheral nervous system (PNS), mesenchyme and limb, and liver (Fig. 5c), consistent with the results reported in the original study[12].

Through clustering, while other methods exhibited scattered spatial distributions of spots belonging to the same cluster, failing to accurately identify any spatial domain, INSTINCT performed much better (Fig. 5e, Supplementary Fig. 17). For instance, it accurately identified the liver (cluster 5) in slices of E12.5 and E13.5. In the E15.5 slice, it identified spatial relationships within brain tissue, distinguishing between forebrain (cluster 1), midbrain (cluster 8), and hindbrain (cluster 2).

Additionally, we utilized INSTINCT to integrate section 2 (S2) slices from the three developmental stages, as well as collectively integrating all six slices, and compared with other methods (Supplementary Figs. 18 to 26), obtaining conclusions consistent with those of integrating S1 slices. This indicates that INSTINCT can integrate multiple samples of whole organisms and accurately capture underlying biological variations, demonstrating robustness and effective avoidance of overfitting.

**INSTINCT enables accurate cross-sample annotation and reveals biological implications in annotated domains**

As spATAC-seq data becomes increasingly abundant, the need for effective annotation methods grows more urgent, which is crucial for understanding biological processes across different samples and revealing biological implications based on annotated domains.

However, there is currently no annotation algorithm specifically designed for spATAC-seq data. INSTINCT addresses this gap by enabling accurate cross-sample annotation, enhancing the analysis and interpretation of spatial epigenomic landscapes and facilitating the discovery of biological insights.

Initially, we used the integration results of various methods to perform cross-sample spot-type annotation on the MISAR-seq MB dataset[15] (Methods). It was completed by employing a k-nearest-neighbor (KNN) classifier, a model-free approach whose annotation efficacy heavily depends on the effectiveness of the low-dimensional representations across samples. Each time, we selected one slice as the target and used the others as reference to annotate the target slice (Supplementary Figs. 27 to 30). We evaluated the annotation effectiveness using four metrics (Methods): accuracy, kappa, macro F1-score (mF1), and weighted F1-score (wF1). INSTINCT consistently ranked first on all four metrics (Fig. 6a). The same evaluation procedure was applied to spatial ATAC ME dataset[12], where INSTINCT again outperformed all other methods across all metrics (Supplementary Fig. 31). This preliminary evidence demonstrates that the accurate data integration achieved by INSTINCT facilitates cross-sample annotation.

Next, for the MISAR-seq MB dataset[15], we used the S1 slices with ground truth annotations from four developmental stages to annotate the corresponding unannotated S2 slices, respectively. Before integration, batch effects existed between each S1 and its corresponding S2 slice (Supplementary Fig. 32). Initially, we preprocessed the samples using the preprocessing procedure provided by INSTINCT, and then directly reduced the dimensionality of the data to match the latent representations of spots obtained by INSTINCT using principal component analysis (PCA)[62]. We found that when annotating samples with complex structures, such as slices from E15.5 and E18.5, spots labeled as the same type were

spatially discontinuous, resulting in inaccurate identification of tissue structure (Supplementary Fig. 33). Moreover, due to batch effects, spots of a certain type in S1 slices were not mixed with those labeled as the same type in corresponding S2 slices, rendering the annotation results unreliable.

We then integrated each pair of S1 and S2 slices from each developmental stage separately using INSTINCT and performed annotation using the joint low-dimensional representations. We observed that after integration, batch effects between the two vertically adjacent slices were eliminated (Supplementary Fig. 34), and the spatial structures of the annotations in S2 slices were consistent with those of the corresponding S1 slices (Supplementary Fig. 35). For instance, in E18.5 slices which possessed the most complex structures, the relative spatial relationships between different spatial domains annotated in S2 were highly consistent with those in S1 (Fig. 6b). INSTINCT successfully annotated many important structures such as subpallium_1, DPallv, DPallm, thalamus, and their shapes and volumes were nearly identical to the corresponding regions in S1. Notably, in the S1 slice of E18.5, there were two types of spots with very low proportions, subpallium_2 and muscle. Similar proportions of these two types of spots were also annotated in similar spatial locations in the S2 slice. We conducted gene ontology (GO) analysis (Fig. 6c, Methods) using SRT data paired with the S2 slice of E18.5, and found that for the annotated subpallium_2, there was enrichment in the biological process of forebrain neuron differentiation, consistent with the results obtained using four S1 slices in the original study[15]. For the category annotated as muscle, enrichment was observed in many biological processes related to muscle, including muscle contraction and skeletal muscle contraction, consistent with the enrichments observed in the original study[15].

We further identified spot-type-specific peaks for each annotated spot-type, using the

remaining peaks as background peaks (Methods), and performed expression enrichment analysis for the single nucleotide polymorphisms (SNPs) in each set of spot-type-specific peaks and the set of background peaks, respectively (Fig. 6d, Supplementary Figs. 36 and 37, Methods). We found that genes in spot-type-specific peaks of all cerebral regions were predominantly enriched in brain-related tissues. Conversely, genes in the specific peaks of non-brain-specific tissues, such as cartilage and muscle, did not exhibit enrichment in brain-related tissues. Additionally, genes in the background peaks were generally not enriched in brain-related tissues. This further confirms that the annotation provided by INSTINCT can help distinguish spots from different biological systems with distinct functionalities.

Furthermore, we utilized the SNPs found in each set of spot-type-specific peaks and the set of background peaks to perform partitioned heritability analysis (Methods). We found that in many psychiatric disorders and mental-related phenotypes, the enrichment of heritability in cerebral regions was significantly higher than that in mesenchyme, cartilage, and muscle (Fig. 6e, Supplementary Figs. 38 and 39), and it tended to be biased towards certain specific regions. For instance, attention deficit hyperactivity disorder (ADHD) is associated with the pallium and thalamus[63]. Our result validated this association, as its heritability was significantly enriched in subpallium_2, DPallv, DPallm, and thalamus, while other regions exhibited relatively lower levels of enrichment. For schizophrenia, heritability was significantly enriched in the thalamus, highlighting its importance in the implicated circuitry, which has been validated in previous studies[64,65]. Autism has been linked to the dorsal pallium[66], and the result indicated that the enrichment of heritability in DPallv was significantly higher than in DPallm, aiding in pinpointing a specific location within the tissue. This emphasizes that INSTINCT can reveal biological implications in the annotated domains, thereby guiding and contributing to the studies of biological traits and pathology.

## Discussion

Recent advances in spATAC-seq technologies have led to an increasing amount of spATAC-seq data, which simultaneously provides epigenomic landscapes and spatial information. However, existing integration methods for other omics are not well-suited for integrating this data modality. We propose INSTINCT as the first method capable of effectively integrating spATAC-seq data, demonstrating superiority across multiple tasks and providing a guidance for spATAC-seq data integration. INSTINCT can fully utilize spatial information, allowing our method to identify tissue regions with stronger spatial continuity compared to integration methods developed for scATAC-seq data. Furthermore, the stochastic domain translation step in INSTINCT enables the noise generator to effectively extract noise and batch effects from the low-dimensional representations, allowing the representations of spots to effectively highlight biological variations compared to other methods. We also demonstrated that the outstanding integration results of spATAC-seq data provided by INSTINCT can be utilized for cross-sample annotation and further reveal biological implications in the annotated domains.

To demonstrate the effectiveness of the design adopted in INSTINCT, we conducted a series of ablation experiments on four annotated S1 slices of the MISAR-seq MB dataset 错误! 未找到引用源。. Firstly, we performed ablation experiments on the designed loss functions (Methods). It was observed that the exclusion of any loss function resulted in a degradation of the performance of the model (Supplementary Figs. 40 and 41), thereby substantiating the efficacy of all loss functions. Additionally, since INSTINCT includes a noise generator and discriminator for achieving stochastic domain transformation (Methods), we conducted ablation experiments on the model structure. We compared the integration effectiveness of the complete INSTINCT model with the model where the discriminator was removed, as

well as the model where both the discriminator and noise generator were removed. We did not consider the model where only the noise generator was removed due to the structural constraints of INSTINCT. We found that the complete model achieved the best performance, demonstrating the effectiveness of the discriminator and noise generator modules (Supplementary Figs. 42 and 43). Additionally, we observed that the performance of the model was better when both the discriminator and noise generator were removed compared to when only the discriminator was removed. This suggests that the discriminator plays a crucial role in constraining noise simulation and stochastic domain translation. Lastly, since we employed a clamp strategy in INSTINCT to prevent the erroneous mixing of different types of spots while enhancing the conservation of spot-types unique to a particular slice (Methods), we conducted an ablation experiment for it. we found that adopting this strategy led to improvements in all three aspects including clustering performance, representation quality, and batch correction (Supplementary Figs. 44 and 45). It is noteworthy that employing this strategy enhanced the performance of INSTINCT on isolated label ASW and isolated label F1-score, demonstrating the ability of this strategy to preserve sample-specific spot-types.

Moreover, we envision potential avenues for further optimizing INSTINCT. Firstly, inspired by methods developed for SRT data[31], exploring an efficient cell-type decomposition method and embedding it into INSTINCT could elevate spATAC-seq data to single-cell resolution, enabling more precise exploration of biological properties at different scales. Secondly, constructing associations between different samples through effective alignment algorithms could offer a 3D perspective[29,31,67], facilitating a more comprehensive study of the influence of the microenvironment. Thirdly, the structure of the decoder can be altered to a single linear layer, allowing spot-type-specific peaks or cluster-specific peaks to be defined

in a model-specific way based on the weight parameters of the decoder network[37,68]. This can aid in recognizing spot-types, providing a more intuitive understanding of the mechanisms of regulatory networks and facilitating various downstream tasks. Lastly, we have demonstrated that using cosine similarity to find neighbors is superior to using Euclidean distance for the integration of spATAC-seq data. Other mechanisms more suitable for this data modality should also be gradually explored.

## Methods

### Data preprocessing

A single sample (slice) of spATAC-seq dataset consists of a spot-by-peak data matrix and a corresponding spatial coordinate matrix. We define $E = [e_{ij}] \in \mathbb{N}^{n \times p}$ as the data matrix of a single slice, where each row represents a spot and each column represents a feature, specifically for spATAC-seq data, a peak. $E$ is typically a high-dimensional matrix with high sparsity. We take $D = [d_{ij}] \in \mathbb{R}^{n \times 2}$ as the spatial coordinate matrix, where each row represents a 2D coordinate vector corresponding to a barcoded spot on the sequencing array.

We employed a standardized preprocessing procedure for multi-samples ($E_s = [e_{s_{ij}}] \in \mathbb{N}^{n_s \times p_s}, for\ s \in \{0, 1, \dots, S - 1\}$, where $S$ is the total number of slices, and $n_s$ and $p_s$ are the number of spots and number of peaks for the $s$-th slice, respectively) prior to integration. Given that each sample each possesses its specific set of peaks as features, and these peaks from different peak sets may exhibit overlapping regions but are typically not identical, it become impractical to directly concatenate the data matrices of different samples. Therefore, we initially conducted a peak merging process across samples according to the instructions provided by Signac[69]. We amalgamated peak sets from two slices at a time. Peaks from both slices, exhibiting overlapping regions, were amalgamated into larger merged peaks.

Specifically, if a pair of peaks from distinct slices contained an identical subregion, we

merged them and took the shortest region that encompassed these peaks as the merged peak.

This process persisted until no overlapping peaks remained unmerged. We filtered out the

unmerged peaks as well as peaks with sequence lengths greater than 10,000 bp or less than

20 bp, and the remaining peaks served as the merged peak set. Subsequently, the peak sets of

the remaining slices were alternately merged with the merged peak set in the same manner.

Finally, the values from the original data matrix were transferred to the corresponding merged

peaks and accumulated to form a series of new data matrices ($X_s = \left[ x_{s_{ij}} \right] \in \mathbb{N}^{n_s \times M}, for\ s \in$

$\{0, 1, \dots, S - 1\}$, where $M$ is the number of merged peaks for all slices) with identical peak

set. We then concatenated all the slices to form a larger data matrix $X_{concat} \in \mathbb{N}^{N \times M}$, where

$N = \sum_{s=0}^{S-1} n_s$ is the total number of spots across all slices.

We converted the read count matrix into a fragment count matrix following a previously

established method[70]. It has been proven that the fragment count matrix better preserves

quantitative regulatory information, thereby enhancing the analysis of ATAC-seq data[70].

Subsequently, according to the sequencing depth of different datasets, we filtered out peaks

accessible in spots fewer than a certain proportion and removed spots with zero count using

*EpiScanpy* (v0.4.0) package[71]. For the spatial ATAC ME dataset[12], this proportion was set to

0.003, and for the spatial ATAC-RNA-seq MB dataset[16], this proportion was set to 0.02, while

for other datasets, it was set to 0.03. Then, we applied the term frequency-inverse document

frequency (TF-IDF) transformation, which has been widely used for preprocessing

epigenomics data[7,69,72-74], on the concatenated data matrix $X_{concat}$. Finally, we performed

PCA on the transformed matrix and retained the top 100 principal components as $X_{pca} \in$

$\mathbb{R}^{N \times 100}$. For ease of subsequent elaboration, we refer to $X_{pca}$ as $X$.

**Construction of spatial neighbor graph**

To fully leverage the spatial information provided by spATAC-seq data, we constructed a spatial graph based on the spatial coordinate matrices. Initially, we computed pairwise distances between spots within each individual slice. The thresholds for adjacency determination of each slice was set to 1.5 times the minimum distance between spots within that slice. Pairs of spots closer than the corresponding threshold were deemed adjacent, while those farther apart were considered non-adjacent, resulting in a symmetric adjacency matrix $(A_s = \left[ a_{s_{ij}} \right] \in \mathbb{N}^{n_s \times n_s}, for\ s \in \{0, 1, \dots, S-1\})$ for each slice. Here, $a_{s_{ij}} = 1$ indicates adjacency between the $i$-th and $j$-th spot in the $s$-th slice, while $a_{s_{ij}} = 0$ denotes non-adjacency. These matrices were then arranged diagonally to construct an overall spatial neighbor graph $A \in \mathbb{N}^{N \times N}$, where spots from different slices were considered non-adjacent.

**The model structure of INSTINCT**

INSTINCT is an integration method which takes the first 100 principal components of the TF-IDF-transformed data matrix ($X$) and the overall spatial neighbor graph ($A$) as input, and generate a low-dimensional representation for each spot in all slices where batch effects have been removed. The low-dimensional representations of spots can be used for spatial domain identification, visualization, spot-type annotation and various downstream analyses such as expression enrichment analysis and partitioned heritability analysis to elucidate biological principles and facilitate biological discoveries.

INSTINCT comprises four components: an encoder, a noise generator, a decoder and a discriminator, where the former three constitute the generator. Adversarial training is applied to train the model.

**Encoder.** The encoder adopts a graph attention network (GAT)[34], which, for each input node, integrates not only the information of the node itself but also that of its related nodes based on an input graph. This enables INSTINCT to fully leverage the spatial information provided by spATAC-seq data. The graph attention mechanism of the $l$-th layer is as follows:

$$h_i^{(l)} = \sigma\left(\sum_{j\in\mathcal{N}_i} \alpha_{ij}^{(l)} W^{(l)} h_j^{(l-1)}\right), for\ l \in \{1, 2, \dots, L\}$$

$$\alpha_{ij}^{(l)} = \frac{a_{ij}\exp\left(e_{ij}^{(l)}\right)}{\sum_{k\in\mathcal{N}_i} a_{ik}\exp\left(e_{ik}^{(l)}\right)}$$

$$e_{ij}^{(l)} = \text{sigmoid}\left(v_0^{(l)^T} W^{(l)} h_i^{(l-1)} + v_1^{(l)^T} W^{(l)} h_j^{(l-1)}\right)$$

where $h_i^{(l-1)}$ and $h_j^{(l-1)}$ represent the input vectors of the $i$-th and $j$-th spots, respectively, while $h_i^{(l)}$ denotes the output representation of the $i$-th spot. $h_i^{(0)}$ is equivalent to the $i$-th row of $X$. $W^{(l)}$, $v_0^{(l)}$ and $v_1^{(l)}$ are the trainable parameters of the $l$-th layer. $a_{ij}$ signifies the attention score of the $i$-th and $j$-th spots for the $i$-th spot, measuring the impact of the $j$-th spot on the $i$-th spot. $\sigma(\cdot)$ is the activation function. The output of the $L$-th layer is the low-dimensional latent representation matrix $Z = [z_{ij}] \in \mathbb{R}^{N\times P}$, where $P$ represents the dimensionality of the shared latent space.

**Noise generator.** The noise generator is structured as a linear network comprising only a single layer, dedicated to capturing the pattern of the noise and batch effects for each spot. It takes as input the concatenation of the low-dimensional representation matrix $Z$ and a one-hot matrix $Q \in \mathbb{R}^{N\times S}$, which encodes the slice indices of each spot. Specifically, each row of $Q$ is the one-hot vector indicating the slice index of the corresponding spot, with a dimension equal to the number of slices.

We assume that the noise and batch effects of each spot are influenced by the

characteristics of that spot, the characteristics of its surrounding environment, and the specific attributes of the slice it belongs to. Since the encoder utilizes graph attention mechanism, the representations of spots in $Z$ already incorporates information of each spot and its surrounding environment, while the slice-specific information is encoded in $Q$.

The noise generator outputs a matrix $N_X \in \mathbb{R}^{N \times P}$, identical in size to $Z$, where each row signifies the representation of noise and batch effects for the corresponding spot in the low-dimensional space.

**Decoder.** The decoder is a multi-layer perceptron (MLP)[42] that takes as input the low-dimensional representation matrix containing noise and batch effects, mapping each spot to the domain of a specified slice. Since INSTINCT involves a stochastic domain translation procedure which maps each spot to different domains, a process to be detailed later, the decoder is responsible for both reconstructing spots back to domains of their original slices and mapping spots to the domains of other slices. The output of the decoder is either the reconstructed data matrix $X_{rec} \in \mathbb{R}^{N \times 100}$ or the generated fake data matrix $Y \in \mathbb{R}^{N \times 100}$.

**Discriminator.** The discriminator is an essential component for adversarial learning[75]. We employ the discriminator to simultaneously discern the authenticity of the data and the category of each spot. The discriminator has a shared first layer and separate second layers which result in two outputs: a vector $d_{src} \in \mathbb{R}^N$ for predicting the authenticity of the data and a matrix $D_{cls} \in \mathbb{R}^{N \times S}$ for predicting the category of the data, i.e., the sample index of each spot. In $d_{src}$, higher values indicate that the representation of the corresponding spot is more likely to be from real data. This architecture, widely used in image processing for adversarial training[39,40], offers the advantage of flexible translation between multiple

domains using a single encoder and decoder.

We designate $X$ as real data, and $X_{rec}$ and $Y$ as fake data. The index of the slice where

the spot originally belongs serves as the category label for the corresponding spot in $X$ and

$X_{rec}$. Additionally, we assign the index of the specified target slices chosen during the

stochastic translation procedure as the category label for the corresponding spot in $Y$, which

will be detailed later.

The encoder of INSTINCT utilizes two graph attention layers. The initial layer

diminishes the dimensionality of the input representations of spots to 50, and the subsequent

layer further reduces it to 30. We use ELU[76] as the activation function. Similarly, the decoder

consists of two layers: the first layer expands the dimensionality to 50, and the second layer

further expands it back to 100. The shared first layer of the discriminator reduces the

dimensionality to 50, and the separate second layers decrease the dimensionality to the

dimension of the corresponding output, respectively.

INSTINCT adopts a two-stage strategy. During stage 1, we exclusively utilize the encoder,

decoder, and discriminator to construct the model. The operational procedure unfolds as

follows:

664
$$Z = \mathrm{E}(X, A)$$

665
$$X_{rec} = \mathrm{D}(Z)$$

666
$$d_{src}^{X} = \mathrm{Disc}_{src}(X)$$

667
$$D_{cls}^{X} = \mathrm{Disc}_{cls}(X)$$

668
$$d_{src}^{X_{rec}} = \mathrm{Disc}_{src}(X_{rec})$$

669
$$D_{cls}^{X_{rec}} = \mathrm{Disc}_{cls}(X_{rec})$$

670 where $\mathrm{E}(\cdot)$, $\mathrm{D}(\cdot)$, $\mathrm{Disc}_{src}(\cdot)$ and $\mathrm{Disc}_{cls}(\cdot)$ denote the encoder, the decoder, the part

where the discriminator distinguishes between real and fake data, and the part where the discriminator distinguishes between data categories, respectively. The encoder takes $X$ and $A$ as input, yielding $Z$ as illustrated earlier, while the decoder directly uses $Z$ as input to reconstruct the input data matrix. In this stage, we also pretrain the discriminator to distinguish between the fake data matrix $X_{rec}$ and the real data matrix $X$, as well as to classify the spots based on the indices of their original slices.

We declare that the training in stage 1 serves merely as a pretraining step, aimed at enabling the model to initially learn feature extraction and data reconstruction, and allowing the discriminator to acquire a certain level of data discrimination capability. This is expected to enhance the effectiveness of batch effect removal during training in stage 2. However, the low-dimensional representations learned in this step still contain noise and batch effects.

In stage 2, we add the noise generator module to the model to capture the patterns the of noise and batch effects for each spot in the latent space. Additionally, employing a stochastic domain translation process, each spot is mapped to a specified target domain distinct from its original one. The operational mechanism is detailed below:

$$Z = \mathrm{E}(X, A)$$

$$N_X = \mathrm{NG}([Z \parallel Q])$$

$$N_X \xrightarrow{\mathrm{PNG}} N_Y$$

$$Z_X = Z + N_X$$

$$Z_Y = Z + N_Y$$

$$X_{rec} = \mathrm{D}(Z_X)$$

$$Y = \mathrm{D}(Z_Y)$$

$$d_{src}^X = \mathrm{Disc}_{src}(X)$$

$$D_{cls}^X = \mathrm{Disc}_{cls}(X)$$

$$d_{src}^Y = \text{Disc}_{src}(Y)$$

$$D_{cls}^Y = \text{Disc}_{cls}(Y)$$

where $Q$ is the one-hot matrix indicating the slice indices as stated above, $\text{NG}(\cdot)$ represents the noise generator, and $\parallel$ denotes concatenation. The noise generator output $N_X$, which contains the representations of noise and batch effects for spots, and is subsequently added to $Z$ to generate the latent representations of spots with batch effects, denoted as $Z_X$. $Z_X$ is then utilized to reconstruct the input data matrix via the decoder. Concurrently, through a pseudo-noise generation (PNG) process, we construct a pseudo-noise matrix, denoted as $N_Y$, which simulates the representations of noise and batch effects that could be added to $Z$ to enable spots to be mapped to the domains of other slices.

Specifically, during each training epoch in stage 2, each slice is assigned at random an index, representing the domain of the target slice to which the spots in the source slice (the original slice of spots) should be mapped during that epoch. Subsequently, for the $i$-th source spot in the $s$-th source slice, we find $k$ nearest neighbors within the target slice for the source spot using the representations in $Z$. It is worth noting that we select the nearest neighbors based on the magnitude of cosine similarity rather than Euclidean distance. This strategy has been proven to significantly enhance the integration effectiveness of INSTINCT (Supplementary Figs. 1 and 2). These neighbors are spots within the target slice that are most likely to possess similar biological characteristics as the source spot. We then average the representations of the noise and batch effects corresponding to these spots in $N_X$ to determine the noise and batch effects required for mapping the source spot to the domain of the target slice. By repeating this process for each spot, we obtain $N_Y$, where each row represents the pseudo-noise-and-batch-effects for the corresponding spot. $N_Y$ is then added to $Z$ to yield $Z_Y$, after which the same decoder is used to map each spot to its designated

target domain.

In stage 2, we task the discriminator with distinguishing between the real data matrix $X$ and the fake data matrix $Y$, while also discerning the categories of spots. Here, the categories of spots corresponding to $Y$ are defined by the target domains.

Compared to conventional autoencoders, the incorporation of the noise generator and discriminator modules, along with the utilization of the stochastic domain translation procedure, enables the low-dimensional representations of spots with added noise and batch effects to be seamlessly reconstructed back to the domain of their original slices or generated to any specified domain of another slice. This approach allows the noise generator to effectively capture noise and batch effect patterns across different slices, thereby facilitating the more efficient removal of batch effects from the low-dimensional representations while preserving as much biological variations as possible. After training in both stages, $Z$ can be extracted as the low-dimensional representations of spots without noise and batch effects, suitable for various applications and downstream analyses.

**Loss functions of INSTINCT**

We employed four types of loss functions for training the model, including adversarial loss, classification loss, reconstruction loss, and latent loss.

**Adversarial loss.** In INSTINCT, the generator (comprising the encoder and decoder for stage 1, and including the encoder, decoder, and noise generator for stage 2) attempts to produce realistic data, while the discriminator aims to distinguish between samples generated by the generator and the real data. The adversarial loss is employed to encourage the generator to produce data more closely resembling real samples, thereby enhancing its capability, while

also improving the precision of the discriminator. The form of the adversarial loss for the two stages are as following:

$$Stage\ 1: \mathcal{L}_{adv} = \mathbb{E}[\log \text{Disc}_{src}(x)] + \mathbb{E}\left[\log\left(1 - \log(\text{Disc}_{src}(x'))\right)\right]$$

$$Stage\ 2: \mathcal{L}_{adv} = \mathbb{E}[\log \text{Disc}_{src}(x)] + \mathbb{E}\left[\log\left(1 - \log(\text{Disc}_{src}(y))\right)\right]$$

where $x$, $x'$ and $y$ represent spot representation from $X$, $X_{rec}$ and $Y$, respectively. The discriminator maximizes this function to distinguish between real and fake data, while the generator minimizes it to confound the discriminator.

**Classification loss.** The classification loss consists of the following two components:

$$Stage\ 1\ \&\ 2: \mathcal{L}_{cls}^r = \mathbb{E}[-\log \text{Disc}_{cls}(c|x)]$$

$$Stage\ 1: \mathcal{L}_{cls}^f = \mathbb{E}[-\log \text{Disc}_{cls}(c|x')]$$

$$Stage\ 2: \mathcal{L}_{cls}^f = \mathbb{E}[-\log \text{Disc}_{cls}(c'|y)]$$

where $c$ represents the original category of the spot $x$ or $x'$, and $c'$ is the assigned target category correspond to $y$. We use $\mathcal{L}_{cls}^r$ to train the discriminator to distinguish spots from different slices, and $\mathcal{L}_{cls}^f$ to train the generator to accurately map spots to specified domains.

**Reconstruction loss.** The reconstruction loss is used to enhance the ability of the model to reconstruct the input data. By minimizing the reconstruction loss, the model learns effective data representations and reconstruction techniques, thereby improving the quality and accuracy of its generated data. We used mean square error (MSE) as the reconstruction loss:

$$\mathcal{L}_{rec} = \mathbb{E}[\|x - x'\|_2^2]$$

**Latent loss.** We developed a latent loss to better remove the batch effects between samples:

$$\mathcal{L}_{la} = \mathbb{E}\left[\frac{1}{k}\sum_{j=0}^{k} \max\left(clamp_{max}\left(\left\|z - z_{p_j}\right\|_2 - \left\|z - z_{n_j}\right\|_2, m\right) + \alpha, 0\right)\right]$$

where $z$ is the spot representation from $Z$. For each spot, we identify $k$ positive samples and $k$ negative samples. A positive sample, a negative sample, and the spot form a triplet. We use triplet loss[77] to encourage the spot to be close to its positive samples and far from negative samples in the latent space. The positive samples are the nearest neighbors selected for the spot during the PNG process, while the negative samples are randomly chosen from the same slice as the spot, following the manner of STAligner[28]. Here, $z_{p_j}$ and $z_{n_j}$ represent the $j$-th positive sample and the $j$-th negative sample found for $z$, respectively, and $\alpha$ denotes the hyperparameter of triplet loss.

However, we cannot guarantee that all positive samples found for a particular spot are the same type as that spot. In fact, sometimes, if a spot belongs to a type specific to a certain slice, or if the number of spots of that type in other slices is less than $k$, then some or all of the positive samples found for it will not be of the same type as the spot itself. Using triplet loss to encourage the spot to be close to these incorrect positive samples may result in the inability to identify rare or specific spot-types, which is undesirable. Additionally, since negative samples are randomly selected, it is impossible to avoid selecting samples of the same type as the spot. Therefore, we employ a clamp strategy, denoted as $clamp_{max}(\cdot)$. Triplets where the difference between the distance of the spot to the positive sample and the distance of the spot to the negative sample exceeds a hyperparameter $m$ are not included in the backpropagation process. This strategy helps in removing positive samples that are too distant from the spot (potentially incorrect positives) or negative samples that are too close to the spot (potentially incorrect negatives).

**Total objectives.** The total objectives for the generator and discriminator of the two stages are as displayed below:

$$Stage\ 1\ \&\ 2: \mathcal{L}_{Disc} = -\mathcal{L}_{adv} + \lambda_{cls}\mathcal{L}_{cls}^r$$

$$Stage\ 1: \mathcal{L}_G = \mathcal{L}_{adv} + \lambda_{cls}\mathcal{L}_{cls}^f + \lambda_{rec}\mathcal{L}_{rec}$$

$$Stage\ 2: \mathcal{L}_G = \mathcal{L}_{adv} + \lambda_{cls}\mathcal{L}_{cls}^f + \lambda_{rec}\mathcal{L}_{rec} + \lambda_{la}\mathcal{L}_{la}$$

where $\lambda_{cls}$, $\lambda_{rec}$ and $\lambda_{la}$ are hyperparameters which control the contribution of different loss functions to the total objectives.

For all experiments reported in this paper, we set $\lambda_{cls} = \lambda_{rec} = 10$, $\lambda_{la} = 20$, $k = 50$, $m = 10$, and $\alpha = 1$. The generator was optimized with a learning rate of 0.001, while the discriminator used 0.0005. The number of training epochs for both stages were set to 500.

## Simulations

We conducted extensive simulation experiments to validate the performance of INSTINCT. Inspired by SpatialPCA[43], a method developed for SRT, we adopted a similar approach to generate spATAC-seq datasets. The data information for different scenarios are summarized in Supplementary Tabs. 1 to 6.

Specifically, we utilized simCAS[44], a method for simulating scATAC-seq data, to generate multiple single-cell datasets with batch effects. For the reference data and parameters used in the simulation, we employed the discrete mode of simCAS, with the *tree_text* parameter set to "(((0:0.2, 1:0.2):0.2, 2:0.4):0.5, (3:0.5, 4:0.5):0.4);", and followed its online tutorial. We utilized the summary statistics of the Buenrostro2018 dataset[78] provided at https://github.com/Chen-Li-17/simCAS/tree/main/data as the training data. Besides, simCAS provides an optional step for adding batch effects to simulated data. As per simCAS, batch effects are categorized into those originating from technical variations and

those originating from biological factors. By adjusting the parameters for introducing these two types of batch effects, we produced three scATAC-seq datasets exhibiting varying degrees of batch effects. Each dataset comprises 10,000 cells distributed among five cell-types, with 2,000 cells for each type labeled as 0 to 4, respectively. The parameters for the batch effects added to these three datasets were as follows: for dataset 0, we set the mean of batch effects from biological factors to 0 with a standard deviation of 0.5. For dataset 1, we set the mean of batch effects from technical variations to 0 with a standard deviation of 0.1. For dataset 2, we set the mean of batch effects from technical variations to 0 with a standard deviation of 0.1 and the mean of batch effects from biological factors to 0 with a standard deviation of 0.5.

After that, we pre-defined a single-cell resolution slice of size $120 \times 150$ as a template, where each location corresponds to integer coordinates ranging from 0 to 149 for the x-axis and 0 to 119 for the y-axis. This template was divided into five domains represented by domain 0 to 4, respectively. Each domain contained a different number of locations: 2700, 4500, 3150, 3600, and 4050, each corresponding to a major cell-type with the same number as its index. Initially, each position was assigned a cell-type, where in a specific region, the cell-type corresponding to that region was assigned with a probability of $r$, while the remaining cell-types were each assigned with a probability of $\frac{1-r}{4}$. Subsequently, based on the assigned cell-types for each location, a cell corresponding to that cell-type was randomly selected from the scATAC-seq dataset and allocated to that position. By repeating this process for each simulated scATAC-seq dataset, we obtained three single-cell resolution slices. Each spot, defined as a $3 \times 3$ cell grid, aggregated data from the nine cells within it by summation, with the prevalent cell-type among the nine cells dictating the type of that spot. The coordinates of the spot were determined by the coordinates of the central cell. Thus, three

single-cell resolution slices were transformed into corresponding spatial resolution slices theoretically containing five spot-types, denoted as spot-type 0 to 4. We iteratively decreased the value of $r$, resulting in increasing levels of confusion among spot-types and progressively more complex batch effects across three scenarios, where scenarios 1, 2, and 3 corresponded to $r$ values of 0.8, 0.7, and 0.6, respectively. We point out that the value of $r$ should not be too small because it may lead to a discontinuous ground truth, resulting in pre-defined domains losing their practical significance.

In certain situations, vertically adjacent slices may not exhibit complete overlap, where different slices display varying proportions of spot-types, or a slice may lack some spot-types found in other slices. This issue can also arise in samples from different developmental stages. Therefore, based on scenario 1, we created scenarios 4 and 5 to simulate two cases where the proportions of spot-types differ between slices (Supplementary Fig. 8). In scenario 4, we applied different cropping to the three spatial resolution slices from scenario 1, preserving 80% of the spots on the right side, middle, and left side, respectively, to generate three new slices. In scenario 5, we retained slice 0 and slice 1 from scenario 1 and removed domain 2 from slice 2.

We simulated scenario 6 based on scenario 1 to test the ability of INSTINCT to preserve rare and specific spot-types. We excluded domain 1 from slices 1 and 2 of scenario 1, retaining a $5 \times 5$ area of domain 1 in slice 0 and removing the rest (Supplementary Fig. 8). Only about 25 spots of spot-type 1 were kept across the three slices, mostly concentrated in slice 0. It is worth noting that, due to the specific way we determined spot-types, there are several spots belonging to spot-type 1 in other domains.

**Spatial domain identification**

For datasets with ground truth annotations, we initially applied a Gaussian mixture model-based clustering algorithm implemented in the *scikit-learn* (v1.4.0) package[79], setting the number of components to the number of spot-types, following the approach of STitch3D[31], to cluster the latent representations of spots in $Z$. Subsequently, we utilized the *match_cluster_labels* function provided in the online tutorial of PASTE[80] to map the cluster labels to the ground truth labels. For datasets without annotations, we applied the Leiden algorithm[81] implemented in the *Scanpy* (v1.9.8) package[82] with the default settings to cluster the latent representations of spots.

**Visualization**

We used the UMAP algorithm[46] to further reduce the dimensionality of the spot representations in $Z$ and visualized them in a two-dimensional space. Following Portal[83], the algorithm was run with parameters set to 30 nearest neighbors, a minimum distance of 0.3 and utilizing the correlation metric. For visualizing the raw data, we initially preprocessed the data following the procedure outlined above. Subsequently, we employed UMAP algorithm with the same settings to reduce the dimensionality of the spot representations in $X_{pca}$.

**Spot-type annotation**

We performed spot-type annotation by employing a k-nearest-neighbor (KNN) classifier, a model-free approach whose annotation efficacy heavily depends on the effectiveness of the low-dimensional representations across samples. For each spot in the target sample to be annotated, we selected a specified number of nearest neighbors from the annotated source

sample. The label assigned to each spot was determined by the majority class among its

nearest neighbors. We set the number of nearest neighbors to 20, consistent with the approach

outlined in Portal[83].

**Gene ontology (GO) analysis**

Initially, we conducted differential gene expression analysis using the *rank_genes_groups*

function in the *Scanpy* package with the Wilcoxon test as the method. A gene was considered

differentially expressed if it exhibited a log2-fold change greater than 0.2 and an adjusted *p*-

value (*q*-value) less than 0.05. Subsequently, the differentially expressed genes (DEGs) were

utilized for performing GO analysis via the algorithm provided in the *gseapy* (v1.1.2)

package[84], employing *GO_Biological_Process_2021* as the reference database.

**Determination of spot-type-specific peaks**

We first filtered the peaks according to the preprocessing procedure and applied TF-IDF

transformation to the sample. Then, we utilized the *rank_genes_groups* function in the

*Scanpy* package with the Wilcoxon test as the method. For each spot-type, we filtered out

peaks those exhibited log2-fold change no more than 0.2, and selected 300 peaks with the

lowest adjusted *p*-value (*q*-value) as its specific peaks. For spot-types whose peaks were no

more than 300 after filtering, we kept all peaks as their specific peaks. Peaks not chosen as

specific peaks were retained as background peaks.

**Expression enrichment analysis**

Following EpiAnno[68], we initially mapped the genome coordinates of the spot-type-specific

peaks and background peaks to GRCh37/hg19 using LiftOver[85]. Subsequently, we conducted

expression enrichment analysis utilizing the *SNPsea* (v1.0.3) package[86] with default settings for SNPs in each set of spot-type-specific peaks and the set of background peaks, respectively.

**Partitioned heritability analysis**

We first mapped the genome coordinates of the spot-type-specific peaks and background peaks to GRCh37/hg19. Then, we used *LDSC* (v1.0.1) package[87] to quantified the enrichment of heritability for phenotypes related to brain. The European samples from the 1000 Genomes Project served as the reference panel, and the summary statistics for GWAS were all obtained from https://doi.org/10.5281/zenodo.7768714.

**Data collection and description**

The MISAR-seq MB dataset[15] comprises eight spATAC-seq samples and their corresponding SRT profiles from E11.0, E13.5, E15.5 and E18.5 developmental stages, with two vertically adjacent slices from each stage. The eight spATAC-seq samples include E11_0-S1_atac (1,258 spots and 242,024 peaks), E13_5-S1_atac (1,777 spots and 271,126 peaks), E15_5-S1_atac (1,949 spots and 265,014 peaks), E18_5-S1_atac (2,129 spots and 294,734 peaks), E11_0-S2_atac (1,353 spots and 213,562 peaks), E13_5-S2_atac (2,183 spots and 251,555 peaks), E15_5-S2_atac (1,939 spots and 244,394 peaks), and E18_5-S2_atac (2,248 spots and 223,333 peaks). The eight corresponding SRT samples include E11_0-S1_expr (1,263 spots and 32,285 genes), E13_5-S1_expr (1,777 spots and 32,285 genes), E15_5-S1_expr (1,949 spots and 32,285 genes), E18_5-S1_expr (2,129 spots and 32,285 genes), E11_0-S2_expr (1,353 spots and 32,285 genes), E13_5-S2_expr (2,183 spots and 32,285 genes), E15_5-S2_expr (1,939 spots and 32,285 genes), and E18_5-S2_expr (2,248 spots and 32,285 genes).

The spatial ATAC-RNA-seq MB[16] dataset includes three spATAC-seq samples of mouse postnatal day 21/22 (P21/22), each paired with an SRT sample. The three spATAC-seq samples are GSM6204623 (2,500 spots and 233,155 peaks), GSM6758284 (2,500 spots and 290,047 peaks), and GSM6758285 (9,994 spots and 517,411 peaks). The three SRT samples are GSM6204636 (2,373 spots and 19,859 genes), GSM6753041 (2,498 spots and 20,046 genes), and GSM6753043 (9,215 spots and 22,914 genes). GSM6204623 is paired with GSM6204636, GSM6758284 is paired with GSM6753041, and GSM6758285 is paired with GSM6753043.

The spatial ATAC ME dataset[12] consists of six spATAC-seq samples from E12.5, E13.5, and E15.5 developmental stages. The six samples comprise E12_5-S1 (2,234 spots), E12_5-S2 (2,246 spots), E13_5-S1 (2,970 spots), E13_5-S2 (3,160 spots), E15_5-S1 (3,903 spots), and E15_5-S2 (3,274 spots). The peaks were merged in the original study[12], resulting in 248,047 peaks for each sample.

**Baseline methods**

We compared INSTINCT with six baseline methods: Scanorama[21], SCALEX[25], PeakVI[26], SEDR[27], STAligner[28] and GraphST[29]. Among these, Scanorama, SCALEX, and PeakVI are originally developed for integrating single-cell omics data, while SCALEX and PeakVI are specifically designed for scATAC-seq data. On the other hand, SEDR, STAligner and GraphST are developed for integrating SRT data.

For each method, we conducted eight iterations by varying the random seed. Data was filtered before used. All parameters were kept as their default settings, and we followed the tutorial provided by each method to conduct the experiments.

**Evaluation metrics**

We compared the capabilities of the integration methods in three aspects: clustering performance, representation quality, and batch correction. We regarded clustering performance and representation quality as two separate aspects of biological conservation, noting that representation quality depends solely on the model, whereas clustering performance also relies on the clustering algorithm. We assessed clustering performance using six metrics: Adjusted Rand Index (ARI), Adjusted Mutual Information (AMI), Normalized Mutual Information (NMI), Fowlkes-Mallows Index (FMI), completeness score (Comp), and homogeneity score (Homo), leveraging functions from the *scikit-learn* package. For evaluating representation quality, we employed mean average precision (mAP)[88], average silhouette width across spot-types (spot-type ASW), average silhouette width across isolated labels (isolated label ASW), and isolated label F1-score, utilizing the *scib* (v1.1.4) package[45]. Regarding batch correction assessment, we computed average silhouette width across batches (batch ASW), principal component regression across batches (batch PCR), k-nearest-neighbor batch effect test (kBET), and graph connectivity using the *scib* package. The scores of each group were averaged to obtain the corresponding group-specific overall score. The group-specific overall scores for the three aspects were further averaged to obtain the final overall score.

For the evaluation of spot-type annotation, we calculated accuracy, kappa, macro F1-score (mF1), and weighted F1-score (wF1) using k-fold cross-validation, with each fold corresponding to one slice.

**ARI.** ARI standardizes the Rand Index (RI) to quantify the similarity between two sets of labels. The higher the ARI value, the more similar the two sets of clusters are. The formula

is as follows:

$$\text{ARI} = \frac{\text{RI}(L_T, L_C) - \mathbb{E}[\text{RI}(L_T, L_C)]}{\max[\text{RI}(L_T, L_C)] - \mathbb{E}[\text{RI}(L_T, L_C)]}$$

where $L_T$ and $L_C$ are true labels and clustering labels, respectively. $\text{RI}(\cdot)$ denotes the function of RI, and $\mathbb{E}[\text{RI}]$ represents the expected RI under random assignment.

**AMI.** AMI is a measure used to compare the similarity between two sets of labels by adjusting the Mutual Information (MI) score, and is calculated as follows:

$$\text{AMI} = \frac{\text{MI}(L_T, L_C) - \mathbb{E}[\text{MI}(L_T, L_C)]}{\text{avg}[\text{H}(L_T), \text{H}(L_C)] - \mathbb{E}[\text{MI}(L_T, L_C)]}$$

where $\text{MI}(\cdot)$ denotes the function of MI and $\text{H}(\cdot)$ represents the function of entropy.

**NMI.** NMI calculates the normalization of MI score. Higher NMI score represents better agreement between the two sets of clusters. The formula is as follows:

$$\text{NMI} = \frac{\text{MI}(L_T, L_C)}{\sqrt{\text{H}(L_T)\text{H}(L_C)}}$$

**FMI.** FMI focuses on the precision and recall of the clustering results, with a higher score indicating better clustering performance. It is calculated as follows:

$$\text{FMI} = \frac{TP}{\sqrt{(TP + FP)(TP + FN)}}$$

where $TP$ is the number of pairs of elements that are in the same cluster in both the true labels and the clustering labels, $FP$ is the number of pairs of elements that are in the same cluster of the true labels but in different clusters of the clustering labels, and $FN$ is the number of pairs of elements that are in different clusters of the true labels but in the same clusters of the clustering labels.

**Comp.** Completeness score evaluates whether all members of a specific class of the true labels are grouped into the same cluster and is calculated as follows:

$$\text{Comp} = 1 - \frac{H(L_C|L_T)}{H(L_C)}$$

**Homo.** Homogeneity score evaluates whether a single cluster contains only data samples that are members of a single ground truth class. The formula is as follows:

$$\text{Homo} = 1 - \frac{H(L_T|L_C)}{H(L_T)}$$

**mAP.** mAP quantifies how well the labels of the nearest neighbors of each spot match its own label, with a higher value indicating the better preservation of the consistency. For the calculation methods, we followed its original work[88]. The number of the nearest neighbors considered was set to 30.

**Spot-type ASW.** Spot-type ASW is a measure of how similar a sample is to its own class compared to other classes, and is calculated with respect to spot-type labels. We used the function implemented in the *scib* package[45] and kept its default settings.

**Isolated label ASW.** Isolated label ASW evaluates the conservation of spot-types unique to individual samples by calculating the ASW with respect to these labels. We computed it with the function implemented in the *scib* package, keeping its default settings.

**Isolated label F1-score.** Isolated label F1-score is also a metric developed to assess the preservation of the spot-types unique to individual samples. It calculates the F1-scores[89] for

these isolated labels and takes the mean. We used the function implemented in the *scib* package and kept its default settings.

**Batch ASW.** Batch ASW is a measure of how similar a sample is to its own slice compared to other slices, and is calculated with respect to slice indices. The score is scaled so that a higher value indicates a better mixing of data from different slices. We computed it with the function implemented in the *scib* package, keeping its default settings.

**Batch PCR.** Batch PCR evaluates the effectiveness of batch correction by comparing the variance before and after integration. We used the function implemented in the *scib* package and kept its default settings.

**kBET.** kBET measures the similarity of spot-type distributions between slices. It evaluates batch mixing by comparing local batch distribution to global batch distribution. It is computed by leveraging the function implemented in the *scib* package and the default settings were kept.

**Graph connectivity.** Graph connectivity quantifies the connectivity of the subgraph per spot-type label and takes the average. We computed it with the function implemented in the *scib* package.

**Accuracy.** Accuracy is the proportion of correctly annotated spots to the total number of spots and is calculated as follows:

$$\text{Accuracy} = \frac{TP + TN}{TP + TN + FP + FN}$$

where $TP$, $FP$, $TN$ and $FN$ denote the number true positive samples, false positive samples, true negative samples, and false negative samples, respectively.

**Kappa.** Kappa measures inter-rater agreement for categorical items. It compares the observed agreement between two raters to the agreement that would be expected by chance. The formula is as follows:

$$\text{Kappa} = \frac{p_a - p_e}{1 - p_e}$$

where $p_a$ represents the proportion of observed agreement between two raters, and $p_e$ is the proportion of agreement expected if the raters are to rate completely randomly.

**mF1.** mF1 takes the mean of the F1-scores[89] calculated for all spot-types. The formula is as follows:

$$\text{mF1} = \frac{1}{n} \sum_{i=1}^{n} \text{F1}_i$$

where $n$ is the number of spot-types and $\text{F1}_i$ represents the F1-score of the $i$-th spot-type.

**wF1.** wF1 calculates the weighted average of F1-scores for all spot-types, where the weight of each F1-score is the proportion of the number of spots for that type relative to the total number of spots. It is computed as follows:

$$\text{wF1} = \sum_{i=0}^{n} \frac{N_i}{N} \text{F1}_i$$

where $N_i$ denotes the number of spots of the $i$-th spot-type, and $N$ is the total number of spots.

## Data availability

The MISAR-seq MB dataset[15] can be downloaded from the National Genomics Data Center with accession number OEP003285. The spatial ATAC-RNA-seq MB dataset[16] is available at the Gene Expression Omnibus with accession code GSE205055. The spatial ATAC ME dataset[12] can be obtained at Gene Expression Omnibus using accession code GSE214991.

## Code availability

INSTINCT and all the code for reproducing the analyses and benchmarking can be found at https://github.com/yyLIU12138/INSTINCT.

## Acknowledgements

This work was supported by the National Key Research and Development Program of China (grant nos. 2023YFF1204802 and 2021YFF1200902) and the National Natural Science Foundation of China (grant nos. 62273194).

## Author contributions

R.J. conceived the study and supervised the project. Y.L., Z.L. designed and implemented and validated INSTINCT. X.Y.C., X.J.C. and Z.G. helped with analyzing the results. Y.L., Z.L. and R.J. wrote the manuscript, with input from all the authors.

## Competing interests

The authors declare no competing interests.

# Figures

1097

**Fig. 1 | The overview of INSTINCT. a,** the model structure of INSTINCT. INSTINCT adopts a two-stage strategy. In stage 1, the model of INSTINCT consists of a GAT encoder, a MLP decoder, and a discriminator. Taking the preprocessed data matrix of multiple samples and the concatenated neighbor graph as input, it simply learns to project all the spots from feature space (PCs space) into a low-dimensional shared latent space, while the discriminator is pretrained. Noise and batch effects still persist in the latent representations of spots after the training in stage 1. During stage 2, a noise generator is added to the model. After the extraction of biological variations performed by the encoder, the noise generator captures the original pattern of batch effects of each spot in the latent space, and a pseudo-noise generation (PNG) process simulates a different pattern of batch effects for each spot. These two simulated batch effects are then added to the corresponding spot separately, enabling reconstruction back to its origin domain or conversion to a new domain, referred to as the stochastic domain translation process. After training, the latent representation matrix $Z$ can be extracted and utilized for various applications and downstream analyses. **b,** the sketch figure of the stochastic domain translation procedure, using the integration process of two samples as an illustration. The GAT encoder, MLP decoder, and discriminator respectively implement the processes of biological variations extraction, reconstruction or generation, as well as discrimination and classification. The batch effects addition procedure is completed by the noise generator and the PNG process. Specifically, the encoder extracts the biological variations within the data and projects it to the latent space. Subsequently, two different patterns of batch effects are added to the low-dimensional representation of each spot, enabling reconstruction to its original domain or generation to the target domain through the decoder. The raw data and the generated data are then discriminated and classified. **c,** the applications of the integration results of INSTINCT.

1116
**Fig. 2 | The simulation procedure and the performance of INSTINCT in simulated scenarios. a,** the predefined template
comprises five spatial domains, named by integers 0 to 4, each possesses a major cell-type within cell-type 0 to 4. In each domain,
the corresponding major cell-type is assigned to each location with a probability of $r$, and other cell-types are each assigned with
a probability of $\frac{1-r}{4}$. Three simulated single-cell datasets with batch effects are generated by simCAS, each used to generate one

spatial slice. Initially, each location in all slice is assigned with a cell-type, and a cell belongs to this cell-type in the corresponding
single-cell dataset is randomly selected and allocated to the location to form three cell-level slices with single-cell resolution.
Subsequently, with $3 \times 3$ region defined as a spot, the three slices are merged into spot-level slices, respectively, to simulate the
resolution of spATAC-seq techniques. The category of each spot is determined by the cell-type that predominantly occupied that
spot. Evident batch effects between simulated slices can be observed. **b,** the overview table of metrics calculated in scenario 1
demonstrates the superior performance of INSTINCT. INSTINCT reaches the highest score in 9 of the 12 metrics, and ranks
second in 2 metrics. It also ranks first on the group-specific overall scores of clustering performance and representation quality,
as well as the final overall score, and ranks second on the overall score of batch correction. **c,** the probability $r$ is reduced
continuously across the first three scenarios, resulting in increasing levels of confusion among spot-types and progressively more
complex batch effects. INSTINCT consistently outperforms the benchmarking methods. **d,** two different transformations are
applied to the data of scenario 1 to generate scenario 4 and 5, in order to simulate scenarios where the proportions of spot-types
differ between slices. INSTINCT results in satisfying integration outcomes. **e,** based on scenario 1, scenario 6 is generated to test
the ability of INSTINCT to preserve rare spot-types. INSTINCT succeed in preserving the rare spot-type which mainly exists in
the first slice.
1135

1136

1137 **Fig. 3 | The comparison of evaluation metrics, visualizations and clustering results on the MISAR-seq MB dataset. a,** the

1138 individual scores of each evaluation metric for INSTINCT and benchmarking methods. Two new metrics assessing the

preservation of sample-specific spot-types, isolated label ASW and isolated label F1-score, are added to the group of representation quality. INSTINCT ranks first in 10 out of 14 evaluation metrics and second in one. **b,** the group-specific overall scores and the final overall score. INSTINCT ranks first in clustering performance, representation quality, as well as the final overall score, and second in batch correction. **c,** UMAP visualization of data from different developmental stages before integration. **d,** UMAP visualization of data from different developmental stages after integration performed by INSTINCT. **e,** UMAP visualization of integration results, with spots colored according to ground truth annotations. Results of the four methods that rank in the top for the final overall score are displayed. **f,** the ground truth annotations of different slices. **g,** the clustering results of the four methods that rank in the top for the final overall score, with identified clusters first mapped to true labels and then colored accordingly.

1148
**Fig. 4 | Integration results of INSTINCT on samples with different tissue scales from the spatial ATAC-RNA-seq MB**
**dataset. a,** Nissl (left) and anatomical annotations (right) from the Allen Mouse Brain Atlas and Allen Reference Atlas – Mouse
Brain. Allen Mouse Brain Atlas, mouse.brain-map.org and atlas.brain-map.org. **b,** UMAP visualizations of the integration outcome
and clustering results of INSTINCT. Biological variations of different spot-types are highlighted and 13 clusters are identified
through clustering. **c,** spots are colored according to their specific clusters and displayed by their spatial arrangement. The clusters
exhibit strong continuity in space, indicating the identification of spatial domains. **d,** Visualizations of DEGs for identified clusters
based on their expression levels.
1156

1157

**Fig. 5 | The comparison of evaluation metrics, visualizations and clustering results on the spatial ATAC ME dataset. a,** the

group-specific overall scores and the final overall score. INSTINCT ranks first in clustering performance, representation quality,

as well as the final overall score. **b,** UMAP visualization of integration results, with spots colored according to slice affiliation.

Results of the three methods that rank in the top for the final overall score are displayed. **c,** UMAP visualization of integration

results, with spots colored according to the ground truth annotations. **d,** the ground truth annotations of different slices. **e,** the

clustering results of the three methods that rank in the top for the final overall score, with identified clusters first mapped to true

labels and then colored accordingly.

1165

1166
1167 **Fig. 6 | Annotation results and downstream analyses on the MISAR-seq MB dataset. a,** comparison on four metrics: accuracy,
1168 kappa, mF1, and wF1, for evaluating annotation results using cross-validation. INSTINCT consistently ranks first on all the

metrics. **b,** the ground truth of the S1 slice of developmental stage E18.5 and the annotation outcome of the corresponding S2 slice based on the integration results of INSTINCT. **c,** GO analysis performed for the two rare spot-types, subpallium_2 and muscle, identified in S2 slice. **d,** results of expression enrichment analysis on the spot-type-specific peaks of diencephalon_and_hindbrain and background peaks. The top 30 significantly enriched tissues are displayed. **e,** the results of partitioned heritability analysis performed on the spot-type-specific peaks and background peaks for three diseases, including ADHD, schizophrenia, and autism.

# References

1. Tsompana, M. & Buck, M. J. Chromatin accessibility: a window into the genome. *Epigenetics Chromatin* **7**, 1-16 (2014).

2. Klemm, S. L., Shipony, Z. & Greenleaf, W. J. Chromatin accessibility and the regulatory epigenome. *Nat. Rev. Genet.* **20**, 207-220 (2019).

3. Buenrostro, J. D. et al. Single-cell chromatin accessibility reveals principles of regulatory variation. *Nature* **523**, 486-490 (2015).

4. Rotem, A. et al. Single-cell ChIP-seq reveals cell subpopulations defined by chromatin state. *Nat. Biotechnol.* **33**, 1165-1172 (2015).

5. Jin, W. et al. Genome-wide detection of DNase I hypersensitive sites in single cells and FFPE tissue samples. *Nature* **528**, 142-146 (2015).

6. Lareau, C. A. et al. Droplet-based combinatorial indexing for massive-scale single-cell chromatin accessibility. *Nat. Biotechnol.* **37**, 916-924 (2019).

7. Cusanovich, D. A. et al. Multiplex single-cell profiling of chromatin accessibility by combinatorial cellular indexing. *Science* **348**, 910-914 (2015).

8. Sinha, S. et al. Profiling chromatin accessibility at single-cell resolution. *Genomics Proteomics Bioinformatics* **19**, 172-190 (2021).

9. Bressan, D., Battistoni, G. & Hannon, G. J. The dawn of spatial omics. *Science* **381**, eabq4964 (2023).

10. Browaeys, R., Saelens, W. & Saeys, Y. NicheNet: modeling intercellular communication by linking ligands to target genes. *Nat. Methods* **17**, 159-162 (2020).

11. Brücher, B. L. & Jamall, I. S. Cell-cell communication in the tumor microenvironment, carcinogenesis, and anticancer treatment. *Cell. Physiol. Biochem.* **34**, 213-243 (2014).

12. Llorens-Bobadilla, E. et al. Solid-phase capture and profiling of open chromatin by spatial ATAC. *Nat. Biotechnol.* **41**, 1085-1088 (2023).

13. Deng, Y. et al. Spatial profiling of chromatin accessibility in mouse and human tissues. *Nature* **609**, 375-383 (2022).

14. Xie, Y. et al. Spatial chromatin accessibility sequencing resolves high-order spatial interactions of epigenomic markers. *Elife* **12**, RP87868 (2024).

15. Jiang, F. et al. Simultaneous profiling of spatial gene expression and chromatin accessibility during mouse brain development. *Nat. Methods* **20**, 1048-1057 (2023).

16. Zhang, D. et al. Spatial epigenome–transcriptome co-profiling of mammalian tissues. *Nature* **616**, 113-122 (2023).

17. Argelaguet, R. et al. Computational principles and challenges in single-cell data integration. *Nat. Biotechnol.* **39**, 1202-1215 (2021).

18. Butler, A. et al. Integrating single-cell transcriptomic data across different conditions, technologies, and species. *Nat. Biotechnol.* **36**, 411-420 (2018).

19. Polański, K. et al. BBKNN: fast batch alignment of single cell transcriptomes. *Bioinformatics* **36**, 964-965 (2020).

20. Korsunsky, I. et al. Fast, sensitive and accurate integration of single-cell data with Harmony. *Nat. Methods* **16**, 1289-1296 (2019).

21. Hie, B., Bryson, B. & Berger, B. Efficient integration of heterogeneous single-cell transcriptomes using Scanorama. *Nat. Biotechnol.* **37**, 685-691 (2019).

22. Dries, R. et al. Advances in spatial transcriptomic data analysis. *Genome Res.* **31**, 1706-1718 (2021).

23. Liu, Y. & Yang, C. Computational methods for alignment and integration of spatially resolved transcriptomics data. *Comput. Struct. Biotechnol. J.* **23**, 1094-1105 (2024).

24. Velten, B. & Stegle, O. Principles and challenges of modeling temporal and spatial omics data. *Nat. Methods* **20**, 1462-1474 (2023).

25. Xiong, L. et al. Online single-cell data integration through projecting heterogeneous datasets into a common cell-embedding space. *Nat. Commun.* **13**, 6118 (2022).

26. Ashuach, T. et al. PeakVI: A deep generative model for single-cell chromatin accessibility analysis. *Cell Rep.*

*Methods* **2**, 100182 (2022).

27.   Xu, H. et al. Unsupervised spatially embedded deep representation of spatial transcriptomics. *Genome Med.* **16**,
12 (2024).

28.   Zhou, X., Dong, K. & Zhang, S. Integrating spatial transcriptomics data across different conditions, technologies
and developmental stages. *Nat. Comput. Sci.* **3**, 894-906 (2023).

29.   Long, Y. et al. Spatially informed clustering, integration, and deconvolution of spatial transcriptomics with GraphST.
*Nat. Commun.* **14**, 1155 (2023).

30.   Guo, T. et al. SPIRAL: integrating and aligning spatially resolved transcriptomics data across different experiments,
conditions, and technologies. *Genome Biol.* **24**, 241 (2023).

31.   Wang, G. et al. Construction of a 3D whole organism spatial atlas by joint modelling of multiple slices with deep
neural networks. *Nat. Mach. Intell.* **5**, 1200-1213 (2023).

32.   Kingma, D. P. & Welling, M. Auto-encoding variational bayes. Preprint at https://arxiv.org/abs/1312.6114 (2013).

33.   Kipf, T. N. & Welling, M. Semi-supervised classification with graph convolutional networks. Preprint at
https://arxiv.org/abs/1609.02907 (2016).

34.   Velickovic, P. et al. Graph attention networks. *Stat* **1050**, 10-48550 (2017).

35.   Rao, A. et al. Exploring tissue architecture using spatial transcriptomics. *Nature* **596**, 211-220 (2021).

36.   Chen, H. et al. Assessment of computational methods for the analysis of single-cell ATAC-seq data. *Genome Biol.*
**20**, 1-25 (2019).

37.   Xiong, L. et al. SCALE method for single-cell ATAC-seq analysis via latent feature extraction. *Nat. Commun.* **10**,
4576 (2019).

38.   Stuart, T. et al. Comprehensive integration of single-cell data. *Cell* **177**, 1888-1902. e1821 (2019).

39.   Choi, Y. et al. Stargan: Unified generative adversarial networks for multi-domain image-to-image translation. In:
*Proceedings of the IEEE conference on computer vision and pattern recognition*) (2018).

40.   Choi, Y. et al. Stargan v2: Diverse image synthesis for multiple domains. In: *Proceedings of the IEEE/CVF*
*conference on computer vision and pattern recognition*) (2020).

41.   Romero, A. et al. Smit: Stochastic multi-label image-to-image translation. In: *Proceedings of the IEEE/CVF*
*International Conference on Computer Vision Workshops*) (2019).

42.   Popescu, M.-C. et al. Multilayer perceptron and neural networks. *WSEAS Trans. Circuits Syst.* **8**, 579-588 (2009).

43.   Shang, L. & Zhou, X. Spatially aware dimension reduction for spatial transcriptomics. *Nat. Commun.* **13**, 7203
(2022).

44.   Li, C. et al. simCAS: an embedding-based method for simulating single-cell chromatin accessibility sequencing
data. *Bioinformatics* **39**, btad453 (2023).

45.   Luecken, M. D. et al. Benchmarking atlas-level data integration in single-cell genomics. *Nat. Methods* **19**, 41-50
(2022).

46.   McInnes, L., Healy, J. & Melville, J. Umap: Uniform manifold approximation and projection for dimension reduction.
Preprint at https://arxiv.org/abs/1802.03426 (2018).

47.   Sunkin, S. M. et al. Allen Brain Atlas: an integrated spatio-temporal portal for exploring the central nervous system.
*Nucleic Acids Res.* **41**, D996-D1008 (2012).

48.   Thomas, E. A., Danielson, P. E. & Sutcliffe, J. G. RGS9: a regulator of G-protein signalling with specific expression
in rat and mouse striatum. *J. Neurosci. Res.* **52**, 118-124 (1998).

49.   Bonate, R. et al. Phosphodiesterase 10A (PDE10A): Regulator of dopamine agonist-induced gene expression in
the striatum. *Cells* **11**, 2214 (2022).

50.   Danielson, P. E. et al. Chromosomal mapping of mouse genes expressed selectively within the central nervous
system. *Genomics* **19**, 454-461 (1994).

51.   Desplats, P. A., Lambert, J. R. & Thomas, E. A. Functional roles for the striatal-enriched transcription factor, Bcl11b,
in the control of striatal gene expression and transcriptional dysregulation in Huntington's disease. *Neurobiol.*
*Dis.* **31**, 298-308 (2008).

52.   Tamura, S. et al. Foxp1 gene expression in projection neurons of the mouse striatum. *Neuroscience* **124**, 261-

267 (2004).

53.    Cheung, M. et al. Roles of Sox4 in central nervous system development. *Mol. Brain Res.* **79**, 180-191 (2000).

54.    Bulfone, A. et al. Spatially restricted expression of Dlx-1, Dlx-2 (Tes-1), Gbx-2, and Wnt-3 in the embryonic day
1274           12.5 mouse forebrain defines potential transverse and longitudinal segmental boundaries. *J. Neurosci.* **13**, 3155-
1275           3172 (1993).

1276    55.    Doeppner, T. R. et al. Zbtb20 regulates developmental neurogenesis in the olfactory bulb and gliogenesis after
1277           adult brain injury. *Mol. Neurobiol.* **56**, 567-582 (2019).

1278    56.    Anderson, C. et al. PLP1 and CNTN1 gene variation modulates the microstructure of human white matter in the
1279           corpus callosum. *Brain Struct. Funct.* **223**, 3875-3887 (2018).

1280    57.    Vincze, A. et al. A correlative light and electron microscopic study of postnatal myelination in the murine corpus
1281           callosum. *Int. J. Dev. Neurosci.* **26**, 575-584 (2008).

1282    58.    Wang, H.-F. & Liu, F.-C. Developmental restriction of the LIM homeodomain transcription factor Islet-1
1283           expression to cholinergic neurons in the rat striatum. *Neuroscience* **103**, 999-1016 (2001).

1284    59.    Stanley, G. et al. Continuous and discrete neuron types of the adult murine striatum. *Neuron* **105**, 688-699. e688
1285           (2020).

1286    60.    Phillips III, R. A. et al. Transcriptomic characterization of human lateral septum neurons reveals conserved and
1287           divergent    marker    genes    across    species.    Preprint    at
1288           https://www.biorxiv.org/content/10.1101/2024.04.22.590602v1.abstract (2024).

1289    61.    Leifer, D. et al. MEF2C, a MADS/MEF2-family transcription factor expressed in a laminar distribution in cerebral
1290           cortex. *Proc. Natl. Acad. Sci. U.S.A.* **90**, 1546-1550 (1993).

1291    62.    Maćkiewicz, A. & Ratajczak, W. Principal components analysis (PCA). *Comput. Geosci.* **19**, 303-342 (1993).

1292    63.    Wang, X.-K. et al. Gray matter network associated with attention in children with attention deficit hyperactivity
1293           disorder. *Front. Psychiatry.* **13**, 922720 (2022).

1294    64.    Andreasen, N. C. The role of the thalamus in schizophrenia. *Can. J. Psychiatry.* **42**, 27-33 (1997).

1295    65.    Byne, W. et al. The thalamus and schizophrenia: current status of research. *Acta Neuropathol.* **117**, 347-368 (2009).

1296    66.    Qiu, S. et al. Circuit-specific intracortical hyperconnectivity in mice with deletion of the autism-associated Met
1297           receptor tyrosine kinase. *J. Neurosci.* **31**, 5855-5864 (2011).

1298    67.    Li, Z. et al. Latent feature extraction with a prior-based self-attention framework for spatial transcriptomics.
1299           *Genome Res.* **33**, 1757-1773 (2023).

1300    68.    Chen, X. et al. Cell type annotation of single-cell chromatin accessibility data via supervised Bayesian embedding.
1301           *Nat. Mach. Intell.* **4**, 116-126 (2022).

1302    69.    Stuart, T. et al. Single-cell chromatin state analysis with Signac. *Nat. Methods* **18**, 1333-1341 (2021).

1303    70.    Martens, L. D. et al. Modeling fragment counts improves single-cell ATAC-seq analysis. *Nat. Methods* **21**, 28-31
1304           (2024).

1305    71.    Danese, A. et al. EpiScanpy: integrated single-cell epigenomic analysis. *Nat. Commun.* **12**, 5228 (2021).

1306    72.    Li, Z. et al. Chromatin-accessibility estimation from single-cell ATAC-seq data with scOpen. *Nat. Commun.* **12**,
1307           6386 (2021).

1308    73.    Liu, Q. et al. Simultaneous deep generative modelling and clustering of single-cell genomic data. *Nat. Mach. Intell.*
1309           **3**, 536-544 (2021).

1310    74.    Chen, S. et al. RA3 is a reference-guided approach for epigenetic characterization of single cells. *Nat. Commun.*
1311           **12**, 2177 (2021).

1312    75.    Lowd, D. & Meek, C. Adversarial learning. In: *Proceedings of the eleventh ACM SIGKDD international conference*
1313           *on Knowledge discovery in data mining*) (2005).

1314    76.    Clevert, D.-A., Unterthiner, T. & Hochreiter, S. Fast and accurate deep network learning by exponential linear units
1315           (elus). Preprint at https://arxiv.org/abs/1511.07289 (2015).

1316    77.    Schroff, F., Kalenichenko, D. & Philbin, J. Facenet: A unified embedding for face recognition and clustering. In:
1317           *Proceedings of the IEEE conference on computer vision and pattern recognition*) (2015).

78.    Buenrostro, J. D. et al. Integrated single-cell analysis maps the continuous regulatory landscape of human

hematopoietic differentiation. *Cell* **173**, 1535-1548. e1516 (2018).

79.    Pedregosa, F. et al. Scikit-learn: Machine learning in Python. *J. Mach. Learn. Res.* **12**, 2825-2830 (2011).

80.    Zeira, R. et al. Alignment and integration of spatial transcriptomics data. *Nat. Methods* **19**, 567-575 (2022).

81.    Traag, V. A., Waltman, L. & Van Eck, N. J. From Louvain to Leiden: guaranteeing well-connected communities. *Sci.*
*Rep.* **9**, 5233 (2019).

82.    Wolf, F. A., Angerer, P. & Theis, F. J. SCANPY: large-scale single-cell gene expression data analysis. *Genome Biol.*
**19**, 1-5 (2018).

83.    Zhao, J. et al. Adversarial domain translation networks for integrating large-scale atlas-level single-cell datasets.
*Nat. Comput. Sci.* **2**, 317-330 (2022).

84.    Fang, Z., Liu, X. & Peltz, G. GSEApy: a comprehensive package for performing gene set enrichment analysis in
Python. *Bioinformatics* **39**, btac757 (2023).

85.    Raney, B. J. et al. The UCSC Genome Browser database: 2024 update. *Nucleic Acids Res.* **52**, D1082-D1088 (2024).

86.    Slowikowski, K., Hu, X. & Raychaudhuri, S. SNPsea: an algorithm to identify cell types, tissues and pathways
affected by risk loci. *Bioinformatics* **30**, 2496-2497 (2014).

87.    Finucane, H. K. et al. Partitioning heritability by functional annotation using genome-wide association summary
statistics. *Nat. Genet.* **47**, 1228-1235 (2015).

88.    Cao, Z.-J. & Gao, G. Multi-omics single-cell data integration and regulatory inference with graph-linked
embedding. *Nat. Biotechnol.* **40**, 1458-1466 (2022).

89.    Powers, D. M. Evaluation: from precision, recall and F-measure to ROC, informedness, markedness and correlation.
Preprint at https://arxiv.org/abs/2010.16061 (2020).

1339