1 Article

Brown Adipose Tissue and Skeletal Muscle Coordinately Contribute to Thermogenesis in Mice

Yuna Izumi-Mishima,[1] Rie Tsutsumi,[1] Tetsuya Shiuchi,[2] Saori Fujimoto,[1] Momoka Taniguchi,[1] Yuko Okamatsu-Ogura,[3] Takeshi Yoneshiro,[4] Masashi Kuroda,[1] Kazuhiro Nomura,[1] and Hiroshi Sakaue[1,5,*]

[1]Department of Nutrition and Metabolism, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima 770-8503, Japan
[2]Department of Integrative Physiology, Institute of Biomedical Sciences, Tokushima University Graduate School, Tokushima 770-8503, Japan
[3]Laboratory of Biochemistry, Faculty of Veterinary Medicine, Hokkaido University, Sapporo 060-0818, Japan
[4]Division of Molecular Physiology and Metabolism, Tohoku University Graduate School of Medicine, Sendai 980-8575, Japan
[5]Lead contact
*Correspondence: hsakaue@tokushima-u.ac.jp

Abstract

22
Endotherms increase the rate of metabolism in metabolic organs as one
strategy to cope with a decline in temperature of the external environment.
However, an additional major contributor to maintenance of body temperature in
a cold environment is contraction-based thermogenesis in skeletal muscle. Here
we show that impairment of hind limb muscle contraction by cast immobilization
suppresses skeletal muscle thermogenesis and activates brown adipose tissue
(BAT) thermogenesis as a compensatory mechanism. BAT utilizes free
branched-chain amino acids (BCAAs) derived from skeletal muscle as an
energy substrate for thermogenesis, and interleukin-6 released by skeletal
muscle stimulates BCAA production in muscle for support of BAT
thermogenesis. Our findings suggest that BAT and skeletal muscle cooperate to
maintain body temperature in endotherms.
35
36

Introduction

Endotherms rely on the regulation of heat generation by metabolism in tissues of the body to maintain a stable core body temperature in the face of a fluctuating temperature of the external environment and thereby to ensure optimal physiological function.[1] Small mammals adjust their metabolic rate to cope with temperature fluctuations and to support their survival during seasonal acclimation. Some mammalian species are able to lower their metabolic rate and enter a state of torpor or hibernation in order to conserve energy and ensure survival.[2] However, small mammals that do not hibernate must increase their rate of metabolism, including obligatory thermogenesis in metabolic organs such as the liver and skeletal muscle as well as regulatory thermogenesis (shivering or nonshivering thermogenesis) in skeletal muscle and brown adipose tissue (BAT), in order to maintain thermal homeostasis in a cold environment.[1,3,4]

Under normothermic conditions, core body temperature is determined by basal metabolic organs such as the brain, heart, liver, and skeletal muscle.[1] Skeletal muscle is the largest weight-bearing organ in humans and consumes a substantial amount of energy, even at rest. Shivering is the major source of heat production in acute cold exposure before sufficient amount of UCP1 (uncoupling protein 1) can be recruited in BAT[1,5,6]. Shivering is gradually replaced by UCP1 dependent thermogenesis and nonshivering thermogenesis is necessary for long-term adaptation to a cold environment.[5] Skeletal muscle and BAT thermogenesis can compensate in part each other to maintain core body temperature is widely recognized[5-8], and recent study suggests that suppression of BAT thermogenesis has been found to promote thermogenesis in skeletal muscle.[8]

While skeletal muscle is the critical metabolic organ, it also stores an abundance of amino acids as energy substrates in the form of muscle proteins.[9] Skeletal muscle also has the plasticity for adapting to environmental conditions and shows metabolic flexibility in response to the substrate requirements of other organs and thereby contributes to the maintenance of energy balance.[9,10] Activation of BAT themogenesis is associated with increased lipid and glucose catabolism.[11-13] Succinate and branched-chain amino acids (BCAAs) have also been found to support BAT thermogenesis, however.[14–16] Although succinate may be provided by muscle contraction,[14,17] the primary source of BCAAs for BAT thermogenesis has remained unclear. Indeed, mitochondrial BCAA

catabolism in brown adipocytes promotes systemic BCAA clearance. Skeletal

muscle may therefore also respond to the energy demands of BAT.

Whereas body temperature and energy metabolism in metabolic organs are

closely related, how they are regulated in a coordinated manner is poorly

understood. Several endocrine factors alter metabolism in abdominal organs.[18]

In particular, interleukin (IL)–6 is an endocrine factor that plays multiple roles in

the response of the body to infection, exercise, and stress as well as in

inflammation.[19–21] IL-6 induces various fever responses, including nonshivering

thermogenesis in BAT, by promoting prostaglandin E2 (PGE2) synthesis in

central vascular endothelial cells.[22] On the other hand, IL-6 directly affects

energy metabolism in various organs, including skeletal muscle, liver, and

adipose tissue.[23–27] These multiple roles suggest that IL-6 is a key molecule

linking thermogenesis and energy metabolism.

Here we show that suppression of skeletal muscle thermogenesis by cast

immobilization in mice activates a compensatory thermoregulatory system

dependent on BAT thermogenesis. We also show that free BCAAs derived from

skeletal muscle support BAT thermogenesis, and we identify IL-6 as a key

regulator of this system. Our findings reveal the importance of skeletal muscle

as a source of amino acids and uncover a previously unrecognized

thermoregulatory system mediated by BAT and skeletal muscle metabolism.

Results

Cast immobilization suppresses muscle thermogenesis and activates a

compensatory thermoregulatory system

We first examined the role of skeletal muscle in endothermic homeostasis by

performing an acute cold tolerance test at 4°C for 4 h in mice with both hind

limbs immobilized in casts. We confirmed that ~50% of systemic skeletal

muscle was immobilized (Table S1), and that the immobilization induced

skeletal muscle atrophy within 3 to 5 days (Figure 1_figure supplement 1A–1C).

Consistent with previous findings,[28] we found that gene expression for the

ubiquitin ligases Atrogin-1 and MuRF-1 was increased during the early phase

(days 1–3) of cast immobilization (Figure 1_figure supplement 1D and 1E). We

also found that expression of the *Ppargc1a* (PGC-1α), a master regulator of

mitochondrial biogenesis and oxidative metabolism, was decreased from 10 h

after cast immobilization and the amount of *Tfam* (transcription factor A,

mitochondrial) mRNA was subsequently decreased in immobilized muscle

(Figure 1_figure supplement 1F and 1G). Metabolomic profiling of the

immobilized muscle revealed decreases in the amounts of fumaric acid and

malic acid within 1 day of cast immobilization (Figure 1_fugure supplement 1H).

Mice with both hind limbs immobilized for 7 days showed a significantly lower

core body temperature after cold exposure compared with control mice (Figure

1A). Cast immobilization for 24 h also induced cold intolerance with

preservation of muscle mass (Figure 1B and Figure 1_figure supplement 1A-

1C). To investigate whether cold intolerance depends on the immobilized

muscle mass, we examined the change in core body temperature in mice with

unilateral immobilization. Systemic locomotor activity during cold exposure was

similar for mice with unilateral or bilateral immobilization and for control mice

(Figure 1_figure supplement 1I). However, the decrease in body temperature

induced by cold exposure in mice with unilateral immobilization was attenuated

compared with that in mice subjected to bilateral immobilization (Figure 1C).

These findings suggested that immobilization of skeletal muscle suppresses

thermogenesis in this organ and it may negatively affects the maintenance of

body temperature under cold environments independently of locomotor activity.

Mice subjected to cast immobilization showed cold intolerance (Figure 1A

and B), but the core body temperature of cast-immobilized mice and control

mice were little different under room temperature (Figure 1D). We then

examined the effects of such immobilization on interscapular BAT (iBAT) to

investigate the mechanism underlying the maintenance of core body

temperature in mice with cast immobilization. Expression of UCP1 at the mRNA

and protein levels as well as the abundance of PGC-1α, a transcriptional

cofactor that promotes the expression of thermogenesis-related genes, showed

transient increases in iBAT after cast immobilization (Figures 1E–1H).

Expression of *Ucp1* mRNA in iBAT was significantly increased after cold

exposure compared to control mice at room temperature, whereas those

expression were not induced by cole exposure in cast immobilized mice (Figure

1_figure supplement 1J). Small and multilocular lipid droplets were apparent in

adipocytes of iBAT within 24 h of cast immobilization (Figure 1_figure

supplement 1K). However, mice subjected to cast immobilization showed no

increase in food consumption or activity levels (Figure 1_figure supplement 1L–

1N). These results indicated that the suppression of muscle thermogenesis by

cast immobilization induces BAT thermogenesis without an increase in food

consumption or activity levels. On the other hand, the amount of *Sln* (sarcolipin)

and *Camk2a* (Ca^{2+}- and calmodulin-dependent protein kinase II) mRNA were

not affected in immobilized muscle (Figure 1_figure supplement 1O and 1P).

These findings thus suggested that nonshivering thermogenesis was induced

minimally in cast-immobilized skeletal muscle. Cast immobilization thus

preferentially activated nonshivering thermogenesis in BAT, not in skeletal

muscle.

152 Cast immobilization promotes BAT thermogenesis via sympathetic

153 activation

We next explored the mechanism underlying the activation of BAT

thermogenesis by cast immobilization. We first focused on the sympathetic

nervous system, a general inducer of BAT thermogenesis,[29] and found that the

norepinephrine concentration in iBAT was tended to increased after 24 h of cast

immobilization (Figure 1I). We then subjected mice to sympathetic denervation

of iBAT. Expression of *Ucp1* mRNA in iBAT was significantly attenuated by such

denervation in mice with or without subsequent cast immobilization (Figure 1J).

Whereas adipocytes of iBAT showed small and multilocular lipid droplets after 7

days of cast immobilization in sham-operated mice, iBAT of mice subjected to

surgical denervation was hypertrophied and did not show such lipid droplets

(Figure 2_figure supplement 2A). Moreover, mice with denervated iBAT

manifested a lower core body temperature compared with sham-operated mice

after cold exposure at 4°C for 1 h, and cast immobilization further increased the

extent of hypothermia in the denervated mice (Figure 1K). These results

suggested that BAT thermogenesis is activated via sympathetic nerves to

maintain core body temperature in cast-immobilized mice.

171 Cast immobilization alters systemic metabolic dynamics associated with

172 BAT thermogenesis

We next examined the metabolic dynamics of iBAT as well as systemic

metabolic changes in mice subjected to cast immobilization. Metabolomics

analysis revealed increases in the amounts of carbohydrate metabolites and

amino acids in iBAT that were apparent as early as 24 h after cast

immobilization, and that such increases were not apparent in mice with iBAT

denervation (Figures 2A and 2B). Brown adipocytes incorporate fatty acids and

various other energy substrates to initiate thermogenesis and contribute to the

maintenance of core body temperature.[30] We found that expression of the

genes for fatty acid (Figure 2C) and succinate (Figure 2D) transporters in iBAT

was induced by cast immobilization and that such induction was prevented by
iBAT denervation. The concentration of succinate was also increased in iBAT of
cast-immobilized mice in a manner sensitive to iBAT denervation (Figure 2E).
Consistent with these changes in metabolite levels, oxygen consumption and
systemic lipid utilization were increased by cast immobilization in intact mice
(Figures 2F and 2G) but not in those subjected to iBAT denervation (Figure
supplement 1A and 1B).
The weight of epididymal white adipose tissue (eWAT) and hepatic
glycogen content were reduced by cast immobilization (Figures 2H and 2I).
Expression of the genes for glucose-6-phosphatase (G6Pase) and
phosphoenolpyruvate carboxykinase (PEPCK) in the liver increased gradually
after cast immobilization, peaking at day 5 (Figure 2_figure supplement 1C and
1D), suggesting that gluconeogenesis was slowly activated in the cast-
immobilized mice. Metabolomic profiling of the liver revealed marked increases
in the amounts of 3-phosphoglyceric acid and 2-phosphoglyceric acid within 1
day of cast immobilization (Figure 2_figure supplement 1E), suggesting that
glycerol influx was also enhanced in the liver. Together, these results thus
indicated that cast immobilization affects metabolic dynamics in the liver and
WAT, and that these changes might occur after the activation of BAT
thermogenesis.
The serum concentration of norepinephrine was increased within 1 day of cast
immobilization (Figure 2_figure supplement 1F), but these metabolic changes
induced by cast immobilization were not associated with by an increase in the
serum concentration of corticosterone or in expression of the gene for
corticotropin-releasing hormone (CRH) in the hypothalamus (Figure 2_figure
supplement 1G and 1H). These results suggest that cast immobilization may
activates BAT thermogenesis and systemic metabolic changes through lower
body temperature resulting from suppression of muscle thermogenesis rather
than stress.
211
Free amino acids are transferred from skeletal muscle to BAT and the liver
for energy homeostasis
We next focused on metabolism of amino acids in BAT and skeletal muscle in
mice after cast immobilization. As shown above, amino acid concentrations in
iBAT were increased by cast immobilization in a manner sensitive to
denervation of iBAT (Figure 2B). The amounts of free amino acids were also

increased in both soleus and extensor digitorum longus (EDL) muscles of cast-

immobilized mice (Figure 3A), whereas those in serum and the liver were not

increased by cast immobilization (Figure 3B). We then examined gene

expression profiles for solute carrier (SLC) transporters that mediate the

transport of amino acids across cell membranes as well as for mitochondrial

BCAA catabolic enzymes (BCAT2, branched-chain aminotransferase 2;

BCKDHA, branched-chain keto acid dehydrogenase E1 subunit α) in iBAT, liver,

and soleus. The expression of these genes in iBAT was increased after cast

immobilization for 10 to 24 h, whereas that in the liver was increased after 5 to 7

days (Figure 3C). In particular, expression of the genes for SLC1A5 (a sodium-

dependent glutamate transporter) and SLC38A2 (a sodium-coupled neutral

amino acid transporter) was significantly increased in iBAT after cast

immobilization for 10 h (Figure 3_figure supplement 1A–1F). In the liver,

expression of the gene for SLC38A2, which is upregulated by extracellular

amino acid deprivation,[31,32] and of that for BCKDHA, which is thought to be

upregulated under fasting conditions to oxidize BCAAs for gluconeogenesis,[33,34]

was significantly increased by cast immobilization (Figure 3C and

Figure3_figure supplement 1G–1K). On the other hand, gene expression for

amino acid transporters (SLC1A5, SLC7A5, SLC38A2) and BCAA catabolic

enzymes (BCAT2, BCKDHA) in soleus was significantly decreased after cast

immobilization (Figure 3C and Figure 3_figure supplement 1L–1Q). Expression

of the gene for SLC43A1, which is upregulated by food deprivation and is

thought to transport amino acids from cells to the external environment,[35] was

significantly increased in soleus by cast immobilization (Figure 3C and Figure

3_figure supplement 1O).

We next investigated amino acid incorporation in tissues by

administration of [³H]leucine into the tail vein of mice. Incorporation of

[³H]leucine was not increased in skeletal muscle of mice subjected to cast

immobilization for 1 or 3 days (data not shown), whereas it was significantly

increased in iBAT, liver, and kidney of those subjected to cast immobilization for

1 day (Figure 3D). Gene expression for SLC25A44, a mitochondrial BCAA

transporter that contributes to BAT thermogenesis,[15] was increased in iBAT at

24 h after cast immobilization (Figure 3E and Figure 3_figure supplement 1R–

1T). However, the early increases in expression of the genes for SLC25A44,

SLC1A5, and SLC38A2 induced by cast immobilization in iBAT of intact mice

were prevented by denervation of iBAT (Figure 3E and Figure 3_figure

supplement 1U–1AA). Our results thus suggested that free amino acids are
utilized as an energy substrate in BAT and the liver rather than in skeletal
muscle of cast-immobilized mice.
257
Cast immobilization upregulates IL-6 gene expression in BAT and skeletal
muscle
We next set out to identify regulatory molecules that might contribute to the
maintenance of body temperature and to the systemic metabolic changes in
cast-immobilized mice. Gene expression analysis revealed that the IL-6 gene
tended to be induced earlier and to a greater extent in iBAT and soleus than
were genes for other cytokines including tumor necrosis factor–α (TNF-α),
monocyte chemoattractant protein–1 (MCP-1), IL-1β, IL-10, and IL-15; for
batokines including fibroblast growth factor 21 (FGF21) and bone
morphogenetic protein 8B (BMP8B); or for myokines including FGF21 and irisin
(Figures 4A–4D and Figure 4_figure supplement 1A–1N). Whereas the increase
in the amount of *Il6* mRNA was apparent as early as 6 to 10 h in iBAT, that in
soleus was more gradual (Figures 4C and 4D). IL-6 has been found to be
released by various organs, including adipose tissue and skeletal muscle, in
response to inflammation, stress, or exercise.[18-21] The abundance of *Il6* mRNA
in liver, eWAT, or spleen was not affected by cast immobilization (Figure
4_figure supplement 1O–1Q). Expression of the genes for serum amyloid A3
(*Saa3*) and suppressor of cytokine signaling 3 (*Socs3*), both of which are
regulated by IL-6 signaling, was also increased in iBAT and soleus at the same
time as was that of *Il6* (Figures 4E–4H). Expression of these genes was also
increased in the liver or eWAT at 10 or 24 h after cast immobilization (Figures
4I–4L). However, expression of *Il6* was not induced by cast immobilization in
denervated iBAT (Figures 4M). The serum concentration of IL-6 was
significantly increased from 10 h after cast immobilization (Figure 4N), whereas
that of TNF-α was not altered in cast-immobilized mice (Figure 4_figure
supplement 1R). Together, these results suggested that cast immobilization
induces *Il6* expression in iBAT and skeletal muscle, and that induction of *Il6*
expression by cast immobilization in iBAT is dependent on the sympathetic
nervous system.
287
IL-6 affects energy metabolism in BAT and skeletal muscle of cast-
immobilized mice
We investigated the role of IL-6 in the skeletal muscle–BAT
thermoregulatory system with the use of subjected IL-6 knockout (KO) mice to

cast immobilization. Expression of *Ucp1* in iBAT was significantly increased at
24 h after cast immobilization in both WT and IL-6 KO mice (Figure 5_figure
supplement 1A). However, core body temperature of cast-immobilized IL-6 KO
mice was decreased to a greater extent by cold exposure at 4°C for 4 h
compared with that of cast-immobilized WT mice (Figure 5A). Furthermore,
unlike that in WT mice, oxygen consumption was not increased in IL-6 KO mice
after cast immobilization (Figure 5_figure supplement 1B and 1C).
Metabolomics analysis revealed that the amounts of carbohydrate metabolites
including glucose 1-phosphate, glucose 6-phosphate, and phosphoenolpyruvic
acid in iBAT were higher for IL-6 KO mice than for WT mice, and were not
increased substantially by cast immobilization in the mutant mice (Figure 5B).
Whereas the concentrations of amino acids such as Ser, Asn, Lys, Met, His,
Phe, and Tyr in iBAT of WT mice were increased markedly at 24 h after cast
immobilization, such was not the case for IL-6 KO mice (Figure 5C). Expression
of the genes for SLC25A44, BCKDHA, and CD36 in iBAT was higher for IL-6
KO mice than for WT mice, but expression of the genes for SLC25A44 and
CD36 in iBAT of IL-6 KO mice was not increased further at 24 h after cast
immobilization (Figures 5D–5F). Incorporation of [^{3}H]leucine was also not
increased in iBAT of IL-6 KO mice after cast immobilization for 24 h (Figure 5G).
In addition, adipocytes with small and multilocular lipid droplets were observed
in iBAT of IL-6 KO mice with or without cast immobilization (Figure 5_figure
supplement 1D). The respiratory exchange ratio (RER) of WT mice was
significantly decreased in both the light and dark phases for the first 5 days of
cast immobilization, whereas similar differences were apparent only for the first
2 days in IL-6 KO mice (Figure 5_figure supplement 1E and 1F). IL-6 has been
shown to promote lipolysis and fatty acid oxidation in WAT.[26] We found that,
unlike in WT mice, cast immobilization did not result in a loss of eWAT mass in
IL-6 KO mice (Figure 5_figure supplement 1G). Collectively, these data
suggested that utilization of energy substrates was not substantially altered by
cast immobilization in IL-6 KO mice.
The amounts of BCAAs in soleus were significantly increased after cast
immobilization in WT mice but not in IL-6 KO mice (Figures 5H–5J). Serum
BCAA concentrations were decreased after cast immobilization in both WT and
IL-6 KO mice (Figure 5_figure supplement 1H–1J). These findings suggested
that deficiency of IL-6 impairs the maintenance of core body temperature and
alters energy metabolism in BAT and skeletal muscle in mice subjected to cast
immobilization.
329

Administration of IL-6 increases BCAA concentrations in skeletal muscle of cast-immobilized IL-6 KO mice

We examined whether administration of exogenous IL-6 might increase amino acid concentrations in skeletal muscle and BAT of IL-6 KO mice with cast immobilization. Administration of IL-6 restored the increase in the amounts of amino acids including BCAAs in soleus of IL-6 KO mice with cast immobilization (Figures 6A–6C and Figure 6_figure supplement 1A). Treatment with IL-6 also increased expression of *Saa3* and *Socs3* (Figure 6_figure supplement 1B and 1C) as well as the amounts of BCAAs (Figure 6D) in cultured mouse C2C12 myotubes. However, the amounts of carbohydrate metabolites in iBAT of cast-immobilized IL-6 KO mice were not increased by IL-6 administration (Figure 6_figure supplement 1D). In contrast to its effects on soleus, administration of IL-6 also did not increase the amounts of amino acids in iBAT of IL-6 KO mice with cast immobilization (Figure 6_figure supplement 1A and 1E). In addition, exogenous IL-6 did not increase the expression of UCP1, SLC25A44, and BCKDHA genes in iBAT of these mice (Figure 6_figure supplement 1F–1H). On the other hand, administration of IL-6 ameliorated the cold intolerance of IL-6 KO mice with cast immobilization (Figure 6E). We therefore examined whether IL-6 might increase core body temperature via the sympathetic nervous system with the use of C57BL/6J mice with denervated iBAT. Core body temperature of cold-exposed mice with cast immobilization and denervated iBAT was not significantly increased by administration of IL-6 (Figure 6F).

We also found that the core body temperature of IL-6 KO mice was lower than that of WT mice after cold exposure at 4°C for 4 h (Figure 6_figure supplement 1I) whereas the amount of *Ucp1* mRNA in iBAT was significantly increased for IL-6 KO mice compared with WT mice at room temperature (Figure 6_figure supplement 1J). We then investigated the possible direct effects of IL-6 on energy metabolism in BAT with the use of cultured mouse brown adipocytes. Expression of the SLC25A44, BCKDHA, and UCP1 genes in the brown adipocytes were not increased by treatment with IL-6 (Figure 6_figure supplement 1K–1M). In contrast, treatment with the β_3-adrenergic receptor agonist CL316 243 increased the expression of these genes, and this upregulation was suppressed by treatment with IL-6 (Figure 6_figure supplement 1K–1M). Long-term administration of CL316 243 in mice was previously shown to increase expression of *Ucp1*, *Slc25a44*, and *Bckdha* in BAT.[16] We found that administration of CL316 243 for 7 days significantly increased the amounts of *Ucp1*, *Slc25a44*, and *Bckdha* mRNAs in iBAT of IL-6

KO mice but not in that of WT mice (Figure 6_figure supplement 1N–1P).

Together, these data suggested that IL-6 may suppress β-adrenergic receptor–

mediated thermogenesis and BCAA metabolism in brown adipocytes.

Nevertheless, IL-6 induces fever response through stimulation of the central[21,22]

and uptake of energy-rich BCAA delivered to BAT from skeletal muscle may

facilitate a net induction of BAT thermogenesis.

Skeletal muscle–derived IL-6 increases BCAA concentrations in skeletal

muscle for BAT thermogenesis

Given that our data suggested that an elevated blood concentration of IL-6

contributes to maintenance of body temperature in cast-immobilized mice in a

manner dependent on the sympathetic nervous system (Figures 6E and 6F), we

sought to determine the main source of circulating IL-6 in cast-immobilized

mice. We first focused on the acute expression of the IL-6 gene apparent in

iBAT after cast immobilization (Figure 4C), and we subjected mice to surgical

removal of iBAT. The serum IL-6 concentration of such mice was not increased

after cast immobilization (Figure 6G). We also found that the reduction in eWAT

mass after cast immobilization in C57BL/6J mice was suppressed by surgical

removal of iBAT (Figure 6H). These results suggested that BAT-derived IL-6

induces lipolysis in WAT.

We further investigated the effects of BAT-derived IL-6 on skeletal

muscle. Whereas the serum concentration of IL-6 in mice depleted of iBAT was

not affected by cast immobilization (Figure 6G), the concentrations of BCAAs in

soleus were increased after cast immobilization in such mice (Figures 6I–6K).

The amount of *Slc43a1* mRNA in soleus of mice depleted of iBAT was

significantly increased by cast immobilization (Figure 6_figure supplement 2A).

Gene expression for other amino acid transporters (such as SLC1A5 and

SLC7A5) and for BCAA catabolic enzymes (BCAT2 and BCKDHA) in soleus

also showed a similar pattern of changes after cast immobilization in iBAT-

depleted mice (Figure 6_figure supplement 2B–2E) as in intact mice (Figure

3_figure supplement Fig. 1L, 1M, 1P, and 1Q). Moreover, cast immobilization for

7 days induced similar losses of both soleus and gastrocnemius mass in mice

with or without removal of iBAT (Figure 6_figure supplement 2F and Table S2).

These data thus suggested that IL-6 secretion in an autocrine or paracrine

manner may regulate BCAA metabolism in skeletal muscle independently of IL-

6 derived from iBAT.

Acute cold exposure also induced IL-6 from skeletal muscle to supply

BCAA for BAT thermogenesis

We further found that expression of *Il6* was also increased in not immobilized

fore limb muscles (biceps brachii muscle; BBM) after cast immobilization

(Figure 6_figure supplement 2G). Expression of *Saa3* and *Socs3* were

increased in BBM at the same time as was that of *Il6* after cast immobilization

(Figure 6_figure supplement 2H and 1I). Furthermore, gene expression for the

ubiquitin ligases Atrogin-1 and MuRF-1 in BBM was increased during the early

phase of cast immobilization (Figure 6_figure supplement 2J and 2K). BCAA

concentrations in BBM were also increased after cast immobilization (Figure

6_figure supplement 2L-2N). These data suggest that cast immonbilization

stimulates BCAA metabolits in not immobilized fore limb muscles as well as

immobilized muscle.

Finally, we investigated whether the thermoregulatory system through amino

acid metabolism in the interaction between BAT and skeletal muscle also plays

an important role in other conditions besides cast immobilization. Acute cold

exposure also increased expression of *Il6* only in iBAT and skeletal muscle in

mice (Figure 6L). Expression of *Saa3* and *Socs3* were increased in soleus

(Figures 6M and 6N), and both soleus and gastrocnemius mass were

decreased by cold exposure at 4°C for 4 h (Figure 6_figure supplement 2O and

2P). Furthermore, the amounts of valine in soleus were significantly increased

by acute cold exposure in WT mice but not in IL-6 KO mice (Figures 6O and

6P). BCAA concentrations in soleus were also increased after cold exposure at

4°C for 4 h in iBAT-depleted mice (Figure 6_figure supplement 2Q). Together,

thermoregulatory system through amino acid metabolism in skeletal muscle and

BAT may be activated via sympathetic nervous system under cold temperature.

Discussion

We have here shown that cast immobilization suppressed skeletal muscle

thermogenesis, resulting in failure to maintain core body temperature in a cold

environment. Cast immobilization also activated BAT thermogenesis via the

sympathetic nervous system and triggered systemic changes in energy

metabolism associated with BAT thermogenesis. In addition, we found that free

BCAAs derived from skeletal muscle serve as substrates for energy metabolism

in BAT, and that skeletal muscle–derived IL-6 promotes this provision of BAT

with amino acids from muscle.

Exercise-induced muscle contraction generates large amounts of heat

which dependents on hydrolysis of ATP, and skeletal muscle, as the first organ

to be recruited for thermogenesis, plays an important role in maintenance of

body temperature in endotherms.[36] Muscle thermogenesis can account for up to

90% of systemic oxygen consumption during periods of maximal recruitment of

muscle contraction, such as during exercise or an intense bout of shivering.[36,37]

We found that cast immobilization suppressed mitochondria related genes

expression and metabolites of TCA cycle from early stage of immobilization,

suggesting that metabolic rate is rapidly decreased in immobilized muscle. Thus

our results of cast immobilization leading cold intolerance in mice show that

metabolic thermogenesis in skeletal muscle plays a key role in mammals even

in BAT-enriched mammals such as rodents.

Thermoregulatory system in endotherms cannot be explained by

thermogenesis based on muscle contraction alone, with nonshivering

thermogenesis being required as a component of the ability to tolerate cold

temperatures in the long term[1]. Our results now show that expression of the

gene for sarcolipin, a key regulator of the sarco/endoplasmic reticulum Ca^{2+}-

ATPase (SERCA) and an important mediator of muscle nonshivering

thermogenesis, was not increased in muscle after cast immobilization. On the

other hand, BAT thermogenesis was activated via the sympathetic nervous

system even when skeletal muscle was immobilized. It is a well-recognized fact

that BAT and skeletal muscle function in an orchestrated manner to maintain

core body temperature in endotherms[5-7]. A bit surprisingly, BAT capacity itself

was not sufficient to maintain stable body temperature under acute cold

exposure, whereas compensatory BAT recruitment protected cast immobilized

mice from hypothermia at room temperature.

We also found that changes in systemic energy metabolism induced by

cast immobilization were associated with the activation of BAT thermogenesis.

Activation of BAT results in the tissue becoming a high consumer of a variety of

energy substrates including lipids, glucose, BCAAs, succinate, and lactate.[30] In

addition, our results suggest that skeletal muscle is a source of free amino acids

for BAT thermogenesis or hepatic gluconeogenesis. In response to metabolic

demands imposed by starvation, exercise, or cold exposure, for example,

skeletal muscle manifests metabolic flexibility in order to meet the demands of

other organs and is an important determinant of metabolic homeostasis.[9,10] In

contrast, the reduction in metabolic rate may contribute to protein conservation

and maintain skeletal muscle mass under hibernation.[9] Our findings now

provide new insight into the importance of skeletal muscle as a source of amino

acids, with a recent study also suggesting that a highly active proteolysis

system in the heart provides substantial amounts of amino acids for distribution

to other organs via the bloodstream.[38] The heart may therefore be another

source of free BCAAs for BAT thermogenesis in cast-immobilized mice.

We found that muscle-derived IL-6 directly increased the abundance of

BCAAs in skeletal muscle after cast immobilization. Previous studies have

shown that excessive IL-6 induces amino acid catabolism in skeletal

muscle,[39,40] and that IL-6 is one of the factors responsible for the induction of

muscle atrophy in cast-immobilized mice.[41] IL-6 is a secreted factor induced by

several mechanisms including muscle contraction, intracellular calcium

signaling, and inflammation.[20,42] However, inflammation associated with

macrophage infiltration is not thought to be induced in the early stages of

muscle atrophy[28], and thus inflammation-induced IL-6 expression may therefore

not contribute to the changes in amino acid metabolism in cast-immobilized

muscle. In addition, whereas intracellular calcium concentrations may increase

after muscle immobilization for 2 weeks,[43] they appear to decrease in response

to short-term cast immobilization.[41] Cast immobilization–induced *Il6* expression

may therefore also not be regulated by calcium signaling. Upregulation of IL-6

expression by cast immobilization was previously shown to be mediated by a

Piezo1-KLF15 axis,[41] whereas our observation that *Il6* expression were

increased in non immobilized fore limb muscles. Furthermore, acute cold

exposure tended to increase *Il6* expression in skeletal muscle, suggesting that

muscle-derived IL-6 for thermoregulatory system may also be regulated by the

central nervous system. A recent study showed that motor circuits modulate the

production of neutrophil-induced chemokines in skeletal muscle after acute

stress.[44] The regulation of IL-6 production in muscle by the central nervous

system warrants further investigation.

The central nervous system tightly controls core body temperature

through integrates information about external temperature, humidity, and

thermal sensation to induce an adaptive response.[1] Disappointingly, our study

was inconclusive as to whether the trigger for BAT thermogenesis after cast

immobilization was hypothermia associated with supression of muscle

thermogenesis or stress. However, we found that acute cold exposure also

recruits substrate supply from skeletal muscle for BAT thermogenesis. Possibly,

thermoregulatory system through amino acid metabolism in skeletal muscle and

BAT may also be an important metabolic strategy even in the case of fever

responce induced by stress or infection. Furthermore, inflammation, infection,

and acute stress trigger rapid increases in the circulating IL-6 concentration[21,45]

and induce a fever response by promoting PGE2 synthesis. Mitochondrial

BCAA oxidation in BAT was recently shown to be increased to support

thermogenesis induced by PGE2.[16] We found that BAT-derived IL-6 increased

blood IL-6 levels after cast immobilization and administration of exogenous IL-6

ameliorated cold intolerance in cast-immobilized IL-6 KO mice via the

sympathetic nervous system. It may modulate thermal and energy homeostasis

through the fever response and metabolic regulation in multiple organs.

In conclusion, we have shown that cast immobilization induced

thermogenesis in BAT that was dependent on the utilization of free amino acids

derived from skeletal muscle, and that muscle-derived IL-6 stimulated BCAA

metabolism in skeletal muscle. Our findings may provide new insights into the

significance of skeletal muscle as a large reservoir of amino acids in the

regulation of body temperature. In addition, given that circulating levels of IL-6

or BCAAs are associated with obesity and diabetes,[46,47] IL-6–mediated BCAA

metabolism in skeletal muscle and BAT may also be associated with muscle

atrophy in metabolic diseases. Further investigation of IL-6–dependent amino

acid metabolism in BAT and skeletal muscle may inform the development of

preventive or therapeutic interventions for some forms of muscle atrophy.

Limitations of the study

Given that rodents are BAT-enriched mammals, whether BAT

thermogenesis is similarly activated in humans after skeletal muscle

immobilization remains to be investigated. Our study also did not determine the

mechanism underlying the induction of *Il6* expression in skeletal muscle by cast

immobilization. In addition, whereas muscle-derived IL-6 may stimulate protein

catabolism in immobilized muscle, the mechanism by which IL-6 increases

BCAA concentrations in skeletal muscle requires further investigation.

Methods

Mice

All experiments were approved by the Animal Care and Use Committee of Tokushima University and were conducted in accordance with the guidelines for the care and use of animals approved by the Council of the Physiological Society of Japana and ARRIVE guidelenies. Every effort was made to minimize animal suffering and to reduce the number of animals used in the experiments. All mice were housed at a constant room temperature of 23° ± 1°C and with a 12-h-light/12-h-dark cycle (lights on at 8.00 a.m.). They were fed standard nonpurified chow (Oriental Yeast) and had free access to both food and water. Our study examined male mice because male animals exhibited less variability in phenotype. Male mice at 9 to 13 weeks of age were studied. The mice were randomly assigned to experimental groups at the time of purchase or weaning. C57BL/6J mice were obtained from Japan SLC (Shizuoka, Japan), and IL-6 KO mice (B6;129S2-Il6<tm1kopf>) from RIKEN BRC through the National Bioresource Project of MEXT/AMED.[46] The IL-6 KO mice were maintained on the C57BL/6J background.

Cells

The mouse myoblast cell line C2C12 was obtained from American Type Culture Collection (CRL-1772™) and was maintained in Dulbecco's modified Eagle's medium (DMEM) containing 25 mM glucose (#D6429, Sigma) and supplemented with 10% fetal bovine serum (535-94155, BioSera) and 1% penicillin-streptomycin (15140-122, Gibco). After the cells had achieved 95% to 100% confluence, the culture medium was changed to DMEM containing 25 mM glucose and supplemented with 2% horse serum (16050-130, Thermo Fisher Scientific) and the cells were cultured for 5 days to promote their differentiation from myoblasts into myotubes. The myotubes were washed with phosphate-buffered saline (PBS) and then incubated in DMEM containing 25 mM glucose and supplemented with 1% bovine serum albumin (A8806-5G, Sigma) and 1% penicillin-streptomycin for 16 h before exposure to recombinant mouse IL-6 (rIL-6) at 50 ng/ml (575702, BioLegend) or vehicle (PBS) in the same medium for 1 h. The treated cells were washed with PBS or 5% mannitol solution and collected for RNA or metabolite extraction.

Mouse preadipocytes were obtained from Dr. Takeshi Yoneshiro the University of Tohoku, Tohoku, Japan[48] and were maintained in DMEM

containing 25 mM glucose and supplemented with 10% fetal bovine serum and
1% penicillin-streptomycin. After the cells had achieved 95% to 100%
confluence, they were induced to differentiate into brown adipocytes by
incubation in the same medium that was also supplemented with 1 nM
triiodothyronine (45006-44, Nacalai Tesque), insulin (093-06473, Wako) at 5
µg/ml, dexamethasone (11107-51, Nacalai Tesque) at 2 µg/ml, 0.5 mM
isobutylmethylxanthine (15879, Sigma), and 125 µM indomethacin (19233-51,
Nacalai Tesque). After culture for 2 days, the medium was replaced with DMEM
containing 25 mM glucose and supplemented with 10% fetal bovine serum, 1%
penicillin-streptomycin, and insulin (5 µg/ml), and the cells were maintained in
this medium with medium replenishment every 2 days. The cells were fully
differentiated at 6 to 7 days after the induction of differentiation. For
experiments, the brown adipocytes were washed with PBS at 7 days after
differentiation initiation and then incubated in DMEM containing 25 mM glucose
and supplemented with 1% bovine serum albumin and 1% penicillin-
streptomycin for 12 h before exposure to rIL-6 (50 ng/ml), 1 µM CL316 243
(Ab144605, Abcam), both agents, or vehicle (PBS). The treated cells were
collected after 24 h for RNA extraction.
598

Cast immobilization
Mice were subjected to bilateral or unilateral cast immobilization of hind limbs,
and unfixed mice were studied as controls. In brief, mice were lightly
anesthetized with inhalational isoflurane (MSD Animal Health), and the hind
limbs were fixed in a natural position, with care taken to avoid skin irritation and
congestion. After casting, the mice were housed individually with free access to
standard mouse chow and water. The mice out of cast immobilization were
excluded from experiments. In all experiments, mice were killed by cardiac
puncture after food deprivation for 3 h and tissue samples were collected.
608

Surgical denervation of iBAT
Surgical denervation of iBAT was performed on 9-week-old male mice as
described previously.[49] Animals were anesthetized by intraperitoneal injection of
domitor (Zenoaq) at a dose of 0.75 mg/kg, midazolam (Wako) at 4 mg/kg, and
butorphanol tartrate (Meiji Seika Pharma) at 5 mg/kg, and the incision site was
shaved and then disinfected with 0.05% chlorhexidine gluconate (5% Hibitane,
Sumitomo Pharma). A midline skin incision was made in the interscapular

region to expose both iBAT pads, the nerve fibers (five nerves per pad) were
identified and cut with sterile scissors, and the incision was closed with sutures.
For sham surgery, a skin incision was made and the medial side was gently
exposed. All mice were then individually housed in clean cages at room
temperature and were monitored daily. Cast immobilization was performed 1
week after surgery.

Surgical removal of iBAT
Surgical removal of iBAT was performed on 11-week-old male mice, with cast
immobilization being performed 1 week after surgery. Animals were
anesthetized and the incision site was shaved and disinfected as described for
iBAT denervation. A midline skin incision was made in the interscapular region,
all iBAT was removed, and the incision was closed with sutures. For sham
surgery, a skin incision was made and the medial side was gently exposed. All
mice were then housed individually in clean cages at room temperature.

Acute cold tolerance test
Core body temperature was measured with a Homeothermic Monitor (BWT-
100A, BRC) by gentle insertion of a thermal probe into the rectum of the mouse.
After recording of baseline body temperature at room temperature, animals
were placed in a cold chamber at 4°C for up to 4 h without access to food.
Tissue was then collected and snap-frozen.

Treatment of mice with CL316 243 or rIL-6
IL-6 KO mice with cast immobilization for 24 h were subjected to intraperitoneal
administration of 400 ng of rIL-6 (575702, BioLegend) in saline or of saline
alone. The mice were maintained sedentary for 3 h without access to food
before collection of tissue samples. Acute exposure to a cold environment
chamber was initiated 30 min after intraperitoneal treatment with rIL-6 (400 ng)
or saline in IL-6 KO mice or in C57BL/6J mice with denervated iBAT that had
been subjected to cast immobilization for 24 h. IL-6 KO mice with denervated
iBAT were treated intraperitoneally with CL316 243 (Ab144605, Abcam) at a
dose of 0.1 mg/kg, rIL-6 (400 ng), both agents, or saline vehicle, and tissue was
collected 3 h later. For long-term treatment of IL-6 KO mice with CL316 243, the
animals were injected intraperitoneally with the drug (0.1 mg/kg per day) or
saline for 7 days. Tissue was collected at 3 h after drug treatment on the last

day.

Oxygen consumption and RER measurement

Mice were acclimated to the analysis cage for 3 days before the start of

metabolic measurements. They were housed individually with free access to

food and water. Analysis of respiratory gases was performed with a mass

spectrometer (ARCO-2000, ARCO System). Measurements were performed for

seven consecutive days after cast immobilization.

Locomotor activity recording

The activity level of mice was assessed with an infrared activity monitor

(ACTIMO-100N, Shin Factory). A cage containing one mouse was placed inside

the activity monitor with infrared beams at 20-mm intervals. Mouse movements

were counted every 0.5 s for 24 h. Recording was performed for seven

consecutive days after cast immobilization.

RNA isolation and RT and real-time PCR analysis

Total RNA was extracted from mouse tissues and cells with the use of RNAiso

(#9109, Takara Bio) and subjected to RT with the use of a TaKaRa PrimeScript

II 1st Stand cDNA Synthesis Kit (Takara Bio). The resulting cDNA was subjected

to real-time PCR analysis with Fast SYBR Green Master Mix (3485612, Applied

Biosystems) in a Step One Plus Real-Time PCR System (Applied Biosystems).

The abundance of target mRNAs was normalized by that of *Gapdh* mRNA. The

PCR primers are listed in Table S3.

Protein extraction and immunoblot analysis

Total protein was extracted from mouse tissues by homogenization in a lysis

solution consisting of 1 M Tris-HCL (pH 7.4), 0.1 M EDTA, 1% Nonidet P-40, 5

mM sodium pyrophosphate, and phosphatase (25955-24, Nacalai Tesque) and

protease (07575-51, Nacalai Tesque) inhibitor cocktails. The homogenate was

centrifuged at 15,000 × g for 15 min at 4°C, and the resulting supernatant was

assayed for protein concentration with a Pierce BCA Protein Assay Kit (Thermo

Fisher Scientific) and then subjected to SDS-polyacrylamide gel electrophoresis

on a 10% acrylamide/4% bisacrylamide gel. The separated proteins were

transferred to polyvinylidene difluoride membrane, which was then exposed to

5% dried skim milk for 1 h at room temperature before incubation with primary

antibodies including those to PGC-1α (A12348, Abclonal), to UCP1 (sc-293418,

Santa Cruz Biotechnology), and to GAPDH (CAB932Hu01, Cloud-Clone) for 16

h at 4°C. Immune complexes were detected with horseradish peroxidase–

conjugated secondary antibodies (458, MBL; 7076P2, Cell Signaling

Technology), enhanced chemiluminescence reagents (170-5061, Bio-Rad

Labotatories, Inc.), and an Amersham Imager 600 instrument (GE Healthcare).

Band intensity was quantified with ImageJ software.

Histology

BAT was fixed in formalin, sectioned, and stained with hematoxylin-eosin.

Metabolomics analysis

Targeted profiling of amino acid metabolites was performed by capillary

electrophoresis–time-of-flight mass spectrometry (CE-TOF/MS) with an Agilent

7100 CE system coupled to an Agilent 6230 TOF mass spectrometer (Agilent

Technologies) according to previously described methods.[50]

Measurement of noradrenaline

Tissue samples were homogenized in methanol containing internal standards

(($\pm$)-Norepinephrine-D6 hydrochloride solution, Sigma-Ardrich). The mixture

was centrifuged at 4 °C, and the aqueous fraction was isolated and centrifuged

through an cosmonice filter W (06543-04, nacalai tesque). Noradrenaline

content of the supernatants was measured by high-performance liquid

chromatography (SHIMAZU) with a reversed-phase column (Cadenza CD-C18

MF 100 × 2 mm 3 μm, Imtakt) and tandem mass spectrometry (MS/MS) (API

3200, AB SCIEX). Data are analyzed using AnalystLauncher (AB SCIEX).

Measurement of [³H]leucine uptake

Mice were deprived of food for 3 h before intravenous injection of L-[4,5-

³H(*N*)]leucine (NET1166, PerkinElmer) at a dose of 1 μCi/g. They were then

maintained in a resting state for 40 min until blood and tissue collection, with

perfusion being performed before tissue isolation. Isolated tissue was cut into

small pieces, which were then placed in a round-bottom polypropylene tube

before the addition of 500 μl of distilled water, 150 μl of 30% hydrogen peroxide

(Wako), and 150 μl of 2 M KOH (1310-58-3, Hayashi Pure Chemical) in 2-

propanol (000-64783, Kishida Chemical) and incubation at 65°C for 2 h. The

tissue lysate was neutralized with 30% acetic acid, and 1 ml of liquid scintillation
solubilizer (Soluen-350, PerkinElmer) was added to the sample before
incubation at 50°C for 5 h and the further addition of 5 ml of liquid scintillation
cocktail (Hionic-Flour, PerkinElmer). Leucine uptake in each organ was
quantified by measurement of radioactivity with a scintillation counter (LSC-
7400, Hitachi).
730
731   **Hepatic glycogen assay**
732   The liver was homogenized in ice-cold citrate buffer (pH 4.2) containing NaF
733   (2.5 g/l), the homogenate was centrifuged at 14,000 × g for 5 min at 4°C, and
734   the resulting supernatant was assayed for glycogen with an EnzyChrom
735   Glycogen Assay Kit (BioAssay Systems).
736
ELISAs
Blood samples were collected from the heart of mice that had been
anesthetized by intraperitoneal injection of domitor (Zenoaq) at a dose of 0.75
mg/kg, midazolam (Wako) at 4 mg/kg, and butorphanol tartrate (Meiji Seika
Pharma) at 5 mg/kg after food deprivation for 3 h, and they were centrifuged at
7500 × g for 10 min at 4°C to isolate serum. Serum IL-6 and TNF-α
concentrations were determined with enzyme-linked immunosorbent assay
(ELISA) kits (Legend Max Mouse IL-6 ELISA Kit, BioLegend, and Mouse TNF
Alpha Uncoated ELISA, Invitrogen, respectively). Serum noradrenaline was
extracted with the use of a cis-diol–specific affinity gel, acylated, enzymatically
converted, and then measured with an ELISA kit (Noradrenaline Research
ELISA, ImmuSmol). Serum corticosterone was determined with a
Corticosterone ELISA Kit (Enzo Life Sciences).
750
Statistical analysis
Data are presented as means ± SEM unless indicated otherwise and were
analyzed with statistical software (JMP or Statcel–The Useful Add-in Forms on
Excel 4th ed.). Comparisons between two groups were performed with the
paired or unpaired t test, as appropriate. Those among more than two groups
were performed with Dunnett's test, by one-way analysis of variance (ANOVA)
followed by the Tukey-Kramer test, or by two-way ANOVA followed by Tukey's
post hoc test or the post hoc paired/unpaired t test with Bonferroni's correction.
A two-tailed p value was calculated for all experiments, and a value of <0.05

was considered statistically significant.

Acknowledgements
We thank members of the Radioisotope Center of Tokushima University
Graduate School for technical support with the tracer experiments; members of
the Special Mission Center for Metabolome Analysis, School of Medical
Nutrition, Faculty of Medicine, Tokushima University, for technical support with
the metabolomics analysis; Yoichiro Iwakura (University of Tokyo) for
permission to study the IL-6 KO mice; Kohta Onishi (University of Tokushima)
for technical support with the LC/MS/MS analysis; and the Support Center for
Advanced Medical Sciences at Tokushima University Graduate School of
Biomedical Sciences for general technical support. This study was supported by
Funds for the Development of Human Resources in Science and Technology
from the Ministry of Education, Culture, Sports, Science, and Technology of
Japan (MEXT) through the Home for Innovative Researchers and Academic
Knowledge Users consortium, and by KAKENHI grants (21J14844 to Y.I.-M. ,
23K19926 to Y.I.-M., 23K19926 to Y.I.-M., 22H03535 to H.S., and 21K11724 to
R.T.) from the Japan Society for the Promotion of Science.
778

779 **Auther Contributions**
780 Y.I.-M., T.S., S.F., and M.T. conducted the experiments; R.T. and H.S. designed
781 the experiments; Y.I.-M., R.T., and H.S. wrote the manuscript; Y.O.-O., T.Y.,
782 K.N., and M.K. provided conceptual advice on data interpretation.
783

784 **Decompeting interests**
785 The authors declare no competing interests.
786

787 **Additional information**
788 **Supplementary data** is available for this paper at https://
789

790 **Correspondence and requests for materials** should be addressed to Hitoshi
791 Sakaue.
792

References

References

1. Tansey, E.A., Johnson, C.D. (2015) Recent advances in thermoregulation.

Adv Physiol Educ. 39 (3):139-48

https://doi.org/10.1152/advan.00126.2014

2. Piscitiello, E., Herwig, A., Haugg, E., Schröder, B., Breves G., *et al*. (2021)

Acclimation of intestinal morphology and function in Djungarian hamsters

(Phodopus sungorus) related to seasonal and acute energy balance. J Exp Biol.

16;224(Pt 4): jeb232876

https://doi.org/10.1242/jeb.232876

3. Cannon, B., Nedergaard, J. (2004) Brown adipose tissue: function and

physiological significance. Physiol Rev. 84(1): 277-359

https://doi.org/10.1152/physrev.00015.2003

4. Nowack, J., Giroud, S., Arnold, W., Ruf, T. (2017) Muscle Non-shivering

Thermogenesis and Its Role in the Evolution of Endothermy. Front Physiol. 8:

889

https://doi.org/10.3389/fphys.2017.00889

5. Rowland, L. A., Bal, N. C., Kozak,L. P., Periasamy, M. (2015) Uncoupling

Protein 1 and Sarcolipin Are Required to Maintain Optimal Thermogenesis, and

Loss of Both Systems Compromises Survival of Mice under Cold Stress. J Biol

Chem. 8;290(19):12282-9

https://doi.org/10.1074/jbc.M115.637603

6. Golozoubova, V., Hohtola, E., Matthias, A., Jacobsson, A., Cannon, B.,

Nedergaard, J. (2001) Only UCP1 can mediate adaptive nonshivering

thermogenesis in the cold. FASEB J. 15(11):2048-50

https://doi.org/10.1096/fj.00-0536fje

7. Janovska, P., Zouhar, P., Bardova, K., Otahal, J., Vrbacky, M., *et al*. (2013)

Impairment of adrenergically-regulated thermogenesis in brown fat of obesity-

resistant mice is compensated by non-shivering thermogenesis in skeletal

muscle. Mol Metab. 69:101683

https://doi.org/10.1016/j.molmet.2023.101683

8. Blondin, D.P., Frisch, F., Phoenix, S., Guérin, B., Turcotte, É. E., *et al*. (2017)

Inhibition of Intracellular Triglyceride Lipolysis Suppresses Cold-Induced Brown

Adipose Tissue Metabolism and Increases Shivering in Humans. Cell Metab.

25(2): 438-447

https://doi.org/10.1016/j.cmet.2016.12.005

9. Bertile, F., Habold, C., Maho, Y. L., Giroud, S. (2021) Body Protein Sparing in

Hibernators: A Source for Biomedical Innovation. Front Physiol. 18:12:634953

https://doi.org/10.3389/fphys.2021.634953

10. Holeček, M. (2018) Branched-chain amino acids in health and disease:

metabolism, alterations in blood plasma, and as supplements. Nutr Metab

(Lond). 15:33

https://doi.org/10.1186/s12986-018-0271-1

11. Heine, M., Fischer, A.W., Schlein, C., Jung, C., Straub, L.G., *et al*. (2018)

Lipolysis Triggers a Systemic Insulin Response Essential for Efficient Energy

Replenishment of Activated Brown Adipose Tissue in Mice

Cell Metab. 28(4): 644-655.e4

https://doi.org/10.1016/j.cmet.2018.06.020

12. Padmini, P., Khedoe, S.J., Hoeke, G., Kooijman, S., Dijk, W. *et al*. (2015)

Brown adipose tissue takes up plasma triglycerides mostly after lipolysis. J Lipid

Res. 56(1): 51-9

https://doi.org/10.1194/jlr.M052746

13. Olsen, J.S., Aslund, A., Bokhari, M.H., Hutchison, D.S., Bengtsson, T.

(2019) Acute β-adrenoceptor mediated glucose clearance in brown adipose

tissue; a distinct pathway independent of functional insulin signaling. Mol Metab.

30:240-249

https://doi.org/10.1016/j.molmet.2019.10.004

14. Millis, E.L., Pierce, K.A., Jedrychowski, M.P., Garrity, R., Winther, S. *et al*.

(2018) Accumulation of succinate controls activation of adipose tissue

thermogenesis. Nature. 560(7716): 102-106

https://doi.org/10.1038/s41586-018-0353-2

15. Yoneshiro, T., Wang, Q., Tajima, K., Matsushita, M., Maki, H., *et al*. (2019)
BCAA catabolism in brown fat controls energy homeostasis through SLC25A44.
Nature. 572(7771): 614-619
https://doi.org/10.1038/s41586-019-1503-x

16. Yoneshiro, T., Kataoka, N., Walejko, J.M., Ikeda, K., Brown, Z. *et al*. (2021)
Metabolic flexibility via mitochondrial BCAA carrier SLC25A44 is required for
optimal fever. Elife. 10: e66865
https://doi.org/10.7554/elife.66865

17. Reddy, A., Bozi, L.H.M., Yaghi, O.K., Millis, E.L., Xiao, H., *et al*. (2020) pH-
Gated Succinate Secretion Regulates Muscle Remodeling in Response to
Exercise. Cell. 183(1):62-75. e17
https://doi.org/10.1016/j.cell.2020.08.039

18. de Oliveira Dos Santos A.R. de Oliveira Zanuso B. Miola V.F.B. *et al*. (2021)
Adipokines, Myokines, and Hepatokines: Crosstalk and Metabolic
Repercussions. Int J Mol Sci. 22(5): 2639
https://doi.org/10.3390/ijms22052639

19. Tanaka, T., Narazaki, M., Kishimoto, T. (2014) IL-6 in inflammation,
immunity, and disease. Cold Spring Harb Perspect Biol. 6(10): a016295
https://doi.org/10.3390/ijms22052639

20. Pedersen, B.K., Febbraio, M.A. (2008) Muscle as an Endocrine Organ:
Focus on Muscle-Derived Interleukin-6. Physiol Rev. 88(4): 1379-406
https://doi.org/10.1152/physrev.90100.2007

21. Qing, H., Desrouleaux, R., Israni-Winger, K., Mineur, Y.S., Fogelman, N., *et*
al. (2020) Origin and Function of Stress-Induced IL-6 in Murine Models. Cell.
182(2): 372-387.e14
https://doi.org/10.1016/j.cell.2020.05.054

22. Chen, Q., Wang, W. C., Bruce, R., Li, H., Schleider, D.M., Mulbury, M.J. *et*

al. (2004) Central role of IL-6 receptor signal-transducing chain gp130 in

activation of L-selectin adhesion by fever-range thermal stress. Immunity. 20(1):

59-70

https://doi.org/10.1016/s1074-7613(03)00358-3

23. Serrano, A. L., Baeza-Raja, B., Perdiguero, E., Muñoz-Cánoves, M.P.

(2008) Interleukin-6 Is an Essential Regulator of Satellite Cell-Mediated Skeletal

Muscle Hypertrophy. Cell Metab. 7(1): 33-44

https://doi.org/10.1016/j.cmet.2007.11.011

24. Ellingsgaard, H., Hauselmann, I., Schuler, B., Habib, A.M., Baggio, L.L., *et*

al. (2011) Interleukin-6 enhances insulin secretion by increasing glucagon-like

peptide-1 secretion from L cells and alpha cells. Nat Med. 17(11): 1481-9

https://doi.org/10.1038/nm.2513

25. Knudsen, J. G., Murholm, M., Carey, A.L., Beins, R.S., Basse, A.L.,et al.

(2014) Role of IL-6 in exercise training- and cold-induced UCP1 expression in

subcutaneous white adipose tissue. PLoS One. 9(1): e84910

https://doi.org/10.1371/journal.pone.0084910

26. Hall, G. V., Steensberg, A., Sacchetti, M., Fischer, C., Keller, C., *et al*.

(2003) Interleukin-6 stimulates lipolysis and fat oxidation in humans. J Clin

Endocrinol Metab. 88(7): 3005-10

https://doi.org/10.1210/jc.2002-021687

27. Nonogaki, K., Fuller, G.M., Fuentes, N.L., Moser, A.H., Staprans, I., *et al*.

(1995) Interleukin-6 stimulates hepatic triglyceride secretion in rats.

Endocrinology. 136(5): 2143-9

https://doi.org/10.1210/endo.136.5.7720663

28. Kawanishi, N., Funakoshi, T., Machida, S. (2018) Time-course study of

macrophage infiltration and inflammation in cast immobilization-induced

atrophied muscle of mice. Muscle Nerve. 57(6): 1006-1013

https://doi.org/10.1002/mus.26061

29. Bartness, T.J., Vaughan, C.H., Song, C.K. (2010) Sympathetic and sensory

innervation of brown adipose tissue. Int J Obes (Lond). 34 Suppl 1(0 1): S36-42

https://doi.org/10.1038/ijo.2010.182

30. Wang, Z., Wang, Q.A., Liu, Y., Jiang, L. (2021) Energy metabolism in brown

adipose tissue. FEBS J. 288(12): 3647-3662

https://doi.org/10.1111/febs.16015

31. Gazzola, R.F., Sala, R., Bussolati, O., Visigalli, R., Dall'Asta, V., *et al*. (2001)

The adaptive regulation of amino acid transport system A is associated to

changes in ATA2 expression. FEBS Lett. 490(1-2): 11-4

https://doi.org/10.1016/s0014-5793(01)02126-3

32. Gjymishka, A., Palii, S.S., Shan, J., Kilberg, M.S. (2008) Despite increased

ATF4 binding at the C/EBP-ATF composite site following activation of the

unfolded protein response, system A transporter 2 (SNAT2) transcription activity

is repressed in HepG2 cells. J Biol Chem. 283(41): 27736-27747

https://doi.org/10.1074/jbc.m803781200

33. HUANG, Y. and CHUANG, D.T. (1999) Down-regulation of rat mitochondrial

branched-chain 2-oxoacid dehydrogenase kinase gene expression by

glucocorticoids. Biochem. J. 339: 503–510

34. Kobayashi, R., Shimomura, Y., Murakami, T., Nakai, N., Otsuka, M., *et al*.

(1999) Hepatic branched-chain α-keto acid dehydrogenase complex in female

rats: activation by exercise and starvation. J Nutr Sci Vitaminol 45: 303-9

https://doi.org/10.3177/jnsv.45.303

35. Fukuhara, D., Kanai, Y., Chairoungdua, A., Babu, E., Bessho, F., *et al*.

(2007) Protein characterization of Na+-independent system l amino acid

transporter 3 in mice: a potential role in supply of branched-chain amino acids

under nutrient starvation. Am. J. Pathol. 170 (3): 888-89

https://doi.org/10.2353/ajpath.2007.060428

36. Rowland, L.A., Bal, N.C., Periasamy, M. (2015) The role of skeletal-muscle-

based thermogenic mechanisms in vertebrate endothermy. Biol Rev Camb

Philos Soc. 90(4): 1279-97

https://doi.org/10.1111/brv.12157

37. Zurlo, F., Larson, K., Bogardus, C., Ravussin, E. (1990) Skeletal muscle
metabolism is a major determinant of resting energy expenditure. J Clin Invest.
86(5): 1423-7
https://doi.org/10.1172/jci114857

38. Murashige, D., Jang, C., Neinast, M., Edwards, J.J., Cowan, A., *et al*. (2020)
Comprehensive quantification of fuel use by the failing and nonfailing human
heart. Science. 370(6514): 364-368
https://doi.org/10.1126/science.abc8861

39. Bonetto, A., Aydogdu, T., Jin, X., Zhang, Z., Zhan, R.,*et al*. (2012)
JAK/STAT3 pathway inhibition blocks skeletal muscle wasting downstream of
IL-6 and in experimental cancer cachexia. Am J Physiol Endocrinol Metab 303:
E410–E421
https://doi.org/10.1152/ajpendo.00039.2012

40. Zanders, L., Kny, M., Hahn, A., Schmidt, S., Wundersitz, S., *et al*. (2022)
Sepsis induces interleukin 6, gp130/JAK2/STAT3, and muscle wasting. J
Cachexia Sarcopenia Muscle. 2022; 13(1): 713-727
https://doi.org/10.1002/jcsm.12867

41. Hirata, Y., Nomura, K., Kato, D., Tachibana, Y., Niikura, T., *et al*. (2022) A
Piezo1/KLF15/IL-6 axis mediates immobilization-induced muscle atrophy. J Clin
Invest. 132(10): 1-13
https://doi.org/10.1172/jci154611

1000

1001   42. Bustamante, M., Fernández-Verdejo, R., Jaimovich, E., Buvinic, S. (2014)
1002   Electrical stimulation induces IL-6 in skeletal muscle through extracellular ATP
1003   by activating Ca (2+) signals and an IL-6 autocrine loop. Am J Physiol
1004   Endocrinol Metab. 306(8): E869-82
1005   https://doi.org/10.1152/ajpendo.00450.2013

1006

1007   43. Tomiya, S., Tamura, Y., Kouzaki, K., Kotani, T., Wakabayashi, Y., *et al*.
1008   (2019) Cast immobilization of hindlimb upregulates sarcolipin expression in

atrophied skeletal muscles and increases thermogenesis in C57BL/6J mice. Am
J Physiol Regul Integr Comp Physiol. 317(5): R649-R661
https://doi.org/10.1152/ajpregu.00118.2019

44. Poller, W.C., Downey, J., Mooslechner, A.A., Khan, N., Li, L., *et al*. (2022)
Brain motor and fear circuits regulate leukocytes during acute stress. Nature.
607(7919): 578-584
https://doi.org/10.1038/s41586-022-04890-z

45. Cheng, Y., Jope, R.S., Beurel, E. (2015) A pre-conditioning stress
accelerates increases in mouse plasma inflammatory cytokines induced by
stress. BMC Neurosci. 16:31
https://doi.org/10.1186/s12868-015-0169-z

46. Wallenius, V., Wallenius, K., Ahrén, B., Rudling, M., Carlsten, H., *et al*.
(2002) Interleukin-6-deficient mice develop mature-onset obesity. Nat Med. 8(1):
75-9
https://doi.org/10.1038/nm0102-75

47. Wang, T.J., Larson, M.G., Vasan, R.S., Cheng, S., Rhee, E.P. *et al*. (2011)
Metabolite profiles and the risk of developing diabetes. Nat Med. 17(4): 448-53
https://doi.org/10.1038/nm.2307

48. Shinoda, K., Luijten, I.H.N., Hasegawa, Y., Hong, H., Sonne, S.B., *et al*.
(2015) Genetic and functional characterization of clonally derived adult human
brown adipocytes. Nat Med. 21(4):389-94
https://doi.org/10.1038/nm.3819

49. Vaughan, C.H., Zarebidaki, E., Ehlen, J.C., Bartness, T.J. (2014) Analysis
and measurement of the sympathetic and sensory innervation of white and
brown adipose tissue. Methods Enzymol. 537: 199-225
https://doi.org/10.1016/b978-0-12-411619-1.00011-2

50. Okamatsu-Ogura, Y., Kuroda, M., Tsutsumi, R., Tsubota, A., Saito, M., *et al*.
(2020) UCP1-dependent and UCP1-independent metabolic changes induced by
acute cold exposure in brown adipose tissue of mice. Metabolism. 113: 154396

1045   https://doi.org/10.1016/j.metabol.2020.154396
1046
1047
1048

Figure Legends

Figure 1. Cast immobilization suppresses muscle thermogenesis and induces BAT thermogenesis via sympathetic activation

(A) Time course of rectal core body temperature of cast-immobilized (gypsum, n = 6) and control (n = 5) mice subjected to a cold (4°C) challenge test. The test was performed 7 days after bilateral immobilization of hind limbs.

(B) Rectal core body temperature of cast-immobilized (n = 5) and control (n = 5) mice during cold (4°C) exposure at 24 h after bilateral cast immobilization.

(C) Decline in rectal temperature of cast-immobilized and control (n = 9) mice after cold (4°C) exposure for 4 h performed 2 h after unilateral (UL, n =8) or bilateral (BL, n = 9) immobilization.

(D) Rectal core body temperature of cast-immobilized (n = 6) and control (n = 4) mice at room temperature after bilateral cast immobilization.

(E–G) Immunoblot analysis of PGC-1α (n = 3 or 4 per group) and UCP1 (n = 6 per group) in iBAT of control mice and mice subjected to bilateral cast immobilization for the indicated times. GAPDH was examined as a loading control. Representative blots (E) and densitometric quantitation of the PGC-1α/GAPDH (F) and UCP1/GAPDH (G) band intensity ratios are shown.

(H) Reverse transcription (RT) and real-time polymerase chain reaction (PCR) analysis of the *Ucp1/Gapdh* mRNA abundance ratio in iBAT of control mice and mice subjected to bilateral cast immobilization for the indicated times (n = 7 to 13 per group).

(I) Noradrenaline concentrations in iBAT of cast-immobilized (n = 4) and control (n = 4) mice at 24 h after subsequent biteral cast immobilization.

(J) RT and real-time PCR analysis of the *Ucp1/Gapdh* mRNA abundance ratio in iBAT subjected to surgical denervation or sham surgery, either in control mice or in mice at 24 h after subsequent bilateral cast immobilization (n = 6–9 per group).

(K) Rectal core body temperature of mice as in (I) subjected to cold (4°C) exposure for 1 h (n = 3 or 4 per group).

All quantitative data are means $\pm$ SEM. *p < 0.05, **p < 0.01 as determined by two-way ANOVA followed by Tukey's post hoc test or the post hoc paired/unpaired t test with Bonferroni's correction (A, B, D, and K), by one-way ANOVA followed by Tukey's post hoc test (C and J), by Dunnett's test (F-H), or by the unpaired t test (I)

See also Figure 1_figure supplement 1 and 2, and Table S1.

Figure 2. Cast immobilization alters systemic metabolic dynamics associated with BAT thermogenesis

(A and B) Concentrations of carbohydrate metabolites (A) and amino acids (B) in sham-operated or denervated iBAT of mice without (control) or at the indicated times after bilateral cast immobilization. The metabolomics data are presented as heat maps corresponding to the $\log_{10}$ value of fold change in cast-immobilized mice relative to the corresponding control mice and are means of four or five mice in each group. TCA, tricarboxylic acid cycle.

(C and D) RT and real-time PCR analysis of mRNA abundance for the fatty acid transporter CD36 and the succinate transporter SLC25A10 in sham-operated or denervated iBAT of control mice or mice subjected to bilateral cast immobilization for 24 h (n = 6 per group).

(E) Succinate content of sham-operated or denervated iBAT of control mice or mice subjected to bilateral cast immobilization for 24 h (n = 3–5 per group).

(F and G) Oxygen consumption rate (VO_2) and respiratory exchange ratio (RER), respectively, for control mice and mice subjected to bilateral cast immobilization for 7 days (n = 6 per group).

(H) Weight of eWAT for control mice and mice subjected to bilateral cast immobilization for the indicated times (n = 5–7 per group).

(I) Hepatic glycogen content for control mice and mice subjected to bilateral cast immobilization for the indicated times (n = 6–10 per group).

Date in (C) through (I) are means $\pm$ SEM. *$p < 0.05$, **$p < 0.01$ as determined by one-way ANOVA followed by Tukey's post hoc test (C–E), by two-way ANOVA followed by Tukey's post hoc test or the unpaired post hoc t test with Bonferroni's correction (F and G), or by Dunnett's test (H and I).

See also Figure 2_figure supplement 1.

Figure 3. Free amino acids are transferred from skeletal muscle to BAT and the liver for maintenance of energy homeostasis

(A and B) Metabolomics analysis of amino acid concentrations in soleus and EDL muscles (A) as well as in serum and the liver (B) of control mice and mice subjected to bilateral cast immobilization for the indicated times. Results are presented as heat maps of the $\log_{10}$ value of fold change for cast-immobilized mice relative to control mice and are means of three mice per group.

(C) RT and real-time PCR analysis of gene expression for SLC transporters and

mitochondrial BCAA catabolic enzymes in iBAT, liver, and soleus of control mice and mice subjected to bilateral cast immobilization for the indicated times. Results are presented as heat maps of the $\log_{10}$ value of fold change for cast-immobilized mice relative to control mice and are are shown individually for three to six mice of each group.

(D) Organ-specific [^{3}H]leucine uptake in control mice and mice subjected to bilateral cast immobilization for 24 h (n = 4–6 per group). sWAT, subcutaneous WAT.

(E) RT and real-time PCR analysis of SLC25A44 gene expression in sham-operated or denervated iBAT of control mice and mice subjected to subsequent bilateral cast immobilization for 24 h (n = 6 per group).

Data in (D) and (E) are means ± SEM. *$p < 0.05$, **$p < 0.01$ as determined by the unpaired t test (D) or by one-way ANOVA followed by Tukey's post hoc test (E).

See also Figure 3_figure supplement 1.

Figure 4. Cast immobilization upregulates *Il6* expression in BAT and skeletal muscle

(A and B) RT and real-time PCR analysis of IL-1β, IL-6, IL-10, IL-15, TNF-α, MCP-1, BMP8B, FGF21, and irisin gene expression in iBAT (A) or soleus (B) of control mice or mice at 10 h (iBAT) or 24 h (soleus) after bilateral cast immobilization (n = 5 or 6 per group).

(C and D) RT and real-time PCR analysis of *Il6* expression in iBAT (C) and soleus (D) of control mice and mice at the indicated times after bilateral cast immobilization (n = 3–7 per group).

(E–L) RT and real-time PCR analysis of *Saa3* (E, G, I, and K) and *Socs3* (F, H, J, and L) mRNA abundance in iBAT (E and F), soleus (G and H), liver (I and J), and eWAT (K and L) of control mice and mice at the indicated times after bilateral cast immobilization (n = 3–7 per group)

(M) RT and real-time PCR analysis of *Il6* expression, respectively, in denervated iBAT of control mice or mice at the indicated times after bilateral cast immobilization (n = 4-7 per group).

(N) Serum IL-6 concentration in control mice or mice at the indicated times after bilateral cast immobilization (n = 6 per group).

All data are means ± SEM. *$p < 0.05$, **$p < 0.01$, NS (not significant) as determined by the unpaired t test (A and B) or by Dunnett's test (C–N).

1157 See also Figure 4_figure supplement 1.

1158

Figure 5. IL-6 affects energy metabolism in BAT and skeletal muscle of
cast-immobilized mice
(A) Change in rectal body temperature during cold exposure at 4°C for 4 h for
WT (n = 4 or 5) and IL-6 KO (n = 4 or 5) mice that had been subjected (or not)
to bilateral cast immobilization for 24 h.
(B and C) Concentrations of carbohydrate metabolites (B) and amino acids (C)
in iBAT of IL-6 KO (–/–) and WT (+/+) mice without (control) or with bilateral cast
immobilization for the indicated times. Results are presented as heat maps of
the $\log_{10}$ value of the fold change relative to control WT mice and are means of
four mice in each group.
(D–F) RT and real-time PCR analysis of the expression of SLC25A44 (D),
BCKDHA (E), and CD36 (F) genes in iBAT of IL-6 KO and WT mice without
(control) or with bilateral cast immobilization for 24 h (n = 5 or 6).
(G) Incorporation of [^{3}H]leucine in iBAT of IL-6 KO and WT mice without
(control) or with bilateral cast immobilization for 24 h (n = 4 or 5 per group).
(H–J) BCAA (Val, Ile, and Leu, respectively) concentrations in soleus of IL-6 KO
and WT mice without (control) or with bilateral cast immobilization for 1 or 7
days (n = 4 per group).
Data in (A) and (D) through (J) are means $\pm$ SEM. *p < 0.05, **p < 0.01 as
determined by one-way ANOVA followed by Tukey's post hoc test (D–G), by
Dunnett's test (H–J), or by two-way ANOVA followed by Tukey's post hoc test
(A).
See also Figure 5_figure supplement 1.

1182

Figure 6. Skeletal-muscle-derived IL-6 increases BCAA concentrations in
skeletal muscle for support of BAT thermogenesis
(A–C) Concentrations of BCAAs (Val, Ile, and Leu, respectively) in the soleus of
control IL-6 KO mice and at 3 h after intraperitoneal administration of
recombinant IL-6 (rIL-6, 400 ng) or vehicle in IL-6 KO mice that had been
subjected to bilateral cast immobilization for 24 h (n = 4 or 5 per group).
(D) BCAA concentrations in C2C12 myotubes incubated in the absence or
presence of rIL-6 (50 ng/ml) for 1 h (n = 6 independent experiments) .
(E) Rectal core body temperature during cold exposure at 4°C for 1 h for control
IL-6 KO mice and for IL-6 KO mice subjected to bilateral cast immobilization for

24 h and injected intraperitoneally with rIL-6 (400 ng) or vehicle 30 min before

cold challenge (n = 5 per group).

(F) Rectal core body temperature during cold exposure at 4°C for 1 h for

C57BL/6J mice with denervated iBAT subjected or not to bilateral cast

immobilization and treatment with rIL-6 or vehicle as in (E) (n = 4 per group).

(G) Serum IL-6 concentration of mice subjected to surgical removal of iBAT and

then subjected (or not, control) to bilateral cast immobilization for the indicated

times (n = 4-6 per group).

(H) Weight of eWAT in mice subjected to surgical removal of iBAT (or to sham

surgery) followed (or not, control) by bilateral cast immobilization for 7 days (n =

5-7 per group).

(I-K) Concentrations of BCAAs (Val, Ile, and Leu, respectively) in soleus of mice

subjected to surgical removal of iBAT and then subjected (or not, control) to

bilateral cast immobilization for the indicated times (n = 4 per group).

(L) RT and real-time PCR analysis of *Il6* expression in iBAT, soleus, liver, eWAT,

and spleen of mice subjected (or not) to cold (4°C) exposure for 4 h (control, n =

6–9; cold exposure, n = 5 or 6).

(M and N) RT and real-time PCR analysis of *Saa3* (M) and *Socs3* (N) mRNA

abundance in soleus of mice subjected (or not) as (L) (control, n = 6; cold

exposure, n = 5).

(O) Amino acid concentrations in soleus of WT (+/+) and IL-6 KO (–/–) mice

subjected (or not) to cold exposure at 4°C for 4 h. Results are presented as

heat maps of the $\log_{10}$ value of fold change relative to WT control mice and are

means of four mice in each group.

(P) Valine concentration in soleus of WT and IL-6 KO mice subjected (or not) to

cold exposure at 4°C for 4 h (n = 4 per group).

All data other than those in (O) are means ± SEM. *$p < 0.05$, **$p < 0.01$, NS (not

significant) as determined by Dunnett's test (A–C and G and I-K), or by the

unpaired t test (D and L-N) or by one-way ANOVA followed by Tukey's post hoc

test (H and P), or by two-way ANOVA followed by the post hoc paired/unpaired t

test with Bonferroni's correction (E and F),

See also Figure 6_figure supplement 1 and 2.

1225

Fig.1

A Day 7

B Day 1

C Cold challenge

D Room temperature

E iBAT

F PGC-1α/GAPDH

G UCP1/GAPDH

H iBAT

I iBAT

J iBAT

K Cold challenge

Fig.2

A Carbohydrate metabolites in iBAT

Sham Denervated

Glucose 6-phosphate
Fructose 6-phosephate
Fructose 1,6-bisphosphate
3-Phosphoglyceric acid
2-Phosphoglyceric acid
Lactic acid
6-Phosphogluconic acid
Ribulose 5-phosphate
Sedoheptulose 7-phosphate
Citric acid
Succinic acid
Malic acid
Glucose 1-phosphate
Glucose 3-phosphate

Glycolytic pathway
Pentose phosphate pathway
TCA cycle
Glycogen metabolism
Lipid metabolism

Control Day 1 Day 3 Day 5

Log₁₀ (fold change)
-0.2   0.2

B Amino acids in iBAT

Sham Denervated

Gly
Ala
Ser
Pro
Val
Thr
Ile
Leu
Asn
Asp
Gln
Lys
Glu
Met
His
Phe
Arg
Tyr
Trp

Free amino acids

Control Day 1 Day 3 Day 5

Log₁₀ (fold change)
-0.15   0.15

Fig.3

Fig.4

Fig.5

Fig.6

Figure 1_figure supplement 2

A

Figure 4_figure supplement 1

Figure 6_figure supplement 1

Figure 6_figure supplement 2