1   Evolution of invertebrate *Cdk4/6-Cdkn2* system

Shiori Yuki[1], Shunsuke Sasaki[2], Yuta Yamamoto[2], Fumika Murakami[1], Kazumi Sakata[1,2]

and Isato Araki[1,2]* (1: Graduate School of Arts and Sciences, 2: Faculty of Science and

Engineering, Iwate University, Morioka, Japan)

* Corresponding author

Isato Araki

Faculty of Science & Engineering, Iwate University

3-18-33 Ueda, Morioka, Iwate 020-8550, Japan

phone/fax: +81-(0)19-621-6909

e-mail: iaraki@iwate-u.ac.jp

Short title: Evolution of invertebrate Cdk4/6-Cdkn2

Keywords: cell cycle regulation, comparative genomics, cyclin-dependent kinase,

cyclin-dependent kinase inhibitor, evolution, invertebrates, synteny, tumor suppressor

Abstract

Cell cycle is driven by cyclin-dependent kinase (Cdk). The decision whether the cell

cycle proceeds is made during G1 phase when Cdk4/6 functions. Cyclin-dependent

kinase inhibitor 2 (Cdkn2) is the specific inhibitor for Cdk4/6, and the interaction

depends on D84 in Cdkn2 and R24/31 in Cdk4/6. These knowledge have been obtained

mainly from studies in mammalian cells. Here, we comprehensively analyzed *Cdk4/6*

and *Cdkn2* in invertebrates, and found that while *Cdk4/6* exist in most of investigated

phyla, there is presence bias of *Cdkn2*. Interestingly, the positive charge in Cdk4/6

R24/R31 was conserved in all analyzed species in phyla with Cdkn2, and the presence

of *Cdkn2* and the conservation of the positive charge were statistically correlated. We

also found that *Cdkn2* has been tightly linked with *Fas associated factor 1* (*Faf1*) during

evolution. These genome-wide data suggest that 1) *Cdkn2* might have been lost multiple

times during evolution, 2) the interaction between Cdkn2 D84 and Cdk4/6 R24/31 might

impose constraint on the evolution of cell cycle regulation mechanisms, 3) the loss of

Cdkn2 might loosen the constraint, and 4) there might be potential interaction between

Cdkn2 and *Faf1*.

1   Introduction

Regulation of cell cycle is closely associated with ontogeny, tissue regeneration, cell

senescence, pathogenesis, especially carcinogenesis and evolution. Cyclin-dependent

kinases (Cdk's) and their regulatory subunits, cyclins, promote cell cycle while Cdk

inhibitors (Cdkn's) inhibit the cycle by repressing the function of Cdk's (Reviewed in

Weinberg, 2014). In mammalian cells Cdk4 or Cdk6, and cyclin D's drive cell cycle in

G1 phase, when cells are sensitive to extracellular regulatory signals on the proliferation.

Fish have Cdk21, the third member of Cdk4/6 subgroup of the Cdk family (Webster *et*

al., 2018). Mammals have three types of structurally different Cdkn's, Cdkn1, Cdkn2

and Cdkn3. They possess three Cdkn1's, Cdkn1a, Cdkn1b and Cdkn1c (aka $p21^{Cip1/Waf1}$,

$p27^{Kip1}$ and $p57^{Kip2}$, respectively; reviewed in Mühleder *et al*., 2021). Cdkn1's are

preferential inhibitors for Cdk1 and Cdk2, but can also inhibit Cdk4/6. Cdkn1's are

tumor suppressors or the candidate (El-Deiry *et al*., 1993; Chen *et al*., 1996; Zhang *et*

al., 1997).

Cdk inhibitor 2's (Cdkn2's) are specific inhibitors for Cdk4/6. Mammals possess four

Cdkn2's, Cdkn2a, Cdkn2b, Cdkn2c and Cdkn2d (aka $p16^{Ink4a}$, $p15^{Ink4b}$, $p18^{Ink4c}$ and $p18^{Ink4d}$,

respectively). Mammalian Cdkn2a and Cdkn2b possess four ankyrin repeats while

Cdkn2c and Cdkn2d do five (Baumgartner *et al*., 1998). Cdkn2's are known as tumor

suppressors (Kamb *et al*., 1994; Jen *et al*. 1994; Husemann *et al*., 1999; Gluick *et al*.,

2013) as Cdkn1's. It has been also shown that Cdkn2's accumulate in mammalian cells

during cell senescence (Serrano *et al*., 1997; Kita *et al*., 2022; Liu *et al*., 2023). It has

been thought that electrostatic interaction between R24/R31 of Cdk4/6 and D84 of

Cdkn2's plays a significant role in the formation of Cdk4/6-Cdkn2 complex, and the

Cdkn2 binding to R24/R31 causes structural change preventing cyclin and ATP binding

to Cdk4/6 and its following activation (Russo *et al*., 1998; Brotherton *et al*., 1998;

reviewed in Baker & Reddy 2013) *. R24C and R24H mutations in CDK4 has been

reported to cause familial melanoma, suggesting that strong positive charge at the residue

is necessary for the complex formation (Wölfel *et al*., 1995; Zuo *et al*., 1996; Tsao *et al*.,

1998; Soufir *et al*., 1998; Molven *et al*., 2005; Veinalde *et al*., 2013). This interaction

mechanism is also applied to a cell immortalization protocol, K4DT method (Shiomi *et

al*., 2011).

Cdkn3 (aka Cdk2-associated dual specificity phosphatase, Kap or Cip2) is a

phosphatase and specifically inhibits Cdk2 (reviewed in Patterson *et al*., 2009). It has

been suggested that Cdkn3 could act an oncogene, not a tumor suppressor gene, as

opposed to Cdkn1's and Cdkn2's (Liu *et al*., 2019). No direct interaction between Cdk4/6

and Cdkn3 has been reported.

In contrast to mammals the molecular mechanisms of cell cycle have been poorly

studied especially in invertebrates other than *Caenorhabditis elegans* and *Drosophila*

melanogaster (reviewed in Kipreos & van den Heuvel, 2019). *C. elegans* and *D.*

melanogaster have no Cdkn2 gene; instead, it is thought that the Cdkn1 subrogates the

Cdk4/6 function in G1 phase. A few pioneering small-scale genome studies reported the

existence of *Ckd4/6* genes in invertebrates Chordata (*Ciona intestinalis*, and

Branchiostoma floridae), Echinodermata (*Strongylocentrotus purpuratus*), Arthropoda

(*Drosophila melanogaster*), Platyhelminthes (*Schmidtea mediterranea*), Cnidaria

(*Nematostella vectensis* and *Hydra vulgaris*), and Placozoa (*Trichoplax adhaerens*)

while no *Ckd4/6* gene in Porifera (*Amphimedon queenslandica*) nor *Cdkn2* gene in

Platyhelminthes (*Schmidtea mediterranea*) and Cnidaria (*Hydra vulgaris*) (Cao *et al*.,

2014; Espinal-Centeno *et al*., 2020). However, considering the diversity of invertebrates

more systematic and comprehensive investigation on the regulatory mechanisms of the

cell cycle has been needed. From this point of view, we performed a large-scale

investigation on the existence of *Cdk4/6* and *Cdkn2* genes in invertebrates.

Footnote

The numbering of the amino acids is based on the human proteins. D84 in human

CDKN2A correspond to D86 in CDKN2B, D76 in CDKN2C and D80 in CDKN2D,

1 respectively. To avoid complexities, we use D84 for the D84/D86/D76/D80 in this paper.

Results

The presence of Cdkn2 in invertebrate was biased

First, we systematically searched *Cdkn2* in the genomes of all invertebrate phyla.

Even in the initial keyword and blastp searches followed by more thorough tblastn

searches on whole-genome shotgun contigs databases, we noticed that the presence bias

of *Cdkn2* among invertebrates. We concluded that *Cdkn2* was detected only in seven

phyla (Figures 1, 2, S1, Tables S1, S2). The species in the most of phyla in invertebrate

Deuterostomia, except in Urochordata and Xenacoelomorpha, possess *Cdkn2*. Some

phyla in Lophotrochozoa also possess *Cdkn2*; Mollusca, Annelida, and

Brachiopoda/Phoronida. In contrast we could not find *Cdkn2* in Ecdysozoa except in

Onychophora. We could not find the *Cdkn2* in the basal invertebrates, neither. We also

found that the *Cdkn2* presence in Mollusca is biased. For example, we could not detect

Cdkn2 in Bivalvia, Heterobranchia, and Coleoidea (Table 1).

During in the tblastn search we noticed that there were sporadic "weakly positive"

sequences in phyla where majority of the search result suggest that the phylum may lack

Cdkn2 (shown in red in Table S2b). Some of them apparently originated from

contaminated DNA of other species. To obtain more standards other than blast search

to distinguish whether the "weakly positive" sequences, excluding evident contaminated

DNA, are *bona fide* Cdkn2 or genes for other ankyrin repeat containing-proteins, we

first analyzed the gene structure, and found that the exon junction between Q50 and V51

is conserved in all species analyzed in this study. We also extracted conserved amino

acid residues among invertebrate Cdkn2's (Figure S1c). Based on the conserved exon

junction, presence of the conserved residue, and linkage to *Faf1* (see below) as well as

blast search results we concluded that the "weakly positive" sequences are not *bona fide*

Cdkn2 (for details, see the notes in Table S2b).

We also noticed that there are invertebrate sequences potentially misannotated as

Cdkn2 in the NCBI Protein database (Table S4). Some of them were apparently Cdkn1.

We judged that the sequences other than Cdkn1 homologs are not *bona fide* Cdkn2 but

other ankyrin repeat containing proteins based on the criteria mentioned above (for details,

see the notes in Table S4).

The corresponding residue to the D84 in Cdkn2 was conserved in most of invertebrates,

but not in Crinoid

Next, we investigated the evolutionary conservation of the corresponding residue to

Cdkn2 D84 in invertebrates. We found that it was conserved in most of (95.1%, 58/61)

the detected invertebrate *Cdkn2*; however, Cdkn2 sequences from two Crinoid species

(D84Q and D84N) as well as one of the two Cdkn2 in *Phoronis ovalis* (D84N) lacked

the negative charge (Figures 2, S1b), although their Cdk4/6's retained R24/R31 (see

below). In vertebrates as a control, the negative charge was conserved in almost all

Cdkn2 sequences (99.9% 1849/1850; Figure S3).

Cdkn2 gene has been strongly linked to Faf1 gene during evolution

We also analyzed the microsynteny around the *Cdkn2* locus in invertebrates. The

results showed that all analyzed invertebrate *Cdkn2* genes are linked with *Faf1* (*Fas*

associated factor 1) as same as vertebrate *Cdkn2c* (Figure 3, Tables S1, S5). In addition,

cephalochordate *Cdkn2* genes are linked with *Mtap* (*methylthioadenosine*

phosphorylase) as same as vertebrate *Cdkn2a/b* (Figure 3).

Microsynteny analysis around Faf1 locus in the species lacking Cdkn2

To obtain clues on the origin and the loss of *Cdkn2* we analyzed microsynteny around

Faf1 in the species without *Cdkn2*. While we could not detect any evolutionary

conservation of linkages comparable to the one between *Cdkn2* and *Faf1*, potentially

interesting is that the linkage between *Faf1* and *Kank1* (*KN Motif and Ankyrin Repeat*

Domains 1) is conserved among some species in Anthozoa of Cnidaria (Figure 4, Table

S6). Considering the evolutionary position of Cnidaria, *Cdkn2* in Bilateria and *Kank1* in

Anthozoa might share the origin in ancient metazoans.

Analysis of the presence of Cdkn1 in invertebrate phyla

It is thought that the Cdkn1 subrogates the Cdk4/6 function in G1 phase in *C. elegans*

and *D. melanogaster* (reviewed in Kipreos & van den Heuvel, 2019). A recent study

suggested that *Cdkn1* compensates the lack of *Cdkn2* also in an Urochordata *Ciona*

embryos (Kobayashi *et al*., 2022), while another study reported that planarians lack

Cdkn1's and Cdh1 is important for exiting cell cycle (Sato *et al*., 2022). To more precise

distribution of Cdkn1 among invertebrates we performed comprehensive search on

invertebrate genomes. We could find at least one *Cdkn1* gene in all invertebrate phyla

except Dicyemida and Orthonectida (Table 1; Table S7). In Platyhelminthes we could

find Cdkn1 in Cestoda, but not in Rhabditophora nor Trematoda.

All investigated invertebrates except Porifera, Dicyemida and Orthonectida *possessed*

Cdk4/6 gene

Then, we systematically searched *Cdk4/6* in invertebrate genomes. We could find

Cdk4/6 in all invertebrate phyla except in Porifera, as previously reported (Cao *et al*.,

2014), Dicyemida and Orthonectida (Table 1, Table S8). The lack of *Cdk4/6*, as well as

Cdkn1 mentioned above, in Dicyemida and Orthonectida may be due to the shrunk

genome size (Lu *et al.*, 2019; Slyusarev *et al.*, 2020). Thirty species including the ones in

Arachnida whose genome analyses suggested an ancient whole-genome duplication event

possess more than one *Cdk4/6* (Figure S3e, Table S8; Schwager *et al.*, 2017).

The extent of the evolutionary conservation of the corresponding residue to the R24/R31

in Cdk4/6 varied among phyla

Next, we investigated the evolutionary conservation of the corresponding residue to

the R24/R31 in invertebrates. Among 1296 investigated sequences of Cdk4/6 from 1262

invertebrate species in 20 phyla the positive charge in R24/R31 was conserved in 1148

sequences (88.6%) (Tables 2, S8). In 215 sequences (16.6%) the arginine was

substituted with lysine while the strong positive charge is lost in other 148 sequences

(11.4%). In invertebrate Deuterostomia two *Ciona* species in Urochordata and one

species in Xenacoelomorpha lacked the strong positive charge at the residue

corresponding to the Cdk4/6 R24/R31 (3/43, 7.0%) (Figures 5, S3a). This was especially

in contrast to *Corella inflata* and *Ascidia aspersa*, species belonging to the same order

Phlebobranchia as *Ciona*, in which the charge was conserved, although it has been

substituted with lysine (Shito & Hotta, personal communication). In Lophotrochozoa

39.1% (52/133) of the sequences investigated the strong positive charge was not

conserved (Table 2, Figure S3b, c). Strikingly, 97.9% (46/47) of the species in

Platyhelminthes lacked the charge and the corresponding residues are mostly threonine's.

In Ecdysozoa 8.0% (85/1058) of the sequences investigated the charge was not

conserved (Table 2, Figure S3d, e). A significant ratio of species (63.9%, 46/72) in

Nematoda also lacked the charge and the corresponding residues are mostly, again,

threonine's. In basal invertebrates 9.7% (6/62) of the sequences investigated the strong

charge was not conserved (Table 2, Figure S3f). These data in invertebrate Cdk4/6 are

contrast to the one in vertebrate Cdk4/6/21, in which the charge is conserved in 1249

out of 1252 investigated sequences (99.8%) from 572 species (Figure S4, Table S9).

Comparison between the presence bias of Cdkn2 and the differential conservation of the

corresponding residue to the R24/R31 in Cdk4/6 among phyla

While species in most of animal phyla possess *Cdk4/6*, the extent of the evolutionary

conservation of the corresponding residue to the R24/R31 greatly differed among phyla.

Since we speculated that the presence of *Cdkn2* might affect the extent, we compared

them in each phylum, and summarized the result in Figure 1. In all analyzed species in

phyla with *Cdkn2* gene(s) the strong positive charge in the corresponding residue to the

R24/R31 of Cdk4/6 homologs was conserved, even in the two species in Crinoid where

the negative charge in the corresponding residue to Cdkn2 D84 has been lost. On the other

hand, the evolutionary conservation of the strong positive charge in the Cdk4/6 homologs

in phyla where we could detect no *Cdkn2* gene showed extreme difference. For example,

while only small number of species in Cnidaria, Arthropoda, and Urochordata lacked the

strong positive charge (10.0%, 3.9%, 14.3%, respectively), many or almost all species in

Nematoda, Rotifera and Platyhelminthes seemed to lack the strong positive charge

(63.9%, 44.4%, 97.9%, respectively). It is interesting that many species in the

evolutionary distant two phyla, Platyhelminthes and Nematoda, both of which contains

large numbers of parasites, share a threonine instead of R24/R31, although the

significance is unknown.

Finally, we analyzed the statistical correlation between the presence of *Cdkn2* and the

conservation of the strong positive charge in the corresponding residue at the

corresponding residue to the R24/R31 in Cdk4/6. We performed the analysis on the nine

phyla in which more than five species were included in this study (highlighted yellow in

the Figure 1). We used the percentage of the number of species in which both Cdkn2 and

Cdk4/6 to the number of species in which Cdk4/6 was detected ("**[2]/[1]** in %" in Figure

1   S5) , and the percentage of number of species in which the strong positive charge was

2   conserved in each of the phyla ("% of -R, K at Cdk4/6 R24/31" in Figure S5) to determine

3   the Spearman's rank correlation coefficient. The coefficient was 0.816 (P value =

4   0.00735), suggesting that the presence of *Cdkn2* and the conservation of the positive

5   charge were strongly correlated with statistical significance.

Discussion

Evolution of Cdkn2 gene

Based on the data on the presence bias of *Cdkn2* in invertebrate genomes, we postulate

a scenario of the evolution of *Cdkn2* (Figure 6a). In this scenario the ancestral *Cdkn2* gene

first appears from one of the ankyrin repeat-containing genes during the early evolution

of Bilateria. During the diversification of Bilateria *Cdkn2* gene has been lost in some

phyla, although the extent differs among Ecdysozoa, Lophotrochozoa, and Deuterostomia.

The observation that some invertebrates possess *Cdkn2* while the other not might be

obscurely explained that each of the strategies are beneficial to survive or reproduce more

efficiently in each group. Since Cdkn2's regulate the cell cycle in mammals, the cause of

the presence bias of *Cdkn2* in invertebrates might be related to cell proliferation, cell

differentiation or cell aging. However, it should be noted that even in deuterostomal

invertebrates the mechanisms of cell aging seem to be different from the ones in mammals

(reviewed in Amir *et al*., 2020). Species in phyla without *Cdkn2*, like Platyhelminthes,

Nemertea, Bryzoa and Urochordata as well as non-Bilateria, are known to their extreme

regenerative ability; however, species in phyla with *Cdkn2*, like Phoronida, Annelida and

Echinodermata, are also the same (reviewed in Bely & Nyberg, 2010). Thus, *Cdkn2* is

not another example of genes like JmjC domain encoding gene which has been shown to

be highly linked to extreme regeneration ability (Cao *et al.*, 2019). It seems that longevity

may not the cause of the presence bias of *Cdkn2* in invertebrates because even if you

exclude non-Bilateria which has species with ultra-extreme longevity or theoretical

immortality (Jochum *et al.*, 2012; Piraino *et al.*, 1996; Brien, 1953), both a sea urchin,

Mesocentrotus franciscanus (aka *Strongylocentrotus franciscanus*), in which we could

detect a *Cdkn2* gene (Figure S1), and a clam, *Arctica islandica*, which belong to Bivalvia

where we could not detect *Cdkn2* (Table 1), are known to their longevity, although the

latter nuclear genome has not yet been published (Ebert & Southon, 2003; Schöne *et al.*,

2005). Thus, we have currently no compelling clear-cut hypothesis to explain this issue.

Strong evolutionary conservation of the microsynteny in Cdkn2-Faf1 locus

Our synteny analysis revealed that *Cdkn2* has been strongly linked with *Faf1*. Based

on the data we postulate a scenario of the evolution of the linkage around Cdkn2 locus in

invertebrate (Figure 6b). In this scenario the ancestral *Cdkn2* gene first appeared linking

with *Faf1*. In the ancestral chordate the *Cdkn2* gene has begun to link also with *Mtap*.

During the twice duplication of the genome in the ancestral vertebrate one of the *Cdkn2*,

i.e., *Cdkn2c*, kept linked with *Faf1* while another one, i.e., *Cdkn2b*, kept linked with *Mtap*.

One of other *Cdkn2*'s became *Cdkn2d*, and the other one "*Cdkn2e*" was lost. Later, the

duplication of *Cdkn2b* created *Cdkn2a*. It is interesting that in human *Faf1*, *Mtap* and

Cdkn2 are all known as tumor suppressors, and the genetic linkage between *Cdkn2* and

Faf1 among them can be traced back into the appearance of *Cdkn2* (Christopher *et al*.,

2002; Menges *et al*., 2009; Kamb *et al*., 1994; Jen *et al*., 1994; Husemann *et al*., 1999;

Gluick *et al*., 2013).

Why has the microsynteny in *Cdkn2-Faf1* locus been conserved during bilaterian

evolution? It has been suggested that the microsynteny in a locus with multiple genes may

be conserved by *cis*-regulatory constraints between the genes and/or the functional

interdependency of the gene products (Imai *et al*., 2011; Irimia *et al*., 2012; Simakov &

Kawashima, 2017; Cai *et al*., 2010). Although both *Cdkn2*'s and *Faf1* are categorized to

tumor suppressor genes in mammals as mentioned above, they function in different

cellular systems, i.e., Cdkn2's regulate cell cycle while Faf1 does apoptosis, and their

physical interaction has not yet been reported. Thus, *cis*-regulatory constraints between

them might keep the microsynteny. In that context it is interesting that the expression of

a lncRNA gene, which is on the immediate opposite side of *C1orf185* gene from *Cdkn2c*,

can affect *Faf1* expression in *Homo sapiens*, although the linkage between *C1orf185* and

Cdkn2c-Faf1 locus is not evolutionary conserved even in vertebrates (Wan *et al*., 2014;

Araki, unpublished data). However, it is also noteworthy that a conservation of

microsynteny might reflect a covert physical interaction (Moretti *et al*., 2017).

Potential interactions between Cdkn2 and Cdk4/6 evolutions

The results of the comparative analysis between the presence bias of *Cdkn2* and the differential conservation of the strong positive charge at the corresponding residue to the R24/R31 in Cdk4/6 among phyla suggest potential interactions between *Cdkn2* and *Cdk4/6* evolutions. The fact all investigated species in Annelida, Mollusca and Echinodermata keep the charge and possess *Cdkn2* while all investigated species in Platyhelminthes, Rotifera and Nematoda lack *Cdkn2* and high percentage of the species also lack the charge imply that the existence of *Cdkn2* might constrain the corresponding residue to the R24/R31 from amino acid substitution or loss.

On the other hand, most of the species in Cnidaria, Arthropoda and Urochordata keep the charge while they lack *Cdkn2*. Furthermore, Cdkn2 sequences from two Crinoid species lack the negative charge in the corresponding residue to D84, although their Cdk4/6's retain R24/R31. In these species the Cdk4/6 sequences might be still on the way to amino acid substitution. Alternatively, the strong positive charge in the Cdk4/6 homologs may be conserved for the maintenance of the three-dimensional structure, and/or protein interaction with other than Cdkn2.

Cell cycle regulation without Cdkn2

What substitutes for Cdkn2 in the species without *Cdkn2*? It has been shown that

Cdkn1 is a preferential inhibitor for Cdk1/2, but can also inhibit Cdk4/6 in mammalian

cells, although the interaction between Cdk4/6 and Cdkn1 has not yet been solved by X-

ray crystallography (reviewed in Wood & Endicott, 2018, and Mühleder *et al.*, 2021).

Thus, it is easy to speculate that Cdkn1 substitutes for Cdkn2 in the species without *Cdkn2*,

and, in fact, it has been reported that Cdkn1 can substitute for Cdkn2 in some invertebrates

including *Ciona* as mentioned above (Kobayashi *et al.*, 2022; reviewed in Kipreos & van

den Heuvel, 2019). However, it might be necessary to consider cautious to extrapolate

this mechanism to other urochordates, or, species lacking *Cdkn2* in other phyla because

Ciona is the only urochordates investigated lacking the strong positive charge at the

residue corresponding to Cdk4/6 R24/R31. Since the extent of the evolutionary

conservation of the positive charge greatly differs among phyla lacking *Cdkn2*, the

conserved strong positive charge in Cnidaria, Arthropoda and Urochordata might not be

involved in the potential interaction with Cdkn1, and the positive charge might be

necessary for interaction with other than Cdkn2 as mentioned above.

Conclusion

In this research we performed the first comprehensive investigations on the distribution

of *Cdk4/6* and *Cdkn2* in invertebrate genomes and the evolutionary conservation of the

key charged residue as well as the synteny analysis on *Cdkn2*. We believe that the result

should widen our knowledge on the mechanisms of the cell cycle regulation and their

evolution in Metazoan.

Experimental procedures

Taxonomy

We used the taxonomy of invertebrates by Telford *et al*. (2015) for phyla, and the one

in the NCBI Taxonomy database for subclassification in this study. There has been

controversy on the evolutionary position on Xenacoelomorpha (Cannon *et al*., 2016;

Philippe *et al*., 2019). In the paper we tentatively categorize it as a diverged deuterostome.

Database search

To analyze the presence of *Cdkn2* and *Cdk4/6* genes, and to obtain their amino acid

sequences, we performed keyword and blastp search in the NCBI Protein database,

Ensembl Metazoa, Ensembl Rapid Release, Ghost Database: *Ciona intestinalis* genomic

and cDNA resources, Aniseed - Ascidian Network for In Situ Expression and

Embryological Data, PlanMine and OIST Marine Genomics Unit Genome Browser

(Sayers *et al*., 2024; Yates *et al*., 2022; Satou *et al*., 2005; Dardaillon *et al*., 2020;

Rozanski *et al*., 2019; Shinzato *et al*., 2020, 2021; Yoshioka *et al*., 2022; Khalturin *et*

al., 2019; Luo *et al*., 2018; Takeuchi *et al*., 2022; Arimoto *et al*., 2019; Simakov *et al*.,

2015). We also performed tblastn search in the NCBI Whole-genome shotgun contigs

Database. To confirm the identity of the sequences obtained as results, we performed

blastp search against vertebrate sequences. We also collected vertebrate *Cdkn2* and

Cdk4/6/21 sequences as controls. During the process we noticed that there are

misannotations on the vertebrate genes in the NCBI databases. For example, a platypus

Cdkn2 (XP_028910528.1) is annotated as "cyclin-dependent kinase 4 inhibitor B-like",

but the microsynteny suggests that it is Cdkn2a; thus, we treated it as Cdkn2a in this

paper. Since 1) the paralogs are specific to vertebrates, thus, the misannotation does not

affect the results in this study, and 2) it is out of scope in this paper, we will publish the

details of the analysis elsewhere. The database ID of the analyzed sequences were

tabulated in the Tables S1, S2, S8, S9).

As the result of the search 61 Cdkn2 sequences from 61 invertebrate species, and

1330 Cdk4/6 sequences from 1264 invertebrate species were included in this analysis.

In addition, 1850 Cdkn2 sequences and 1252 Cdk4/6/21 sequences from 572 vertebrate

species were included as controls in the analysis.

Protein sequence multiple alignment

We used NCBI COBALT (Constraint-based Multiple Alignment Tool) for protein

sequence multiple alignment (Papadopoulos & Agarwala, 2007). When the alignment

was questionable, we corrected it manually. We used WebLogo

(https://weblogo.berkeley.edu/logo.cgi) to create a consensus sequence logo, and

MOTIF Search (https://www.genome.jp/tools/motif/) to identify protein motives.

Synteny analysis

We used the NCBI gene database for synteny analysis if the sequence had a NCBI

gene ID (Brown *et al*., 2015). If the sequence had no gene ID, and the long enough

genomic sequence was available, we analyzed the annotation, if available, or performed

blastp search against the surrounding sequence. If the surrounding genes have been

annotated, we performed blastp against vertebrates to confirm the identities. We looked

up GeneCards (https://www.genecards.org) for abbreviation of gene name.

Statistical analysis

We used a statistical software EZR to determine Spearman's rank correlation

coefficient and create the scatter plot (Kanda, 2013).

Acknowledgement

We thank Drs. Yutaka Satou (Kyoto Univ) and Takeshi Kawashima (Inuyama Bio) for

critically reading the manuscript, and Takuya Minokawa (Tohoku Univ), Kazuya

Kobayashi (Hirosaki Univ), Hiroshi Wada (Tsukuba Univ), and Koji Tamura (Tohoku

Univ) for helpful discussion. We also thank Drs. Takumi Shito and Kohji Hotta (Keio

Univ) for sharing their unpublished data. This research was financially supported by

Iwate University.

References

Amir, Y., Insler, M., Giller, A., Gutman, D. & Atzmon, G. (2020). Senescence and

Longevity of Sea Urchins. *Genes* (Basel), *11*, 573.

https://doi.org/10.3390/genes11050573.

Arimoto, A., Hikosaka-Katayama, T., Hikosaka, A., Tagawa, K., Inoue, T., Ueki, T., ...

Satoh, N. (2019). A draft nuclear-genome assembly of the acoel flatworm

Praesagittifera naikaiensis. *Gigascience*, 8, 1-8.

https://doi.org/10.1093/gigascience/giz023.

Baker, S. J. & Reddy, E. P. (2013). CDK4: A Key Player in the Cell Cycle, Development,

and Cancer. *Genes and cancer*, *3*, 658-669.

https://doi.org/10.1177/1947601913478972.

Baumgartner, R., Fernandez-Catalan, C., Winoto, A., Huber, R., Engh, R. A. & Holak, T.

A. (1998). Structure of human cyclin-dependent kinase inhibitor p19[INK4d]: comparison

to known ankyrin-repeat-containing structures and implications for the dysfunction of

tumor suppressor p16[INK4a]. *Structure*, *6*, 1279-1290. https://doi.org/10.1016/s0969-

2126(98)00128-2.

Bely, A. E. & Nyberg, K. G. (2010). Evolution of animal regeneration: re-emergence of

a field. *Trends in ecology and evolution*, *25*, 161-170.

https://doi.org/10.1016/j.tree.2009.08.005.

Brien, P. (1953). La pérennité somatique. *Biological Reviews*, *28*, 308-349.

https://doi.org/10.1111/j.1469-185x.1953.tb01381.x.

Brotherton, D. H., Dhanaraj, V., Wick, S., Brizuela, L., Domaille, P. J., Volyanik, E., ...

Laue, E. D. (1998). Crystal structure of the complex of the cyclin D-dependent kinase

Cdk6 bound to the cell-cycle inhibitor p19[INK4d]. *Nature*, *395*, 244-250.

https://doi.org/10.1038/26164.

Brown, G. R., Hem, V., Katz, K. S., Ovetsky, M., Wallin, C., Ermolaeva, O., ... Murphy,

T. D. (2015). Gene: a gene-centered information resource at NCBI. *Nucleic Acids Res*,

43, D36-42. https://doi.org/10.1093/nar/gku1055.

Cai, W. W., Wang, L. & Chen, Y. (2010). Aspartyl aminopeptidase, encoded by an

evolutionarily conserved syntenic gene, is colocalized with its cluster in secretory

granules of pancreatic islet cells. *Bioscience, Biotechnology, and Biochemistry*, *74*,

2050-2055. https://doi.org/10.1271/bbb.100349.

Cannon, J. T., Vellutini, B. C., Smith, 3rd, J., Ronquist, F., Jondelius, U. & Hejnol, A.

(2016). Xenacoelomorpha is the sister group to Nephrozoa. *Nature*, *530*, 89-93.

https://doi.org/10.1038/nature16520.

Cao, L., Chen, F., Yang, X., Xu, W., Xie, J. & Yu, L. (2014). Phylogenetic analysis of

CDK and cyclin proteins in premetazoan lineages. *BMC Ecology and Evolution*, *14*,

10. https://doi.org/10.1186/1471-2148-14-10.

Cao, P.-L., Kumagai, N., Inoue, T., Agata, K. & Makino, T. (2019). JmjC Domain-

Encoding Genes Are Conserved in Highly Regenerative Metazoans and Are

Associated with Planarian Whole-Body Regeneration. Genome Biology and Evolution,

11, 552-564. https://doi.org/10.1093/gbe/evz021.

Chen, J., Willingham, T., Shuford, M. & Nisen, P. D. (1996). Tumor suppression and

inhibition of aneuploid cell accumulation in human brain tumor cells by ectopic

overexpression of the cyclin-dependent kinase inhibitor p27^{KIP1}. *Journal of Clinical*

Investigation, *97*, 1983-1988. https://doi.org/10.1172/JCI118631.

Christopher, S. A., Diegelman, P., Porter, C. W. & Kruger, W. D. (2002).

Methylthioadenosine phosphorylase, a gene frequently codeleted with p16$^{cdkN2a/ARF}$,

acts as a tumor suppressor in a breast cancer cell line. Cancer Research, 62, 6639-6644.

Dardaillon, J., Dauga, D., Simion, P., Faure, E., Onuma, T. A., DeBiasse, M. B., ...

Lemaire, P. (2020). ANISEED 2019: 4D exploration of genetic data for an extended

range of tunicates. *Nucleic Acids Res*, 48, D668-D675. https://doi.org/

10.1093/nar/gkz955.

Ebert, T. A. & Southon, J. R. (2003). Red sea urchins (*Strongylocentrotus franciscanus*)

can live over 100 years: Confirmation with A-bomb ^{14}carbon. *Fishery Bulletin*, *101*,

915-922.

El-Deiry, W. S., Tokino, T., Velculescu, V. E., Levy, D. B., Parsons, R., Trent, J. M., ...

Vogelstein, B. (1993). *WAF1*, a potential mediator of p53 tumor suppression. *Cell*, *75*,

817-825. https://doi.org/10.1016/0092-8674(93)90500-p.

Espinal-Centeno, A., Dipp-Álvarez, M., Saldaña, C., Bako, L. & Cruz-Ramírez, A.

(2020). Conservation analysis of core cell cycle regulators and their transcriptional

behavior during limb regeneration in *Ambystoma mexicanum*. *Mechanisms of*

Development, *164*, 103651. https://doi.org/10.1016/j.mod.2020.103651.

Gluick, T., Yuan, Z., Libutti, S. K. & Marx, S. J. (2013). Mutations in *CDKN2C* (*p18*)

and *CDKN2D* (*p19*) may cause sporadic parathyroid adenoma. *Endocrine-Related*

Cancer, *20*, L27-29. https://doi.org/10.1530/ERC-13-0445.

Husemann, K., Wolter, M., Büschges, R., Boström, J., Sabel, M. & Reifenberger, G.

(1999). Identification of two distinct deleted regions on the short arm of chromosome

1 and rare mutation of the *CDKN2C* gene from 1p32 in oligodendroglial tumors.

Journal of Neuropathology & Experimental Neurology, *58*, 1041-1050.

Imai, K. S., Daido, Y., Kusakabe, T. G. & Satou Y. (2012) Cis-acting transcriptional

repression establishes a sharp boundary in chordate embryos. *Science*, 337, 964-967.

https://doi.org/10.1126/science.1222488.

Irimia, M., Tena, J. J., Alexis, M. S., Fernandez-Miñan, A., Maeso, I., Bogdanovic, O., ...

Fraser, H. B. (2012) Extensive conservation of ancient microsynteny across metazoans

due to cis-regulatory constraints. *Genome Research*, *22*, 2356-2367.

https://doi.org/10.1101/gr.139725.112.

Jen, J., Harper, J. W., Bigner, S. H., Bigner, D. D., Papadopoulos, N., Markowitz, S., ...

Vogelstein, B. (1994). Deletion of *p16* and *p15* genes in brain tumors. *Cancer Research*,

54, 6353-6358.

Jochum, K. P., Wang, X., Vennemann, T. W., Sinha, B. & Müller, W. E. G. (2012)

Siliceous deep-sea sponge *Monorhaphis chuni*: A potential paleoclimate archive in

ancient animals. *Chemical Geology*, *300-301*, 143–151.

https://doi.org/10.1016/j.chemgeo.2012.01.009.

Khalturin, K., Shinzato, C., Khalturina, M., Hamada, M., Fujie, M. Koyanagi, R., ... Satoh,

N. (2019). Medusozoan genomes inform the evolution of the jellyfish body plan.

Nature Ecology & Evolution, 3, 811-822. https://doi.org/10.1038/s41559-019-0853-y.

Kamb, A., Gruis, N. A., Weaver-Feldhaus, J., Liu, Q., Harshman, K., Tavtigian, S. V., ...

Skolnick, M. H. (1994). A cell cycle regulator potentially involved in genesis of many

tumor types. *Science*, *264*, 436-440. https://doi.org/10.1126/science.8153634.

Kanda, Y. (2013). Investigation of the freely available easy-to-use software 'EZR' for medical statistics. *Bone Marrow Transplantation*, 48, 452–458. https://doi.org/10.1038/bmt.2012.244.

Kipreos, E. T. & van den Heuvel, S. (2019). Developmental Control of the Cell Cycle: Insights from *Caenorhabditis elegans*. *Genetics*, *211*, 797-829. https://doi.org/10.1534/genetics.118.301643.

Kita, A., Saito, Y., Miura, N., Miyajima, M., Yamamoto, S., Sato, T., ... Chikenji, T. S. (2022). Altered regulation of mesenchymal cell senescence in adipose tissue promotes pathological changes associated with diabetic wound healing. *Communications Biology*, *5*, 310. https://doi.org/10.1038/s42003-022-03266-3.

Kobayashi, K., Tokuoka, M., Sato, H., Ariyoshi, M., Kawahara, S., Fujiwara, S., ... Satou, Y. (2022). Regulators specifying cell fate activate cell cycle regulator genes to determine cell numbers in ascidian larval tissues. *Development*, *149*, dev201218. https://doi.org/10.1242/dev.201218.

Liu, J., Min, L., Zhu, S., Guo, Q., Li, H., Zhang, Z., ... Zhang, S. (2019). Cyclin-Dependent Kinase Inhibitor 3 Promoted Cell Proliferation by Driving Cell Cycle from G1 to S Phase in Esophageal Squamous Cell Carcinoma. *Journal of Cancer*, *10*, 1915-1922. https://doi.org/10.7150/jca.27053.

Liu, J., Lin, X., McDavid, A., Yang, Y., Zhang, H., Boyce, B. F. & Xing, L. (2023).

Molecular signatures distinguish senescent cells from inflammatory cells in aged

mouse callus stromal cells. *Frontiers in Endocrinology* (Lausanne), *14*, 1090049.

https://doi.org/10.3389/fendo.2023.1090049.

Lu, T.-M., Kanda, M., Furuya, H. & Satoh, N. (2019). Dicyemid Mesozoans: A Unique

Parasitic Lifestyle and a Reduced Genome. *Genome Biology and Evolution*, *11*, 2232-

2243. https://doi.org/10.1093/gbe/evz157.

Luo, Y.-J., Kanda, M., Koyanagi, R., Hisata, K., Akiyama, T., Sakamoto, H., ... Satoh, N.

(2018). Nemertean and phoronid genomes reveal lophotrochozoan evolution and the

origin of bilaterian heads. *Nature Ecology & Evolution*, 2, 141-151.

https://doi.org/10.1038/s41559-017-0389-y.

Menges, C. W., Altomare, D. A. & Testa, J.R. (2009). FAS-associated factor 1 (FAF1):

diverse functions and implications for oncogenesis. *Cell Cycle*, *8*, 2528-2534.

https://doi.org/10.4161/cc.8.16.9280.

Molven, A., Grimstvedt, M. B., Steine, S. J., Harland, M., Avril, M.-F., Hayward, N. K.

& Akslen, L. A. (2005). A large Norwegian family with inherited malignant melanoma,

multiple atypical nevi, and *CDK4* mutation. *Genes, Chromosomes & Cancer*, *44*, 10-

18. https://doi.org/10.1002/gcc.20202.

Moretti, A. I. S., Pavanelli, J.C., Nolasco, P., Leisegang, M. S., Tanaka, L. Y., Fernandes,

C.G., ... Laurindo, F. R. M. (2017). Conserved Gene Microsynteny Unveils Functional

Interaction Between Protein Disulfide Isomerase and Rho Guanine-Dissociation

Inhibitor Families. *Scientific Reports*, *7*, 17262. https://doi.org/10.1038/s41598-017-

16947-5.

Mühleder, S., Fernández-Chacón, M., Garcia-Gonzalez, I. & Benedito, R. (2021).

Endothelial sprouting, proliferation, or senescence: tipping the balance from

physiology to pathology. *Cellular and Molecular Life Sciences*, *78*, 1329-1354.

https://doi.org/10.1007/s00018-020-03664-y.

Papadopoulos, J. S, & Agarwala, R. (2007). COBALT: constraint-based alignment tool

for multiple protein sequences. *Bioinformatics*, 23, 1073-1079.

Patterson, K. I., Brummer, T., O'Brien, P. M. & Daly, R. J. (2009). Dual-specificity

phosphatases: critical regulators with diverse cellular targets. *Biochemical Journal*,

418, 475-489. https://doi.org/10.1042/bj20082234.

Philippe, H., Poustka, A. J., Chiodin, M., Hoff, K. J., Dessimoz, C., Tomiczek, B., ...

Telford, M. J. (2019). Mitigating Anticipated Effects of Systematic Errors Supports

Sister-Group Relationship between Xenacoelomorpha and Ambulacraria. *Current*

Biology, *29*, 1818-1826.e6. https://doi.org/10.1016/j.cub.2019.04.009.

Piraino, S., Boero, F., Aeschbach, B. & Schmid, V. (1996). Reversing the Life Cycle: Medusae Transforming into Polyps and Cell Transdifferentiation in *Turritopsis nutricula* (Cnidaria, Hydrozoa). *Biology Bulletin*, *190*, 302-312. https://doi.org/10.2307/1543022.

Rozanski, A., Moon, H., Brandl, H., Martín-Durán, J. M., Grohme, M. A., Hüttner, K., ... Rink, J. C. (2019). PlanMine 3.0-improvements to a mineable resource of flatworm biology and biodiversity. *Nucleic Acids Res*, 47, D812-D820. https://doi.org/10.1093/nar/gky1070.

Russo, A. A., Tong, L., Lee, J. O., Jeffrey, P. D. & Pavletich, N. P. (1998). Structural basis for inhibition of the cyclin-dependent kinase Cdk6 by the tumour suppressor p16[INK4a]. *Nature*, *395*, 237-243. https://doi.org/10.1038/26155.

Sato, Y., Umesono, Y., Kuroki, Y., Agata, K. & Hashimoto, C. (2022). Proliferation maintains the undifferentiated status of stem cells: The role of the planarian cell cycle regulator Cdh1. Developmental Biology, 482, 55–66. https://doi.org/10.1016/j.ydbio.2021.12.006

Satou, Y., Kawashima, T., Shoguchi, E., Nakayama, A. & Satoh, N. (2005). An Integrated Database of the Ascidian, *Ciona intestinalis*: Towards Functional Genomics. *Zool Sci*, 22, 837-843. https://doi.org/10.2108/zsj.22.837.

Sayers, E. W., Beck, J., Bolton, E. E., Brister, J. R., Chan, J., Comeau, D. C., ... Sherry, S. T. (2024). Database resources of the National Center for Biotechnology Information. *Nucleic Acids Res*, 52, D33-D43. https://doi.org/10.1093/nar/gkad1044.

Schöne, B. R., Fiebig, J., Pfeiffer, M., Gless, R., Hickson, J., Johnson, A. L. A., ... Oschmann, W. (2005). Climate records from a bivalved Methuselah (*Arctica islandica*, mollusca; Iceland). *Palaeogeography, Palaeoclimatology, Palaeoecology*, *228*, 130-148. https://doi.org/10.1016/j.palaeo.2005.03.049.

Schwager, E. E., Sharma, P. P., Clarke, T., Leite, D. J., Wierschin, T., Pechmann, M., ... McGregor, A. P. (2017). The house spider genome reveals an ancient whole-genome duplication during arachnid evolution. *BMC Biology*, *15*, 62. https://doi.org/10.1186/s12915-017-0399-x.

Serrano, M., Lin, A. W., McCurrach, M. E., Beach, D. & Lowe, S. W. (1997). Oncogenic *ras* provokes premature cell senescence associated with accumulation of p53 and p16[INK4a]. *Cell*, *88*, 593-602. https://doi.org/10.1016/s0092-8674(00)81902-9.

Shinzato, C., Khalturin, K., Inoue, J., Zayasu, Y., Kanda, M., Kawamitsu, M., ... Satoh, N. (2020). Eighteen coral genomes reveal the evolutionary origin of Acropora strategies to accommodate environmental changes. *Molecular Biology and Evolution*, 38, 16-30. https://doi.org/10.1093/molbev/msaa216.

Shinzato, C., Takeuchi, T., Yoshioka, Y., Tada, I., Kanda, M., Broussard, C., ... Inoue,

M. (2021). Whole-Genome Sequencing Highlights Conservative Genomic Strategies

of a Stress-Tolerant, Long-Lived Scleractinian Coral, *Porites australiensis* Vaughan,

1918.   *Genome   Biology   and   Evolution*,   13,   evab270.

https://doi.org/10.1093/gbe/evab270.

Shiomi, K., Kiyono, T., Okamura, K., Uezumi, M., Goto, Y., Yasumoto, S., ... Hashimoto,

N. (2011). CDK4 and cyclin D1 allow human myogenic cells to recapture growth

property without compromising differentiation potential. *Gene Therapy*, *18*, 857-866.

https://doi.org/10.1038/gt.2011.44.

Simakov, O., Kawashima, T., Marlétaz, F., Jenkins, J., Koyanagi, R., Mitros, T., ...

Gerhart, J. (2015). Hemichordate genomes and deuterostome origins. *Nature*, 527, 459-

465. https://doi.org/10.1038/nature16150.

Simakov, O. & Kawashima, T. (2017). Independent evolution of genomic characters

during   major   metazoan   transitions.   *Developmental   Biology*,   *427*,   179-192.

https://doi.org/10.1016/j.ydbio.2016.11.012.

Slyusarev, G. S., Starunov, V. V., Bondarenko, A. S., Zorina, N. A. & Bondarenko, N. I.

(2020). Extreme Genome and Nervous System Streamlining in the Invertebrate

Parasite   *Intoshia   variabili*.   *Current   Biology*,   30,   1292-1298.   doi:

10.1016/j.cub.2020.01.061.

Soufir, N., Avril, M. F., Chompret, A., Demenais, F., Bombled, J., Spatz, A., ... Bressac-de Paillerets, B. (1998). Prevalence of *p16* and *CDK4* germline mutations in 48 melanoma-prone families in France. The French Familial Melanoma Study Group. *Human Molecular Genetics*, *7*, 209-216. https://doi.org/10.1093/hmg/7.2.209.

Takeuchi, T., Suzuki, Y., Watabe, S., Nagai, K., Masaoka, T., Fujie, M., ... Myers, E. W. (2022). A high-quality, haplotype-phased genome reconstruction reveals unexpected haplotype diversity in a pearl oyster. *DNA Research*, 29, 1-13. https://doi.org/10.1093/dnares/dsac035.

Telford, M. J., Budd, G. E., & Philippe, H. (2015). Phylogenomic Insights into Animal Evolution. *Current Biology*, *25*, R876-887. https://doi.org/10.1016/j.cub.2015.07.060.

Tsao, H., Benoit, E., Sober, A. J., Thiele, C. & Haluska, F. G. (1998). Novel mutations in the *p16/CDKN2A* binding region of the *cyclin-dependent kinase-4* gene. *Cancer Research*, 58, 109-113.

Veinalde, R., Ozola, A., Azarjana, K., Molven, A., Akslen, L. A., Doniņa, S., ... Pjanova, D. (2013). Analysis of Latvian familial melanoma patients shows novel variants in the noncoding regions of *CDKN2A* and that the *CDK4* mutation R24H is a founder

mutation. *Melanoma Research*, *23*, 221-226. https://doi.org/10.1097/CMR.0b013e3283608695.

Wan, Z.-Y., Song, F., Sun, Z., Chen, Y.-F., Zhang, W.-L., Samartzis, D., ... Luo, Z.-J. (2014). Aberrantly expressed long noncoding RNAs in human intervertebral disc degeneration: a microarray related study. *Arthritis Research and Therapy*, *16*, 465. https://doi.org/10.1186/s13075-014-0465-5.

Webster, K. A., Henke, K., Ingalls, D. M., Nahrin, A., Harris, M. P. & Siegfried, K. R. (2018). Cyclin-dependent kinase 21 is a novel regulator of proliferation and meiosis in the male germline of zebrafish. *Reproduction*, *157*, 383-398. https://doi.org/10.1530/REP-18-0386.

Weinberg, R. A. (2014). *The Biology of Cancer*, *2nd edition*. New York City: Garland Science.

Wölfel, T., Hauer, M., Schneider, J., Serrano, M., Wölfel, C., Klehmann-Hieb, E., ... Beach, D. (1995). A p16[INK4a]-insensitive CDK4 mutant targeted by cytolytic T lymphocytes in a human melanoma. *Science*, *269*, 1281-1284. https://doi.org/10.1126/science.7652577.

Wood, D. J. & Endicott, J. A. (2018). Structural insights into the functional diversity of the CDK-cyclin family. *Open Biology*, *8*, 180112. https://doi.org/ 10.1098/rsob.180112.

Yates, A. D., Allen, J., Amode, R. M., Azov, A. G., Barba, M., Becerra, A., ... Flicek, P.

(2022). Ensembl Genomes 2022: an expanding genome resource for non-vertebrates.

https://doi.org/10.1093/nar/gkab1007.

Yoshioka, Y., Suzuki, G., Zayasu, Y., Yamashita, H. & Shinzato, C. (2022). Comparative

genomics highlight the importance of lineage-specific gene families in evolutionary

divergence of the coral genus, *Montipora*. *BMC Ecology and Evolution*, 22, 71.

https://doi.org/10.1186/s12862-022-02023-8.

Zhang, P., Liégeois, N. J., Wong, C., Finegold, M., Hou, H., Thompson, J. C., ... Elledge,

S. J. (1997). Altered cell differentiation and proliferation in mice lacking p57[KIP2]

indicates a role in Beckwith-Wiedemann syndrome. *Nature*, *387*, 151-158.

https://doi.org/10.1038/387151a0.

Zuo, L., Weger, J., Yang, Q., Goldstein, A. M., Tucker, M. A., Walker, G. J., ... Dracopoli,

N. C. (1996). Germline mutations in the p16[INK4a] binding domain of CDK4 in familial

melanoma. *Nature Genetics*, *12*, 97-99. https://doi.org/10.1038/ng0196-97.

Figure legends

FIGURE 1 Distribution of *Cdkn2*, *Cdk4/6* and *Cdkn1*, and the extent of the

conservation of the key charged amino acids in all invertebrate phyla. The phylogenetic

tree was modified from Telford *et al*. (2015). The rows for phyla in which more than

five species were included in this analysis are highlighted with yellow. *The N-terminal

sequence including the corresponding residue to R24/31 in the contig was lacking. ND:

not determined because of the lack of the genome data. N/A: not applied.

FIGURE 2 Amino acid alignment around the residue corresponding to Cdkn2 D84 in

representative invertebrate species from seven phyla where *Cdkn2* was detected. The

residue corresponding to Cdkn2 D84 is colored with light green when the negative

charge is conserved. If not, it is colored with magenta. Human CDKN2's are aligned as

references. **a** and **b** represent amino acids of α-helix and β-turn in CDKN2A,

respectively (Baumgartner *et al*., 1998).

FIGURE 3 *Cdkn2* and *Faf1* are tightly linked during evolution. Arrows show the

direction of transcription. The number of the chromosome or the linkage group is shown

when available. Microsynteny around human *CDKN2* loci are shown as references.

Genes encoding ankyrin repeat-containing proteins other than Cdkn2 are represented in

red.

FIGURE 4 Microsynteny around the *Faf1* locus in representative invertebrates without

Cdkn2. Arrows show the direction of transcription. The number of the chromosome or

the linkage group is shown when available. Genes encoding ankyrin repeat-containing

proteins other than Cdkn2 are represented in red.

FIGURE 5 Amino acid alignment around the residue corresponding to Cdk4/6 R24/31

in representative invertebrate species of Deuterostomata. The residue corresponding to

Cdk4/6 R24/31 is colored with light green when the strong positive charge was

conserved. If not, it is colored with magenta. Human CDK4 and CDK6 are aligned as

references.

FIGURE 6 Hypothesis on the evolution of *Cdkn2* locus. (a) Loss of *Cdkn2* occurred

multiple times. (b) Evolution of microsynteny at *Cdkn2* locus. *Cdkn2e* is a hypothetical

transient gene. For details, see the main text.

Supplementary material

Table S1 The list of the analyzed invertebrate *Cdkn2*'s.

Table S2 Results of blast search on invertebrate Cdkn2. (a) Result of blastp on

refseq_protein with Cdkn2. (b) Results of tblastn on wgs database with Cdkn2. (a) and

(b) are put together in one file with the worksheet tab.

Table S3 The list of the analyzed vertebrate *Cdkn2*'s.

Table S4 Microsynteny around *Cdkn2* locus in invertebrates.

Table S5 Microsynteny around *Faf1* locus in invertebrates. (a) Microsynteny around *Faf1*

in invertebrates other than Arthropoda. (b) Microsynteny around *Faf1* in Arthropoda. (a)

and (b) are put together in one file with the worksheet tab.

Table S6 The list of the identified invertebrate *Cdkn1*'s. (a) The list of the *Cdkn1* in the

species other than Arthropoda. (b) The list of the *Cdkn1* in Arthropoda. (a) and (b) are

put together in one file with the worksheet tab.

Table S7 The list of the analyzed invertebrate *Cdk4/6*'s. The tables contain the database

ID number and the scientific name of the species in the deuterostome invertebrates (A),

Platyhelminthes (B), other Lophotrochozoa (C), Nematoda (D), other Ecdysozoa (E), and

basal invertebrates (F).

Table S8 The list of the analyzed vertebrate *Cdk4/6*'s.

Table S9 The list of possibly misannotated invertebrate genes as *Cdkn2*.

Figure S1 Amino acid alignment of invertebrate Cdkn2's. (a) The alignment of the N-

terminal half of Cdkn2. (b) The alignment of the C-terminal half. The residue

corresponding to Cdkn2 D84 is colored with light green when the negative charge is

conserved. If not, it is colored with magenta. (c) Consensus residues in invertebrate

Cdkn2. **a** and **b** represent amino acids of α-helix and β-turn in *Homo sapiens* CDKN2's,

respectively (Baumgartner *et al*., 1998).

Figure S2 Cdkn2 amino acid alignment around the residue corresponding to D84 in

vertebrates; Cdkn2a in mammals (a), and in non-mammalian vertebrates (b); Cdkn2b in

mammals (c), in non-mammalian/non-fish vertebrates (d), and in fish (e); Cdkn2c in

mammals (f), in non-mammalian/non-fish vertebrates (g), and in fish (h); Cdkn2d in

mammals (i) and in non-mammalian vertebrates (j). The residue corresponding to

Cdkn2 D84 is colored with light green when the negative charge is conserved. If not, it

is colored with magenta.

Figure S3 Cdk4/6 amino acid alignment around the residue corresponding to R24/R31

in invertebrates; deuterostome invertebrates (a), Nematoda (b), other Ecdysozoa (c),

Platyhelminthes (d), other Lophotrochozoa (e), and basal invertebrates (f). The residue

corresponding to Cdk4/6 R24/31 is colored with light green when the strong positive

charge is conserved. If not, it is colored with magenta.

Figure S4 Cdk4 amino acid alignment around the residue corresponding to R24/R31 in

vertebrates; Cdk4 in mammals (a), in non-mammalian/non-fish vertebrates (b), and in

fish (c); Cdk6 in mammals (d), in non-mammalian/non-fish vertebrates (e), and in fish

(f); Cdk21 in fish (g). The residue corresponding to Cdk4/6 R24/31 is colored with light

green when the strong positive charge is conserved. If not, it is colored with magenta.

Figure S5 The data to determine the Spearman's rank correlation coefficient. (a) "**[2]/[1]**

in %" and "% of -R, K at Cdk4/6 R24/31" were used to determine the Spearman's rank

correlation coefficient. (b) The scatter plot of "**[2]/[1]** in %" and " of -R, K at Cdk4/6

R24/31".

FIGURE 1 Distribution of *Cdkn2*, *Cdk4/6* and *Cdkn1*, and the extent of the conservation of the key charged amino acids in all invertebrate phyla.

```
                                                  ankyrin repeat 3
                                    aa          b          bb  aaaaaaaa  aaaaaaaaaa
        Homo sapiens CDKN2A         LHGAEPNCADPA----------TLTRPVHDAAREGFLDTLVVLHRAGARL
        Homo sapiens CDKN2B         LHGAEPNCADPA----------TLTRPVHDAAREGFLDTLVVLHRAGARL
        Homo sapiens CDKN2C         LRGANPDLKDRT-----------GFAVIHDAARAGFLDTLQTLLEFQADV
        Homo sapiens CDKN2D         KQGASPNVQDTS-----------GTSPVHDAARTGFLDTLKVLVEHGADV

Cephalochrdata
        Branchiostoma belcheri      ENGADPNVRDPGR---------KGQYPVHDAARGGWDEALKLLFDYHAEL
Hemichordata
        Ptychodera flava            RRGADPNVADST-----------GRAPIHDAAEGGYVNTVQVFVEYGGDL
        Saccoglossus kowalevskii    RRGADPNVADST-----------GRAPIHDAAEGGYVNTVQVFVEYGGDL
Echinodarmata
        Strongylocentrotus purpuratus  DHGADLDRPDPS----------VQRTILHDIAERGCIETLNLLLEHGANA
        Apostichopus japonicus      QKGANPNVYDMNP---DSDHRPFLRTPLHDALHHGNEDSAILLIQHGADL
        Acanthaster planci          KSGADPNVRDRD----------VGRTPLHDAAEHGHEDTVRLLLKHGADP
        Ophioderma brevispina       DEGADPNVRDPD----------VGRTPLHDAVEYGIPSTVMCLLQGGADV
        Anneissia japonica          EYGANPNLKEPV----------MGGIPLHQAAREGATATLELMLQFGANV
        Nesometra (Dorometra) sesokonis EYGANPHLPEPV----------MGGIPLHNAAREGATTTLELLLQYGSNV
Onychophora
        Epiperipatus broadwayi      KAGANVNVSDV-----------TLRTPVHDAAEAGFLDVVQNLIAYGADI
        Euperipatoides rowelli      SAGANPNICDR-----------TLRTPVHDAAEAGFLDVLQCLLANGGVI
Annelida
        Lamellibrachia satsuma      RNGADPNVRDRS--YP--GHQDTLVTPAHDVARAGYTDVLKCLLRYGADV
        Dimorphilus gyrociliatus    KAGANPNVQDRG--IG--NGRCLLLTPGHDAARVGYVRTLKVLLKYGADT
        Sipunculus nudus            EAGADPNVVDHS--KN--NKKDTLLTPAHDVARQGYLDTLKVLLWYGADI
Mollusca
        Pomacea canaliculata        EAGGDPNALDIG--KR--TDTSTLRTVAHDVAASGYTDVLRCLLRHGADI
        Nodopelta heminoda          RAGADPNVKDVS--LRQDGGSSTFRTICHDVAESGALSTLRCLLRHGADI
        Patella vulgata             HSGANPNNQDIS--TRDTDCLSTYRTICHDVAASGYLETLHVLLKYGTDT
        Haliotis rufescens          CAGADPNVKDIS--HRHIGGTSTYRTICHDVAESGSLPTLQCLLRHGADI
        Nautilus pompilius          RAGADPNLMEGD----------TRRTICHDVAESGYLDTLRCLLKHGADL
Brachiopoda/Phoronida
        Lingula anatina             KHGADPNIQDKH--FQD-LHLSSYRTPAHDIAEQGFLDTMKVLLKHGADV
        Phoronis ovalis             EHGADPNLKERSPHHEADPSRSLLRTPAHNTCCEGFLDTLKTLLRCGANI
        Phoronis pallida            MRGADPNIQDT----------STLRTILHDVCEFGYMDTLQCLISFGANL
```

FIGURE 2 Amino acid alignment around the residue corresponding to Cdkn2 D84 in representative invertebrate species from seven phyla where *Cdkn2* was detected.

FIGURE 3 *Cdkn2* and *Faf1* are tightly linked during evolution.

FIGURE 4 Microsynteny around the *Faf1* locus in representative invertebrates without *Cdkn2*.

```
Cdk4
Homo sapiens              RYEPVAEIGVGAYGTVYKARDPHSG---HFVALKSVRVPN
Cdk6
Homo sapiens              QYECVAEIGEGAYGKVFKARDLKNG--GRFVALKRVRVQT

Cdk4/6
Urochordata
Phallusia fumigata        RYMPLEEIGVGAYGIVYKARDTSPDTDHGVVAMKCLRVEN
Ciona intestinalis        RYHTLSEIGVGAYGVVFKAQDMQSDDENKFVAIKCVRVEN
Ciona savignyi            RYHTLTEIGVGAYGVVFKAQDRHSEEENKFVAMKCIRVEN
Corella inflata           RYKPVEKIGVGAYGVVFKAKDLRKD--GEFVAMKCLRVEN
Molgula occulta           KYEKDEEIGVGAYGTVYRGKDLANL--GKYVALKCLRVQT
Botrylloides schlosseri   NYEMVAEIGVGAYGVVYRARDLEND--RKLVALKCLRVQN
Styela clava              NYEMVAEIGVGAYGIVYRARDLENN--GRFVALKCLRVQN
Halocynthia roretzi       NYEMVAEIGVGAYGIVYRARDLEND--NRYVALKCLRVQN
Oikopleura dioica         DWIDMEEIGGGAYGVVFRARDRQHD--NRVVALKRLRVQA
Cephalochordata
Branchiostoma belcheri    EYEKVAKIGTGAYGTVYKARDLKNG--GEFVALKRVRVQT
Asymmetron lucayanum      QYEKVAKIGTGAYGTVYKARDLKNG--GEFVALKRVRVQT
Hemichordata
Ptychodera flava          RYEELAEIGTGAYGIVYKARDVESG---RIVALKKVRVQT
Saccoglossus kowalevskii  KYEEVAEIGTGAYGTVYKARDLETG---RYVALKKVRVQT
Echinodermata
Strongylocentrotus purpuratus EYVQAAEIGSGAYGIVYKARDTETG---HFVALKSVRIPI
Apostichopus japonicus    EFSIIEEIGTGAYGTVYKARSKATR---EFVALKKIIVSH
Acanthaster planci        NYAVISEIGTGAYGTVYKARDLETG---KVVALKKVRLHT
Ophionereis fasciata      NYSVMAEIGTGAYGTVFKARDKTNG---HIVALKKVRIQT
Anneissia japonica        DFQKIEKIGEGTYGVVYKARDKISG---EVVALKKIRLDT
Xenacoelomorpha
Hofstenia miamia          HYSVLEKVGVGAYGEVYRAKDPRTG---GHVALKRVRVFD
Praesagittifera naikaiensis NYSIVESVGKGAYAKVYKAVNPQNPT--EIVALKQINIAS
```

FIGURE 5 Amino acid alignment around the residue corresponding to Cdk4/6 R24/31 in representative invertebrate species of Deuterostomata.

FIGURE 6 Hypothesis on the evolution of *Cdkn2* locus.

TABLE 1 Detailed distribution of Cdkn2 and Cdk4/6 in the subtaxons of the seven phyla where Cdkn2 was detected.

Phyla/subtaxons	Total number of analyzed species	The number of species where *Cdk4/6* was detected	The number of species where *Cdkn2* was detected
Cephalochordata	4	4	3
Hemichordata	2	2	2
Echinodermata	21	21	21
Echinozoa (incl. Holothuroidea)	9	9	9
Asterozoa (incl. Ophiuroidea)	9	9	9
Pelmatozoa	2	2	2
Onychophora	2	1	2
Annelida	13	7	6
Polychaeta (incl. Pogonophora)	9	6	5
Sipuncula	1	0	1
Clitellata	1	1	0
Mollusca	65	63	22
Bivalvia	30	30	0
Gastropoda	30	29	21
Caenogastropoda	11	10	10
Heterobranchia	6	6	0
Neomphalina	2	2	1
Patellogastropoda	4	4	3
Vetigastropoda	7	7	7
Cephalopoda	5	5	1
Coleoidea	4	4	0
Nautiloidea	1	1	1
Brachiopoda/Phoronida	4	4	4

TABLE 2 Number of species with each type of substituted amino acids for CDK4/6 R24/R31.

Phyla	Total number of analyzed sequences	R	K	H	A	V	I	L	T	S	N	Q	E	Y	F	C	% of non-R, K
Deuterostome Invertebrates	**43**	**36**	**4**			**1**							**2**					**7.0%**
Urochordata	14	9	3										2					14.3%
Cephalochordata	4	4																0.0%
Hemichordata	2	2																0.0%
Echinodermata	21	21																0.0%
Xenacoelomorpha	2		1			1												50.0%
Ecdysozoa	**1058**	**772**	**199**	**3**	**1**	**2**	**3**	**2**	**56**	**2**	**5**	**2**	**2**			**7**	**8.0%**
Arthropoda	979	753	186	3	1	1	1	2	17		4	1	1			7	3.9%
Onychophora	2	2																0.0%
Tardigrada	2	1				1												50.0%
Nematomorpha	3	3																0.0%
Nematoda	72	13	13				2		39	2	1	1	1				63.9%
Lophotrochozoa	**133**	**72**	**9**	**1**		**2**	**5**	**3**	**35**	**1**		**1**		**3**	**1**		**39.1%**
Rotifera	9	5					1							3			44.4%
Annelida	7	6	1															0.0%
Mollusca	63	56	7															0.0%
Nemertea	2						2											100.0%
Brachiopoda /Phoronida	4	4																0.0%
Bryozoa	1	1																0.0%
Platyhelminthes	47		1	1		2	2	3	35	1		1			1		97.9%
Basal Invertebrates	**62**	**53**	**3**			**2**	**1**	**1**							**2**		**9.7%**
Cnidaria	60	51	3			2	1	1							2		10.0%
Ctenophora	1	1																0.0%
Placozoa	1	1																0.0%