Parental assigned chromosomes for cultivated cacao provides insights into genetic architecture underlying resistance to vascular streak dieback

Authors: Peri A. Tobias[1]*, Jacob Downs[1], Peter Epaina[1,5], Gurpreet Singh[1], Robert F. Park[1], Richard J. Edwards[2], Eirene Brugman[3], Andi Zulkifli[6], Junaid Muhammad[3], Agus Purwantara[4], David I. Guest[1]

*Author for correspondence: peri.tobias@sydney.edu.au

Affiliations: [1]School of Life and Environmental Sciences, The University of Sydney, Camperdown, NSW, Australia, [2] Minderoo OceanOmics Centre at UWA, Oceans Institute, The University of Western Australia, Perth, WA, Australia, [3]Cocoa Research Center Faculty of Agriculture, Universitas Hasanuddin, Makassar, Indonesia, [4]Mars Cocoa Research Station, Pangkep, Indonesia, [5]Cocoa Board of Papua New Guinea, Kokopo, East New Britain Province, Papua New Guinea, [6]Megarezky University, Makassar, Indonesia

Keywords: *Theobroma cacao, Ceratobasidium theobromae,* VSD*,* chromosome-level genome, resistance genes, NLR

Abstract

Diseases of *Theobroma cacao* disrupt cocoa bean supply and economically impact growers. Vascular streak dieback (VSD), caused by *Ceratobasidium theobromae,* is a new encounter disease of cacao currently contained to southeast Asia and Melanesia. Resistance to VSD has been tested with large progeny trials in Sulawesi, Indonesia, and in Papua New Guinea with the identification of informative quantitative trait loci (QTL). Using a VSD susceptible progeny tree (clone 26), derived from a resistant and susceptible parental cross, we assembled the genome to chromosome-level and discriminated alleles inherited from either resistant or susceptible parents. The parentally phased genomes were annotated for all predicted genes and then specifically for resistance genes of the *nucleotide-binding site leucine-rich repeat*

30   class (NLR). On investigation, we determined the presence of NLR clusters at informative

31   QTLs, as well as other potential disease response gene candidates. Understanding the

32   genetics underlying resistance and susceptibility to VSD will accelerate the breeding cycle by

33   providing clear targets for molecular screening. Additionally, we provide the first diploid, fully

34   scaffolded and parentally phased genome resource for *Theobroma cacao* L.

Introduction

Cacao, *Theobroma cacao* L. (Malvaceae) is an important perennial tree crop producing beans

that are used as raw materials in the chocolate, beverage, and cosmetic industries (Schnell *et*

al., 2007). Much of the world's cocoa bean production comes from small-holder, mixed

cropping farms in some fifty countries that depend on trade with developed countries where

the raw product is manufactured into chocolate (Daymond *et al.*, 2022). Cacao is a difficult

crop to grow profitably due to several endemic problems. Alongside other problems, diseases

disrupt the supply of cocoa and have serious economic impacts to growers (Marelli *et al.*,

2019). Responses to disease can lead to tree removal and replacement with more profitable

and reliable crops (Daymond *et al.*, 2022). The centre of origin for *T. cacao* is the Upper

Amazon, South America, where evolutionary studies indicate divergence from close common

ancestors around 9.9 million years ago (Richardson *et al.*, 2015). While genetic diversity exists

within the geographic centre of origin, cultivation largely relies on a limited genetic pool of *T.*

cacao (Motamayor *et al.*, 2008). Southeast (SE) Asia is one of the major suppliers of cocoa

beans and cultivation has historically been relatively disease-free due to the lack of co-

evolved pathogens (Bailey and Meinhardt, 2016). However in the 1960s, vascular streak

dieback (VSD) emerged as a new encounter disease of cacao and caused devastating crop

losses in Papua New Guinea (PNG) (Guest and Keane, 2007).

The causal agent of VSD is *Ceratobasidium theobromae*, a fastidious biotrophic fungus, first

discovered and described by Philip Keane (1972). *Ceratobasidium theobromae* then spread to

other SE Asian and Melanesian cacao regions as an emergent disease on cacao (Guest and

Keane, 2007). In Indonesia, the pathogen has spread into Sumatera Regions (Trisno *et al.*,

2016) and was subsequently confirmed in a new area of Barru District, Sulawesi (Junaid *et al.*,

2020). The pathogen is currently contained to the SE Asia region but presents a serious

biosecurity threat to cacao in West Africa and Latin America. Symptoms include green spotted

chlorosis in older leaves and vascular browning in the petioles and stems that often lead to

rapid branch or whole tree death in susceptible genotypes (Guest and Keane, 2007; Samuels

et al., 2012). Some genotypes of *T. cacao* that were resistant to disease 50 years ago remain

resistant, indicating that durable resistance is highly heritable and therefore may be exploited

in breeding to introgress resistance into new, improved cacao genotypes (Tan and Tan, 1988).

Varieties of cacao that show resilience to infection by *C. theobromae* have been studied in

both PNG and Sulawesi, Indonesia. Results of these studies identified informative quantitative

trait loci (QTL) that correlate with both qualitative and quantitative resistance in progeny

populations (Epaina, 2012; Singh, 2021). The two independent studies used progenies derived

from crosses between Trinitario **KA2-101** (resistant to VSD) and **K82** (susceptible to VSD) in

PNG (Epaina, 2012), and between Amelonado **S1** (resistant to VSD) with Iquitos, Criollo and

Amelonado **RUQ 1347** (susceptible to VSD and formerly known as CCN 51) in Sulawesi,

Indonesia (Singh, 2021). The progeny trials used 346 and 130 progeny that segregated in

response to VSD in PNG and Sulawesi, respectively. The informative QTLs associated with

resistance indicated a likely polygenic response and were mapped onto nine of the ten *T.*

cacao chromosomes from the PNG progeny trial (Epaina, 2012) and three in the Sulawesi trial

(Singh, 2021). The two studies used either simple sequence repeat (SSR) or single nucleotide

polymorphism (SNP) markers and compared linkage maps with established disease resistance

mapping developed by Lanaud et al. (2009). The available genomes for Matina 1-6

(Motamayor *et al.*, 2013) and Criollo (Argout *et al.*, 2011) were used to inform the linkage

maps. Interestingly, VSD QTLs co-localized with a region identified for resistance to

Phytophthora Pod Rot (PPR) and Frosty Pod Rot (FP) (causal agents *Phytophthora palmivora*

and *Moniliophthora roreri*, respectively) on chromosome 8, while others were common for

response to both VSD and PPR on chromosomes 3, 4 and 5 in only one study. Both studies

identified VSD QTLs within similar regions on chromosomes 3, 8 and 9. These apparent shared

QTL regions indicate hotspots for disease resistance.

Classical selection and breeding for resistance, particularly in perennial trees, can be a long

and difficult process (Sniezko, 2006). The research is further complicated for resistance to *C.*

theobromae as the pathogen cannot be cultured in vitro and is extremely slow growing within

the host plant before visible symptoms develop. Traditional Koch's postulates cannot be

satisfied for biotrophic organisms, however molecular approaches are becoming increasingly

important in validation studies (Bhunjun *et al.*, 2021). Understanding the genetics underlying

resistance and susceptibility to VSD is likely to accelerate the breeding cycle by providing clear

targets for molecular screening.

Effective plant responses to disease at the molecular level are well established as involving

resistance genes of the *nucleotide-binding site leucine-rich repeat* family (NLRs) (Fluhr *et al.*,

2001). NLR genes are known to be numerous, clustered, and highly polymorphic both

between and within species (Barragan and Weigel, 2020; Santos *et al.*, 2022; Weyer *et al.*,

2019; Van de Weyer *et al.*, 2019). We therefore approached the problem of understanding

the underlying genetics of resistance to VSD by building on the structural knowledge of NLR

genes and from the progeny QTL studies of Epaina (2012) and Singh (2021). We aimed to

leverage the latest sequencing and bioinformatic approaches to gain a clearer insight into the

genetic regions of interest. Our premise for the work was that (i) QTLs are informative for

predicting resistance, (ii) genetic resistance hotspots are present, (iii) QTLs may indicate the

presence of NLR gene clusters, (iv) polymorphic NLR genes within clusters, from contrasting

genotypes, may explain disease phenotype. The questions we therefore posed were, can we

assemble the cacao genome and successfully discriminate alleles inherited from parents of

known VSD phenotype within an F1 tree progeny, and if so, can we determine differences in

NLR clusters, type, and specific gene sequences at the QTL locations, based on parentage? An

additional question that we tested was, using new sequencing technologies, can we assemble

and scaffold to chromosome level both pathogen (~30 Mb) and host (~380 Mb) genomes from

a single sample? This would circumvent the inability to culture the biotrophic *C. theobromae*

and provide insight into the genomics of a compatible interaction. This final aim was not

successful due to the limited pathogen DNA we were able to extract, but future studies could

work to improve on this approach.

Taking these premises and goals, we built a high-quality parental phased genome resource

for a susceptible cacao (clone 26) of known parental cross, **S1 maternal** and **RUQ 1347**

paternal. We annotated the genome for all predicted genes, and then specifically for only

NLR-type genes, and conducted fine scale investigation of genomic regions associated with

resistance to *C. theobromae* to determine the resistance gene complements inherited from

both parents. We made comparative studies into the NLR genes, structurally, quantitatively,

and qualitatively. Our investigations show structural differences in NLR presence within

parental chromosomes and provide insights into resistance to VSD and potentially other

serious diseases of cacao. Additionally, we provide the first diploid, fully scaffolded and

phased genome resource for *Theobroma cacao* L.

Results and Discussion

Near complete parental assigned chromosomes for cultivated *Theobroma cacao*

We assembled and parentally phased the chromosomes of a VSD susceptible *T. cacao* (clone

26), the progeny of a cross between a maternal resistant (S1, haplotype A) with a paternal

susceptible (RUQ 1347, haplotype B) tree. All but three inherited chromosomes (two S1, one

RUQ 1347) show telomeric caps (black dots in Figure 1). The genomes show few gaps (red

plus symbol in Figure 1), high levels of contiguity and completeness, as well as chromosomal

synteny (Edwards *et al.*, 2022) based on conserved single copy orthologs (Simão *et al.*, 2015).

Minor inverted translocations on chromosomes four and nine (pink synteny lines) are likely

to be biologically real considering the lack of flanking gaps that would indicate mis-scaffolding.

The phased genomes are both highly complete according to assessment of conserved single

copy orthologs and basic statistics (Table 1).

Figure 1 Chromsyn (Edwards *et al.*, 2022) **synteny plots of Theobroma cacao parentally phased genome, haplotype A (top)**
and B (below). Synteny blocks of collinear "Complete" BUSCO genes (Simão *et al.*, 2015) **link scaffolds from adjacent**
assemblies: blue, same strand; red, inverse strand. Yellow triangles mark "Duplicated" BUSCOs. Filled circles mark
telomere predictions from Telociraptor (black) (Edwards, 2023). Assembly gaps are marked as dark red + signs.

The goal for genome assemblies of gapless and telomere-to-telomere (T2T) chromosomes has

recently been achievable with the latest sequencing technologies (Sergey *et al.*, 2022). The

resulting genome assemblies permit greater depth in understanding structural and

evolutionary biology (Mao and Zhang, 2022). Recently, chromosome-level genomes for three

wild cacao species from the Upper Amazon were made public (Nousias *et al.*, 2024). These

provide the first cacao genomes produced using long read sequence technology and indicate

an evolutionary divergence, based on conserved orthologs, from cultivated Criollo and Matina

reference genomes (Argout *et al.*, 2011; Motamayor *et al.*, 2013) at ~ 1.83 and ~ 1.34 million

years ago. The wild cacao assemblies represent the collapsed haploid state for genomes and
are larger in both chromosome and genome base pair (bp) lengths compared with earlier
reference genomes. Each of our parentally phased genome assemblies are comparable in size
to the wild cacao, at around 380 Mb, however we annotated more predicted protein coding
genes at around 26,000 (Table 1) compared to around 21,000, and higher percentage of
repetitive sequences at 62% compared to 53% (Nousias *et al.*, 2024). We annotated all genes
for the predicted nucleotide binding site (NBARC) domain type as well as predicted 'complete'
NLRs that incorporate NBARC plus leucine-rich repeat (LRR) domains (Table 1). Of the
complete NLRs we determined only 11 and 15 Toll/Interleukin-1 receptor-type TNL-type, and
only 5 and 12 novel integrated domains (ID) within NLRs per haplotype.

Table 1 **Raw data statistics from NanoPlot (De Coster and Rademakers, 2023) and genome statistics for both sets of**
parentally inherited *Theombroma cacao* chromosomes using Quast and BUSCOs (Gurevich *et al.*, 2013; Simão *et al.*, 2015)**.**
Haplotype A chromosomes derived from maternal plant S1 and haplotype B from paternal plant RUQ 1347. Annotation
statistics based on FindPlantNLRs and Funannotate (Chen *et al.*, 2023; Palmer and Stajich, 2023)**. NLR annotated resistance**
genes were classified and largely conformed to coiled-coil class with only small numbers of Toll/Interleukin-1 receptor-
type (TIR) and integrated domain type NLRs.

Raw HiFi data		
Total bases	30,259,224,851	
Mean read length	14,413.4	
Mean read quality	35.6	
Genome statistic	**Haplotype A**	**Haplotype B**
Total genome length (bp)	374,523,129	381,631,648
chromosome length (bp)	338,223,858	340,140,478
No. of chromosomes	10	10
N50 (bp)	38,945,508	39,156,173
L50	5	5
No. of contigs	165	315
GC (%)	33.73	33.67
BUSCO complete (genome; *n* = 1,614)	**98.2%**	**98.4%**
Single-copy	97.5%	97.8%
Duplicated	0.7%	0.6%
BUSCO fragmented	0.6%	0.9%
BUSCO missing	0.3%	0.7%

Protein-coding genes (funannotate)	26,199	26,321
NBARCs (FindPlantNLRs annotation)	342	370
NLRs	293	326
TIR-NLR (TNL)	11	15
Integrated domain (ID) NLRs	5	12
Repeats	**62.0%**	**62.4%**

Low structural diversity of NLRs between *Theobroma cacao* alleles

The NLR-type resistance genes were annotated, as described in the experimental procedures, and the parentally inherited chromosomes were compared, noting that haplotype A (red annotations (Figure 2Figure 2) aligned with the resistant maternal and haplotype B (yellow annotations) with susceptible paternal trees. We found that the predicted NLR-type genes varied in quantity and types between the two sets of chromosomes. Of the 342 and 370 predicted genes that contained NBARC domains (Table 1), one of the key identifiers of this class of NLR-type resistance genes, our research determined a discrepancy of 293 and 326 complete gene models within Haplotype A and B respectively. Using the web tool version of OrthoVenn3 (Sun *et al.*, 2023) with the OrthoMLC algorithm, using an expect value of 1e-15 and inflation value 1.5, we found that NLR amino acid sequences clustered into 83 orthologous groups (orthogroups). Of these orthogroups, 82 clusters were shared between the two haplotypes with the largest cluster including 55 proteins, 28 and 27 from haplotype A and B respectively. Forty-four orthogroups contained a single NLR protein from each haplotype, meaning that the matching homolog was present in each assembly. Nevertheless, aligned sequences for homologous proteins showed some variations and there were also 2.4% single NLRs, numbering 4 and 11 from haplotype A and B respectively.

Figure 2 Physical locations of predicted NLR genes on the paired chromosomes of *Theobroma cacao* (A) haplotype A (red) and (B) haplotype B (yellow) generated using ChromoMap (Anand and Rodriguez Lopez, 2022) **in RStudio.**

Low numbers of *Theobroma cacao* NLR-type genes compared to other tree species

Resistance genes of the NLR-type are numerous and highly polymorphic both within and between plant species (Van de Weyer *et al.*, 2019). Trees have generally been shown to have higher numbers of NLR-type genes than herbaceous species, and this is suggested to accomodate a changing pathogen environment over the life of a tree (Tobias and Guest, 2014). More recently, analyses of gene content in diploid genomes have been possible due to advances in sequencing and computational technologies. Notably, tree resistance genes have been investigated within the diploid genome for *Melaleuca quinquenervia* (Chen *et al.*, 2023), highlighting variation in numbers (763 and 733) between haplotypes and across chromosomes for this class of gene. Our analysis of the NLR complement within *T. cacao* showed similar representation to an earlier genome study (Argout *et al.*, 2011) that determined just 297 non-redundant NLR-type genes. Our work, however, presents the diploid content of NLRs as 293 and 326 for haplotype A and B respectively, permitting deeper investigation of homologs at loci. We confirmed low numbers and percentages of the Toll/Interleukin-1 receptor (TIR) NLR class (termed TNLs) of resistance genes within *T. cacao* (Figure 3) compared to many trees within the Rosid clade, in which they comprise 40 - 60% (Arya *et al.*, 2014; Christie *et al.*, 2016; Kohler *et al.*, 2008). This was noted in earlier molecular and genomic studies where only 4% of NLRs contained the TIR domain (Argout *et al.*, 2011; Kuhn *et al.*, 2006), with our classification indicating 3.8% and 5% for haplotype A and B. Unlike the previous study with primers for molecular markers that found TNLs on chromosome three and five (Kuhn *et al.*, 2006), we determined TNLs within clusters on chromosome 4 (three genes), 5 (two genes) and 6 (three and five genes on haplotype A and B), plus a single TNL

gene on chromosome 2, 4 and 6 of each haplotype. Our finding of only five and 12 NLR-type

genes with non-canonical integrated domains in the two haplotypes also suggests that this

integration is less important as a strategy for pathogen recognition and response in cultivated

T. cacao, compared to other species (Marchal *et al.*, 2022).

Figure 3 The predicted NLR gene complement in the parentally phased *Theobroma cacao* genome. The two sets of
chromosomes corresponding to (above) Haplotypes A and (below) B were independently classified (Chen *et al.*, 2023) **and**
visualised to present the domain classes using SankeyMatic (Bogart, 2014) including novel integrated domains (IDs) with
abbreviations derived from Pfam database (Mistry *et al.*, 2021). **NBARC = Nucleotide Binding Domain, TIR =**
Toll/Interleukin-1 receptor, Rx = Potato CC-NB-LRR protein Rx, CC=Coil-Coil Domain, RPW8 = RESISTANCE TO POWDERY
MILDEW 8-like coiled-coil

Theobroma cacao NLR-type genes are present and polymorphic at QTLs for VSD

resistance

Informative QTL regions were investigated for resistance gene homologs across the parental

inherited chromosomes. The most closely aligned VSD QTLs between PNG and Sulawesi *T.*

cacao progeny trials were on chromosomes 3 and 9, at around 29 and 31 Mb, while

chromosome 8 harboured a region between 6 – 7 Mb (Table 3, experimental procedures). We

checked these regions based on our annotations for NLRs and identified an NLR cluster of two

predicted genes at around 29 Mb on **chromosome 3** within both parentally inherited alleles.

The location matches almost exactly with our calculated region based on the reversed start position for the cM QTL (Table 3). When we aligned the four amino acid sequences for these NLRs with ClustalW (Thompson *et al.*, 1994) and visualised in BioEdit (Hall, 1991), we see close homology across the cluster but variation in one haplotype A protein, notably an insertion and deletion in the sequence for gene identity g2391.t1 (Figure 4, above). We found an NLR cluster of three genes on **chromosome 8** at around 3.8 Mb which falls within the 2 cM interval specified by Singh (2021) of 2 - 4 Mb from the QTL locus midpoint. The NLR cluster formed two orthogroups, one group of two head-to-head Rx-NLR predicted proteins from each haplotype (g1318.t1, g1320.t1 and g3681.t1, g3683.t1) and the other of single, nearly exactly homologous proteins, with one amino acid variant, namely proline (g1316.t1, Haplotype A) to arginine (g3679.t1, Haplotype B). Between the region 27 – 36 kb on **chromosome 9** we determined four NLRs from Haplotype A and three from Haplotype B. Paired homologs closest to the calculated QTL mid-point location at around 31 Mb on chromosome 9 were aligned and show an insertion/deletion as well as five single amino acid substitutions (Figure 4, below).The single predicted Rx-NLR gene on chromosome 9 of Haplotype A at 27 Mb has no homolog in Haplotype B.

The QTLs on chromosome 3 and 8 accounted for 11 and 15% phenotypic variance in the Sulawesi progeny trials and were informative for VSD and PPR resistance in the PNG trial (Table 3). The chromosome 9 QTL accounted for 16 - 18% phenotypic variance in the Sulawesi trial. All putative resistance genes at QTLs were annotated as Rx NLR-type resistance genes. The Rx-NLR coded protein from potato (*Solanum tuberosum*) has been identified as conferring resistance to the potato virus X by direct recognition. Recent work has clarified the interaction of sensor Rx-NLRs with downstream helper NLR protein oligomerisation (Contreras *et al.*, 2023), showing a distinct mechanism for immune responses in plants. While our work has uncovered the arrangement and sequence differences of predicted NLR clusters at QTLs for resistance to VSD, confirmation of these genes is needed across multiple progenies that have clear phenotypes for resistance/susceptibility. Amplification of DNA regions around these clusters will provide additional clarity but was beyond the scope of the current study. Functional validation would also be beneficial to determine mechanisms of recognition and response as it is known that paired head to head NLRs, as we identified on chromosome 8, can determine resistance to more than one pathogen (Xi *et al.*, 2022), potentially explaining the results from the study conducted in PNG.

Figure 4 Amino acid sequence alignment of an NLR cluster matching the QTL mapped regions at around 29 Mb on chromosome 3 (above) and single NLR homologs at 31 Mb chromosome 9 (below) within both parentally inherited alleles, A and B, for *Theobroma cacao* clone 26. All predicted NLRs are annotated as Rx-NLRs (Chen *et al.*, 2023) **named for resistance to potato virus X.**

Annotation statistics show high homology between allelic pairs in *Theobroma cacao*

The orthology of all annotated proteins was analysed with OrthoVenn3 as described previously. The predicted proteins indicated 22,486 shared clusters, of which 21,800 included a single predicted homologous gene from each haplotype. The largest cluster had 44 predicted proteins and there were 9.76% singletons. In total there are 2,928 (2,896), 1,865 (1,798), 3,278 (3,315) predicted genes for Haplotype A (and B) on chromosomes 3, 8 and 9 respectively. Except for chromosome 9, with around 76 predicted genes within ~2 Mb of the QTL informing resistance to VSD, the locations on chromosomes 3 and 8 are in highly dense gene-encoding regions, with 325 and 370 predicted genes. We investigated the predicted genes at the QTLs and found that homologs were present between haplotypes (Table 2) and we reviewed their predicted protein functions based on Pfam database (Mistry *et al.*, 2021) annotations. Due to the density of predicted genes, we limited our investigations to

immediate proximity around NLRs that corresponded to QTLs. Around the predicted QTL for resistance to VSD on chromosome 3, the Rx-NLR gene cluster is interspersed with two predicted genes coding for protein kinases (Table 2). The kinases have fused LRR domains and predicted transmembrane motifs indicating extracellular receptors and a putative role in defence response. The importance of this class of receptor-like kinase (RLK) has been understood in rice with studies showing *Xa21* confers resistance to *Xanthomonas* sp. and multiple other pathogens (Ercoli *et al.*, 2022). The finding of kinases so closely associated with the Rx-NLR genes on chromosome 3 in both parental alleles, at the predicted region for VSD resistance QTL, suggests a potential role in disease response and a target for further research. The NLR QTL regions associated with resistance on chromosomes 8 and 9 did not have associated disease response-like genes, however it is likely that multiple genes are involved in the VSD resistance response observed.

Table 2 Predicted and annotated genes that are in proximity to the QTLs for resistance to *Ceratobasium theobromae*, causal agent of vascular streak dieback in cultivated *Theobroma cacao*. The annotated genomes are available at NCBI: PRJNA1083235 and PRJNA1086984.

Haplotype A	Haplotype B	Pfam ID	Annotation
NLR QTL chromosome 3 (40 kb region)			
V9T21_008377	WDX93_008321	PF00334	Nucleoside diphosphate kinase
V9T21_008378	WDX93_008322	PF01625	Peptide methionine sulfoxide reductase
V9T21_008380	WDX93_008324	PF07714	Protein tyrosine and serine/threonine kinase
V9T21_008380	WDX93_008324	PF13855	Leucine rich repeat
V9T21_008382	WDX93_008326	PF07714	Protein tyrosine and serine/threonine kinase
V9T21_008382	WDX93_008326	PF13855	Leucine rich repeat
V9T21_008382	WDX93_008326	PF08263	Leucine rich repeat N-terminal domain
V9T21_008383	WDX93_008327	PF02298	Plastocyanin-like domain
NLR QTL chromosome 8 (10 kb region)			
V9T21_020396	WDX93_020310	PF00745	Glutamyl-tRNAGlu reductase, dimerisation
V9T21_020396	WDX93_020310	PF01488	Shikimate / quinate 5-dehydrogenase
V9T21_020397	WDX93_020311	PF01293	Phosphoenolpyruvate carboxykinase
V9T21_020399	WDX93_020313	PF08694	Ubiquitin-fold modifier-conjugating enzyme1
V9T21_020401	WDX93_020315	PF12295	Symplekin tight junction protein C terminal
V9T21_020401	WDX93_020315	PF11935	Symplekin/PTA1 N-terminal
V9T21_020402	WDX93_020316	PF11935	Symplekin/PTA1 N-terminal
V9T21_020404	WDX93_020318	PF00719	Inorganic pyrophosphatase
NLR QTL chromosome 9 (3 kb region)			
V9T21_023576	WDX93_023431	PF00514	Armadillo/beta-catenin-like repeat
V9T21_023578	WDX93_023433	PF02341	RbcX protein
V9T21_023579	WDX93_023434	PF00076	RNA recognition motif

Concluding remarks

A recent study, using single nucleotide polymporphisms, confirmed low genetic diversity due to selection for domestication in cultivated cacao (Orduña *et al.*, 2024). The results of that work may explain why similar loci are apparent across varieties for resistance to different diseases, as was found in the two studies, in Indonesia and in PNG, that informed the current research. Genome resources have, until recently, been presented as collapsed haploid genomes and the allocation of QTLs has been limited due to loss of valuable information from the two allele sets. Technologies are rapidly improving and permitting far more detailed comparative studies than were possible only a few years ago. Our original goal of assembling both the host and compatible pathogen genomes from one sample was unsuccessful due to low sequence coverage, only 1X HiFi reads, for *C. theobromae* reads. However future methods could use adaptive sequencing with the Oxford Nanopore Technologies metagenome sampling tools (Martin *et al.*, 2022). This approach is rapidly developing, with new models incorporating Bayesian dynamic adaptive sampling (Weilguny *et al.*, 2023), and providing promising tools to investigate the biology of unculturable biotrophic pathogens.

For this research we employed recent advances in technology and software to assemble and assign parental chromosomes for a VSD susceptible *T. cacao* genotype (clone 26) from the Mars Cocoa Research Centre in Pangkep, Sulawesi, Indonesia. We annotated the two parental chromosome sets independently and specifically we characterised all the predicted NLR-type resistance genes. The resulting curated and annotated genomes were compared at QTL locations that were identified to explain resistant phenotypes to VSD and PPR. Our results show that the QTLs may indeed resolve to NLR-type resistance genes, and that these may be important to investigate further for breeding trees with resistance to VSD, PPR, and potentially other diseases of cacao. Our work provides a more comprehensive analysis of the predicted NLRs and other genes on each chromosome and shows the variation in structure and type of these genes within a VSD susceptible individual.

Future work could develop molecular screening around these specific genes across both resistant and susceptible trees for a range of diseases of cacao. The current work has developed foundational genome resources to facilitate such analyses that may be used to assist future breeding programs for resistance.

Experimental procedures

High Molecular Weight DNA extraction protocol testing

Prior to our work in Sulawesi, Indonesia, we ran exhaustive high molecular weight (HMW) DNA extraction protocol testing on cacao leaf samples from plants maintained in the University of Sydney greenhouses. A persistent problem remained for clean extractions due to high contaminant levels that had the properties of a jelly-like substance, presumably pectin, though not confirmed. We attempted to resolve the nature of the contaminant with enzymatic testing using pectinase. We also tested our contaminant DNA with nuclear magnetic resonance spectrometer (NMR). Data was collected on a 600 MHz spectrometer equipped with triple resonance CryopProbe and the following buffer conditions: 10 mM Tris; 10% D2O; pH 8.0, in 3 mm short NMR tubes (volume ~160 µl). We ran controls of 10 mM Tris buffer and clean HMW DNA in Tris buffer. Test samples included pectin, pectin with pectinase treatment, contaminated DNA with pectinase treatment, contaminated DNA with macerozyme (incl. pectinase) treatment. Unfortunately the signals in the aromatic/amide regions were beyond detection, suggesting a large DNA-contaminant (potentially covalent) complex form in this sample. Young leaves were found to produce cleaner DNA but did not show any symptoms of VSD, so we aimed to extract samples from older infected leaves using a combined protocol described below that improved on but did not resolve the contaminant problem.

Cacao leaf material collected from field trial tree

We collected fully expanded leaf material on the 10 January 2023, during wet weather, from the Mars Cocoa Research Institute field trials in Pangkep, South Sulawesi, Indonesia. The leaves were found on T. *cacao* F1 generation progeny, VSD susceptible clone number 26, from Sulawesi 1 – S1 (resistant maternal), crossed with RUQ 1347 (MIS_GBR207 CCN 51) (susceptible paternal) parental plants (Turnbull et al., 2017). The leaves were symptomatic of VSD with browning of the leaf apex and vascular death visible in the petiole (Figure 5). Leaves were placed in plastic bags and transported to the laboratories at Universitas Hasannudin, Makassar and refrigerated overnight.

Figure 5 Leaf from *Theobroma cacao*, VSD susceptible clone 26, growing at the Mars Cocoa Research Institute field trials,
Pangkep South Sulawesi, Indonesia. (Left) The symptomatic leaf apex dieback and some chlorosis indicating *Ceratobasium*
theobromae*. (Right) Visible vascular death caused by the pathogen. Scale bar left ~5 cm.

High Molecular Weight DNA extracted from cacao leaves

The following day while leaves were fresh, HMW DNA was extracted using a protocol adapted

from the sorbitol wash method developed by Jones et al. (2021), and the HMW plant DNA

CTAB extraction method developed by Hilario (2018). Small (1 cm^2) pieces of the leaves were

placed into 1.5 mL centrifuge tubes with a small amount (a few mg) of PVP40000. The tissue

was flash frozen in liquid nitrogen and ground into a powder with micropestle while frozen.

Three or more rounds of sorbitol wash were used on the samples, in accordance with the

protocol (Jones et al., 2021), until the supernatant was no longer viscous. Then DNA was

extracted, according to the protocol (Hilario, 2018), using CTAB buffer with an incubation step

at 56 °C for 2 hours, in a water bath. Apart from an extended RNase A incubation step, of 10

mins, the protocol was followed until the precipitation step. The DNA formed a visible

gelatinous precipitate at this stage and was left to continue precipitating at -20 °C overnight.

The following day the DNA was collected by centrifugation at 3000 rpm for 25 min at room

temperature and cleanup steps followed according to the protocol before dissolving in 10

mM Tris-HCL pH 8.0. The DNA was quantified, and purity was checked using a NanoDrop™

Lite Plus spectrophotometer and a QuBit™ 2.0 Fluorometer. Purified HMW DNA (10 x 1.5 µL

tubes) was couriered under an Australian Department of Agriculture, Water, and the

Environment import permit to the Australian Genome Research Facility (AGRF), Brisbane,

Australia, for cleanup, size selection, library prep and sequencing on Pacific Biosciences of

California, Inc. (PacBio) Sequel II.

Formaldehyde cross-linking of leaves for HiC sequencing

374
We prepared samples according to the manufacturer's protocols for chromatin crosslinking
library prep and sequencing at Phase Genomics (Seattle, Washington, US) to obtain HiC reads.
Five grams of leaf material was immersed in 2% (v/v) sodium hypochlorite for 2 mins for
surface sterilisation and then washed with ultrapure water. The leaves were cut into strips
and immersed in 1% (v/v) formaldehyde for 40 minutes with intermittent mixing. The
crosslinking reaction was quenched with the addition of glycine to a final concentration of
125 mM for 15 minutes. The crosslinked leaves were washed with ultrapure water, dried,
then ground to a powder after freezing with liquid nitrogen.

Data processing and genome assembly

384 Details of all the software, versions, parameters and scripts used to process the data are
385 available here: https://github.com/peritob/Theobroma-cacao-genome/tree/main

Preparing the raw data for processing

We downloaded the 23 GB (ccs.bam) HiFi sequence read data from AGRF, filtered the reads
using HiFiAdapterFilt (Sim *et al.*, 2022) and checked the statistics using Nanoplot (De Coster
and Rademakers, 2023). The HiC paired-end read (25 GB) mapping statistics against the cacao
Criollo reference genome (GCF_000208745.1) were good for informative read pairs (12.18%)
but not against the pathogen genome (GCA_012932095.1) at (0.03%), although 4.28% same
strand high quality reads were mapped. We mapped all the filtered HiFi reads to the *C.*
theobromae genome (Ali *et al.*, 2019) with Minimap2 (Li, 2018) to filter the pathogen from
host. We presumed all the mapped reads to be pathogen and all the unmapped reads to be
from the host and proceeded accordingly. Additionally, we ran a local blast with internal
transcribed spacer (ITS) primers on the VSD mapped raw sequence reads. This confirmed a
single exact *in silico* hit using the ITS regions using forward primers ITS4 (White *et al.*, 1990)
and reverse ITS5A (Stanford *et al.*, 2000). Raw Illumina data available for the parental plants
was trimmed of adaptors and low quality reads with Fastp (Chen *et al.*, 2018).

Genome assembly and chromosome scaffolding

401 We attempted several assembly approaches with Hifiasm software (Cheng *et al.*, 2021).
402 Previous experience with HiC scaffolding has shown the possibility to phase the separate

nuclear compartments (Tobias *et al.*, 2022). We therefore attempted to build a plant-pathogen hybrid genome to scaffold with HiC and visually identify and separate the two organisms using the heatmaps. The hybrid genome did not complete, likely due to insufficient overlaps for the pathogen reads. We tried to assemble all the pathogen reads into consensus sequences using Flye (Kolmogorov *et al.*, 2019) and Hifiasm without success. We then tested the trio-binning option for Hifiasm of the diploid *T. cacao* incorporating the Illumina parental reads after running Yak (Li, 2020). This was unsuccessful, so we assembled the diploid *T. cacao* genome using HiC integration with Hifiasm followed by HiC scaffolding of the resulting genomes independently with Juicer (Neva C. Durand *et al.*, 2016), 3D-DNA (Dudchenko *et al.*, 2017) and manual scaffolding with Juicebox (Neva C Durand *et al.*, 2016).

Theobroma cacao genome curation

We used the Matina 1-6 cacao genome (GCF_000403535.1) to anchor our genome scaffolds with PAFScaff (Field *et al.*, 2020) based on Minimap2 (Li, 2018) mapping. We screened the resulting fasta files for non-nuclear genomic contaminants with FCS-GX (Astashyn *et al.*, 2024) and Tiara (Karlicki *et al.*, 2022). In each case, the two haplotypes were run independently. FCS-GX was run against the NCBI gxdb (build date 2023-01-24; downloaded 2023-09-12). Additional checks of contamination were executed with Taxolotl (Tobias *et al.*, 2022). Taxolotl was provided with the annotation for each haplotype and run with taxwarnrank=class using mmseqs2 against the NCBI nr database (compiled 2022-05-19). In addition to generating results for all sequences in each haplotype, analysis of subsets was performed for (1) chromosome scaffolds only, (2) pure FCS-GX scaffolds, (3) contaminated FCS-GX scaffolds, (4) Tiara "eukarya" scaffolds, and (5) scaffolds not classified as "eukarya" by Tiara. Results were visualised with Pavian. Lists were made of non-plant contaminant contigs from the fcs_gx_reports and were removed, as well as plastid contigs flagged in the tiara outputs. We then ran Telociraptor (Edwards, 2023) followed by Chromsyn (Edwards *et al.*, 2022) on the chromosome-level contigs to compare synteny and scaffolds tweaked according to the contig graphs and figure outputs. We used the parental data available within our lab (Singh, 2021) to assign scaffolds and contigs. We mapped the maternal S1 data to the concatenated haplotype A and B genomes using Hisat2 (Kim *et al.*, 2019) and then similarly mapped the RUQ 1347 data. We assumed that the most reads mapped were the likely parental chromosome/contig and curated each genome. Of the chromosomes, 3, 5 and 8 were

switched accordingly (Figure 6). We designated the S1 maternally inherited contigs as genome T_cacao_v1.3_A, and RUQ 1347 paternally inherited as T_cacao_v1.3_B.

Figure 6 *Theobroma cacao* **resistant (S1) and susceptible (RUQ 1347) parental paired-end (PE) Illumina sequence reads mapped to the concatenated Hap A and B progeny chromosomes. The reads that were mapped in greater numbers to chromosomes and contigs were used to determine parental assignment and swapped accordingly.**

Parental assigned genomes were annotated independently

To repeat mask the genome for annotation we ran both genomes through Repeatmasker and Repeatmodeler. We then mapped publicly available RNAseq data from CCN51 *Theobroma cacao* from NCBI (PRJNA933172) to the soft-masked genomes independently with hisat2. We trimmed the RNAseq data with Fastp (Chen *et al.*, 2018) prior to mapping and processed the outputs with Samtools (Li *et al.*, 2009) to make bam files. The mapped RNAseq bam files were used as evidence to predict protein coding genes within the funannotate (Palmer and Stajich, 2023) *predict* pipeline with the BUSCO embryophyte (Manni *et al.*, 2021; Simão *et al.*, 2015) database. We used Interproscan5 (Jones *et al.*, 2014) independently and then ran funannotate *annotate* to obtain final predicted proteins and coding sequences. In order to make GenBank (Sayers *et al.*, 2023) compatible annotations, we ran the resulting files with Table2asn (GenBank, 2023).

NLR-type resistance gene annotation

We annotated the resistance gene complement with the FindPlantNLRs pipeline (Chen *et al.*, 2023), that incorporates NLR-Annotator (Steuernagel *et al.*, 2020), HMMER (Finn *et al.*, 2015; Johnson *et al.*, 2010) and BLAST (Altschul *et al.*, 1990) components as well as Interproscan5

(Jones *et al.*, 2014). Unlike many annotation methods, the pipeline takes an unmasked genome as a starting point to avoid masking, and hence not annotating, repetitive LRR regions that are characteristic of NLR-type genes. The last stage of the pipeline determines NLR gene classes, including predicted integrated domains, and locations within each parental haplotype. All NLRs were plotted to visualise on chromosomes with Chromomap (Anand and Rodriguez Lopez, 2022) and classes, based on domain Pfams, were visualised with Sankeymatic (Bogart, 2014).

QTL locations on chromosomes predicted and compared against NLR-type gene predictions

Informative QTLs for resistance to VSD were obtained from data of two previous studies (Epaina, 2012; Singh, 2021). Using the chromosome lengths for our parental genome haplotypes we calculated the predicted locations for VSD resistance QTLs. In brief, we took averages of Linkage Group (LG) map lengths (Epaina, 2012), and divided the chromosome base pair length from assembled genomes (current study) by centimorgan (cM). We then determined the expected mid-point (in base pairs) for the resistance QTLs. Singh (2021) predicted QTLs confidence intervals at 3-4 Mb and determined three significant QTLs; on chromosome 3, 8 and 9. Initially our calculations for the QTL locations from the PNG progeny study showed a significant region at around 18.5 Mb on chromosome 8 (Table 3). This was not concordant with the Sulawesi study, at between 5.5 and 8.8 Mb, leading us to investigate if our calculations were misrepresenting the 'start' position for chromosome eight. When recalculated based on the reverse orientation, we found that the QTL matched more closely, at around 7 Mb. We also noted that the QTL identified on chromosome 3, when start positions were reversed, indicated a cluster of NLR-type genes at around 30 Kb. Several significant QTLs were associated with resistance to other pathogens including PPR (Table 3) in the study by Epaina (2012). QTLs that were in similar regions within both studies were regarded as likely harbouring useful candidate genes for breeding resistance. The most closely aligned VSD QTLs between studies were on chromosomes 3, 8 and 9. Detailed comparisons for the VSD significant QTLs across the two studies were undertaken to investigate resistance gene homologs and other predicted genes across the parental chromosomes.

Table 3 QTLs for vascular streak dieback (VSD) resistance were calculated using average *Theobroma cacao* Linkage Group (LG) lengths (Epaina, 2012) and dividing chromosome lengths (bp) by LG lengths (in cM) to determine 1 cM in bp (per chromosome). QTLs (cM) were then multiplied by bp length for 1 cM to obtain the chromosome locations for the diploid genome. Green cells and brown cells are VSD and *Phytophthora* pod rot (PPR) resistance QTL physical locations on chromosomes.

Theobroma cacao A (S1 inherited)

Chromosomes (LG)		1	2	3	4	5	6	7	8	9	10
cM (average)		99.9	107.2	91.6	79.7	89.8	75.8	56.6	60	106.1	62.4
bp per chromosome (this study)		39,005,981	41,965,441	38,945,508	33,857,508	40,959,067	25,938,454	26,721,266	25,579,687	40,159,845	25,091,101
bp per cM		390,450	391,469	425,169	424,812	456,114	342,196	472,107	426,328	378,509	402,101
Singh (2021)	VSD QTL (bp)			9,353,725 (*29,591,783)					8,782,359	25,170,874	
Epaina (2012)	VSD QTL (bp)	11,270,737	26,861,405	8,531,872 (*30,413,636)	27,163,747	7,976,071			18,514,151 (*7,065,536)	31,452,994	
Epaina (2012)	PPR QTL (bp)			8,531,872 (*30,413,636)	23,158,620	6,937,499			18,514,151 (*7065,536)		

Theobroma cacao B (RUQ 1347 inherited)

Chromosomes (LG)		1	2	3	4	5	6	7	8	9	10
bp per chromosome (this study)		39,156,173	42,761,000	39,170,102	34,430,140	41,360,500	29,284,707	25,927,970	21,951,902	41,168,245	24,929,739
bp per cM		391,954	398,890	427,621	431,997	460,585	386,342	458,091	365,865	388,014	399,515
Singh (2021)	VSD QTL (bp)			9,407,666 (*29,762,436)					7,536,820	25,802,906	
Epaina (2012)	VSD QTL (bp)	11,314,135	27,370,630	8,581,075 (*30,589,027)	27,623,167	8,054,243			15,888,421 (*6,063,481)	32,242,768	
Epaina (2012)	PPR QTL (bp)			8,581,075 (*30,589,027)	23,550,302	7,005,492			15,888,421 (*6,063,481)		

*QTL determined with start position on chromosome reversed.

Acknowledgements

The Joint Cocoa Research Fund, European Cocoa Association supported this research. The authors acknowledge the facilities, and the scientific and technical assistance of the Sydney Informatics Hub at the University of Sydney and, in particular, access to the high-performance computing facility Artemis. We acknowledge the support of Sydney Analytical for running samples and interpretation on the University of Sydney Nuclear Magnetic Resonance (NMR) Core Facilities. Thank you to Mark Powrie who generously assisted during the work in Sulawesi.

Author contributions

DG and PT initiated and led the project. PT and JD collected the material from established clonal trials developed and managed by AP. JM and AP supported and facilitated the research in Sulawesi. JD, PT, AZ, EB extracted HMW DNA and cross-linked samples. PT and EB organised sample deliveries. PE and GS provided results of QTL trials that were supported by DG and RP. GS provided parental data. PT assembled, curated, annotated the genomes, and interpreted results, RE ran contamination screening of genomes. PT submitted the data to NCBI and wrote the manuscript. All authors contributed comments and edits to the manuscript.

Data availability statement

All data that support the findings of this research are publicly available. Raw data is available at National Centre for Biotechnology Information (NCBI) under the following biosample accession: SAMN40241783 and bioproject accession: PRJNA1083235. Annotated diploid *Theobromae cacao* (clone 26) genomes are available here: PRJNA1083235 (maternally inherited), PRJNA1086984 (paternally inherited). NLR predicted coding and amino acid sequences and gff3 data is available at https://github.com/peritob/Theobroma-cacao-genome/tree/main.

References

Ali, S.S., Asman, A., Shao, J., et al. (2019) Draft genome sequence of fastidious pathogen Ceratobasidium theobromae, which causes vascular-streak dieback in Theobroma cacao. *Fungal Biol. Biotechnol.* **6**, 1–10.

Altschul, S.F., Gish, W., Miller, W., Myers, E.W. and Lipman, D.J. (1990) Basic local alignment search tool. *J. Mol. Biol.* **215**, 403–10.

519   **Anand, L. and Rodriguez Lopez, C.M.** (2022) ChromoMap: an R package for interactive
520   visualization of multi-omics data and annotation of chromosomes. *BMC Bioinformatics*
521   **23**, 1–9.

522   **Argout, X., Salse, J., Aury, J.-M., et al.** (2011) The genome of Theobroma cacao. *Nat. Genet.*
523   **43**, 101–108.

524   **Arya, P., Kumar, G., Acharya, V. and Singh, A.K.** (2014) Genome-wide identification and
525   expression analysis of NBS-encoding genes in *Malus* x *domestica* and expansion of *NBS*
526   genes family in Rosaceae. *PLoS One* **9**, e107987.

527   **Astashyn, A., Tvedte, E.S., Sweeney, D., et al.** (2024) Rapid and sensitive detection of genome
528   contamination at scale with FCS-GX. *Genome Biol.* **25**, 1–25.

529   **Bailey, B.A. and Meinhardt, L.W. eds** (2016) Cacao Diseases: A History of Old Enemies and
530   New Encounters. In Cacao Diseases: A History of Old Enemies and New Encounters.,
531   Heidelberg: Springer International Publishing.

532   **Barragan, A.C. and Weigel, D.** (2020) Plant NLR Diversity : The Known Unknowns of Pan-
533   NLRomes A Brief History of Plant NLRs. *Plant Cell*.

534   **Bhunjun, C.S., Phillips, A.J.L., Jayawardena, R.S., Promputtha, I. and Hyde, K.D.** (2021)
535   Importance of molecular data to identify fungal plant pathogens and guidelines for
536   pathogenicity testing based on Koch'S postulates. *Pathogens* **10**, 1–18.

537   **Bogart, S.** (2014) SankeyMatic. https://sankeymatic.com/build/

538   **Chen, S., Zhou, Y., Chen, Y. and Gu, J.** (2018) Fastp: An ultra-fast all-in-one FASTQ
539   preprocessor. *Bioinformatics* **34**, i884–i890.

540   **Chen, S.H., Martino, A.M., Luo, Z., Schwessinger, B., Jones, A., Tolessa, T., Bragg, J.G.,**
541   **Tobias, P.A. and Edwards, R.J.** (2023) A high-quality pseudo-phased genome for
542   Melaleuca quinquenervia shows allelic diversity of NLR-type resistance genes.
543   *Gigascience* **12**, 1–18.

544   **Cheng, H., Concepcion, G.T., Feng, X., Zhang, H. and Li, H.** (2021) Haplotype-resolved de novo
545   assembly using phased assembly graphs with hifiasm. *Nat. Methods* **18**, 170–175.

546   **Christie, N., Tobias, P.A., Naidoo, S. and Külheim, C.** (2016) The *Eucalyptus grandis NBS-LRR*
547   gene family: physical clustering and expression hotspots. *Front. Plant Sci.* **6**, 1–16.

548   **Contreras, M.P., Pai, H., Tumtas, Y., et al.** (2023) Sensor NLR immune proteins activate
549   oligomerization of their NRC helpers in response to plant pathogens . *EMBO J.* **42**.

550   **Coster, W. De and Rademakers, R.** (2023) NanoPack2: population-scale evaluation of long-
551   read sequencing data. *Bioinformatics* **39**, 0–2.

552   **Daymond, A.J., Mendez, D.G., Hadley, P., Bastide, P., Abdoellah, S., Acheampong, K.,**
553   **Amores-Puyutaxi, F.M., Anhert, D. and Konan, D.C.** (2022) *A Global Review of Cocoa*
554   *Farming Systems,*.

555   **Dudchenko, O., Batra, S.S., Omer, A.D., et al.** (2017) De novo assembly of the Aedes aegypti
556   genome using Hi-C yields chromosome-length scaffolds. *Science (80-.).* **356**, 92–95.

557   **Durand, Neva C, Robinson, J.T., Shamim, M.S., et al.** (2016) Juicebox Provides a Visualization
558   System for Hi-C Contact Maps with Unlimited Zoom Tool. *Cell Syst.* **3**, 99–101.

559   **Durand, Neva C, Shamim, M.S., Machol, I., Rao, S.S.P., Huntley, M.H., Lander, E.S. and**
560   **Aiden, E.L.** (2016) Juicer Provides a One-Click System for Analyzing Loop-Resolution Hi-C
561   Experiments. *Cell Syst.* **3**, 95–98.

562   **Edwards, R.J.** (2023) Telociraptor. https://github.com/slimsuite/telociraptor

563   **Edwards, R.J., Dong, C., Park, R.F. and Tobias, P.A.** (2022) A phased chromosome-level
564   genome and full mitochondrial sequence for the dikaryotic myrtle rust pathogen,
565   Austropuccinia psidii. *bioRxiv*, 2022.04.22.489119.

Epaina, P. (2012) *Identification of Molecular Markers and Quantitative Trait Loci linked to*
resistance to Vascular Streak Dieback and Phytophthora Pod Rot of Cacao (Theobroma
cacao L .). The University of Sydney.

Ercoli, M.F., Luu, D.D., Rim, E.Y., et al. (2022) Plant immunity: Rice XA21-mediated resistance
to bacterial infection. *Proc. Natl. Acad. Sci. U. S. A.* **119**.

Field, M.A., Rosen, B.D., Dudchenko, O., et al. (2020) Canfam-GSD: De novo chromosome-
length genome assembly of the German Shepherd Dog (Canis lupus familiaris) using a
combination of long reads, optical mapping, and Hi-C. *Gigascience* **9**, 1–12.

Finn, R.D., Clements, J., Arndt, W., Miller, B.L., Wheeler, T.J., Schreiber, F., Bateman, A. and
Eddy, S.R. (2015) HMMER web server: 2015 Update. *Nucleic Acids Res.* **43**, W30–W38.

Fluhr, R., King, S.I., Vi, H. and Shakespeare, W. (2001) Sentinels of Disease . Plant Resistance
Genes. **127**, 1367–1374.

GenBank (2023) Table2asn. https://www.ncbi.nlm.nih.gov/genbank/table2asn

Guest, D. and Keane, P. (2007) Vascular-streak dieback: A new encounter disease of cacao in
Papua New Guinea and Southeast Asia caused by the obligate basidiomycete
Oncobasidium theobromae. *Phytopathology* **97**, 1654–1657.

Gurevich, A., Saveliev, V., Vyahhi, N. and Tesler, G. (2013) QUAST: Quality assessment tool
for genome assemblies. *Bioinformatics* **29**, 1072–1075.

Hall, T.A. (1991) BioEdit: a user-friendly biological sequence alignment editor and analysis
program for Windows 95/98/NT. *Nucleic Acids Symp. Ser.* **41**, 95–98.

Hilario, E. (2018) Plant nuclear genomic DNA preps. *protocols.io:*

Johnson, L.S., Eddy, S.R. and Portugaly, E. (2010) Hidden Markov model speed heuristic and
iterative HMM search procedure. *BMC Bioinformatics* **11**.

Jones, A., Torkel, C., Stanley, D., Nasim, J., Borevitz, J. and Schwessinger, B. (2021) High-
molecular weight DNA extraction, cleanup and size selection for long-read sequencing.
PLoS One **16**, 1–6.

Jones, P., Binns, D., Chang, H.Y., et al. (2014) InterProScan 5: Genome-scale protein function
classification. *Bioinformatics* **30**, 1236–1240.

Junaid, M., Purwantara, A. and Guest, D. (2020) First report of vascular streak dieback
symptom of cocoa caused by Ceratobasidium theobromae in Barru District, South
Sulawesi. *IOP Conf. Ser. Earth Environ. Sci.* **486**.

Karlicki, M., Antonowicz, S. and Karnkowska, A. (2022) Tiara: deep learning-based
classification system for eukaryotic sequences. *Bioinformatics* **38**, 344–350.

Keane, P.J., Flentje, N.J. and Lamb, K.P. (1972) Investigation of vascular-streak dieback of
cocoa in papua new guinea. *Aust. J. Biol. Sci.* **25**, 553–564.

Kim, D., Paggi, J.M., Park, C., Bennett, C. and Salzberg, S.L. (2019) Graph-based genome
alignment and genotyping with HISAT2 and HISAT-genotype. *Nat. Biotechnol.* **37**, 907–
915.

Kohler, A., Rinaldi, C., Duplessis, S., Baucher, M., Geelen, D., Duchaussoy, F., Meyers, B.C.,
Boerjan, W. and Martin, F. (2008) Genome-wide identification of NBS resistance genes
in Populus trichocarpa. *Plant Mol. Biol.* **66**, 619–636.

Kolmogorov, M., Yuan, J., Lin, Y. and Pevzner, P.A. (2019) Assembly of long, error-prone
reads using repeat graphs. *Nat. Biotechnol.* **37**, 540–546.

Kuhn, D.N., Narasimhan, G., Nakamura, K., Brown, J.S., Schnell, R.J. and Meerow, A.W.
(2006) Identification of cacao TIR-NBS-LRR resistance gene homologues and their use as
genetic markers. *J. Am. Soc. Hortic. Sci.* **131**, 806–813.

Lanaud, C., Fouet, O., Clément, D., Boccara, M., Risterucci, A.M., Surujdeo-Maharaj, S.,

Legavre, T. and Argout, X. (2009) A meta-QTL analysis of disease resistance traits of Theobroma cacao L. *Mol. Breed.* **24**, 361–374.

Li, H. (2018) Minimap2: Pairwise alignment for nucleotide sequences. *Bioinformatics* **34**, 3094–3100.

Li, H. (2020) Yak.

Li, H., Handsaker, B., Wysoker, A., Fennell, T., Ruan, J., Homer, N., Marth, G., Abecasis, G. and Durbin, R. (2009) The Sequence Alignment/Map format and SAMtools. *Bioinformatics* **25**, 2078–2079.

Manni, M., Berkeley, M.R., Seppey, M., Simão, F.A. and Zdobnov, E.M. (2021) BUSCO Update: Novel and Streamlined Workflows along with Broader and Deeper Phylogenetic Coverage for Scoring of Eukaryotic, Prokaryotic, and Viral Genomes. *Mol. Biol. Evol.* **38**, 4647–4654.

Mao, Y. and Zhang, G. (2022) A complete, telomere-to-telomere human genome sequence presents new opportunities for evolutionary genomics. *Nat. Methods* **19**, 635–638.

Marchal, C., Michalopoulou, V.A., Zou, Z., Cevik, V. and Sarris, P.F. (2022) Show me your ID: NLR immune receptors with integrated domains in plants. *Essays Biochem.* **66**, 527–539.

Marelli, J.P., Guest, D.I., Bailey, B.A., Evans, H.C., Brown, J.K., Junaid, M., Barreto, R.W., Lisboa, D.O. and Puig, A.S. (2019) Chocolate under threat from old and new cacao diseases. *Phytopathology* **109**, 1331–1343.

Martin, S., Heavens, D., Lan, Y., Horsfield, S., Clark, M.D. and Leggett, R.M. (2022) Nanopore adaptive sampling: a tool for enrichment of low abundance species in metagenomic samples. *Genome Biol.* **23**, 1–27.

Mistry, J., Chuguransky, S., Williams, L., et al. (2021) Pfam: The protein families database in 2021. *Nucleic Acids Res.* **49**, D412–D419.

Motamayor, J.C., Lachenaud, P., Silva e Mota, J.W. da, Loor, R., Kuhn, D.N., Brown, J.S. and Schnell, R.J. (2008) Geographic and genetic population differentiation of the Amazonian chocolate tree (Theobroma cacao L). *PLoS One* **3**.

Motamayor, J.C., Schnell, R., Kuhn, D., et al. (2013) The genome sequence of the most widely cultivated cacao type and its use to identify candidate genes regulating pod color. *Genome Biol.* **14**, r53.

Nousias, O., Zheng, J., Li, T., et al. (2024) Three de novo assembled wild cacao genomes from the Upper Amazon. *Sci. data* **11**, 1–9.

Orduña, H.E.N., Müller, M., Krutovsky, K. V and Gailing, O. (2024) Genotyping of cacao (Theobroma cacao L .) germplasm resources with SNP markers linked to agronomic traits reveals signs of selection. *Tree Genet. Genomes*.

Palmer, J.M. and Stajich, J.E. (2023) Funannotate.

Richardson, J.E., Whitlock, B.A., Meerow, A.W. and Madriñán, S. (2015) The age of chocolate: A diversification history of Theobroma and Malvaceae. *Front. Ecol. Evol.* **3**, 1–14.

Samuels, G.J., Ismaiel, A., Rosmana, A., et al. (2012) Vascular Streak Dieback of cacao in Southeast Asia and Melanesia: In planta detection of the pathogen and a new taxonomy. *Fungal Biol.* **116**, 11–23.

Santos, M. de L., Resende, M.L.V. de, Alves, G.S.C., Huguet-Tapia, J.C., Resende, M.F.R. de J. and Brawner, J.T. (2022) Genome-Wide Identification, Characterization, and Comparative Analysis of NLR Resistance Genes in Coffea spp. *Front. Plant Sci.* **13**, 1–16.

Sayers, E.W., Cavanaugh, M., Clark, K., Pruitt, K.D., Sherry, S.T., Yankie, L. and Karsch-Mizrachi, I. (2023) GenBank 2023 update. *Nucleic Acids Res.* **51**, D141–D144.

Schnell, R.J., Kuhn, D.N., Brown, J.S., Olano, C.T., Amores, F.M. and Motamayor, J.C. (2007) Development of a Marker Assisted Selection Program for Cacao. In Cacao Diseases : Important Threats to Chocolate Production Worldwide., Phytopathology.

Sergey, N., Sergey, K., Arang, R., et al. (2022) The complete sequence of a human genome. *Science (80-.).* **376**, 44–53.

Sim, S.B., Corpuz, R.L., Simmonds, T.J. and Geib, S.M. (2022) HiFiAdapterFilt, a memory efficient read processing pipeline, prevents occurrence of adapter sequence in PacBio HiFi reads and their negative impacts on genome assembly. *BMC Genomics* **23**.

Simão, F.A., Waterhouse, R.M., Ioannidis, P., Kriventseva, E. V. and Zdobnov, E.M. (2015) BUSCO: Assessing genome assembly and annotation completeness with single-copy orthologs. *Bioinformatics* **31**, 3210–3212.

Singh, G. (2021) *Identification of Quantitative Trait Loci (QTL) associated with resistance to Vascular Streak Dieback disease of cacao*. The University of Sydney.

Sniezko, R.A. (2006) Resistance breeding against nonnative pathogens in forest trees — current successes in North America 1. *Can. J. Plant Pathol.* **279**, 270–279.

Stanford, A.M., Harden, R. and Parks, C.R. (2000) Phylogeny and biogeography of Juglans (Juglandaceae) based on matK and ITS sequence data. *Am. J. Bot.* **87**, 872–882.

Steuernagel, B., Witek, K., Krattinger, S.G., et al. (2020) The NLR-annotator tool enables annotation of the intracellular immune receptor repertoire. *Plant Physiol.* **183**, 468–482.

Sun, J., Lu, F., Luo, Y., Bie, L., Xu, L. and Wang, Y. (2023) OrthoVenn3: An integrated platform for exploring and visualizing orthologous data across genomes. *Nucleic Acids Res.* **51**, W397–W403.

Tan, G.Y. and Tan, W.K. (1988) Genetic variation in resistance to vascular-streak dieback in cocoa (Theobroma cacao). *Theor. Appl. Genet.* **75**, 761–766.

Thompson, J.D., Higgins, D.G. and Gibson, T.J. (1994) CLUSTAL W: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* **22**, 4673–4680.

Tobias, P.A., Edwards, R.J., Surana, P., Mangelson, H., Inácio, V., Céu Silva, M. do, Várzea, V., Park, R.F. and Batista, D. (2022) A chromosome-level genome resource for studying virulence mechanisms and evolution of the coffee rust pathogen Hemileia vastatrix *bioRxiv*, 2022.07.29.502101.

Tobias, P.A. and Guest, D.I. (2014) Tree immunity: growing old without antibodies. *Trends Plant Sci.* **19**, 367–370.

Trisno, J., Reflin, R. and Martinius, M. (2016) Vascular Streak Dieback: Penyakit Baru Tanaman Kakao di Sumatera Barat. *J. Fitopatol. Indones.* **12**, 142.

Weilguny, L., Maio, N. De, Munro, R., Manser, C., Birney, E., Loose, M. and Goldman, N. (2023) Dynamic, adaptive sampling during nanopore sequencing using Bayesian experimental design. *Nat. Biotechnol.* **41**, 1018–1025.

Weyer, A.-L. van de, Monteiro, F., Furzer, O.J., et al. (2019) The Arabidopsis thaliana pan-NLRome. *bioRxiv*, 1–32.

Weyer, A.L. Van de, Monteiro, F., Furzer, O.J., et al. (2019) A Species-Wide Inventory of NLR Genes and Alleles in Arabidopsis thaliana. *Cell* **178**, 1260-1272.e14.

White, Bruns T, Lee S and Taylor, J. (1990) Amplification and direct sequencing of fungal ribosomal RNA genes for phylogenetics. In PCR Protocols: A Guide to Methods and Application., pp. 315–322. San Diego, California: Academic Press.

Xi, Y., Cesari, S. and Kroj, T. (2022) Insight into the structure and molecular mode of action of plant paired NLR immune receptors. *Essays Biochem.* **66**, 513–526.

Figure legends

Figure 1 Chromsyn (Edwards *et al.*, 2022) synteny plots of *Theobroma cacao* parentally phased genome, haplotype A (top) and B (below). Synteny blocks of collinear "Complete" BUSCO genes (Simão *et al.*, 2015) link scaffolds from adjacent assemblies: blue, same strand; red, inverse strand. Yellow triangles mark "Duplicated" BUSCOs. Filled circles mark telomere predictions from Telociraptor (black) (Edwards, 2023). Assembly gaps are marked as dark red + signs**.**

Figure 2 Physical locations of predicted NLR genes on the paired chromosomes of *Theobroma cacao* (A) haplotype A (red) and (B) haplotype B (yellow) generated using ChromoMap (Anand and Rodriguez Lopez, 2022) in RStudio.

Figure 3 The predicted NLR gene complement in the parentally phased *Theobroma cacao* genome. The two sets of chromosomes corresponding to (above) Haplotypes A and (below) B were independently classified (Chen *et al.*, 2023) and visualised to present the domain classes using SankeyMatic (Bogart, 2014) including novel integrated domains (IDs) with abbreviations derived from Pfam database (Mistry *et al.*, 2021). NBARC = Nucleotide Binding Domain, TIR = Toll/Interleukin-1 receptor, Rx = Potato CC-NB-LRR protein Rx, CC=Coil-Coil Domain, RPW8 = RESISTANCE TO POWDERY MILDEW 8-like coiled-coil**.**

Figure 4 Amino acid sequence alignment of an NLR cluster matching the QTL mapped regions at around 29 Mb on chromosome 3 (above) and single NLR homologs at 31 Mb chromosome 9 (below) within both parentally inherited alleles, A and B, for *Theobroma cacao* clone 26. All predicted NLRs are annotated as Rx-NLRs (Chen *et al.*, 2023) named for resistance to potato virus X.

Figure 5 Leaf from *Theobroma cacao*, VSD susceptible clone 26, growing at the Mars Cocoa Research Institute field trials, Pangkep South Sulawesi, Indonesia. (Left) The symptomatic leaf apex dieback and some chlorosis indicating *Ceratobasium theobromae*. (Right) Visible vascular death caused by the pathogen. Scale bar left ~5 cm**.**

739 Figure 6 *Theobroma cacao* resistant (S1) and susceptible (RUQ 1347) parental paired-end (PE)

740 Illumina sequence reads mapped to the concatenated Hap A and B progeny chromosomes.

741 The reads that were mapped in greater numbers to chromosomes and contigs were used to

742 determine parental assignment and swapped accordingly.

Abbreviations

NLR – nucleotide binding site leucine-rich repeat gene models

LRR – leucine-rich repeat domain

GB – gigabyte

Gb – Gigabase

Mb – Megabase

kb – kilobase

VSD – vascular streak dieback, causal agent *Ceratobasidium theobromae*

PPR – Phytophthora Pod Rot, causal agent *Phytophthora palmivora*