1 **Title: Distinct mechanisms of non-autonomous UPR[ER] mediated by GABAergic,**
2 **glutamatergic, and octopaminergic neurons.**
3

4 Aeowynn J. Coakley[1,*], Adam J. Hruby[1,*], Jing Wang[1], Andrew Bong[1], Tripti Nair[1], Carmen M.
5 Ramos[1], Athena Alcala[1], Daniel Hicks[1], Maxim Averbukh[1], Naibedya Dutta[1], Darius Moaddeli[1],
6 Cynthia Siebrand[1,2], Mattias de los Rios Rogers[3], Arushi Sahay[4], Sean P. Curran[1], Peter J.
7 Mullen[5], Bérénice A Benayoun, Gilberto Garcia[1,^], Ryo Higuchi-Sanabria[1,^].
8
[1]Leonard Davis School of Gerontology, University of Southern California, Los Angeles, CA
90089, United States.
[2]Current: Buck Institute for Research on Aging, Novato, CA, 94945, USA.
[3]Department of Molecular, Cell, and Developmental Biology, University of California, Los
Angeles, CA 90095, United States.
[4]Department of Cell & Molecular Biology, University of Pennsylvania, Philadelphia, PA 19104,
United States.
[5]Department of Molecular Microbiology and Immunology, Keck School of Medicine, University of
Southern California, Los Angeles, CA 90089, United States.
*These authors contributed equally to the work.
^Correspondence to ryo.sanabria@usc.edu, ggarcia1@usc.edu

20
21 **Keywords:** Aging, stress, ER, neurons.
22

Abstract

The capacity to deal with stress declines during the aging process, and preservation of cellular
stress responses is critical to healthy aging. The unfolded protein response of the endoplasmic
reticulum (UPR[ER]) is one such conserved mechanism, which is critical for the maintenance of
several major functions of the ER during stress, including protein folding and lipid metabolism.
Hyperactivation of the UPR[ER] by overexpression of the major transcription factor, *xbp-1s*, solely
in neurons drives lifespan extension as neurons send a neurotransmitter-based signal to other
tissue to activate UPR[ER] in a non-autonomous fashion. Previous work identified serotonergic,
dopaminergic, and tyraminergic neurons in this signaling paradigm. To further expand our
understanding of the neural circuitry that underlies the non-autonomous signaling of ER stress,
we activated UPR[ER] solely in glutamatergic, octopaminergic, and GABAergic neurons in *C.*
elegans and paired whole-body transcriptomic analysis with functional assays. We found that
UPR[ER]-induced signals from glutamatergic neurons increased expression of canonical protein
homeostasis pathways and octopaminergic neurons promoted pathogen response pathways;
while minor, statistically significant changes were observed in lipid metabolism-related genes
with GABAergic UPR[ER] activation. These findings provide further evidence for the distinct role
neuronal subtypes play in driving the diverse response to ER stress.

Introduction

43
Regulation of organelle homeostasis is essential for maintenance of cellular health, which has
direct implications for organismal health and longevity. The endoplasmic reticulum (ER) is one
such organelle, which processes about a third of the proteins and lipids in cells and has
dedicated quality control machineries to preserve these and numerous other functions. One

primary quality control machinery is the ER unfolded protein response (UPR[ER]), a transcriptional
response to ER damage or stress, which activates genes essential for maintenance of proper
ER function[1]. Activation of the UPR[ER] involves the ER-membrane protein, inositol-requiring
enzyme 1 (IRE-1), which dimerizes upon sensing ER stress to splice X-box binding protein
(*xbp-1*) mRNA into *xbp-1s*. *xbp-1s* mRNA encodes a functional transcription factor, XBP-1s,
which activates genes essential for restoring ER homeostasis, including protein chaperones,
autophagy, and the ubiquitin proteosome system, among others. This transcriptional response
to stress is essential for maintaining proper function of the ER and has direct implications in
longevity. Specifically, UPR[ER] function has been shown to decline with age, and heightened
activation of UPR[ER] maintained stress resilience at old age in *C. elegans*[2].

Overexpression of *xbp-1s* solely in neurons is sufficient to enhance *C. elegans* lifespan due to a
whole-body UPR[ER] activation through neuron-to-body communication mediated by
neurotransmitter signaling[2]. Upon neuronal UPR[ER] activation, a complex signaling event
mediated by a combination of dopamine, serotonin[3], and tyramine[4], results in dramatic
remodeling of peripheral cells. Specifically, intestinal cells can activate proteostasis[4], lipid
metabolism[5,6], and lysosomal function[7] to drive longevity. These studies revealed numerous
neuronal subtypes and distinct mechanistic pathways, including chaperone induction
downstream of serotonergic signaling, lipid remodeling through lipophagy downstream of
dopaminergic signaling, and proteostasis machinery through tyramine signaling from RIM and
RIC neurons, all of which are essential to promote longevity. Finally, this UPR[ER] signaling is not
limited to neurons, as several glial subtypes were also capable of eliciting a glia-to-body UPR[ER]
signaling event to promote longevity[8].

Similar homeostatic benefits of UPR[ER] in neurons are observed in mammals, wherein *Xbp1s*
expression in pro-opiomelanocortin (POMC) neurons has been shown to protect against diet-
induced obesity by improving leptin and insulin sensitivity under ER stress[9]. While all these
studies utilized an artificial transgenic expression system, two recent studies have shown that
neuron-to-body UPR[ER] signaling is also occurs in endogenous pathways. In mice, olfactory
perception of food is sufficient to promote POMC *Xbp1s* expression and activation of post-
prandial liver ER adaption[10]. In *C. elegans*, chemosensation of pathogenic bacteria was found to
promote neuronal *xbp1-s* expression, leading to UPR[ER] activation in peripheral tissues and
extension of lifespan[11]. These studies revealed that endogenous neuron-to-body signaling
utilized similar mechanistic pathways to *xbp-1s* overexpression paradigms, which highlight the
translatability of using transgenic approaches to dissect the neuronal circuitry of UPR[ER]
signaling.

Building on previous research, we were interested in understanding whether other neuronal
subtypes are involved in neuron-to-body UPR[ER] activity. We sought to determine whether
glutamatergic, GABAergic, and octopaminergic neurons are necessary and/or sufficient to drive
neuron-to-body communication of the UPR[ER] in *C. elegans*. We accomplished this by *xbp-1s*
overexpression in these neuronal subtypes and assessing measurements of general health,
such as lifespan, healthspan, ER function, and stress resilience. Further, we performed a
comprehensive transcriptomic analysis to identify potential mechanistic pathways that drive
phenotypic outcomes in these neuronal subtype UPR[ER] paradigms.

Materials and Methods

C. elegans maintenance

All strains utilized in this investigation are derived from the N2 wild-type worm sourced from the
Caenorhabditis Genetics Center (CGC) and are detailed below. The worms are maintained at
15°C, fed with OP50 *E. coli* B strain. Animals are bleached and L1 arrested as described below
for all experimentation and transferred to growth conditions at 20°C, utilizing HT115 *E. coli* K
strain for all experiments. Experiments employed HT115 bacteria carrying an empty pL4440
vector referred to as empty vector (EV). An important note is that the *tbh-1p::xbp-1s* animals
represent a bimodal population, where some animals have severely stunted growth. Those with
stunted growth also exhibit a significant decrease in lifespan, so for the majority of the
manuscript, we opted to select for the animals that are similar in size to wild-type animals, which
we call "normal" sized for ease. Wherever relevant, data is represented as "mixed" when we did
not separate the "normal" sized versus the "small" sized animals and labeled as "small" when
animals with stunted growth were isolated. Wherever data is presented simply as "*tbh-1*", these
are the "normal" sized animals.
110

Plates
112 Standard NGM plates for maintenance using OP50 contained the following: Bacto-Agar (Difco)
113 2% w/v, Bacto Peptone 0.25% w/v, $NaCl_2$ 0.3% w/v, 1 mM $CaCl_2$, 5 µg/ml cholesterol, 0.625
114 mM KPO_4 pH 6.0, 1 mM $MgSO_4$.
115
116 Standard NGM plates for experimental condition using HT115 bacteria contained the following:
117 2% RNAi plates for experiments contained the following: Bacto-Agar (Difco) 2% w/v, Bacto
118 Peptone 0.25% w/v, $NaCl_2$ 0.3% w/v, 1 mM $CaCl_2$, 5 µg/ml cholesterol, 0.625 mM KPO_4 pH 6.0,
119 1 mM $MgSO_4$, 100 µg/mL carbenicillin, 1 mM IPTG.
120
121 For all aging experiments, 100 µL of 10 mg/mL (+)-5-Fluorodeoxyuridine (FUDR) was placed
122 directly on the bacterial lawn and animals are moved onto FUDR-containing plates at day 1 of
123 adulthood. FUDR is not used for plates containing tunicamycin, as tunicamycin prohibits growth
124 of progeny and thus use of FUDR is not necessary[12].
125

Bleaching
Experiments were conducted on animals of the same age, synchronized using a standard
bleaching protocol. Worms were collected into a 15 mL conical tube using M9 solution (22 mM
KH_2PO_4 monobasic, 42.3 mM $NaHPO_4$, 85.6 mM NaCl, 1 mM $MgSO_4$) and subjected to a
bleaching solution (1.8% sodium hypochlorite, 0.375 M NaOH in M9) until complete digestion of
carcasses. Intact eggs were then washed four times with M9 solution by centrifugation at 1,100
x g for 30 seconds. After the final wash, animals were L1 arrested by incubating overnight in M9
at 20°C on a rotator for a maximum of 24 hours.
134

Transgenic strain synthesis
The sequence for *xbp-1s* expression was defined as previously described[2] and is provided
below. The *xbp-1s* coding sequence was cloned from cDNA synthesized via reverse
transcriptase using RNA isolated from N2 worms, the endogenous *eat-4*, *tbh-1*, and *unc-25*
promoter was cloned from gDNA isolated from N2 worms, and an *unc-54* 3′UTR was cloned
from gDNA isolated from N2 worms. Plasmids were injected into N2 worms using a standard
microinjection protocol as described[13] with 10 ng/µl of overexpression plasmid, 2.5 ng/µl of *myo-*
2p::mCherry or *myo-2p::GFP* as a co-injection marker. Both injections and integration of
constructs were performed by SUNY Biotech. All integrated animals were then backcrossed to
our N2 lines >8 times to eliminate mutations and create an isogenic line. All sequences used in
this manuscript are as follows:
146

xbp-1s
ATGAGCAACTATCCAAAACGTATTTATGTGCTCCCAGCACGCCACGTGGCAGCGCCACAG
CCTCAGAGAATGGCTCCCAAGCGTGCACTTCCAACAGAACAAGTTGTCGCACAACTTCTTG
GCGATGATATGGGACCATCTGGGCCACGCAAAAGAGAACGACTGAATCATTTGAGTCAGG
AGGAGAAAATGGATCGTCGGAAACTTAAAAATCGAGTCGCAGCCCAAAATGCTAGAGACAA
AAAGAAGGAAAGATCAGCAAAGATCGAGGATGTGATGCGCGATCTGGTGGAGGAGAACCG
CCGGCTCCGCGCTGAAAACGAACGTCTTCGCCGTCAAATAAAAATCTTATGAACCAGCAG
AACGAGTCCGTCATGTATATGGAAGAGAACAACGAAAACTTGATGAACAGCAATGATGCAT
GCATCTACCAGAACGTCGTCTACGAAGAAGAAGTCGTCGGTGAGGTTGCACCAGTTGTCG
TCGTCGGAGGAGAGGATCGCCGTGCCTTTGAATCAGCAGTGGGAACAGGCCCGATCCAC
CTCCATCAACAACAACATCAGCAACCAACTCCGTCGTATGGATTCCAAGAAGAACAACACA
ATCAGTGTGGATATGTATCTAACTATCATCTCGATTCTATGCAACCACATGGATCGCAACAA
GAAGATGGACACCTCGAACAAATCCTCGAACATCTCAAGAGCCCAAGCGGAGAGTTCGAT
CGATTCGTTGCTGGCTACATTGAGGAAGGAGCAGACGGTTATGCAGCGTCTTGTTCAAGC
GGATCCATGTACACATCTTCAGAAACGCGTGAAACACTTTCGCCGAATTCCCTAGCCATGT
CCCCGTCGATGAGCAGCTCGAGCACTGACTGGGATGATGAGCTTTTGGGATGTGGAACCG
AAACTGGAACTGGAACCGACGAGCTGCTTACCGACCCCGGAAACTGGAACTTTGAAACTTT
CGACGAAAATTCAATCGACCTAAATTTCTTCCAAAATTAA
165
unc-54 UTR
CATCTCGCGCCCGTGCCTCTGACTTCTAAGTCCAATTACTCTTCAACATCCCTACATGCTCT
TTCTCCCTGTGCTCCCACCCCCTATTTTTGTTATTATCAAAAAACTTCTCTTAATTTCTTTGTT
TTTTAGCTTCTTTTAAGTCACCTCTAACAATGAAATTGTGTAGATTCAAAAATAGAATTAATT
CGTAATAAAAAGTCGAAAAAAATTGTGCTCCCTCCCCCCATTAATAATAATTCTATCCCAAA
ATCTACACAATGTTCTGTGTACACTTCTTATGTTTTTTACTTCTGATAAATTTTTTTGAAACAT
CATAGAAAAAACCGCACACAAAATACCTTATCATATGTTACGTTTCAGTTTATGACCGCAAT
TTTTATTTCTTCGCACGTCTGGGCCTCTCATGACGTCAAATCATGCTCATCGTGAAAAAGTT
TTGGAGTATTTTTGGAATTTTTCAATCAAGTGAAAGTTTATGAAATTAATTTTCCTGCTTTTG
CTTTTTGGGGTTTCCCCTATTGTTTGTCAAGATTTCGAGGACGGCGTTTTTCTTGCTAAAAT
CACAAGTATTGATGAGCACGATGCAAGAAAGATCGGAAGAAGGTTTGGGTTTGAGGCTCA
GTGGAAG
178
eat-4p
ATTTCTAATAAAACGGTCTACCATTTTGAGTCTATTATAGCCGAAAATCTCCAATGTGACTGT
GACTTCTTAAAACTACTAAAACATTATTTGTCCATTTACATCTTCCTAAAACCGTATATCATC
AAAAACATTCACAAAATCCGAAAAATGAGACAAAATTTTTTTTTGATTGTTATTGCAATAAA
TCTAATAAAAATATTCATATATTGCCTGGCGCCCCCCATATCTCCATTTCCGGTCCCATCAC
CCCCACACCTCCAAGATTGATAGGTGGCTATAAGCATTTTTTGCATTTGAATGTGTTGCACC
AGTAGTCATCATCATCATTATCTAAACTGACGTGATAGTAGGGGGCTTTCTAGAAGTCGATT
TTCTATTAATGTCAACTTCATTCGTTGTCCCTTCCTTTCCCCGTCTTCCCTCACTTCCTTTTT
TCTATTTTTTCCAGTGGTCCGTAGTGGGCGGCACCCGATTTTGACTTGAAATCAGACCCGT
TTCCGGTTCTTTTGGTAGTTGTTAAGTTCTGATTCTATGACGTGGAGTGAAACAAAGAACGA
CCATATTTCATGTGTTGTGTTTTCTAGGCAGTCTAGGCAGGCAGTAGGCAGATACTGTCAA
AATTGGAATATTTCCATCTTCTAAATACCCTCAACTATTGTTAGCGCGTTTGAAACAATAAT
TGCAAAACATTTTTTTCGCATTGATTGGGGCATTTTGAAATTTAGAGTAAAATCGACTATCAA
CTGTCATTCCATAATAATTGGCAGAATTATTTTGGTTATGCCACCAAATAATCAATAAAGAAA
GATTTCTGTCCTTATTAAACTAAAAATTGAAGCAACGGTAGAGTTGCTGAACACAGTCGCCG
GCAAAATTTTTTAATTTTCTGCAAATTTGAAATTCTTTTGCGGGTTTTTAGTTATGGTTCTAAT
AGTGGTTAAAAGTCATTATAAAACACTTCAATTTTTTTGTAATGCTTTCATTCGCAGTCGTGAA
GTCGTGAAACTCAGTTTTCACCTATCCTAACTTTGGAAGTCGTCCAAAAAATTATTTTAATT
CTACAATTTTATATTTCTTTTTAAACATACGATGTGATCAATCCTACATCAATTCTGCAAATTT

```
198    CTCACATTCTTGGAAGCTTGGTAATTTATCAGACTTTGACTGAAAATTTGAAAAAAAATTCAG
199    TAATTTTTGGAGATCCCTTTAAATATTATTCTAGCATTGCCATATAGAATAATTGCAAAATTC
200    AATTGGTTTCCTAGAAAGAAAATTAGATAATCTTATGAAAGAGAACCTAACCACAACAGGTG
201    TTAAATATTGATTTAAACACTAAAAAAAGTCTCTCCTTCTACCTCTCTTCTCATAGCTTTCAT
202    GTCTTGAAGCTTTTCGGTTAATTCGAGCGGAAGACTTATCAAGGTACGTCATTTTCAATTAC
203    TTGTATACATCTCTCAGCTTCTTCTAATTTTCCATGTACATCTTAGATATTCTTTTTTCAGCGA
204    CTTGATATTAGGAAGTTTTGTGTTTTCAATTTTAAAGTTGGATTAATATAGAATACCAGTCTT
205    TAAACACAAACCAACAAGGGTTCAATATCAAAATAGAAACCGAAAAAAAATATCGAAAACTG
206    AAAAAATCAAATACATATCTAAAGCAAGCTATTCGAGAAATATTTATGCATTATAACAACTAT
207    GCAGCGCTTATATCTTTATTTTTCAACAAGTGTTCCAGCAACGAGAGTCTCTTCACCAAAAA
208    GCCATCTATCAAAAACCAGGCAGTGAGTCCTAGAACCAAGCTTGTCAGAAGACAAGTGCAT
209    GTATAAATAAGTAGTAAAAGACGGGTGGGACCCAGCGCGGTGAGTAGTAGATCATAAAAAA
210    GTGACTGAAAGAAGGGGCGTTTCCTTTTCTTTATTACCTCTTCCTCTACTTCTTCTCATCTC
211    AACTTGCTTTTTCCTTCCTCTTACAACAACTCCATCATCATCATCATCATTTTCAGAAACC
212
213    tbh-1p
214    GAAATGTGTGTAGTGTCATGCTCAAAATAAAATCTGGCTATAGAGGTTACAAAAAGTATGGT
215    TGCAATGATGTTACATAAATTGTGCCCACATGAAATTATGCAAGATAATTGGTAATGGGGTT
216    GCTCATGTGATGACGGAAAATGCATAAATATATTTTGTTCACTTTTCCCCTATTCATGTGTGT
217    AGAAGAATGTAAAAATACGCATGCATGTTTTGCATTTTTCAACATTTTTCGGCATTTTTGTAC
218    ATTTTTCTACATCAGCCGACAGCCCACTTCACATGCTGAAGGCGAGAAACGCAGTGAGATG
219    TGCCCCGCAATGTCGAAAAATGCCGAAAAATGCAGAACTGCATGCGTTTTTTTACATTT
220    TACTACACACATGGCTAGGGGAAAAAAAAATGTAAAAATACATTTTTGCAATTTTCGTAATAA
221    CATAAATAACCCCATAAATGTCTTTTTTATAACTAGAATTCTAGAATTTCCTTAAAGCCACAA
222    AGATATTCCTCTGAAATTTTATTTTTATCTGACTGTGACTTATCTTGTTTCTATTATGTGCATT
223    TTTTGTCGTTAGCTTTGTCGGCTCCAAATTAAATTTCATAAATTATCAATAAATTATCATTGCA
224    CAAATATGTAACAGTATGCAATATTCACATTACAATTTTATTTTGATTTCTCTACTGGTAGAG
225    TTGATGGTTGCGTTGGTGCAGCTGCAGTAGTTCTTACATATTTTGAAGAGCGAAGAGCCTC
226    GGCGGATAAAAGTAGTGTCCATTTTCTCGTCGGATACTCCGGAGCAGCGCAGTTTGATGT
227    ATCAGCTTCTCTGCTGGACATGTTTGAATCTAAATTTTCAGAACTTTCAAAAGAACTAACAAT
228    TTTCATATCCCACCAGCTTTTGACGTTTTCCAATCTCGAGCTCTTGAAACAAATTCCTATATG
229    TTGTTTGCGTGTCAAGCGGCACATTTGATCTGAAAAATGAAATTCTTGATAAGAATTCAATT
230    ATCATGGAATGCTTCATTACACTGGCAAAGCGGATGCTTTTCTCGAAAGGCGTGTTCTGGC
231    GACGGATGCATCAAATGGTTGATATTGGCTACGGTGGGTGGTTTCAGAATAAAAGGGGATT
232    TTCGGAGCATCTCCTGTTTCAGATTCAGTTATTTCTCTGAAAATAAAAACTTATCAACTTTTT
233    CATTTCATGTGTAAGATTTGGTGGTTATCAGGATCAATACTATAAATCTTAAAAAAAGTAAA
234    TTTATTTTTATTTTTTAAATATTTCCGAAAAATAGTTTTTTGTTGAGTGGTTTTTCACAACCGC
235    GATGTCTCTTCATTTTATTACACAAGGTATCACTTCTTTCCTTTCTTGTGTTTTTATGACTAG
236    CATATTCTTTTTTTGAAAGATGTGCAGTTGACAGTTTTCAAATACTCCCTGTCATTGTCACTT
237    GTCATTTTCATACCTCTTCTCCCTTGCGTGCGGCACTTTCCACGTATAAGCGGCGTTGTTTG
238    TGGTGCGCCCGTAGA
239
240    unc-25p
241    TCCTACCTTTTCTATCCCACTTTTTCCCCGGAATTTTGAGATTTGACATGATTTTTCAGAGAA
242    TTTCGAGTTTTGAGAAAAAAAAATCAAAAAGCGATTTTTTGGTCAAAGCCGAAATTTAAA
243    GCTAGTTTTTTTGGAAATATCGATTTTTTTTTTGAGAATTTTTCGATTTTCCAAAAAAAAAATC
244    GAAAGTTGTTTCCCGAAAAGCCGAAATTTTTTTTTGAGATATCGAAAAATCGGAGAATTTTTC
245    GATGTTTCCCAGAAATACTAAAATTTTGAATTTTCCGGAAAAAAAATCGTTAAAAAATTAGTT
246    TCTTTTAAAGACCGAAATTTAAAGTTTTTTTTTTTGGAAATATCGATTTTTTCCGAGATATCTAA
247    AATTGAAAGAAATTTTCGATTTTCTGAGAAAAATACGAAAAAATCGAAAAAAAAACCCAAATT
248    TCGGTTTTTTCGTAAAAAAAGTCGTAAAAATGTAATTTTTTTCCTGAAAAATCGGAATTTTTTA
```

CAAAATATCGGAAAATACTCAAAAAAAAGCTGAAAATTTCGATTTTCCGGAGAAAAATCTTT
TTAAAAAAATATTTTTTTTTCAGAAAATAAGAAAAGCCGAAATTTATAACTATTTCTCCGGAA
ATTCGAAATTTTTAACGAAATATCGGAATAATTTTTAGATTTTTCAAGTTTGACTTTGCGAGA
AAAAAATCGGAAAAACCTCGATTCGACGCCGAAAAATGCTCCTTTTCGAAAAGATTTTTGAA
ATTTCAGAAAATCGACATGCAAGCGCGCTCCACGGCGAAATGACAACGATGATCCACCGC
CCTCAAAAAGTTGGGTCTCGTTAGGTATTTGGCGGTAAAACTGGTAAAACTCCAGTTTTGC
CTCCAACGAGACCCAATTTTTTGGGGCGGTGGTGGAGCGCGCTTGCACAAGCTGAAAGCA
TTTTTCTGCGACTCGATAATATTTTGAAAACCTGTGTCAATTCTCGAAATTCTTTTTTTAAAAA
ATAATCCCGAGCTTCTCTCAGTCCTCCTCTATGAGGATGTTCCTTTTTTTTGGTTTTTCAATT
TTTTTTAAAATTCCAAATTTCTGTTGTGCAATTCACTTCCCCCCAAGAAATCCCAAAAATCCC
CAGTTTTCCCCAAAAATGTTCCGTTTTCATGTGATTTTTCCCCCATTTTTAAAACATTTTTTTG
ACTTTTTTTTAAAATGATTATTATTATTGTTTTCTATTTCATGGCCGGTAAATTATTTTTTTTCT
TTCTTTTTTTTTGCTCTTTTTTTTTCAAGAATTTTCGAATTGTTTGAAGGGCTGCTCATCTAATC
TTTTGTCATTTTGTTCTGATGCCATCATTTCTGAGAGGACCTTTGAAGACTCGTCACGAAAC
GGGAGGGGGGCTCAAGTGAGCATTATTATTATTATTGTCGCAAAAAGTTTACCCCGGG
CTCCCCCTGGCTCCCCTCTTTGAGCAAGGGTTTAAGGGCTCATTTTGATGACGAATTGCTC
ATTGGGATTATAGTCACGCCCCTCTTTTGGAGCAACTACACAACTGAGCCACAGTAATCCT
TGGGGGCGGGGTCAGTAGGACCCCCTCCGGAATAGGGAAAGCTCAGTTCACCGCCAAA
A

Neuron Count

Worm Atlas (https://www.wormatlas.org/neurotransmitterstable.htm) summarizes the proposed neuron location for glutamatergic, octopaminergic, and GABAergic neurons in hermaphrodites.

Neuron Type	Proposed Number	Location by Body Region
Glutamatergic	79	Head: ADA, ADL, AFD, AIB, AIM, AIZ, ASE, ASG, ASH, ASK, AQR, AUA, AWC, BAG, FLP, IL1, OLL, OLQ, RIA, RIG, RIM, URY Pharynx: M3, M1, I2, I5 Ventral nerve cord and body: ALM, AVM, CP0, CP5, CP6, CP7 Tail: DVC, LUA, PHA, PHB, PHC, PLM, PVD, PVQ, PQR, PVR
Octopaminergic	2	Head: RIC
GABAergic	32	Head: AVI, DD1-6, DVB, RIB, RIS, RME, SMDD/V, VD1-13 Ventral nerve cord and body: DD2-4, VD3-11 Tail: DD6, DVB, VD12-13

C. elegans microscopy

Stereoscope

For whole-worm imaging of *vha-6p::Q40::YFP*, DHS-3::GFP, and VIT-2::GFP strains, synchronized animals were grown on RNAi plates seeded with EV bacteria or RNAi bacteria. Animals were imaged at day 1 of adulthood and for aging experiments also imaged at day 5 and

day 9 of adulthood. 10+ animals were placed in a pool of 100 mM sodium azide in M9 on
standard NGM plates without bacteria to induce paralysis. Paralyzed animals were then lined
alongside each other and imaged on a Leica M205FCA automated fluorescent
stereomicroscope running LAS X software and equipped with a standard GFP filter, Leica LED3
light source, and Leica K5 camera. For all imaging experiments, 3 biological replicates were
performed with 2 technical replicates each, and 1 representative image was chosen for use in
figures. For *vha-6p::Q40::YFP* quantification, Fiji[14] was used to draw a region of interest along
the posterior half of each group of worms, and integrated density was measured. Graphing and
statistical analysis were performed with GraphPad Prism 10 software using a Mann-Whitney
test unless otherwise stated. *vha-6p::Q44::YFP* quantification was performed by counting the
number of protein aggregates in individual worms and statistical analysis was performed with
GraphPad Prism 10 software using a Mann-Whitney test.
291

Widefield and confocal imaging

Widefield imaging utilized a Leica THUNDER Imager equipped with a 63x/1.4 Plan
AproChromat objective, standard dsRed filter (11525309), Leica DFC9000 GT camera, a Leica
LED5 light source, and run on LAS X software. For high-resolution imaging of DHS-3::GFP and
ER morphology, imaging was performed using a Leica Stellaris 5 confocal microscope equipped
with a white light laser source and spectral filters, HyD detectors, 63x/1.4 Plan ApoChromat
objective, and run on LAS X software. Animals were placed in 100 mM sodium azide solution on
a glass slide to induce paralysis and imaged within 5 minutes of slide preparation to prevent
artifacts from prolonged exposure to sodium azide. For imaging experiments, 3 biological
replicates were performed with 1 or 2 technical replicates each, and 1 representative image was
chosen for use in figures.
303

Intestinal bacteria invasion assay

Assessing intestinal bacteria invasion was performed as previously described[15]. Animals were
L1 synchronized via bleaching and plated on RNAi plates containing a bacterial lawn derived
from a mixture of 80% HT115 bacteria containing EV and 20% HT115 bacteria expressing
mCherry. Once at the desired age, animals were manually transferred onto a standard OP50
plate and allowed to feed on OP50 for 2 hours at 20°C to facilitate clearance of mCherry
bacteria. For imaging, worms were paralyzed by exposure to M9 solution containing 100 mM
sodium azide and arranged on a standard NGM plate without bacteria. Images were captured
using a Leica M205FCA fluorescent stereomicroscope equipped with a standard dsRed filter as
described above. For each of the 3 biological replicates, 2 technical replicates with 13 animals
per replicate were performed and 1 representative image was used for figures. The percentage
of animals exhibiting bacterial invasion was quantified and plotted with GraphPad Prism 10
software for each technical and biological replicate. Statistical analysis was conducted across all
replicates using a Mann-Whitney test with GraphPad Prism 10 software.
318

ER Secretion Assay

Assaying of ER secretory function was performed as described previously[16]. Transgenic control
animals and *eat-4p::xbp-1s*, *tbh-1p::xbp-1s*, and *unc-25p::xbp-1s* animals expressing VIT-
2::GFP were bleached to obtain eggs. Eggs were then placed on glass slides and imaged using
a Leica THUNDER Imager equipped with a 63x/1.4 Plan AproChromat objective, standard GFP
filter (11525309), Leica DFC9000 GT camera, a Leica LED5 light source, and run on LAS X
software. Images were quantified using Fiji and drawing a region of interest around each
individual egg to obtain an integrated density value. 4 independent biological replicates were
performed. SuperPlots[17] were created using GraphPad Prism 10 software where large dots
represent the median value of each biological replicate and small dots represent single eggs

with different intensities of colors representing eggs from the same biological replicate; lines
indicate median and interquartile range. All statistical analyses were conducted using Mann–
Whitney testing with GraphPad Prism 10 software. For whole worm imaging, animals were
raised to day 3 of adulthood and imaged using the Leica M205FCA fluorescent
stereomicroscope equipped with a standard GFP filter as described above.
334

Oil Red O Staining

Oil Red O fat staining was performed as previously described[18]. Briefly, worms were bleached,
and eggs were plated to obtain a synchronous population. Worms were grown on RNAi plates
with a lawn of EV bacteria and aged to day 3 of adulthood. Aging was performed in the absence
of FUDR by gravity settling in M9 solution and aspirating to remove progeny. For staining,
worms were washed off plates using a PBS + 0.01% Triton solution, rocked for 3 minutes in
40% isopropyl alcohol, pelleted, and then stained with Oil Red O in diH$_2$O for 2 hours while
rocking at room temperature. Worms were pelleted and washed in PBS + 0.01% Triton for 30
min before being imaged at 20× magnification with a Leica THUNDER Imager Flexacam C3
color camera and run on LAS X software. To quantify somatic fat depletion, worms were scored
as previously described[19]. The level and distribution of fat was placed into categories of non-
somatic lipid depletion, displaying no loss of fat and being darkly stained throughout the body,
and somatic lipid depletion, being stained largely in the germ cells. At least 100 worms were
scored for each condition over 3 biological replicates.
349

C. elegans RT-qPCR and RNA-seq analysis

For collection of RNA, we used *glp-4(bn2)* animals to eliminate progeny. After bleaching and L1
arresting, all animals were raised at 22°C (the restrictive temperature for our backcrossed *glp-*
4(bn2) strain) for 3 days to collect animals at day 1 of adulthood. Approximately 1000 animals
were used per condition. Worms were collected using M9 and transferred to TRIzol solution and
underwent 3 freeze/thaw cycles between liquid nitrogen and a 37°C bead bath with a 30-second
vortexing step between each cycle to lyse worms. Following the final thaw, chloroform was
added at a ratio of 1:5 chloroform/TRIzol, and aqueous separation of RNA was achieved by
centrifugation using a heavy gel phase-lock tube (VWR, 10847–802). The aqueous phase was
mixed with isopropanol at a 1:1 ratio and applied to a QuantaBio Extracta Plus RNA kit (95214)
for RNA purification according to the manufacturer's instructions.
361

Library preparation and sequencing was conducted at Novogene using their standard pipeline
using paired-end, polyA selection, first-strand synthesis, and an Illumina NovaSeq6000. Each
condition was measured with 3 biological replicates. Gene expression levels were quantified
using STAR-2.7.3a[20] with WBcel235 as the reference genome. Fold changes were determined
using DESeq2[21]. Gene targets of XBP-1 were defined based on previous experimental
findings[22]. Gene Ontology (GO) enrichment analysis was performed using WormEnrichr[23,24].
rgef-1p, *dat-1p*, *tph-1p*, *eat-4p*, *tbh-1p*, and *unc-25p* driven expression of *xbp-1s* was compared
to the N2 wild-type control.
370

371 For RT-qPCR, cDNA synthesis was conducted using qScript cDNA SuperMix (QuantaBio,
372 101414-102) with 500 ng of RNA. RT-qPCR was performed using NEB Q5 DNA polymerase
373 following the manufacturer's guidelines and utilizing the primers listed below. Each condition
374 was assessed using 3 biological replicates. QuantStudio 3 (Thermo Fisher) was used for
375 quantification using a standard curve method.
376

Lifespan measurements

For lifespan experiments, animals were grown on RNAi plates on either EV or
RNAi bacteria from L1 stage. At day 1 of adulthood, animals were washed off plates with M9
solution and then moved to plates containing 100 µL of 100 mg/mL FUDR to eliminate progeny.
One replicate of lifespan assays was performed using a lifespan machine[25] with others being
done by hand. Tunicamycin survival assays were performed based on established protocols[26].
For tunicamycin assays, animals were moved onto plates supplemented with 25 µg/mL
tunicamycin in DMSO directly in the plate. Animals were grown at 20°C and checked every 2
days for viability. Animals were considered dead if they did not exhibit any movement when
prodded with a platinum wire at both the head and the tail. Animals that exhibited bagging,
intestinal leakage, desiccation on the side of the plate, or other deaths unrelated to aging were
scored as censored. All lifespans were performed on >3 biological replicates. Lifespan assay
survival curves were plotted using GraphPad Prism 10 software and statistics were performed
using a Log-Rank test in GraphPad Prism 10. Representative data are depicted in figures and a
table of all lifespan assays performed is available in **Table S7**.
392

Caenorhabditis elegans brood size assay

Brood assays were measured as previously described[26]. Bleaching was used to obtain a
synchronized population of animals, and 10 L4 stage animals were transferred onto individual
plates. Every 12 hours, the animals were moved onto new plates, while plates containing eggs
were stored in a 15°C incubator for 2–3 days. All surviving progeny on each egg-laying plate
were counted and totaled to determine the brood size. SuperPlots[17] were created using
GraphPad Prism 10 software where large dots represent the median value of each biological
replicate and small dots represent single animals with different intensities of colors representing
animals from the same biological replicate; lines indicate median and interquartile range. All
statistical analyses were conducted using Mann–Whitney testing with GraphPad Prism 10
software.
404

Caenorhabditis elegans thrashing assay

Thrashing assays were conducted on animals synchronized via bleaching and aged on RNAi
plates containing FUDR from day 1 of adulthood. Upon reaching the desired age, plates
containing adult animals were flooded with 100 µL of M9 solution, and 30-second videos were
recorded using an M205FCA stereomicroscope equipped with a Leica K5 microscope running
LAS X software. Thrashing movements were manually recorded over a 10-second period. A
bending of more than 50% of the animal's body in the opposite direction was deemed a single
thrash. Representative data from 3 independent biological replicates are presented.
SuperPlots[17] were created using GraphPad Prism 10 software where large dots represent the
median value of each biological replicate and small dots represent single animals with different
intensities of colors representing animals from the same biological replicate; lines indicate
median and interquartile range. All statistical analyses were conducted using Mann–Whitney
testing with GraphPad Prism 10 software.
418

Fast kill assay

Fast kill assays were performed as previously described with minor modifications[27,28].
Pseudomonas aeruginosa (PA14) cultures were grown overnight at 37°C for 14-15 hours. 5 µL
of overnight culture was spread over 3.5 cm peptone glucose media plates (1% Bacto-Peptone,
1% NaCl, 1% glucose, 1.7% Bacto Agar) containing 0.15 M sorbitol, using a spreader made
from an open loop tipped glass pasture pipette. The plates were incubated at 37°C for 24 hours
and then at 25°C for 48 hours. Following this, 30 to 40 synchronized L4 animals were placed on
each plate. Assays were performed at 25°C. Survival of animals was plotted over a period of 8

hours with intervals of 2 hours. An animal was deemed dead when it no longer responded to
touch. 3 biological replicates with 3 technical replicates were performed for a total of 9
replicates. Survival rates were measured wherein 100% survival was indicated as an integer "1"
and the fraction of survival populations at every time point was represented as decimal values.
431

Forced food choice

Forced food choice assays were performed as described previously[29,30]. Plates were made using
the same recipe for NGM plates aside from the addition of 0.35% peptone. A single colony of
Pseudomonas aeruginosa (PA14) and *E. coli* OP50 bacteria were inoculated into separate 3 mL
of LB for overnight primary culture at 37°C. The following day, the OD_{600} of each of the cultures
was diluted to an OD_{600} of 1.0. A culture of PA14 was transferred as a line along the center of an
NGM plate using a glass Pasteur pipette bent at a 90-degree angle. Next, 15 μL of OP50 culture
was seeded as a dot onto the plate 2.5 cm away from the center and 0.5 cm away from the edge
of the plate. The plates were dried and transferred to 37°C for 24 hours, followed by incubation at
25°C for 48 hours. On the day of the assay, the plates were removed from the 25°C incubator
and allowed to reach room temperature. Worms were washed three times in M9 solution before
being placed onto the assay plate diametrically opposite to the OP50 dot at 2.5 cm away from the
center. The proportion of worms found on or off each food was recorded after 1 hour, 2 hours, 4
hours, 6 hours, and 8 hours. After each time point, the population was scored in which -1
represented 100% of the population on PA14 and +1 represented 100% of the population on
OP50. The movement index was then calculated towards the OP50 dot using the formula:

448
449 **("A" population of worm on OP50 dot – "B" population of worms on PA14 line)**
450 **("A" population of worm on OP50 dot + "B" population of worms on PA14 line)**
451 Each assay was done in biological triplicate with technical triplicates for a total of 9 replicates.
452 Statistical analyses were conducted using Mann–Whitney testing with GraphPad Prism 10
453 software.

Statistics

All statistical analyses were performed using GraphPad Prism 10 software. No assumptions
were made about data distribution. Mann-Whitney testing was used for most comparative
analyses with p-values less than 0.05 considered significant unless otherwise specified. To
determine statistical significance of lifespan data, a log-rank test was performed with p-values
less than 0.05 considered significant. At least 3 biological replicates were performed for each
experiment unless otherwise noted.

Figure captions

464
Fig. 1. Dopaminergic and serotonergic *xbp-1s* together do not recapitulate pan-neuronal
xbp-1s overexpression. A comparison of differentially expressed genes (p-value ≤ 0.01)
between worms expressing *xbp-1s* pan-neuronally (*rab-3p*), and in either **(A)** dopaminergic (*dat-*
1p) neurons, **(B)** serotonergic (*tph-1p*) neurons, or **(C)** concurrently in both dopaminergic and
serotonergic (*dat-1/tph-1p*) neurons. For a complete list of differentially expressed genes see
Table S3. For a complete list of genes represented in Venn Diagrams, see **Table S4**. **(D)** Heat
map of differentially expressed genes in worms expressing *xbp-1s* pan-neuronally with
corresponding expression levels in serotonin, dopamine, and both serotonin and dopaminergic
xbp-1s expressing animals. Warmer colors indicate increased expression, and cooler colors
indicate decreased expression. See **Table S5** for a list of gene names and expression values.

(E) Schematic of each neuronal type explored in this study: glutamatergic, *eat-4p* (n = 79,
location: head, pharynx, ventral nerve cord and body, tail), octopaminergic, *tbh-1p* (n = 2,
location: head), GABAergic, *unc-25p* (n = 32, location: head, ventral nerve cord and body, tail).
(F) qPCR of transcripts of glutamatergic *xbp-1s* (green), octopaminergic *xbp-1s* (yellow), and
GABAergic *xbp-1s* (pink) animals grown on empty vector from hatch. RNA was isolated in day 1
adults and data are compared using a standard curve with data represented as relative fold
change against control. Each dot represents a biological replicate averaged across three
technical replicates per sample. Lines represent geometric mean with geometric standard
deviations.
484
Fig. 2. Glutamatergic, octopaminergic, and GABAergic *xbp-1s* modulate distinct
transcriptional pathways. Volcano plots of whole-body genome-wide changes in gene
expression upon *xbp-1s* overexpression in **(A)** glutamatergic, **(B)** octopaminergic, and **(C)**
GABAergic neurons. Red dots indicate significantly differentially expressed genes with p-value ≤
0.01. See **Table S3**. **(D)** Comparison of differentially expressed genes (p-value ≤ 0.01) between
worms expressing *xbp-1s* in glutamatergic, octopaminergic, and GABAergic neurons. For a
complete list of differentially expressed genes see **Table S4**. **(E)** Heat map of XBP-1s target
gene [31] expression under neuronal, glutamatergic, octopaminergic, and GABAergic *xbp-1s*.
Warmer colors indicate increased expression, and cooler colors indicate decreased expression.
See **Table S5**. Top ten most enriched gene ontology terms of differentially expressed genes
upon *xbp-1s* overexpression in **(F)** glutamatergic, **(G)** octopaminergic, and **(H)** GABAergic
neurons. See **Table S6**.
497
Fig. 3. Octopaminergic *xbp-1s*, but not glutamatergic or GABAergic *xbp-1s*, is sufficient
to extend lifespan. (A) Lifespan measurements of control (blue) and a mixed population of
both "normal" and "stunted" growth octopaminergic *xbp-1s* (yellow, *tbh-1p*, mixed) animals. **(B)**
Lifespan measurements of control (blue) and octopaminergic *xbp-1s* animals. Octopaminergic
xbp-1s animals were separated into normal size (yellow, *tbh-1p*) and stunted growth (purple,
tbh-1p, small). **(C)** Lifespan measurements of control (blue, light blue) and octopaminergic *xbp-*
1s animals (*tbh-1*, yellow, orange) grown on either EV or *xbp-1* RNAi. **(D)** Lifespan
measurements of control (blue, light blue), glutamatergic *xbp-1s* animals (*eat-4p*, green), and
GABAergic *xbp-1s* (*unc-25p,* pink) animals. Lifespans were scored every 2 days and data is
representative of 3 biological replicates (N). Sample size (n) is written next to each condition
followed by significance measured using Log-Rank testing: n.s. = not significant, * = p < 0.05,
*** = p < 0.001. All statistical analysis is available in **Table S7**.
510
Fig. 4. Glutamatergic, octopaminergic, and GABAergic *xbp-1s* enhance pathogen
resistance and increases pathogen apathy. (A) Heat map of immune response
(GO:0006955) gene expression under pan-neuronal (*rgef-1p*), glutamatergic (*eat-4p*),
octopaminergic (*tbh-1p*), and GABAergic (*unc-25p*) *xbp-1s* overexpression. Warmer colors
indicate increased expression, and cooler colors indicate decreased expression. See **Table S6**.
(B) Survival analysis of control (N2, blue), glutamatergic *xbp-1s* (green, *eat-4p*), octopaminergic
xbp-1s (yellow, *tbh-1p*), or GABAergic *xpb-1s* (pink, *unc-25p*) on PA14 fast kill assay plates for
2, 4, 6, and 8 hours. Each fast kill assay is comprised of 3 technical replicates per biological
replicate and at least 3 biological replicates per condition. Results were analyzed via two-way
ANOVA test; **(p<0.01) ***(p<0.001) ****(p<0.0001). **(C)** Pathogen avoidance behavior of
control (N2, blue), glutamatergic *xbp-1s* (green, *eat-4p*), octopaminergic *xbp-1s* (yellow, *tbh-1p*),
or GABAergic *xpb-1s* (pink, *unc-25p*) during "forced" food choice assays measured at 1, 2, 3, 6,
and 8 hour time points. Each forced food choice assay is comprised of 3 technical replicates per
biological replicate and at least 3 biological replicates per condition. Results were analyzed via
two-way ANOVA test; **(p<0.01) ***(p<0.001) ****(p<0.0001). **(D)** Representative brightfield and

fluorescent images of adult worms grown on bacteria expressing mCherry. Animals are moved to OP50 plates for two hours to remove mCherry expressing bacteria from the intestine before imaging. Any remaining mCherry signal after OP50 clarification are signs of bacterial colonization. **(E)** Quantification of the percent of animals displaying intestinal bacterial colonization was performed across 2 technical replicates for each of 3 biological replicates for a total of 6 replicates. Lines represent mean and standard deviation. * = p ≤ 0.05, ** = p ≤ 0.01, ns = p > 0.05 using a Mann-Whitney test.

Fig. 5. Glutamatergic, octopaminergic, and GABAergic *xbp-1s* results in depletion of lipids. (A) Heat map of lipid homeostasis (GO:0055088) gene expression under pan-neuronal (*rgef-1p*), glutamatergic (*eat-4p*), octopaminergic (*tbh-1p*), and GABAergic (*unc-25p*) *xbp-1s* overexpression. Warmer colors indicate increased expression, and cooler colors indicate decreased expression. See **Table S6**. **(B)** Representative fluorescent micrographs of day 3 adult animals of control, glutamatergic *xbp-1s* (*eat-4p*), octopaminergic *xbp-1s* (*tbh-1p*), and GABAergic *xpb-1s* (*unc-25p*) taken on a stereomicroscope and on a confocal microscope (bottom)**.** All images are contrast matched. Scale bar represents 10 um. **(C)** Representative images of day 3 adult animals of control, glutamatergic *xbp-1s* (*eat-4p*), octopaminergic *xbp-1s* (*tbh-1p*), and GABAergic *xpb-1s* (*unc-25p*) of ORO-stained lipids. Quantification of lipid staining as non-lipid depletion (black) and lipid depletion (gray). **(D)** Representative fluorescent micrographs of day 3 adult mRuby::HDEL of control, glutamatergic *xbp-1s* (*eat-4p*), and GABAergic *xbp-1s* (*unc-25p*), animals (top) or day 3 adult mCherry::HDEL of control and octopaminergic *xbp-1s* (*tbh-1p*), animals (bottom). Images are representative of three independent biological replicates and are independently contrast enhanced for each individual image. Scale bar represents 10 µm.

Fig. 6. Glutamatergic, octopaminergic, and GABAergic *xbp-1s* enhance proteostasis. (A) Representative images of protein aggregation in animals expressing intestinal polyglutamine repeats (*vha-6p::polyQ40::YFP*) [32] in glutamatergic (*eat-4p*), octopaminergic (*tbh-1p*), or GABAergic (*unc-25p*) *xbp-1s* animals. All animals were imaged on day 1, 5, and 9 of adulthood. Images were captured using a Leica M205 stereo microscope. **(B)** Quantification of fluorescence integrated density normalized to area was performed across 2 technical replicates each of 3 biological replicates for a total of 6 replicates. Lines represent mean and standard deviation. * = p ≤ 0.05, ** = p ≤ 0.01, ns = p > 0.05 using a Mann-Whitney test. **(C)** Representative images of a second distinct integration line of glutamatergic, octopaminergic, or GABAergic *xbp-1s* animals expressing intestinal polyglutamine repeats (*vha-6p::polyQ40::YFP*) grown on EV or *xbp-1s* RNAi and imaged as per (A). **(D)** Quantification of fluorescence integrated density normalized to area was performed across 3 biological replicates. A Shapiro-Wilk test was used to determine normality and a student's t-test was used to assess significance.

Results and discussion

Overexpression of xbp-1s in glutamatergic, octopaminergic, and GABAergic neurons.
In previous studies, serotonergic, dopaminergic, and RIM/RIC neurons have been identified to be involved in neuron-to-body communication of UPR[ER][3,4]. However, these four neuron subtypes make up only ~18 of the 302 neurons in *C. elegans*, raising the question of what other neuronal subtypes may be involved in neuron-to-body UPR[ER] communication. Indeed, transcriptomic analysis of previously published datasets[3,8] reveals only a minor overlap of differentially expressed genes between worms expressing *xbp-1s* in dopaminergic and serotonergic neurons as compared to pan-neuronal expression (**Fig. 1A-C**). In addition, the

576 expression levels of differentially expressed genes under pan-neuronal *xbp-1s* expression are
577 largely dissimilar when *xbp-1s* is expressed in serotoninergic or dopaminergic neurons, or both
578 concurrently (**Fig. 1D**). This strongly suggests that other neuronal subtypes are involved in non-
579 autonomous UPR^{ER} signaling.
580
Previously, we performed a screen of several neurotransmitter signaling pathways involved in
neuronal communication of UPR^{ER}, which revealed glutamatergic, octopaminergic, and
GABAergic neurons as candidates involved in this signaling event[3]. Glutamate is a widely
utilized, excitatory neurotransmitter in both invertebrate and vertebrate systems[33]; octopamine is
a *C. elegans*-specific neurotransmitter similar to the mammalian norepinephrine, and is involved
in immune response[34]; and gamma-aminobutyric acid (GABA) is a widely utilized
neurotransmitter that has been found to function as both an excitatory and an inhibitory signal in
C. elegans[35]. *C. elegans* possess 79 glutamatergic neurons, 2 octopaminergic neurons, and 32
GABAergic neurons in hermaphrodites (**Fig. 1E**) (Loer CM, Worm Atlas,
https://www.wormatlas.org/neurotransmittercriteria.htm).
591
To determine the potential involvement of glutamatergic, octopaminergic, and GABAergic
neurons in neuron-to-body communication of UPR^{ER}, we overexpressed *xbp-1s* in each
neuronal subtype using the *eat-4* promoter for *xbp-1s* overexpression in glutamatergic
neurons[36] (hereafter referred to as glutamatergic *xbp-1s*); *tbh-1* promoter for *xbp-1s*
overexpression in octopaminergic neurons[37] (hereafter referred to as octopaminergic *xbp-1s*);
and the *unc-25* promoter for *xbp-1s* overexpression in GABAergic neurons[38] (hereafter referred
to as GABAergic *xbp-1s*). We confirmed by quantitative PCR (qPCR) that all three subtypes
display an increase in *xbp-1s* mRNA, although our data did not reach statistical significance
(**Fig. 1F**).
601
Glutamatergic, octopaminergic, and GABAergic xbp-1s alter distinct transcriptional
pathways.
To more thoroughly investigate the impact of neuronal subtype UPR^{ER} on the periphery, we
performed whole-worm RNA sequencing on animals overexpressing *xbp-1s* in glutamatergic,
octopaminergic, and GABAergic neurons. Glutamatergic and octopaminergic *xbp-1s* resulted in
sizable changes to gene expression, while more mild changes occurred with GABAergic *xbp-1s*
(**Fig. 2A-C**). Interestingly, the majority of differentially expressed genes were unique to each
condition, suggesting distinct responses were induced by each neuronal subtype (**Fig. 2D**). This
adds more insight into a previous study that identified distinct pathways activated downstream
of serotonergic and dopaminergic *xbp-1s* [3].
612
To further characterize the similarities and differences between peripheral response to neuronal
subtype UPR^{ER}, we directly compared our glutamatergic, octopaminergic, and GABAergic *xbp-*
1s animals to previously published RNA-seq datasets [3,8]. First, we sought to determine the
overlap between neuronal subtype *xbp-1s* overexpression with pan-neuronal *xbp-1s*
overexpression (hereafter referred to as neuronal *xbp-1s*), as we would expect that neuronal
xbp-1s includes each neuronal subtype. We compared neuronal *xbp-1s* using two different
promoters, *rab-3p* and *rgef-1p* and were surprised to find that while there was significant overlap
between these two neuronal *xbp-1s* strains, a majority of differentially expressed genes were
not shared (**Fig. S1**). Since this could potentially be due to leakiness of the *rab-3p* compared to
the *rgef-1p*[39,40], in our subsequent studies, we focused on making comparisons to results from
the *rgef-1p::xbp-1s* strain (which we will continue to refer to as neuronal *xbp-1s*).
624
625

As expected, neuronal *xbp-1s* animals display altered expression of a large number of direct
XBP-1s targets[31]. Interestingly, we see that glutamatergic *xbp-1s* induces many of these same
XBP-1s targets and to an even greater extent than neuronal *xbp-1s* (**Fig. 2E**). These data
suggest that glutamatergic *xbp-1s* activates a more canonical UPR^{ER} signature involved in
conventional protein processing pathways. Gene ontology (GO) enrichment analysis supported
this idea, as the most enriched biological processes included pathways related to ER function
and protein homoeostasis, including ER to Golgi vesicle-mediated transport, protein N-linked
glycosylation, endoplasmic-reticulum-associated protein degradation (ERAD) pathway, and
proteolysis (**Fig. 2F**). However, a majority of differentially expressed genes in glutamatergic *xbp-*
1s are still distinct from neuronal *xbp-1s* (**Fig. S2A**), suggesting that these protein homeostatic
pathways are being regulated in different ways in each condition. Interestingly, glutamatergic
xbp-1s transcriptionally regulates an entirely different set of genes than serotonergic *xbp-1s*
(**Fig. S2E**), although these animals were also shown to induce canonical protein homeostasis
pathways[3]. Altogether, these data show that even amongst neuronal subtypes that share a
similar peripheral response (e.g., protein homeostasis), the specific genes targeted in these
similar pathways are distinct, highlighting the fact that non-autonomous UPR^{ER} is dramatically
different based on which neuronal subtype is involved.
643
Octopaminergic *xbp-1s* showed smaller gene expression changes to XBP-1s targets in
comparison to glutamatergic *xbp-1s*, being more reminiscent of the levels found in neuronal
xbp-1s (**Fig. 2E**). However, similar to glutamatergic *xbp-1s*, when all differentially expressed
genes for octopaminergic *xbp-1s* were compared to neuronal *xbp-1s*, the majority of
differentially expressed genes were distinct (**Fig. S2B**). The differentially expressed genes
identified were entirely different from those found in serotonergic and dopaminergic *xbp-1s* (**Fig.**
S2D-E). GO analysis identified that the most dramatic changes in gene expression in
octopaminergic *xbp-1s* were defense response pathways, particularly those involved in immune
response (**Fig. 2G**). These data are consistent with previous findings that showed pathogen
response in *C. elegans* is associated with UPR^{ER} induction[41] and a role for non-autonomous
signaling in this response[11], potentially through octopaminergic signaling[34]. These data add an
additional downstream function of non-autonomous UPR^{ER} in regulation of immune response,
potentially downstream of octopaminergic neurons.
657
Finally, GABAergic *xbp-1s* activation caused minimal changes in gene expression overall, with
very little overlap with other neuronal subtype *xbp-1s* (**Fig 2E, S2C-E**). Although the gene
expression changes were minor, GO analysis did reveal some pathways previously associated
with UPR^{ER} induction, including lipid remodeling[3,5] (**Fig. 2H**). When looking at all genes related
to canonical UPR^{ER} (**Fig. S2F**) and XBP-1s targets (**Fig. S2G**), glutamatergic *xbp-1s* animals
displayed the most significantly differentially expressed genes, while octopaminergic *xbp-1s* had
more subtle effects and GABAergic *xbp-1s* displayed no major differences. Interestingly,
octopaminergic *xbp-1s* also had significantly differentially expressed genes for the UPR^{MT} (**Fig.**
S2H), which further adds evidence to its role in immune response as UPR^{MT} activation has been
directly linked to response to pathogens[42]. Glutamatergic *xbp-1s* animals also displayed
significantly differentially expressed genes for the heat-shock response (HSR, **Fig. S2I**) and
oxidative stress response (OxSR, **Fig. S2J**), which is consistent with previous reports that
suggested some overlap between UPR^{ER} and the HSR [43] and OxSR [44]. Finally, octopaminergic
xbp-1s animals show significantly differentially expressed genes for genes related to translation
(**Fig. S2K**), another feature often correlated with UPR^{MT} activation[45]. Altogether, our data adds
more evidence to the previously proposed model[46] that specific neuronal subtypes participate in
activation of unique downstream pathways in response to stress.
675

676 *Glutamatergic, octopaminergic, and GABAergic xbp-1s do not alter general organismal*
677 *health, and only octopaminergic xbp-1s is sufficient to extend longevity.*
678 Next, we sought to test the impact of neuronal subtype *xbp-1s* on general organismal health, as
679 previous studies have shown that neuronal *xbp-1s* results in a significant improvement in
680 longevity and animal health, with a reduction in reproductive health [2,37]. Interestingly, we found
681 that octopaminergic *xbp-1s* animals had a bimodal population of lifespan (**Fig. 3A**), which
682 correlates with animal size (**Fig. 3B**). Specifically, a proportion of octopaminergic *xbp-1s*
683 animals display a stunted growth phenotype, and these animals tend to have a mildly reduced
684 lifespan compared to control animals. In contrast, animals that display regular size display a
685 significant lifespan extension. Since the reduction in lifespan for the stunted growth animals can
686 be due to a number of pleiotropic and unrelated reasons, we opted to perform all further
687 analyses on octopaminergic *xbp-1s* animals on the long-lived, "normal" sized animals.
688 Importantly, this lifespan extension of octopaminergic *xbp-1s* animals was fully dependent on
689 *xbp-1* (**Fig. 3C**), similar to all other neuronal *xbp-1s* paradigms previously established[2,3].
690
691 Interestingly, glutamatergic or GABAergic *xbp-1s* animals did not display any increase in
692 lifespan, and glutamatergic *xbp-1s* animals actually had a mild decrease in lifespan (**Fig. 3D**). In
693 addition, while we saw a mild decrease in brood size in glutamatergic, octopaminergic, or
694 GABAergic *xbp-1s* animals, these differences were not statistically significant (**Fig. S3A-C**).
695 Finally, general organismal health was also unchanged as no change in motility was observed,
696 except a mild but statistically not significant increase in day 1 thrashing of octopaminergic *xbp-*
697 *1s* animals (**Fig. S3D-F**). Thus, only octopaminergic *xbp-1s* was sufficient to promote longevity,
698 with only minor – if any – changes in other healthspan metrics.
699
700 *Glutamatergic, octopaminergic, and GABAergic xbp-1s* **improve immune function.**
701 To further investigate a potential mechanism whereby octopaminergic *xbp-1s* animals promote
702 longevity, we next measured common features of UPR[ER] induction. Our transcriptomics analysis
703 revealed that glutamatergic, octopaminergic, and GABAergic *xbp-1s* displayed significant
704 changes in immune response related genes (**Fig. 4A**), with octopaminergic *xbp-1s* animals
705 having defense response against bacteria as one of the most significantly enriched GO terms
706 (**Fig 2G**). Therefore, we measured the impact of *xbp-1s* overexpression on innate immune
707 response using multiple methods. First, we used a standard pathogen resistance assay using
708 exposure to *Pseudomonas aeruginosa* (PA14)[47]. Using a canonical PA14 fast kill assay, we
709 found that glutamatergic, octopaminergic, and GABAergic *xbp-1s* animals all displayed a
710 significant increase in survival against PA14, with the octopaminergic *xbp-1s* displaying the
711 most significant increase in survival even after 8 hours (**Fig. 4B**). This is consistent with the
712 octopaminergic *xbp-1s* animals having the greatest change in expression of genes associated
713 with immune response and previous reports that indicate a functional role for octopamine
714 signaling in innate immunity in *C. elegans*[34]. In addition, since octopaminergic *xbp-1s* animals
715 are the only condition that display an improved lifespan, it is possible that the improvement in
716 pathogen resistance of octopaminergic *xbp-1s* animals reaches a critical level to also impact
717 longevity.
718
719 *C. elegans* also utilize their nervous system for aversive learning behavior to avoid pathogenic
720 bacteria[48]. This avoidance behavior is mediated by several neurotransmitters, including
721 serotonin[49] and octopamine[50], and certain strains with heightened stress responses have been
722 shown to lack this typical avoidance behavior[29]. Here, we used a previously validated forced
723 exposure method[29] to determine the impact of neuronal *xbp-1s* overexpression on pathogen
724 apathy. Similar to pathogen resistance, glutamatergic, octopaminergic, and GABAergic *xbp-1s*
725 animals all displayed increased apathy to pathogens, with glutamatergic animals having the
726 mildest phenotype (**Fig. 4C**). Thus, it is likely that the heightened resistance to pathogens is

directly correlated with a lack of urgency to escape these pathogens. While we did observe an
increase in expression of innate immune response genes, it is also possible that the increase in
pathogen resistance is due to an increase in gut barrier integrity, as age-associated loss of gut
barrier integrity results in infiltration of bacteria and bacterial colonization in the gut[51,52].
Interestingly, glutamatergic, octopaminergic, and GABAergic *xbp-1s* animals all showed similar
breakdown of gut barrier integrity and age-associated bacterial colonization in the gut compared
to wild-type controls (**Fig. 4D-E**). These data suggest that the pathogen resistance and apathy
of glutamatergic, octopaminergic, and GABAergic *xbp-1s* animals is likely due to a heightened
immune response, rather than a gut-barrier-related phenotype.
736
***Glutamatergic, octopaminergic, and GABAergic xbp-1s* display reduced lipid levels, but**
have minor changes to ER morphology.
Next, we sought to determine whether neuron subtype-specific *xbp-1s* overexpression altered
lipid levels[5,6]. Indeed, we found that *xbp-1s* overexpression resulted in changes to lipid related
gene expression (**Fig. 5A**). Therefore, we measured lipid levels using DHS-3::GFP an abundant
protein on the surface of *C. elegans* intestinal lipid droplets[53,54]. Interestingly, we saw a significant
decrease in lipid droplet abundance in glutamatergic, octopaminergic, or GABAergic *xbp-1s*
animals, despite no major changes in lipid droplet size or morphology (**Fig. 5B**). To further
evaluate changes in lipid content, we utilized a more comprehensive dye, Oil Red O (ORO), which
stains neutral lipids, cholesteryl esters, and lipoproteins[55]. Consistent with lipid droplet imaging,
we observed a significant decrease in lipid content in glutamatergic, octopaminergic, or
GABAergic *xbp-1s* animals using ORO (**Fig. 5C**). These data suggest that similar to other
paradigms of neuronal *xbp-1s* overexpression[3,5,6], glutamatergic, octopaminergic, or GABAergic
xbp-1s animals results in depletion of neutral lipids, likely resulting in improved lipid homeostasis.
751
Previous studies have shown that neuronal *xbp-1s* animals exhibit changes in ER morphology
associated with a general increase in secretory capacity of the ER and depletion of lipids,
potentially through an increase in lipophagy[6]. Therefore, we next sought to determine whether
changes in lipid levels found in glutamatergic, octopaminergic, and GABAergic *xbp-1s* animals
are also correlated with changes to ER morphology and secretory capacity. To measure general
changes to the ER, we first performed imaging of the ER using an mRuby::HDEL fused to an
HSP-4 signal sequence to localize the fluorophore to the ER[6]. Since we could not successfully
make homozygous octopaminergic *xbp-1s* animals with this mRuby::HDEL marker, we used an
mCherry::HDEL fused to a SEL-1 signal sequence, which previous studies have shown display
similar ER morphology[6]. Using these ER-localized fluorophores, we did not observe major
changes to ER morphology in glutamatergic, octopaminergic, or GABAergic *xbp-1s* animals (**Fig.**
5D). Next, to measure ER secretory capacity, we utilized the yolk protein marker VIT-2::GFP. This
maternal yolk protein is secreted by the intestinal ER in adults and subsequently endocytosed by
developing eggs and is a commonly used marker for secretory capacity[56,57]. Although fluorescent
levels appear to be higher in intact animals (**Fig. S4A**), when we quantitatively measured
fluorescent levels in isolated eggs, there was no significant change in VIT-2::GFP signal (**Fig.**
S4B-C), suggesting that there are no changes to ER secretory capacity in glutamatergic,
octopaminergic, or GABAergic *xbp-1s* animals.
770
***Glutamatergic xbp-1s* promotes protein homeostasis and ER stress resilience.**
Finally, we measured the impact neuronal subtype *xbp-1s* on protein homeostasis, as UPR[ER]
activation is also directly linked to improved protein homeostasis[7]. We crossed glutamatergic,
octopaminergic, and GABAergic overexpressing *xbp-1s* strains into animals expressing
fluorescently-tagged aggregation-prone polyglutamine repeats in the intestine[58] and assessed
the extent of aggregation as these animals aged. Strikingly, a significant decrease in
fluorescence intensity, demonstrating a reduction of polyQ40 aggregation, was observed at

days 1 and 5 of adulthood in all neuronal subtype *xbp-1s* animals as compared to controls (**Fig**
6A-B). Moreover, this was not due to artifacts in our transgenic animal synthesis, as
independently synthesized transgenic lines recapitulated these phenotypes (**Fig. 6C-D**). These
data suggest that while only glutamatergic *xbp-1s* improved ER proteotoxic stress resistance, all
neuronal subtype *xbp-1s* animals have improved peripheral protein homeostasis.

To investigate the physiological impact of the improved protein homeostasis in these animals,
we next measured the impact of neuronal subtype *xbp-1s* on animal survival upon polyQ40
overexpression. Interestingly, octopaminergic *xbp-1s* animals do display a significant increase in
lifespan in animals overexpressing polyQ40 (**Fig. S5A**). Similar to standard lifespan extension,
this resistance to polyQ overexpression was specific to "normal" sized octopaminergic *xbp-1s*
animals, as stunted growth octopaminergic *xbp-1s* animals had reduced lifespan. In addition,
glutamatergic and GABAergic *xbp-1s* animals also displayed reduced lifespan in polyQ
overexpression conditions. While these data may suggest that octopaminergic *xbp-1s* is the
only condition that protects against polyQ overexpression, we found that intestinal polyQ40
overexpression did not consistently reduce lifespan in wild-type animals (**Fig. S5B**). Therefore, it
is entirely possible that polyQ40 lifespans do not directly correlate with protein homeostasis
capacity.

Therefore, we next measured the ability of neuronal subtype *xbp-1s* animals to resist against
protein misfolding stress in the ER. Tunicamycin is a well-characterized ER stressor which
blocks N-linked glycosylation in the ER[59], and animals with neuronal *xbp-1s* exhibit an increased
resistance to tunicamycin[2]. Consistent with our transcriptomics data, glutamatergic *xbp-1s*
resulted in a small but significant increase in tunicamycin resistance (**Fig. S5C**). Interestingly,
octopaminergic and GABAergic *xbp-1s* had no effect on resistance to ER stress (**Fig. S6C-D**),
suggesting that even though polyQ aggregation load is decreased in these animals, they do not
display a significant impact on proteotoxic stress resilience.

Conclusions

The UPR[ER] is involved in diverse cellular processes that impact organismal health, including
proteostasis[2,60], autophagy[7], lipid metabolism[5,6], and immune response[61]. Many of these
functions can occur in a non-autonomous fashion, whereby neural cells with XBP-1s activation
signal to the body to coordinate a homeostatic response[46]. Numerous neural circuits have been
implicated in this response, including serotonin, dopamine[3], tyramine[4], RIM/RIC interneurons[11],
and glial cells[8]. This study further elucidates this complex neural circuitry, identifying additional
functional roles for glutamatergic, octopaminergic, and GABAergic neurons in non-autonomous
UPR[ER] signaling.

The overexpression of *xbp-1s* in glutamatergic, octopaminergic, and GABAergic neurons
requires several considerations. First, promoter strength and neuron number can drive different
phenotypes across each neuronal *xbp-1s* overexpression paradigm, which is unrelated to the
biological significance of a particular neuronal identity. This is especially important to consider
as using two different neuron-specific promoters, *rab-3p* and *rgef-1p*, display dramatically
different downstream transcriptional responses. However, our data show that *xbp-1s* expression
level alone does not purely drive phenotypic outcome. While previous studies utilizing pan-
neuronal, serotonergic, or dopaminergic *xbp-1s* showed a significant increase in *xbp-1s*
overexpression[3], in our study, although we can see a trend for an increase in whole-body *xbp-
1s* expression in glutamatergic, octopaminergic, and GABAergic neuron-driven *xbp-1s*
overexpression, these data did not reach statistical significance. Despite this lack of a significant
increase in *xbp-1s* levels, we still observed dramatic transcriptomic changes, especially in

glutamatergic and octopaminergic *xbp-1s*, with changes in expression in canonical XBP-1s
targets[62], protein homeostasis[2], and immune response[41] pathways. Further, while there are 79
glutamatergic and 34 GABAergic neurons and glutamate and GABA are the most prevalent
excitatory and inhibitory neurotransmitters, respectively, here we only find increased longevity
when *xbp-1s* is expressed in a pair of octopaminergic neurons. This finding is consistent with
other analogous non-autonomous signaling pathways that have system-wide effects dependent
on just a handful of neurons. For example, longevity conferred by dietary restriction, one of the
best conserved lifespan-extending paradigms across diverse animal species, is dependent on
the two ASI neurons in *C. elegans*[63]. Similarly, non-autonomous *xbp-1s* signaling has been
shown to extend lifespan when expressed in a relatively small numbers of cells, including: the
four cephalic sheathe glia[8], the four neurons expressing the tyramine synthesis gene[4], the six
serotonergic neurons, the eight dopaminergic neurons[3], and in this work, the two
octopaminergic neurons. A possible explanation for this is that: 1) it would be inefficient for
numerous neurons to redundantly carry out the same function and 2) in order to enact broad
effects on metabolism, proteostasis, etc., a sufficiently specific signal is preferred to one that
might interfere with the diverse signaling performed by the many glutamatergic and GABAergic
neurons. Thus, if we were to overexpress *xbp-1s* in specific subtypes of glutamatergic or
GABAergic neurons, we may observe more robust phenotypic effects. Taken together, these
data argue that neuron number and promoter strength alone do not drive phenotypic outcomes,
and neuronal identity is a critical factor in non-autonomous signaling, even when using an
artificial system such as *xbp-1s* overexpression.
849
Moreover, serotonergic and dopaminergic *xbp-1s*, both of which have lifespan extensions, were
shown to be beneficial by eliciting distinct responses in the periphery. In contrast to
dopaminergic *xbp-1s* driving lipid depletion, serotonergic *xbp-1s* resulted in an increase in lipids.
These two neuronal subtypes, at least in terms of lipid remodeling, have opposing effects.
Beyond neural *xbp-1s*, a similar phenomenon is observed in whole-body *xbp-1s* overexpression
wherein no extension in lifespan occurs, which may be due to the cumulative effects of muscle
xbp-1s shortening lifespan while intestinal and neuronal *xbp-1s* extend lifespan. Altogether,
these data suggest that overexpression of *xbp-1s* in a smaller subset of cell types may be more
effective at revealing specific downstream pathways, avoiding pleiotropic effects which can
occur in more broad *xbp*-1s overexpression.
860
In a previous study, it was found that overexpressing *xbp-1s* driven by the tyramine synthesis
gene, *tdc-1*, was sufficient to extend longevity and drive non-autonomous UPR^{ER}[4]. Two pairs of
neurons express tyramine as tyramine is the chemical precursor of octopamine: the
octopaminergic RIC neurons and the tyraminergic RIM neurons. It was determined that UPR^{ER}
was upregulated in a tyraminergic signaling specific manner. Whether lifespan extension
depended on tyramine, octopamine, or a combination of these, however, was left ambiguous as
the *tdc-1* promoter used to express *xbp-1s* targets both RIM and RIC neurons. In this study, a
potential resolution to this ambiguity is presented. Here, we overexpress *xbp-1s* exclusively in
the RIC neurons as opposed to the RIM neurons which signal through both tyramine and
glutamate. Thus, our glutamatergic neuron overexpression of *xbp-1s* also targets the
tyraminergic RIM neurons, to the exclusion of octopaminergic RIC neurons[33,37]. Here, we find
that glutamatergic overexpression of *xbp-1s* increases resilience to tunicamycin, a phenotype
related to UPR^{ER}. Thus, our findings on glutamatergic *xbp-1s* may be phenotypes driven by the
tyraminergic RIM neurons, rather than other glutamatergic neuron subtypes. Further supporting
this idea, previous work using a transcriptional reporter in a pan-neuronal *xbp-1s*
overexpression animal has shown the tyramine synthesis gene, but not the glutamatergic *vglut*
gene, was required for upregulation of UPR^{ER}[4]. As this condition leaves tyraminergic signaling,
but not glutamatergic signaling, of RIM neurons intact, it suggests that RIM neurons drive

UPR^{ER} in a tyramine-dependent manner, but not lifespan, which results from octopaminergic
signaling by RIC neurons.
881
Interestingly, we find that although GABAergic *xbp-1s* had minimal changes to gene expression,
these animals still displayed significant changes to organismal health, including improved
proteostasis and immune response. While transcriptional changes are not the only change that
could translate to physiology, as altered protein function, organelle dynamics, and metabolism
can all occur in the absence of transcriptional change, technical limitations could also be
responsible for the lack of difference observed in GABAergic *xbp-1s*. Here, we used whole-
worm transcriptomics, and it is entirely possible that opposite changes in gene expression in
different tissues could result in a net result of no change. Indeed, in terms of *xbp-1s*
overexpression, this is observed wherein whole-body overexpression of *xbp-1s* does not result
in lifespan extension, likely due to the summation of negative effects in the muscle and positive
effects in the intestine and neurons[2]. Thus, it is entirely possible that GABAergic *xbp-1s* may
drive differential effects in different tissue, as it does drive depletion of lipids and increased
protein homeostasis in the intestine. Future tissue-specific studies can reveal whether these
physiological outputs are dependent on gene expression changes in the intestine in these
animals.
897
A potential limitation of our findings is whether ectopic gene overexpression correlates with the
endogenous roles these neuronal subtypes play in signaling or gene regulation. Previous work
in numerous animal models provide sufficient evidence that this is the case. In *C. elegans*,
olfactory sensation of pathogenic bacteria utilize neuron-to-body XBP-1s signaling through TGF-
β signaling to improve longevity and healthspan[11]. In mice, Xbp1s overexpression in POMC
neurons promotes adipose tissue UPR^{ER} to improve metabolic health[9]. Similarly, hepatic *Xbp1s*
activation in mice promotes metabolic health downstream of food perception[10]. In *D.*
melanogaster, glutamate signaling can promote lipid mobilization as a systemic metabolite,
altering lipid metabolism[64]. These reports suggest that even ectopic genetic models can provide
mechanistic insight into important endogenous physiological processes.
908
Lastly, in addition to neuronal identity, there exist complex neuronal circuits that involve
signaling between different neuronal subtypes – some of which are discussed here – which may
contribute to phenotypic differences. In this study, we are generalizing neuronal signals as
separate entities, although neuronal circuits are often intertwined and complex [65]. While our
previous study has separated the utility of dopamine in serotonergic *xbp-1s* signaling and vice
versa [3], this does not preclude the convergence of other neuronal subtypes. For example, we
find that although glutamatergic *xbp-1s* animals display mostly changes to protein homeostasis-
related pathways, they also show improved immune function and increased lipid depletion.
While the improvement in protein homeostasis could be responsible for increased immune
function [41], it is entirely possible that glutamatergic neurons can also recruit octopaminergic or
GABAergic signaling to alter immune response and lipid metabolism. Future studies mapping
the neural circuitry across subtypes will be necessary to develop a full neural map of non-
autonomous XBP-1s signaling. Overall, our study further builds on the complex literature of non-
autonomous XBP-1s signaling, adding three additional neuronal subtypes to this rapidly
expanding map.
924
925

Acknowledgements

927   A.J.C., A.J.H., M.A., and G.G. are supported by T32AG052374; A.J.H. is supported by the NSF
928   GRFP; S.P.C. is supported by NIH R01AG058610 and Hevolution Foundation award 748 HF

AGE-004; B.A.B. is supported by the Simons Collaboration on Plasticity in the Aging Brain grant SF811217; P.J.M. is supported by the Impetus Grants 014746-00001 and the Baxter award; and R.H.S. is supported by R00AG065200 and R01AG079806 from the National Institute on Aging and the Glenn Foundation for Medical Research and AFAR Grant for Junior Faculty Award. Some strains were provided by the CGC, which is funded by the NIH Office of Research Infrastructure Programs (P40 OD010440). Some gene analysis was performed using Wormbase, which is funded on a U41 grant HG002223.

Competing Financial Interests

All authors of the manuscript declare that they have no competing interests.

Data Availability

All data required to evaluate the conclusions in this manuscript are available within the manuscript and Supporting Information. All strains synthesized in this manuscript are derivatives of N2 or other strains from CGC and are either made available on CGC or available upon request. All raw RNA-seq datasets are available through Annotare 2.0 Array Express Accession E-MTAB-14132.

References

1. Dutta, N., Garcia, G. & Higuchi-Sanabria, R. Hijacking Cellular Stress Responses to Promote Lifespan. *Front. Aging* **3**, 860404 (2022).

2. Taylor, R. C. & Dillin, A. XBP-1 Is a Cell-Nonautonomous Regulator of Stress Resistance and Longevity. *Cell* **153**, 1435–1447 (2013).

3. Higuchi-Sanabria, R. *et al.* Divergent Nodes of Non-autonomous UPRER Signaling through Serotonergic and Dopaminergic Neurons. *Cell Reports* **33**, 108489 (2020).

4. Özbey, N. P. *et al.* Tyramine Acts Downstream of Neuronal XBP-1s to Coordinate Inter-tissue UPRER Activation and Behavior in C. elegans. *Developmental Cell* **55**, 754-770.e6 (2020).

5. Imanikia, S., Sheng, M., Castro, C., Griffin, J. L. & Taylor, R. C. XBP-1 Remodels Lipid Metabolism to Extend Longevity. *Cell Reports* **28**, 581-589.e4 (2019).

6. Daniele, J. R. *et al.* UPRER promotes lipophagy independent of chaperones to extend life span. *Science Advances* **6**, eaaz1441 (2020).

7. Imanikia, S., Özbey, N. P., Krueger, C., Casanueva, M. O. & Taylor, R. C. Neuronal XBP-1 Activates Intestinal Lysosomes to Improve Proteostasis in C. elegans. *Current Biology* **29**, 2322-2338.e7 (2019).

8. Frakes, A. E. *et al.* Four glial cells regulate ER stress resistance and longevity via neuropeptide signaling in C. elegans. *Science* **367**, 436–440 (2020).

9. Williams, K. W. *et al.* Xbp1s in Pomc Neurons Connects ER Stress with Energy Balance and Glucose Homeostasis. *Cell Metabolism* **20**, 471–482 (2014).

10. Brandt, C. *et al.* Food Perception Primes Hepatic ER Homeostasis via Melanocortin-Dependent Control of mTOR Activation. *Cell* **175**, 1321-1335.e20 (2018).

11. De-Souza, E. A., Thompson, M. A. & Taylor, R. C. Olfactory chemosensation extends lifespan through TGF-β signaling and UPR activation. *Nat Aging* **3**, 938–947 (2023).

12. Bar-Ziv, R. *et al.* Measurements of Physiological Stress Responses in C. Elegans. *JoVE (Journal of Visualized Experiments)* e61001 (2020) doi:10.3791/61001.

13. Garcia, G., Homentcovschi, S., Kelet, N. & Higuchi-Sanabria, R. Imaging of Actin Cytoskeletal Integrity During Aging in C. elegans. in *Cytoskeleton : Methods and Protocols* (ed. Gavin, R. H.) 101–137 (Springer US, New York, NY, 2022). doi:10.1007/978-1-0716-1661-1_5.

14. Schindelin, J. *et al.* Fiji: an open-source platform for biological-image analysis. *Nat Methods*
9, 676–682 (2012).

15. Egge, N. *et al.* Age-Onset Phosphorylation of a Minor Actin Variant Promotes Intestinal
Barrier Dysfunction. *Developmental Cell* **51**, 587-601.e7 (2019).

16. Grant, B. & Hirsh, D. Receptor-mediated Endocytosis in the Caenorhabditis elegans Oocyte.
MBoC **10**, 4311–4326 (1999).

17. Lord, S., Velle, K., Mullins, D. & Fritz-Laylin, L. SuperPlots: Communicating reproducibility
and variability in cell biology. *Journal of Cell Biology* **219**, e202001064 (2020).

18. Stuhr, N. *et al.* Rapid Lipid Quantification in Caenorhabditis elegans by Oil Red O and Nile
Red Staining. *BIO-PROTOCOL* **12**, (2022).

19. Lynn, D. A. *et al.* Omega-3 and -6 fatty acids allocate somatic and germline lipids to ensure
fitness during nutrient and oxidative stress in Caenorhabditis elegans. *Proceedings of the*
National Academy of Sciences **112**, 15378–15383 (2015).

20. Dobin, A. *et al.* STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**, 15–21 (2013).

21. Love, M. I., Huber, W. & Anders, S. Moderated estimation of fold change and dispersion for
RNA-seq data with DESeq2. *Genome Biol* **15**, 550 (2014).

22. Calfon, M. *et al.* IRE1 couples endoplasmic reticulum load to secretory capacity by
processing the XBP-1 mRNA. *Nature* **415**, 92–96 (2002).

23. Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list enrichment analysis
tool. *BMC Bioinformatics* **14**, 1–14 (2013).

24. Kuleshov, M. V. *et al.* Enrichr: a comprehensive gene set enrichment analysis web server
2016 update. *Nucleic Acids Research* **44**, W90–W97 (2016).

25. Stroustrup, N. *et al.* The Caenorhabditis elegans Lifespan Machine. *Nature Methods* **10**,
665–670 (2013).

26. Torres, T. C. *et al.* Surveying Low-Cost Methods to Measure Lifespan and Healthspan in
Caenorhabditis elegans. *JoVE (Journal of Visualized Experiments)* e64091 (2022)
doi:10.3791/64091.

27. Cezairliyan, B. *et al.* Identification of Pseudomonas aeruginosa Phenazines that Kill
Caenorhabditis elegans. *PLOS Pathogens* **9**, e1003101 (2013).

28. Mahajan-Miklos, S., Tan, M.-W., Rahme, L. G. & Ausubel, F. M. Molecular Mechanisms of
Bacterial Virulence Elucidated Using a Pseudomonas aeruginosa– Caenorhabditis elegans
Pathogenesis Model. *Cell* **96**, 47–56 (1999).

29. Nair, T., Weathers, B. A., Stuhr, N. L., Nhan, J. D. & Curran, S. P. Serotonin deficiency from
constitutive SKN-1 activation drives pathogen apathy. Preprint at
https://doi.org/10.1101/2024.02.10.579755 (2024).

30. Stuhr, N. L., Ramos, C. M., Turner, C. D., Soukas, A. A. & Curran, S. P. C. elegans display
antipathy behavior towards food after contemporaneous integration of nutritional needs and
dietary lipid availability. 2024.02.23.581740 Preprint at
https://doi.org/10.1101/2024.02.23.581740 (2024).

31. Urano, F. *et al.* A survival pathway for Caenorhabditis elegans with a blocked unfolded
protein response. *Journal of Cell Biology* **158**, 639–646 (2002).

32. Bar-Ziv, R. *et al.* Glial-derived mitochondrial signals affect neuronal proteostasis and aging.
Science Advances **9**, eadi1411 (2023).

33. Serrano-Saiz, E. *et al.* Modular Control of Glutamatergic Neuronal Identity in C. elegans by
Distinct Homeodomain Proteins. *Cell* **155**, 659–673 (2013).

34. Sellegounder, D., Yuan, C.-H., Wibisono, P., Liu, Y. & Sun, J. Octopaminergic Signaling
Mediates Neural Regulation of Innate Immunity in Caenorhabditis elegans. *mBio* **9**, e01645-
18 (2018).

35. McIntire, S. L., Jorgensen, E. & Horvitz, H. R. Genes required for GABA function in
Caenorhabditis elegans. *Nature* **364**, 334–337 (1993).

36. Lee, R. Y. N., Sawin, E. R., Chalfie, M., Horvitz, H. R. & Avery, L. EAT-4, a Homolog of a
Mammalian Sodium-Dependent Inorganic Phosphate Cotransporter, Is Necessary for
Glutamatergic Neurotransmission in Caenorhabditis elegans. *J Neurosci* **19**, 159–167
(1999).
37. Alkema, M. J., Hunter-Ensor, M., Ringstad, N. & Horvitz, H. R. Tyramine Functions
Independently of Octopamine in the Caenorhabditis elegans Nervous System. *Neuron* **46**,
247–260 (2005).
38. Jin, Y., Jorgensen, E., Hartwieg, E. & Horvitz, H. R. The Caenorhabditis elegans Gene unc-
25Encodes Glutamic Acid Decarboxylase and Is Required for Synaptic Transmission But
Not Synaptic Development. *J Neurosci* **19**, 539–548 (1999).
39. Gelino, S. *et al.* Intestinal Autophagy Improves Healthspan and Longevity in C. elegans
during Dietary Restriction. *PLoS Genet* **12**, e1006135 (2016).
40. Yang, Y. *et al.* Autophagy protein ATG-16.2 and its WD40 domain mediate the beneficial
effects of inhibiting early-acting autophagy genes in C. elegans neurons. *Nat Aging* **4**, 198–
212 (2024).
41. Richardson, C. E., Kooistra, T. & Kim, D. H. An essential role for XBP-1 in host protection
against immune activation in C. elegans. *Nature* **463**, 1092–1095 (2010).
42. Kwon, S., Kim, E. J. E. & Lee, S.-J. V. Mitochondria-mediated defense mechanisms against
pathogens in Caenorhabditis elegans. *BMB Rep* **51**, 274–279 (2018).
43. Liu, Y. & Chang, A. Heat shock response relieves ER stress. *EMBO J.* **27**, 1049–1059
(2008).
44. Glover-Cutter, K. M., Lin, S. & Blackwell, T. K. Integration of the Unfolded Protein and
Oxidative Stress Responses through SKN-1/Nrf. *PLOS Genetics* **9**, e1003701 (2013).
45. Anderson, N. & Haynes, C. M. Folding the Mitochondrial UPR into the Integrated Stress
Response. *Trends Cell Biol* **30**, 428–439 (2020).
46. Homentcovschi, S. & Higuchi-Sanabria, R. A neuron's ambrosia: non-autonomous unfolded
protein response of the endoplasmic reticulum promotes lifespan. *Neural Regen Res* **17**,
309–310 (2022).
47. Tan, M.-W., Mahajan-Miklos, S. & Ausubel, F. M. Killing of Caenorhabditis elegans by
Pseudomonas aeruginosa used to model mammalian bacterial pathogenesis. *Proceedings
of the National Academy of Sciences* **96**, 715–720 (1999).
48. Meisel, J. D. & Kim, D. H. Behavioral avoidance of pathogenic bacteria by *Caenorhabditis
elegans*. *Trends in Immunology* **35**, 465–470 (2014).
49. Zhang, Y., Lu, H. & Bargmann, C. I. Pathogenic bacteria induce aversive olfactory learning
in Caenorhabditis elegans. *Nature* **438**, 179–184 (2005).
50. O'Donnell, M. P., Fox, B. W., Chao, P.-H., Schroeder, F. C. & Sengupta, P. A
neurotransmitter produced by gut bacteria modulates host sensory behaviour. *Nature* **583**,
415–420 (2020).
51. Egge, N. *et al.* Age-Onset Phosphorylation of a Minor Actin Variant Promotes Intestinal
Barrier Dysfunction. *Developmental Cell* **51**, 587-601.e7 (2019).
52. Garcia, G. *et al.* Large-scale genetic screens identify BET-1 as a cytoskeleton regulator
promoting actin function and life span. *Aging Cell* **22**, e13742 (2023).
53. Na, H. *et al.* Identification of lipid droplet structure-like/resident proteins in Caenorhabditis
elegans. *Biochimica et Biophysica Acta (BBA) - Molecular Cell Research* **1853**, 2481–2491
(2015).
54. Zhang, P. *et al.* Proteomic study and marker protein identification of Caenorhabditis elegans
lipid droplets. *Mol. Cell Proteomics* **11**, 317–328 (2012).
55. Du, J., Zhao, L., Kang, Q., He, Y. & Bi, Y. An optimized method for Oil Red O staining with
the salicylic acid ethanol solution. *Adipocyte* **12**, 2179334.
56. Grant, B. & Hirsh, D. Receptor-mediated endocytosis in the Caenorhabditis elegans oocyte.
Mol. Biol. Cell **10**, 4311–4326 (1999).

57. Stevens, J. & Spang, A. N-glycosylation is required for secretion and mitosis in C. elegans.
PLoS ONE **8**, e63687 (2013).
58. Morley, J. F., Brignull, H. R., Weyers, J. J. & Morimoto, R. I. The threshold for
polyglutamine-expansion protein aggregation and cellular toxicity is dynamic and influenced
by aging in Caenorhabditis elegans. *Proceedings of the National Academy of Sciences* **99**,
10417–10422 (2002).
59. Heifetz, A., Keenan, R. W. & Elbein, A. D. Mechanism of action of tunicamycin on the UDP-
GlcNAc:dolichyl-phosphate Glc-NAc-1-phosphate transferase. *Biochemistry* **18**, 2186–2192
(1979).
60. Frakes, A. E. & Dillin, A. The UPR(ER): Sensor and Coordinator of Organismal
Homeostasis. *Mol. Cell* **66**, 761–771 (2017).
61. Grootjans, J., Kaser, A., Kaufman, R. J. & Blumberg, R. S. The unfolded protein response in
immunity and inflammation. *Nat Rev Immunol* **16**, 469–484 (2016).
62. Shen, X., Ellis, R. E., Sakaki, K. & Kaufman, R. J. Genetic interactions due to constitutive
and inducible gene regulation mediated by the unfolded protein response in C. elegans.
PLoS Genet. **1**, e37 (2005).
63. Bishop, N. A. & Guarente, L. Two neurons mediate diet-restriction-induced longevity in C.
elegans. *Nature* **447**, 545–549 (2007).
64. Zhao, X. & Karpac, J. Glutamate metabolism directs energetic trade-offs to shape host-
pathogen susceptibility in Drosophila. *Cell Metabolism* **33**, 2428-2444.e8 (2021).
65. Randi, F., Sharma, A. K., Dvali, S. & Leifer, A. M. Neural signal propagation atlas of
Caenorhabditis elegans. *Nature* **623**, 406–414 (2023).

1100

Supporting Information

1102

Fig. S1. A comparison of pan-neuronal *xbp-1s* expression via two different promoters reveals differences in whole-body transcriptomic changes. Volcano plots of whole-body genome-wide changes in gene expression upon pan-neuronal *xbp-1s* overexpression driven by **(A)** *rab-3* promoter or **(B)** *rgef-1* promoter. Red dots indicate significantly differentially expressed genes with p-value ≤ 0.01. See **Table S3** for a list of differentially expressed genes and expression values. **(C)** Comparison of differentially expressed genes (p-value ≤ 0.01) between worms expressing *xbp-1s* pan-neuronally driven by *rab-3p* or *rgef-1p*. For a complete list of differentially expressed genes in each group, see **Table S4**. **(D)** Heat map of common differentially expressed genes upon pan-neuronal *xbp-1s* expression under control of *rab-3* or *rgef-1* promoter. Warmer colors indicate increased expression, and cooler colors indicate decreased expression. See **Table S5** for a list of genes and values. **(E)** Heat map of UPR[ER] related gene (GO:0030968) expression under pan-neuronal *xbp-1s* driven by *rab-3* or *rgef-1* promoter. Warmer colors indicate increased expression, and cooler colors indicate decreased expression. See **Table S5** for a list of genes and values. **(F)** Heat map of XBP-1s target gene [31] expression under pan-neuronal *xbp-1s* driven by *rab-3* or *rgef-1* promoter. Warmer colors indicate increased expression, and cooler colors indicate decreased expression. See **Table S5** for a list of genes and values.

1120

Figure S2. *xbp-1s* overexpression in *C. elegans* glutamatergic, octopaminergic, and GABAergic neurons drives differential gene expression. Comparison of differentially expressed genes (p-value ≤ 0.01) in neuronal *xbp-1s* driven by *rgef-1p* and **(A)** glutamatergic (*eat-4p*), **(B)** octopaminergic *(tbh-1p)*, and **(C)** GABAergic (*unc-25p*) *xbp-1s* expression. For a complete list of differentially expressed genes in each group, see **Table S3**. Comparison of differentially expressed genes (p-value ≤ 0.01) in glutamatergic, octopaminergic, GABAergic,

and **(D)** dopaminergic or **(E)** serotonergic neurons. See **Table S4**. Gene expression changes in
groups of genes related to **(F)** UPR[ER] (GO:0030968), **(G)** XBP-1s targets [31], **(H)** mitochondrial
unfolded protein response (GO:0034514), **(I)** heat shock response (GO:0009408), **(J)** oxidative
stress response (GO:0006979), and **(K)** translation (GO:0006412). Significance was determined
using a one-sample Wilcoxon test. * = $p \le 0.05$, ** = $p \le 0.01$, *** = $p \le 0.001$, **** = $p \le 0.0001$,
ns = $p > 0.05$

1133

Fig. S3. Glutamatergic, octopaminergic, and GABAergic *xbp-1s* does not increase
healthspan. Measurements of fecundity of control (blue) and **(A)** glutamatergic *xbp-1s* (green,
eat-4p), **(B)** octopaminergic *xbp-1s* (yellow, *tbh-1p*), and **(C)** GABAergic *xbp-1s* (pink, *unc-25p*)
animals. Total number of eggs that hatched were counted per animal. Measurements of
thrashing of control (blue) and **(D)** glutamatergic *xbp-1s* (green, *eat-4p*), **(E)** octopaminergic
xbp-1s (yellow, *tbh-1p*), and **(F)** GABAergic *xbp-1s* (pink, *unc-25p*) animals. Number of thrashes
was assessed over a 10 second period in animals at day 1 adult (young), day 4-5 adult (middle),
and day 9 adult (old) in M9 solution with each thrash being counted as a movement from a
concave to a convex formation. For SuperPlots, each small dot represents a single animal with
various intensities of colors representing independent biological replicates and each large dot is
the median value of each biological replicate. Lines represent the median across all biological
replicates and whiskers indicate interquartile range. Statistical analysis was performed using a
Mann-Whitney test.

1147

Fig. S4. Glutamatergic, octopaminergic, and GABAergic *xbp-1s* does not alter ER
secretory capacity. (A) Representative fluorescent micrographs of day 3 adult animals of
VIT2::GFP in control, glutamatergic *xbp-1s* (*eat-4p*), octopaminergic *xbp-1s* (*tbh-1p*), or
GABAergic *xpb-1s* (*unc-25p*). All images are contrast-matched. **(B)** Representative fluorescent
micrographs of eggs collected using a standard bleaching protocol of day 3 adult animals of
VIT2::GFP in control, glutamatergic *xbp-1s* (*eat-4p*), octopaminergic *xbp-1s* (*tbh-1p*), or
GABAergic *xpb-1s* (*unc-25p*). All images are contrast-matched. Scale bar represents 500 μm.
(C) Quantification of eggs from (B) using measurements of integrated intensity. For SuperPlots,
each small dot represents a single animal with various intensities of colors representing
independent biological replicates and each large dot is the median value of each biological
replicate. Lines represent the median across all biological replicates and whiskers indicate
interquartile range. Statistical analysis was performed using a Mann-Whitney test.

1160

Fig. S5. Octopaminergic *xbp-1s* promotes lifespan in polyQ40 expression animals, but
not on tunicamycin. (A) Lifespan measurements of control (here, *vha-6p::polyQ*), blue),
glutamatergic *xbp-1s* animals (*eat-4p*, green), octopaminergic *xbp-1s* separated for normal
sized (*tbh-1,* yellow) or stunted growth (*tbh-1,* small, purple), and GABAergic *xbp-1s* (*unc-25p,*
pink) animals expressing polyQ40::YFP in the intestine (*vha-6p::polyQ*). **(B)** Median lifespan
measurements from 5 replicates of wild-type N2 animals either expressing *vha-6p::polyQ40*
(light blue, *vha-6p::polyQ*) or not (dark blue, control). Dots indicate each biological replicate and
lines represent median plus interquartile range. **(C)** Lifespan measurements of control (blue,
light blue), glutamatergic *xbp-1s* animals (*eat-4p*, green), and GABAergic *xbp-1s* (*unc-25p,* pink)
animals moved onto 25 mg/mL tunicamycin (TM) plates starting from day 1 of adulthood.
Lifespans were scored every 2 days and data is representative of 3 biological replicates (N). **(D)**
Lifespan measurements of control (blue) and octopaminergic *xbp-1s* animals moved onto 25
mg/mL tunicamycin (TM) plates starting from day 1 of adulthood. Octopaminergic *xbp-1s*
animals were separated into normal size (yellow, *tbh-1p*) and stunted growth (purple, *tbh-1p,*
small). Sample size (n) is written next to each condition followed by significance measured

using Log-Rank testing: n.s. = not significant, * = p < 0.05, *** = p < 0.001. All statistical analysis
is available in **Table S7**.

1178
1179   **Table S1. Strains used in this study.**

1180
1181   **Table S2. Primers used in this study.**

1182
1183   **Table S3. List of differentially expressed genes.**

1184
1185   **Table S4. List of all genes in Venn Diagrams.**

1186
1187   **Table S5. List of all genes used in heat maps.**

1188
1189   **Table S6. List of all gene ontologies.**

1190
1191   **Table S7. Lifespan statistics.**

A *rab-3p* *dat-1p*
745 4 75

B *rab-3p* *tph-1p*
745 4 100

C *rab-3p* *dat-1p/tph-1p*
669 83 1438

D *rab-3p* DEGs comparison

log₂(fold change)

rab-3p *dat-1p* *tph-1p* *dat-1p/tph-1p*

E

glutamatergic = 79 neurons

octopaminergic = 2 neurons

GABAergic = 32 neurons

F

Fold Change *xbp-1s* (Log2)

eat-4p *tbh-1p* *unc-25p*

A Lipid Regulation

B DHS-3::GFP

C control, eat-4p, tbh-1p, unc-25p; Proportion of worms

D mRuby::HDEL; mCherry::HDEL