Title: **Mechanism of RSL3-Induced Ferroptotic Cell Death in HT22 Cells: Crucial Role of Protein Disulfide Isomerase**

Authors: Ming-Jie Hou[1], Xuanqi Huang[1] and Bao Ting Zhu[1,2]

Affiliations: [1]Shenzhen Key Laboratory of Steroid Drug Discovery and Development, School of Medicine, The Chinese University of Hong Kong, Shenzhen, Guangdong, 518172, China, and [2]Shenzhen Bay Laboratory, Shenzhen 518172, China

Correspondence: Bao Ting Zhu, School of Medicine, The Chinese University of Hong Kong, 2001 Longxiang Blvd., Longgang District, Shenzhen, Guangdong 518172, China.

E-MAIL: **BTZhu@CUHK.edu.cn**. PHONE: **+086-755-84273851**.

Running title: Role of PDI in RSL3-Induced Ferroptosis

Abbreviations: PDI, protein disulfide isomerase; nNOS, neuronal nitric oxide synthase; iNOS, inducible nitric oxide synthase; NO, nitric oxide; ROS, reactive oxygen species; TrxR1, thioredoxin reductase 1

Conflict of interest: The authors declare no conflict of interest.

1 ABSTRACT

Protein disulfide isomerase (PDI) was recently shown to be an upstream mediator

of erastin-induced, glutathione depletion-associated ferroptosis through its catalysis of

nitric oxide synthase (NOS) dimerization and nitric oxide (NO) accumulation. A recent

study reported that RSL3, a known ferroptosis inducer and glutathione peroxidase 4

(GPX4) inhibitor, can inhibit thioredoxin reductase 1 (TrxR1). The present study seeks

to test a hypothesis that RSL3 may, through its inhibition of TrxR1, facilitate PDI

activation (*i.e.*, in a catalytically-active, oxidized state), thereby enhancing RSL3-induced

ferroptosis through NOS dimerization and NO accumulation. Using the HT22 mouse

neuronal cells as an *in-vitro* model, we show that treatment of these cells with RSL3 can

strongly increase NOS protein level, and the PDI-mediated NOS dimerization is activated

by RSL3, resulting in NO accumulation. Mechanistically, we find that PDI is activated in

cells treated with RSL3 resulting from its inhibition of TrxR1, and the activated PDI then

catalyzes NOS dimerization, which is followed by accumulation of cellular NO, ROS and

lipid-ROS, and ultimately ferroptotic cell death. Genetic or pharmacological inhibition of

PDI or TrxR1 partially abrogates RSL3-induced NOS activation and the subsequent

accumulation of cellular NO, ROS/lipid-ROS, and ultimately ferroptosis in HT22 cells.

The results of this study clearly show that PDI activation resulting from RSL3 inhibition

of the TrxR1 activity contributes crucially to RSL3-induced ferroptosis in a cell culture

model through the PDI◻NOS◻NO◻ROS/lipid-ROS pathway, in addition to its known

inhibition of the GPX4 activity.

Keywords: Protein disulfide isomerase; Thioredoxin reductase 1; Neuronal nitric

oxide synthase; Nitric oxide; Reactive oxygen species; Lipid reactive oxygen species.

1 INTRODUCTION

Ferroptosis is a form of oxidative cell death [1], and is morphologically distinct from

apoptosis-associated characteristics [1, 2]. Mechanistically, ferroptosis can be

selectively induced through cellular glutathione (GSH) depletion and/or through

suppression of glutathione peroxidase 4 (GPX4), an enzyme involved in the reduction of

lipid peroxides. RSL3 is a prototypical inducer of ferroptosis [3], and over the past few

years, it has been widely thought that RSL3 induces ferroptosis primarily through its

suppression of GPX4 function, which then leads to accumulation of cellular lipid reactive

oxygen species (lipid-ROS) [4]. However, a recent study has reported that RSL3 can also

strongly inhibit the catalytic activity of thioredoxin reductase 1 (TrxR1) [5]. Because

TrxR1 uses NADPH as a cofactor to reduce the active-site disulfide bonds in

thioredoxins, when TrxR1 is inhibited by an inhibitor (such as RSL3), the active site of

the thioredoxins would stay in the oxidized state (*i.e.*, containing a disulfide bond in

their active sites) [6].

Protein disulfide isomerase (PDI or PDIA1) is the prototype of the PDI family

proteins, which belong to the ubiquitous dithiol/disulfide oxidoreductases of the

thioredoxin superfamily [7-9]. PDI is primarily localized in the endoplasmic reticulum of

mammalian cells, although a small fraction of this protein is also found in the nucleus,

cytosol, mitochondria, plasma membrane and extracellular space [10]. PDI is involved in

protein processing by catalyzing the formation of intra- and inter-molecular disulfide

bridges in proteins [11]. Our recent studies have shown that PDI plays an important

role in mediating glutamate- and erastin-induced, GSH depletion-associated oxidative

cytotoxicity through PDI-mediated NOS activation (*i.e.*, homodimer formation through a

disulfide bond linkage) which is followed by the accumulation of cellular NO, ROS and

lipid-ROS, and ultimately the induction of ferroptotic cell death [12].

In the present study, we sought to examine the role of PDI in mediating RSL3-

induced ferroptosis in HT22 neuronal cells and the mechanism of PDI activation. We

found that PDI plays a crucial role in mediating RSL3-induced ferroptotic cell death in

these cells through its ability to catalyze NOS dimerization, which is followed by cellular

accumulation of NO, ROS and lipid-ROS, and ultimately ferroptotic cell death. Genetic or

pharmacological inhibition of PDI's function could abrogate RSL3-induced ferroptosis.

The mechanism by which PDI is activated in RSL3-treated cells is caused by TrxR1

inhibition by RSL3, and this inhibition leaves more PDI proteins in the oxidized state,

which is the active form to catalyze the dimerization of NOS (including both nNOS and

iNOS) in HT22 cells.

MATERIALS AND METHODS

Reagents

RSL3 (#SML2234) and *S*-nitroso-*N*-acetylpenicillamine (SNAP, #N3398) were

purchased from Sigma-Aldrich (St. Louis, MO, USA). CPZ (chlorpromazine, #S5749),

ferrostatin-1 (Fer-1, #S7243), cystamine (#S3695) and 3-(4,5-dimethylthiazol-2-yl)-

2,5-diphenyltetrazolium bromide (MTT, #S6821) were purchased from Selleck

Chemicals (Houston, TX, USA). Diaminofluorescein-FM diacetate (DAF-FM-DA, #S0019)

and diaminofluorescein-2 diacetate (DCF-DA, #S0033) were obtained from Beyotime

Biotechnology (Shanghai, China), and C11-BODIPY-581/591 (#D3861) from Thermo

Fishers (Waltham, MA, USA). Anti-PDI (#3501), anti-TrxR1 (#6925) and anti-β-actin

(#4970) antibodies were purchased from Cell Signaling Technology (Danvers, MA, USA);

anti-GPX4 (#ab125066), anti-nNOS (#ab1376) and anti-iNOS (#ab3523) antibodies

from Abcam (Waltham, MA, USA). TrxR1-siRNAs (#sc-36751) and iNOS-siRNAs (#sc-

36092) were obtained from Santa Cruz Biotechnology (Dallas, TX, USA); PDI-siRNAs and

nNOS-siRNAs were from Ruibo Biotechnology (Guangdong, China). The siRNA

sequences targeting the mouse TrxR1 (siTrxR1) are UUCAAGUAGAUUAGCCAAGtt (#1),

UACGUUAUGAACAGCUUCCtt (#2), and UUACGUUAUGAACAGCUUCtt (#3). The siRNA

sequences targeting the mouse iNOS (si-iNOS) are UUUCCUUUGUUACAGCUUCtt (#1),

UUCAUGAUAACGUUUCUGGtt (#2), and UAGUAGUCCACAAUAGUACtt (#3). The siRNA

sequences targeting the mouse PDI (siPDI) are CCAAGTACCAGCTGGACAA (#1),

GAACGGTCATTGATTACAA (#2), and TGCTAAGATGGACTCAACA (#3). The siRNA

sequences targeting the mouse nNOS (sinNOS) are GCTGCCATCCCATCACATA (#1),

CCTCGTGAATGCACTCATT (#2), and GCGACAATTTGACATCCAA (#3). The siRNA

sequences for negative control (siCon) are AUCCGCGCGAUAGUACGUAtt (#1),

UUACGCGUAGCGUAAUACGtt (#2), and UAUUCGCGCGUAUAGCGGUtt (#3). The MDA

(malondialdehyde) Assay Kit (#M496) was purchased from Dojindo Molecular

Technologies (Kumamoto, Japan), and the TrxR1 Activity Assay Kit (#BC1155) from

Solarbio (Beijing, China).

Cell culture and cell viability assay

HT22 cells were obtained from the Cell Bank of the Chinese Academy of Sciences

(Shanghai, China). The HT22 murine hippocampal neuronal cells were cultured in

DMEM supplied with 10% (*v/v*) fetal bovine serum and 1% penicillin and streptomycin,

and incubated at 37°C under 5% CO_2. Cells were sub-cultured and used for subsequent

experiments when they reached about 80% confluence.

The MTT reduction assay was used to determine the change in gross cell viability.

The HT22 cells were usually seeded in 96-well plates at a density of 2,000 cells/well 24

h prior to receiving different experimental treatments. To test the effect of a modulating

compound, cells were jointly treated with RSL3 and the compound for 24 h. Afterwards,

MTT at a final concentration of 0.5 mg/mL was added to each well, and incubated for 2.5

h at 37°C under 5% CO_2. After incubation, the MTT-containing medium was removed

and 100 μL DMSO was added to each well to dissolve the MTT formazan. Absorbance of

the MTT formazan was measured with a UVmax microplate reader (Molecular Device,

Palo Alto, CA, USA) at 560 nm.

siRNA transfection

For siRNA transfection, the procedures were described in our earlier study [12].

Briefly, the siRNAs (at a final concentration of 60 nM) for targeted genes (PDI, iNOS,

nNOS, and TrxR) were transfected into HT22 cells 24 h after seeding with Lipofectamine

RNA iMAX (Invitrogen). Twenty-four h after siRNA transfection, cells were treated with

selected drugs, and subsequently processed for cell viability measurement, fluorescence

staining and measurement, and immunoblot analysis.

Measurement of cellular NO, ROS and lipid-ROS levels

The cellular NO, ROS and lipid-ROS levels were jointly determined using analytical

flow cytometry and fluorescence/confocal microscopy. The HT22 cells were seeded in

24-well plates at a density of 5×10^4 cells/well 24 h before treatment. Afterwards, the

cells were treated with RSL3 ± different modulating compounds for 6 h or as indicated.

It is of note that the 6-h RSL3 treatment time was selected based on our initial analysis

of the time course of NO, ROS and lipid-ROS accumulation. Prior to staining with

fluorescent probes, cells were washed with HBSS twice. The cellular levels of NO, ROS

and lipid-ROS were detected after addition of respective cell-permeant fluorescent

probes (10 μM DAF-FM-DA for NO detection, 10 μM DCFH-DA for ROS detection, and 10

μM BODIPY^{TM}-581/591-C11 for lipid-ROS detection), which were dissolved in 200 μL

phenol red- and serum-free DMEM. The cells were incubated at 37°C for an additional

30 min, and washed twice with HBSS. The fluorescence images of cellular NO and ROS

were taken using a fluorescence microscope (AXIO, Carl Zeiss Corporation, Germany).

The fluorescence images of the cellular lipid-ROS were captured using a confocal laser

scanning microscope (LSM 900; Carl Zeiss, Oberkochen, Germany) and analyzed with

the Zen software (Carl Zeiss). The quantitative fluorescence intensity reflecting the

cellular accumulation NO, ROS and lipid-ROS was detected using a flow cytometer

(Beckman Coulter, Brea, CA, USA). Three or more experiments were usually conducted

to confirm the observations, and the average fluorescence intensity (mean ± S.D.) based

on the replicate measurements in one representative experiment was calculated with

the FlowJo software (FlowJo, LLC, Ashland, USA) and selected for presentation.

For measuring intracellular MDA levels, a MDA Assay Kit was used, and all

procedures followed manufacturer's instructions.

Measurement of TrxR1 enzyme activity

The TrxR1 enzyme activity was determined using a commercial kit which is based

on the measurement of NADPH-dependent reduction of the disulfide bond in the

substrate 5,5'-dithiobis-2-nitrobenzoic acid (DTNB). To determine the effect by RSL3 or

auranofin on TrxR1's enzyme activity in cultured HT22 cells, the cells were treated with

RSL3 (100 nM) or auranofin (1 μM) for 6 h, and then washed with PBS twice, lysed and

centrifuged at 9500 g for 10 min (4°C) to collect cellular supernatants. Aliquots of the

supernatants were properly diluted and then added into the 96-well plate for

determination of TrxR1 enzyme activity according to manufacturer's instructions. The

change in absorbance at 412 nm was recorded using a UVmax microplate reader

(Molecular Device, Palo Alto, CA, USA). The relative TrxR1 enzyme activity was

calculated as a percentage of the vehicle-treated control cells. To determine the direct

inhibition of TrxR1's catalytic activity by RSL3 and auranofin in the *in-vitro* enzymatic

assay, the untreated HT22 cells were washed with PBS twice, lysed and then centrifuged

at 9500 g for 10 min (4°C) to prepare cellular supernatants which contain the TrxR1

enzyme. Different concentrations of RSL3 and auranofin were included in the

incubation, and the enzyme activity in each treatment group was determined by

monitoring the change in absorbance at 412 nm once every 2 min for up to 150 min.

Here it should be noted that in this enzyme activity assay, it is assumed that the NADPH-

dependent specific catalytic activity of TrxR1 is inhibited by high concentrations of

auranofin (its IC_{50} for TrxR1 is approximately 1.1 nM as determined in this study), and

the baseline reading at absorbance 412 nm which cannot be inhibited by 1000 nM

auranofin is subtracted.

Immunoblot analysis

After treatment with RSL3 for different durations as indicated, the HT22 cells were

collected by trypsinization and centrifugation, and then lysed in a RIPA buffer

containing 1% protease inhibitor cocktail on ice for 30 min. Protein concentrations

were determined using the BCA Assay Kit (ThermoFisher). For total nNOS and iNOS

analysis (including both monomeric and dimeric forms of nNOS and iNOS), samples

were heated at 95°C for 5 min with a reducing buffer before loaded onto the

electrophoresis gel. To analyze the monomeric and dimeric forms of nNOS and iNOS, the

samples were prepared with a non-reducing buffer and were not heated, and the

temperature of the gel was maintained below 15°C during electrophoresis. The proteins

were separated using 10% SDS-PAGE gel (for total nNOS and iNOS analysis) or 6% SDS-

PAGE gel (for monomeric and dimeric nNOS and iNOS analysis), and then transferred to

PVDF membranes. The membranes were incubated for 1 h in 5% skim milk at room

temperature, and then incubated with the primary antibody overnight. Afterwards, the

membranes were washed three times with TBST at room temperature (10 min each

time). The membranes were then incubated with the secondary antibody for 1 h at

room temperature and washed with TBST three times before visualization. Desired

protein bands were visualized using the chemiluminescence imaging system (Tanon

5200, Shanghai, China).

Statistical analysis

All quantitative experiments and data described in this study were repeated at least

three times. The data were usually presented as mean ± S.D. of multiple replicate

measurements from one selected representative experiment. Statistical analysis was

conducted using one-way ANOVA, followed by Dunnett's *post-hoc* tests for multiple

comparisons as needed (GraphPad Prism 7, GraphPad Software, La Jolla, CA). Statistical

significance was denoted by $P < 0.05$ (* or #) and $P < 0.01$ (** or ##) for significant and

very significant differences, respectively. In most cases, * and ** denote the comparison

for statistical significance between the control group (cells treated with vehicle only)

and the cells treated with a chemical (such as RSL3), whereas # and ## denote the

comparison between the cells treated with RSL3 alone and the cells jointly treated with

RSL3 and another compound. For Western blot quantification, one representative data

is shown.

RESULTS

Role of cellular NO, ROS and lipid-ROS accumulation in RSL3-induced ferroptotic

cell death

Treatment of HT22 cells with RSL3-induced cell death in time- and dose-dependent

manners (**Fig. 1A⬚1C**). Ferrostatin-1 (Fer-1), a commonly-used lipid-ROS scavenger [13,

14], showed a strong protection against RSL3-induced cytotoxicity in these cells (**Fig.**

1D), which confirms earlier observations [15, 16] reporting that RSL3 induces

ferroptotic cell death in cultured HT22 cells.

Our recent studies showed that during erastin-induced, GSH depletion-associated

ferroptotic cell death in HT22 cells, there was a time-dependent sequential induction of

NO and ROS/lipid-ROS accumulation [12]. To probe whether RSL3-induced ferroptosis

in HT22 cells is also related to cellular NO and ROS/lipid-ROS accumulation, we

determined the time-dependent change in cellular NO, ROS and lipid-ROS levels

following RSL3 treatment (**Fig. 1E⬚1J**). Based on measurements using fluorescence

microscopy and flow cytometry, we found that treatment of HT22 cells with RSL3

elicited a time-dependent increase in cellular NO level (**Fig. 1E, 1G, Fig. S1A**). Similarly,

the cellular levels of ROS and lipid-ROS were also increased in a time- and dose-

dependent manner in RSL3-treated cells (**Fig. 1E, 1H, Fig. S1B** for ROS; **Fig. 1F, 1I** for

lipid-ROS). Malondialdehyde (MDA), one of the end products of lipid peroxidation and

often used as an indicator of cellular lipid peroxidation [17], was increased by RSL3

treatment in a time-dependent manner (**Fig. 1J**). It was observed that NO accumulation

occurs at a very similar time as does ROS/lipid-ROS accumulation.

To determine whether NO accumulation subsequently leads to ROS and lipid-ROS

accumulation and ultimately ferroptotic cell death, we studied the effect of 2-(4-

carboxyphenyl)-4,4,5,5-tetramethylimidazoline-1-oxyl-3-oxide (cPTIO), a NO scavenger,

in RSL3-treated cells [18]. cPTIO effectively suppressed RSL3-induced accumulation of

NO (**Fig. 2A**), ROS (**Fig. 2B**) and lipid-ROS (**Fig. 2C**) in HT22 cells. In addition, cPTIO also

attenuated RSL3-induced cell death (**Fig. 2D**).

6-Hydroxy-2,5,7,8-tetramethylchroman-2-carboxylic acid (Trolox), which has a

strong ROS-scavenging activity plus a weaker NO-scavenging activity [19], partially

abrogated RSL3-induced accumulation of NO (**Fig. 2E**) but strongly abrogated the

accumulation of ROS (**Fig. 2F**) and lipid-ROS (**Fig. 2G**), and these effects were associated

with a very strong protection against RSL3-induced ferroptotic cell death (**Fig. 2H**).

NAC (*N*-acetyl-*L*-cysteine), another commonly-used antioxidant [20], very strongly

abrogated RSL3-induced accumulation of cellular NO (**Fig. 2I, 2J**), ROS (**Fig. 2K**) and

lipid-ROS (**Fig. 2L**), and these effects were also associated with a strong protection

against RSL3-induced cell death (**Fig. 2M**).

Enhancement of RSL3-induced ferroptotic cell death by sodium nitroprusside

Sodium nitroprusside (SNP) is a NO donor and can rapidly release NO inside the

cells [21]. To provide further support for an upstream role of NO in RSL3-induced

ROS/lipid-ROS accumulation and ferroptosis, we sought to determine the effect of SNP

on RSL3-induced ROS/lipid-ROS accumulation and ferroptosis in HT22 cells. First,

treatment of HT22 cells with SNP alone caused a rapid rise in cellular NO level (**Fig. 3A,**

3B, 3C), which was accompanied by increase in cellular ROS level (**Fig 3A, 3B, 3D**) and

lipid-ROS level (**Fig. 3E**). It is of note while NO accumulation peaked within 0.5 h after

SNP exposure, the accumulation of ROS and lipid-ROS lagged slightly behind NO

increase, reaching the peak at around 1 h after SNP exposure. In addition, treatment of

HT22 cells with SNP alone also elicited a dose-dependent cytotoxicity (**Fig. 3F**).

It is of note that joint treatment of HT22 cells with RSL3 + SNP (200 μM) drastically

sensitized HT22 cells to RSL3-induced ferroptotic cell death (**Fig. 3G**). The observed

enhancement of RSL3's cytotoxicity by SNP was associated in a parallel manner with a

drastically-enhanced accumulation of cellular ROS (**Fig. 3H, 3I**) and lipid-ROS (**Fig. 3J,**

3K, 3L) in cells jointly treated with RSL3 + SNP. Together, these results provide

additional support for the hypothesis that RSL3-induced NO accumulation is an

important determinant for accumulation of cellular ROS and lipid-ROS, and

subsequently ferroptotic cell death.

Role of NOS induction and activation in RSL3-induced ferroptotic cell death

It is known that the generation of cellular NO is mostly catalyzed by the nitric oxide

synthase (NOS), which uses *L*-arginine as its primary substrate [22-24]. There are three

known subtypes of NOS, namely, nNOS, iNOS and eNOS [22-24]. Our earlier study

showed that while nNOS was readily detected in untreated HT22 cells, iNOS was mostly

undetectable [25]. In this study, we found that treatment of HT22 cells with increasing

concentrations of RSL3 for different durations did not significantly affect total cellular

nNOS protein level (including both monomer and dimer forms) (**Fig. 4A**). By contrast,

while little or no iNOS band was detected in vehicle-treated control HT22 cells as

reported earlier [25], treatment of these cells with RSL3 significantly increased the total

cellular iNOS level in a time- and dose-dependent manner (**Fig. 4A**).

The catalytic activity of NOS is activated through the formation of NOS dimers [22-

24], which are stabilized by the formation of an intermolecular disulfide bond [26]. Our

earlier studies showed that the NOS proteins are activated to form dimers in erastin-

treated cells [12, 27]. Next, we sought to determine whether nNOS and iNOS were

activated (*i.e.*, dimerized) in RSL3-treated HT22 cells. For nNOS, treatment of HT22 cells

with RSL3 significantly increased the formation of nNOS dimers (**Fig. 4B**), but the total

nNOS protein level was not significantly changed (**Fig. 4A**). For iNOS, RSL3 not only

increased iNOS activation (dimerization) in a time- and dose-dependent manner (**Fig.**

4B), but it also markedly increased the total iNOS protein level (**Fig. 4A**).

Next, we determined the modulating effect of S-methylisothiourea (SMT), a

preferential inhibitor of iNOS [28, 29], and chlorpromazine (CPZ), a preferential

inhibitor of nNOS [30], on RSL3-induced cytotoxicity in HT22 cells. We found that joint

treatment of HT22 cells with SMT abrogated RSL3-induced accumulation of cellular NO

(**Fig. 4C, 4E**), ROS (**Fig. 4D, 4F**) and lipid-ROS (**Fig. 4G**), and these effects were

associated with a strong protection (~100% restoration of cell viability) against RSL3-

induced ferroptotic cell death (**Fig. 4H**). In comparison, the effect of CPZ against RSL3-

induced accumulation of NO (**Fig. 4C, 4E**), ROS (**Fig. 4D, 4F**) and lipid-ROS (**Fig. 4G**) was

markedly weaker, which is also associated with a much weaker protection against

RSL3-induced cell death (**Fig. 4I**). The combined use of CPZ and SMT at lower

concentrations which only produces a partial cytoprotection when each chemical was

used alone exerted a stronger protective effect (**Fig. 4J**).

Lastly, we also evaluated in this study the contributing role of nNOS and iNOS in

mediating RSL3-induced ferroptotic cell death by using the siRNA approach to

selectively knock down their expression in HT22 cells. The efficiency of iNOS and nNOS

knockdown was confirmed by measuring the change in cellular nNOS and iNOS protein

levels (**Fig. 5A**). Knockdown of iNOS abrogated RSL3-induced increase in iNOS protein

levels (**Fig. 5A**), and it also exerted a significant protection against RSL3-induced

cytotoxicity (**Fig. 5B**). By contrast, selective knockdown of nNOS in HT22 cells did not

show a significant protective effect (**Fig. 5C**). It is of note that the nNOS knockdown

experiment was repeated multiple times, and similar results were obtained which

showed a lack of significant protective effect against RSL3-induced cytotoxicity.

Together, these results indicate that NO accumulation resulting from the induction

and activation (dimerization) of iNOS and nNOS in RSL3-treated cells is involved in

mediating RSL3-induced ferroptosis in HT22 cells. It is apparent that iNOS plays a more

dominant role than nNOS in this process.

PDI mediates NOS dimerization in RSL3-treated cells

In this study, we also sought to determine whether PDI can catalyze nNOS and iNOS

dimerization in RSL3-treated HT22 cells. First, we found that the cellular level of total

PDI protein was not significantly affected by RSL3 treatment (**Fig. 4A**). To provide

support for the hypothesis that PDI is responsible for catalyzing nNOS and iNOS

dimerization, we determined the effect of cystamine on RSL3-induced ferroptosis.

Cystamine is an organic disulfide molecule capable of inhibiting PDI's enzymatic activity

through covalent modification of the cysteine residue(s) in its catalytic site [31, 32]. For

nNOS, it was observed that treatment of HT22 cells with RSL3 stimulated its activation

(dimerization), and joint treatment with cystamine abrogated RSL3-induced nNOS

dimerization (**Fig. 5D**). For iNOS, RSL3 markedly increased iNOS total protein level and

its dimer formation in HT22 cells (**Fig. 5D**), and joint treatment with cystamine

abolished RSL3-induced increase in total iNOS protein level and dimerization (**Fig. 5D**).

These effects of cystamine on nNOS and iNOS dimerization and on total iNOS protein

level in RSL3-treated cells were associated with a strong abrogation of NO (**Fig. 5E**),

ROS (**Fig. 5F**) and lipid-ROS (**Fig. 5G**) accumulation, as well as a strong protection

against RSL3-induced ferroptotic cell death (**Fig. 5H**).

We recently have shown that 4-hydroxyestrone (4-OH-E$_1$) is an endogenous

estrogen metabolite that can effectively inhibit the enzymatic activity of PDI [33]. We

found that joint treatment of HT22 cells with 4-OH-E$_1$ abrogated RSL3-induced increase

in total iNOS protein levels (**Fig. 6A**). Because the nNOS dimers are relatively easier to

detect than the iNOS dimers, we used the nNOS dimers as a representative parameter to

reflect PDI-mediated NOS dimerization (activation) in RSL3-treated HT22 cells. We

found that joint treatment of HT22 cells with 4-OH-E$_1$ abrogated RSL3-induced nNOS

dimerization (**Fig. 6A**). This effect was associated with decreased accumulation of

cellular NO (**Fig. 6B**), ROS (**Fig. 6C**) and lipid-ROS (**Fig. 6D**), together with increased cell

survival (**Fig. 6E**).

Our earlier study showed that PDI in untreated HT22 cells is mostly present in the

catalytically-inactive *S*-nitrosylated form, and it undergoes *S*-denitrosylation following

GSH depletion [25]. To probe the involvement of PDI *S*-denitrosylation in mediating

RSL3-induced cytotoxicity in HT22 cells, we studied the effect of *S*-nitroso-*N*-

acetylpenicillamine (SNAP), a protein *S*-nitrosylating agent [34]. We found that joint

treatment of HT22 cells with SNAP abrogated RSL3-induced NOS activation (based on

detection of nNOS dimer; **Fig. 6F**), and this effect was associated with reduced

accumulation of NO (**Fig. 6G**), ROS (**Fig. 6H**), lipid-ROS (**Fig. 6I**), and protection against

ferroptotic cell death (**Fig. 6J**). These results indicate that *S*-nitrosylation of PDI is

indeed associated with suppression of PDI activity, reduction in NOS dimerization,

reduction in cellular NO, ROS and lipid-ROS levels, and ultimately, protection from

ferroptosis.

Based on the consistent observations with different PDI inhibitors, next we also

tried to probe PDI's effect by using specific siRNAs to selectively knockdown its

expression. The effectiveness of PDI knockdown was assessed by Western blotting of its

cellular protein levels (**Fig. 6K**). As expected, PDI knockdown elicited a partial

protection against RSL3-induced cytotoxicity (**Fig. 6L**), but the degree of protection in

PDI-knockdown cells is smaller than expected, which likely is due to the fact that a

transient PDI knockdown may not last long enough for the cell viability experiment.

Collectively, these results still clearly show that PDI is a key upstream mediator of

RSL3-induced ferroptosis by catalyzing iNOS dimerization in RSL3-treated HT22 cells,

which then leads to accumulation of cellular NO and ROS/lipid-ROS and ultimately,

oxidative cell death.

Role of TrxR1 in RSL3-induced ferroptosis

A recent study reported that RSL3 can inhibit the catalytic activity of TrxR1 [5].

Since PDI belongs to the thioredoxin superfamily and a substrate of TrxR1 [35, 36], next

we sought to determine the potential role of TrxR1 in RSL3-induced ferroptotic cell

death. First, we used the TrxR1-siRNAs to transiently knock down its expression in

HT22 cells. The effectiveness of TrxR1 knockdown was confirmed by Western blot

analysis of TrxR1 protein levels (**Fig. 7A**). While TrxR1 knockdown alone (in the

absence of RSL3) reduced the baseline cell viability compared to cells transfected with

the control siRNAs alone (**Fig. S2A**), TrxR1 knockdown markedly reduced the

sensitivity of cells to the cytotoxicity induced by RSL3 and auranofin (a prototypical

inhibitor of TrxR1 [5, 37, 38]) (**Fig. 7B**).

We also determined whether the TrxR1 activity in HT22 cells was inhibited by

treatment of cells with RSL3 or auranofin. The HT22 cells were first treated with 0.1 μM

RSL3 or 1 μM auranofin for 5 h, then the cells were washed and harvested for

preparation of cell lysates. Even after all these preparative steps, the TrxR1 activity

contained in the cell lysates was still markedly reduced in RSL3- or auranofin-treated

cells (**Fig. 7C**); in comparison, the TrxR1 protein levels remained largely unchanged or

slightly increased in these cells (**Fig. 7D, 7E, 7G, 7H**). Jointly, these observations suggest

that RSL3 and auranofin can bind to TrxR1 very tightly (most likely covalently) to exert

a strong and sustained inhibition of the enzyme activity.

Next, we determined the iNOS dimer level in RSL3-treated HT22 cells with prior

TrxR1 knockdown. As shown in **Fig. 7F**, TrxR1 knockdown alone (without RSL3

treatment) caused a significant increase in iNOS dimer level, but it significantly

abrogated RSL3-induced increase in iNOS dimer formation. Notably, the cellular PDI

level appeared to be also reduced in TrxR1-knockdown cells when it was treated with

RSL3 (**Fig. 7F**). These results suggest that TrxR1 plays an important role in meditating

PDI-catalyzed iNOS dimerization and subsequent ferroptotic cell death in HT22 cells.

It is of note that in this study, we also probed the effect of jointly knocking down PDI

and TrxR1 in HT22 cells and then determined its effect on RSL3-induced cell death. We

found that joint knockdown of PDI and TrxR1 reduced the sensitivity to RSL3

cytotoxicity in a similar manner as what was seen with PDI knockdown alone (**Fig. S2B**).

This result is expected because PDI is a substrate of TrxR1, and when PDI is mostly

absent (resulting from its knockdown), there is not much effect left to be seen even by

further knocking down TrxR1.

Next, we determined the ability of RSL3 and auranofin to directly inhibit the

enzymatic activity of TrxR1 in biochemical assays. The whole cell lysates were prepared

from untreated HT22 cells, and then the lysates (which contain TrxR1 enzyme) were

used to assay the TrxR1 enzyme activity in the presence or absence of RSL3 or

auranofin. We found that the TrxR1 enzyme activity could be very potently inhibited in

a concentration-dependent manner by the presence of auranofin, with an apparent IC_{50}

of approximately 1.1 nM (**Fig. 7I**). It is of note that the TrxR1 enzyme inhibition by

auranofin became stronger but with prolonged incubation time until it reached a

plateau (**Fig. 7J**). This pattern of inhibition is consistent with the earlier observation

that auranofin can covalently inhibit TrxR1 [39]. Similarly, the TrxR1 enzyme activity

was also dose-dependently inhibited by RSL3, with an apparent IC_{50} of approximately 4

μM (**Fig. 7K**), which is in agreement with the results from a recent study when the

purified TrxR1 enzyme was used in the *in vitro* assay [5]. It is of note that the overall

pattern of TrxR1 inhibition by RSL3 is similar to the inhibition by auranofin, *i.e.*, a

stronger inhibition was observed with prolonged reaction time, until it reached a

plateau (**Fig. 7L**).

Next, we determined whether auranofin can induce cytotoxicity in HT22 cells.

Treatment of HT22 cells with auranofin alone elicited a concentration-dependent

cytotoxicity (**Fig. 8A**). The IC_{50} of auranofin's cytotoxicity is 0.8 μM, which is much

higher than its IC_{50} for TrxR1 inhibition. In addition, auranofin-induced cell death was

not protected by Fer-1 (**Fig. 8B**) or the iron chelator DFO (**Fig. 8C**), two commonly-used

protective agents for chemically-induced ferroptosis. Similarly, while auranofin (at the 1

μM concentration) elicited accumulation of cellular NO (**Fig. 8D**) and ROS (**Fig. 8E**), it

did not elicit a significant accumulation of lipid-ROS (**Fig. 8F**). Together, these

observations suggest that the cell death induced by auranofin at high concentrations

(much higher than its IC_{50} for TrxR1 inhibition) is not ferroptosis. In addition, the

apoptosis inhibitor z-VAD-FMK and necrosis inhibitor Nec-1 also failed to prevent

auranofin-induced cell death (**Fig. 8G, 8H**), indicating that auranofin-induced

cytotoxicity in HT22 cells is also not typical apoptosis or necrosis.

Based on the above observations, we also investigated the effect of jointly treating

HT22 cells with auranofin + BSO, which is a GSH synthase inhibitor [40, 41]. It is

expected that BSO will facilitate the induction of GSH depletion-associated ferroptosis

when they were jointly treated with auranofin. In this experimental design, BSO was

used at a concentration (such as 6 μM) that did not cause significant cytotoxicity when

present alone (**Fig. 9A**). We found that joint treatment of HT22 cells with very low

concentrations of auranofin + 6 μM BSO caused a concentration-dependent reduction in

cell viability (**Fig. 9B**). Interestingly, the GPX4 protein level was also decreased in the

cells jointly treated with auranofin + BSO (**Fig. 9C**). Notably, a similar decrease in GPX4

protein level was observed in cells treated with RSL3 alone (**Fig. 7D**). The cytotoxicity of

auranofin + BSO was partially rescued by Fer-1 (**Fig. 9D**). Cystamine, an inhibitor of PDI

which can effectively prevent RSL3-induced ferroptosis (**Fig. 5H**), also exerted a partial

protection against the cell death jointly induced by auranofin and BSO (**Fig. 9E**).

Similarly, 4-OH-E$_1$, which can effectively rescue RSL3-induced cell death (**Fig. 6E**), also

exerted a partial protection against the cell death jointly induced by auranofin and BSO

(**Fig. 9E**). All these observations suggest that when very low concentrations of auranofin

(which are close to its IC_{50} for TrxR1 inhibition) were used under conditions where

cellular GSH is depleted, the cell death induced by auranofin is largely ferroptosis, and

as expected, the cell death can be partially prevented by both ferroptosis inhibitors and

PDI inhibitors.

To provide further support for the above observations, we also analyzed changes in

cellular NO, ROS and lipid-ROS accumulation and NOS dimerization following treatment

of cells with very low concentrations of auranofin + BSO. We found that under these

joint treatment conditions, there was a time-dependent accumulation of cellular NO (**Fig.**

9F), ROS (**Fig. 9G**) and lipid-ROS (**Fig. 9H**). In addition, the dimeric nNOS (detected as a

marker for NOS dimerization) was also significantly increased in a time-dependent

manner (**Fig. 9C**). Joint treatment of these cells with cystamine partially abrogated the

accumulation of NO (**Fig. 9I**), ROS (**Fig. 9J**) and lipid-ROS (**Fig. 9K**) induced by

treatment of auranofin + BSO. Similar observations on the abrogation of NO, ROS and

lipid-ROS accumulation were also made when the HT22 cells were jointly treated with

4-OH-E$_1$ ± auranofin and BSO (**Fig. 9I**$\square$**9K**).

DISCUSSION

RSL3 is a prototypical ferroptosis inducer, and its ferroptosis-inducing ability is thought to be associated with inhibition of GPX4 [4, 42, 43]. However, a recent study showed that in addition to GPX4 inhibition, RSL3 can also inhibit the enzymatic activity of TrxR1 [5], which would shift the pool of thioredoxins to their oxidized forms. PDI is a member of thioredoxin superfamily, and we hypothesize that RSL3 might, through its inhibition of TrxR1's enzymatic activity [5], keep more cellular PDI proteins in the oxidized form, thereby facilitating RSL3-induced ferroptosis through catalyzing NOS dimerization, followed by cellular ROS and lipid-ROS accumulation, and ultimately ferroptotic cell death (summarized in **Fig. 10**). As discussed below, large amount of experimental data is provided in this study which jointly offers strong support for this hypothesis.

We find that during the development of RSL3-induced ferroptosis in HT22 cells, NO accumulation is an early event which subsequently leads to the formation and accumulation of ROS and lipid-ROS, contributing to the development of ferroptotic cell death. The supporting evidence includes: First, we find that RSL3 can induce the accumulation of both NO and ROS/lipid-ROS in a time- and dose-dependent manner, although there appears to be no clear time differential in their buildup (the reason for this is provided later). The observations made with SNP, which can release NO [21], offer support for this suggestion. We find that SNP can quickly lead to accumulation of cellular ROS and lipid-ROS when present alone in HT22 cells, and it also dramatically sensitizes the cells to RSL3-induced ROS/lipid-ROS accumulation and ferroptosis. Here it is of note that our earlier study showed that during erastin-induced ferroptosis, NO accumulation occurred earlier than did ROS/lipid-ROS accumulation in HT22 cells [12].

In addition, when the HT22 cells were jointly treated with erastin + SNP, the NO

accumulation also occurred much earlier (due to the rapid release of NO from SNP), and

in parallel manner, the ROS/lipid-ROS accumulation also occurred much earlier, lagging

shortly behind NO accumulation [12]. These experimental observations jointly suggest

that during chemically-induced ferroptosis, NO accumulation occurs first, which then

leads to ROS/lipid-ROS accumulation. The second line of evidence comes from the study

of NO scavengers and NOS inhibitors. We find that cPTIO, a NO scavenger, can abrogate

RSL3-induced NO accumulation, and this effect is accompanied by abrogation of

ROS/lipid-ROS accumulation and ferroptosis. In addition, SMT, an iNOS inhibitor, exerts

a strong protection against RSL3-induced NO accumulation, which is accompanied by

abrogation of ROS/lipid-ROS accumulation and ferroptotic cell death. Together, these

observations suggest that RSL3-induced NO accumulation contributes to the

subsequent accumulation of ROS and lipid-ROS, and ultimately the induction of

ferroptotic cell death.

Although RSL3-induced NO accumulation leads to cellular ROS/lipid-ROS

accumulation in HT22 cells, it is interesting that RSL3-induced buildup of cellular NO

and cellular ROS (which includes lipid-ROS) do not display a distinguishable time

difference; in fact, they occur nearly at the same time. This unexpected experimental

observation likely is attributable to the fact that RSL3 can increase both NO and lipid-

ROS accumulation at the same time. RSL3 increases NO biosynthesis through its direct

inhibition of TrxR1, which then leads to PDI activation and NOS dimerization; on the

other hand, RSL3 also directly increases cellular lipid-ROS accumulation (independent

of cellular NO accumulation) due to its inhibition of GPX4. Therefore, the buildup of

cellular lipid-ROS (which is part of the cellular ROS) in RSL3-treated cells is the result of

both GPX4 inhibition (which likely occurs first) and NO accumulation (which likely

occurs at a slightly later time). Because of these reasons, the accumulation of cellular NO

and ROS/lipid-ROS occurs rapidly and almost at the same time after exposure to RSL3.

The dimeric NOS is catalytically active and catalyzes NO formation [12-16]. In this

study, we show that RSL3 exposure increases the formation of nNOS and iNOS dimers in

a time- and dose-dependent manner in HT22 cells. The results from iNOS and nNOS

inhibitor experiments support the idea that both iNOS and nNOS are involved in RSL3-

induced ferroptosis, but the results with iNOS and nNOS knockdown experiments are

inclusive. It is shown that iNOS knockdown is associated with a partial abrogation of

RSL3's cytotoxicity, but nNOS knockdown fails to exert a significant protection (**Fig. 5**).

Two potential reasons may underlie the inability to observe a significant protective

effect with nNOS knockdown: First, it is likely that the relative contribution of cellular

nNOS (which is not induced following RSL3 exposure) to the cellular NO pool is

markedly smaller than the contribution of cellular iNOS (which is strongly-induced by

RSL3). The other reason might be due to the low effectiveness of a sustained nNOS

protein knockdown in live HT22 cells. Together, it is apparent that iNOS clearly plays a

more important role than nNOS in mediating RSL3-induced NO accumulation and

ferroptosis in this neuronal cell culture model.

By jointly using chemical inhibitors or PDI knockdown, strong evidence is provided

in this study to show that PDI is involved in catalyzing NOS dimer formation in RSL3-

treated HT22 cells. For instance, joint treatment of HT22 cells with cystamine (a PDI

inhibitor) strongly reduces RSL3-induced dimerization of both nNOS and iNOS, the

accumulation of NO, and ROS/lipid-ROS, and the oxidative cytotoxicity. Another

example involves the *S*-nitrosylation of the cysteine residues in PDI's active site, which

inactivates PDI's enzyme activity [44, 45]. Our earlier study showed that PDI in

untreated HT22 cells was mostly present in the *S*-nitrosylated form, but following

glutamate treatment (which induces cellular GSH depletion), PDI underwent *S*-

denitrosylation [25]. We find in this study that RSL3-induced NO, ROS and lipid-ROS

accumulation is abrogated by SNAP (which causes PDI *S*-nitrosylation), and this effect

associated with a protection against cell death. Together, these observations offer

support for the notion that PDI plays an important role in catalyzing NOS dimerization

in RSL3-treated HT22 cells, which then leads to accumulation of NO, ROS and lipid-ROS,

and ultimately ferroptotic cell death.

ERO1 is an enzyme that oxidizes PDI by forming disulfide bonds in its active site

[46]. Our recent work showed that EN460 (an inhibitor of ERO1) has a strong protective

effect against erastin-induced ferroptosis in HT22 cells [12]. However, EN460 only

shows a very weak protective effect against RSL3-induced ferroptosis as observed in

this study (**Fig. S3A–S3D**). This difference is because RSL3 is an inhibitor of TrxR1,

which catalyzes the reduction of the disulfide bonds in PDI's active site. When the

cellular TrxR1 activity is inhibited by RSL3, the majority of PDI will be in the oxidized

form; as such, the presence of ERO1 will not add significantly more oxidized PDI

(because not much reduced PDI is left as substrate for ERO1). This is why it will be

difficult to display a strong protective effect of EN460 when TrxR1 is already inhibited

by RSL3. This lack of a significant protective effect of EN460, in fact, also supports the

important role of PDI in driving RSL3-mediated ferroptosis.

Based on the findings of this study, it is evident that RSL3-induced ferroptosis

involves two important mechanistic components: one is mediated by the activation of

PDI (resulting from TrxR1 inhibition by RSL3), and other component is the inhibition of

the GPX4-mediated protection against lipid-ROS. Because of the combination of these

two effects, a PDI inhibitor which does not have a direct antioxidant activity is expected

to be less efficacious in protecting against RSL3-induced ferroptosis compared to a PDI

inhibitor which also has a strong antioxidant activity. For instance, 4-OH-E$_1$, which was

recently identified as an effective PDI inhibitor [33], is found to have a strong protective

effect against RSL3-induced ferroptotic cell death. The protective effect of 4-OH-E$_1$

against RSL3-induced cell death is more potent than its protection against erastin-

induced cell death in some of the cell lines (unpublished data). This observation

suggests that the direct antioxidant activity of 4-OH-E$_1$ against lipid-ROS also

contributes, to some degrees, to its cytoprotective effect. This explanation is also in line

with the observation with cystamine, which can inhibit PDI function but does not have

any antioxidant property. Cystamine exerts a less efficacious protection against RSL3-

induced ferroptosis in HT22 cells compared to 4-OH-E$_1$; however, both cystamine and 4-

OH-E$_1$ exert a similarly strong protection (90□100% protection) against erastin-

induced ferroptosis in the same cells (data not shown).

In this study, we find that the TrxR1's enzyme activity can be directly and potently

inhibited by auranofin (with an apparent IC_{50} of approximately 1.1 nM), which is slightly

lower than the reported IC_{50} (~20 nM) for human TrxR1 [47]. Notably, it is observed in

this study that when the *in vitro* reaction time is prolonged, the reduction in product

formation becomes progressively greater, and there is a sustained reduction in product

formation (reflecting a sustained TrxR1 inhibition). This observation suggests that

auranofin inhibits TrxR1's catalytic activity resulting from its covalent modification of

the enzyme. Indeed, earlier studies have reported that auranofin can inactivate TrxR1

by forming covalent diselenide bridges with its Sec498 residue [39, 48].

We find that the potency of RSL3 to inhibit TrxR1 *in vitro* is markedly lower than

that of auranofin. RSL3's apparent IC_{50} (approximately 4 μM) detected in this study is in

line with the results of an earlier study [5]. Based on the similar patterns of TrxR1

inhibition by RSL3 and auranofin observed in this study, it is suggested that RSL3 may

also inhibit TrxR1 in a covalent manner. This suggestion is partially supported by the

observation made in HT22 cells showing a high-potency inhibition of the TrxR1 activity

by RSL3. It should be noted that the apparent discrepancy between the high-potency

RSL3 inhibition in intact cells and the low-potency inhibition of the TrxR1's catalytic

activity in an *in vitro* enzymatic assay may suggest that RSL3 cannot readily access the

catalytic site of TrxR1 in an aqueous reaction system *in vitro*. A similar phenomenon

was also observed in an earlier study with the flavonoid-type compounds: while these

compounds can strongly activate the cyclooxygenase's catalytic activity in live cells in

culture when present at low nM concentrations, they needed to be present at 100-200

μM concentrations in order to exert a similar activation of the pure enzymes in the *in*

vitro enzyme assays [49].

In conclusion, the results of our present study demonstrate that PDI plays a critical

role in mediating RSL3-induced ferroptotic cell death in HT22 cells by catalyzing the

dimerization of both nNOS and iNOS, followed by cellular accumulation of NO, ROS and

lipid-ROS, and ultimately ferroptotic cell death (summarized in **Fig. 10**). The discovery

of an important role of PDI in mediating chemically-induced ferroptosis suggests that

PDI might serve as a potential drug target for protection against ferroptosis-related

neuronal degeneration.

Data Availability Statement: The authors confirm that the data supporting the

findings of this study are available within the article and its supplementary materials.

Funding: This work was supported, in part, by research grants from Shenzhen Peacock

Plan (No. KQTD2016053117035204), the National Natural Science Foundation of China

(No. 81630096), Shenzhen Key Laboratory of Steroidal Drug Research (No.

ZDSYS20190902093417963) and Shenzhen Bay Laboratory (No. SZBL2019062801007).

REFERENCES

1. Dixon, S.J., et al., *Ferroptosis: an iron-dependent form of nonapoptotic cell death.* Cell, 2012. **149**(5): p. 1060-72.

2. Stockwell, B.R., *Ferroptosis turns 10: Emerging mechanisms, physiological functions, and therapeutic applications.* Cell, 2022. **185**(14): p. 2401-2421.

3. Yang, W.S. and B.R. Stockwell, *Synthetic lethal screening identifies compounds activating iron-dependent, nonapoptotic cell death in oncogenic-RAS-harboring cancer cells.* Chem Biol, 2008. **15**(3): p. 234-45.

4. Yang, W.S., et al., *Regulation of ferroptotic cancer cell death by GPX4.* Cell, 2014. **156**(1-2): p. 317-331.

5. Cheff, D.M., et al., *The ferroptosis inducing compounds RSL3 and ML162 are not direct inhibitors of GPX4 but of TXNRD1.* Redox Biol, 2023. **62**: p. 102703.

6. Mustacich, D. and G. Powis, *Thioredoxin reductase.* Biochem J, 2000. **346 Pt 1**(Pt 1): p. 1-8.

7. Tasanen, K., et al., *Characterization of the human gene for a polypeptide that acts both as the beta subunit of prolyl 4-hydroxylase and as protein disulfide isomerase.* J Biol Chem, 1988. **263**(31): p. 16218-24.

8. Pihlajaniemi, T., et al., *Molecular cloning of the beta-subunit of human prolyl 4-hydroxylase. This subunit and protein disulphide isomerase are products of the same gene.* EMBO J, 1987. **6**(3): p. 643-9.

9. Freedman, R.B., T.R. Hirst, and M.F. Tuite, *Protein disulphide isomerase: building bridges in protein folding.* Trends Biochem Sci, 1994. **19**(8): p. 331-6.

10. Turano, C., et al., *Proteins of the PDI family: unpredicted non-ER locations and functions.* J Cell Physiol, 2002. **193**(2): p. 154-63.

11. Cai, H., C.C. Wang, and C.L. Tsou, *Chaperone-like activity of protein disulfide isomerase in the refolding of a protein with no disulfide bonds.* J Biol Chem, 1994. **269**(40): p. 24550-2.

12. Hou, M.J., P. Wang, and B.T. Zhu, *Biochemical mechanism of erastin-induced ferroptotic cell death in neuronal cells.* Acta Biochim Biophys Sin (Shanghai), 2023. **55**(5): p. 853-865.

13. Liu, P., et al., *Ferrostatin-1 alleviates lipopolysaccharide-induced acute lung injury via inhibiting ferroptosis.* Cell Mol Biol Lett, 2020. **25**: p. 10.

14. Miotto, G., et al., *Insight into the mechanism of ferroptosis inhibition by ferrostatin-1.* Redox Biol, 2020. **28**: p. 101328.

15. Peng, W., et al., *L-F001, a Multifunctional Fasudil-Lipoic Acid Dimer Prevents RSL3-Induced Ferroptosis via Maintaining Iron Homeostasis and Inhibiting JNK in HT22 Cells.* Front Cell Neurosci, 2022. **16**: p. 774297.

16. Kato, K., et al., *Quercetin and resveratrol inhibit ferroptosis independently of Nrf2-ARE activation in mouse hippocampal HT22 cells.* Food Chem Toxicol, 2023. **172**: p. 113586.

17. Ayala, A., M.F. Munoz, and S. Arguelles, *Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal.* Oxid Med Cell Longev, 2014. **2014**: p. 360438.

18. Maeda, H., et al., *Multiple functions of nitric oxide in pathophysiology and microbiology: analysis by a new nitric oxide scavenger.* J Leukoc Biol, 1994. **56**(5): p. 588-92.

19. Castro, I.A., et al., *Free radical scavenger and antioxidant capacity correlation of alpha-tocopherol and Trolox measured by three in vitro methodologies.* Int J Food Sci Nutr, 2006. **57**(1-2): p. 75-82.

20. Aldini, G., et al., *N-Acetylcysteine as an antioxidant and disulphide breaking agent: the reasons why.* Free Radic Res, 2018. **52**(7): p. 751-762.

21. Zoupa, E. and N. Pitsikas, *The Nitric Oxide (NO) Donor Sodium Nitroprusside (SNP) and Its Potential for the Schizophrenia Therapy: Lights and Shadows.* Molecules, 2021. **26**(11).

22. Andrew, P.J. and B. Mayer, *Enzymatic function of nitric oxide synthases.* Cardiovasc Res, 1999. **43**(3): p. 521-31.

23. Forstermann, U. and W.C. Sessa, *Nitric oxide synthases: regulation and function.* Eur Heart J, 2012. **33**(7): p. 829-37, 837a-837d.

24. Alderton, W.K., C.E. Cooper, and R.G. Knowles, *Nitric oxide synthases: structure, function and inhibition.* Biochem J, 2001. **357**(Pt 3): p. 593-615.

25. Okada, K., M. Fukui, and B.T. Zhu, *Protein disulfide isomerase mediates glutathione depletion-induced cytotoxicity.* Biochem Biophys Res Commun, 2016. **477**(3): p. 495-502.

26. Crane, B.R., et al., *Structure of nitric oxide synthase oxygenase dimer with pterin and substrate.* Science, 1998. **279**(5359): p. 2121-6.

27. Wang, H., P. Wang, and B.T. Zhu, *Mechanism of Erastin-Induced Ferroptosis in MDA-MB-231 Human Breast Cancer Cells: Evidence for a Critical Role of Protein Disulfide Isomerase.* Mol Cell Biol, 2022. **42**(6): p. e0052221.

28. Narayanan, K. and O.W. Griffith, *Synthesis of L-thiocitrulline, L-homothiocitrulline, and S-methyl-L-thiocitrulline: a new class of potent nitric oxide synthase inhibitors.* J Med Chem, 1994. **37**(7): p. 885-7.

29. Balaganur, V., et al., *Effect of S-methylisothiourea, an inducible nitric oxide synthase inhibitor, in joint pain and pathology in surgically induced model of osteoarthritis.* Connect Tissue Res, 2014. **55**(5-6): p. 367-77.

30. Palacios, M., et al., *Chlorpromazine inhibits both the constitutive nitric oxide synthase and the induction of nitric oxide synthase after LPS challenge.* Biochem Biophys Res Commun, 1993. **196**(1): p. 280-6.

31. Fujita, I., et al., *Cystamine-mediated inhibition of protein disulfide isomerase triggers aggregation of misfolded orexin-A in the Golgi apparatus and prevents extracellular secretion of orexin-A.* Biochem Biophys Res Commun, 2017. **489**(2): p. 164-170.

32. Hoffstrom, B.G., et al., *Inhibitors of protein disulfide isomerase suppress apoptosis induced by misfolded proteins.* Nat Chem Biol, 2010. **6**(12): p. 900-6.

33. Wang, H., et al., *Strong Protection by 4-Hydroxyestrone against Erastin-Induced Ferroptotic Cell Death in Estrogen Receptor-Negative Human Breast Cancer Cells: Evidence for Protein Disulfide Isomerase as a Mechanistic Target for Protection.* Biochemistry, 2024. **63**(8): p. 984-999.

34. Mallis, R.J. and J.A. Thomas, *Effect of S-nitrosothiols on cellular glutathione and reactive protein sulfhydryls.* Arch Biochem Biophys, 2000. **383**(1): p. 60-9.

35. Ali Khan, H. and B. Mutus, *Protein disulfide isomerase a multifunctional protein with multiple physiological roles.* Front Chem, 2014. **2**: p. 70.

36. Lundstrom, J. and A. Holmgren, *Protein disulfide-isomerase is a substrate for thioredoxin reductase and has thioredoxin-like activity.* J Biol Chem, 1990. **265**(16): p. 9114-20.

37. Hwang-Bo, H., et al., *Auranofin, an inhibitor of thioredoxin reductase, induces apoptosis in hepatocellular carcinoma Hep3B cells by generation of reactive oxygen species.* Gen Physiol Biophys, 2017. **36**(2): p. 117-128.

38. Zhang, X., et al., *Repurposing of auranofin: Thioredoxin reductase remains a primary target of the drug.* Biochimie, 2019. **162**: p. 46-54.

39. Becker, K., et al., *Thioredoxin reductase as a pathophysiological factor and drug target.* Eur J Biochem, 2000. **267**(20): p. 6118-25.

40. Drew, R. and J.O. Miners, *The effects of buthionine sulphoximine (BSO) on glutathione depletion and xenobiotic biotransformation.* Biochem Pharmacol, 1984. **33**(19): p. 2989-94.

41. Tagde, A., et al., *The glutathione synthesis inhibitor buthionine sulfoximine synergistically enhanced melphalan activity against preclinical models of multiple myeloma.* Blood Cancer J, 2014. **4**(7): p. e229.

42. Yang, W.S. and B.R. Stockwell, *Ferroptosis: Death by Lipid Peroxidation.* Trends Cell Biol, 2016. **26**(3): p. 165-176.

43. Friedmann Angeli, J.P., et al., *Inactivation of the ferroptosis regulator Gpx4 triggers acute renal failure in mice.* Nat Cell Biol, 2014. **16**(12): p. 1180-91.

44. Uehara, T., et al., *S-nitrosylated protein-disulphide isomerase links protein misfolding to neurodegeneration.* Nature, 2006. **441**(7092): p. 513-7.

45. Kaplan, A., et al., *Small molecule-induced oxidation of protein disulfide isomerase is neuroprotective.* Proc Natl Acad Sci U S A, 2015. **112**(17): p. E2245-52.

46. Araki, K. and K. Nagata, *Functional in vitro analysis of the ERO1 protein and protein-disulfide isomerase pathway.* J Biol Chem, 2011. **286**(37): p. 32705-12.

47. Gromer, S., et al., *Human placenta thioredoxin reductase. Isolation of the selenoenzyme, steady state kinetics, and inhibition by therapeutic gold compounds.* J Biol Chem, 1998. **273**(32): p. 20096-101.

48. Fang, J. and A. Holmgren, *Inhibition of thioredoxin and thioredoxin reductase by 4-hydroxy-2-nonenal in vitro and in vivo.* J Am Chem Soc, 2006. **128**(6): p. 1879-85.

49. Bai, H.W. and B.T. Zhu, *Strong activation of cyclooxygenase I and II catalytic activity by dietary bioflavonoids.* J Lipid Res, 2008. **49**(12): p. 2557-70.

FIGURE LEGENDS

Figure 1. Time- and dose-dependent accumulation of NO, ROS and lipid-ROS in
RSL3-induced ferroptosis in HT22 cells.

A, B. Time- and dose-dependent effect of RSL3 on cell viability (MTT assay, n = 5). The
RSL3 concentration used is 100 nM (**A**) and the treatment duration is 24 h (**B**).

C. Time-dependent change in the gross morphology in RSL3 (100 nM)-treated cells
(scale bar = 45 μm).

D. Protective effect of Fer-1 against RSL3-induced cytotoxicity following treatment of
cells with RSL3 (100 nM) ± Fer-1 (1 μM) for 24 h (MTT assay, n = 5).

E. Time-dependent accumulation of cellular NO and ROS (fluorescence microscopy
images, scale bar = 60 μm) following RSL3 treatment, and the quantitative intensity
values are shown in the right panel (n = 3).

F. Time-dependent accumulation of cellular lipid-ROS (confocal microscopy images) in
RSL3-treated cells (scale bar = 10 μm).

G, H, I. Time-dependent accumulation of cellular NO (**G**), ROS (**H**) and lipid-ROS (**I**) in
RSL3-treated cells (analytical flow cytometry). The quantitative intensity values are
shown in the right panel (n = 3).

J. Time-dependent change in cellular MDA levels in RSL3-treated cells (n = 3).

Each value represents the mean ± S.D. (* P < 0.05; ** or $^{\#\#}P$ < 0.01; n.s. not significant).

Figure 2. Effect of free radical scavengers on RSL3-induced cytotoxicity and
cellular accumulation of NO, ROS and lipid-ROS in HT22 cells.

A, B, C. Abrogation by cPTIO of RSL3-induced accumulation of cellular NO (**A**), ROS (**B**)
and lipid-ROS (**C**). Cells were treated with RSL3 (100 nM) ± cPTIO (200 μM) for 6 h, and
then stained for NO, ROS and lipid-ROS (**A, B** for fluorescence microscopy; scale bar = 60
μm; **C** for analytical flow cytometry). The quantitative intensity values are shown in the
lower panel (n = 3; the mean value for the control group is labeled next to the bar).

D. Protective effect of cPTIO against RSL3-induced cytotoxicity following treatment of
cells with RSL3 (100 nM) ± cPTIO (200 μM) for 24 h (MTT assay, n = 5).

E, F, G. Abrogation by Trolox of RSL3-induced accumulation of cellular NO (**E**), ROS (**F**)
and lipid-ROS (**G**). Cells were treated with RSL3 (100 nM) ± Trolox (100 μM) for 6 h, and
then stained for NO, ROS and lipid-ROS (**E, F** for fluorescence microscopy; scale bar = 60
μm; **G** for analytical flow cytometry). The quantitative intensity values are shown in the
lower panel (n = 3; the mean value for the control group is labeled next to the bar).

H. Protective effect of Trolox against RSL3-induced cytotoxicity following treatment of
cells with RSL3 (100 nM) ± Trolox (100 μM) for 24 h (MTT assay, n = 5).

I, J, K, L. Abrogation by NAC of RSL3-induced accumulation of cellular NO (**I, J**), ROS (**K**)
and lipid-ROS (**L**). Cells were treated with RSL3 (100 nM) ± NAC (800 μM) for 6 h, and
then stained for NO, ROS and lipid-ROS (**J, K** for fluorescence microscopy; scale bar = 60
μm; **I, L** for analytical flow cytometry). The quantitative intensity values are shown in
the lower panel (n = 3; the mean value for the control group is labeled next to the bar).

M. Protective effect of NAC against RSL3-induced cytotoxicity following treatment of cells with RSL3 (100 nM) ± NAC (200, 400 and 800 μM) for 24 h (MTT assay, n = 5).

Each value represents the mean ± S.D. (** or $^{##}$ P < 0.01).

Figure 3. Effect of SNP on RSL3-induced NO, ROS and lipid-ROS accumulation in HT22 cells.

A, B. Levels of NO and ROS after treatment with SNP (200 μM) for different durations (0.5, 1, 2, 4 and 6 h; scale bar = 50 μm) (**A**). The quantitative intensity values are shown in the right panel (n = 3) (**B**).

C, D, E. Time-dependent accumulation of cellular NO (**C**), ROS (**D**) and lipid-ROS (**E**) following SNP treatment (analytical flow cytometry) and the quantitative intensity values are in the right panel (n = 3).

F. Dose-dependent effect of SNP on cell viability (MTT assay, n = 5).

G. Cell viability change following treatment with SNP (200 μM) and different concentrations of RSL3 for 24 h (MTT assay, n = 5).

H, I, J, K, L. Accumulation of ROS (**H, I**) and lipid-ROS (**J, K, L**) after treatment with RSL3 (100 nM) ± SNP (200 μM) (**H** for fluorescence microscopy; scale bar = 60 μm; **J, K, L** for analytical flow cytometry). The quantitative intensity values for **H** are shown in the lower panel (**I**) (n = 3).

Each value represents the mean ± S.D. (* or $^{#}$ P < 0.05; ** P < 0.01; n.s., not significant).

Figure 4. RSL3-induced NOS activation in HT22 cells.

A. Levels of total iNOS and nNOS proteins following treatment of cells with RSL3 (100 nM) for the indicated time intervals, or with increasing concentrations of RSL3 for 6 h.

B. Levels of the dimer and monomer iNOS and nNOS proteins following treatment of cells with RSL3 (100 nM) for the indicated time intervals, or with increasing concentrations of RSL3 for 6 h.

C, D, E, F, G. Abrogation by SMT and CPZ of RSL3-induced accumulation of cellular NO (**C, E**), ROS (**D, F**) and lipid-ROS (**G**). Cells were treated with RSL3 (100 nM) ± SMT (50 μM) or RSL3 (100 nM) ± CPZ (40 μM) for 6 h, and then stained for NO, ROS and lipid-ROS (**C, D** for fluorescence microscopy; scale bar = 60 μm; **E, F, G** for analytical flow cytometry). The quantitative intensity values are shown on the right panel (n = 3; the mean value for the control group is labeled next to the bar).

H. Protective effect of SMT against RSL3-induced cytotoxicity following treatment of cells with RSL3 (100 nM) ± SMT (25, 50, 100 μM) for 24 h (MTT assay, n = 4).

I. Protective effect of SMT against RSL3-induced cytotoxicity following treatment of cells with RSL3 (100 nM) ± CPZ (40 μM) for 24 h (MTT assay, n = 5).

J. Protective effect of SMT + CPZ against RSL3-induced cytotoxicity following treatment of cells with RSL3 (100 nM) ± SMT (25 μM) and CPZ (12.5 μM) for 24 h (MTT assay, n = 4).

Each value represents the mean ± S.D. (** or $^{##}$ P < 0.01).

Figure 5. Protective effect of cystamine (a PDI inhibitor) against RSL3-induced ferroptosis in HT22 cells.

A. Effectiveness of iNOS-siRNAs and nNOS-siRNAs in reducing cellular iNOS and nNOS protein levels, respectively. Cells were transfected with iNOS-siRNAs or nNOS-siRNAs for 48 h prior to treatment of cells with RSL3 (50 and 100 nM) for an additional 6 h.

B, C. Effect of iNOS (**B**) and nNOS (**C**) knockdown on RSL3-induced cytotoxicity. Cells were transfected with iNOS-siRNAs or nNOS-siRNAs for 24 h prior to treatment of cells with RSL3 (100 nM) for an additional 24 h.

D. Levels of the dimer and monomer iNOS and nNOS proteins, and the levels of the total iNOS and nNOS proteins after treatment of cells with RSL3 (100 nM) ± cystamine (100 nM) for 6 h.

E, F, G. Abrogation by cystamine of RSL3-induced accumulation of cellular NO (**E**), ROS (**F**) and lipid-ROS (**G**). Cells were treated with RSL3 (100 nM) ± cystamine (100 μM) for 6 h, and then stained for NO, ROS and lipid-ROS (**E, F** for fluorescence microscopy; scale bar = 60 μm; **G** for analytical flow cytometry). The quantitative intensity values are shown in the lower panel (n = 3; the mean value for the control group is labeled next to the bar).

H. Protective effect of cystamine against RSL3-induced cytotoxicity following treatment of cells with RSL3 (100 nM) ± cystamine (100 μM) for 24 h (MTT assay, n = 4).

Each value represents the mean ± S.D. (** or ## P < 0.01; n.s. not significant).

Figure 6. 4-OH-E$_1$ and SNAP (a *S*-nitrosylating agent) protect against RSL3-induced ferroptosis in HT22 cells.

A. Levels of the dimer and monomer nNOS proteins, and the levels of total iNOS and nNOS proteins after treatment of cells with RSL3 (100 nM) ± 4-OH-E$_1$ (5 μM) for 6 h.

B, C, D. Abrogation by 4-OH-E$_1$ of RSL3-induced accumulation of cellular NO (**B**), ROS (**C**) and lipid-ROS (**D**). Cells were treated with RSL3 (100 nM) ± 4-OH-E$_1$ (5 μM) for 6 h, and then stained for NO, ROS and lipid-ROS (**B, C** for fluorescence microscopy; scale bar = 65 μm; **D** for analytical flow cytometry). The quantitative intensity values are shown in the lower panel (n = 3; the mean value for the control group is labeled next to the bar).

E. Protective effect of 4-OH-E$_1$ against RSL3-induced cytotoxicity following treatment of cells with RSL3 (100 nM) ± 4-OH-E$_1$ (1.25, 2.5 and 5 μM) for 24 h (MTT assay, n = 4).

F. Levels of the dimer and monomer nNOS proteins and the total nNOS protein after treatment of cells with RSL3 (100 nM) ± SNAP (200 μM) for 6 h.

G, H, I. Abrogation by SNAP of RSL3-induced accumulation of cellular NO (**G**), ROS (**H**) and lipid-ROS (**I**). Cells were treated with RSL3 (100 nM) ± SNAP (200 μM) for 6 h, and then stained for NO, ROS and lipid-ROS (**G, H** for fluorescence microscopy; scale bar = 65 μm; **I** for analytical flow cytometry). The quantitative intensity values are shown in the right panel (n = 3; the mean value for the control group is labeled next to the bar).

J. Protective effect of SNAP against RSL3-induced cytotoxicity following treatment of cells with RSL3 (100 nM) ± SNAP (200 μM) for 24 h (MTT assay, n = 5).

K. Effectiveness of PDI-siRNAs in reducing cellular PDI protein level. Cells were
transfected with PDI-siRNAs for 24 and 48 h, respectively, and the PDI protein levels
were determined by Western blotting.

L. Protective effect of PDI knockdown on RSL3-induced cytotoxicity. Cells were
transfected with PDI-siRNAs for 24 h prior to treatment of cells with RSL3 (100 nM) for
an additional 24 h.

Each value represents the mean ± S.D. ($^{\#}P$ < 0.05; ** or $^{\#\#}P$ < 0.01).

Figure 7. Role of TrxR1 in RSL3-induced ferroptosis in HT22 cells.

A. Effectiveness of TrxR1-siRNAs in reducing cellular TrxR1 protein level. Cells were
transfected with TrxR1-siRNAs for 24, 48 and 72 h, and the TrxR1 protein levels were
determined by Western blotting.

B. Effect of TrxR1 knockdown on RSL3- and auranofin-induced cytotoxicity. Cells were
transfected with TrxR1-siRNAs for 24 h prior to the treatment of RSL3 (100 nM) or
auranofin (1 μM) for an additional 24 h, and then cell viability was determined (MTT
assay, n = 4).

C. Levels of TrxR1 enzyme activity in cells following treatment with RSL3 (100 nM) or
auranofin (1 μM) for 6 h. Cellular supernatants were prepared from different treatment
groups (using same numbers of cells), and then the TrxR1 enzyme activity was assayed.
The relative TrxR1 activity in the RSL3 and auranofin groups was presented as a
percentage of the relative TrxR1 activity in the vehicle-treated control group.

D, E. Levels of TrxR1, GPX4 and total nNOS proteins following treatment with 100 nM
RSL3 (**D**) or 1 μM auranofin (**E**) for different durations as indicated.

F. Levels of iNOS monomer, iNOS dimer, TrxR1 and PDI proteins in RSL3-treated cells
with TrxR1 knockdown. Cells were transfected with control-siRNAs or TrxR1 siRNAs for
24 h and then treated with 100 nM RSL3 for an additional 6 h.

G, H. Relative protein band intensity of TrxR1, GPX4 and total nNOS proteins following
treatment with 100 nM RSL3 (**G**) or 1 μM auranofin (**H**) for different durations as
indicated. The protein bands were determined by densitometry counting (the ImageJ
software). The relative intensity of a given protein was normalized to the corresponding
β-actin intensity, and the relative protein level in a given treatment group is presented
relative to the vehicle-treated control group.

H, I, J, K. Concentration- and incubation time-dependent inhibition of TrxR1's enzyme
activity *in vitro* by auranofin (**H, I**) and RSL3 (**J, K**). The TrxR1 enzyme activity in the
presence of an inhibitor (auranofin or RSL3) was presented as % of the control activity
in the absence of an inhibitor.

Each value is the mean ± S.D. (*P < 0.05; ** or $^{\#\#}P$ < 0.01; n.s., not significant).

Figure 8. Effect of auranofin on cell viability and cellular levels of NO, ROS and
lipid-ROS in HT22 cells.

A. Dose-dependent effect of auranofin on cell viability (MTT assay, n = 5).

B, C. Effect of Fer-1 (**B**) and DFO (**C**) on auranofin-induced cytotoxicity. The cells were

treated with auranofin (1 μM) ± Fer-1 or DFO at indicated concentrations for 24 h, and
change in cell viability was determined by MTT assay (n = 5).

D, E, F. Time-dependent accumulation of cellular NO (**D**), ROS (**E**) and lipid-ROS (**F**) in
auranofin-treated cells (analytical flow cytometry). The quantitative intensity values are
shown in the respective right panels (n = 3).

G. H. Effect of z-VAD-FMK (**G**) and Nec-1 (**H**) on auranofin-induced cytotoxicity. The
cells were treated with auranofin (1 μM) ± z-VAD-FMK or Nec-1 at indicated
concentrations for 24 h, and change in cell viability was determined by MTT assay (n =
5).

Each value represents the mean ± S.D. (*$P < 0.05$; **$P < 0.01$; n.s., not significant).

Figure 9. PDI mediates the cytotoxicity jointly induced by auranofin and BSO in
HT22 cells.

A. Dose-dependent effect of BSO on cell viability (MTT assay, n = 4).

B. Dose-dependent effect of auranofin ± BSO on cell viability. The cells were jointly
treated with auranofin (25, 50 and 100 nM) ± BSO (2, 4 and 6 μM) for 24 h, and then cell
viability was determined by MTT assay (n = 4).

C. Protein levels of the monomer and dimer nNOS, total nNOS, TrxR1 and GPX4 in cells
jointly treated with auranofin (50 nM) + BSO (6 μM) for different time intervals as
indicated. The relative protein levels are shown in the right panels. The relative
intensity of protein bands was normalized to β-actin, and calculated relative to the
vehicle (control) group.

D. Protective effect of Fer-1 against cytotoxicity jointly induced by auranofin and BSO.
The cells were treated with auranofin (50 nM) + BSO (6 μM) alone or in combination
with Fer-1 (1 μM) for 24 h, and cell viability was determined by MTT assay (n = 4).

E. Protective effect of cystamine and 4-OH-E$_1$ against cell death jointly induced by
auranofin + BSO. The cells were treated with auranofin (50 nM) and BSO (6 μM) alone
or in combination with cystamine (12.5 μM) or 4-OH-E$_1$ (2 μM) for 24 h, and cell
viability was determined by MTT assay (n = 4).

F, G, H. Time-dependent accumulation of cellular NO (**F**), ROS (**G**) and lipid-ROS (**H**) in
cells jointly treated with auranofin and BSO (analytical flow cytometry). The
quantitative intensity values are shown in the respective right panels (n = 3).

I, J, K. Abrogation by cystamine and 4-OH-E$_1$ of auranofin + BSO-induced accumulation
of cellular NO (**I**), ROS (**J**) and lipid-ROS (**K**). Cells were treated with auranofin (50 nM)
and BSO (6 μM) alone or in combination with cystamine (12.5 μM) or 4-OH-E$_1$ (2 μM)
for 6 h, and then stained for NO, ROS and lipid-ROS (analytical flow cytometry). The
quantitative intensity values are shown in the respective right panels (n = 3).

Each value represents the mean ± S.D. (* $P < 0.05$; ** or [##] $P < 0.01$; n.s., not significant).

Figure 10. Schematic illustration of the mechanism by which PDI plays an
important contributing role in mediating RSL3-induced ferroptotic cell death in
HT22 cells. It is proposed that PDI plays an important role in mediating RSL3-induced
ferroptotic cell death. PDI is activated in RSL3-treated cells resulting from its inhibition

of TrxR1, which subsequently leads to PDI-mediated NOS dimerization (activation) and
accumulation of cellular NO, ROS and lipid-ROS, and ultimately ferroptotic cell death.

Supplementary Figure S1. Additional data comparing the time-dependent
accumulation of cellular NO (A) and ROS (B) in RSL3-treated cells (analytical flow
cytometry). All the experimental conditions were the same as described for **Fig. 1G**
and 1H, except that a 3-h time point was added in this experiment. Each value is the
mean ± S.D. (n = 3).

Each value represents the mean ± S.D. (**$P < 0.01$; n.s., not significant).

Supplementary Figure S2. Effect of TrxR1 ± PDI knockdown by siRNAs on RSL3-
induced ferroptosis in HT22 cells.

A. Effect of TrxR1 knockdown by TrxR1-siRNAs on the viability of HT22 cells (MTT
assay, n = 5). Cells were transfected with the control and TrxR1-specific siRNAs for 24 h,
and then the cells were treated with a vehicle (without RLS3) for 24 h. Then the MTT
assay was performed, and the absolute absorbance values (after subtracting the
background value) are shown.

B. Effect of joint knockdown of PDI and TrxR1 by siRNAs on the sensitivity to 100 nM
RSL3-induced cell death (MTT assay, n = 5). The viability of vehicle-treated cells in the
siCon group or in the siPDI + siTrxR1 group are individually set as 100% when
compared with the respective RSL3-treated cells.

Each value represents the mean ± S.D. (** or ##$P < 0.01$).

Supplementary Figure S3. Effect of EN460 (a ERO1 inhibitor) on RSL3-induced
ferroptosis in HT22 cells.

A. Effect of EN460 on RSL3-induced cytotoxicity following treatment of cells with RSL3
(100 nM) ± EN460 (5, 10 and 20 μM) for 24 h (MTT assay, n = 4).

B, C, D. Effect of EN460 on RSL3-induced accumulation of cellular NO (**B**), ROS (**C**) and
lipid-ROS (**D**). Cells were treated with RSL3 (100 nM) ± EN460 (20 μM) for 6 h, and then
stained for NO, ROS and lipid-ROS (**B, C** for fluorescence microscopy; scale bar = 65 μm;
D for confocal microscopy, scale bar = 10 μm).

Each value represents the mean ± S.D. (** or ##$P < 0.01$; n.s., not significant).

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Figure 10