Title: Oxydifficidin, a potent *Neisseria gonorrhoeae* antibiotic due to DedA assisted uptake and ribosomal protein RplL sensitivity

Authors: Jingbo Kan[1, 2, 3], Adrian Morales[1], Yozen Hernandez[1], Melinda A. Ternei[1], Christophe Lemetre[1], Logan W. MacIntyre[1], Nicolas Biais[2, 3, 4, *], Sean F. Brady[1, *]

[1] Laboratory of Genetically Encoded Small Molecules, The Rockefeller University, 1230 York Avenue, New York, NY 10065.
[2] Graduate Center, City University of New York, New York, NY 10016.
[3] Brooklyn College, City University of New York, Brooklyn, NY 11210.
[4] Laboratoire Jean Perrin, UMR 8237 Sorbonne Université/CNRS, Paris, France.

*Corresponding Authors: Nicolas Biais and Sean F. Brady.
Contact: Laboratory of Genetically Encoded Small Molecules
The Rockefeller University
1230 York Avenue
New York, NY 10065
Phone: 212-327-8280
Fax: 212-327-8281
Email: sbrady@rockefeller.edu

Acknowledgements: We thank Nathan Weyand at Ohio University for providing Neisseria stains, the Bacillus Genetic Stock Center for providing *Bacillus amyloliquefaciens* strains, Rinat Abzalimov and James Aramini at Advanced Science Research Center (ASRC) for performing initial LCMS and NMR, David Dubnau and Jeanette Hahn at Rutgers New Jersey Medical School for assistance with Bacillus genetics, and Hilda Amalia Pasolli at The Rockefeller University Electron Microscopy Resource Center for collecting scanning electron microscope images. We acknowledge the members of the MMBL lab at Brooklyn College for their help and input in this work. This work was supported by 5R35GM122559 (SFB), NIH AI116566 (NB) and iBio Initiative (NB).

Data availability: Source data for this paper is available upon request from Sean Brady.
Supporting information: The following are provided as supporting information: ^{13}C and ^{1}H NMR spectra and chemical shift data for oxydifficidin. MS/MS data and analysis for oxydifficidin.
Code availability: No custom code was used for this study.
Competing Interest Statement: The authors declare no competing financial interests.
Keywords: Oxydifficidin; *Neisseria gonorrhoeae*; DedA; L7/L12 (RplL); ribosome; antibiotic; antibiotic resistance.

Abstract

Gonorrhea, which is caused by *Neisseria gonorrhoeae*, is the second most prevalent sexually transmitted infection worldwide. The increasing appearance of isolates that are resistant to approved therapeutics raises the concern that gonorrhea may become untreatable. Here, we serendipitously identified oxydifficidin as a potent *N. gonorrhoeae* antibiotic through the observation of a *Bacillus amyloliquefaciens* contaminant in a lawn of *N. gonorrhoeae*. Oxydifficidin is active against both wild-type and multidrug-resistant *N. gonorrhoeae*. It's potent activity results from a combination of DedA-assisted uptake into the cytoplasm and the presence of an oxydifficidin-sensitive ribosomal protein L7/L12 (RplL). Our data indicates that oxydifficidin binds to the ribosome at a site that is distinct from other antibiotics and that L7/L12 is uniquely associated with its mode of action. This study opens a potential new avenue for addressing antibiotic resistant gonorrhea and underscores the possibility of identifying overlooked natural products from cultured bacteria, particularly those with activity against previously understudied pathogens.

Main Text

Introduction

Gonorrhea is the second most prevalent sexually transmitted infection worldwide, its causative agent is the bacterium *Neisseria gonorrhoeae*. According to the World Health Organization (WHO) approximately 82.4 million new adult gonorrhea infections occurred globally in 2020.[1] The high dose (500 mg) of the cephalosporin ceftriaxone is currently the only recommended therapy for treating gonorrhea infections.[2] The growing instances of drug resistant "superbugs", together with the limited clinical treatment options, underscore the urgent need for additional antibiotics that target *N. gonorrhoeae*.[3-8] The characterization of antibacterial active natural products inspired the development of both ceftriaxone and azithromycin.[9, 10] Bacterial natural products have been a key source of antibiotics with diverse modes of action and the most fruitful source of therapeutically useful antibiotics.[11-13] Here we describe the serendipitous identification of the natural product oxydifficidin[14], [15] as a potent *N. gonorrhoeae* active antibiotic and show that this activity arises from a combination of DedA flippase assisted uptake and ribosomal protein L7/L12 (RplL) sensitivity. Oxydifficidin provides a new therapeutic lead structure for addressing the growing problem of antibiotic resistant gonorrhea. Over the last century, bacteria were extensively examined for antibiotic production in screens that often focused on a small number of pathogens. This study suggests that reexamining cultured bacteria for antibiotics active against today's emerging pathogens may be fruitful as metabolites with specific potent activity against historically less problematic pathogens may have been overlooked.

Results

Oxydifficidin isomers selectively and potently inhibit *N. gonorrhoeae*

In our day to day experiments we regularly use agar plates containing lawns of pathogenic bacteria. During these experiments we often find random environmental bacteria growing on these plates. On one lawn of *N. gonorrhoeae* we observed an environmental contaminant that was surrounded by a zone of growth inhibition suggesting that it produced an anti-*N. gonorrhoeae* metabolite (**Figure 1a**). When we screened this contaminant for antibacterial activity against lawns of other Gram-negative bacteria it did not produce a zone of growth of inhibition against any of the bacteria we tested (*e.g., Escherichia coli*, *Vibrio cholerae*, *Caulobacter crescentus*). Since antibiotics that preferentially inhibit the growth *N. gonorrhoeae* are rare, we looked at this contaminant in more detail. Sequencing of the contaminant's genome and genome clustering analysis (**Figure S1a**) revealed that it was most closely related to *Bacillus amyloliquefaciens*, which is a root-colonizing bacterium that is used as a biocontrol agent.[16] We named the anti-*N. gonorrhoeae* contaminant *Bacillus amyloliquefaciens* BK.

To identify the biosynthetic gene cluster (BGC) responsible for the observed antibiosis we screened *B.*
amyloliquefaciens BK transposon mutants for strains that no longer produced anti-*N. gonorrhoeae*
activity. The sequencing of one non-producer strain revealed that it surprisingly contained four
transposon insertions and one frame shift mutation (**Figure S1b**). The frame shift mutation and one
transposon insertion were predicted to each disrupt unique BGCs. The transposon inserted into the
bacillomycin (mycosubtilin) BGC, while the frame shift mutation was predicted to disrupt the
(oxy)difficidin BGC. To determine which of these two BGCs was responsible for the *N. gonorrhoeae*
activity we tested *B. amyloliquefaciens* strains with targeted disruptions of each BGC for activity against
N. gonorrhoeae.(17) Only disruption of the (oxy)difficidin BGC eliminated the anti-*N. gonorrhoeae*
activity (**Figure S1c**). To confirm the identity of the *N. gonorrhoeae* active antibiotic we carried out a
bioassay guided fractionation of *B. amyloliquefaciens* BK culture broth.(14), (15) HRMS and NMR data
from the major active peak we isolated were consistent with its being an oxydifficidin isomer (**Figure**
1b, Figure S5 - 11, and Table S2). Oxydifficidin contains a 27-carbon polyketide backbone that is
cyclized through a terminal carboxylic acid and an oxidation at position 21. This hydrophobic core is
phosphorylated at C16. Oxydifficidin occurs naturally as a collection of interconverting thermal isomers
(**Figure 1b**). As reported previously, we observed an interconversion of isomers with the compound we
purified from *B. amyloliquefaciens* BK cultures.(14), (15) All assays were performed using the mixture
of interconverting oxydifficidin isomers we obtained from *B. amyloliquefaciens* BK cultures.

We tested oxydifficidin for activity against diverse bacterial pathogens. Oxydifficidin showed only weak
activity against most pathogens, however we observed potent activity against *N. gonorrhoeae* (**Figure**
1c). When we examined other *Neisseria* species, we found that oxydifficidin was consistently more
active against *Neisseria* than any of the other bacteria we tested. Among *Neisseria* spp., oxydifficidin
was most active against *N. gonorrhoeae* underscoring a unique narrow spectrum of potent activity. A
key issue with the current treatment of *N. gonorrhoeae* infections is the development of resistance to
existing therapeutics. Resistance to the standard of care cephalosporins is particularly problematic.
Oxydifficidin was more potent against wild-type (WT) *N. gonorrhoeae* than almost all other antibiotics
we tested. In fact, it was only slightly less active than the highly optimized third-generation
cephalosporin, ceftazidime.(18) However, unlike third-generation cephalosporins, oxydifficidin retained
activity against the multidrug resistant H041 clinical isolate (**Table 1**).(4) H041 is resistant to the
"standard of care" cephalosporin ceftriaxone (2 µg/mL) as well as a number of other antibiotics that are
normally active against *N. gonorrhoeae* (penicillin G, 4 µg/mL; cefixime, 8 µg/mL; levofloxacin, 32
µg/mL).

Oxydifficidin's structure is interesting as phosphorylated antibiotics, and moreover natural products in
general, remain rare.(19, 20) Its structure together with its specific and potent activity against drug resist
N. gonorrhoeae suggested it might have a unique mode of action (MOA). This is appealing from the
perspective of developing therapeutics that are capable of circumventing clinically problematic
resistance mechanisms and therefore we focused on characterizing the mechanism of oxydifficidin's
potent anti-*N. gonorrhoeae* activity.

DedA assists the uptake of oxydifficidin into *N. gonorrhoeae*
As a first step to understanding the MOA of oxydifficidin, we raised resistant mutants by directly plating
N. gonorrhoeae cultures on antibiotic containing plates (1 µg/ml, 4x MIC) (**Figure 2a**). Out of the
>1.5X10^{10} cells we screened, only one resistant mutant appeared. Oxydifficidin's MIC against this
mutant increased by 8-fold (2 µg/ml). No increase in MIC was observed for any other antibiotic we
tested (**Figure 2a**). Sequencing and comparison of this mutant's genome to the sensitive parent
genome revealed a single mutation that introduced a frame shift in one of the three predicted *dedA*
genes found in the *N. gonorrhoeae* MS11 (NCBI:txid528354 *dedA* NGFG_RS04905(21)). To confirm
that DedA was necessary for oxydifficidin's potent activity, we created a *dedA* deletion mutant in a clean
N. gonorrhoeae background. This mutant showed the same 8-fold increase in oxydifficidin's MIC,

confirming that the deletion of *dedA* is sufficient to reduce oxydifficidin's potent activity. Interestingly,
not only was *dedA* deficient *N. gonorrhoeae* less susceptible to oxydifficidin, oxydifficidin also kills this
mutant more slowly (**Figure 2b**) than WT *N. gonorrhoeae*.
105
The DedA protein superfamily is highly conserved, with examples in almost every sequenced genome
across all domains of life.(22) DedA family members are predicted to be transmembrane proteins with
still largely, poorly defined, functions. However, a few recent studies indicate that DedA homologs are
flippases. They have been reported to flip phospholipids (phosphatidylethanolamine,
phosphatidylserine) or phospholipid like structures (C55-isoprenyl pyrophosphate) across prokaryotic
and eukaryotic lipid bilayers.(23-26) Although oxydifficidin is not a phospholipid, its overall structure
resembles that of reported DedA protein substrates, especially C55-isoprenyl pyrophosphate (**Figure
3a**). Interestingly, among the two characterized bacterial family members, the DedA protein associated
with oxydifficidin potency is most closely related to the C55-isoprenyl pyrophosphate flippase YghB
from *Vibrio cholerae* (**Figure S2**).(23) As oxydifficidin's activity was not completely abrogated in the
dedA knockout we postulated that DedA was not the direct target of oxydifficidin, but it instead acted to
increase oxydifficidin's potency in *N. gonorrhoeae*. The structural similarity between oxydifficidin and
the known substrates of DedA homologs led us to explore the possibility that DedA was responsible for
assisting with oxydifficidin uptake into *N. gonorrhoeae*.
120
We examined the effect of DedA on antibiotic accumulation by comparing the amount of compound
found in the cell pellet collected from antibiotic treated cultures of wild type and *dedA* knockout *N.
gonorrhoeae* strains (**Figure 3b**). In the case of oxydifficidin we saw 6 times more antibiotic in cells
from wild type cultures than from *dedA* deletion strain cultures. For the other antibiotics we tested
(tetracycline and chloramphenicol) this ratio was less than two. Based on DedA homologs flipping
phospholipid-like structures across a lipid bilayer our data is consistent with DedA flipping oxydifficidin
across the inner membrane to increase its cytoplasmic concentration and in turn increase its potency
against *N. gonorrhoeae*. A DedA assisted uptake mechanism could also explain the slower rate of
killing we observed for oxydifficidin against *dedA* deficient *N. gonorrhoeae* compared to wild type *N.
gonorrhoeae*. While we cannot definitely rule out the possibility that DedA accumulation of oxydifficidin
in the membrane could also have a direct toxicity effect, we did not detect any cell lysis or membrane
depolarization at even 100 times oxydifficidin's MIC (**Figure S3**).
133
Oxydifficidin inhibits protein synthesis by interacting with RplL
To look for an intracellular target of oxydifficidin we carried out a second round of resistant mutant
screening. In this case, we used the *N. gonorrhoeae dedA* deletion strain and searched for colonies
with an even higher tolerance to oxydifficidin. From ~1x10^{10} cells plated on 8 µg/ml oxydifficidin (4x
MIC for *N. gonorrhoeae dedA* deletion strain*)* we identified 12 resistant mutants. In each case the
oxydifficidin MIC increased to 16 µg/ml. Sequencing of these strains revealed that each contained a
point mutation in the gene encoding for the large ribosomal protein(s) L7/L12 (*rpl*L). Eleven strains
contained the same R76C mutation and one contained a K84E mutation (**Table S1**). To determine if
mutations in the *rpl*L gene alone were sufficient to confer oxydifficidin resistance, we created an RplL
R76C mutant in a wild type *N. gonorrhoeae* background (*i.e.,* non *dedA* deletion). This mutant exhibited
an 8-fold increase in oxydifficidin's MIC (2 µg/ml) compared to the parent strain, confirming that the
R76C mutation in L7/L12 alone was sufficient to increase the MIC of oxydifficidin. Ribosomal proteins
L7 and L12 have the same sequence, however L12 has an N-terminal acetylation.(27) L7/L12 is part
of the L10/L7 stalk of the large (50S) subunit of the bacterial ribosome and is critical to a number of
processes including GTP hydrolysis.(27, 28)
149
150 The appearance of resistance mutations in *rpl*L suggested that oxydifficidin inhibited protein synthesis,
151 which would be consistent with isotope feeding studies performed with difficidin and *E. coli*.(29) We
152 initially tested this hypothesis *in vitro* using a coupled transcription/translation system. While this
153 reaction mixture contained all the components necessary to produce a protein from DNA, no protein

was produced in the presence of oxydifficidin (**Figure 3c**). Using a coupled system, it was not possible to distinguish between inhibition of RNA or protein synthesis, and therefore, to rule out inhibition of transcription, we next looked directly at RNA synthesis *in vitro*. In this case we saw no inhibition of RNA synthesis, even at the highest oxydifficidin concentration we tested (13.4 µg/ml) (**Figure 3d**). Taken together these two experiments indicate that oxydifficidin inhibits translation but not transcription and are consistent with *rplL* mutations providing resistance to oxydifficidin.

To the best of our knowledge *rplL* mutations have not been previously associated with antibiotic resistance and no characterized antibiotics have been found to bind L7/L12. When we screened for cross resistance, the L7/L12 R76C mutation did not confer resistance to any other antibiotics we tested (**Table 2**). These included ribosome targeting antibiotics with diverse binding sites. Antibiotics that bind different regions of the 30S decoding center, including tetracycline, gentamicin and spectinomycin showed no increase in MIC with the L7/L12 R76C mutation.(30, 31) Similarly, the L7/L12 R76C mutation did not confer resistance to chloramphenicol or erythromycin, which interact with distinct regions of the 50S peptidyl transferase center.(30, 31) Antibiotics that bind away from these two hot spots also showed no cross resistance.(30-32) The activity of avilamycin, which binds in a unique site in the 50S subunit at the entrance to the A-site tRNA accommodating corridor was also not affected by the L7/L12 R76C mutation.(33) In the case of thiostrepton A(34), which targets the L11 protein GTPase-associated center and is part of the only class of antibiotics known to bind the ribosome in the vicinity of L10/L7 stalk, we also observed no change in MIC for *N. gonorrhoeae* with the L7/L12 R76C mutation. The inability of the L7/L12 R76C mutation to confer cross resistance to known antibiotics suggests that oxydifficidin binds to a different site on the ribosome and that L7/L12 is uniquely associated with oxydifficidin's mode of action.

To determine whether the DedA and RplL mutations we observed were specific to oxydifficidin resistance in *N. gonorrhoeae*, we selected for resistant mutants using two other *Neisseria* species: *Neisseria subflava* and *Neisseria cinerea*. Both strains were natively less susceptible to oxydifficidin than *N. gonorrhoeae*. In the case of *N. cinerea* all resistance mutants contained the RplL K84E mutation that we first observed in *N. gonorrhoeae,* and in the case of *N. subflava* all resistance mutants contained mutations in *dedA* (**Table S1**). Other *Neisseria* species likely show higher MICs than *N. gonorrhoeae* because of either the absence of DedA assisted uptake or reduced RplL protein oxydifficidin sensitivity. The model that arises from our studies is that high oxydifficidin sensitivity results from a combination of DedA assisted oxydifficidin uptake into the cytoplasm and the presence of an oxydifficidin sensitive RplL protein (**Figure 3e**). *N. gonorrhoeae* is uniquely sensitive to oxydifficidin because it contains both. DedA is critical to oxydifficidin's potent activity as it not only sensitizes *N. gonorrhoeae* to oxydifficidin but also decreases its kill time, both of which are appealing from a clinical development perspective.

Discussion

Almost 60% of clinically approved antibiotics target the ribosome.(35) Although it is the most common target for natural product antibiotics, most of these molecules inhibit the ribosome by binding to only a small number of sites.(30, 36) Clinically approved antibiotics generally inhibit protein synthesis by binding in either the ribosomal decoding center, peptidyl transfer center or the peptide exit tunnel. Targeting unexploited sites of the ribosome is considered a key step to developing next generation antibiotics that can circumvent the existing antibiotic resistance mechanisms. Our data suggest that oxydifficidin has a distinct binding site in the ribosome compared to other clinically used antibiotics making it an appealing therapeutic candidate. In our resistance mutant screening experiments with *N. gonorrhoeae* the frequency of mutations in either *dedA* or *rplL* was quite low (~10^{-9}). The *dedA* deletion mutant also showed an altered cell morphology with reduced membrane integrity and lower formation of micro-colonies (**Figure S4**), indicating that it should show reduced pathogenesis and fitness, and, as a result, not accumulate in a clinical setting, which adds to the therapeutic appeal of oxydifficidin.

While most natural product research efforts are focused on identifying novel chemical entities, the work
presented here suggests that reexamining old sources through the lens of today's most important
pathogens may also be a productive approach for identifying therapeutically appealing antibiotics.
Increasing rates of antibiotic resistance present a significant clinical threat as they have the potential of
rendering gonorrhea infections untreatable. The recent appearance of *N. gonorrhoeae* "superbugs"
with high-level resistance to all currently recommended drugs treating gonorrhea infections, is
particularly concerning.(4, 5) One appealing feature in the original reports of the discovery of
oxydifficidin was its activity in an animal model, a key hurdle for many natural products.(15)
Unfortunately oxydifficidin's clinical significance was limited because of the weak activity it exhibited
against most bacterial pathogens. Our identification of oxydifficidin as being specifically potent against
N. gonorrhoeae including multi-drug resistant *N. gonorrhoeae*, provides a potential new path forward
for this structurally interesting natural antibiotic.
219
220
221 **Materials and Methods**
222
Bacterial Strains and Cultivation
Neisseria strains were grown at 37 °C with 5% CO_2 in GCB medium(37) (VWR CA90002-016). *Bacillus*
amyloliquefaciens BK was isolated from a lab agar plate, *Bacillus amyloliquefaciens* FZB42 wild type
(WT) and various mutant strains were provided by Bacillus Genetic Stock Center. All Bacillus strains
were grown at 30 °C in Luria-Bertani (LB) medium. *Escherichia coli* DH5α, *Klebsiella pneumoniae*
ATCC 10031, *Enterobacter cloacae* ATCC 13047, *Acinetobacter baumannii* ATCC 17978,
Pseudomonas aeruginosa PAO1, *Morganella morganii* ATCC 25830 and *Staphylococcus aureus*
USA300 were grown at 37 °C in Tryptic Soy Broth (TSB).
231
Crude Extraction and Disc Diffusion Test
Bacillus strains were grown overnight at 30 °C on LB agar. 3 ml of methanol was added to the top of
the plate and spread uniformly by gentle shaking. The supernatant was then collected, centrifuged at
15,000 rpm for 2 min and filtered through a 0.2 μm cellulose acetate membrane (VWR). Then, 10 μl of
the crude extract was added onto a paper disc (VWR) and after 5 min drying the paper disc was
transferred on top of GCB agar plate lawned (swabbed from a single colony) with *N. gonorrhoeae* cells.
The plate was incubated at 37 °C with 5% CO_2 overnight and the inhibition zone was visually inspected.
239
Fermentation of Bacillus amyloliquefaciens BK
A single colony of Bacillus *amyloliquefaciens* BK was grown in 250 ml LB in a 1 L flask at 30 °C on a
rotary shaker (200 rpm) overnight, then 50 ml of the overnight culture was transferred to 1 L modified
Landy medium (20 g glucose, 5 g glutamic acid, 1 g yeast extract, 1 g K_2HPO_4, 0.5 g $MgSO_4$, 0.5 g
KCl, 1.6 mg $CuSO_4$, 1.2 mg $Fe_2(SO_4)_3$, 0.4 mg $MnSO_4$ per 1 L deionized H_2O, pH adjusted to 6.5 before
autoclaving) in a 2.8 L triple baffled flask and fermented at 30 °C on a rotary shaker (200 rpm) for 3
days.
247
Isolation of Oxydifficidin
The isolation process followed the published protocol(14) with adjustments. The fermentation culture
(5 L) was centrifuged at 4,000 rpm for 10 min and the supernatant was acidified to pH 3.0 before
extraction with 5 L of ethyl acetate. The crude extract was concentrated under reduced pressure (60
mBar) on a rotary evaporator to a final volume of about 20 ml. The concentrated sample was then
loaded onto Diaion HP-20 resin (Sigma-Aldrich) and the resin was then washed by 500 ml of water,
500 ml of methanol-water (3:7) and 500 ml of methanol-water (7:3). The oxydifficidin rich cut was then
obtained by eluting the resin with 500 ml of methanol. This rich cut was concentrated and absorbed
onto DEAE-Sephadex A-25 (Cl⁻) resin (Sigma-Aldrich). The resin was washed with 250 ml of methanol-
water (2:8), 500 ml of methanol-water (9:1), and a rich cut was then obtained by eluting the resin with

100 ml methanol-water (2:8) with 3% ammonium chloride. The rich cut from A-25 resin was then diluted
with 300 ml of water and adsorbed onto Amberlite XAD-16N resin (Sigma-Aldrich) and washed
successively with 500 ml of water, 250 ml of 0.075 M K_2HPO_4 (pH 7) and 250 ml of methanol-water
(2:8), followed by elution with 100 ml of methanol. This oxydifficidin-containing eluate was diluted with
100 ml of 0.075 M K_2HPO_4 (pH 7) and applied to a RediSep Rf C18 column (Teledyne ISCO, 100 Å,
20 - 40 µm, 300 ± 50 m²/g) with a reverse-phase linear gradient system (5-100% 0.075 M
K_2HPO_4/MeOH, 45 min). Fractions containing oxydifficidin was monitored by anti-*N. gonorrhoeae*
activity and further purified by semipreparative HPLC using a reverse-phase C18 column (Waters,130
Å, 5 µm, 10 mm × 150 mm) with an isocratic system (0.075 M K_2HPO_4/MeOH (32:68)). The purified
oxydifficidin was then desalted by Amberlite XAD-16N resin (Sigma-Aldrich).
268
Electrospray Ionization LC-MS and NMR Analysis
LC–high-resolution mass spectrometry (HRMS) data were acquired using a SCIEX ExionLC HPLC
system coupled to an X500R QTOF mass spectrometer (The Rockefeller University). The system was
equipped with a Phenomenex Kinetex PS C18 100 Å column (150 mm × 2.1 mm, 2.6 µm) and operated
with SCIEX OS v.2.1 software. The following chromatographic conditions were used for LC–HRMS:
5% B to 2.0 min; 5 - 95% B from 2.0 to 30.0 min; 95% B from 30.0 to 37.0 min; 95 - 5% B from 37.0 to
38.0 min; and 5% B from 38.0 to 45.0 min (flow rate of 0.25 ml/min, 3 µl injection volume, A/B:
water/acetonitrile, supplemented with 0.1% (v/v) formic acid). Both electrospray ionization modes (ESI)
Full HRMS spectra were acquired in the range m/z 100-1000, applying a declustering potential of 80
V, collision energy of 5 V, source temperature of 500 °C and a spray voltage of 5500 V. A maximum of
7 candidate ions from every Full HRMS event were subjected to Q2-MS/MS experiments in the range
m/z 60-1000, applying a collision energy of 35 ± 10 V for both ESI modes. Full HRMS and the most
intense MS/MS spectra were analyzed with MestReNova software (14.3.0). Nuclear magnetic
resonance (NMR) spectra were acquired on a Bruker Avance DMX 800MHz spectrometer equipped
with cryogenic probes (New York Structural Biology Center). All spectra were recorded at 298 K in
$CD_3OD – D_2O$ (1:1). Chemical shift values are reported in parts per million (ppm) and referenced to
residual solvent signals: 3.31 ppm (1H) and 49.0 ppm (C). Spectra analysis and visualization were
carried out in TopSpin (3.6.0) and MestReNova (14.3.0).
287
Minimum Inhibitory Concentration (MIC) Assay
Neisseria strains were grown on GCB agar plate overnight, cells were collected by a polyester swab
tip (Puritan) and resuspended in 1 ml GCB medium followed by a 1 in 5,000-fold dilution. The stock
solution of antibiotics was serial diluted 2-fold in a 96-well plate (ThermoFisher Scientific) containing
GCB liquid medium, then an equal amount of diluted bacterial culture was added to each well and
mixed by pipetting. The plates were incubated at 37 °C with 5% CO_2 for 16 hours. *E. coli*, *K.*
pneumoniae and *M. morganii* strains were grown at 37 °C in Tryptic Soy Broth (TSB) medium overnight.
The stock solution of antibiotics was serial diluted 2-fold in a 96-well plate containing TSB agar medium,
then 10^4 cells were spotted on each well. The plates were incubated at 37 °C for 16 hours. The top and
bottom rows of plates contained an equal volume of media alone to prevent edge effect. The last column
did not contain antibiotic to serve as a control for cell viability. The MIC of antibiotics was determined
by visual inspection as the concentration in the well that prevents bacterial growth compared to control
wells. The MIC assays were performed in duplicate.
301
Phylogenetic Tree Building
The phylogenetic trees for bacterial genomes were built using Genome Clustering function with default
parameters in MicroScope(38). The bacterial DedA family protein sequences were downloaded from a
previous study(39) and aligned using MUSCLE v5(40) with the "Super5" algorithm. The phylogenetic
tree was built using FastTree(41) with the Jones-Taylor-Thornton (default) mode of amino acid
evolution. All trees were visualized and edited in iTOL.
308
Transposon Mutagenesis

Transposon mutagenesis was performed by electroporation using the EZ-Tn5 <KAN-2> Tnp
Transposome kit (Lucigen). Briefly, an overnight culture of *Bacillus amyloliquefaciens* BK was diluted
100 fold in NCM medium(42) and grown at 37 °C on a rotary shaker (200 rpm) until an OD600nm of
0.5. The cell culture was then supplemented with glycine, DL-threonine and Tween 80 at a final
concentration of 3.89%, 1.06% and 0.03% respectively, and grown at 37 °C on a rotary shaker (200
rpm) for 1 h. The culture was then cooled on ice for 30 min and collected by centrifugation (4,000 g, 10
min) at 4 °C. The cell pellet was washed three times with ETM(43) buffer. The washed cell pellet was
resuspended in 100 µl of ETM buffer containing 0.25 mM KH_2PO_4 and 0.25 mM K_2HPO_4 and 1 µl of
the EZ-Tn5 transposome was added. The cell mixture was transferred to a 1 mm electroporation
cuvette and pulsed using a Gene Pulser Xcell Electroporation System (Bio-Rad) using 2.1 kV/cm, 150
Ω and 36 µF. The mixture was immediately and gently mixed with 1 ml NCM medium containing 0.38
M mannitol and recovered in a round bottom tube (VWR) at 37 °C on a rotary shaker (120 rpm) for 3 h.
After recovery, the cells were concentrated by centrifugation and spread on a LB plate with 50 µg/ml
kanamycin and incubated at 37 °C overnight and single colonies collected for subsequent analysis.
324
Mutant development
For natural mutagenesis, single colonies of *N. gonorrhoeae* cells were swabbed on GCB agar plates
and grown at 37 °C with 5% CO_2 for 16 hours. The grown cells were collected by polyester swab tip
(Puritan) in GCB medium and cell number was calculated by measuring optical density (OD, estimated
value of OD 0.7 = 5 x 10^8 CFU (Colony-forming unit)/ml). Cells were then uniformly spread out on GCB
agar plate containing 4x MIC of oxydifficidin (1 µg/ml for WT and 8 µg/ml for ΔdedA) and grown at 37
°C with 5% CO_2 until mutant colonies are observed.
332
Mutant construction
For *dedA* gene deletion, the 3' (Primer: *dedA* 3' overhang F/R) and 5' (Primer: *dedA* 5' overhang F/R)
overhang regions of *N. gonorrhoeae dedA* gene and a kanamycin resistance cassette (Primer: kan^R
(kanamycin resistance) cassette F/R) were amplified using GoTaq master mix (Promega). The
fragments were then assembled using NEBuilder HiFi DNA assembly master mix (NEB) to place the
cassette sequence positioned in the middle. Transformation of the construct to *N. gonorrhoeae* was
done by using spot transformation protocol(44). For mutant *rplL*_R76C, the mutated *rplL* gene including
the 3' and 5' overhangs (Primer: *rplL*_R76C F/R) was amplified using *N. gonorrhoeae* MS11 Δ*dedA*
*rplL*_R76C as the PCR template. Another construct incorporating a kanamycin resistance cassette with
the 3' (Primer: *trpB-lga* 3' overhang F/R) and 5' (Primer: *trpB-lga* 5' overhang F/R) overhangs from
locus 272,353 bp (*trpB-lga*) of *N. gonorrhoeae* WT genome was assembled using HiFi assembly. These
2 constructs were co-transformed(45) into *N. gonorrhoeae* WT cells. The sequence verification was
carried out using Sanger sequencing services provided by Genewiz.
346
Genomic Sequencing and Mutational Analysis
Genomic DNA from Neisseria WT and resistance mutants was isolated using PureLink Microbiome
DNA purification kit (Invitrogen) according to manufacturer's instructions. The whole-genome
sequencing was performed by a MiSeq Reagent Kit v3 using Illumina MiSeq system following
manufacturer's instructions. Genomes were assembled and mapped using corresponding reference
genomes. The single nucleotide polymorphism (SNP) was detected by aligning the mutant's
sequencing reads to the genomic sequence of the WT strain.
354
Time Dependent Killing Assay
N. gonorrhoeae cells were grown on a GCB agar plate at 37 °C with 5% CO_2 for 16 hours and the
overnight cells were normalized to ~10^5 CFU/ml with GCB medium in a 14 ml round-bottom tube (VWR).
The normalized cell culture was supplemented with 8x MIC of oxydifficidin (2 µg/ml for WT and 16 µg/ml
for ΔDedA) and incubated at 37 °C with 5% on a rotary shaker (200 rpm). 100 µl of the culture was
collected and mixed thoroughly into 10 ml of GCB medium at 5 min, 10 min, and 30 min respectively,

then 200 µl of the diluted culture was spread on an GCB agar plate and grown at 37 °C with 5% CO_2 overnight. Colonies were counted the following day. The killing assays were done in triplicate.

Cell Lysis assay
The cell lysis assay was conducted using SYTOX green (Invitrogen) following manufacturer's instructions. Briefly, 1 ml of cell suspension in Live Cell Imaging Solution at OD 0.7 was stained by adding 0.4 µl of 5 mM SYTOX solution and incubated in the dark at RT for 5 min. Subsequently, 30 µl of the stained culture was added into a 384-well Flat Clear Bottom Black plate (Corning) and the fluorescence signal was recorded by an Infinite M NANO$^+$ (TECAN) with Excitation/Emission = 488/523 nm. The interval time was set to 7 s. Once the signal reached equilibrium, 30 µl of antibiotics diluted in Live Cell Imaging Solution at 16x MIC was added to the culture and mixed thoroughly. The fluorescence signal monitoring was then continued for 1 h. The cell lysis assays were done in triplicate.

Cell Depolarization assay
The cell depolarization assay was conducted using $DiSC_3(5)$ dye (Invitrogen) following manufacturer's instructions. Briefly, 1 ml of cell suspension in Live Cell Imaging Solution at OD 0.7 was stained by adding 1 µl of 2 mM SYTOX solution and incubated in the dark at RT for 15 min. Then, 30 µl of the stained culture was added into a 384-well Flat Clear Bottom Black plate (Corning) and the fluorescence signal was recorded by an Infinite M NANO$^+$ (TECAN) with Excitation/Emission = 622/675 nm. The interval time was set to 7 s. Once the signal reached equilibrium, 30 µl of antibiotics diluted in Live Cell Imaging Solution at16xMIC was added to the culture and mixed thoroughly. The fluorescence signal monitoring was then continued for 1 h. The cell depolarization assays were done in triplicate.

Drug Accumulation Assay
N. gonorrhoeae cells were grown on a GCB agar plate at 37 °C with 5% CO_2 for 9 hours and the overnight cells were normalized to ~10^8 CFU/mL with GCB medium in a 14 ml round-bottom tube (VWR). Cells were incubated with 0.125 µg/ml of oxydifficidin at 37 °C with 5% CO_2 on a rotary shaker (200 rpm) for 3 hours. Cell pellets were collected by centrifugation at RT, washed twice with fresh GCB medium, and resuspended in an equal volume of GCB medium to that of the supernatant. The cell suspension and supernatant were then separately extracted by ethyl acetate in a 1:1 ratio. The extracts were evaporated to dryness under vacuum, resuspended in 100 µl of methanol and then 3 µl of the resuspension was injected to LC-HRMS for the quantification of oxydifficidin using target peak area. Experiment 1 (oxydifficidin and tetracycline) was done in duplicate, and experiment 2 (oxydifficidin and chloramphenicol) was done in triplicate.

In Vitro Transcription and Translation Inhibition Assays
The *in vitro* transcription assay was conducted using the HiScribe T7 High Yield RNA Synthesis Kit (NEB) following the manufacturer's instructions. 24 µM of oxydifficidin was added to the transcription mixture, and 20 mM of EDTA was added to provide a positive control for inhibition. The yielded RNA was detected by a BioAnalyzer (Agilent). The *in vitro* translation assay was performed using PURExpress In Vitro Protein Synthesis Kit (NEB). 24 µM of antibiotics were added individually to the translation mixture and incubated at 37 °C for 4 hours, the samples were then analyzed by SDS-PAGE in a 5 – 20% gradient gel.

Membrane Integrity Assay
The membrane integrity assay was conducted using BacLight Bacterial Viability Kit (Invitrogen) following manufacturer's instructions. Briefly, 1 ml of cell suspension at OD 0.7 was stained by adding 3 µl of the dye mixture and incubated in the dark at RT for 15 min. Subsequently, 1 µl of the stained culture was added onto the center of a round cover glass (Warner Instruments), a 1 mm × 1 mm GCB agar pad was overlaid on the culture. The sample was then imaged by an Eclipse Ti Microscope with a 60x objective (Nikon). DIC, GFP and Texas Red fluorescence channels were applied to the imaging. ImageJ was used to analyze all images. Membrane integrity was assessed by calculating the ratio of

the number of green cells to red cells. Boiled cells were served as the negative control. The killing assays were done in 10 replicates.

Microcolony Formation Assay
100 µl of cell suspension at OD 0.7 were added to 1 ml of fresh GCB medium in a 6-well plate (VWR) and incubated at 37 °C with 5% CO_2 for 3 hours. Pictures were then taken using a 20x objective (Nikon).

SEM of Neisseria gonorrhoeae
30 µl of cell suspension at OD 0.7 was spotted onto 12 mm glass or Aclar film round coverslips coated with 0.1% Poly-lysin. A drop of fixative (100 µl of 2% glutaraldehyde, 4% formaldehyde in 0.1 M sodium cacodylate buffer pH 7.2) was added on top of the coverslip for 30 min. Additional fixative was then added to the dish containing the coverslips and left it in the fridge overnight. Samples were gently washed three times with buffer for 5 - 10 min each time and postfixed with osmium tetroxide 1% in 0.1 M sodium cacodylate buffer pH 7.2 for 1 hour at RT. After rinsing three times with buffer, samples were dehydrated in a graded series of ethanol concentrations 30%, 50%, 70%, 90% for 10 min each and three times in 100% ethanol with molecular sieves for 15 min each and critical point dried in a Tousimis Autosamdri 931. Samples were coated with 10 nm of iridium using a Leica ACE600 sputter coater. Imaging was done in a JEOL JSM-IT500HR at 5.0 kV.

Statistical Analysis
Statistical analysis was performed using Prism 10 (GraphPad). Group data are presented as means with SEM. The significance was determined using One-way ANOVA (and nonparametric or mixed). P values less than 0.05 were considered significant.

References

1.   G. World Health Organization, Prevention and control of sexually transmitted infections (STIs) in the era of oral pre-exposure prophylaxis (PrEP) for HIV. *Licence: CC BY-NC-SA 3.0 IGO.* (2019).
2.   S. St Cyr *et al.*, Update to CDC's Treatment Guidelines for Gonococcal Infection, 2020. *MMWR Morb Mortal Wkly Rep* **69**, 1911-1916 (2020).
3.   M. Unemo *et al.*, High-level cefixime- and ceftriaxone-resistant Neisseria gonorrhoeae in France: novel penA mosaic allele in a successful international clone causes treatment failure. *Antimicrob Agents Chemother* **56**, 1273-1280 (2012).
4.   M. Ohnishi *et al.*, Ceftriaxone-resistant Neisseria gonorrhoeae, Japan. *Emerg Infect Dis* **17**, 148-149 (2011).
5.   M. Day *et al.*, Detection of 10 cases of ceftriaxone-resistant Neisseria gonorrhoeae in the United Kingdom, December 2021 to June 2022. *Euro Surveill* **27** (2022).
6.   A. Derbie, D. Mekonnen, Y. Woldeamanuel, T. Abebe, Azithromycin resistant gonococci: a literature review. *Antimicrob Resist Infect Control* **9**, 138 (2020).
7.   P. Sawatzky *et al.*, Increasing Azithromycin Resistance in Neisseria gonorrhoeae Due to NG-MAST 12302 Clonal Spread in Canada, 2015 to 2018. *Antimicrob Agents Chemother* **66**, e0168821 (2022).
8.   B. Shaskolskiy, I. Kandinov, E. Dementieva, D. Gryadunov, Antibiotic Resistance in Neisseria gonorrhoeae: Challenges in Research and Treatment. *Microorganisms* **10** (2022).
9.   D. M. Richards, R. C. Heel, R. N. Brogden, T. M. Speight, G. S. Avery, Ceftriaxone. A review of its antibacterial activity, pharmacological properties and therapeutic use. *Drugs* **27**, 469-527 (1984).
10.   A. H. Bakheit, B. M. Al-Hadiya, A. A. Abd-Elgalil, Azithromycin. *Profiles Drug Subst Excip Relat Methodol* **39**, 1-40 (2014).
11.   D. J. Newman, G. M. Cragg, Natural Products as Sources of New Drugs over the Nearly Four Decades from 01/1981 to 09/2019. *J Nat Prod* **83**, 770-803 (2020).
12.   D. A. Dias, S. Urban, U. Roessner, A historical overview of natural products in drug discovery. *Metabolites* **2**, 303-336 (2012).
13.   K. Lewis, The Science of Antibiotic Discovery. *Cell* **181**, 29-45 (2020).
14.   K. E. Wilson *et al.*, Difficidin and oxydifficidin: novel broad spectrum antibacterial antibiotics produced by Bacillus subtilis. II. Isolation and physico-chemical characterization. *J Antibiot (Tokyo)* **40**, 1682-1691 (1987).
15.   S. B. Zimmerman *et al.*, Difficidin and oxydifficidin: novel broad spectrum antibacterial antibiotics produced by Bacillus subtilis. I. Production, taxonomy and antibacterial activity. *J Antibiot (Tokyo)* **40**, 1677-1681 (1987).
16.   M. S. Ngalimat *et al.*, A Review on the Biotechnological Applications of the Operational Group Bacillus amyloliquefaciens. *Microorganisms* **9** (2021).

17.　A. Koumoutsi *et al.*, Structural and functional characterization of gene clusters directing nonribosomal synthesis of
bioactive cyclic lipopeptides in Bacillus amyloliquefaciens strain FZB42. *J Bacteriol* **186**, 1084-1096 (2004).
18.　D. M. Richards, R. C. Heel, Ceftizoxime. A review of its antibacterial activity, pharmacokinetic properties and
therapeutic use. *Drugs* **29**, 281-329 (1985).
19.　J. J. Petkowski, W. Bains, S. Seager, Natural Products Containing 'Rare' Organophosphorus Functional Groups.
Molecules **24** (2019).
20.　Y. Cao, Q. Peng, S. Li, Z. Deng, J. Gao, The intriguing biology and chemistry of fosfomycin: the only marketed
phosphonate antibiotic. *RSC Adv* **9**, 42204-42218 (2019).
21.　F. E. Jen, J. M. Atack, J. L. Edwards, M. P. Jennings, Complete Genome Sequences of Seven Neisseria
gonorrhoeae Clinical Isolates from Mucosal and Disseminated Gonococcal Infections. *Microbiol Resour Announc*
10, e0073421 (2021).
22.　W. T. Doerrler, R. Sikdar, S. Kumar, L. A. Boughner, New functions for the ancient DedA membrane protein family.
J Bacteriol **195**, 3-11 (2013).
23.　B. Sit *et al.*, Undecaprenyl phosphate translocases confer conditional microbial fitness. *Nature* **613**, 721-728 (2023).
24.　I. J. Roney, D. Z. Rudner, The DedA superfamily member PetA is required for the transbilayer distribution of
phosphatidylethanolamine in bacterial membranes. *P Natl Acad Sci USA* **120** (2023).
25.　D. Huang *et al.*, TMEM41B acts as an ER scramblase required for lipoprotein biogenesis and lipid homeostasis.
Cell Metab **33**, 1655-+ (2021).
26.　Y. E. Li *et al.*, TMEM41B and VMP1 are scramblases and regulate the distribution of cholesterol and
phosphatidylserine. *J Cell Biol* **220** (2021).
27.　M. Diaconu *et al.*, Structural basis for the function of the ribosomal L7/12 stalk in factor binding and GTPase
activation. *Cell* **121**, 991-1004 (2005).
28.　M. A. Carlson *et al.*, Ribosomal protein L7/L12 is required for GTPase translation factors EF-G, RF3, and IF2 to
bind in their GTP state to 70S ribosomes. *Febs J* **284**, 1631-1643 (2017).
29.　M. M. Zweerink, A. Edison, Difficidin and oxydifficidin: novel broad spectrum antibacterial antibiotics produced by
Bacillus subtilis. III. Mode of action of difficidin. *J Antibiot (Tokyo)* **40**, 1692-1697 (1987).
30.　J. Lin, D. Zhou, T. A. Steitz, Y. S. Polikanov, M. G. Gagnon, Ribosome-Targeting Antibiotics: Modes of Action,
Mechanisms of Resistance, and Implications for Drug Design. *Annu Rev Biochem* **87**, 451-478 (2018).
31.　H. Paternoga *et al.*, Structural conservation of antibiotic interaction with ribosomes. *Nat Struct Mol Biol* **30**, 1380-
1392 (2023).
32.　Y. S. Polikanov, N. A. Aleksashin, B. Beckert, D. N. Wilson, The Mechanisms of Action of Ribosome-Targeting
Peptide Antibiotics. *Front Mol Biosci* **5**, 48 (2018).
33.　S. Arenz *et al.*, Structures of the orthosomycin antibiotics avilamycin and evernimicin in complex with the bacterial
70S ribosome. *Proc Natl Acad Sci U S A* **113**, 7527-7532 (2016).
34.　J. M. Harms *et al.*, Translational regulation via L11: molecular switches on the ribosome turned on and off by
thiostrepton and micrococcin. *Mol Cell* **30**, 26-38 (2008).
35.　L. Zhang *et al.*, Ribosome-targeting antibacterial agents: Advances, challenges, and opportunities. *Med Res Rev*
41, 1855-1889 (2021).
36.　J. Poehlsgaard, S. Douthwaite, The bacterial ribosome as a target for antibiotics. *Nature Reviews Microbiology* **3**,
870-881 (2005).
37.　N. E. Freitag, H. S. Seifert, M. Koomey, Characterization of the pilF-pilD pilus-assembly locus of Neisseria
gonorrhoeae. *Mol Microbiol* **16**, 575-586 (1995).
38.　D. Vallenet *et al.*, MicroScope: an integrated platform for the annotation and exploration of microbial gene functions
through genomic, pangenomic and metabolic comparative analysis. *Nucleic Acids Res* **48**, D579-D589 (2020).
39.　H. Todor, N. Herrera, C. Gross, Three bacterial DedA subfamilies with distinct functions and phylogenetic
distribution. *bioRxiv* 10.1101/2023.01.04.522824, 2023.2001.2004.522824 (2023).
40.　R. C. Edgar, Muscle5: High-accuracy alignment ensembles enable unbiased assessments of sequence homology
and phylogeny. *Nat Commun* **13**, 6968 (2022).
41.　M. N. Price, P. S. Dehal, A. P. Arkin, FastTree 2--approximately maximum-likelihood trees for large alignments.
PLoS One **5**, e9490 (2010).
42.　M. Ito, M. Nagane, Improvement of the electro-transformation efficiency of facultatively alkaliphilic Bacillus
pseudofirmus OF4 by high osmolarity and glycine treatment. *Biosci Biotechnol Biochem* **65**, 2773-2775 (2001).
43.　G. Q. Zhang *et al.*, Enhancing electro-transformation competency of recalcitrant Bacillus amyloliquefaciens by
combining cell-wall weakening and cell-membrane fluidity disturbing. *Anal Biochem* **409**, 130-137 (2011).
44.　J. P. Dillard, Genetic Manipulation of Neisseria gonorrhoeae. *Curr Protoc Microbiol* **Chapter 4**, Unit4A 2 (2011).
45.　A. B. Dalia, Natural Cotransformation and Multiplex Genome Editing by Natural Transformation (MuGENT) of Vibrio
cholerae. *Methods Mol Biol* **1839**, 53-64 (2018).
527
528

Figures and Tables

Figure 1. Oxydifficidin isomers inhibit the growth of *N. gonorrhoeae*. a. Discovery of a contaminant (*Bacillus amyloliquefaciens* BK) that inhibited the growth of *N. gonorrhoeae*. **b.** Example of known oxydifficidin thermal isomers. **c.** MIC of oxydifficidin against different bacteria (n = 2). Genome-based phylogenetic tree built using Genome Clustering of MicroScope.

Figure 2. Oxydifficidin-resistant *N. gonorrhoeae* mutant development and corresponding susceptibilities. a. Schematic representation of *N. gonorrhoeae* mutant development that identified *dedA*. Activity of different antibiotics against WT and *dedA* gene disrupted *N. gonorrhoeae*. **b.** Time-dependent antibiotic killing assay of *N. gonorrhoeae* strains. Each antibiotic was tested at 8x its MIC for the specific strain being examined (WT Ampicillin: 8 µg/ml; WT Oxydifficidin: 2 µg/ml; ΔdedA Oxydifficidin: 16 µg/ml). ΔdedA indicates *N. gonorrhoeae dedA* deletion mutant. (n = 3)

Figure 3. Oxydifficidin's anti-*N. gonorrhoeae* activity arises from a combination of DedA flippase assisted uptake and ribosomal protein L7/L12 (RplL) sensitivity. a. Structure of oxydifficidin compared to that of the known substrates for DedA homologs. **b.** Comparison of antibiotic accumulation in WT and *dedA* knockout cells. Blue and yellow highlighted sections represent independent experiments. (Oxydifficidin and Tetracycline: n = 2; Oxydifficidin and chloramphenicol: n = 3) **c.** *In vitro* coupled transcription/translation assay. The effect of oxydifficidin and other antibiotics on *in vitro* protein production using a coupled transcription/translation system was monitored by SDS-PAGE. Red dots indicate *in vitro* production of dihydrofolate reductase (18 kDa) from the *DHFR* gene. MW std: kDa molecular weight standard. **d.** *In vitro* transcription assay. Red dots indicate *in vitro* production of a 1704 bp RNA from the *FLuc* gene. A reaction containing 20 mM of EDTA was used as an inhibition control. MW std: bp molecular weight standard. **e.** Model explaining oxydifficidin's potent activity in *N. gonorrhoeae*. In this model DedA flips oxydifficidin across the inner membrane to assist its uptake and oxydifficidin then inhibits protein synthesis through either a direct or indirect interaction with L7/L12 (RplL). Two spontaneous mutations (K84E and R76C) in the RplL (L7/L12) protein were found to confer resistance to oxydifficidin. Image was generated by BioRender.

Table 1. Susceptibilities of *N. gonorrhoeae* to antibiotics.

Antibiotic	MIC (µg/ml)*	
	MS11	**H041**
Oxydifficidin	0.25	0.125
Ceftazidime	0.125	16
Ampicillin	1	8
Chloramphenicol	4	4
Rifampicin	0.25	0.25
Tetracycline	1	1
Nalidixic acid	16	16
Irgasan	0.5	0.5
Vancomycin	>64	>64
Polymyxin B	>64	>64
Melittin	2	2
Bacitracin	4	4
Daptomycin	>64	>64

*n=2

Table 2. Activity of antibiotics against *N. gonorrhoeae rplL* mutant.

Antibiotic	MIC (µg/ml)*	
	WT	RplL_R76C (Engineered)
Oxydifficidin	0.25	2
Ampicillin	1	1
Ceftazidime	0.125	0.125
Bacitracin	4	4
Ribosome-targeting antibiotic		
Chloramphenicol	4	4
Spectinomycin	16	16
Tetracycline	1	1
Erythromycin	0.5	0.5
Gentamicin	8	8
Avilamycin	4	4
Thiostrepton	0.125	0.125

*n=2

a

Bacillus subtilis 168
Bacillus amyloliquefaciens DSM 7
Bacillus amyloliquefaciens LL3
Bacillus amyloliquefaciens XH7
Bacillus amyloliquefaciens LFB112
Bacillus velezensis GA1
Bacillus amyloliquefaciens subsp. plantarum
Bacillus amyloliquefaciens subsp. plantarum
Bacillus amyloliquefaciens BK
Bacillus amyloliquefaciens FZB42
Bacillus amyloliquefaciens subsp. plantarum
Bacillus amyloliquefaciens subsp. plantarum

0.2

c

Strain	Gene Descriptor	Halted production
10A6	Wild type	None
10A7	Δ*bmyA*::EmR	Bacillomycin (mycosubtilin)
10A15	Δ*pks3KS1*::ermAM	Difficidin/Oxydifficidin

b

Transposon insertion site	Protein encoded
1	Phage SPbeta protein
2	Asp tRNA
3	Mycosubtilin synthase B
4	Transpeptidase penicillin -binding protein spoVD

WT	ATTTCCATT-AAATTGTCCC
Tn5 mutant	ATTTCCATTTAAATTGTCCC

Fig. S1. Mutagenesis results of oxydifficidin-producing Bacillus spp. a. Genome-based phylogenetic tree containing *Bacillus amyloliquefaciens* BK and closely related Bacillus spp.. The tree was built using Genome Clustering of MicroScope. **b.** Disc diffusion assay of a methanol extract from cultures of WT *Bacillus amyloliquefaciens* BK (WT) and a Tn5 mutant. The test lawn was *N. gonorrhoeae*. The table shows all transposon insertion sites in the Tn5 mutant strain. The Tn5 also contains a frame-shift mutation in the *difF* gene. The red box highlights the location of frame-shift mutation in the oxydifficidin BGC. **c.** Disc diffusion assay of a methanol extract from cultures of WT and BGC knockout strains of *Bacillus amyloliquefaciens* FZB42. The test lawn was *N. gonorrhoeae*. Strain genotypes are shown in the table. Red arrow indicates that only strain 10A15 no longer produce the anti-*N. gonorrhoeae* compound.

Fig. S2. Phylogenetic tree of 15,825 bacterial DedA family proteins. The tree was built by MUSCLE v5 and FastTree and visualized using iTOL. *N. gonorrhoeae* NGFG_RS04905 highlighted in red represents the DedA gene associated with the activity of oxydifficidin. *N. gonorrhoeae* NGFG_RS07420 and *N. gonorrhoeae* NGFG_RS00755 represents 2 other DedA family proteins in *N. gonorrhoeae*.

Fig. S3. Oxydifficidin does not lyse or depolarize the membrane of *N. gonorrhoeae*. a. Lysis assay using SYTOX green dye and 8x the MIC of each antibiotic. **b.** Depolarization assay using DiSC$_3$(5) dye and 8x the MIC of each antibiotic. **c.** Lysis assay using SYTOX green dye with 100x the MIC of oxydifficidin and 32x the MIC of nisin. **d.** Depolarization assay using DiSC$_3$(5) dye with 100x the MIC of oxydifficidin and the 32x the MIC of melittin. (n = 3 for all assays)

Fig. S4. Mutations in *dedA* affect cell morphology and pili functionality of *N. gonorrhoeae*. a. Membrane integrity assay of *N. gonorrhoeae* WT and *dedA* deletion mutant (Δ*dedA*) cells using SYTO 9 and propidium iodide. Cell integrity was assessed using the ratio of green-stained cell count to red-stained cell count. *: $p < 0.05$ **b.** Scanning electron microscope pictures of *N. gonorrhoeae* WT and Δ*dedA* cells. **c.** Micro-colony formation assay of *N. gonorrhoeae* WT and Δ*dedA* cells.

Fig. S5. ¹H-NMR spectrum of oxydifficidin (800 MHz, 298 K, CD₃OD - D₂O (1:1)).

Fig. S6. ¹³C-NMR spectrum of oxydifficidin (800 MHz, 298 K, CD₃OD - D₂O (1:1)).

Fig. S6. ^{13}C-NMR spectrum of oxydifficidin (800 MHz, 298 K, CD$_3$OD - D$_2$O (1:1)).

Fig. S7. edHSQC spectrum (800 MHz, 298 K, CD₃OD - D₂O (1:1)) of oxydifficidin.

Fig. S8. COSY spectrum (800 MHz, 298 K, CD₃OD - D₂O (1:1)) of oxydifficidin.

Fig. S9. ^{1}H-^{13}C HMBC spectrum (800 MHz, 298 K, CD$_3$OD - D$_2$O (1:1)) of oxydifficidin.

Fig. S10. Partial COSY (left) and ROESY (right) comparison and key ROESY correlations of oxydifficidin (800 MHz, 298 K, CD₃OD - D₂O (1:1)).

	Ion	Formula	m/z calc.	m/z obs.	RDB	Δ ppm
1	[M-H]⁻	$C_{31}H_{44}O_7P^-$	559.2819	559.2816	11.0	-0.64

Fig. S11a. Full HRMS and annotation of oxydifficidin [M–H]⁻ parental ion.

	Ion	Formula	m/z calc.	m/z obs.	RDB	Δ ppm
	[M-H]$^-$	C$_{31}$H$_{44}$O$_7$P$^-$	559.2830	559.2907	10.5	13.7
a	[X-H-H$_2$O]$^-$	C$_{30}$H$_{44}$O$_5$P$^-$	515.2932	515.3023	9.5	17.6
b	[X-H]$^-$	C$_{25}$H$_{35}$O$_6$P$^-$	463.2255	463.2328	8.5	15.8
c	[X-H-H$_2$O]$^-$	C$_{25}$H$_{34}$O$_5$P$^-$	445.2149	445.2214	9.5	14.5
d	[X-H]$^-$	H$_2$O$_4$P$^-$	96.9696	96.9683	0.5	13.9
e	[X-H]$^-$	O$_3$P$^-$	78.9591	78.9574	0.5	-21.5

Fig. S11b. -ESI MS/MS spectrum and fragment annotation of oxydifficidin [M–H]$^-$ ion.

Table S1. Activity of oxydifficidin against of Neisseria mutants.

Organism	MIC (μg/ml)*
N. gonorrhoeae MS11	0.25
N. gonorrhoeae MS11 Δ*dedA*	2
N. gonorrhoeae MS11 Δ*dedA rplL*_R76C	16
N. gonorrhoeae MS11 Δ*dedA rplL*_K84E	16
N. cinerea ATCC 14685	2
N. cinerea ATCC 14685 *rplL*_K84E	16
N. subflava NJ 9703	8
N. subflava NJ 9703 *dedA*_I59-D82Del	64
N. subflava NJ 9703 *dedA*_A78-K212Del	64
N. subflava NJ 9703 *dedA*_A160V	64
N. subflava NJ 9703 *dedA*_L53-K212Del	64
N. subflava NJ 9703 *dedA*_Q22*	64
N. subflava NJ 9703 *dedA*_G49S	64
N. subflava NJ 9703 *dedA*_C37*	64

Table S2. ^{1}H and ^{13}C NMR data of oxydifficidin (800 MHz, 298 K, CD$_3$OD - D$_2$O (1:1))

Position	Type	^{13}C		^{1}H	multiplicity, J (Hz)
1	C	174.1		-	
2	CH$_2$	42.3	a	3.22	d, 15.5
			b	2.93	d, 15.5
3	C	145.2		-	
3-CH$_2$	CH$_2$	113.9	a	4.97	s
			b	4.84	s
4	CH	47.2		2.50	m
4-CH$_3$	CH$_3$	16.5		1.16	d, 7.2
5	CH	75.7		4.31	*br* t, 3.5
6	CH	137.3		5.80	dd, 5.5; 15.0
7	CH	125.4		6.75	dd, 11.6; 14.9
8	CH	129.3		6.02	m
9	CH	125.4		6.39	t, 11.4
10	CH	125.4		6.47	t, 11.4
11	CH	134.2		5.41	dt, 5.9; 10.0; 10.0
12	CH$_2$	26.7	a	2.64	m
			b	2.04	m
13	CH$_2$	24.8	a	1.67	m
			b	1.32	m
14	CH$_2$	34.2	a	1.71	br
			b	1.42	m
15	CH	73.1		5.22	dt, 6.3; 8.2; 8.2
16	C	141.8		-	-
16-CH$_3$	CH$_3$	18.9		1.81	s
17	CH	121.7		6.02	m
18	CH	128.0		6.33	t, 11.5
19	CH	124.8		5.16	dt, 5.3; 11.0; 11.0
20	CH$_2$	31.5	a	2.51	m
			b	2.25	dt, 4.0; 4.0; 14.5
21	CH	75.4		4.86	m
22	CH$_2$	31.7	a	1.63	m
			b	1.50	m
23	CH$_2$	36.6		2.00	m
24	C	139.2		-	-
24-CH$_3$	CH$_3$	16.7		1.72	s
25	CH	126.9		5.77	d, 8.2
26	CH	134.5		6.56	dt, 10.3; 10.3; 16.7
27	CH$_2$	115.5	a	5.05	dd, 1.7; 16.7
			b	4.97	dd. 1.7, 10.3

^{13}C NMR chemical shifts were obtained by the interpretation of ^{13}C, HSQC and HMBC experiments.

Table S3. Primer Sequences used in this study.

Amplicon	Primer	Sequence (5'-3')
dedA 5' overhang	F	CCCTTTCTGCCTGTACTTCGACTCAAG
	R	CATAAAGTGTCAAGCCCTCGAGGGGTTTTCCAAAACACAATGTCGAGG
dedA 3' overhang	F	GCCGTCTGAAGTTTAAACATCGATTTGTTGGAAATTGACATTATGAATATATTATCCG
	R	CTTGGTAATCGCGCAACAGATCTTCAAGC
trpB-lga 5' overhang	F	AACGCCATCGGTTTGTTCTATC
	R	TAAAGTGTCAAGCCCTCGAGGAGTCAAGCTTCGGACGGCATTTT
trpB-lga 3' overhang	F	GCCGTCTGAAGTTTAAACATCGTTCAGACGGCATTTTATTTTGC
	R	CTTGAGAAGCCGGTTACAAACG
*rplL*_R76C	F	GTGCAACTGGGAAACAATCACA
	R	GTCTTTTTAGGTTACCGCGCTG
kan^R cassette	F	CTCGAGGGCTTGACACTTTATG
	R	ATCGATGTTTAAACTTCAGACGGC