Horizontal gene transfer of a functional *cki* homolog in the human pathogen *Schistosoma*

mansoni

George Wendt, James J. Collins III*

Department of Pharmacology, University of Texas Southwestern Medical Center

* To whom correspondence should be addressed

Abstract

Schistosomes are parasitic flatworms responsible for the neglected tropical disease

schistosomiasis, causing devastating morbidity and mortality in the developing world. The

parasites are protected by a skin-like tegument, and maintenance of this tegument is controlled

by a schistosome ortholog of the tumor suppressor TP53. To understand mechanistically how

p53-1 controls tegument production, we identified a cyclin dependent kinase inhibitor homolog

(*cki*) that was co-expressed with *p53-1*. RNA interference of *cki* resulted in a hyperproliferation

phenotype, that, in combination with *p53-1* RNA interference yielded abundant tumor-like growths,

indicating that *cki* and *p53-1* are bona fide tumor suppressors in *Schistosoma mansoni*.

Interestingly, *cki* homologs are widely present throughout parasitic flatworms but evidently absent

from their free-living ancestors, suggesting this cki homolog came from an ancient horizontal gene

transfer event. This in turn implies that the evolution of parasitism in flatworms may have been

aided by a highly unusual means of metazoan genetic inheritance.

Introduction

Parasitic flatworms are a diverse clade of organisms that include pathogens of great medical and

agricultural purposes. Interestingly, all parasitic flatworms come from a single speciation event,

and therefore are a derived clade relative to all free-living flatworms[1]. As such, common traits

shared across parasitic flatworms could represent early adaptations that facilitated a free-living

ancestor in their transition to parasitic lifestyle. One such common trait is the presence of a

tegument, a syncytial skin-like tissue that acts as the interface between the host and the parasite.

The tegument is thought to be essential for parasitic flatworms, facilitating processes such as

immune evasion, nutrient acquisition, and protection from physical and chemical insults[2-4].

The most medically important parasitic flatworms are the blood-dwelling parasites known as

schistosomes. These parasites infect over 200 million of the world's poorest people, killing tens

of thousands every year[5]. They reside in the vasculature of the host, where they can survive

for decades, laying eggs that ultimately cause the pathology of schistosome infection[6]. The

ability of the parasite to thrive within their host's bloodstream for decades on end is thought to be

modulated by the parasite's stem cells and their tegument. Previous work has shown that in adult

worms, stem cells are almost entirely dedicated to maintaining the schistosome's tegument[7, 8].

This tegument maintenance is regulated by a number of different factors including flatworm-

specific transcription factors[8] and a parasite ortholog of *TP53* termed *Sman-p53-1* (henceforth

p53-1 for brevity, gene symbol Smp_139530)[9].

Despite being most widely recognized as a tumor suppressor, *TP53* homologs appear to play a

different role in flatworms. Most free-living flatworms have a single *TP53* ortholog that is essential

for the maintenance of a sub-population of their stem cells responsible for producing the worm's

epidermis[10, 11]. Schistosomes have two TP53 homologs: a TP53 ortholog (*p53-1*) that is

responsible for maintaining the stem cells responsible for producing the parasite's tegument and

a second p53 molecule (*p53-2*, gene symbol Smp_136160) that is paralogous to *p53-1* that is

only found in parasitic flatworms. Interestingly, the schistosome *p53-2* is required for the

parasite's response to genotoxic stress[9]. Mechanistically, it is unclear how either of these TP53

homologs function.

One of the most well characterized mechanisms of TP53 function involves its regulation of the

expression of the cyclin dependent kinase inhibitor (CKI) *P21/CDKN1A[12]*. In response to insults

such as genotoxic stress, TP53 induces the expression of P21. P21 then binds to and inhibits

cyclin dependent kinase (CDK)/cyclin complexes, resulting in the hypophosphorylation of RB,

preventing progression through the G1-S checkpoint[12, 13]. While direct regulation of CKIs by

P53 homologs is best characterized in vertebrates[14, 15], the interactions between CKIs, CDKs,

cyclins, and RB homologs seems to be broadly conserved across all metazoans[16, 17].

In this study, we attempted to understand how *p53-1* regulates tegument production in

schistosomes. In doing so, we identified a cyclin dependent kinase inhibitor that we termed *Sman-*

cki (henceforth *cki* for brevity, gene symbol Smp_199050). Functional studies of *cki* showed that

it is required for normal stem cell proliferation and differentiation. We also show that *cki* and *p53-*

1 cooperate as tumor suppressors in *Schistosoma mansoni*. Interestingly, *cki* homologs were

present in almost every parasitic flatworm genome examined, but entirely absent from free-living

flatworms. Given the evolutionary relationship between free-living and parasitic flatworms, this

makes horizontal gene transfer (HGT) the likely origin of *cki* in parasitic flatworms.

Results

Having previously found that *p53-1*, the schistosome homolog of TP53, plays an important role in

tegument development[9], we sought to identify the targets of this transcription factor to

understand how *p53-1* controls tegument development. One common mechanism by which TP53

homologs function across different animals is by inducing the expression of cyclin dependent

kinase inhibitors such as *CDKN1A[15, 18]*. We identified a distant homolog of *CDKN1A* in the

Schistosoma mansoni genome (*Sman-cki*, henceforth *cki*) that was expressed in many of the

same cells as *p53-1*, including tegument progenitor cells and stem cells, suggesting it could play

a role in regulating tegument production (**Figure 1A-C**). Though this homolog did not possess an

annotated cyclin dependent kinase inhibitor (CDI) domain (InterPro entries

IPR044898/IPR003175[19]), it did bear high similarity and identity to other CDI-domain containing

proteins such as Drosophila Dap and human CDKN1B (**Figure S1A-C**). Additionally, the

AlphaFold[20] predicted structure of the n-terminus of CKI conformed to the structure of CDKN1B

when superimposed on the solved structure of CDK4/CCND1/CDKN1B[21] (**Figure S1C**).

We next tested the function of *cki* in the parasite. In other organisms, *cki* homologs act as tumor

suppressors in part by binding to and inhibiting the cyclin/cdk complexes that control the cell cycle,

halting proliferation[22]. As such, loss of function of *cki* homologs results in hyperproliferation in a

variety of animals in a variety of contexts[16, 17, 23-26]. When we knockdown *cki* using RNAi, we

similarly observe a hyperproliferation phenotype (**Figure 2, Figure S2A-B**). While there is nearly

100% penetrance of the hyperproliferation phenotype in the parasite's head (**Figure 2, top**), there

is variable penetrance of the phenotype in the body of the worm (**Figure 2, bottom**). This increase

in proliferation is also accompanied by a loss of production of *tsp-2*$^+$ tegument progenitor cells

(**Figure 2, Figure S2A-B**) as well as new tegument cells (**Figure S2C-D**), demonstrating that the

increase in proliferation is accompanied by a loss of differentiation. Additionally, we see an

approximately five-fold increase in apoptotic cells in the heads of *cki(RNAi)* parasites (**Figure

S2E-F]**). Together, this suggests that *cki* RNAi causes stem cells to engage in hyperproliferation

at the expense of differentiation, ultimately resulting in stem cell death.

Human *cki* homologs are widely recognized as tumor suppressors[22, 27], but no studies of

invertebrate *cki* homologs have demonstrated spontaneous tumor formation in *cki* loss-of-function

models[16, 23-25]. Unlike invertebrate models tested to date, schistosomes have lifespans

measured in decades[6] and rely upon proliferative somatic cells throughout these long lives[28].

As such, we wondered whether *cki* could function as a tumor suppressor in schistosomes.

Knockdown of *cki* alone causes hyperproliferation (**Figure 2**), but never results in robust

tumorigenesis. Given our original model that *cki* expression might be induced by *P53-1* (**Figure

3A**), we next wondered whether knocking down *p53-1* in combination with *cki* would create a

more permissive environment for tumor formation in the parasite. We found that knocking down

p53-1 in conjunction with *cki* does in fact result in the appearance of abnormal proliferative tumor-

like masses in the parasite's head with nearly 100% penetrance (**Figure 3B**). This suggests that

cki and *p53-1* cooperate as tumor suppressors in *Schistosoma mansoni* (**Figure 3C**).

Interestingly, we were able to readily identify *cki* homologs in nearly all parasitic flatworm genomes

(present in 43 out of 47 examined) (**Table S1**) but were unable to find any *cki* homologs in any

free-living flatworm genomes (present in 0 out of 19 genomes examined across 9 taxonomic

orders) (**Figure 4A**). This is surprising because phylogenetic analysis suggests that all parasitic

flatworms arose from a common free-living ancestor approximately 200 million years ago[1]. As

such, any gene found broadly across parasitic clades should be present at some frequency in the

extant free-living flatworms. The fact that we do not observe this suggests either 1) the parasite

cki homologs arose from de novo gene birth, 2) *cki* homologs were present broadly throughout all

flatworms but were independently lost in all extant free-living flatworm lineages that we have

genomes for, or 3) the parasite *cki* homolog was not inherited "vertically" and therefore may have

originated from horizontal gene transfer in an ancestral parasitic flatworm. Given that parasite *cki*

homologs have sequence-level similarity to other metazoan *cki* homologs (**Figure S1A-B**), de

novo gene birth can be ruled out. While we cannot rule out multiple independent losses of *cki*

homologs across extant free-living flatworms, it would be incredibly unlikely that *cki* homologs

were independently lost in 19 different animals across 9 taxonomic orders, leaving HGT as a

plausible explanation for the origin of parasitic flatworm *cki*.

To search for evidence of a more typical (i.e., 'vertical') inheritance of *cki*, we decided to examine

sequence changes across evolution in genes that are likely to have co-evolved with *cki*. In this

case, we chose to examine cyclin dependent kinase 4 (*cdk4*) and cyclin d (*ccnd*) because they

interact with each other and with *cki* homologs across diverse metazoans[17, 29]. They are also

all present in the three groups of animals we chose to examine: parasitic flatworms, the relatively

closely related molluscs, and the outgroup bony fish (**Figure 4A**). We reasoned that phylogenetic

analysis of the sequences of 'vertically' inherited genes should recapitulate the phylogeny as we

know it to exist (that is to say, our bootstrap analysis should indicate that parasitic flatworm genes

are more similar to the closely related mollusc genes than they are to their bony fish counterparts).

This is indeed what we see when we analyze the sequences of CDK4 and CCND homologs, but

examination of CKI homologs is unable to robustly identify the correct phylogeny (**Figure 4B,**

Figures S3-S7). Combined with the phylogenetic relationship between free-living and parasitic

flatworms, this further supports HGT as the likely origin of *cki* in parasitic flatworms.

Discussion

Tumor suppressors such as TP53 and CDKN1A are some of the most widely studied molecules

on the planet, but their functions in animals that do not generally develop tumors are largely

unexplored. Here, we show that schistosomes have a *CDKN1A* homolog, *cki*, that acts as a tumor

suppressor together with the parasite's *TP53* ortholog, *p53-1*. Knocking down this *cki* homolog

results in a dramatic increase in stem cell proliferation with a concomitant decrease in

differentiation. Knocking down *cki* in combination with *p53-1* results in spontaneous and robust

formation of abnormal proliferative masses of cells, making *cki* and *p53-1* bona fide tumor

suppressors in schistosomes. Further studies of these tumors could better describe their

molecular identity and their cell of origin.

One exciting implication of this study is that schistosomes (and likely other parasitic flatworms)

possess a functional *cki* homolog, implying that cyclin-dependent kinases may be important in

regulating cellular proliferation in parasitic flatworms. One potential new area for schistosomacidal

drug development is the targeting of the parasite's stem cells. Previous studies have shown that

disrupting schistosome tissues maintained by stem cells is capable of injuring parasites[8] or

preventing disease pathology[30]. One of the greatest challenges in targeting parasite stem cells,

however, is identifying a way to specifically target proliferative cells in the parasite while sparing

proliferating cells in the host. Future work determining exactly which CDKs are involved in

regulating stem cells in schistosomes could reveal novel drug targets that could have good

therapeutic properties; CDK4/6 inhibitors, for example, are a relatively new family of anti-tumor

drugs that are generally well tolerated by patients[31, 32]. Schistosome CDK4 is relatively

divergent compared to human CDK4. The schistosome CDK4 kinase domain has 41.2%/63.9%

identity/similarity to human CDK4, comparable to the difference between human CDK4 and

human CDK1 (43.4%/62.0%), and CDK4 inhibitors are generally very specific (the CDK4/6

inhibitor Palbociclib has >1000 fold selectivity for CDK4 over CDK1[33]), suggesting that

schistosome CDK4 has potential to be a selective drug target. Further studies of *cki* and its

interacting molecules could therefore have significant therapeutic implications for combatting

schistosomiasis.

Perhaps most interestingly, one of the most likely origins of the parasite's *cki* homolog is horizontal

gene transfer. Given that parasitic flatworms all derive from a common free-living ancestor[1],

anything ubiquitous in parasitic flatworms should be present at some frequency in extant free-

living flatworms. That *cki* is absent from free-living flatworms led us to 3 very unlikely explanations

for its origin. First, we believe we can safely rule out de novo gene birth given that parasite *cki* is

related to other metazoan *cki* homologs on the sequence level (**Figure S1A-B**). That leaves us

with either 1) a situation where *cki* homologs were broadly present throughout all flatworms, but

then sometime after parasitic flatworms diverged 200 million years ago, the ancestors of all extant

free-living flatworm lineages lost their *cki* homologs while parasitic flatworms retained theirs or 2)

a situation where an early ancestral parasitic flatworm acquired a *cki* homolog via horizontal gene

transfer. Definitive proof of horizontal gene transfer would require us to be able to identify the non-

flatworm animal that has a *cki* homolog with sequence similarity to the flatworm molecule.

Unfortunately, we have been unable to find such an animal, possibly because this animal would

have had to been alive approximately 200 million years ago (in order to transmit its *cki* homolog

to the ancestral parasitic flatworm) and may not have survived to create modern descendants. As

such, rather than stating that parasitic flatworm *cki* came from horizontal gene transfer, we instead

propose that horizontal gene transfer is one of the most plausible origins for this *cki*.

Further complicating the assumption that parasite *cki* came from horizontal gene transfer, HGT in

metazoans, especially between metazoans, is relatively rare[34, 35]. Still, HGT in parasitic

flatworms might be more common than in other metazoans. HGT between multicellular organisms

has been frequently identified between parasitic plants and their host plants[36-39]. Aside from

the analogous relationship between parasitic flatworms and their metazoan hosts, parasitic

flatworms may represent especially molecularly fertile ground for HGT. In addition to their syncytial

tegument, all parasitic flatworms are also united by their loss of the piRNA pathway[40] that,

combined with apparent the lack of DNA methylation across flatworms[41-43], makes the parasitic

flatworms deficient in two key pathways used to suppress mobile selfish genetic elements like

transposons and endogenous retroviruses[34, 44, 45]. Indeed, this type of HGT has been

reported between parasitic lamprey and their host[46], both of which have intact defenses against

selfish genetic elements. Selfish genetic element mediated HGT could therefore be unusually

successful in parasitic flatworms.

Origin of *cki* gene notwithstanding, the fact that it is nearly ubiquitous across all parasitic flatworms

means that it has been present in parasitic flatworms from early in their evolution and has likely

been retained because it plays an important role in these animals. Understanding that role could

teach us not only about the evolution of parasitism in these fascinating animals but could also

reveal ways to combat the diseases that they cause.

Figure Legends

Figure 1. Expression pattern of *cki.* A) Uniform manifold approximation plots showing expression

patterns of *p53-1* and *cki* in adult schistosomes. Red indicates high expression, orange indicates

medium expression, and blue indicates low/no expression. B) Colorimetric WISH showing

expression patterns of *cki* in male (left) and female (right) worms. H, head; T, testes; O, ovary; V,

vitellaria. C) Double FISH experiment showing expression of *cki* relative to the tegument

progenitor marker *tsp-2*, the proliferative cell marker *h2b*, and *p53-1*. Scale bars: (B) 500 µm, (C)

5 µm.

Figure 2. *cki* RNAi results in an increase in proliferation at the expense of differentiation. FISH

experiment in conjunction with EdU detection showing expression pattern of the tegument

progenitor marker *tsp-2* (green) in conjunction with the presence of proliferative EdU$^+$ cells

(yellow) under the indicated RNAi conditions. The fraction indicates the number of worms that are

similar to the representative image. Data are from 3 biological replicates. Scale bar: 50 µm.

Figure 3. *cki* and *p53-1* cooperate as tumor suppressors in *Schistosoma mansoni*. (A) Cartoon

depicting model of *p53-1* and *cki* interaction in normal parasites. In proliferative cells, P53-1

activity is low and *cki* is not transcribed, resulting in active cyclin/CDK complexes, promoting

proliferation. When proliferative cells commit to differentiating, P53-1 activity is high and *cki* is

transcribed, resulting in inactive cyclin/CDK/CKI complexes, promoting differentiation. (B) EdU

labelled neoblasts in parasites treated witih control RNAi, *p53-1* RNAi, *cki* RNAi, or a combination

of *p53-1* and *cki* RNAi. (C) Cartoon depicting consequences of *cki* RNAi. By knocking down *cki*,

cyclin/CDK complexes are inappropriately active, leading to increased proliferation and

decreased differentiation. (D) Cartoon depicting consequences of knocking down *cki* and *p53-1*

simultaneously. Knocking down *cki* directly depletes CKI protein. Knocking down *p53-1* decreases

P53-1-dependent induction of *cki* expression. This combination may lower CKI levels low enough

that proliferating cells become abnormal and form tumors. The fraction indicates the number of

worms that are similar to the representative image. Scale bar: 50 µm.

Figure 4. Phylogeny of CKI. (A) Top. Cladogram of *cki* homologs across select metazoans.

Bottom. Chart indicating the presence or absence of *cdk4*, *ccnd*, and *cki* homologs in the above

species. Green circles indicate presence and red x's indicate absence. (B) Simplification of

phylogenetic analysis of aligned sequences of CDK4, CCND, and CKI homologs across bony

fish, molluscs, and parasitic flatworms. Circle with the number indicates the bootstrap value of the

bifurcation between molluscs and parasitic flatworms. The color of the circle indicates relative

confidence of tree structure, where green is high and red is low.

Figure S1. Sman-CKI protein domains/structure. (A) Clustal alignment of CKI sequences from a

variety of parasitic flatworms as well as human CDKN1B. Dashed lines indicate parts of

alignments that fall within a CDI domain (InterPro IPR044898). The number at top of alignment

corresponds to position in the alignment relative to the *Schistosoma mansoni* CKI. Protein

sequences used for alignment and species abbreviations can be found in **Table S2** (B) Schematic

of the domain structure of CKI homologs from *S. mansoni*, *E. granulosus*, *D. melanogaster*, and

H. sapiens along with the percentage identity/similarity relative to the *S. mansoni* CKI homolog.

(C) Superimposition of the AlphaFold indicated residues of *S. mansoni* CKI (accession:

AF_A0A3Q0KUT0-F1-model_v4) onto the solved structure of *H. sapiens* CDKN1B, CCND1, and

CDK4 (PDB accession: 6p8e). Hash-shaded portion from (B) corresponds to the superimposition

in (C).

Figure S2. *cki* RNAi results in an increase in proliferation at the expense of differentiation. (A)

qPCR showing expression of *cki*, the stem cell marker *nanos2*, and the tegument progenitor

marker *tsp-2* under indicated RNAi conditions. 'cki(RNAi)(alt)' indicates knockdown performed

with dsRNA corresponding to a non-overlapping sequence relative to the original dsRNA

sequence. (B) FISH experiment in conjunction with EdU detection showing expression pattern of

the tegument progenitor marker *tsp-2* (green) in conjunction with the presence of proliferative

EdU$^+$ cells (yellow) under the indicated RNAi conditions. Note that the micrographs for

control(RNAi) are duplicates of the micrographs for control(RNAi) from Figure 3 (top) as these

came from the same experiment. (C) FISH of the tegument marker *calp* after control or *cki* RNAi.

Neoblast progeny are labeled with the EdU via a pulse-chase experiment. (D) Quantification of

(C). (E) Fluorescent TUNEL assay labeling DNA breaks in worms under the indicated RNAi

conditions. T, testes. Scale bar: 50 µm. Note signal in the testes is a normal occurrence in the

parasite's proliferative germline. (F) Quantification of (E). Scale bars: (B) 50 µm, (C) 10 µm. (E)

50 µm.

Figure S3. Phylogeny of CDK4 excluding free-living flatworms. Phylogenetic tree based on

analysis of aligned protein sequences of CDK4 homologs across a sampling of bony fish,

molluscs, and parasitic flatworms. The bifurcation point of molluscs and parasitic flatworms is

indicated with a circle. The tree is arbitrarily rooted to *D. rerio*. Species name abbreviations and

protein sequences are available in **Table S3**. Numbers indicate bootstrap values. Scale bar: 1

substitution/site.

Figure S4. Phylogeny of CDK4 including free-living flatworms. Phylogenetic tree based on

analysis of aligned protein sequences of CDK4 homologs across a sampling of bony fish,

molluscs, free-living, and parasitic flatworms. Numbers indicate bootstrap values. The tree is

arbitrarily rooted to *D. rerio*. Species name abbreviations and protein sequences are available in

Table S4. Numbers indicate bootstrap values. Scale bar: 1 substitution/site.

Figure S5. Phylogeny of CCND excluding free-living flatworms. Phylogenetic tree based on

analysis of aligned protein sequences of CCND homologs across a sampling of bony fish,

molluscs, and parasitic flatworms. The bifurcation point of molluscs and parasitic flatworms is

indicated with a circle. Numbers indicate bootstrap values. The tree is arbitrarily rooted to *D. rerio*.

Species name abbreviations and protein sequences are available in **Table S5**. Numbers indicate

bootstrap values. Scale bar: 1 substitution/site.

Figure S6. Phylogeny of CCND including free-living flatworms. Phylogenetic tree based on

analysis of aligned protein sequences of CCND homologs across a sampling of bony fish,

molluscs, and parasitic flatworms. The bifurcation point of molluscs and parasitic flatworms is

indicated with a circle. Numbers indicate bootstrap values. The tree is arbitrarily rooted to *D. rerio*.

Species name abbreviations and protein sequences are available in **Table S6**. Numbers indicate

bootstrap values. Scale bar: 1 substitution/site.

Figure S7. Phylogeny of CKI. Phylogenetic tree based on analysis of aligned protein sequences

of CKI homologs across a sampling of bony fish, molluscs, and parasitic flatworms. The bifurcation

point of molluscs and parasitic flatworms is indicated with a circle. Numbers indicate bootstrap

values. The tree is arbitrarily rooted to *D. rerio*. Species name abbreviations and protein

sequences are available in **Table S7**. Numbers indicate bootstrap values. Scale bar: 1

substitution/site.

Materials and Methods

Phylogenetic analysis

Identification of orthologs of *Schistosoma mansoni cki* homologs was carried out using Wormbase

Parasite (https://parasite.wormbase.org/)(version 19.0 (March 2024))[47, 48]. Examination of

Sman-cki (Smp_199050) homologs initially only revealed *cki* homologs in trematodes (present in

27/28 unique genomes) and monogeneans (present in 1/3 unique genomes). A reciprocal BLAST

search of the *Hymenolepis diminuta* genome identified the *cki* homolog WMSIL1_LOCUS6166

("CDI domain-containing protein"). Wormbase Parasite then identified 12 orthologs of

WMSIL1_LOCUS6166 in 11 unique cestode genomes, each of which was confirmed to be a *cki*

ortholog via a reciprocal BLAST search. Additional BLAST searches of cestode genomes

revealed 4 additional *cki* homologs in cestodes (in total present in 15/16 unique genomes), with

two being incorrectly included in the gene model of TIMM21 homologs (see **Table S2**). PlanMine

v3.0 (https://planmine.mpinat.mpg.de/planmine/)[49] was utilized in order to search 19 free-living

flatworm genomes for *cki* homologs. Domain structure of CKI homologs was obtained using

SMART v9.0 (http://smart.embl-heidelberg.de/)[50, 51].

Clustal alignment of CKI sequences was performed using Clustal Omega Multiple Sequence

Alignment[52] with default settings except ORDER was set to 'input'[53, 54]. Protein sequences

used for alignment and species abbreviations can be found in **Table S2**.

Phylogenetic trees were generated from CKI protein sequences by first using Clustal Omega[52]

to perform multiple sequence alignments using default settings. Aligned sequences were then

analyzed using RAxML (Version 8.2.12[55, 56]]) via raxmlGUI (version 2.0.10[57]) with the

following parameters:

-T 10 -f a -x 756002 -p 756002 -N 1000 -m PROTGAMMAJTTF -k -O (for analysis of CKI

homologs),

-T 10 -f a -x 756002 -p 756002 -N 1000 -m PROTGAMMAIJTTF -k -O (for analysis of CDK4

homologs both including and excluding free-living flatworms)

-T 10 -f a -x 296744 -p 296744 -N 1000 -m PROTGAMMAIJTTF -k -O (for analysis of CCND

homologs including free-living flatworms)

-T 10 -f a -x 756002 -p 756002 -N 1000 -m PROTGAMMAIJTTF -k -O (for analysis of CCND

homologs excluding free living flatworms)

The best model was determined using ModelTest-NG (version 0.1.7[58]). The best maximum-

likelihood tree was visualized using Interactive Tree of Life[59] and rooted to *Danio rerio* for visual

consistency.

All sequences used for phylogenetic analysis are available in **Tables S3-S7**.

Parasite acquisition and culture

Adult S. mansoni (NMRI strain, 6–7 weeks post-infection) were obtained from infected female

mice by hepatic portal vein perfusion with 37°C DMEM (Sigma-Aldrich, St. Louis, MO) plus 10%

Serum (either Fetal Calf Serum or Horse Serum) and heparin. Parasites were cultured as

previously described[7]. Unless otherwise noted, all experiments were performed with male

parasites in order to maximize the amount of somatic tissue present and to avoid additional

experimental modifications that have to be undertaken for successful in vitro female parasite

culture[60]. Experiments with and care of vertebrate animals were performed in accordance with

protocols approved by the Institutional Animal Care and Use Committee (IACUC) of UT

Southwestern Medical Center (approval APN: 2017-102092).

Labeling and imaging

Schistosome colorimetric and fluorescence in situ hybridization analyses were performed as

previously described[7-9, 28] with the following modification. To improve signal-to-noise for

colorimetric and fluorescence in situ hybridization, worms were incubated at 95°C for 15m in

10mM sodium citrate, pH6.0 prior to proteinase K digestion. *in vitro* EdU labeling and detection

was performed as previously described[61]. TUNEL labeling was carried out as follows:

formaldehyde-fixed worms stored in 100% MeOH were rehydrated in 50% MeOH/50% PBS +

0.3% TritonX-100 pH7.4 (PBSTx), then rinsed in PBSTx. Parasites were then photobleached as

described previously[7]. After photobleaching, parasites were rinsed in PBSTx, then incubated at

95°C for 15m in 10mM sodium citrate pH6.0. Following this, parasites were rinsed in PBSTx then

permeablized in 5 μg/ml proteinase K (Ambion AM2546). Following proteinase K treatment,

parasites were post-fixed in 4% formaldehyde in PBSTx. Parasites were then rinsed in PBSTx

then incubated at 37°C in TdT reaction buffer from Click-iT™ Plus TUNEL Assay for In Situ

Apoptosis Detection kit (Thermo Fisher Scientific C10617) for one hour. Parasites were then

incubated at 37°C in TdT reaction mix for one hour per manufacturer's instructions. Parasites

were then thoroughly rinsed and EdU detection was carried out as previously described[7]. All

fluorescently labeled parasites were counterstained with DAPI (1 μg/ml), cleared in 80% glycerol,

and mounted on slides with Vectashield (Vector Laboratories).

Confocal imaging of fluorescently labeled samples was performed on a Zeiss LSM900 Laser

Scanning Confocal Microscope. Unless otherwise mentioned, all fluorescence images represent

maximum intensity projection plots. To perform cell counts, cells were manually counted in

maximum intensity projection plots derived from confocal stacks. Brightfield images were acquired

on a Zeiss AxioZoom V16 equipped with a transmitted light base and a Zeiss AxioCam 105 Color

camera.

RNA interference

Schistosome RNAi: All RNAi experiments utilized freshly perfused male parasites between 6-7

weeks post-infection (separated from females). dsRNA treatments were all carried out at 30 µg/ml

in Basch Media 169. Double knockdown experiments were carried out at 60 µg/ml (30 µg/ml of

each dsRNA) for the first 4 days, then switched to 30 µg/ml (15 µg/ml of each dsRNA) for the

remainder of the experiment. dsRNA was generated by in vitro transcription and was replaced

daily for 3 days and then every other day until the end of the experiment (14 days for all

experiments except for EdU pulse-chase experiments, which were EdU pulsed at day 14 then

carried out for 8 more days). EdU pulses were performed at 5µM for 4 hours before either fixation

or chase as previously described.

As a negative control for RNAi experiments, we used a non-specific dsRNA containing two

bacterial genes(48). cDNAs used for RNAi and in situ hybridization analyses were cloned as

previously described(48); oligonucleotide primer sequences are listed in **Table S8**.

Statistical analysis

Statistical analysis was carried out using one-way ANOVA tests (qPCR experiments) or two-way

ANOVA with Tukey's multiple comparisons tests (all other experiments). All statistical tests were

performed using GraphPad Prism v10.2.3.

Acknowledgments

Schistosome infected mice and *B. glabrata* snails were provided by the National Institute of Allergy and Infectious Diseases Schistosomiasis Resource Center of the Biomedical Research Institute (Rockville, MD, USA) through National Institutes of Health National Institute of Allergy and Infectious Diseases Contract HHSN272201700014I for distribution through BEI Resources.

Funding: This work was supported by the National Institutes of Health R01AI121037 (J.J.C.) and R01AI150776(J.J.C.), as well as the Welch Foundation I-1948-20180324 (J.J.C.).

References

1. Laumer, C.E., A. Hejnol, and G. Giribet, *Nuclear genomic signals of the 'microturbellarian' roots of platyhelminth evolutionary innovation.* Elife, 2015. **4**.

2. Alan Wilson, R. and M.K. Jones, *Fifty years of the schistosome tegument: discoveries, controversies, and outstanding questions.* Int J Parasitol, 2021.

3. Fairweather, I., L.T. Threadgold, and R.E. Hanna, *Development of Fasciola hepatica in the mammalian host.* Fasciolosis, 1999. **1**: p. 47-111.

4. Brehm, K. and U. Koziol, *Echinococcus-Host Interactions at Cellular and Molecular Levels.* Adv Parasitol, 2017. **95**: p. 147-212.

5. Colley, D.G., et al., *Human schistosomiasis.* Lancet, 2014. **383**(9936): p. 2253-64.

6. Basch, P.F., *Schistosomes: Development, Reproduction, and Host Relations.* 1991, New York: Oxford University Press. 248.

7. Collins, J.J., III, et al., *Stem cell progeny contribute to the schistosome host-parasite interface.* Elife, 2016. **5**.

8. Wendt, G.R., et al., *Flatworm-specific transcriptional regulators promote the specification of tegumental progenitors in Schistosoma mansoni.* Elife, 2018. **7**.

9. Wendt, G.R., et al., *Convergent evolution of a genotoxic stress response in a parasite-specific p53 homolog.* Proc Natl Acad Sci U S A, 2022. **119**(37): p. e2205201119.

10. Cheng, L.C., et al., *Cellular, ultrastructural and molecular analyses of epidermal cell development in the planarian Schmidtea mediterranea.* Dev Biol, 2018. **433**(2): p. 357-373.

11. Pearson, B.J. and A. Sánchez Alvarado, *A planarian p53 homolog regulates proliferation and self-renewal in adult stem cell lineages.* Development, 2010. **137**(2): p. 213-21.

12. Engeland, K., *Cell cycle regulation: p53-p21-RB signaling.* Cell Death Differ, 2022. **29**(5): p. 946-960.

13. Harper, J.W., et al., *The p21 Cdk-interacting protein Cip1 is a potent inhibitor of G1 cyclin-dependent kinases.* Cell, 1993. **75**(4): p. 805-16.

14. Berghmans, S., et al., *tp53 mutant zebrafish develop malignant peripheral nerve sheath tumors.* Proc Natl Acad Sci U S A, 2005. **102**(2): p. 407-12.

15. el-Deiry, W.S., et al., *WAF1/CIP1 is induced in p53-mediated G1 arrest and apoptosis.* Cancer Res, 1994. **54**(5): p. 1169-74.

16. de Nooij, J.C., M.A. Letendre, and I.K. Hariharan, *A cyclin-dependent kinase inhibitor, Dacapo, is necessary for timely exit from the cell cycle during Drosophila embryogenesis.* Cell, 1996. **87**(7): p. 1237-47.

17.   Boxem, M. and S. van den Heuvel, *lin-35 Rb and cki-1 Cip/Kip cooperate in developmental*
regulation of G1 progression in C. elegans. Development, 2001. **128**(21): p. 4349-59.
18.   Langheinrich, U., et al., *Zebrafish as a model organism for the identification and*
characterization of drugs and genes affecting p53 signaling. Curr Biol, 2002. **12**(23): p. 2023-
8.
19.   Paysan-Lafosse, T., et al., *InterPro in 2022*. Nucleic Acids Research, 2022. **51**(D1): p. D418-
D427.
20.   Varadi, M., et al., *AlphaFold Protein Structure Database in 2024: providing structure coverage*
for over 214 million protein sequences. Nucleic Acids Research, 2023. **52**(D1): p. D368-D375.
21.   Guiley, K.Z., et al., *p27 allosterically activates cyclin-dependent kinase 4 and antagonizes*
palbociclib inhibition. Science, 2019. **366**(6471).
22.   Csergeová, L., D. Krbušek, and R. Janoštiak, *CIP/KIP and INK4 families as hostages of*
oncogenic signaling. Cell Div, 2024. **19**(1): p. 11.
23.   Buck, S.H., D. Chiu, and R.M. Saito, *The cyclin-dependent kinase inhibitors, cki-1 and cki-2,*
act in overlapping but distinct pathways to control cell cycle quiescence during C. elegans
development. Cell Cycle, 2009. **8**(16): p. 2613-20.
24.   Fukuyama, M., et al., *Essential embryonic roles of the CKI-1 cyclin-dependent kinase*
inhibitor in cell-cycle exit and morphogenesis in C elegans. Dev Biol, 2003. **260**(1): p. 273-86.
25.   Lane, M.E., et al., *Dacapo, a cyclin-dependent kinase inhibitor, stops cell proliferation during*
Drosophila development. Cell, 1996. **87**(7): p. 1225-35.
26.   Deng, C., et al., *Mice lacking p21CIP1/WAF1 undergo normal development, but are defective*
in G1 checkpoint control. Cell, 1995. **82**(4): p. 675-84.
27.   Abukhdeir, A.M. and B.H. Park, *P21 and p27: roles in carcinogenesis and drug resistance*.
Expert Rev Mol Med, 2008. **10**: p. e19.
28.   Collins, J.J., III, et al., *Adult somatic stem cells in the human parasite Schistosoma mansoni*.
Nature, 2013. **494**(7438): p. 476-9.
29.   Baker, S.J., et al., *CDK4: a master regulator of the cell cycle and its role in cancer*. Genes
Cancer, 2022. **13**: p. 21-45.
30.   Wendt, G., et al., *A single-cell RNA-seq atlas of Schistosoma mansoni identifies a key*
regulator of blood feeding. Science, 2020. **369**(6511): p. 1644-1649.
31.   Gharbi, S.I., et al., *Crystal structure of active CDK4-cyclin D and mechanistic basis for*
abemaciclib efficacy. NPJ Breast Cancer, 2022. **8**(1): p. 126.
32.   Gharbi, S.I., et al., *Author Correction: Crystal structure of active CDK4-cyclin D and*
mechanistic basis for abemaciclib efficacy. NPJ Breast Cancer, 2022. **8**(1): p. 133.
33.   Fry, D.W., et al., *Specific inhibition of cyclin-dependent kinase 4/6 by PD 0332991 and*
associated antitumor activity in human tumor xenografts. Mol Cancer Ther, 2004. **3**(11): p.
1427-38.
34.   Soucy, S.M., J. Huang, and J.P. Gogarten, *Horizontal gene transfer: building the web of life*. Nat
Rev Genet, 2015. **16**(8): p. 472-82.
35.   Pandey, T., et al., *Acquired stress resilience through bacteria-to-nematode interdomain*
horizontal gene transfer. Embo j, 2023. **42**(24): p. e114835.
36.   Yoshida, S., et al., *Horizontal gene transfer by the parasitic plant Striga hermonthica*. Science,
2010. **328**(5982): p. 1128.
37.   Xi, Z., et al., *Horizontal transfer of expressed genes in a parasitic flowering plant*. BMC
Genomics, 2012. **13**: p. 227.
38.   Zhang, Y., et al., *Evolution of a horizontally acquired legume gene, albumin 1, in the parasitic*
plant Phelipanche aegyptiaca and related species. BMC Evol Biol, 2013. **13**: p. 48.

39.   Zhang, D., et al., *Root parasitic plant Orobanche aegyptiaca and shoot parasitic plant*
Cuscuta australis obtained Brassicaceae-specific strictosidine synthase-like genes by
horizontal gene transfer. BMC Plant Biol, 2014. **14**: p. 19.

40.   Tsai, I.J., et al., *The genomes of four tapeworm species reveal adaptations to parasitism.*
Nature, 2013. **496**(7443): p. 57-63.

41.   Raddatz, G., et al., *Dnmt2-dependent methylomes lack defined DNA methylation patterns.*
Proc Natl Acad Sci U S A, 2013. **110**(21): p. 8627-31.

42.   Cox, G.S., C.K. Phares, and R.A. Schmidt, *Molecular characterization of the Spirometra*
mansonoides genome: renaturation kinetics, methylation, and hybridization to human cDNA
probes. Biochim Biophys Acta, 1990. **1049**(2): p. 134-44.

43.   Fantappié, M.R., E.R. Gimba, and F.D. Rumjanek, *Lack of DNA methylation in Schistosoma*
mansoni. Exp Parasitol, 2001. **98**(3): p. 162-6.

44.   Li, D., D.H. Taylor, and J.C. van Wolfswinkel, *PIWI-mediated control of tissue-specific*
transposons is essential for somatic cell differentiation. Cell Rep, 2021. **37**(1): p. 109776.

45.   Goodier, J.L., *Restricting retrotransposons: a review.* Mob DNA, 2016. **7**: p. 16.

46.   Zhang, H.H., et al., *Recurrent horizontal transfers of Chapaev transposons in diverse*
invertebrate and vertebrate animals. Genome Biol Evol, 2014. **6**(6): p. 1375-86.

47.   Howe, K.L., et al., *WormBase ParaSite - a comprehensive resource for helminth genomics.*
Mol Biochem Parasitol, 2017. **215**: p. 2-10.

48.   Howe, K.L., et al., *WormBase 2016: expanding to enable helminth genomic research.* Nucleic
Acids Res, 2016. **44**(D1): p. D774-80.

49.   Rozanski, A., et al., *PlanMine 3.0-improvements to a mineable resource of flatworm biology*
and biodiversity. Nucleic Acids Res, 2019. **47**(D1): p. D812-d820.

50.   Letunic, I. and P. Bork, *20 years of the SMART protein domain annotation resource.* Nucleic
Acids Research, 2017. **46**(D1): p. D493-D496.

51.   Letunic, I., S. Khedkar, and P. Bork, *SMART: recent updates, new developments and status in*
2020. Nucleic Acids Res, 2021. **49**(D1): p. D458-d460.

52.   Madeira, F., et al., *Search and sequence analysis tools services from EMBL-EBI in 2022.*
Nucleic acids research, 2022. **50**(W1): p. W276-W279.

53.   Yachdav, G., et al., *MSAViewer: interactive JavaScript visualization of multiple sequence*
alignments. Bioinformatics, 2016. **32**(22): p. 3501-3503.

54.   Lemoine, F., et al., *NGPhylogeny.fr: new generation phylogenetic services for non-specialists.*
Nucleic Acids Res, 2019. **47**(W1): p. W260-w265.

55.   Stamatakis, A., *RAxML version 8: a tool for phylogenetic analysis and post-analysis of large*
phylogenies. Bioinformatics, 2014. **30**(9): p. 1312-3.

56.   Kozlov, A.M., et al., *RAxML-NG: a fast, scalable and user-friendly tool for maximum likelihood*
phylogenetic inference. Bioinformatics, 2019. **35**(21): p. 4453-4455.

57.   Edler, D., et al., *raxmlGUI 2.0: A graphical interface and toolkit for phylogenetic analyses using*
RAxML. Methods in Ecology and Evolution, 2021. **12**(2): p. 373-377.

58.   Darriba, D., et al., *ModelTest-NG: A New and Scalable Tool for the Selection of DNA and*
Protein Evolutionary Models. Mol Biol Evol, 2020. **37**(1): p. 291-294.

59.   Letunic, I. and P. Bork, *Interactive Tree of Life (iTOL) v6: recent updates to the phylogenetic*
tree display and annotation tool. Nucleic Acids Research, 2024.

60.   Wang, J., R. Chen, and J.J. Collins, III, *Systematically improved in vitro culture conditions*
reveal new insights into the reproductive biology of the human parasite Schistosoma
mansoni. PLoS Biol, 2019. **17**(5): p. e3000254.

61.   Collins, J.J., III and P.A. Newmark, *It's no fluke: the planarian as a model for understanding*
schistosomes. PLoS Pathog, 2013. **9**(7): p. e1003396.

Figure 1: Expression

Figure S1: Protein Domains/Structure

Figure 2: RNAi

Figure 3: Tumor

Figure 4: CKI phylogeny

Figure S3: CDK4 phylogeny without free-living flatworms

Figure S4: CDK4 phylogeny with free-living flatworms

Figure S5: CCND phylogeny without free-living flatworms

Figure S6: CCND phylogeny with free-living flatworms

Figure S7: CKI phylogney