Title

Substantial Braking Force of Anterior Tibial Translation Induces the Spontaneous
Healing of Murine Anterior Cruciate Ligament

Author
Riku Saito[1], Kazuki Nakayama[1], Yuna Usami[1], Saaya Enomoto[2], Koyo Nogi[1], Takanori
Kokubun[1, 3]
8

9 1. Graduate School of Health, Medicine, and Welfare, Saitama Prefectural University,
10   Saitama, Japan
11 2. Saitama Rehabilitation Center
12 3. Department of Physical Therapy, School of Health and Social Services, Saitama
13   Prefectural University, Saitama, Japan
14

15 **Author Contributions**
16   All authors approved the final version to be published. Study design: RS and TK;
17   data collection and statistical analysis: RS, KN, YU, SE, KN; manuscript
18   composition RS, and TK.
19

Corresponding author
Takanori Kokubun, P.T., Ph.D.
Affiliation: Graduate School of Saitama Prefectural University
Postal address: 820 San-nomiya, Koshigaya-Shi, Saitama, 343-8540, Japan
Phone number: +81-48-973-4176
Fax number: +81-48-973-4176
Email address: kokubun-takanori@spu.ac.jp
27
28

29 **Running Title**
30 Braking force affects ACL healing.
31

Abstract

The anterior cruciate ligament (ACL) has been recognized to have no

spontaneous healing capacity after complete rupture due to an intra-articular

environment. Meanwhile, our group reported that a completely ruptured ACL could

heal spontaneously by restabilizing the femur-tibia joint. Several clinical studies have

recently reported that ruptured ACL could heal spontaneously by controlling abnormal

joint movement. Appropriate mechanical stimulation on the ruptured ACL promotes

spontaneous healing, whereas excessive mechanical stress may inhibit the healing

response. Here, we investigated how much braking force to control anterior tibial

translation (ATT) affects the healing response of the ACL using two different braking

forces for controlled ATT (CATT) models. Male and female C57BL/6 mice were

ruptured ACL and randomly classified into Tight-CATT, Loose-CATT, and ACL-

Rupture (ACL-R) groups. The Tight-CATT group showed a higher success rate of ACL

healing than the Loose-CATT group at 4 and 8 weeks. However, collagen maturation

and mechanical properties on healed ACL did not differ between the Tight-CATT and

Loose-CATT groups. Our results suggested that tight ATT braking immediately after

injury is essential for successfully healing the completely ruptured ACL. On the other

hand, tightly ATT in the early healing phase was insufficient to improve mechanical

strength and collagen maturation. Therefore, we should control mechanical stress tightly

in the early phase of ACL spontaneous healing, and it would be necessary to apply

higher mechanical stress in the later phases to achieve greater healing of the ACL.

Study Design: A controlled laboratory study.

Key terms:

Anterior cruciate ligament; spontaneous healing; conservative treatment; braking;

mechanical stress

Introduction

The knee joint has unstable bone shape and structures; hence, the anterior cruciate ligament (ACL) provides joint stability when the knee joint moves. The ACL is the only tissue that controls the anterior translation of the tibia; therefore, once the ACL is injured, the knee joint loses the proper bone position and kinematics during joint motion in daily activities and sports. ACL injuries occur in a variety of sports, but ACL injury rates are reported to be 2-6 times higher in female athletes than in males due to sex differences in kinematics and hormonal factors.[1–4]

ACL is an intra-articular tissue; the injured ACL has low spontaneous healing capacity. The inside of the joint capsule is filled with synovial fluid and poor vascular supply,[5] leading to poor fibroblast activity in the healing site of the injured ACL.[6] Therefore, ACL reconstruction is recommended and the most commonly selected treatment for ACL-injured patients. However, patients with a low activity level sometimes choose conservative treatment without ACL healing. Regarding these treatments, a 2-year follow-up comparing ACL reconstruction and conservative treatment reported no clear differences in knee joint function, covering pain, and difficulty in sports and recreational activities evaluated with the Knee Injury and

Osteoarthritis Outcome Score (KOOS).[7] Thus, optimal treatment after ACL injury is

debatable.

Several clinical studies have reported that ruptured ACL could heal

spontaneously, as confirmed by MRI.[8–11] However, there is insufficient knowledge of

appropriate post-injury management of knee joints to lead ACL to healing. Recently,

Filbay et al. reported spontaneous ACL healing with the Cross Bracing Protocol (CBP),

which constantly fixed the knee at 90 degrees in patients with ACL injuries.[12] They

showed that 90% of patients had evidence of healing on MRI after a three-month post-

injury. Our previous report using the rats model elucidated that the ACL could heal

spontaneously due to only controlling abnormal anterior tibial translation after ACL

injury.[13] Considering the role of mechanical stress, we should not immobilize the knee

joint after ACL injury because appropriate mechanical loading promotes collagen

remodeling during the ligament healing process.[14–16] On the other hand, excessive

mechanical stress inhibits the healing of tendons and ligaments.[17] Thus, understanding

the acceptable range of mechanical stress, especially during the early phase of the ACL

spontaneous healing response, is important for establishing non-surgical treatment and

appropriate management of ACL-injured patients.

In this study, we created two controlled anterior tibial translation (CATT)

models with altered anterior tibial instability using two different diameters of nylon

sutures. We investigated the effect of braking force to control the anterior tibial

translation (ATT) in mice on the healing response of the injured ACL. Our goal in the

current study is to elucidate the acceptable mechanical stresses during the early phase of

spontaneous healing because excessive mechanical stress during the ACL healing

process may inhibit healing, while stress restriction may make the healing ligaments

fragile.

Methods

Experimental Design

The University Animal Experiment Ethics Committee approved all

experiments. We used male and female 8-9 weeks old C57BL/6 mice (n=125). A total

of 125 mice were ruptured the left ACL and then randomly classified into three groups:

Tight-CATT group, controlled ATT with 3-0 nylon suture (n=53), Loose-CATT group,

controlled ATT with 4-0 nylon suture (n=52), and ACL-R (ACL-Rupture) group; not

controlled ATT (n=20). We created the CATT and ACL-R model for the left hindlimb

in all mice, with the right hindlimb used as the intact group. All mice were housed in

plastic cages under a 12-h light/ dark cycle. The mice were allowed unrestricted

movement within the cage and had free access to food and water. Mice were sacrificed

immediately after injury (n=20) for anterior drawer testing and histology, at 4 (n=51)

and 8 weeks (n=54) for anterior drawer testing, histology, immunofluorescence, and

mechanical testing (Figure 1A).

Nylon Suture Experiments

Before creating the CATT models, we measured the mechanical properties of

nylon sutures using a mechanical testing machine, Univert (CellScale Biomaterial

Testing, Canada). The purpose of this experiment was to understand the difference in

mechanical properties between 3-0 and 4-0 nylon sutures (Akiyama Medical MFG, Co.,

LTD., Tokyo, Japan). Each nylon suture was measured in a double layer since a nylon

suture was looped outside the joint to control ATT in the CATT model. In the failure

testing, we stretched the nylon suture at a preload of 0.1 N and a rate of 20 mm/min

until the nylon sutures ruptured. During the test, the load (N) and displacement (mm)

were recorded, and we measured the load to failure (N) and stiffness (N/mm). Then, we

performed the cyclic test to evaluate viscoelasticity. Nylon sutures were repeated with 5

seconds of stretching up to 90%, 80%, 70%, and 60% of the failure load of each nylon

suture, followed by 5 seconds of relaxation (1 cycle: 10 sec). We recorded the number

of cycles at the point of rupture.

Procedure of Creating the Model

All mice were anesthetized with a combination of anesthetics (medetomidine,

0.375 mg/kg; midazolam, 2.0 mg/kg; and butorphanol, 2.5 mg/kg) for sedation and pain

control, and then we ruptured the ACL non-invasively according to a previous study.[18]

We fixed the hindlimb with the knee joint at 90 degrees and ruptured the ACL by

pushing the femur backward against the tibia (Figure 1B-C). Then, we created bone

holes at the femoral condyle and proximal tibia using a 26G needle (Terumo Co., Ltd.,

Tokyo, Japan). In the Tight- and Loose-CATT groups, we tightly looped a single 3-0 or

4-0 nylon suture through the two holes to control the ATT (Figure 1F). In the ACL-R

group, we loosely looped a nylon suture through the two holes not to control the ATT.

Anterior Drawer Testing

We performed anterior drawer testing to evaluate the braking force to control

the ATT of the knee. We harvested knee joint samples immediately after the mice were

sacrificed. We removed the muscle from the knee while retaining the suture used to

control ATT. Then, we fixed the knee joint at 90 degrees on the self-made anterior

drawer testing device (Figure 3A). The tibia was pulled forward with a 0.05 kgf

constant force spring (Sanko Spring Co., Ltd., Fukuoka, Japan), and X-ray images of

the sagittal plane of the knee joint were taken (Figure 3B-E). We measured ATT from

the X-ray images using ImageJ (ImageJ; National Institutes of Health, Bethesda, MD,

USA) to evaluate the braking force by nylon sutures. The ATT was defined as the

perpendicular line from the most prominent point of the femoral condyle to the long

axis of the tibia. We then removed the nylon suture and again measured the ATT to

evaluate the anterior tibial instability of the ACL healed knees.

Histology

After anterior drawer testing, knee samples were immediately fixed (24-48h) in

4% paraformaldehyde and decalcified with 10% ethylenediaminetetraacetic acid

(Sigma-Aldrich Japan, Tokyo, Japan) for 2 weeks. Finally, they were immersed in 5%

and 30% sucrose for 4 and 12 hours, respectively, and then freeze-embedded using OCT

compound (Sakura Finetek Japan Co., Ltd., Tokyo, Japan). Each sample was sectioned

at 12 μm using a Cryostar NX70 (Thermo Fisher Scientific, Kanagawa, Japan). We

performed Hematoxylin & Eosin (HE) staining to observe ACL continuity and

morphological features, and Picrosirius Red (PsR) staining to evaluate the collagen

maturity of the healed ACL. We observed the ACL with a BZ-X710 microscope

(Keyence, Osaka, Japan). We defined the healed ACL if the ACL reconnection was

confirmed in the HE image. PsR images converted to grayscale images (256 degrees)

showed collagenous tissue as 1-255 on the grayscale and non-collagenous tissue as 0 on

the grayscale (Figure 5A-C). A higher gray scale indicates higher ACL collagen

maturation. Using the BZ-X710 analysis software, the entire ACL areas of the PsR

polarization images were specified as Region of Interest (ROI) and then converted to

grayscale images (Figure 5D). We used the average of the grayscale in the ROI as the

representative value of the healed ACL.

Immunofluorescence

We performed immunofluorescence to evaluate the expression of type 1

collagen in the healed ACL. ACL cryosections were blocked at room temperature for 1

hour using DAKO Antibody Diluent with Background Reducing Components (Agilent

Technologies, Inc., USA). The primary antibody was incubated with Goat polyclonal

antibody (AB758, Sigma-Aldrich) at 4°C overnight. The secondary antibody was

incubated with anti-Goat Rabbit antibody Alexa Flour 488 (ab150141, Abcam) for 4

hours at room temperature. All the stained sections were observed macroscopically with

a microscope BZ-X 710 under controlled imaging conditions.

Biomechanical Testing

We performed failure testing to evaluate the mechanical strength of the healed

ACL. The harvested knee joints were removed from the muscle and frozen at -70°C.

After thawing, we removed all soft tissues except the ACL and fixed the femur and tibia

in 1.5-mm tubes. The tubes were then fixed to the mechanical testing machine Univert

in the extended position (Figure 7A). The ACL was stretched at a preload of 0.1 N and a

rate of 10 mm/min until it was ruptured. We recorded the load to failure of the ACL.

Statistical Analysis

We performed statistical analysis on the results of nylon suture experiments,

anterior drawer testing, grayscale, and mechanical testing using R. The Shapiro-Wilk

test was used to verify the normality of all the analyzed data. Mann-Whitney U-tests

were performed for the data from the nylon suture experiments. The two-way analysis

of variance (ANOVA) was then performed to confirm significant differences between

the group and sex differences in the other results. Because all data were non-normally

distributed, an Aligned Rank Transform (ART)[19,20] was performed before ANOVA.

Based on the results of ANOVA, the Steel-Dwass test was performed as a multiple

comparison between all groups when the interaction was found. When no interaction

was found, multiple comparisons were performed within factors for which a main effect

was found. Statistical significance was set at $p < 0.05$.

Results

Mechanical Properties of Two Nylon Sutures

The results showed that the 3-0 nylon sutures had better mechanical properties.

The load to failure and stiffness were significantly higher for 3-0 nylon sutures ($p <$

0.05) (Figure 2A, B). In cyclic testing, the 3-0 nylon sutures showed higher

viscoelasticity and resistance to elongation. The 4-0 nylon sutures ruptured in 3 cycles

at 90% of the maximum load, 12 cycles at 80%, 107 cycles at 70%, and 791 cycles at

60%. The double 3-0 nylon sutures ruptured in 6 cycles at 90% and in 197 cycles at

80%. 3-0 nylon sutures did not rupture even after 2,000 cycles at loads lower than 70%

(Figure 2C).

Magnitude of Anterior Translation of Tibia

First, in the knee joints with nylon sutures for controlling ATT, interactions

between groups and sex were observed at all timepoints ($p < 0.01$). Immediately after

injury, the ATT was smallest in the intact and Tight-CATT groups. The Loose-CATT

group showed a higher ATT than the Tight-CATT groups, and the ACL-R group

showed the highest ATT. However, there were no significant differences among all

groups and sex (Figure 3F). At 4 and 8 weeks after injury, the ATT was smallest in the

intact group, followed by the Tight-CATT and Loose-CATT groups, which showed

similar ATT. The ACL-R group had the highest ATT. However, there were no

significant differences among all groups and sex (Figure 3G, H).

Secondly, for anterior tibial instability of the ACL-healed knees, interactions

between groups and sex were observed at 4 weeks and 8 weeks ($p < 0.001$). In males at

4 and 8 weeks, intact had the smallest ATT, and the Tight-CATT and Loose-CATT

groups had similar ATT but higher ATT than intact. ACL-R showed the highest ATT.

However, there were no significant differences among all groups and sex (Figure 3I, J).

Histology

First, we showed the success rates of ACL healing with both Tight- and Loose-

CATT groups at 4- and 8-weeks post-injury (Table 1). The Tight-CATT group showed

a higher healing rate for both males and females (42.9~100%), while the Loose-CATT

group showed less healing rate (17~57.1%) and no healed ACL in the ACL-R group

(0%). Next, we showed HE-stained images of the healed ACL (Figure 4). Compared to

the intact ACL with parallel and regular nuclear arrangement, the healed ACL showed

heterogeneous nuclear aggregation not only in the healed region but also in the remnant

ACL region. It was also observed at 4- and 8-weeks post-injury.

In collagen maturation of the healed ACL, there are no interactions observed at

both 4- and 8-weeks post-injury. On the other hand, a significant main effect was

observed in the group: the Tight-CATT group had a significantly lower grayscale than

intact ($p < 0.001$) at 4 weeks, and the Tight- and Loose-CATT groups showed a

significantly lower grayscale than intact ($p < 0.01$) at 8 weeks (Figure 5D, E).

Type I Collagen content of Healing ACL

We performed immunofluorescence of type I collagen, the main component of the

ECM, to evaluate the ECM of the healed ACL. The green area indicates type I collagen,

and the blue area indicates the nucleus (Figure 6). In the intact ACL, we observed type I

collagen orientation and nuclear alignment along the ACL. On the other hand, the

healed ACL of Tight- and Loose-CATT groups showed an irregular orientation of type I

collagen compared to intact. In addition, the Tight- and Loose-CATT groups showed

higher and irregular nuclei aggregation throughout the ACL.

Mechanical Properties of Healed ACL

We performed mechanical testing to evaluate the mechanical strength of the

healed ACL. No interaction was observed at both 4- and 8-weeks post-injury. On the

other hand, a significant main effect was observed in the group: at 4 weeks, intact had a

significantly higher mechanical strength than the Tight-CATT ($p < 0.01$) and Loose-

CATT group ($p < 0.05$), and at 8 weeks, intact had a significantly higher mechanical

strength than the Tight-CATT group ($p < 0.01$). However, the healed ACLs in the

Tight- and Loose-CATT groups showed similar mechanical strength and were

approximately 50% of the normal ACL (Figure 7).

Discussion

The first objective of this study was to elucidate whether the difference in braking force to control the ATT affects the healing response of the completely injured ACL. The results of the current study showed that the Tight-CATT group had a higher success rate of ACL healing. The Tight-CATT group showed a similar braking force of the ATT as the intact group, which had a trend higher braking force than the Loose-CATT group immediately after injury. However, after 4 weeks post-injury, the Tight-CATT group showed decreased braking force, which was not significantly different from the Loose-CATT group. These results indicate that controlling the ATT immediately after ACL injury is one of the most critical key factors for spontaneous healing of the ruptured ACL. A clinical study of ACL non-surgical treatment reported that 90% of ACL-injured patients had healed ACL with knee joints immobilized at 90 degrees by orthotics.[12] On the other hand, another study showed a lower rate of successful healing in younger ACL-injured patients because they have higher activity levels compared with middle-aged and older patients.[21] In addition, it was reported that early exercise intervention during the supraspinatus tendon healing process induced tendon degeneration.[22] Accordingly, the lower ACL healing success rate in the Loose-CATT group is probably caused by the mechanical stress associated with increased ATT. During the healing

process of the MCL, vascular-rich granulation tissue invades the rupture area of the

ligament to close and then gradually replace with fibrous tissue.[23] The increase in ATT

probably led to the failure of healing by pulling apart the ACL remnant and inhibiting

the invasion of granulation tissue or by re-rupturing the very fragile granulation tissue.

Therefore, we showed that the tight braking force of ATT is important in the early

healing process after ACL injury.

In this study, we found sex differences in the success rate of ACL healing. It is

known that there are sex differences in ACL injury mechanisms, and the ACL injury

rate in women is 2~6 times higher than that in men.[1] Women are more likely to have

knee ligament laxity, which can lead to decreased knee joint stability during exercise.[24]

However, the ACL healing success rate of the Tight-CATT group was higher in females

than in males. The ATT braking by the nylon suture may have loosened more easily in

males, which would have affected these results. The similar ATT braking force in male

and female Tight-CATT group showed immediately after injury. However, the ATT

braking of the male Tight-CATT group decreased after 4 weeks post-injury, which was

lower than that of female. It may be caused by the difference in cage activity between

male and female mice. Previous studies have reported that the amount of spontaneous

exercise in wheel running in C57BL/6 mice is higher in females.[25] In addition, a study

investigating the amount of home-cage activity showed that the amount of activity in

the dark zone was significantly higher in females.[26] In contrast, males exhibit more

aggressive behaviors, such as fighting and mounting, when kept with 2 or more

animals.[27] In this study, we housed 2-4 mice in 1 cage for both sexes. Based on this

knowledge, male mice may show more intense activity than females in a cage, which

may induce the CATT to loosen and high mechanical stress on the ACL in male mice.

The second objective of this study was to elucidate whether the difference in

mechanical stress in the early phase affects collagen maturation and mechanical strength

of the healed ACL in mice. As mentioned above, we suggested that tight ATT braking

is necessary to achieve a higher spontaneous healing success rate after ACL injury.

Thus, in terms of healing success rate, increased mechanical stress in the early phase of

the healing process is negative. However, we observed ACL healing in several knees of

the Loose-CATT group, although the success rate was low. In general, mechanical

stress on the ligaments promotes fibroblast activation and ECM synthesis, such as

collagen.[15] It is known that the promotion of ECM synthesis improves the mechanical

strength of the ligaments. Based on this knowledge, the healed ACL in the Loose-CATT

group might have a greater quality of the healed ACL, such as ECM maturation and

mechanical strength, although it had a lower healing success rate. Once the injured ACL

in the Loose-CATT group was fortunately healed, it was exposed to higher mechanical

stress during the early ACL healing process. Therefore, we performed ACL mechanical

testing, picrosirius red staining, and immunofluorescence at 4 and 8 weeks to

understand the quality of the healed ACL in each group. As the results, the healed ACLs

in Tight- and Loose-CATT groups showed similar collagen maturation and mechanical

strength, and which were lower quality of the ACL compared to the intact ACL.   In

addition, the expression of type I collagen, a protein that contributing to the mechanical

properties in the ligaments and tendons showed that it was localized along the ACL in

the intact ACL, while irregular and poor collagen localization was observed in both

Tight- and Loose-CATT groups. To summarize these results, against our expectations,

there was no difference in the quality of the healed ACL between the Tight- and Loose-

CATT groups. The reason for these results is probably that the mechanical stress on the

healed ACL was similar in the Tight- and Loose-CATT groups after 4 weeks post-

injury. Anterior drawer testing showed ATT was increased in both the Tight- and

Loose-CATT groups after 4 weeks post-injury compared with immediately after injury.

Increased ATT after 4weeks post-injury should led to mechanical stress on the healed

ACL, but it was insufficient to promote healing of the ACL. Low-intensity exercise

during the healing process of the supraspinatus tendon is reported to be beneficial in

maintaining structure and promoting repair in the early stages of healing but is

insufficient to promote long-term healing.[28] In addition, there is an optimal timing for

mechanical stress loading to promote tendon and ligament healing. In ACL-

reconstructed mice, it was reported that proper timing of mechanical stress was more

effective in promoting healing than immediate postoperative and delayed mechanical

stress.[25] Thus, there is an optimal amount and timing of mechanical stress loading

during tendon and ligament healing processes. In future studies, we need to investigate

the influence of exercise intervention, i.e., rehabilitation, on promoting healing after

spontaneous ACL healing.

In this study, we have several limitations. First, the present data were evaluated

immediately after injury and at 4 and 8 weeks after injury. It was impossible to evaluate

differences in ATT and the healing process from 0 to 4 weeks. Therefore, we will

analyze at finer time points in the future to identify when CATT is no longer necessary.

We will then be able to determine the period when orthotic management is necessary

after ACL injury. Second, differences in the braking force of the ATT in anterior drawer

testing are ex vivo data and do not completely show the differences in stress on the

ACL in vivo. Although we evaluated the mechanical stress load by the difference of

ATT, we will evaluate the knee joint motion during walking and the amount of activity

in the cage as a future study. Then, we would analyze in detail the mechanical stress

loaded on the ACL by the difference in braking force. Third, in this study, mechanical

stress on the healed ACL did not improve mechanical strength sufficiently. Therefore,

applying a higher load of mechanical stress using exercise would be necessary in future

studies.

In this study, we investigated the effects of different ATT braking forces on the

ACL spontaneous healing success rate and mechanical strength of the healed ACL in

mice CATT models with different braking sutures. The results showed that tight ATT

braking immediately after injury is necessary for the successful healing of the injured

ACL. On the other hand, increased ATT in the early phase was insufficient to improve

mechanical strength and collagen maturation. This study suggested that tight ATT

braking is required on ACL healing at the early phase of post-injury, and then higher

mechanical loading may be necessary to achieve higher ACL properties at the

remodeling phase when the ACL has healed. We provide insight into the appropriate

management after ACL-injured patients in this study.

Acknowledgments

This work was supported by JSPS KAKENHI (21H03306, 19KK0411), the grant from
Tokyo Physical Therapist Association (05094) and the attack grant from Saitama
Prefectural University.

Reference

1. Spindler KP, Wright RW. 2008. Anterior Cruciate Ligament Tear. N Engl J Med
359(20):2135–2142.

2. Ford KR, Myer GD, Hewett TE. 2003. Valgus Knee Motion during Landing in
High School Female and Male Basketball Players. Medicine & Science in Sports
& Exercise 35(10):1745.

3. Jordan SS, DeFrate LE, Nha KW, et al. 2007. The in vivo kinematics of the
anteromedial and posterolateral bundles of the anterior cruciate ligament during
weightbearing knee flexion. Am J Sports Med 35(4):547–554.

4. Shultz SJ, Sander TC, Kirk SE, Perrin DH. 2007. Sex differences in knee joint
laxity change across the female menstrual cycle. J Sports Med Phys Fitness
45(4):594-603

5. Bray RC, Leonard CA, Salo PT. 2003. Correlation of healing capacity with
vascular response in the anterior cruciate and medial collateral ligaments of the
rabbit. J. Orthop. Res. 21(6):1118–1123.

6. Menetrey J, Laumonier T, Garavaglia G, et al. 2011. α-Smooth muscle actin and
TGF-β receptor I expression in the healing rabbit medial collateral and anterior
cruciate ligaments. Injury 42(8):735–741.

7. Frobell RB, Roos EM, Roos HP, et al. 2010. A Randomized Trial of Treatment for
Acute Anterior Cruciate Ligament Tears. N Engl J Med 363(4):331–342.

8. Ihara H, Miwa M, Deya K, Torisu K. 1996. MRI of anterior cruciate ligament
healing. J Comput Assist Tomogr 20(2):317–321.

9. Fujimoto E, Sumen Y, Ochi M, Ikuta Y. 2002. Spontaneous healing of acute
anterior cruciate ligament (ACL) injuries - conservative treatment using an
extension block soft brace without anterior stabilization. Archives of Orthopaedic
and Trauma Surgery 122(4):212–216.

10. Previ L, Monaco E, Carrozzo A, et al. 2023. Spontaneous healing of a ruptured
anterior cruciate ligament: a case series and literature review. J EXP ORTOP
10(1):11.

11. Filbay SR, Roemer FW, Lohmander LS, et al. 2023. Evidence of ACL healing on
MRI following ACL rupture treated with rehabilitation alone may be associated
with better patient-reported outcomes: a secondary analysis from the KANON
trial. Br J Sports Med 57(2):91–99.

12. Filbay SR, Dowsett M, Chaker Jomaa M, et al. 2023. Healing of acute anterior
cruciate ligament rupture on MRI and outcomes following non-surgical
management with the Cross Bracing Protocol. Br J Sports Med :bjsports-2023-
106931.

13. Kokubun T, Kanemura N, Murata K, et al. 2016. Effect of Changing the Joint
Kinematics of Knees With a Ruptured Anterior Cruciate Ligament on the
Molecular Biological Responses and Spontaneous Healing in a Rat Model. Am J
Sports Med 44(11):2900–2910.

14. Gomez MA, Woo SL-Y, Amiel D, et al. 1991. The effects of increased tension on
healing medial collateral ligaments. Am J Sports Med 19(4):347–354.

15. Zhang J, Wang JH-C. 2013. The Effects of Mechanical Loading on Tendons - An
In Vivo and In Vitro Model Study. PLoS ONE 8(8):e71740.

16. Eliasson P, Andersson T, Aspenberg P. 2012. Achilles tendon healing in rats is
improved by intermittent mechanical loading during the inflammatory phase:
TENDON HEALING AND EARLY LOADING. J. Orthop. Res. 30(2):274–279.

17. Wada S, Lebaschi AH, Nakagawa Y, et al. 2019. Postoperative Tendon Loading
With Treadmill Running Delays Tendon-to-Bone Healing: Immunohistochemical
Evaluation in a Murine Rotator Cuff Repair Model. J. Orthop. Res. 37(7):1628–
1637.

18. Takahata K, Arakawa K, Enomoto S, et al. 2023. Joint instability causes catabolic
enzyme production in chondrocytes prior to synovial cells in novel non-invasive
ACL ruptured mouse model. Osteoarthritis and Cartilage 31(5):576–587.

19. Wobbrock JO, Findlater L, Gergle D, Higgins JJ. 2011. The aligned rank
transform for nonparametric factorial analyses using only anova procedures. In:
Proceedings of the SIGCHI Conference on Human Factors in Computing Systems.
New York, NY, USA: Association for Computing Machinery. p 143–146
Available from: https://doi.org/10.1145/1978942.1978963.

20. Jordão S, Cortes N, Gomes J, et al. 2022. Synchronization performance affects gait
variability measures during cued walking. Gait & Posture 96:351–356.

21. Ihara H, Kawano T. 2017. Influence of Age on Healing Capacity of Acute Tears of
the Anterior Cruciate Ligament Based on Magnetic Resonance Imaging
Assessment. Journal of Computer Assisted Tomography 41(2):206.

22. Chen Y, Zhang T, Wan L, et al. 2021. Early treadmill running delays rotator cuff
healing via Neuropeptide Y mediated inactivation of the Wnt/β-catenin signaling.
Journal of Orthopaedic Translation 30:103–111.

23. Chamberlain CS, Crowley E, Vanderby R. 2009. The spatio-temporal dynamics of
ligament healing. Wound Repair and Regeneration 17(2):206–215.

24. Park S-K, Stefanyshyn DJ, Loitz-Ramage B, et al. 2009. Changing Hormone
Levels during the Menstrual Cycle Affect Knee Laxity and Stiffness in Healthy
Female Subjects. Am J Sports Med 37(3):588–598.

25. Chen H, Li S, Xiao H, et al. 2021. Effect of Exercise Intensity on the Healing of
the Bone-Tendon Interface: A Mouse Rotator Cuff Injury Model Study. Am J
Sports Med 49(8):2064–2073.

26. Bains RS, Forrest H, Sillito RR, et al. 2023. Longitudinal home-cage automated
assessment of climbing behavior shows sexual dimorphism and aging-related
decrease in C57BL/6J healthy mice and allows early detection of motor
impairment in the N171-82Q mouse model of Huntington's disease. Front. Behav.
Neurosci. 17:1148172.

27. Lidster K, Owen K, Browne WJ, Prescott MJ. 2019. Cage aggression in group-
housed laboratory male mice: an international data crowdsourcing project. Sci Rep
9(1):15211.

28. Camp CL, Lebaschi A, Cong G-T, et al. 2017. Timing of Postoperative
Mechanical Loading Affects Healing Following Anterior Cruciate Ligament
Reconstruction: Analysis in a Murine Model. The Journal of Bone and Joint
Surgery 99(16):1382–1391.

 Table Legend

 Table 1. We defined ACL spontaneous healing from HE stained images. Tight-CATT

 group showed higher healing success rate than Loose-CATT group at 4 and 8 weeks

 after injury.

	4 weeks				8 weeks			
	3-0 CATT		4-0 CATT		3-0 CATT		4-0 CATT	
	Male	**Female**	**Male**	**Female**	**Male**	**Female**	**Male**	**Female**
Healed	5	4	2	1	3	4	1	4
Non-healed	2	1	3	3	4	0	5	3
Success rates of ACL healing (%)	71.4	80.0	40.0	25.0	42.9	100.0	16.7	57.1

Figure Legend

Figure 1. C57BL/6 mice were ruptured the ACL and classified to Tight-CATT, Loose-

CATT and ACL-R group. Mice were analyzed via anterior drawer testing and histology

immediately after injury, and anterior drawer testing, histology, and mechanical testing

at 4- and 8-weeks post-injury (A). Non-invasive ACL-rupture. We fixed the hindlimb

with the knee joint at 90 degrees and ruptured the ACL by pushing the femur backward

against the tibia. (B, C) HE images of intact ACL (D) and immediately after ACL

rupture (E) (Scale bar = 300 μm). Controlled Anterior Tibial Translation (CATT)

model. Tightly looped by single 3-0 or 4-0 nylon suture through the femoral and tibial

bone holes to control the ATT (F).

Figure 2. The load to failure (A) and stiffness (B) of the nylon suture in failure testing,

and the number of cycles at the point of suture rupture in cyclic testing (C) were

recorded. (*$p < 0.05$, **$p < 0.01$)

Figure 3. Anterior drawer testing was performed with the knee joint fixed at 90° via the

self-made anterior drawer device (A). Representative X-ray images of each group (B-

E). We evaluated the braking force of the ATT for each group immediately after injury

(F) and at 4 (G) and 8 weeks (H), and anterior tibial instability of ACL healed knee at 4

(I) and 8 weeks (J).

Figure 4. HE stained images of the Tight-CATT (A, B, C, and D), Loose-CATT (E, F,

G, and H), ACL-R groups (I, J, K, and L) at 4 and 8 weeks (Scale bar = 300 μm). The

arrows point to the healing region of the healed ACL.

Figure 5. Picrosirius red stained (A) and polarized image (B) of the ACL. Picrosirius

red images were converted to grayscale images (C) and specified ACL area as ROI (D).

Collagen maturation of the ACL was evaluated at 4 (E) and 8 weeks (F) after injury

(Scale bar = 300 μm).

Figure 6. Immunofluorescence images of the intact ACL, Tight-CATT, Loose-CATT

groups at 4 and 8 weeks: DAPI (blue) and Collagen type I (green). The images were

taken with a 20× magnification lens (Scale bar = 300 μm).

511 **Figure 7.** To perform biomechanical testing, the soft tissues of the knee except the ACL

512 were dissected and the femur and tibia were fixed in tubes (A). The load to failure (A)

513 of the healed ACL at 4 (B) and 8 (C) weeks after injury were recorded.