Title

The braking force to Control the Abnormal Tibial Translation Affects the

Spontaneous Healing Response for Complete Ruptured Anterior Cruciate

Ligament in Mice Model

Author

Riku Saito[1], Kazuki Nakayama[1], Yuna Usami[1], Saaya Enomoto[2], Koyo Nogi[1], Takanori

Kokubun[1, 3]

1.   Graduate School of Health, Medicine, and Welfare, Saitama Prefectural University,

Saitama, Japan

2.   Saitama Rehabilitation Center

3.   Department of Physical Therapy, School of Health and Social Services, Saitama

Prefectural University, Saitama, Japan

Author Contributions

All authors approved the final version to be published. Study design: RS and TK;

data collection and statistical analysis: RS, KN, YU, SE, KN; manuscript

composition RS, and TK.

Corresponding author

Takanori Kokubun, P.T., Ph.D.

Affiliation: Graduate School of Saitama Prefectural University

Postal address: 820 Sannomiya, Koshigaya-Shi, Saitama, 343-8540, Japan

Phone number: 048-973-4176

Fax number: 048-973-4176

Email address: kokubun-takanori@spu.ac.jp

Running Title

Braking force affects ACL healing

Abstract

The anterior cruciate ligament (ACL) has been considered to not heal spontaneously

after complete injury due to its unique intra-articular environment. However, several

clinical and animal studies have reported that ruptured ACL could heal spontaneously

by controlling abnormal joint movement. On the other hand, heled ACL has lower

collagen maturity and mechanical properties than intact ACL. Appropriate mechanical

stress on the ligament could promote spontaneous healing, whereas excessive

mechanical stress probably inhibits the healing response. We investigated the effect of

braking force to control anterior tibial translation (ATT) on the healing response of the

ACL using two different controlled anterior tibial translation (CATT) models. Male and

female C57BL/6 mice were ruptured ACL and randomly classified to CATT with 3-0

nylon suture (3-0 CATT) group, CATT with 4-0 nylon suture (4-0 CATT) group, ACL-

rupture (ACL-R) group. We evaluated magnitude of ATT, success rate of the ACL

healing, collagen maturity and mechanical property immediately after injury, at 4- and

8-weeks post-injury. 3-0 CATT group had better braking force of ATT than 4-0 CATT

group immediately after injury and showed higher success rate than 4-0 CATT group

females at 4 and 8 weeks. In contrast, collagen maturation and mechanical strength of

the healed ACL showed no difference between 3-0 CATT and 4-0 CATT groups. Our

results suggested that firm ATT braking immediately after injury is necessary for

successful healing of the injured ACL. On the other hand, increased ATT in the early

phase was insufficient to improve mechanical strength and collagen maturation, and

future studies is needed to elucidate the optimal timing of mechanical stress.

Study Design: A controlled laboratory study.

Key terms:

Anterior cruciate ligament; spontaneous healing; conservative treatment; braking;

mechanical stress

Introduction

The anterior cruciate ligament (ACL) is essential to the knee joint stability and

movement. The knee joint has unstable bone shape and structures, and the ACL is the

only soft tissue that controls the anterior translation of the tibia.[1] Therefore, once ACL

injured, the knee loses the proper bone position and kinematics during joint motion in

daily activities and playing sports. ACL injuries used to be young and athlete injuries,

but recently, the number of middle-aged or older ACL-injured patients has increased.[2]

These patients need any type of treatment strategy, not only surgical reconstruction.

ACL in the intra-articular tissue, thus the injured ACL has low self-healing

capacity. The environment of the joint capsule is filled with synovial fluid and poor

vascular supply,[3] these leads poor fibroblast activity in the healing site of injured ACL.[4]

Therefore, ACL reconstruction is recommended and most commonly selected treatment

for ACL-injured patients. However, patients with a low activity level sometimes choose

conservative treatment without ACL healing. Regarding these treatments, a 2-year

follow-up comparing ACL reconstruction and conservative treatment reported no clear

differences in knee joint function, covering pain, and difficulty in sports and

recreational activities evaluated with the Kenn Injury and Osteoarthritis Outcome Score

(KOOS).[5] Thus, optimal treatment after ACL injury is debatable.

Several clinical studies have reported that ruptured ACL could heal

spontaneously, as confirmed by MRI.[6–9] However, there is insufficient knowledge of

appropriate post-injury management of the ACL and knee joint to lead them to healing.

Filbay et al. reported spontaneous ACL healing with the Cross Bracing Protocol (CBP),

which constantly fixed the knee at 90 degrees in patients with ACL injuries.[10] They

showed 90% of patients had evidence of healing on three-month MRI. In our previous

report using the rats model, we elucidated that the ACL could heal spontaneously due to

only controlling abnormal anterior tibial translation after ACL injury.[11] Regarding

mechanical stress, we should not immobilize the knee joint after ACL injury because

appropriate mechanical loading promotes collagen remodeling during the ligament

healing process.[12–14] On the other hand, excessive mechanical stress inhibits the healing

of tendons and ligaments.[15] Thus, understanding the acceptable range of mechanical

stress, especially during the early phase of the ACL spontaneous healing response is

important for establishing non-surgical treatment and appropriate management of the

injured ACL.

In this study, we aimed to investigate the effect of braking force to control the

anterior tibial translation (ATT) in mice on the healing response of the injured ACL. We

created two controlled anterior tibial translation (CATT) models with altered anterior

tibial instability by controlling the anterior tibial translation using two different

diameters of nylon sutures. Thus, we aimed to elucidate the acceptable mechanical

stresses during early phase of the ACL spontaneous healing and to provide insight into

the appropriate management after ACL injured patients.

Methods

Experimental Design

The University Animal Experiment Ethics Committee approved all

experiments. We used male and female 8-9 weeks old C57BL/6 mice (n=125). A total

of 125 mice were ruptured the left ACL and then randomly classified into three groups:

3-0 CATT group; controlled ATT with 3-0 nylon suture (n=53), 4-0 CATT group;

controlled ATT with 4-0 nylon suture (n=52), and ACL-R (ACL-Rupture) group; not

controlled ATT (n=20). We created the CATT and ACL-R model for the left hindlimb

in all mice, with the right hindlimb used as the intact group. Mice were sacrificed

immediately after injury (n=20) for anterior drawer testing and histology, at 4 (n=51)

and 8 weeks (n=54) for anterior drawer testing, histology, immunofluorescence, and

mechanical testing (Figure 1).

Nylon Suture Experiments

Prior to creating the CATT models, we measured the mechanical properties of

nylon sutures using a mechanical testing machine, Univert (CellScale Biomaterial

Testing, Canada). The purpose of mechanical testing was to understand the difference of

mechanical properties between 3-0 and 4-0 nylon sutures (Akiyama Medical MFG, Co.,

LTD., Tokyo, Japan). Each nylon suture was measured in a double layer, since a nylon

suture was looped outside the joint to control ATT in the CATT model. In the failure

testing, we stretched the nylon suture at preload of 0.1 N and a rate of 20 mm/min until

the nylon sutures ruptured. During the test, the load (N) and displacement (mm) were

recorded, and we measured the load to failure (N) and stiffness (N/mm). Then we

performed the cyclic test to evaluate viscoelasticity. Nylon sutures were repeated 5

seconds of stretching up to 90%, 80%, 70%, and 60% of the failure load of each nylon

sutures, followed by 5 seconds of relaxation (1 cycle: 10 sec). We recorded the number

of cycles at the point of rupture.

Procedure of Creating the Model

All mice were anesthetized with a combination of anesthetics (medetomidine,

0.375 mg/kg; midazolam, 2.0 mg/kg; and butorphanol, 2.5 mg/kg) for sedation and pain

control, and then we ruptured the ACL non-invasively (Figure 2A-B) according to a

previous study.[16] We fixed the hindlimb with the knee joint at 90 degrees and ruptured

the ACL by pushing the femur backward against the tibia (Figure 2A-B). Then, we

created bone holes at the femoral condyle and proximal tibia using a 26G needle

(Terumo Co., Ltd., Tokyo, Japan). In the 3-0 and 4-0 CATT groups, we tightly looped a

single 3-0 or 4-0 nylon suture through the two holes to control the ATT (Figure 2E). In

the ACL-R group, we loosely looped a nylon suture through the two holes not to control

the ATT.

Anterior Drawer Testing

We performed anterior drawer testing to evaluate the braking force to control

ATT of the knee. We harvested knee joint samples immediately after the mice were

sacrificed. We removed the muscle from the knee while retaining the suture used for

controlling ATT. Then, we fixed the knee joint in 90-degree on the self-made anterior

drawer testing device (Figure 4A). The tibia was pulled forward with a 0.05 kgf

constant force spring (Sanko Spring Co., Ltd., Fukuoka, Japan), and X-ray images of

the sagittal plane of the knee joint were taken. We measured ATT from the X-ray

images using ImageJ (ImageJ; National Institutes of Health, Bethesda, MD, USA) to

evaluate the braking force by nylon sutures. The ATT was defined as the perpendicular

line from the most prominent point of the femoral condyle to the long axis of the tibia

(Figure 4B). We then removed the nylon suture and again measured the ATT to

evaluate the anterior tibial instability of the ACL healed knees.

Histology

After anterior drawer testing, knee samples were immediately fixed (24-48h) in

4% paraformaldehyde and decalcified with 10% ethylenediaminetetraacetic acid

(Sigma-Aldrich Japan, Tokyo, Japan) for 2 weeks. Finally, they were immersed in 5%

and 30% sucrose for 4 and 12 hours, respectively, and then freeze-embedded using OCT

compound (Sakura Finetek Japan Co., Ltd., Tokyo, Japan). Each sample was sectioned

at 12 μm using a Cryostar NX70 (Thermo Fisher Scientific, Kanagawa, Japan). We

performed Hematoxylin & Eosin (HE) staining to observe ACL continuity and

morphological features and Picrosirius Red (PsR) staining to evaluate collagen maturity

of the healed ACL. We observed the ACL with a BZ-X710 microscope (Keyence,

Osaka, Japan). We defined the healed ACL if the ACL reconnection was confirmed in

the HE image. PsR images converted to grayscale images (256 degrees) showed

collagenous tissue as 1-255 on the grayscale and non-collagenous tissue as on the

grayscale. Higher gray scale indicates higher ACL collagen maturation. Using the BZ-

X710 analysis software, the entire ACL areas of the PsR polarization images were

specified as Region of Interest (ROI) and then converted to grayscale images. We used

the average of the grayscale in the ROI as the representative value of the healed ACL.

Immunofluorescence

We performed immunofluorescence to evaluate the expression of type 1

collagen in the healed ACL. ACL cryosections were blocked at room temperature for 1

hour using DAKO Antibody Diluent with Background Reducing Components (Aglient

Technologies, Inc., USA). The primary antibody was incubated with Goat polyclonal

antibody (AB758, Sigma-Aldrich) at 4°C overnight. Secondary antibody was incubated

with anti-Goat Rabbit antibody Alexa Flour 488 (ab150141, abcam) for 4 hours at room

temperature. All the stained sections were observed macroscopically with a microscope

BZ-X 710 under controlled imaging conditions.

Biomechanical Testing

We performed failure testing to evaluate the mechanical strength of the healed

ACL. The harvested knee joints were removed from the muscle and frozen at -70°C.

After thawing, we removed all soft tissues except the ACL and fixed the femur and tibia

in 1.5-mm tubes. The tubes were then fixed to the mechanical testing machine Univert

in the extended position. The ACL was stretched at a preload of 0.1 N and a rate of 10

mm/min until it was ruptured. We recorded the load to failure of the ACL.

Statistical Analysis

We performed statistical analysis on the results of nylon suture experiments,

anterior drawer testing, gray scale, and mechanical testing using R. The Shapiro-Wilk

test was used to verify the normality of all the analyzed data. Mann-Whitney U-tests

were performed for the data from the nylon suture experiments. The two-way analysis

of variance (ANOVA) was then performed to confirm for significant differences

between the group and sex difference the other results. Because all data were non-

normally distributed, an Aligned Rank Transform (ART)[17,18] was performed before

ANOVA. Based on the results of ANOVA, the Steel-Dwass test was performed as a

multiple comparison between all groups when the interaction was found. When no

interaction was found, multiple comparisons were performed within factors for which a

main effect was found. Statistical significance was set at $p < 0.05$.

Results

Mechanical Properties of Two Nylon Sutures

208   The results showed that the 3-0 nylon sutures had better mechanical properties.

The load to failure and stiffness were significantly higher for 3-0 nylon sutures ($p <$

0.05) (Figure3A, B). In cyclic testing, the 3-0 nylon sutures showed higher

viscoelasticity and resistance to elongation. The 4-0 nylon sutures ruptured in 3 cycles

at 90% of the maximum load, 12 cycles at 80%, 107 cycles at 70%, and 791 cycles at

60%. The double 3-0 nylon sutures ruptured in 6 cycles at 90% and in 197 cycles at

80%. 3-0 nylon sutures did not rupture even after 2,000 cycles at loads lower than 70%

(Figure 3C).

Magnitude of Anterior Translation of Tibia

218   First, in the knee joints with nylon sutures for controlling ATT, interactions

between groups and sex were observed at all timepoint ($p < 0.01$). Immediately after

injury, the ATT was smallest in the intact and 3-0 CATT groups. The 4-0 CATT group

showed the higher ATT compared with intact and 3-0 CATT group, and the ACL-R

group showed the highest ATT. However, there were no significant differences among

all groups and sex (Figure 4D). At 4 and 8 weeks after injury, the ATT was smallest in

the intact group, followed by the 3-0 CATT and 4-0 CATT groups, which showed

similar ATT. The ACL-R group had the highest ATT. However, there were no

significant differences among all groups and sex (Figure 4E, F).

Secondly, for anterior tibial instability of the ACL healed knees, interactions

between groups and sex were observed at 4 weeks and 8 weeks ($p < 0.001$). In males at

4 and 8 weeks, intact had the smallest ATT, and the 3-0 CATT and 4-0 CATT groups

had similar ATT but higher ATT than intact. ACL-R showed the highest ATT.

However, there were no significant differences among all groups and sex (Figure 4G,

H).

Histology

First, we showed the success rates of the ACL healing with both 3-0 and 4-0

CATT groups at 4- and 8-weeks post-injury (Table 1). The 3-0 CATT group showed a

higher healing rate for both males and females (42.9~100%), while 4-0 CATT group

showed less heling rate (17~57.1%), and no healed ACL in ACL-R group (0%). Next,

we showed HE-stained images of the healed ACL (Figure 5). We observed ACL

spontaneously healing at 4 weeks after injury in both males and females. Compared to

the intact ACL with parallel and regular nuclear arrangement, the healed ACL showed

heterogeneous nuclear aggregation not only at the healed region but also the remnant

ACL region. It was also observed at 4- and 8-weeks post-injury.

In collagen maturation of the healed, there are no interaction was observed at

both 4- and 8-weeks post-injury. On the other hand, a significant main effect was

observed in the group: at 4 weeks, the 3-0 CATT group had a significantly lower gray

scale than intact ($p < 0.001$), and at 8 weeks, still the 3-0 and 4-0 CATT groups showed

a significantly lower gray scale than intact ($p < 0.01$) (Figure 6D, E).

Type I Collagen content of Healing ACL

We performed immunofluorescence of type I collagen, the main component of the

ECM, to evaluate the ECM of the healed ACL. The green area indicates type I collagen

and the blue area indicates the nucleus (Figure 7). In the intact ACL, we observed type I

collagen orientation and nuclear alignment along the ACL. On the other hand, the

healed ACL of 3-0 CATT and 4-0 CATT groups showed an irregular orientation of type

I collagen compared to intact. In addition, the 3-0 and 4-0 CATT groups showed higher

and irregular nuclei aggregation throughout the ACL.

Mechanical Properties of Healed ACL

We performed mechanical testing to evaluate the mechanical strength of the

healed ACL. No interaction was observed at both 4- and 8-weeks post-injury. On the

other hand, a significant main effect was observed in the group: at 4 weeks, intact had a

significantly higher mechanical strength than the 3-0 CATT ($p < 0.01$) and 4-0 CATT

group ($p < 0.05$), and at 8 weeks, intact had a significantly higher mechanical strength

than the 3-0 CATT group ($p < 0.01$). However, the healed ACLs in the 3-0 CATT and

4-0 CATT groups showed similar mechanical strength and were approximately 50% of

the normal ACL (Figure 8).

Discussion

The objectives of this study were to elucidate whether the difference of braking

force to control the ATT affect the healing response of the completely injured ACL.

And also, whether the difference of mechanical stress in the early phase affects to

collagen maturation and mechanical strength of the healed ACL in mice. The results of

the current study showed that the 3-0 CATT group with firmly braked ATT had a higher

success rate of ACL healing. This means that the control of the ATT after ACL injured

knee is one of the most critical key factors for the spontaneous healing of the ruptured

ACL. On the other hand, the 4-0 CATT group with increased mechanical stress in the

early phase did not show improvement in the quality of the healed ACL such as

collagen maturation and mechanical strength.

In this study, we created the CATT model using 3-0 and 4-0 nylon sutures. We

then confirmed the braking force with these sutures of the ATT ex vivo using anterior

drawer testing. As a result, the 3-0 CATT group showed the similar braking force of the

ATT as the intact group, which had a trend higher braking force than the 4-0 CATT

group immediately after injury. However, at 4- and 8-weeks post-injury, the 3-0 CATT

group showed a trend toward decreased braking force, which was no difference to the 4-

0 CATT group. Prior to mice experiments, we evaluated the mechanical properties of

each nylon suture. The 3-0 nylon sutures with larger diameters showed higher

mechanical strength in failure testing and better viscoelastic properties in cyclic testing.

These results indicate that the difference in nylon suture elasticity is probably one of the

factors that caused the difference in braking force between the 3-0 and 4-0 CATT

groups. In addition, differences in the diameter of the nylon suture relative to the bone

hole probably also affect the difference in braking force. We used the same 26G needle

to create the bone holes in the femur and tibia in the 3-0 CATT and 4-0 CATT models.

The 3-0 CATT group with a larger diameter suture had a smaller gap after passing

through the bone holes, whereas the 4-0 CATT group had a larger gap. This may have

been another reason for the difference in braking force between the 3-0 and 4-0 CATT.

Since the result of HE-staining showed that the 3-0 CATT group had a higher success

rate of ACL healing than 4-0 CATT group at 4 weeks and 8 weeks, we found that firm

ATT braking at early phase post-injury is one of the most important factors for

spontaneous healing of the ACL. Even mild anterior tibial instability, such as in the 4-0

CATT group, inhibited spontaneous healing of the ACL. In a clinical study of ACL

conservative treatment, it was reported that 90% of ACL-injured patients had healed

ACL with knee joints immobilized at 90 degrees by orthotics.[10] On the other hand,

another study showed a lower rate of successful healing in younger ACL-injured

patients because they have higher activity levels compared with middle-age and older

patients.[19] In addition, it was reported that early exercise intervention during the

supraspinatus tendon healing process induced degeneration of the tendon.[20] Therefore,

the mechanical stress on the ACL probably needs to be controlled during the early

phase of the ACL spontaneous healing. During the healing process of the MCL,

vascular-rich granulation tissue invades the rupture area of the ligament to close and

then gradually replace with fibrous tissue.[21] Thus, the increase in ATT probably led to

the failure of healing by pulling apart the ACL remnant and inhibiting the invasion of

granulation tissue, or by re-rupturing the very fragile granulation tissue. Therefore, we

showed that firm braking of the ATT is important in the early healing process after ACL

injury.

It is known that there are sex differences in ACL injury mechanisms, and the

ACL injury rate in women is 2~8 times higher than that in men.[22] Women are more

likely to have knee ligament laxity, which can lead to decreased knee joint stability

during exercise.[23] These sex differences in ACL injury mechanisms[24–26] may also affect

the ACL healing process, so we also investigated sex differences in the ACL healing.

The ACL healing success rate of the 3-0 CATT group was higher in females than in

males. The ATT braking by the nylon suture may have loosened more easily in male,

which would have affected these results. Anterior drawer testing with nylon suture

showed that the ATT braking force in the 3-0 CATT group both males and females was

similar to that of intact immediately after injury. On the other hand, the ATT of male 3-

0 CATT group was higher than that of female at 4 weeks post-injury, although there

was no significant differences. These results suggested that the nylon sutures of the

male CATT group were more likely to loosen, and it had a negative effect on the ACL

healing. It may be caused by the difference in cage activity between male and female

mice. Previous studies have reported that the amount of spontaneous exercise in wheel

running in C57BL/6 mice is higher in females.[27] In addition, a study investigating the

amount of home-cage activity showed that the amount of activity in the dark zone was

significantly higher in females.[28] In contrast, male is known to exhibit more aggressive

behaviors such as fighting and mounting when kept with 2 or more animals.[29] In this

study, we housed both each sex mice with 2-4 mice. Based on these knowledges, male

mice may have been higher intense activity than female in cage, which may have caused

the CATT to loosen and higher mechanical stress on the ACL in male.

As mentioned above, we suggested that firm ATT braking is necessary to achieve

a higher spontaneous healing success rate after ACL injury. Thus, in terms of healing

success rate, increased mechanical stress in the early phase of healing process had a

negative effect. On the other hand, we observed healing of the ACL even in the 4-0

CATT group with higher ATT. In general, mechanical stress on the ligaments promotes

fibroblast activation and ECM synthesis such as collagen.[13] It is known that the

promotion of ECM synthesis improves the mechanical strength of the ligaments. Based

on these knowledges, once the injured ACL in 4-0 CATT group was healed, it exposed

higher mechanical stress during the early ACL healing process. Healed ACL in 4-0

CATT group might have grater quality of the healed ACL such as ECM maturation and

mechanical strength, although had a lower healing success rate. Therefore, we

performed ACL mechanical testing, picrosirius red staining, and immunofluorescence at

4 and 8 weeks to understand the mechanical strength and collagen maturation of the

ACL in each group. We evaluated the maturity of collagen, the main component of the

ECM, from the grayscale of the picrosirius red staining. In addition, we evaluated

expression of type I collagen, the most abundant ECM in intact ligaments, with

immunofluorescence. The intact ACL showed that type I collagen was localized along

the ACL, while irregular and poor collagen localization was observed in both 3-0 and 4-

0 CATT groups. In addition, the mechanical testing results showed that the 3-0 and 4-0

CATT groups had similar mechanical strength of the healed ACL and only 50%

strength compared with intact ACL at both 4 and 8 weeks after injury. To summarize

these results, against our expectations, there was no difference in collagen maturation

and mechanical strength between the 3-0 and 4-0 CATT groups. The reason for these

results is probably that the mechanical stress on the healed ACL was similar in the 3-0

and 4-0 CATT groups after 4 weeks post-injury. Anterior drawer testing showed that

ATT was lower in the 3-0 CATT group than in the 4-0 CATT group immediately after

injury, while ATT increased in both the 3-0 and 4-0 CATT groups after 4 weeks post-

injury, which indicated no difference in mechanical stress in each group. Moreover, the

ACL mechanical strength of both groups was lower than intact. Increased ATT after

4weeks post-injury lead mechanical stress on the healed ACL, but it was not sufficient

to promote healing of the ACL. Low-intensity exercise during the healing process of the

supraspinatus tendon is reported to be beneficial in maintaining structure and promoting

repair in the early stages of healing but is insufficient to promote long-term healing.[30] In

addition, there is an optimal timing for mechanical stress loading to promote tendon and

ligament healing. In ACL-reconstructed mice, it was reported that moderate timing of

loading was more effective in promoting healing than immediate postoperative and

delayed mechanical stress.[27] Thus, there is an optimal amount and timing of mechanical

stress loading during tendon and ligament healing processes. In this study, it was found

that mechanical stress by increasing ATT in the early phase was not sufficient to

promote healing. Therefore, in the future study, we need to investigate the influence of

exercise intervention, i.e., rehabilitation, on promoting healing after spontaneous ACL

healing.

In this study, we have several limitations. First, the present data were evaluated

immediately after injury, and early phase of ACL healing at 4 weeks, and at 8 weeks. It

is not possible to evaluate differences in ATT and the healing process from 0 to 4

weeks. Therefore, we will analyze at finer time points in the future to identify when

CATT is no longer necessary. We will then be able to determine the period when

orthotic management is necessary after ACL injury. Second, differences in the braking

force of the ATT in anterior drawer testing are ex vivo data and do not completely show

the differences in stress on the ACL in vivo. Although we evaluated the mechanical

stress load by the difference of ATT, we will evaluate the knee joint motion during

walking and the amount of activity in the cage as a future study. Then, we can analyze

in detail the mechanical stress loaded on the ACL by the difference in braking force.

Third, in this study, mechanical stress on the healed ACL did not improve mechanical

strength sufficiently. Therefore, it would be necessary to apply a higher load of

mechanical stress using exercise in future studies.

In this study, we investigated the effects of different ATT braking forces on the

ACL spontaneous healing success rate and mechanical strength of the healed ACL in

mice CATT model with different braking sutures. The results showed that firm ATT

braking immediately after injury is necessary for successful healing of the injured ACL.

On the other hand, increased ATT in the early phase was insufficient to improve

mechanical strength and collagen maturation. These results suggested that firm ATT

braking is required on ACL healing at the early phase of post-injury, and then higher

mechanical loading may be necessary to achieve higher ACL properties at the

remodeling phase when the ACL has healed.

Acknowledgments

This work was supported by JSPS KAKENHI (21H03306, 19KK0411), grants from

Tokyo Physical Therapist Association (05094) and the attack grant from Saitama

Prefectural University.

Reference

1.   Woo SL-Y, Vogrin TM, Abramowitch SD. 2000. Healing and Repair of Ligament
Injuries in the Knee: Journal of the American Academy of Orthopaedic Surgeons
8(6):364–372.

2.   Granan L-P, Bahr R, Steindal K, et al. 2008. Development of a National Cruciate
Ligament Surgery Registry: The Norwegian National Knee Ligament Registry.
Am J Sports Med 36(2):308–315.

3.   Bray RC, Leonard CA, Salo PT. 2003. Correlation of healing capacity with
vascular response in the anterior cruciate and medial collateral ligaments of the
rabbit. J. Orthop. Res. 21(6):1118–1123.

4.   Menetrey J, Laumonier T, Garavaglia G, et al. 2011. α-Smooth muscle actin and
TGF-β receptor I expression in the healing rabbit medial collateral and anterior
cruciate ligaments. Injury 42(8):735–741.

5.   Frobell RB, Roos EM, Roos HP, et al. 2010. A Randomized Trial of Treatment for
Acute Anterior Cruciate Ligament Tears. N Engl J Med 363(4):331–342.

6.   Ihara H, Miwa M, Deya K, Torisu K. 1996. MRI of anterior cruciate ligament
healing. J Comput Assist Tomogr 20(2):317–321.

7.   Fujimoto E, Sumen Y, Ochi M, Ikuta Y. 2002. Spontaneous healing of acute
anterior cruciate ligament (ACL) injuries - conservative treatment using an
extension block soft brace without anterior stabilization. Archives of Orthopaedic
and Trauma Surgery 122(4):212–216.

8.   Previ L, Monaco E, Carrozzo A, et al. 2023. Spontaneous healing of a ruptured
anterior cruciate ligament: a case series and literature review. J EXP ORTOP
10(1):11.

9.   Filbay SR, Roemer FW, Lohmander LS, et al. 2023. Evidence of ACL healing on
MRI following ACL rupture treated with rehabilitation alone may be associated
with better patient-reported outcomes: a secondary analysis from the KANON
trial. Br J Sports Med 57(2):91–99.

10. Filbay SR, Dowsett M, Chaker Jomaa M, et al. 2023. Healing of acute anterior
439   cruciate ligament rupture on MRI and outcomes following non-surgical
440   management with the Cross Bracing Protocol. Br J Sports Med :bjsports-2023-
441   106931.

442   11. Kokubun T, Kanemura N, Murata K, et al. 2016. Effect of Changing the Joint
443   Kinematics of Knees With a Ruptured Anterior Cruciate Ligament on the
444   Molecular Biological Responses and Spontaneous Healing in a Rat Model. Am J
445   Sports Med 44(11):2900–2910.

446   12. Gomez MA, Woo SL-Y, Amiel D, et al. 1991. The effects of increased tension on
447   healing medial collateral ligaments. Am J Sports Med 19(4):347–354.

448   13. Zhang J, Wang JH-C. 2013. The Effects of Mechanical Loading on Tendons - An
449   In Vivo and In Vitro Model Study. PLoS ONE 8(8):e71740.

450   14. Eliasson P, Andersson T, Aspenberg P. 2012. Achilles tendon healing in rats is
451   improved by intermittent mechanical loading during the inflammatory phase:
452   TENDON HEALING AND EARLY LOADING. J. Orthop. Res. 30(2):274–279.

453   15. Wada S, Lebaschi AH, Nakagawa Y, et al. 2019. Postoperative Tendon Loading
454   With Treadmill Running Delays Tendon-to-Bone Healing: Immunohistochemical
455   Evaluation in a Murine Rotator Cuff Repair Model. J. Orthop. Res. 37(7):1628–
456   1637.

457   16. Takahata K, Arakawa K, Enomoto S, et al. 2023. Joint instability causes catabolic
458   enzyme production in chondrocytes prior to synovial cells in novel non-invasive
459   ACL ruptured mouse model. Osteoarthritis and Cartilage 31(5):576–587.

460   17. Wobbrock JO, Findlater L, Gergle D, Higgins JJ. 2011. The aligned rank
461   transform for nonparametric factorial analyses using only anova procedures. In:
462   Proceedings of the SIGCHI Conference on Human Factors in Computing Systems.
463   New York, NY, USA: Association for Computing Machinery. p 143–146

464   18. Jordão S, Cortes N, Gomes J, et al. 2022. Synchronization performance affects gait
465   variability measures during cued walking. Gait & Posture 96:351–356.

466   19.   Ihara H, Kawano T. 2017. Influence of Age on Healing Capacity of Acute Tears of
467   the Anterior Cruciate Ligament Based on Magnetic Resonance Imaging
468   Assessment. Journal of Computer Assisted Tomography 41(2):206.

469   20.   Chen Y, Zhang T, Wan L, et al. 2021. Early treadmill running delays rotator cuff
470   healing via Neuropeptide Y mediated inactivation of the Wnt/β-catenin signaling.
471   Journal of Orthopaedic Translation 30:103–111.

472   21.   Chamberlain CS, Crowley E, Vanderby R. 2009. The spatio-temporal dynamics of
473   ligament healing. Wound Repair and Regeneration 17(2):206–215.

474   22.   Spindler KP, Wright RW. 2008. Anterior Cruciate Ligament Tear. N Engl J Med
475   359(20):2135–2142.

476   23.   Park S-K, Stefanyshyn DJ, Loitz-Ramage B, et al. 2009. Changing Hormone
477   Levels during the Menstrual Cycle Affect Knee Laxity and Stiffness in Healthy
478   Female Subjects. Am J Sports Med 37(3):588–598.

479   24.   Ford KR, Myer GD, Hewett TE. 2003. Valgus Knee Motion during Landing in
480   High School Female and Male Basketball Players. Medicine & Science in Sports
481   & Exercise 35(10):1745.

482   25.   Jordan SS, DeFrate LE, Nha KW, et al. 2007. The in vivo kinematics of the
483   anteromedial and posterolateral bundles of the anterior cruciate ligament during
484   weightbearing knee flexion. Am J Sports Med 35(4):547–554.

485   26.   Shultz SJ, Sander TC, Kirk SE, Perrin DH. 2007. Sex differences in knee joint
486   laxity change across the female menstrual cycle.

487   27.   Chen H, Li S, Xiao H, et al. 2021. Effect of Exercise Intensity on the Healing of
488   the Bone-Tendon Interface: A Mouse Rotator Cuff Injury Model Study. Am J
489   Sports Med 49(8):2064–2073.

490   28.   Bains RS, Forrest H, Sillito RR, et al. 2023. Longitudinal home-cage automated
491   assessment of climbing behavior shows sexual dimorphism and aging-related
492   decrease in C57BL/6J healthy mice and allows early detection of motor
493   impairment in the N171-82Q mouse model of Huntington's disease. Front. Behav.
494   Neurosci. 17:1148172.

29. Lidster K, Owen K, Browne WJ, Prescott MJ. 2019. Cage aggression in group-
housed laboratory male mice: an international data crowdsourcing project. Sci Rep
9(1):15211.

30. Camp CL, Lebaschi A, Cong G-T, et al. 2017. Timing of Postoperative
Mechanical Loading Affects Healing Following Anterior Cruciate Ligament
Reconstruction: Analysis in a Murine Model. The Journal of Bone and Joint
Surgery 99(16):1382–1391.

Table Legend

Table 1. We defined ACL spontaneous healing from HE stained images. 3-0 CATT

group showed higher healing success rate than 4-0 CATT group at 4 and 8 weeks after

injury.

	4 weeks				8 weeks			
	3-0 CATT		4-0 CATT		3-0 CATT		4-0 CATT	
	Male	Female	Male	Female	Male	Female	Male	Female
Healed	5	4	2	1	3	4	1	4
Non-healed	2	1	3	3	4	0	5	3
Success rates of ACL healing (%)	71.4	80.0	40.0	25.0	42.9	100.0	16.7	57.1

Figure Legend

Figure 1. Experimental Design. Male and female 8-9 weeks old C57BL/6 mice were

ruptured the ACL and classified to 3-0 CATT, 4-0 CATT and ACL-R group. Mice were

analyzed via anterior drawer testing and histology immediately after injury, and anterior

drawer testing, histology, and mechanical testing at 4- and 8-weeks post-injury.

Figure 2. Non-invasive ACL-rupture. We fixed the hindlimb with the knee joint at 90

degrees and ruptured the ACL by pushing the femur backward against the tibia. (A, B)

HE images of intact ACL (C) and immediately after ACL rupture (D) (Scale bar = 300

μm). Controlled Anterior Tibial Translation (CATT) model. Tightly looped by single 3-

0 or 4-0 nylon suture through the femoral and tibial bone holes to control the ATT.

Figure 3. Nylon suture experiments. The load to failure (A) and stiffness (B) of the

nylon suture in failure testing, and the number of cycles at the point of suture rupture in

cyclic testing (C) were recorded. (*$p < 0.05$, **$p < 0.01$)

Figure 4. Anterior drawer testing was performed with the knee joint fixed at 90° via the

self-made anterior drawer device (A). Representative X-ray images of 3-0 CATT group

(B) and ACL-R group (C). We evaluated the braking force of the ATT for each groups

immediately after injury (D) and at 4 (E) and 8 weeks (F), and anterior tibial instability

of ACL healed knee at 4 (G) and 8 weeks (H).

Figure 5. HE stained images of the 3-0 CATT (A, B, C, and D), 4-0 CATT (E, F, G, and H), ACL-R groups (I, J, K, and L) at 4 and 8 weeks (Scale bar = 300 μm).

Figure 6. Picrosirius red stained (A) and polarized image (B) of the ACL. Picrosirius

red images were converted to grayscale images (C) and specified ACL area as ROI (D).

Collagen maturation of the ACL was evaluated at 4 (E) and 8 weeks (F) after injury

(Scale bar = 300 μm).

Figure 7. Immunofluorescence images of the intact ACL, 3-0 CATT, 4-0 CATT groups

at 4 and 8 weeks: DAPI (blue) and Collagen type I (green). The images were taken with

a 20× magnification lens (Scale bar = 300 μm).

Figure 8. To perform biomechanical testing, the soft tissues of the knee except the ACL

were dissected and the femur and tibia were fixed in tubes (A). The load to failure (A)

of the healed ACL at 4 (B) and 8 (C) weeks after injury were recorded.