# Title Page Template

**Title of the Manuscript**

**Early and Adequate Braking of Abnormal Tibial Translation Determines Spontaneous Healing Success of the Anterior Cruciate Ligament in a Murine Model.**

---

**Authors**

Riku Saito[1], Kazuki Nakayama[1], Yuna Usami[1], Saaya Enomoto[2], Koyo Nogi[1], Takanori Kokubun[1, 3]

---

**Affiliations**

1. Graduate School of Health, Medicine, and Welfare, Saitama Prefectural University, Saitama, Japan

2. Saitama Rehabilitation Center

3. Department of Physical Therapy, School of Health and Social Services, Saitama Prefectural University, Saitama, Japan

---

**Corresponding Author**

**Name:** Takanori Kokubun, P.T., Ph.D.

**Email:** kokubun-takanori@spu.ac.jp

**Address:**

Saitama Prefectural University

Graduate School of Health, Medicine, and Welfare

820 San-nomiya, Koshigaya-Shi,

Saitama, 343-8540, Japan

______________

**Keywords**

Anterior cruciate ligament; spontaneous healing; conservative treatment; braking;
mechanical stress

______________

**Declarations**

*(When not applicable, please indicate "N/A")*

• **Data availability statement:** The data that support the findings of this study
are available on request from the corresponding author. The data are not
publicly available due to privacy or ethical restrictions.

• **Funding statement:** This work was supported by JSPS KAKENHI
(21H03306, 19KK0411), the grant from Tokyo Physical Therapist Association
(05094), and the attack grant from Saitama Prefectural University.

• **Conflict of interest disclosure:** N/A

• **Authors' contributions:** All authors approved the final version to be
published. Study design: RS and TK; data collection and statistical analysis:
RS, KN, YU, SE, KN; manuscript composition RS, and TK.

• **Patient consent statement** (if applicable): N/A

• **Permission to reproduce material from other sources** (if applicable): N/A

• **Ethics statement:** The University Animal Experiment Ethics Committee
approved all experiments (2021-9).

• **For clinical trials** (if applicable)**:** N/A

- **Level of evidence:** N/A

________________________________________

- **List of abbreviations in the text: List of Abbreviations**

ACL: Anterior Cruciate Ligament

ACL-R: Anterior Cruciate Ligament Rupture

ANOVA: Analysis of Variance

ART: Aligned Rank Transform

ATT: Anterior Tibial Translation

CATT: Controlled Anterior Tibial Translation

CBP: Cross Bracing Protocol

COL1A1: Type I Collagen

ECM: Extra-Cellular Matrix

HE: Hematoxylin and Eosin

MCL: Medial Collateral Ligament

PsR: Picrosirius Red

________________________

*Abstract*

Purpose: The purpose of this study was to elucidate the acceptable range of mechanical stresses

during the early phase of spontaneous anterior cruciate ligament (ACL) healing. We developed

models with varying degrees of controlled anterior tibial translation (CATT) using nylon sutures of

two diameters.

Methods: Male and female C57BL/6 mice were ruptured ACL and randomly classified into three

groups: Tight-CATT; controlled anterior tibial translation (ATT) with 3-0 nylon suture, Loose-

CATT; controlled ATT with 4-0 nylon suture, and ACL-Rupture (ACL-R) group; not controlled

ATT. Mice were sacrificed and analyzed immediately after injury and at 4 and 8 weeks. We

evaluated the braking force to control the ATT of each knee by anterior drawer testing, the success

rate of the ACL healing, collagen maturation and COL1A1 expression with immunofluorescence in

the healed ACL, and the mechanical strength of the healed ACL.

Results: The Tight-CATT group showed a higher success rate of ACL healing than the Loose-CATT

group at 4 and 8 weeks. However, collagen maturation and mechanical properties of healed ACL did

not differ between the Tight-CATT and Loose-CATT groups.

Conclusion: Our results suggested that loose ATT braking immediately after injury is a negative

factor for the healing of the completely ruptured ACL. On the other hand, tightly ATT in the early

healing phase was insufficient to improve mechanical strength and collagen maturation. Therefore,

our results suggest that tightly controlling abnormal tibial translation in the early phase promotes

spontaneous ACL healing, and that it may be necessary to apply higher mechanical stress in the later

stages to achieve greater healing.

*Introduction*

Anterior cruciate ligament (ACL) rupture is one of the most common knee injuries across

all generations. In the United States, it is estimated to occur in more than 100,000 cases per year [1,

2] and impose a substantial economic burden, with annual treatment and related societal costs

estimated in the multi-billion-dollar range [3,4]. For a long time, researchers considered that injured

ACLs had a low capacity for spontaneous healing due to the unique environment within the joint

capsule [5–7]. Therefore, ACL reconstruction is the current gold standard treatment. ACL

reconstruction can restore static and dynamic knee stability following ACL injury and facilitate

return to pre-injury levels of activity [8]. However, several reviews have reported that ACL

reconstruction does not yield superior patient-reported outcomes compared with rehabilitation alone

and provides little benefit in preventing the development of knee osteoarthritis [9,10]. Recently, the

proportion of older patients sustaining ACL injuries has increased relative to that of younger patients

[11–13]. ACL reconstruction in older individuals has been reported to be associated with an

increased risk of developing knee osteoarthritis, and the appropriateness of surgical treatment in this

population therefore remains controversial [14,15].

Over the past several decades, growing evidence has suggested that the ACL has the

potential to heal spontaneously without surgical intervention [16–20]. Ihara et al. reported that ACLs

were healed spontaneously by minimizing abnormal sagittal deviation between the femur and tibia

using a knee brace following ACL injury [16,17]. Subsequently, Filbay et al. reported clinical

outcomes of spontaneous ACL healing using the Cross Bracing Protocol (CBP), which maintained

the knee at 90 degrees in patients with ACL injuries [21]. They showed that 90% of patients had

evidence of healing on MRI after a three-month post-injury period. In addition, we previously

showed that controlling anterior tibial translation following ACL injury enables spontaneous ACL

healing in a rodent model [22]. Therefore, non-surgical treatment will probably be a valuable

therapeutic option, particularly for patients who prefer to avoid surgical treatment. However,

spontaneously healed ACLs that had undergone anterior tibial displacement formed scar tissue and

exhibited lower mechanical strength than normal tissue. The next step is to identify the conditions

that promote healing of the spontaneously healed ACL. It is well established that appropriate

mechanical stress during ligament and tendon healing promotes tissue maturation and recovery of

mechanical strength [23–25]. Thus, understanding the acceptable range of mechanical stress,

especially during the early phase of the ACL spontaneous healing response, is important for

establishing non-surgical treatment and appropriate management for ACL-injured patients.

Therefore, the purpose of this study was to elucidate the acceptable braking condition after

ACL injury. To achieve this goal, we developed two types of controlled anterior tibial translation

(CATT) models using nylon sutures of two different diameters. Suture differences conferred varying

degrees of anterior tibial translation (ATT) stability in murine knees. We hypothesized that a mild

increase in mechanical stress during the early phase of ACL healing does not inhibit healing but

rather contributes to improved tissue maturation and mechanical strength recovery of the healed

ACL.

*Methods*

*Experimental Design*

The University Animal Experiment Ethics Committee approved all experiments. We used

male (n=66) and female (n=66) 8-9 weeks old C57BL/6 mice (n=132). A total of 132 mice were

ruptured the left ACL and then were randomly classified into three groups: Tight-CATT group

(Figure 1B); controlled ATT with 3-0 nylon suture (n=57), Loose-CATT group (Figure 1C);

controlled ATT with 4-0 nylon suture (n=55), and ACL-R (ACL-Rupture) group (Figure 1D); not

controlled ATT (n=20). We used the right hindlimb as the intact group. All mice were housed in

plastic cages under a 12-h light/dark cycle. The mice were allowed unrestricted movement within the

cage and had free access to food and water. Three animals from both the Tight-CATT and Loose-

CATT groups were excluded from the experiment due to tibial fractures when the ACL ruptured,

and the abnormal findings during housing. Thus, a total of 126 mice were included in this study, and

were sacrificed immediately after injury for anterior drawer testing and histology (n=20); at 4 (n=52)

and 8 weeks (n=54), for anterior drawer testing, histology, immunofluorescence, and mechanical

testing.

*Nylon Suture Experiments*

Before creating the CATT models, we measured the mechanical properties of each nylon

suture using a mechanical testing machine (Univert. CellScale Biomaterial Testing, Canada). The

purpose of this experiment was to investigate the differences in mechanical properties between 3-0

and 4-0 nylon sutures (Akiyama Medical MFG, Co., Ltd., Tokyo, Japan). Each nylon suture was

measured in a double layer since a nylon suture was looped outside the joint to control ATT in the

CATT model. In the failure testing, we stretched the nylon sutures at a preload of 0.1 N and a rate of

20 mm/min until they ruptured. During the test, load (N) and displacement (mm) were recorded, and

we measured the load at failure (N) and the stiffness (N/mm). Then, we performed the cyclic test to

evaluate viscoelasticity. Nylon sutures were repeated with 5 seconds of stretching up to 90%, 80%,

70%, and 60% of the failure load of each nylon suture, followed by 5 seconds of relaxation (1 cycle:

10 seconds). We recorded the number of cycles at the point of rupture.

*Procedure for Creating the Model*

159  All mice were anesthetized with a combination of anesthetics (medetomidine, 0.375 mg/kg;

160  midazolam, 2.0 mg/kg; and butorphanol, 2.5 mg/kg) for sedation and pain control, and then we

161  ruptured the ACL non-invasively according to a previous study [26]. We fixed the hindlimb with the

162  knee joint at 90 degrees and ruptured the ACL by pushing the femur backward against the tibia. In

163  non-invasive ACL ruptures, the ligament is generally ruptured in the mid-substance region (Figure

164  1E). Then, we created bone tunnels at the femoral condyle and proximal tibia using a 26G needle

165  (Terumo Co., Ltd., Tokyo, Japan). In the Tight- and Loose-CATT groups, we tightly looped a single

166  3-0 or 4-0 nylon suture through the two tunnels to control the ATT. In the ACL-R group, we loosely

167  looped a 3-0 nylon suture through the two tunnels not to control the ATT.

169  *Anterior Drawer Testing*

170  We performed anterior drawer testing to evaluate the braking force to control the ATT of

171  each knee, carefully removing the muscle from the knee while retaining the suture. Then, we fixed

172  the knee joint at 90 degrees on the self-made anterior drawer testing device. The tibia was pulled

173  forward with a 0.05 kgf constant force spring (Sanko Spring Co., Ltd., Fukuoka, Japan), and X-ray

174  images of the sagittal plane of the knee joint were taken. We measured ATT from X-ray images

175  using ImageJ (National Institutes of Health, Bethesda, MD, USA) to evaluate the braking force

176  exerted by nylon sutures. The ATT was defined as the perpendicular line from the most prominent

point of the femoral condyle to the long axis of the tibia. We then removed the nylon suture and

again measured the ATT to evaluate the potential characteristics of the knee stability with healed

ACL. The knees in which the ACL was not healed in histology were excluded from the latter

analysis.

*Histology*

After anterior drawer testing, knee samples were immediately fixed (24-48h) in 4%

paraformaldehyde and decalcified with 10% ethylenediaminetetraacetic acid (Sigma-Aldrich Japan,

Tokyo, Japan) for 2 weeks. Finally, they were immersed in 5% and 30% sucrose for 4 and 12 hours,

respectively, and then freeze-embedded using OCT compound (Sakura Finetek Japan Co., Ltd.,

Tokyo, Japan). Each sample was sectioned at 12μm using a Cryostar NX70 (Thermo Fisher

Scientific, Kanagawa, Japan). We performed Hematoxylin & Eosin (HE) staining to observe ACL

continuity and morphological features, and Picrosirius Red (PsR) staining to evaluate the collagen

maturity of the healed ACL. We observed the stained sections with the BZ-X710 microscope

(Keyence, Osaka, Japan). We determined whether the ACL was healed from HE images obtained at

4 or 8 weeks post-injury. We defined ACLs with observed continuity as "healed" (Figure 1F), and

those with lost continuity as "non-healed" (Figure 1G). The success rate of ACL healing in each

group was calculated based on the HE images. PsR images converted to grayscale images (256

degrees) showed collagenous tissue as 1-255 on the grayscale and non-collagenous tissue as 0 on the

grayscale. A higher gray scale indicates higher ACL collagen maturation [27,28]. Using the BZ-

X710 analysis software, the entire ACL areas in the PsR polarization images were defined as

Regions of Interest (ROIs) and then converted to grayscale. We used the average grayscale value in

the ROI as the representative value of the healed ACL.

*Immunofluorescence*

We performed immunofluorescence to evaluate the expression of type I collagen

(COL1A1) in the healed ACL. ACL cryosections were blocked at room temperature for 1 hour using

DAKO Antibody Diluent with Background Reducing Components (Agilent Technologies, Inc., CA,

USA). The anti-COL1A1 primary antibody was incubated with a Goat polyclonal antibody (1:200;

AB758, Merck, Sigma-Aldrich, Taufkirchen, Germany) at 4°C overnight. The secondary antibody

was incubated with anti-Goat Rabbit antibody Alexa Fluor 488 (1:500, ab150141, Abcam,

Cambridge, UK) for 4 hours at room temperature. All sections were examined macroscopically

using a BZ-X710 microscope under controlled imaging conditions.

*Biomechanical Testing*

The failure load testing was performed to evaluate the mechanical strength of the healed

ACL. The harvested knee joints were dissected and frozen at -70°C. After thawing, all soft tissues

were removed except the ACL, and the femur and tibia were fixed in 1.5-mm tubes. The tubes were

then fixed to the mechanical testing machine Univert in the extended position. The ACL was

stretched at a preload of 0.1 N and a rate of 10 mm/min until it was ruptured. We recorded the load-

to-failure force of the ACL.

*Statistical Analysis*

We performed statistical analyses of the results from nylon suture experiments, anterior

drawer testing, grayscale analysis, and mechanical testing using R. The Shapiro-Wilk test was used

to verify the normality of all the analyzed data. Mann-Whitney U-tests were performed for the data

from the nylon suture experiments. A two-way analysis of variance (ANOVA) was then performed

to confirm significant differences among groups and to assess sex differences in the other results.

Because all data were non-normally distributed, an Aligned Rank Transform (ART) [29,30] was

performed before ANOVA. Based on the ANOVA results, the Steel-Dwass test was used for

multiple comparisons among all groups when an interaction was found. When no interaction was

found, multiple comparisons were performed within factors for which a main effect was found.

Statistical significance was set at $p < 0.05$.

*Results*

*Mechanical Properties of Two Nylon Sutures*

233   The results showed that the 3-0 nylon sutures had better mechanical properties. The load to

failure and stiffness were significantly higher for 3-0 nylon sutures ($p < 0.05$) (Figure 2A, B). In

cyclic testing, the 3-0 nylon sutures exhibited higher viscoelasticity and greater resistance to

elongation. The 4-0 nylon sutures ruptured in 3 cycles at 90% of the failure load, 12 cycles at 80%,

107 cycles at 70%, and 791 cycles at 60%. The 3-0 nylon sutures ruptured in 6 cycles at 90% and in

197 cycles at 80%. 3-0 nylon sutures did not rupture even after 2,000 cycles at loads lower than 70%

(Figure 2C).

*Magnitude of Anterior Translation of Tibia*

242   First, in the knee joints with nylon sutures for controlling ATT, interactions between groups

and sex were observed at all timepoints ($p < 0.01$). Immediately after injury, the ATT was smallest

in the intact and Tight-CATT groups. The Loose-CATT group showed a higher ATT than the Tight-

CATT group, and the ACL-R group showed the highest ATT. However, there were no significant

differences among the groups or by sex (Figure 3C). In males at 4 and 8 weeks after injury, the ATT

was smallest in the intact group, followed by the Tight-CATT and Loose-CATT groups, which

showed similar ATT. The ACL-R group had the highest ATT. In contrast, in females, the ATT of

the Tight-CATT showed a trend to be smaller than that of the Loose-CATT. However, there were no

significant differences among the groups or by sex (Figures 3D, E).

Secondly, for anterior tibial instability of the knee with a healed ACL, fewer knees were

included in the Loose-CATT group because we excluded knees with non-healed ACLs from the

CATT group in this analysis. Interactions between groups and sex were observed at 4 weeks and 8

weeks ($p < 0.001$). In males at 4 and 8 weeks, intact had the smallest ATT, and the Tight-CATT and

Loose-CATT groups had similar ATT but higher ATT than intact. ACL-R showed the highest ATT.

In females, the ATT of the Tight-CATT showed a trend to be smaller than that of the Loose-CATT.

However, there were no significant differences among the groups or by sex (Figures 3F, G).

*ACL Healing Success Rate and Collagen Maturation of the ACL*

First, we reported the success rates of ACL healing in the Tight- and Loose-CATT groups at

4- and 8-week post-injury (Table 1). The Tight-CATT group showed a higher healing rate for both

males and females (42.9~100%), while the Loose-CATT group showed a lower healing rate

(17~57.1%) and no healed ACL in the ACL-R group (0%). Next, we showed HE-stained images of

the healed ACL. Compared to the intact ACL with parallel and regular nuclear arrangement, the

healed ACL showed heterogeneous nuclear aggregation not only in the healed region but also in the

ACL remnant region. It was also observed at 4- and 8-week post-injury (Figure 4A-H).

In collagen maturation of the healed ACL, there are no interactions observed at both 4- and

8-week post-injury. On the other hand, a significant main effect was observed in the group: the

Tight-CATT group had a significantly lower grayscale than intact ($p < 0.001$) at 4 weeks, and the

Tight- and Loose-CATT groups showed a significantly lower grayscale than intact ($p < 0.01$) at 8

weeks (Figure 5E, F).

*Type I Collagen Content of Healing ACL*

We performed immunofluorescence for COL1A1, a major ECM component, to assess ECM

maturity in the healed ACL. The green area indicates COL1A1, and the blue area indicates the

nucleus. In the intact ACL, we observed COL1A1 expression at the entire ACL region and nuclei

aligned longitudinally of the ACL (Figure 6A, B). On the other hand, the healed ACLs of the Tight-

and Loose-CATT groups showed limited expression outside the healed ACL and an irregular

orientation of COL1A1 compared with the intact ACL. In addition, the Tight- and Loose-CATT

groups showed higher and irregular nuclei aggregation throughout the ACL (Figure 6C-F). These

collagen features of the ACL were commonly observed at 4- and 8-weeks post-injury (Shown in

Figure 6 is at 8 weeks post-injury).

*Mechanical Properties of Healed ACL*

We fixed the femur and tibia in 1.5-mm tubes (Figure 7A) and performed mechanical testing

to evaluate the mechanical strength of the healed ACL. The number of ACLs for testing in the

Loose-CATT group was fewer because only the knees with confirmed ACL re-connections were

included when the knees were dissected. No interaction was observed at both 4- and 8-week post-

injury. On the other hand, a significant main effect was observed in the group: at 4 weeks, intact had

a significantly higher mechanical strength than the Tight-CATT ($p < 0.01$) and Loose-CATT group

($p < 0.05$) (Figure 7B), and at 8 weeks, intact had a significantly higher mechanical strength than the

Tight-CATT group ($p < 0.01$) (Figure 7C). However, the healed ACLs in the Tight- and Loose-

CATT groups exhibited similar mechanical strength and were approximately 50% of normal ACL

strength.

*Discussion*

The most important finding of this study was that loosening of the controlling ATT

immediately after ACL injury is a negative factor for spontaneous healing of the ruptured ACL. The

results of the Loose-CATT group showed that increased mechanical stress immediately after injury

negatively affects ACL healing response. Conversely, the healed ACLs in both the CATT group

showed lower mechanical properties than those in the Intact group, and there were no significant

differences in mechanical properties or histological features between tight and loose conditions.

Overall, our results suggest that appropriate force control may be necessary to improve the success

rate of healing and the mechanical properties of the healed ACL.

In this study, we developed CATT models with varying levels of ATT braking force, using

two types of nylon suture, immediately after ACL injury. Results of the nylon suture experiment

showed that the 4-0 nylon suture used in the Loose-CATT group had higher suture elasticity

compared to the 3-0 nylon suture used in the Tight-CATT group. Subsequently, we confirmed in the

anterior drawer test that the Loose-CATT group exhibited a lower braking force than the Tight-

CATT group immediately after ACL injury. Therefore, the Loose-CATT group was defined as the

group with early exposure to the higher mechanical stress immediately after ACL injury. A clinical

study showed a lower rate of successful healing in younger ACL-injured patients because they have

higher activity levels compared with middle-aged and older patients[17], which supports our

findings. On the other hand, another study reported that 90% of ACL-injured patients had healed

ACLs with knee joints immobilized at 90 degrees using orthotics [21]. Furthermore, while proper

mechanical stress after ligament and tendon injury promotes healing [23,24,31–33], excessive

mechanical loading impairs biomechanical properties and tissue maturation [34–36]. In the ligament-

healing process, vascular-rich granulation tissue invades the ligament rupture site, initially sealing it

and then gradually replacing it with fibrous tissue [37]. The increase in ATT in Loose-CATT may

have increased mechanical stress on the ACL immediately after injury and physiologically inhibited

the formation of a fragile provisional matrix. In addition, loosening of the ATT would induce

abnormal biological modulation in the ACL healing. Excessive mechanical stress on ligaments and

tendons induces fibrous tissue to transition to an abnormal matrix, characterized by increased

proteoglycan content and type III collagen.[38,39]. In the wound-healing process, abnormal

mechanical stress prolongs inflammation, delays wound closure, and is a significant factor in

hypertrophic scar formation [40,41]. In our previous data, we reported that controlling abnormal

tibial motion after ACL injury can reduce MMPs that cause matrix degradation in the ACL

[22,42,43]. Therefore, increased ATT in the early post-injury period may have disrupted the fragile

ligaments physiologically or promoted the catabolic process of the injured ACL biochemically. Our

results suggested that loose braking by ATT inhibits early ACL healing.

Interestingly, we also found sex differences in ACL healing success rates. It is known that

there are sex differences in ACL injury mechanisms, and the ACL injury rate in women is 2~6 times

higher than that in men [3,44,45]. On the other hand, there is insufficient consensus regarding sex

differences in clinical outcomes after injury [46,47]. Surprisingly, our findings showed that the ACL

healing success rate was higher in females than in males. While the similar ATT braking force in the

male and female Tight-CATT group was observed immediately after injury, the ATT braking of the

male Tight-CATT group decreased after 4 weeks post-injury, compared with the female. These

results may be attributable to differences in cage activity between male and female mice. Previous

studies have reported that the amount of spontaneous exercise in wheel-running is higher in females

among C57BL/6 mice [48]. Additionally, a study examining home-cage activity levels found that

females exhibited significantly higher activity in the dark time [49]. In contrast, males exhibit more

aggressive behaviors, such as fighting and mounting, when kept with two or more animals [50]. In

this study, we housed 2-4 mice in one cage for both sexes. Based on this knowledge, male mice

might show more intense activity than females in a cage. Additionally, standing on two limbs in a

fighting or defending motion is more unstable than walking on four limbs [51,52], which would

increase stress on the knee joints, particularly the healing ACL. Thus, the aggressive behavioral

characteristics of male mice may have contributed to the reduced ACL healing rate. This finding

underscores the importance of balancing the quality and quantity of exercise during ACL healing.

Further study is needed to clarify the relationship between behavioral characteristics of the mice and

ACL healing.

We hypothesized that the increased mechanical stress on the healing ACL immediately after

injury would promote collagen maturation and mechanical strength, even if it led to a low success

rate in ACL healing. However, against our expectations, there was no difference in the quality of the

healed ACL between the Tight- and Loose-CATT groups. The healed ACLs in Tight- and Loose-

CATT groups showed similar collagen maturation and mechanical strength, and inferior properties

compared to the intact ACL. Accordingly, tight ATT braking is important for a higher success rate

of ACL healing in the early phase. On the other hand, it is suggested that an increase in mechanical

stress at the level of loosening does not promote ACL healing, because loosening occurs in both

Tight- and Loose-CATT and the quality of the healed ACL shows no difference. Previous studies

have shown that low-intensity exercise during the healing process of the supraspinatus tendon is

beneficial for maintaining structure and promoting repair in the early stages of healing, but is

insufficient to promote long-term healing [25]. In ACL-reconstructed mice, proper timing of

mechanical stress was more effective at promoting healing than immediate postoperative or delayed

mechanical stress [31]. This study revealed that the mild increase in mechanical force induced by

loosening is insufficient to improve ECM synthesis or the mechanical strength of the healed ACL.

Although our results suggest the importance of tight control of abnormal tibial translation in the

367 early phase after ACL injury, the optimal mechanical force for promoting ACL healing remains

368 unclear. Future study will focus on this point.

369  In this study, we have several limitations. First, mice are a commonly used model for

370 research on ligament and tendon healing. However, the structure and alignment of the mouse knee,

371 which is always flexed, differ from those of the human knee, which is typically extended. Therefore,

372 the mice are not a model that completely reconstructs the human condition. Second, the present data

373 were evaluated immediately after injury and at 4 and 8 weeks after injury. It was impossible to

374 evaluate differences in ATT and the healing process from 0 to 4 weeks. Therefore, we will analyze

375 at finer time points in the future to identify when CATT is no longer necessary. We will then be able

376 to determine the period during which orthotic management is necessary following ACL injury.

377 Finally, the current study examined, first, how increased mechanical stress affects ACL healing and

378 subsequent properties. Hence, it was unable to elucidate in detail the mechanisms by which

379 increased mechanical stress impaired ACL healing in the early stages of healing. In the future, we

380 will analyze biochemical data and intracellular changes up to 4 weeks after ACL injury to elucidate

381 the mechanism of mechanical transduction in ACL healing.

382  In this study, we investigated the effects of varying ATT braking forces on the ACL

383 spontaneous-healing success rate and the quality of the healed ACL in mice using CATT models

384 with different sutures. The results showed that loose braking of the ATT immediately after injury is

a negative factor for the healing of the completely injured ACL. Conversely, the increasing

mechanical force induced by loosening is insufficient to improve the ECM synthesis or mechanical

properties of the healed ACL. Still, the mechanism underlying spontaneous ACL healing remains

unclear; controlling the mechanical force on the ACL should play a key role in ACL healing and

maturation. We provided essential insights into the state of ACL healing, which are critical for

elucidating ACL healing mechanisms and informing conservative rehabilitation.

*Acknowledgments*

This work was supported by JSPS KAKENHI (21H03306, 19KK0411), the grant from Tokyo

Physical Therapist Association (05094) and the attack grant from Saitama Prefectural University.

*Reference*

1. Hewett TE, Myer GD, Ford KR, et al. 2016. Mechanisms, prediction, and prevention of ACL injuries: Cut risk with three sharpened and validated tools. Journal Orthopaedic Research 34(11):1843–1855.

2. Hewett TE, Lindenfeld TN, Riccobene JV, Noyes FR. 1999. The Effect of Neuromuscular Training on the Incidence of Knee Injury in Female Athletes. Am J Sports Med 27(6):699–706.

3. Spindler KP, Wright RW. 2008. Anterior Cruciate Ligament Tear. N Engl J Med 359(20):2135–2142.

4. Mather RC, Koenig L, Kocher MS, et al. 2013. Societal and Economic Impact of Anterior Cruciate Ligament Tears. Journal of Bone and Joint Surgery 95(19):1751–1759.

5. Murray MM, Fleming BC. 2013. The Biology of Anterior Cruciate Ligament Injury and Repair: Kappa Delta Ann Doner Vaughn Award Paper 2013. J Orthop Res 31(10):1501–1506.

6. Bray RC, Leonard CA, Salo PT. 2003. Correlation of healing capacity with vascular response in the anterior cruciate and medial collateral ligaments of the rabbit. J. Orthop. Res. 21(6):1118–1123.

7. Menetrey J, Laumonier T, Garavaglia G, et al. 2011. α-Smooth muscle actin and TGF-β receptor I expression in the healing rabbit medial collateral and anterior cruciate ligaments. Injury 42(8):735–741.

8. Migliorini F, Vecchio G, Eschweiler J, et al. 2023. Reduced knee laxity and failure rate following anterior cruciate ligament reconstruction compared with repair for acute tears: a meta-analysis. J Orthop Traumatol 24(1):8.

9. Hohmann E, Tetsworth K, Glatt V. 2023. Anterior cruciate ligament reconstruction results in better patient reported outcomes but has no advantage for activities of daily living or the subsequent development of osteoarthritis. A systematic review and meta-analysis. The Knee 41:137–149.

10. Monk AP, Davies LJ, Hopewell S, et al. 2016. Surgical versus conservative interventions for treating anterior cruciate ligament injuries. Cochrane Database of Systematic Reviews 2016(4) [cited 2023 June 6] Available from: http://doi.wiley.com/10.1002/14651858.CD011166.pub2.

424     11.    Sanders TL, Maradit Kremers H, Bryan AJ, et al. 2016. Incidence of Anterior Cruciate
425            Ligament Tears and Reconstruction: A 21-Year Population-Based Study. Am J Sports Med
426            44(6):1502–1507.

427     12.    Mall NA, Chalmers PN, Moric M, et al. 2014. Incidence and Trends of Anterior Cruciate
428            Ligament Reconstruction in the United States. Am J Sports Med 42(10):2363–2370.

429     13.    Granan L-P, Bahr R, Steindal K, et al. 2008. Development of a National Cruciate Ligament
430            Surgery Registry: The Norwegian National Knee Ligament Registry. Am J Sports Med
431            36(2):308–315.

432     14.    Best MJ, Zikria BA, Wilckens JH. 2021. Anterior Cruciate Ligament Injuries in the Older
433            Athlete. Sports Health: A Multidisciplinary Approach 13(3):285–289.

434     15.    Salzler MJ, Chang J, Richmond J. 2018. Management of Anterior Cruciate Ligament Injuries
435            in Adults Aged >40 Years. JAAOS - Journal of the American Academy of Orthopaedic
436            Surgeons 26(16):553.

437     16.    Ihara H, Miwa M, Deya K, Torisu K. 1996. MRI of anterior cruciate ligament healing. J
438            Comput Assist Tomogr 20(2):317–321.

439     17.    Ihara H, Kawano T. 2017. Influence of Age on Healing Capacity of Acute Tears of the
440            Anterior Cruciate Ligament Based on Magnetic Resonance Imaging Assessment. Journal of
441            Computer Assisted Tomography 41(2):206.

442     18.    Fujimoto E, Sumen Y, Ochi M, Ikuta Y. 2002. Spontaneous healing of acute anterior cruciate
443            ligament (ACL) injuries - conservative treatment using an extension block soft brace without
444            anterior stabilization. Archives of Orthopaedic and Trauma Surgery 122(4):212–216.

445     19.    Previ L, Monaco E, Carrozzo A, et al. 2023. Spontaneous healing of a ruptured anterior
446            cruciate ligament: a case series and literature review. J EXP ORTOP 10(1):11.

447     20.    Filbay SR, Roemer FW, Lohmander LS, et al. [date unknown]. Evidence of ACL healing on
448            MRI following ACL rupture treated with rehabilitation alone may be associated with better
449            patient-r-eported outcomes: a secondary analysis from the KANON trial.

450     21.    Filbay SR, Dowsett M, Chaker Jomaa M, et al. 2023. Healing of acute anterior cruciate
451            ligament rupture on MRI and outcomes following non-surgical management with the Cross
452            Bracing Protocol. Br J Sports Med :bjsports-2023-106931.

22.   Kokubun T, Kanemura N, Murata K, et al. 2016. Effect of Changing the Joint Kinematics of
Knees With a Ruptured Anterior Cruciate Ligament on the Molecular Biological Responses
and Spontaneous Healing in a Rat Model. Am J Sports Med 44(11):2900–2910.

23.   Andarawis-Puri N, Flatow EL. 2018. Promoting effective tendon healing and remodeling.
Journal Orthopaedic Research 36(12):3115–3124.

24.   Killian ML, Cavinatto L, Galatz LM, Thomopoulos S. 2012. The role of mechanobiology in
tendon healing. Journal of Shoulder and Elbow Surgery 21(2):228–237.

25.   Chen H, Li S, Xiao H, et al. 2021. Effect of Exercise Intensity on the Healing of the Bone-
Tendon Interface: A Mouse Rotator Cuff Injury Model Study. Am J Sports Med 49(8):2064–
2073.

26.   Takahata K, Arakawa K, Enomoto S, et al. 2023. Joint instability causes catabolic enzyme
production in chondrocytes prior to synovial cells in novel non-invasive ACL ruptured mouse
model. Osteoarthritis and Cartilage 31(5):576–587.

27.   Uezono K, Ide J, Tokunaga T, et al. 2014. Effect of Postoperative Passive Motion on Rotator
Cuff Reconstruction With Acellular Dermal Matrix Grafts in a Rat Model. Am J Sports Med
42(8):1930–1938.

28.   Koike Y, Trudel G, Uhthoff HK. 2005. Formation of a new enthesis after attachment of the
supraspinatus tendon: A quantitative histologic study in rabbits. J. Orthop. Res. 23(6):1433–
1440.

29.   Wobbrock JO, Findlater L, Gergle D, Higgins JJ. 2011. The aligned rank transform for
nonparametric factorial analyses using only anova procedures. In: Proceedings of the SIGCHI
Conference on Human Factors in Computing Systems. New York, NY, USA: Association for
Computing Machinery. p 143–146 [cited 2023 Dec 18] Available from:
https://doi.org/10.1145/1978942.1978963.

30.   Jordão S, Cortes N, Gomes J, et al. 2022. Synchronization performance affects gait variability
measures during cued walking. Gait & Posture 96:351–356.

31.   Camp CL, Lebaschi A, Cong G-T, et al. 2017. Timing of Postoperative Mechanical Loading
Affects Healing Following Anterior Cruciate Ligament Reconstruction: Analysis in a Murine
Model. The Journal of Bone and Joint Surgery 99(16):1382–1391.

32.    Gomez MA, Woo SL-Y, Amiel D, et al. 1991. The effects of increased tension on healing
483           medial collateral ligaments. Am J Sports Med 19(4):347–354.

484    33.    Zhang J, Wang JH-C. 2013. The Effects of Mechanical Loading on Tendons - An In Vivo and
485           In Vitro Model Study. PLoS ONE 8(8):e71740.

486    34.    Packer JD, Bedi A, Fox AJ, et al. 2014. Effect of Immediate and Delayed High-Strain Loading
487           on Tendon-to-Bone Healing After Anterior Cruciate Ligament Reconstruction. Journal of
488           Bone and Joint Surgery 96(9):770–777.

489    35.    Chen Y, Zhang T, Wan L, et al. 2021. Early treadmill running delays rotator cuff healing via
490           Neuropeptide Y mediated inactivation of the Wnt/β-catenin signaling. Journal of Orthopaedic
491           Translation 30:103–111.

492    36.    Wada S, Lebaschi AH, Nakagawa Y, et al. 2019. Postoperative Tendon Loading With
493           Treadmill Running Delays Tendon-to-Bone Healing: Immunohistochemical Evaluation in a
494           Murine Rotator Cuff Repair Model. J. Orthop. Res. 37(7):1628–1637.

495    37.    Chamberlain CS, Crowley E, Vanderby R. 2009. The spatio-temporal dynamics of ligament
496           healing. Wound Repair and Regeneration 17(2):206–215.

497    38.    Nakamichi R, Asahara H. 2024. The role of mechanotransduction in tendon. Journal of Bone
498           and Mineral Research 39(7):814–820.

499    39.    Aicale R, Tarantino D, Maffulli N. 2018. Overuse injuries in sport: a comprehensive overview.
500           J Orthop Surg Res 13(1):309.

501    40.    Harn HI, Ogawa R, Hsu C, et al. 2019. The tension biology of wound healing. Experimental
502           Dermatology 28(4):464–471.

503    41.    Chiang RS, Borovikova AA, King K, et al. 2016. Current concepts related to hypertrophic
504           scarring in burn injuries. Wound Repair Regeneration 24(3):466–477.

505    42.    Morishita Y, Kanemura N, Kokubun T, et al. 2019. Acute molecular biological responses
506           during spontaneous anterior cruciate ligament healing in a rat model. Sport Sci Health
507           15(3):659–666.

508    43.    Nishikawa Y, Kokubun T, Kanemura N, et al. 2018. Effects of controlled abnormal joint
509           movement on the molecular biological response in intra-articular tissues during the acute

phase of anterior cruciate ligament injury in a rat model. BMC Musculoskelet Disord
19(1):175.

44.  Park S-K, Stefanyshyn DJ, Loitz-Ramage B, et al. 2009. Changing Hormone Levels during the
Menstrual Cycle Affect Knee Laxity and Stiffness in Healthy Female Subjects. Am J Sports
Med 37(3):588–598.

45.  Ford KR, Myer GD, Hewett TE. 2003. Valgus Knee Motion during Landing in High School
Female and Male Basketball Players. Medicine & Science in Sports & Exercise 35(10):1745.

46.  Ott SM, Ireland ML, Ballantyne BT, et al. 2003. Comparison of outcomes between males and
females after anterior cruciate ligament reconstruction. Knee Surg Sports Traumatol Arthrosc
11(2):75–80.

47.  Mok AC, Fancher AJ, Vopat ML, et al. 2022. Sex-Specific Outcomes After Anterior Cruciate
Ligament Reconstruction: A Systematic Review and Meta-analysis. Orthopaedic Journal of
Sports Medicine 10(2):232596712210768.

48.  Nigro P, Middelbeek RJW, Alves CRR, et al. 2021. Exercise Training Promotes Sex-Specific
Adaptations in Mouse Inguinal White Adipose Tissue. Diabetes 70(6):1250–1264.

49.  Bains RS, Forrest H, Sillito RR, et al. 2023. Longitudinal home-cage automated assessment of
climbing behavior shows sexual dimorphism and aging-related decrease in C57BL/6J healthy
mice and allows early detection of motor impairment in the N171-82Q mouse model of
Huntington's disease. Front. Behav. Neurosci. 17:1148172.

50.  Lidster K, Owen K, Browne WJ, Prescott MJ. 2019. Cage aggression in group-housed
laboratory male mice: an international data crowdsourcing project. Sci Rep 9(1):15211.

51.  Adams D. 1986. Ventromedial tegmental lesions abolish offense without disturbing predation
or defense. Physiology & Behavior 38(2):165–168.

52.  Miczek KA, Maxson SC, Fish EW, Faccidomo S. 2001. Aggressive behavioral phenotypes in
mice. Behavioural Brain Research 125(1–2):167–181.

**Fig. 1** Study design: Mice were ruptured ACL non-invasively and created the CATT model

immediately post-injury and sacrificed post-injury immediately, at 4- and 8-weeks post-injury (A).

Mice were randomly classified into the Tight-CATT (B), Loose-CATT (C), and ACL-R (D) groups.

The images of the ACL immediately after injury (E), the healed ACL (F), and the non-healed ACL

(G) (Scale bar = 300 μm). This schematic was made using https://www.biorender.com/

**Fig. 2** The load to failure (A) and stiffness (B) of the nylon suture in failure testing, and the number

of cycles at the point of suture rupture in cyclic testing (C) were recorded (*$p < 0.05$, **$p < 0.01$).

**Fig. 3** Anterior drawer testing was performed with the knee joint fixed at 90° via the self-made

anterior drawer device (A). Representative X-ray images of each group (B). We evaluated the

braking force of the ATT for each group immediately after injury (C), at 4 (D) and 8 weeks (E), and

anterior tibial instability in the ACL-healed knee at 4 (F) and 8 weeks (G).

**Fig. 4** HE stained images of the Tight-CATT (A, B, C, and D), Loose-CATT (E, F, G, and H), ACL-

R groups (I, J, K, and L) at 4 and 8 weeks (Scale bar = 300 μm). The black arrows point to the

healing region of the healed ACL and the white arrows point to the non-healed ACL.

**Fig. 5** Picrosirius red stained (A) and polarized image (B) of the ACL. Picrosirius red images were

converted to grayscale images (C) and the specified ACL area was specified as ROI (D). Collagen

maturation of the ACL was evaluated at 4 (E) and 8 weeks (F) post-injury (Scale bar = 300 μm).

**Fig. 6** Immunofluorescence images targeted COL1A1 of the intact ACL (A, B), healed ACL of the

Tight-CATT (C, D), and Loose-CATT (E, F) groups at 8 weeks (Scale bar = 300µm) and showing

mid-substance of the ACL: COL1A1 (green) and DAPI (blue) (Scale bar = 100µm). Arrows pointed

to areas of high COL1A1 expression.

**Fig. 7** To perform biomechanical testing, the soft tissues of the knee, except the ACL, were

dissected, and the femur and tibia were fixed in tubes (A). The load to failure (A) of the healed ACL

at 4 (B) and 8 (C) weeks after injury was recorded.