An 'Epidemic Diversity' conceptual model explains how host genetic diversity affects variation in parasite success

Paplauskas S., Duthie A. B. and Tinsley M. C.

University of Stirling, Department of Biological and Environmental Sciences, Stirling, FK9 4LA, Scotland, UK

Abstract

According to conventional wisdom, disease transmission rate is usually higher in more genetically homogenous host populations. Previous studies have principally considered how host population genetic diversity impacts the mean of parasite infection performance. However, when considering risks from epidemics and the emergence of novel infectious diseases, variability in parasite success may be just as important as the mean. Here we propose an Epidemic Diversity Model for how host-parasite population genetic diversity influences both the mean and variability in parasite success. We evaluate this conceptual model by re-analysing effect size data from two meta-analyses, including 211 comparisons of high versus low genetic diversity host populations from 48 studies. Our analysis challenges previous understanding by demonstrating that high host population genetic diversity only reduces mean parasite success for specialist parasites with narrow host range, but not for generalist multi-host parasites. We also find that the combination of host range and parasite population genetic diversity determine the effect of host population genetic diversity on the variability in parasite success. These results have important implications for the management of host population genetic diversity for natural populations.

Introduction

Epidemics can be defined as extremely rapid increases in disease prevalence over the endemic (baseline) level of disease (Dicker, 2006). They pose a major threat to food security, biodiversity and human health (Altizer et al., 2003). This is because epidemics can devastate crop harvests (Zhu et al., 2000), threaten local host populations with disease-induced extinction (De Castro & Bolker, 2005) and increase the risk of zoonotic spillover (Jones et al., 2008). Previous research suggests that less genetically diverse host populations tend to experience larger epidemics (King & Lively, 2012). Although this phenomenon is well-established in the agricultural literature and is often referred to as a 'monoculture effect' (Elton, 1958; Garrett & Mundt, 1999; Zhu et al., 2000), it is much less well-understood how the level of genetic diversity in animal and bacterial species (i.e. not-plant hosts) determines epidemic size in nature.

Previous studies by Ekroth et al. (2019) and Gibson and Nguyen (2021) found that higher levels of host genetic diversity reduced the mean for various metrics of parasite success (including disease prevalence, parasite load and virulence) across a range of both animal and bacterial host-parasite systems. This evidence is centred around means, and although disease prevalence is only one measure of parasite success, it appears to support the conventional wisdom that less genetically diverse host populations tend to experience larger epidemics, at least *on average* (King & Lively, 2012). However, despite this important discovery (Ekroth et al., 2019; Gibson & Nguyen, 2021), many host populations in nature experience epidemics that *vary* dramatically in both time and space (Altizer et al., 2004; Aznar et al., 2015; Cáceres et al., 2006; Carlsson-Granér & Thrall, 2002; Ericson et al., 1999; Montano et al., 2016; Thrall et al., 2012; Vergara et al., 2013). Therefore, understanding not only the mean, but also the *variability* of parasite success is important.

Previous research suggests that the understanding the variability of parasite success may be important from the perspective that high levels of variation are akin to particularly large, or severe epidemics. For example, a highly cited paper published by Tarpy (2003) in *Nature* showed that more genetically diverse honeybee colonies had less variable disease prevalence. As such, they suggested that this could potentially protect more diverse honeybee colonies against the occurrence of more severe epidemics. Alternatively, increased understanding of the variability of parasite success could increase our general understanding of the repeatability of disease experiments, which are characterised by large amounts of variation (Paplauskas et al., 2021).

Any relationship between host population genetic diversity and either the mean or variability
in parasite success will likely depend on the corresponding level of parasite population genetic
diversity and infection genetics. For example, assuming that there is some level of specificity
for infection (Schmid-Hempel & Ebert, 2003), where host and parasite genotypes vary in their
susceptibility or infectivity to one another, individual host genotypes are more likely to be
infected by a diverse parasite population. Therefore, the parasite transmission rate and the
overall disease prevalence should be higher for more diverse parasite populations (Boomsma,
1996; Van Baalen & Beekman, 2006). However, the amount of host-parasite specificity for
infection varies between different models of infection genetics (Agrawal & Lively, 2002). In
these models, infection specificity ranges from matching allele paradigms where one parasite
genotype can only infect a single host genotype, to gene-for-gene interactions where
infectious parasite genotypes are able to infect many or all host genotypes (Agrawal & Lively,
2002). In reality, these models represent opposite ends of a continuum and host-parasites
systems vary in where they fall along this continuum (Agrawal & Lively, 2002). Nevertheless,
the strength of specificity clearly has the potential to modulate how host-parasite population
genetic diversity influences both mean and variation in parasite infection success.

Here, we propose an 'Epidemic Diversity' conceptual model to describe how host population
genetic diversity and parasite population genetic diversity may influence both the mean and
variability of parasite success (Fig. 1). We envisage a metapopulation structure composed of
multiple local populations. Local populations vary in their level of genetic diversity (measured
at the inter-, versus intra-, group level; Fig. S1). Low genetic diversity results in local
populations that harbour few host or parasite genotypes, but the identity of these genotypes
may differ between local populations. High genetic diversity increases the number of
genotypes present and local populations are less genetically differentiated. Each host
individual encounters many potentially infectious parasites. Initially we consider a case of high
infection specificity, where close matching is required between host and parasite genotypes
for successful infection. We predict the following patterns in mean and variance of parasite
success across the local populations:

A) When host and parasite population genetic diversity are both low (Fig. 1A), parasite
success will be highly variable because host populations will be composed either of
mostly susceptible or mostly resistant host genotypes. Mean parasite success will be
intermediate, and determined by the relative frequency of resistant and susceptible
populations.

B) When genetic diversity is high in the host population but low in the parasite population
(Fig. 1B) both mean and variability in parasite success will be low because the
parasite's encounter rate with matching host genotypes will be consistently low.
C) When genetic diversity is low in the host population but high in the parasite population
(Fig. 1C) mean parasite success will be high and variability in parasite success low,
because many hosts will encounter matching parasite genotypes and efficient chains
of transmission will establish.
D) When genetic diversity is high in both host and parasite populations (Fig. 1D) the mean
and variability in parasite success will be intermediate because although the probability
of hosts encountering infectious parasite genotypes is high, onward transmission is
impaired by the diversity of host genotypes present.
108

Our model predicts that under this scenario of high genetic specificity for successful infection,
an increase in host population genetic diversity will result in a decline in *mean* parasite
infection success regardless of the parasite population's genetic diversity (Fig. 1E and F).
However, increased host population genetic variation will cause reduced *variability* in parasite
infection success when parasite genetic diversity is low (Fig. 1G) but increased *variability*
when parasite diversity is high (Fig. 1F). All these effects will be sensitive to the genetic
specificity that underpins the infection process. In cases where parasite virulence genotypes
can infect many (or all) host genotypes (sensu gene-for-gene models) these predictions
should break down and the effects of host population genetic diversity on parasite infection
success should disappear.
119

This Epidemic Diversity conceptual model builds on previous work by Bensch et al. (2021),
who considered three out of four of the combinations in Figure 1. We extend this previous
work by considering how high parasite genetic diversity can lead to more consistent parasite
success in genetically depauperate host populations (Fig. 1C). Furthermore, in our work we
consider how these relationships between host-parasite population genetic diversity and
parasite success depend on the strength of genetic specificity for infection success.
126

It is impossible to make an explicit test of the effect of genetic specificity for infection success
in our analysis because genetic specificity is not characterised in most host-parasite systems
(but see Pepijn et al., 2013). However, we investigate the impact of a proxy for genetic infection
susceptibility: parasite host-range. This is because it is only specialist parasites (with a narrow
host range) that are likely to have undergone tight coevolution with their host species of the
type that promotes the maintenance of matching-allele-type genetic variants under negative

frequency dependent selection. Whereas for generalist parasites that infect many host species

(and have not tightly coevolved with any single host), virulence mechanisms are more likely

to be general and confer ability to infect many host genotypes.

Figure 1. A hypothetical 'Epidemic Diversity' model for the combined relationship between host and parasite population genetic diversity and either the mean (μ) or the variability (σ) in parasite success. There are four hypothetical populations for each combination of host and parasite population genetic diversity (dashed circles). The level of population genetic diversity is indicated by the number of unique host and parasite genotypes (large and small shapes respectively) and is the same in each replicate population. The colour of hosts indicates their infection status, such that susceptible hosts are white and infected hosts are black, whereas the parasite is always the same colour (also black). Parasite transmission can only occur between matching host and parasite genotypes (shapes) and is indicated by the red arrows. The resulting frequency distributions of parasite infection success for each set of replicate populations is shown at the bottom of each plot. Notably, this hypothetical model only applies for host--parasite systems that have a high level of genetic specificity for infection (i.e. matching-allele versus gene-for-gene infection genetics (Agrawal & Lively, 2002)). The bottom panel of the facetted grid show the expected result from four comparisons of high versus low genetic diversity host populations (two x low parasite diversity, two x high parasite diversity). The design of these plots mirrors the orchard plots for the meta-analytical models from Figure 5.

To evaluate our proposed Epidemic Diversity conceptual model, we combine the data from
two previous meta-analyses that investigated the effect of host population genetic diversity on
mean metrics of parasite infection success (Ekroth et al., 2019; Gibson & Nguyen, 2021). By
pooling the data from these original meta-analyses, we achieve a bigger sample size so that
we can investigate patterns in parasite infection success variance, as well as the impact of
additional moderator variables. Since we focussed on the re-analysis of previous studies by
Ekroth et al. (2019) and Gibson and Nguyen (2021), which mainly focussed on the effect of
differences in host genetic diversity in animal and bacterial populations, rather than plant
populations, we also focussed our study on animal and bacterial populations. If a researcher
was interested in investigating the effect of differences in host genetic diversity in plant
populations (*sensu* Reiss & Drinkwater, 2018), a much more comprehensive review of the
literature would be required.

Methods

Summary

We combined the data from two previous meta-analyses (Ekroth et al., 2019; Gibson &
Nguyen, 2021) that used the standardised mean difference (SMD) to calculate the significance
of the relationship between host genetic diversity and mean metrics of parasite infection
success. These data were mainly from studies where the metrics of parasite infection success
were measured from replicate host populations that were already classified qualitatively as
having either a 'high' or 'low' level of genetic diversity (27 out a total of 48 independent studies),
including studies of replicate host populations with a high versus low level of inbreeding (e.g.
Baer & Schmid-Hempel, 1999), different combinations of host genotypes (e.g. Altermatt &
Ebert, 2008) or comparisons between wild host populations exposed to different selection
regimes (e.g. a population bottleneck, random genetic drift, e.g. Hale & Briskie, 2007).
However, there were some studies that assessed a continuous measure of host population
genetic diversity (for the absence of any significant correlation between mean metrics of
parasite infection success and host population genetic diversity, involving all of the studies
from this subset of data, see Gibson & Nguyen, 2021), and therefore these data were binned
into 'high' and 'low' categories. Following Ekroth et al., (2019) and Gibson & Nguyen (2021)
we used the SMD to assess factors influencing mean parasite infection success; we then
extended these previous analyses by using the log coefficient of variation ratio (lnCVR) to
assess factors influencing variation in parasite success. In the original studies, this variation
was quantified in various ways: between experimental replicates in a laboratory environment;
between multiple natural host populations with similar genetic diversity; or sometimes between
repeated measures of single populations along a time series.

Data collection

Data collection for each comparison of a group of high versus low genetic diversity host
populations involved five steps (Fig. 2):

1) *Combination*: First, we combined the list of studies from (Gibson & Nguyen, 2021) and
(Ekroth et al., 2019), removed any duplicate studies and added the data used to calculate the
effect size on mean parasite success (SMD) and its sampling variance in the original studies
(mean, standard deviation, sample size), the metric of parasite success and the unique study,
experiment and replicate identifiers used to account for the non-independence of separate
effect sizes. We did not use the parasite success data from Ekroth et al., (2019) because the
original data extracted from each study was not included in their online supplementary material

and this meant we would have been unable to check the data accuracy during the third step
of data collection (validation). However, we extracted any missing data from these studies and
recalculated the mean, standard deviation and sample size ourselves during the fourth and
fifth steps of data collection (extraction and recalculation).

The data used to calculate Fisher's z (an effect size for the difference between two correlation
coefficients) for the observational field studies from Gibson & Nguyen (2021), was not
available in the correct format to calculate either SMD or lnCVR. Therefore, we did not include
this information (from multiple populations with a continuous measure of genetic diversity) at
this stage and instead extracted the data from the original publications ourselves and
recalculated it during steps 4 and 5. Also, we excluded any studies on plants (wild or
agricultural) because a more detailed analysis of the plant literature would require a separate
review.

2) *Screening*: Second, we amended the inclusion criteria used in the original meta-analyses
(Ekroth et al., 2019; Gibson & Nguyen, 2021) (Table S1) and removed any studies,
experiments or comparisons which did not meet these criteria:

(i) 'Parasite success', which we define as the ability of a parasite to spread among hosts
(transmission rate, infection rate, prevalence), replicate on / within hosts (macro /
microparasite load, disease severity) or kill hosts (virulence i.e. host survival / mortality rate)
was measured among replicate populations across time or space.

(ii) Parasite success data was collected from two or more host populations with a difference
in genetic diversity assessed by metrics such as: individual inbreeding status (inbred versus
outbred), genotypic diversity (high versus low) or heterozygosity (high vs low).

(iii) Genetic diversity was quantified at the level of the host population, rather than for
community-level diversity.

(iv) The study focused on an animal or bacterial host species.

(v) The study did not re-analyse the data from a previously published study.

(vi) The parasite success data was not replicated simply by using an alternate way of
measuring host population diversity.

(vii) Figures required to extract parasite success data were clearly legible.

3) *Validation*: Third, we checked the accuracy of the data from the excel spreadsheets used
to calculate the summary of the parasite data for each group of host populations with either
high or low genetic diversity from the online data supplied by one of the previous meta-
analyses (Gibson & Nguyen, 2021) and corrected these in the fourth step of the data collection
before including them in our analysis. Although there 69 errors in total in the data from their
study, it is worth noting that we ran-ran the meta-analytical procedures from their study with
the amended data and found that the major conclusions of their study were unaffected by
these errors. In addition, we were thoroughly impressed by the amount of detail included in
their supplementary files, and we were only able to check their data's accuracy due to their
diligence (cf. Ekroth et al. 2019). The different types of error made by the previous meta-
analysis included (i) 27 comparisons that did not match the published raw data (available in
the main text or online or in the supplementary material of each publication), (ii) 32
comparisons where effect sizes were calculated wrong and (iii) 10 comparisons which had not
been transferred into the final metadata file correctly. There was one study which we could
not check, because the original data was sent by personal communication from (King et al.,
2011) to (Gibson & Nguyen, 2021), nevertheless we included it in our analysis.

4) *Extraction*: Fourth, for those studies or comparisons we had excluded (due to missing data
or data errors) we extracted the data from the main text or supplementary files by going back
to the original publications (we used PlotDigitizer (https://plotdigitizer.com/) to extract the
information for any figures). In addition to the comparisons removed in the third step of data
collection, this also included (i) 26 studies that, despite meeting our inclusion criteria, were
removed because they were either missing the replicate-level raw data (Ekroth et al., 2019)
or they were observational field studies based on multiple populations with a continuous
measure of genetic diversity (Gibson & Nguyen, 2021) and (ii) additional data for 3
comparisons (Agha et al., 2018; Baer & Schmid-Hempel, 2001; Giese & Hedrick, 200) that
were not made in the original analysis by (Gibson & Nguyen, 202).

We also collected information on 10 different moderators (see Table 1) by standardizing or
recoding existing moderator variables used by (Gibson & Nguyen, 202), including host range,
parasite diversity, metric of parasite success, host species, parasite type, source of host
genetic diversity, scale of host diversity, mode of host reproduction, whether the parasite
induces host mortality and whether the study was performed in a laboratory environment.

Parasite diversity was not quantified as a continuous variable in the original studies, nor was
it examined as part of the original experiments in most cases. Therefore, we binned parasite
diversity into 'high' or 'low' groups depending on the following reasoning: 'high parasite
diversity' if the isolate was collected from a natural population for a lab study, if the data was
from an observational or experimental field study, or if more than one genotype had been
identified (but this only applied to a small number of studies); 'low parasite diversity' if it was a
laboratory strain, or only one genotype had been identified (but again, this only applied to a
small number of studies). Where the information on these moderator variables was not already
available from the supplementary material of Gibson & Nguyen (202) and was not available in
the published article, we performed an online search to determine characteristics.

5) *Recalculation*: Fifth, we calculated the mean, standard deviation and sample size for each
comparison of high versus low genetic diversity groups of host populations for the data we
extracted in the fourth step of data collection (Gibson & Nguyen, 202). For certain studies, we
calculated a pooled measure of the mean metric of parasite infection success for each group
of high and low genetic diversity host populations, along with a pooled standard deviation and
a pooled sample size. This included (i) studies with shared comparisons (e.g. multiple high
diversity groups paired to the same low diversity group) (Agha et al., 2018; Schmidt et al.,
2011) or (ii) those based on multiple populations with one or more continuous measures of
genetic diversity (Dagan et al., 2013; Dionne et al., 2009; Ellison et al., 2011; S. G. Field et
al., 2007; Giese & Hedrick, 2003; King et al., 2011; Kyle et al., 2014; Loiseau et al., 2011;
Meagher, 1999; Neumann & Moritz, 2000; Parsche & Lattorff, 2018; Pierce et al., 2014;
Puurtinen et al., 2004; Queirós et al., 2016; Rahn et al., 2016; Savage et al., 2015; Trouvé et
al., 2003; Velavan et al., 2009; Whitehorn et al., 2011, 2014; Whiteman et al., 2006). In the
latter case, the most appropriate measure of population-level genetic diversity was used (e.g.
a measure of population-level genetic diversity based on Hardy-Weiberg equilibrium) and two
separate groups of host and low diversity host populations were made with the same number
of host populations in each group. In addition, host survival was converted into host mortality
in some studies to reflect our definition of parasite infection success (see step two of data
collection). Overall, this fifth step of data collection involved calculating parasite success data
for 130 comparisons.

After finishing all five steps of data collection, there was enough parasite success data to
calculate both the SMD and lnCVR for 211 non-independent comparisons of high versus low
genetic diversity groups of host populations.

Figure 2. How data was collected for each comparison of a parasite infection success metric between high versus low genetic diversity groups of host populations, which was later used in calculating effect sizes. n = number of studies / experiments / comparisons (the multi-level structure of the data make it appear that some sums are incorrect). Adapted from the preferred reporting items for systematic reviews and meta-analyses (PRIMSA) statement (Page et al., 202).

Calculation of effect sizes (SMD and lnCVR)

We calculated two effect size measures: the standardized mean difference (SMD) and the log
coefficient of variation ratio (lnCVR). These quantified the effect of host population genetic
diversity on either the mean or variability in parasite success across different studies
(Borenstein et al., 200). This is because we were interested in comparing both the mean and
variability in various measures of parasite infection success between groups of 'high' versus
'low' genetic diversity host populations.

We selected these specific ways of measuring between-group differences in either the mean
or variability among metrics of parasite success because they took in to account certain
factors: (i) SMD measures the mean difference between two groups (high versus low genetic
diversity) in terms of standard deviations (Borenstein et al., 2009; A. P. Field & Gillett, 201),
so it can be used to compare metrics measured on very different scales (prevalence, load and
virulence) (Higgins & Green, 202); it also corrects for small sample sizes, which is a common
feature of ecological studies (Jennions, 200). Similarly, lnCVR measures the ratio of variability
between two groups adjusted for the size of the group means (Nakagawa et al., 201) and
therefore can be used to compare multiple metrics. Also, it accounts for the possibility that the
magnitude of the variability may scale with the mean, which is relevant for many types of count
data that may be described by a Poisson distribution (e.g. parasite load).

Before calculating our effect sizes, we added a small value (0.001) to the mean and standard
deviation in parasite success for each pair of high and low genetic diversity groups of host
populations to ensure log values were calculated correctly. For consistency, we calculated
SMD and its sampling variance from the formula derived from the supplementary material of
(Gibson & Nguyen, 202), whereas we calculated variability effect sizes (lnCVR) and their
sampling variances using the code from (Nakagawa et al., 201). To account for comparisons
based on a shared reference group of host populations with low genetic diversity (i.e. shared
controls), we calculated the variance-covariance matrix for each effect size, using the
make_VCV_matrix function from the metaAidR package v0.0.0.9000 (Lagisz et al., 202).

All effect size calculations and subsequent calculations were performed in R v4.3.2 (R Core
Team, 202)

Publication bias

Before analysing the data fully, we calculated the overall effect sizes for SMD and lnCVR and
tested for any potential publication bias using funnel plots and Egger's regression (Sutton,
200).

Meta-analytic models were fitted to the data using the rma.mv function from the metafor
package v4.4.0 (Viechtbauer, 201). We included fixed effects for each type of effect size, the
variance-covariance matrix of sampling errors, standard random effects for study and host
genus, and correlated random effects for comparisons taken from the same experiment.
Standard random effects for study and host genus were used to account for the possibility of
non-independence between experiments originating from the same study and potential
correlations between effects from closely related host species. Similarly, correlated random
effects were used to account for potential non-independence of comparisons taken from the
same experiment (multiple timepoints for a single comparison of a high and low genetic
diversity group of host populations, or effect sizes from the same group of populations based
on different measures of parasite success).

Funnel plots were used to identify whether published effect sizes were evenly distributed
around model means by examining how outcomes varied as a function of their precision
(standard error). Visual inspection of funnel plots for the effect of host population diversity on
mean parasite success (Fig. 3A) and its effect on the variability in parasite success (Fig. 3B),
showed no evidence for publication bias. More stringent evaluation showed that there was no
correlation between the size of the effects themselves and their standard error (Egger's test
for both SMD and lnCVR: R = 0.06, 95% CI [-0.24, 0.37], p = 0.67 and R = -0.03, 95% CI [-
0.38, 0.32], p = 0.86, respectively).

Figure 3. Testing for publication bias: the distribution of published effect sizes for our meta-analysis as a function of their precision (standard error). The x-axis in both plots shows effects of an increase in host population genetic diversity (high vs low) on A) the mean difference in parasite success (SMD) and B) the ratio of variation in parasite success (lnCVR). Model means and their 95% confidence intervals are shown by the dashed black lines.

355   Meta-analysis of overall data

356   To test whether there was a significant difference in the mean parasite success (SMD) or
357   variability in parasite success (lnCVR) between host populations with high versus low genetic
358   diversity, we fitted mixed effects meta-analytic models. All of the models used in this paper
359   were based on the same structure as those used for testing the presence of publication bias.
360

Context dependence

Moderators with an interaction - To test if the overall effect of host population genetic diversity
on the mean and variability in parasite success depended on an interaction between host
range and parasite genetic diversity, we introduced an interaction term for these two
moderators in our original meta-analytic models. Therefore, we could compare:
1) High versus low population genetic diversity of host specialist parasites.
2) High versus low population genetic diversity of host generalist parasites.
368

We compared the significance level of each individual predictor within the model, as well as
the contrasts between them using the *glht* function from the *multcomp* package v1.4.25
(Hothorn et al., 200).

Moderators without an interaction – To test our additional hypotheses (Table 1) for the eight
remaining moderator variables, we modelled each moderator separately with its own individual
mixed effects model. Before running the models, we removed redundant moderator categories
with a limited sample size, such as transmission or infection rate (versus prevalence) and
disease severity (versus load) for the metric of parasite success, and prokaryotic (versus
vertebrate or invertebrate) for host species.

We compared the significance level of each individual predictor within the model, as well as
the contrasts between them using ANOVA with a correction for multiple comparisons (Holm's
method).

Table 1. Hypotheses for the influence of additional moderator variables on the nature of the
effect of host population genetic diversity on mean and variability in parasite success.

Moderator	Hypothesis
Metric of parasite success	Our study of 'parasite success' combined data of several types (eg prevalence, virulence, infection load). Using this moderator, we tested if the effects of host population genetic diversity differed between these different metrics.
Host type	The effect of host population genetic diversity may be influenced by the specificity of genetic interactions between host and parasite. These genetic interactions are thought to be more specific in invertebrates than in vertebrates (Dybdahl et al., 201), therefore we tested for inconsistency of the host population genetic diversity effect in these two groups.
Parasite type	Microparasites and macroparasites tend to have contrasting infection biology: varying in the duration of infections and the immune responses they trigger (Sorci, 2014). These differences might drive variation in the impact of host population genetic diversity. Therefore, we tested for inconsistency of the host population genetic diversity effect in these two groups.
Source of host genetic diversity	Studies typically investigate the impact of host genetic diversity by (i) controlling mating strategies to compare hosts with varying degrees of 'relatedness' (e.g. polyandry versus polygyny) or with different levels of inbreeding, (ii) using a suite of wildtype genotypes for controlled experiments with either low genetic diversity or high

	genetic diversity, or (iii) sampling organisms from the wild from populations that have been characterised as having different levels of genetic diversity. We used this moderator to test if these different sources of genetic diversity affected the influence of host population genetic diversity.
Scale of host diversity	Host populations were predetermined as having either high or low diversity (discrete) or we separated them into such categories as part of our data collection (Fig. 2, step 5) because the authors used multiple populations with a continuous measure of diversity. We used this moderator to test if this feature of how studies were designed had an effect on the influence of host population genetic diversity.
Mode of host reproduction	Host species reproduced sexually, asexually or using a combination of the two (i.e. facultatively sexual, such as *Daphnia*). We used this moderator to test if these different modes of host reproduction affected the influence of host population genetic diversity.
Host mortality?	The range of parasites studied can be further categorised by whether or not infection typically kills the host (which may be proxy for virulence). We used this moderator to test if differences in the virulent effects of parasitism affected the influence of host population genetic diversity.
Laboratory?	We used this moderator to test if differences in study setting (laboratory versus field) affected the influence of host population genetic diversity.

Sensitivity analysis of overall effects

To test the robustness of our results for the combined (overall) dataset, we performed a series

of 'leave-one-out' sensitivity analyses. This involved the iterative exclusion of either one

independent comparison (i.e. groups of high diversity host populations that shared a

corresponding group of low diversity host populations were considered grouped together into

a single comparison) or study at a time.

Results

Host population genetic diversity has an overall negative effect on mean parasite success

Our dataset contained 211 effect sizes from 48 different studies, including a range of (mainly
animal) host and parasite species (Table S2). These effect sizes represented individual
estimates of the effect that change in host population genetic diversity has on parasite
success; we assessed this effect on both mean parasite success (SMD) and the variability in
parasite success (lnCVR).

Averaging over the whole data set, there was a significant effect of host population genetic
diversity on mean parasite success (SMD = -0.29, 95% CI = [-0.57, -0.02], p = 0.04; Fig. 4A):
higher levels of host population genetic diversity were associated with lower mean parasite
success. In contrast, across the whole data set, the variability of parasite success was not
significantly affected by host population genetic diversity (lnCVR = 0.02, 95% CI = [-0.30,
0.35], p = 0.89; Fig. 4B).

In these analyses the residual variation (heterogeneity) in the data for both the difference in
the mean and the variability of parasite success was high (I2 = 84.0% & 82.0% respectively).
Most of this variation was explained by the effect of study (84.0% & 80.7%) and only a small
amount was explained by host genus (0.0% & and 3.3%).

Figure 4. The overall effect of host population genetic diversity on the mean and variability in parasite success. The x-axis in each plot shows the effect that an increase in host population genetic diversity had on either A) the mean parasite success (SMD) or B) the variability in parasite success (lnCVR). The dashed line indicates an effect size of zero where host population genetic diversity has no influence. Model means are shown with 95% confidence intervals (black rectangles) and prediction intervals (thin black lines). Circles show individual effect sizes and are scaled according to the inverse of their standard error. n = sample size of the data (the number of effect sizes : the number of studies). The asterisk shows that the model mean is significantly different from zero (p < 0.05). Forest plot alternatives are shown in the online supplementary material (Fig. S2).

Impacts of host population genetic diversity on parasite success differ between specialist and
generalist parasites

Next, we investigated how the effect of host population genetic diversity on parasite infection
success was influenced by two fundamental characteristics of the parasite: the host-specificity
of the parasite and the likely genetic diversity of the parasite population studied.

The negative effect of host population genetic diversity on mean parasite infection success
that we observed across the whole dataset (see above), was actually only significantly
negative for specialist (single-host) parasites. This was true whether the observations for
specialist parasites were for parasite populations with low or high genetic diversity (SMD = -
0.54, 95% CI = [-1.02, -0.06], p = 0.03 & SMD = -0.76, 95% CI = [-1.16, -0.37], p < 0.001
respectively; Fig. 5A). The point estimates for the effect of host population genetic diversity on
mean parasite infection success for generalist (multi-host) parasites were also negative;
however, these estimates were not significant, and also not influenced by parasite isolate
genetic diversity (SMD = -0.42, 95% CI = [-0.91, 0.07], p = 0.09 & SMD = -0.23, 95% CI = [-
0.55, 0.08], p = 0.15 respectively; Fig. 5A). This difference between specialist and generalist
parasites in the effect of host population genetic diversity on mean parasite infection success
was highly significant (glht = 0.47, SE = 0.16, p < 0.01; Fig. 5A).

Similar to the results for mean parasite success, the manner in which *variability* in parasite
infection success was influenced by host population genetic diversity was also influenced by
the characteristics of the parasite population. Although there was no significant relationship
between host population genetic diversity and the variability in parasite success for the global
dataset (see above), the observations for specialist parasites that infect only a single host
species matched the predictions of our Epidemic Diversity conceptual model (Fig. 5B). When
the genetic diversity of specialist parasites was low, an increase in host population genetic
diversity tended to reduce variability in infection success (lnCVR = -0.54; 95% CI = [-1.14,
0.06], p = 0.08; Fig. 5B). Whereas, for these specialist parasites, when parasite genetic
diversity was high, an increase in host population genetic diversity resulted in increased
variation in infection outcome (lnCVR = 0.61, 95% CI = [-0.23, 1.47], p = 0.16; Fig. 5B).
Although these associations were not significant in isolation, there was a significant difference
between them (glht = -1.15, SE = 0.53 and p = 0.03). Also, mirroring our results for the mean
of infection success, variability in parasite infection success was unaffected by host population
genetic diversity in multi-host generalist parasites (Fig. 5B).

Figure 5. The effect of host population genetic diversity on both the mean and variability in metrics of parasite infection success depends on a combination of host range and parasite population genetic diversity. The x-axis in each plot shows the effect of an increase in host population genetic diversity on either A) mean parasite success (SMD) or B) variability in parasite success (lnCVR). The dashed line indicates an effect size of zero where there is no influence of host population genetic diversity on parasite success. Model means are shown with 95% confidence intervals (black rectangles) and prediction intervals (thin black lines). Individual effect sizes (circles) are scaled according to the inverse of their standard error. n = sample size of the data (the number of effect sizes : the number of studies). The significance level of individual model means, as well as any pairwise contrasts, is indicated by one ($p < 0.05$) or three ($p < 0.001$) asterisks.

The effect of host population genetic diversity on parasite success is influenced by a number
of other contextual factors

In addition to measuring the combined effect of parasite population genetic diversity and
parasite host range on the relationship between host population genetic diversity and both the
mean and variability in metrics of parasite infection success (see above), we also separately
investigated how eight other contextual factors (Table 1) affected this relationship (Fig. 6).

Most of the moderator levels were not significantly different from zero and comparisons

between different levels within the same moderator showed that they were often similar in

terms of both the magnitude and direction of their effects. However, the parasite type (Fig.

6C), source of host diversity (Fig. 6D), host mode of reproduction (Fig. 6F), whether the

parasite kills the host or not (Fig. 6G) and whether the study was performed in the laboratory

(Fig. 6H) all seemed to have a significant effect on the relationship between host population

genetic diversity and mean parasite success. First, there was a significant reduction in mean

microparasite success in host populations with higher genetic diversity (SMD = -0.47, 95% CI

= [-0.77, -0.18], $p < 0.01$, Fig. 6C) that was also associated with a significant difference

between the effect of host population genetic diversity on microparasite versus macroparasite

success (QM = 13.2, df = 1, $p < 0.001$, Fig. 6C). Second, there was a significant reduction in

mean parasite success for host populations with lower relatedness (outbred versus inbred)

(SMD = -0.51, 95% CI = [-0.98, -0.03], $p = 0.04$, Fig. 6D), similar to those composed of a

higher number of chosen genotypes (close to the 95% significance threshold; SMD = -0.51,

95% CI = [-1.02, 0.01], $p = 0.05$, Fig. 6D), but in contrast to host populations that were naturally

more genetically diverse (SMD = 0.03, 95% CI = [-0.40, 0.45], $p = 0.90$, Fig. 6D). Third, there

was a significant reduction in mean parasite success in sexually reproducing host populations

with higher genetic diversity (SMD = -0.40, 95% CI = [-073, -0.07], $p = 0.02$, Fig. 6F) which

was absent for both asexually and facultatively sexually reproducing host populations (SMD =

-0.36, 95% CI = [-1.62, 0.91], $p = 0.58$ and SMD = 0.03, 95% CI = [-0.51, 0.57], $p = 0.93$

respectively, Fig. 6F). Fourth, there was a significant reduction in mean parasite success for

host populations with higher genetic diversity where the parasite usually killed the host (SMD

= -0.34, 95% CI = [-0.68, -0.01], $p < 0.05$, Fig. 6G) which was absent for parasites that did not

usually kill the host (SMD = -0.13, 95% CI = -0.56, 0.29], $p = 0.54$, Fig. 6G). Fifth, there was a

significant reduction in mean parasite success for host populations with higher genetic

diversity studied outside of a laboratory environment (SMD = -0.33, 95% CI = [-0.65, -0.01], p

= 0.04, Fig. 6H) which was absent from studies conducted in a laboratory (SMD = -0.24, 95%

CI = [-0.60, 0.13], $p = 0.20$, Fig. 6H).

There was only one moderator, the host mode of reproduction (Fig. 6N), which had a

significant effect on the relationship between host population genetic diversity and the

variability in parasite success. Specifically, there was a significant reduction in the variability

of parasite success for asexually reproducing host populations with higher genetic diversity

(lnCVR = -0.74, 95% CI = [-1.26, -0.21], $p < 0.01$), Fig. 6N) that was also associated with

significant differences to the lack of any effect of host population genetic diversity on the

variability in parasite success for both sexually and facultative-sexually reproducing hosts (QM

= 6.40, df = 1, p = 0.01 & QM = 5.53, df = 1, p = 0.02 respectively).

Figure 6. The context-dependence of the effect of host population genetic diversity on the mean and variability in parasite success. The y-axis in each plot shows the effect of an increase in host population genetic diversity on either the difference in mean parasite success (SMD) (panels A-H), or the difference in the variability in parasite success (lnCVR) (panels I-P). Model means are shown with 95% confidence intervals and are scaled according to the number of experiments. The dashed line indicates an effect size of zero. The significance level of individual model means, as well as any pairwise contrasts, is indicated by one (p < 0.05), two (p < 0.01) or three (p < 0.001) asterisks. The following abbreviations are used; Prev. (Prevalence), Vir. (Virulence), Vert. (Vertebrate), Invert. (Invertebrate), Macro. (Macroparasite), Micro. (Microparasite), Source gen. var. (Source of host genetic diversity), Art. (Artificial), Nat. (Natural), Scale host div. (Scale of

host diversity), Cont. (Continuous), Disc. (Discrete), Host mode repro. (Mode of host reproduction), Sex. (Sexual), Asex. (Asexual), Mixd (Mixed).

Some of our results are more robust than others

To test the robustness our of first analysis, which focussed on the overall effect of host
population genetic diversity on parasite success, we performed a series of 'leave-one-out'
sensitivity analyses. By averaging over multiple iterations, we showed that the overall effect
of host population genetic diversity on both the mean and variability of parasite infection
success was not dependent on the inclusion of any specific study or set of independent
comparisons in our dataset. Specifically, the overall effect of host population genetic diversity
on mean parasite infection success was significant across both sets of models (Table 3., Fig.
S3 and S4) and the overall effect of host population genetic diversity on variability in parasite
infection success was still not significant (Table 3, Fig. S3 and S4).

Regarding the results of our main analysis, which focussed on how the mean and variability
of parasite success was influenced by a three-way interaction between host population genetic
diversity, parasite population genetic diversity and parasite host range, we were unable to
perform any kind of formal sensitivity analysis due to the complexity of the underlying mixed-
effects meta-analytical models. However, it is important to note that a large number of the
effect sizes corresponding to the subset of data that included specialist parasites with low
parasite population genetic diversity all came from a single study of a prokaryotic host (49 out
of 68, Van Houte et al., 2016). Therefore, the results of some of these meta-analytical models
may be quite sensitive to the inclusion of this one study.

Table 3. Results of the leave-one-out sensitivity analyses. Model averages were calculated
from the iterative exclusion of either one study (Leave1studyout) or set of independent
comparisons (Leave1setout). The following abbreviations are used; ES (effect size), SE
(mean standard error across all models), ci.lb and ci.ub (mean lower and upper bounds of
95% confidence intervals across all models respectively).

Method	ES	Estimate	SE	z-value	p-value	ci.lb	ci.ub
Leave1studyout	SMD	-0.29	0.14	-2.05	0.04	-0.57	-0.01
Leave1studyout	lnCVR	0.02	0.17	0.13	0.87	-0.31	0.35
Leave1setout	SMD	-0.29	0.14	-2.07	0.04	-0.56	-0.02
Leave1setout	lnCVR	0.02	0.17	0.13	0.9	-0.3	0.35

Discussion

By re-analysing the data from two previous studies (Ekroth et al., 2019; Gibson &
2021) we show that the conventional theory that less genetically diverse host populations
experience larger epidemics (King & Lively, 2011) is true only under certain circumstances.
Despite finding a reduction in mean parasite infection success in host populations with higher
genetic diversity that was consistent with previous studies (Ekroth et al., 2019; Gibson &
Nguyen, 2021), this result only applied to specialist parasites that infect only a single host
species. In addition, we found support for our Epidemic Diversity conceptual model, which
predicted that the *variability* in parasite success would depend on a combination of the
population genetic diversity and host range of the parasite. For highly specific host-parasite
interactions (as indicated by a narrow host range), we found that increased host population
genetic diversity reduced the *variability* in parasite infection success when the corresponding
parasite genetic diversity was low, but increased the *variability* in parasite infection success
when parasite diversity was high. Therefore, the idea that less genetically diverse host
populations generally experience larger epidemics may not be best described as 'conventional
wisdom' (King & Lively, 2012).

Our results contrast those from previous meta-analytical studies of the effect of host population
genetic diversity on metrics of parasite infection success, which also investigated the host
range of the parasite as part of their analysis (Ekroth et al., 2019; Gibson & Nguyen, 202). As
already eluded to above, we found that the host range of the parasite was a significant
moderator of the effect of host population genetic diversity on mean metrics of parasite
infection success. Specifically, we found that a high level of host population genetic diversity
tended to limit mean metrics of infection success for specialist, but not generalist, parasites
relative to a low level of host population genetic diversity. One possible explanation for this is
the increased statistical power of our study due to a larger number of effect sizes from
combining the effect size data from these previous analyses (Gibson, 2022). This supports
our original suggestion that parasite host range is closely related to the level of genetic
specificity for infection (because specialist parasites are more likely to have evolved highly
specific, matching-allele-type interactions between host resistance and parasite infectivity
alleles than generalist parasites that are less tightly co-evolved to their range of hosts). It also
highlights the susceptibility of host populations with only a small amount of genetic diversity to
consistently high levels of infection success by specialist parasites.

Our finding that more diverse host populations tend to have smaller mean metrics of infection
success for specialist parasites shows how there is slightly more complexity associated with

conventional wisdom than previously thought (King & Lively, 2012). It also has important
implications for how the level of host population genetic diversity is managed in species of
conservation concern (Meuwissen et al., 2020). For example, one approach to species
management may be to prioritise the maintenance or restoration of genetic diversity in host
populations threatened by specialist parasite species, or by finding a safe approach for
broadening a specialist parasite's host range. For example, the introduction of a novel host or
parasite species, as some form of biological control (Stenberg et al., 2021), that can either act
as a catalyst for host-mediated parasite evolution of greater generality (Bull et al., 2022) or
cause a parasite host shift through direct competition for hosts (for a review, see Bashley,
2015) may broaden the host range of a specialist parasite away from its target host to include
a non-target, pest species. In addition, recent empirical work has started to test the theory that
high host population genetic diversity (*sensu* 'resource heterogeneity') selects for the
evolution, or maintenance, of a broader parasite host range (*sensu* 'niche width', Gibson et
al., 2020). Therefore, understanding how host population genetic diversity is linked to the
evolution of parasite host range in a number of different host-parasite systems should be a
priority for future research.

Again, in contrast to the results of previous meta-analytical studies (Ekroth et al., 2019; Gibson
& Nguyen, 2021), we also found that there was a significant difference between the effect of
host population genetic diversity on the variability in metrics of infection success for specialist
parasites with a high level of parasite population genetic diversity and a low level of parasite
population genetic diversity. Specifically, we showed that a high level of both host and parasite
population genetic diversity increased the variability in metrics of infection success for
specialist parasites relative to host populations with a low level of population genetic diversity,
whereas a high level of host population genetic diversity and a low level of parasite population
genetic diversity decreased the variability in metrics of infections success for specialist
parasites relative to host populations with a low level of population genetic diversity. Although
these previous meta-analytical studies focused on the mean, rather than the variability in
metrics of parasite infection success (Ekroth et al., 2019; Gibson & Nguyen, 2021),
nevertheless the authors of both studies had expected to find a significant effect of parasite
population genetic diversity on the relationship between host population genetic diversity and
mean metrics of parasite infection success and were surprised that there was no such
significant result (Ekroth et al., 2019; Gibson & Nguyen, 2021). In addition to their reduced
statistical power (as mentioned above), one possible reason for this could be that only one out
of two of these studies investigated the interaction between different moderators (Gibson &
Nguyen, 2021). On the other hand, the difference we observed in the variability in metrics of
infection success for specialist parasites between host populations with high parasite

population genetic diversity and low parasite population genetic diversity matched the initial
predictions we made in our proposed Epidemic Diversity theoretical model (Fig. 1). This
confirms previous theories that the benefits of host population genetic diversity for resistance
to disease depend on the corresponding parasite population genetic diversity (Bensch et al.,
2021; Boomsma, 1996; Van Baalen & Beekman, 2006).

This idea that the combination of both host and parasite population genetic diversity influence
the variability in metrics of parasite infection success has important implications for host-
parasite systems in general. As already mentioned previously, not only could such variability
in metrics of parasite infection success be important for predicting the occurrence of potentially
severe disease epidemics, which could benefit conservation by informing genetic diversity
management strategies to prioritise at risk host populations (Meuwissen et al., 2020), but it
could also be central to our ability to protect against future emerging diseases (Altizer et al.,
2006) and for understanding the extent to which disease experiments are repeatable. For
example, our results highlight that host populations with a low level of genetic diversity are
particularly susceptible to consistently large disease epidemics caused by specialist parasites
with a high level of diversity. Conversely, the inconsistent levels of parasite success predicted
for combinations of low host x low parasite and high host x high parasite population genetic
diversity suggest that the repeatability of both laboratory and field experiments may be quite
low, since they are often characterised respectively by such combinations of host-parasite
diversity. Similarly, patterns of future disease occurrence (and emergence) may be more
difficult to predict in such systems compared to those with different combinations of diversity.

In addition to our moderator analysis using models with an interaction term, we also
investigated the effects of eight other contextual factors to evaluate our list of hypotheses
(Table 1). These are the same as the moderators used in previous meta-analyses (Ekroth et
al., 2019; Gibson & Nguyen, 2021), but in comparison to the total number of significant effects
they observed in their analysis (two), our results show that there were six moderator levels
that had significant effects. In particular, the effect of host population genetic diversity on mean
metrics of infection success for microparasites was much more negative than for
macroparasites. In agreement with our original hypothesis, this suggests that the difference in
infection durability between micro- and macroparasites (Sorci, 2014) affects the specificity of
their interactions with the host (Schmid-Hempel & Ebert, 2003). Therefore, we suggest that
macroparasites, due the longer-lasting nature of their infections (Sorci, 2014), are less tightly
coevolved with their hosts and thus have a lower genetic specificity for infection. We also
found that there was a significant negative effect of host population genetic diversity on mean

metrics of parasite infection success for comparisons of outbred versus inbred hosts. Although
such an effect was absent for other host population comparisons, such as between naturally
high and low genetic diversity populations of hosts, it was quite similar to the effect for host
populations composed of select genotypes. Therefore, this could suggest that experimental
manipulations of host population genetic diversity had a stronger effect on mean metrics of
parasite infection success than studies using a purely natural source of hosts. However, it is
worth noting that this result is somewhat inconsistent with the significantly negative effect of
host population genetic diversity on mean metrics of parasite infection success observed for
non-laboratory-based studies, for which the opposite effect was observed in one out of the
two previous meta-analyses (Ekroth et al., 2019). As such, an alternative explanation would
be that the effect of host population genetic diversity on mean metrics of parasite infection
success were exacerbated for outbred versus inbred hosts by the increased susceptibility of
inbred hosts to disease (Coltman et al., 1999).

Other notable observations from the individual models include a significant negative effect of
host population genetic diversity on mean metrics of parasite infection success in sexually
reproducing host populations and a significant negative effect of host population genetic
diversity on the variability in metrics of parasite infection success for host populations
reproducing asexually, which was strongly contrasted against the absence of either a sexually
reproducing or facultatively sexually reproducing host. These results suggest that sexual
reproduction might contribute to the strength of how population genetic diversity limits disease
spread due to greater dissimilarity between genotypes from genetic recombination than
achieved by asexual reproduction (Hamilton et al., 1990), but also that asexual reproduction
can lead to greater disparity between the consistency of metrics of parasite infection success
of host populations with high versus low genetic diversity than other forms of host
reproduction. There was also a significant negative effect of host population genetic diversity
on mean metrics of infection success for parasites that typically kill the host. Compared to less
harmful parasites, this suggests that virulent parasites could select for higher levels of
resistance and greater variation of resistance in the host population (Ekroth et al., 2019).

Despite the potentially exciting nature of our results, there are some additional considerations
that should be taken into account. For example, there is a large number of effect sizes for
specialist parasites with low population genetic diversity, but most of these actually come from
a prokaryotic bacterial host study (Van Houte et al., 2016), rather than a vertebrate or
invertebrate host, which is the case for most of our data. In addition, the host range of the
parasite may not be a reliable estimate of the genetic specificity for infection. The host range

of the parasite was used as a proxy for the genetic specificity for infection, as such a detailed

level of information was not available. Therefore, we made the prediction that highly specific

interactions between host and parasite genotypes (Schmid-Hempel & Ebert, 2003) would be

more likely for tightly coevolving pathogens (i.e. following a MAM of infection, Agrawal & Lively,

2002), as might be expected for specialist, but not generalist parasites. Similarly, the results

of our moderator analysis rely on somewhat arbitrary ways of creating data sub-categories. In

the case of parasite population genetic diversity, comparing mainly natural versus laboratory

strains of parasites could be a poor indication of the effect of parasite population genetic

diversity because the exact level of diversity was not actually quantified. In the case of the

host range of the parasite, this measure is subjective and based somewhat on an incomplete

literature (Hyman & Abedon, 2010).

One other final consideration is that the majority of our data concentrates on the effect of host

population genetic diversity on both the mean and variability in metrics of parasite infection

success for spatially replicated groups of host populations (but see Hale & Briskie, 2007).

Although we might expect the temporal pattern of the effect of host population genetic diversity

on metrics of parasite infection success to be similar to that observed across space, we also

predict some key differences. For example, recurrent bouts of parasite-mediated directional

selection have the ability to reduce host and parasite population genetic diversity over time

(Buckling & Rainey, 2002; Obbard et al., 2011), which could be accompanied by a higher

mean and lower variability in metrics of parasite infection success. However, the maintenance

of host and parasite genetic diversity over time depends on the precise nature of selection and

the underlying host-parasite infection genetics (i.e. a MAM versus a GFG model for genetic

specificity, Boots et al., 2014). Although there are some studies which measure metrics of

parasite infection success for host populations with different levels of genetic diversity at

multiple timepoints (e.g. Florian Altermatt1 & Ebert1, 2008), more studies would be required

to provide a comprehensive test of the effect of host population genetic diversity on metrics of

parasite infection success over time.

Summary

In this study, we measured the difference in the mean and variability in metrics of parasite

infection success between host populations with high versus low genetic diversity. After first

challenging so-called 'conventional wisdom' (*sensu* (King & Lively, 2012)) we proposed a

Epidemic Diversity model to better understand the context around how host population genetic

diversity might affect not only the mean, but also the variability in metrics of parasite infection

success. We found that host population genetic diversity affected mean metrics of infection

success for specialist but not generalist parasites. We also found that the effect of host
population diversity on the variability in metrics of parasite infection success depends on a
combination of the host range of the parasite and the parasite population diversity, such that
there is some evidence for a Epidemic Diversity theoretical model, at least for the collection of
studies reviewed in this meta-analysis. Additionally, we found that there was a number of other
context dependent effects of host population genetic diversity on both the mean and variability
in metrics of parasite infection success, such as parasite type. Overall, these findings
represent a change of perspective that could help to protect vulnerable host populations by
prioritizing how genetic diversity within these populations is managed. Future study of the
Epidemic Diversity hypothesis across a range of plant host-parasite systems would help
generalize these findings.
718
719

References

References

Agha, R., Gross, A., Rohrlack, T., & Wolinska, J. (2018). Adaptation of a chytrid

parasite to its cyanobacterial host is hampered by host intraspecific diversity. *Frontiers in*

Microbiology, 9(MAY), 1–10. https://doi.org/10.3389/fmicb.2018.00921

Agrawal, A., & Lively, C. M. (2002). Infection genetics: Gene-for-gene versus

matching-alleles models and all points in between. *Evolutionary Ecology Research*.

Alan Pounds, J., Bustamante, M. R., Coloma, L. A., Consuegra, J. A., Fogden, M. P.

L., Foster, P. N., La Marca, E., Masters, K. L., Merino-Viteri, A., Puschendorf, R., Ron, S. R.,

Sánchez-Azofeifa, G. A., Still, C. J., & Young, B. E. (2006). Widespread amphibian

extinctions from epidemic disease driven by global warming. *Nature*, *439*(7073), 161–167.

https://doi.org/10.1038/nature04246

Altermatt, F., & Ebert, D. (2008). Genetic diversity of *Daphnia magna* populations

enhances resistance to parasites. *Ecology Letters*. https://doi.org/10.1111/j.1461-

0248.2008.01203.x

Altizer, S., Dobson, A., Hosseini, P., Hudson, P., Pascual, M., & Rohani, P. (2006).

Seasonality and the dynamics of infectious diseases. *Ecology Letters*, *9*(4), 467–484.

https://doi.org/10.1111/j.1461-0248.2005.00879.x

Anderson, R. M., May, R. M., Joysey, K., Mollison, D., Conway, G. R., Cartwell, R.,

Thompson, H. V., & Dixon, B. (1986). The invasion, persistence and spread of infectious

diseases within animal and plant communities. *Philosophical Transactions of the Royal*

Society of London. B, Biological Sciences, *314*(1167), 533–570.

https://doi.org/10.1098/rstb.1986.0072

Baer, B., & Schmid-Hempel, P. (2001). Unexpected consequences of polyandry for

parasitism and fitness in the bumblebee, *Bombus terrestris*. *Evolution*.

https://doi.org/10.1111/j.0014-3820.2001.tb00683.x

Bashey, F. (2015). Within-host competitive interactions as a mechanism for the

maintenance of parasite diversity. *Philosophical Transactions of the Royal Society B:*

Biological Sciences, *370*(1675), 20140301. https://doi.org/10.1098/rstb.2014.0301

Bensch, H. M., O'Connor, E. A., & Cornwallis, C. K. (2021). Living with relatives

offsets the harm caused by pathogens in natural populations. *ELife*, *10*.

https://doi.org/10.7554/eLife.66649

Boomsma, J. J. (1996). Paternity in eusocial Hymenoptera. *Philosophical*

Transactions of the Royal Society of London. Series B: Biological Sciences, *351*(1342), 947–

975. https://doi.org/10.1098/rstb.1996.0087

Boots, M., White, A., Best, A., & Bowers, R. (2014). How specificity and epidemiology
drive the coevolution of static trait diversity in hosts and parasites. *Evolution*, *68*(6), 1594–
1606. https://doi.org/10.1111/evo.12393
Borenstein, M., Hedges, L. V., Higgins, J. P. T., & Rothstein, H. R. (2009).
Introduction to meta-analysis. In *Introduction to Meta-Analysis*.
https://doi.org/10.1002/9780470743386
Browning, J. A., & Frey, K. J. (1969). Multiline cultivars as a means of disease
control. *Annual Review of Phytopathology*, *7*(1), 355–382.
https://doi.org/10.1146/annurev.py.07.090169.002035
Buckling, A., & Rainey, P. B. (2002). Antagonistic coevolution between a bacterium
and a bacteriophage. *Proceedings of the Royal Society B: Biological Sciences*, *269*(1494),
931–936. https://doi.org/10.1098/rspb.2001.1945
Bull, J. J., Wichman, H. A., & Krone, S. M. (2022). Modeling the directed evolution of
broad host range Phages. *Antibiotics*, *11*(12), 1709.
https://doi.org/10.3390/antibiotics11121709
Coltman, D. W., Pilkington, J. G., Smith, J. A., & Pemberton, J. M. (1999). Parasite-
mediated selection against inbred Soay sheep in a free-living, island population. In *Evolution*
(Vol. 53, Issue 4). https://about.jstor.org/terms
Dagan, Y., Liljeroos, K., Jokela, J., & Ben-Ami, F. (2013). Clonal diversity driven by
parasitism in a freshwater snail. *Journal of Evolutionary Biology*, *26*(11).
https://doi.org/10.1111/jeb.12245
De Castro, F., & Bolker, B. (2004). Mechanisms of disease-induced extinction.
Ecology Letters, *8*(1), 117–126. https://doi.org/10.1111/j.1461-0248.2004.00693.x
Dionne, M., Miller, K. M., Dodson, J. J., & Bernatchez, L. (2009). MHC standing
genetic variation and pathogen resistance in wild Atlantic salmon. *Philosophical*
Transactions of the Royal Society B: Biological Sciences, *364*(1523).
https://doi.org/10.1098/rstb.2009.0011
Dybdahl, M. F., Jenkins, C. E., & Nuismer, S. L. (2014). Identifying the molecular
basis of host-parasite coevolution: Merging models and mechanisms. *The American*
Naturalist, *184*(1), 1–13. https://doi.org/10.1086/676591
Ekroth, A. K. E., Rafaluk-Mohr, C., & King, K. C. (2019). Host genetic diversity limits
parasite success beyond agricultural systems: a meta-analysis. *Proceedings of the Royal*
Society B: Biological Sciences, *286*(1911), 20191811.
https://doi.org/10.1098/rspb.2019.1811

788   Ellison, A., Cable, J., & Consuegra, S. (2011). Best of both worlds? Association
789 between outcrossing and parasite loads in a selfing fish. *Evolution*, *65*(10), 3021–3026.
790 https://doi.org/10.1111/j.1558-5646.2011.01354.x

791   Elton, C. S. (1958). *The Ecology of Invasions by Animals and Plants*. Springer US.
792 https://doi.org/10.1007/978-1-4899-7214-9

793   Field, A. P., & Gillett, R. (2010). How to do a meta-analysis. *British Journal of*
794 *Mathematical and Statistical Psychology*, *63*(3). https://doi.org/10.1348/000711010X502733

795   Field, S. G., Lange, M., Schulenburg, H., Velavan, T. P., & Michiels, N. K. (2007).
796 Genetic diversity and parasite defense in a fragmented urban metapopulation of
797 earthworms. *Animal Conservation*, *10*(2). https://doi.org/10.1111/j.1469-1795.2006.00084.x

798   Ganz, H. H., & Ebert, D. (2010). Benefits of host genetic diversity for resistance to
799 infection depend on parasite diversity. In *Source: Ecology* (Vol. 91, Issue 5). https://www-
800 jstor-
801 org.ezproxy.stir.ac.uk/stable/pdf/25661177.pdf?refreqid=excelsior%3A5c0f30d81cb9568626
802 bbc1823fa625af

803   Garrett, K. A., & Mundt, C. C. (1999). Epidemiology in mixed host populations.
804 *Phytopathology*, *89*(11), 984–990. https://doi.org/10.1094/PHYTO.1999.89.11.984

805   Gibson, A. K. (2022). Genetic diversity and disease: The past, present, and future of
806 an old idea. *Evolution*, *76*(S1), 20–36. https://doi.org/10.1111/evo.14395

807   Gibson, A. K., Baffoe-Bonnie, H., Penley, M. J., Lin, J., Owens, R., Khalid, A., &
808 Morran, L. T. (2020). The evolution of parasite host range in heterogeneous host
809 populations. *Journal of Evolutionary Biology*, *33*(6), 773–782.
810 https://doi.org/10.1111/jeb.13608

811   Gibson, A. K., & Nguyen, A. E. (2020). Does genetic diversity protect host
812 populations from parasites? A meta-analysis across natural and agricultural systems.
813 *Evolution Letters*, *5*(1). https://doi.org/10.1002/evl3.206

814   Giese, A. R., & Hedrick, P. W. (2003). Genetic variation and resistance to a bacterial
815 infection in the endangered Gila topminnow. *Animal Conservation*, *6*(4).
816 https://doi.org/10.1017/S1367943003003445

817   Hale, K. A., & Briskie, J. V. (2007). Decreased immunocompetence in a severely
818 bottlenecked population of an endemic New Zealand bird. *Animal Conservation*, *10*(1).
819 https://doi.org/10.1111/j.1469-1795.2006.00059.x

820   Hamilton, W. D. (1987). *Kinship, recognition, disease, and intelligence: constraints of*
821 *social evolution*. Japan Scientific Societies Press.

Higgins, J. P. T., Thomas J., Chandler J., Cumpston M., Li T., Page M. J., Welch V.

A. (2024). *Cochrane Handbook for Systematic Reviews of Interventions* (Version 5.) The

Cochrane Collaboration, 2011. www.handbook.cochrane.org

Hothorn, T., Bretz, F., & Westfall, P. (2008). Simultaneous inference in general

parametric models. *Biometrical Journal*, *50*(3), 346–363.

https://doi.org/10.1002/bimj.200810425

Hyman, P., & Abedon, S. T. (2010). Bacteriophage host range and bacterial

resistance. *Advances in Applied Microbiology*, *70*, 217–248. https://doi.org/10.1016/S0065-

2164(10)70007-1

Jennions, M. D. (2003). A survey of the statistical power of research in behavioral

ecology and animal behavior. *Behavioral Ecology*, *14*(3), 438–445.

https://doi.org/10.1093/beheco/14.3.438

Johnson, P. T. J., Ostfeld, R. S., & Keesing, F. (2015). Frontiers in research on

biodiversity and disease. *Ecology Letters*, *18*(10), 1119–1133.

https://doi.org/10.1111/ele.12479

Johnson, P. T. J., Preston, D. L., Hoverman, J. T., Henderson, J. S., Paull, S. H.,

Richgels, K. L. D., & Redmond, M. D. (2012). Species diversity reduces parasite infection

through cross-generational effects on host abundance. *Ecology*, *93*(1), 56–64.

https://doi.org/10.1890/11-0636.1

Keesing, F., Belden, L. K., Daszak, P., Dobson, A., Harvell, C. D., Holt, R. D.,

Hudson, P., Jolles, A., Jones, K. E., Mitchell, C. E., Myers, S. S., Bogich, T., & Ostfeld, R. S.

(2010). Impacts of biodiversity on the emergence and transmission of infectious diseases. In

Nature. https://doi.org/10.1038/nature09575

Keesing, F., Holt, R. D., & Ostfeld, R. S. (2006). Effects of species diversity on

disease risk. *Ecology Letters*, *9*(4), 485–498. https://doi.org/10.1111/j.1461-

0248.2006.00885.x

Keesing, F., & Ostfeld, R. S. (2021). Dilution effects in disease ecology. *Ecology*

Letters, *24*(11). https://doi.org/10.1111/ele.13875

King, K. C., Jokela, J., & Lively, C. M. (2011). Parasites, sex, and clonal diversity in

natural snail populations. *Evolution*, *65*(5). https://doi.org/10.1111/j.1558-5646.2010.01215.x

King, K. C., & Lively, C. M. (2012). Does genetic diversity limit disease spread in

natural host populations? *Heredity*, *109*(4), 199–203. https://doi.org/10.1038/hdy.2012.33

Koricheva, J., & Gurevitch, J. (2014). Uses and misuses of meta-analysis in plant

ecology. *Journal of Ecology*, *102*(4), 828–844. https://doi.org/10.1111/1365-2745.12224

Koskella, B., & Lively, C. M. (2009). Evidence for negative frequency-dependent

selection during experimental coevolution of a freshwater snail and a sterilizing trematode.

Evolution, *63*(9), 2213–2221. https://doi.org/10.1111/j.1558-5646.2009.00711.x

Kyle, C. J., Rico, Y., Castillo, S., Srithayakumar, V., Cullingham, C. I., White, B. N., &

Pond, B. A. (2014). Spatial patterns of neutral and functional genetic variations reveal

patterns of local adaptation in raccoon (*Procyon lotor*) populations exposed to raccoon

rabies. *Molecular Ecology*, *23*(9). https://doi.org/10.1111/mec.12726

Lagisz, M., Senior, A., Nakagawa, S., & Noble, D. (2024). _metaAidR: Functions and

*Data for Aiding Meta-analyses in Ecology and Evolution_* (0.0.0.9000).

Leonard, K. J. (1969). Factors affecting rates of stem rust increase in mixed plantings

of susceptible and resistant oat varieties cultivars. *Phytopathology*, *59*, 1845–1850.

Loiseau, C., Zoorob, R., Robert, A., Chastel, O., Julliard, R., & Sorci, G. (2011).

Plasmodium relictum infection and MHC diversity in the house sparrow (*Passer domesticus*).

Proceedings of the Royal Society B: Biological Sciences, *278*(1709).

https://doi.org/10.1098/rspb.2010.1968

Meagher, S. (1999). Genetic diversity and *capillaria hepatica* (nematoda) prevelance

in Michigan deer mouse populations. *Evolution*, *53*(4), 1318–1324.

https://doi.org/10.2307/2640837

Meuwissen, T. H. E., Sonesson, A. K., Gebregiwergis, G., & Woolliams, J. A. (2020).

Management of genetic diversity in the era of genomics. *Frontiers in Genetics*, *11*.

https://doi.org/10.3389/fgene.2020.00880

Mihaljevic, J. R., Joseph, M. B., Orlofske, S. A., & Paull, S. H. (2014). The scaling of

host density with richness affects the direction, shape, and detectability of diversity-disease

relationships. *PLoS ONE*, *9*(5), e97812. https://doi.org/10.1371/journal.pone.0097812

Nakagawa, S., Noble, D. W. A., Lagisz, M., Spake, R., Viechtbauer, W., & Senior, A.

M. (2023). A robust and readily implementable method for the meta-analysis of response

ratios with and without missing standard deviations. *Ecology Letters*, *26*(2), 232–244.

https://doi.org/10.1111/ele.14144

Nakagawa, S., Poulin, R., Mengersen, K., Reinhold, K., Engqvist, L., Lagisz, M., &

Senior, A. M. (2015). Meta-analysis of variation: ecological and evolutionary applications and

beyond. *Methods in Ecology and Evolution*, *6*(2), 143–152. https://doi.org/10.1111/2041-

210X.12309

Neumann, P., & Moritz, R. F. A. (2000). Testing genetic variance hypotheses for the

evolution of polyandry in the honeybee (*Apis mellifera* L.). *Insectes Sociaux*, *47*(3).

https://doi.org/10.1007/PL00001714

Obbard, D. J., Jiggins, F. M., Bradshaw, N. J., & Little, T. J. (2011). Recent and

recurrent selective sweeps of the antiviral RNAi gene Argonaute-2 in three species of

Drosophila. *Molecular Biology and Evolution*, *28*(2), 1043–1056.

https://doi.org/10.1093/molbev/msq280

O'Brien, S. J., & Evermann, J. F. (1988). Interactive influence of infectious disease

and genetic diversity in natural populations. In *Trends in Ecology and Evolution* (Vol. 3, Issue

10). https://doi.org/10.1016/0169-5347(88)90058-4

Ostfeld, R. S., & Keesing, F. (2012). Effects of host diversity on infectious disease.

Annual Review of Ecology, Evolution, and Systematics, *43*(1), 157–182.

https://doi.org/10.1146/annurev-ecolsys-102710-145022

Page, M. J., McKenzie, J. E., Bossuyt, P. M., Boutron, I., Hoffmann, T. C., Mulrow, C.

D., Shamseer, L., Tetzlaff, J. M., Akl, E. A., Brennan, S. E., Chou, R., Glanville, J.,

Grimshaw, J. M., Hróbjartsson, A., Lalu, M. M., Li, T., Loder, E. W., Mayo-Wilson, E.,

McDonald, S., McGuinness, Luke A., Stewart, L. A., Thomas, J., Tricco, A. C., Welch, V. A.,

Whiting, P., Moher, D. (2021). The PRISMA 2020 statement: an updated guideline for

reporting systematic reviews. *BMJ*, n71. https://doi.org/10.1136/bmj.n71

Paplauskas, S., Duthie, B., & Tinsley, M. C. (2024). Meta-analysis of host population-

level diversity and variation in 'parasite success' [Dataset]. *Dryad*.

https://doi.org/https://doi.org/10.5061/dryad.2bvq83bzq

Parsche, S., & Lattorff, M. H. G. (2018). The relative contributions of host density and

genetic diversity on prevalence of a multi-host parasite in bumblebees. *Biological Journal of*

the Linnean Society, *125*(4). https://doi.org/10.1093/biolinnean/bly151

Pierce, A. A., De Roode, J. C., Altizer, S., & Bartel, R. A. (2014). Extreme

heterogeneity in parasitism despite low population genetic structure among monarch

butterflies inhabiting the Hawaiian Islands. *PLoS ONE*, *9*(6).

https://doi.org/10.1371/journal.pone.0100061

Puurtinen, M., Hytönen, M., Knott, K. E., Taskinen, J., Nissinen, K., & Kaitala, V.

(2004). The effects of mating system and genetic variability on susceptibility to trematode

parasites in a freshwater snail, *Lymnaea stagnalis*. *Evolution*, *58*(12).

https://doi.org/10.1111/j.0014-3820.2004.tb01626.x

Queirós, J., Vicente, J., Alves, P. C., de la Fuente, J., & Gortazar, C. (2016).

Tuberculosis, genetic diversity and fitness in the red deer, *Cervus elaphus*. *Infection,*

Genetics and Evolution, *43*. https://doi.org/10.1016/j.meegid.2016.05.031

R Core Team. (2023). _R: A Language and Environment for Statistical Computing_.

R Foundation for Statistical Computing. https://www.r-project.org/

Rahn, A. K., Krassmann, J., Tsobanidis, K., MacColl, A. D. C., & Bakker, T. C. M.
(2016). Strong neutral genetic differentiation in a host, but not in its parasite. *Infection,*
Genetics and Evolution, *44*. https://doi.org/10.1016/j.meegid.2016.07.011

Savage, A. E., Becker, C. G., & Zamudio, K. R. (2015). Linking genetic and
environmental factors in amphibian disease risk. *Evolutionary Applications*, *8*(6).
https://doi.org/10.1111/eva.12264

Schmid-Hempel, P., & Ebert, D. (2003). On the evolutionary ecology of specific
immune defence. *Trends in Ecology & Evolution*, *18*(1), 27–32.
https://doi.org/10.1016/S0169-5347(02)00013-7

Schmidt, A. M., Linksvayer, T. A., Boomsma, J. J., & Pedersen, J. S. (2011). No
benefit in diversity? The effect of genetic variation on survival and disease resistance in a
polygynous social insect. *Ecological Entomology*, *36*(6). https://doi.org/10.1111/j.1365-
2311.2011.01325.x

Sherman, P. W., Seeley, T. D., & Reeve, H. K. (1988). Parasites, pathogens, and
polyandry in social Hymenoptera. In *Source: The American Naturalist* (Vol. 131, Issue 4).
https://www.jstor.org/stable/pdf/2461747.pdf?refreqid=excelsior%3A10d16706d07e226edd1f
3ace6f23a6b4

Sorci, G. (2014). Immunology of parasitism. In *Reference Module in Biomedical*
Sciences. Elsevier. https://doi.org/10.1016/B978-0-12-801238-3.00253-1

Stenberg, J. A., Sundh, I., Becher, P. G., Björkman, C., Dubey, M., Egan, P. A.,
Friberg, H., Gil, J. F., Jensen, D. F., Jonsson, M., Karlsson, M., Khalil, S., Ninkovic, V.,
Rehermann, G., Vetukuri, R. R., & Viketoft, M. (2021). When is it biological control? A
framework of definitions, mechanisms, and classifications. *Journal of Pest Science*, *94*(3),
665–676. https://doi.org/10.1007/s10340-021-01354-7

Sutton, A. J. (2009). Publication bias. In *The handbook of research synthesis and*
meta-analysis. (2nd ed., pp. 435–452). Russell Sage Foundation.

Tarpy, D. R. (2003). Genetic diversity within honeybee colonies prevents severe
infections and promotes colony growth. *Proceedings of the Royal Society B: Biological*
Sciences, *270*(1510). https://doi.org/10.1098/rspb.2002.2199

Trouvé, S., Degen, L., Renaud, F., & Goudet, J. (2003). Evolutionary implications of
a high selfing rate in the freshwater snail *Lymnaea truncatula*. *Evolution*, *57*(10).
https://doi.org/10.1554/02-452

Van Baalen, M., & Beekman, M. (2006). The costs and benefits of genetic
heterogeneity in resistance against parasites in social insects. In *American Naturalist*.
https://doi.org/10.1086/501169

van der Plank, J. E. (1963). *Plant diseases: epidemics and control*. Academic Press.

Van Houte, S., Ekroth, A. K. E., Broniewski, J. M., Chabas, H., Ashby, B., Bondy-
Denomy, J., Gandon, S., Boots, M., Paterson, S., Buckling, A., & Westra, E. R. (2016). The
diversity-generating benefits of a prokaryotic adaptive immune system. *Nature*, *532*(7599).
https://doi.org/10.1038/nature17436

Velavan, T. P., Weller, S., Schulenburg, H., & Michiels, N. K. (2009). High genetic
diversity and heterogeneous parasite load in the earthworm *Lumbricus terrestris* on a
German meadow. *Soil Biology and Biochemistry*, *41*(8).
https://doi.org/10.1016/j.soilbio.2009.03.026

Viechtbauer, W. (2010). Conducting Meta-Analyses in R with the metafor Package.
Journal of Statistical Software, *36*(3). https://doi.org/10.18637/jss.v036.i03

Whitehorn, P. R., Tinsley, M. C., Brown, M. J. F., Darvill, B., & Goulson, D. (2011).
Genetic diversity, parasite prevalence and immunity in wild bumblebees. *Proceedings of the*
Royal Society B: Biological Sciences, *278*(1709), 1195–1202.
https://doi.org/10.1098/rspb.2010.1550

Whitehorn, P. R., Tinsley, M. C., Brown, M. J. F., Darvill, B., & Goulson, D. (2014).
Genetic diversity and parasite prevalence in two species of bumblebee. *Journal of Insect*
Conservation, *18*(4). https://doi.org/10.1007/s10841-014-9673-1

Whiteman, N. K., Matson, K. D., Bollmer, J. L., & Parker, P. G. (2006). Disease
ecology in the Galápagos Hawk (*Buteo galapagoensis*): host genetic diversity, parasite load
and natural antibodies. *Proceedings of the Royal Society B: Biological Sciences*, *273*(1588),
797–804. https://doi.org/10.1098/rspb.2005.3396

Data accessibility

985   Reviewer URL:
986   https://datadryad.org/stash/share/E2NqLZ8KL2oYaPQLSYatmNNIeUub3aeC4ExfjJvE5Hw
987   Data available from the Dryad Digital Repository: doi:10.5061/dryad.2bvq83bzq (Paplauskas
988   et al., n.d.).

Acknowledgements

991   We are grateful to Assistant Professor Alexander Strauss, Dr Camille Ameline, Dr Peter
992   Thrall, Dr Jason Walsman and Assistant Professor Amanda Kyle Gibson for sharing their
993   raw data and clarifying their methods with us. We also thank Dr Laura Braunholtz, Dr
994   Katherine Raines, Dr Luc Bussière and Kyle Morrison for their general statistical and meta-
995   analytical advice.

Supplementary information

Supplementary table S1. The difference between our amended study inclusion criteria and the original study inclusion criteria.

New study inclusion criteria	Original study inclusion criteria	Why changed
1) 'Parasite success', which we define as the ability of a parasite to spread among hosts (transmission rate, infection rate, prevalence), replicate on / within hosts (macro / microparasite load, disease severity), or kill hosts (virulence i.e. host survival / mortality rate) was measured among replicate populations across time or space.	- Ekroth, Rafaluk-Mohr and King, 2019: Defined parasite success as any measure of a parasite's ability to proliferate within a host population. - Gibson and Nguyen, 2020: Focused on population-level parasitism, including prevalence, load and virulence.	We combined the two previous versions of the study inclusion criteria to include several different measures of parasite success, which were later used for contextual factor analysis.
2) Parasite success data was collected from two or more host populations with any comparable difference in genetic diversity, such as the level of relatedness among individuals (inbred versus outbred), genotypic diversity (high versus low) or heterozygosity.	- Ekroth, Rafaluk-Mohr and King, 2019: Data was collected from any study with two distinct populations and any measured difference in diversity. - Gibson and Nguyen, 2020: Collected data for two or more populations.	We collected data from studies of multiple populations with any comparable difference in genetic diversity to increase our sample size and because there was one study with differences in genetic diversity which were not comparable between all pairwise combinations (Baer 2001).
3) Genetic diversity was measured at the host population level and not community diversity or	- Ekroth, Rafaluk-Mohr and King, 2019: Used the exact same wording.	We followed both Ekroth, Rafaluk-Mohr and King, 2019 and Gibson and

individual-level genetic heterozygosity.	- Gibson and Nguyen, 2020: Stated that host genetic diversity had to be intra-specific.	Nguyen 2020 in this criterion.
4) The study focused on an animal (or bacterial) host species.	- Ekroth, Rafaluk-Mohr and King, 2019: Excluded studies of agricultural systems. - Gibson and Nguyen, 2020: Did not specify the study system.	We did not include any studies of non-animal populations, except for prokaryotic bacteria, because a more detailed analysis of the plant literature would require a separate review.
5) The study does not re-analyze the data from a previously published study.	- Both Ekroth, Rafaluk-Mohr and King, 2019 and Gibson and Nguyen, 2020: Did not include this specification.	We included this specification because Ekroth, Rafaluk-Mohr and King, 2019 included data from two different studies by Baer and Schmid-Hempel which were based on the same dataset.
6) The parasite success data was not replicated simply by using an alternate way of measuring host population diversity.	- Both Ekroth, Rafaluk-Mohr and King, 2019 and Gibson and Nguyen, 2020: Did not include this specification.	We included this specification because there two studies included by the previous meta-analyses (Giese 2003 and Puurtinen 2004) which included parasite success data for the same populations with two different measures of genetic diversity, which was a form of pseudoreplication.

7) An attempt to take the parasite success data from clearly illegible figures was not made.	- Both Ekroth, Rafaluk-Mohr and King, 2019 and Gibson and Nguyen, 2020: Did not include this specification.	We included this specification because Gibson and Nguyen, 2020 had collected data from two studies with illegible figures (Agha, 2018 and van Houte et al. 2016.

1001

1002

	Low host div.		High host div.	
	Replicate 1	Replicate 2	Replicate 1	Replicate 2
High intra-, low inter-				
High inter-, low intra-				

Figure S1. The difference between intra- and inter-group level variation in host population genetic diversity. Each bucket represents a unique host population (e.g. jar of *Daphnia*). Each shape represents a unique host genotype. There are two sets of three host populations per replicate which correspond to the high and low diversity groups. The first two replicates show a combination of high intra-group variation in host population genetic diversity x low inter-group variation in host population genetic diversity. The second set of replicates show a combination of low intra-group variation in host population genetic diversity x high inter-group variation in host population genetic diversity.

Supplementary table S2. The taxonomic distribution of unique host and parasite species
(genera) within the data. The number of unique combinations of host and parasite species are
shown in each cell of the table, along with the number of studies they are sourced from after
a backslash. The total number of studies (59) is higher than the total number of studies in our
dataset (48), because there were some studies with multiple comparisons of unique host and
parasite combinations. The colour coding is based on the number of unique combinations of
host and parasite genera.

1013

		Host group			
		Prokaryote	Invertebrate	Vertebrate	Total
Parasite group	Animal	0 (0) / 0	14 (12) / 6	13 (10) / 8	27 (22) / 14
	Bacteria	0 (0) / 0	2 (2) / 2	16 (13) / 5	18 (15) / 7
	Fungi	1 (1) / 1	18 (15) / 16	1 (1) / 1	20 (17) / 18
	Protozoa	0 (0) / 0	12 (7) / 11	3 (3) / 2	15 (10) / 13
	Virus	1 (1) / 1	7 (3) / 2	4 (4) / 4	12 (7) / 7
	Total	2 (1) / 2	53 (39) / 37	37 (31) / 20	92 (71) / 59

1014
1015
1016
1017

Figure S2. Study effects of host population genetic diversity on the mean and variability in parasite success. The x-axis in each plot shows the effect of increasing host population genetic diversity on either A) the difference in mean parasite success (SMD) or B) the difference in the variability in parasite success (lnCVR). Aggregated effects for each study are shown with 95% confidence intervals. Where the same host genus was studied more than once ('Duplicated Genus'), the colour of the points is white, rather than black, and the specific host genus studied is indicated by its shape (there were only five duplicated host genera). Each point is scaled by the amount of weighting they received in an aggregated mixed effects model, whereas the actual analysis was conducted based on the full set of 211 individual data points. The dashed lines indicate an effect size of zero and the overall model means are shown by the solid grey line with 95% confidence intervals bands in light grey.

Figure S3. The results of the leave-one-study-out method of sensitivity analysis. The x-axis in each plot shows the effect of increasing host population genetic diversity on either A) the difference in mean parasite success (SMD) or B) the difference in the variability in parasite success (lnCVR). The names of the authors and the publication date for the study omitted in each model iteration is shown on the left, with the overall effect size and its confidence interval shown on in the middle. The mean effect size across all models is shown by the vertical line and specific value is shown on the right (with 95% confidence intervals). The size of each point is scaled according to its precision.

Figure S4. The results of the leave-one-independent-comparison-out method of sensitivity analysis visualized using a modified version of an orchard plot. The x-axis in each plot shows the effect of increasing host population genetic diversity on either A) the difference in mean parasite success (SMD) or B) the difference in the variability in parasite success (lnCVR). Unlike traditional orchard plots, which show the distribution of individual effect sizes, the mean effect size for each model iteration is shown by the coloured circles. The size of each point is scaled by its precision (inverse of the standard error). The mean effect size across all models is shown by the dashed line.

1022

1023