1   **Title:** The effect of host population genetic diversity on variation in metrics of

2   parasite success

Authors: Sam Paplauskas[1]*, Brad Duthie[1], Matthew Tinsley[1]

[1] Biological & Environmental Sciences, University of Stirling, Stirling, United Kingdom

* Corresponding author

Email: sam.paplauskas@stir.ac.uk

Keywords: Genetic diversity, intra-specific diversity, disease dilution, monoculture

effect, meta-analysis

ABSTRACT

Conventional wisdom suggests that populations with lower levels of genetic diversity

are at a greater risk of the more harmful effects of disease. However, previous attempts

to qualify this proposition have focused on measuring the mean, rather than the

variability, in metrics of parasite success. Since the ability of host population genetic

diversity to limit the spread of disease requires some specificity between hosts and

parasites, and the benefits of host population genetic diversity in resistance to infection

may depend on the respective parasite population genetic diversity, we propose a

diversity-uncertainty model which predicts that the mean and variability in parasite

success depend on a combination of host range and parasite population genetic

diversity. By re-analyzing a dataset combining 48 studies collected by previous meta-

analyses, we show that the effect of host population genetic diversity reduces the mean

success of single-host, but not host generalist, parasites. We find evidence for our

original hypothesis that the variability of parasite success depends on a combination

of host population genetic diversity, parasite population genetic diversity and host

range. Together, these results challenge conventional wisdom and have important

implications for how genetic diversity can be better managed in host populations.

INTRODUCTION

It is commonly believed that host populations with lower genetic diversity are at a greater risk of experiencing higher parasite success (i.e. disease (1)). This refers to the population-level prevalence (proportion of infected hosts), virulence (parasite-induced loss of fitness) or parasite load (average parasites per host (2–4)).

Previous studies of the generality of this proposed 'conventional wisdom' (1), have often focused on measuring the mean, rather than the variability, of parasite success (5,6). This is surprising, considering the importance of parasitic extremes, in terms of epidemics and whether they cause mass extinction (7,8), the predictably of recurrent bouts of disease across years and the repeatability of disease experiments in general. As a result, the relationship between host diversity and variability in parasite success is poorly understood (9). However, it is central to our ability to protect against future emerging diseases (10).

The implications of host community, species or genetic diversity on infectious diseases is often referred to as 'disease dilution' (11–15), the diversity-disease hypothesis (16–18) or the monoculture effect (19–23). This can be caused by an increase in individual host susceptibility (24), or a variety of population-level effects such as reducing the rate of encounter between susceptible and infectious individuals (encounter reduction), reducing the probability of transmission given an encounter (transmission reduction), decreasing the density of susceptible individuals (susceptible host regulation), increasing the recovery rate (recovery augmentation), or increasing the death rate of infected individuals (infected host mortality) (for a review, see (14)).

Although the exact mechanism is unclear, the negative relationship between host population genetic diversity and disease spread is often attributed to encounter reduction (25). Specifically, assuming that there is some level of matching (or genetic specificity (26)) required for a successful infection to occur (*sensu* matching alleles

model [MAM] (27)), there should be a lower chance of a parasite encountering a

susceptible host as it spreads through a more diverse host population. Since the

strength of genetic specificity varies across different host-parasite systems (27), we

might expect that the effect of host population diversity on parasite success depends

on the level of specificity for infection.

In theory, the effects of host population diversity on parasite success may also depend

on the level of parasite diversity (28,29). For example, if there is a high level of genetic

specificity for infection (*sensu* MAM), then we might expect host populations composed

of a single genotype to be entirely susceptible to a single parasite genotype, which is

much more likely to occur in a population with a high level of parasite diversity (28,29).

One empirical study in a *Daphnia* host-parasite system found that the benefits of host

genetic diversity for resistance to infection were reliant on a high level of parasite

diversity (30).

Therefore, if we assume that there is a high level of genetic specificity for infection

(*sensu* MAM) and both host and parasite populations are characterized by either high

or low levels of genetic diversity, we can predict the following patterns for both the

mean and variability in parasite success (Table 1):

A) Low host x low parasite population genetic diversity (Table 1A): We predict that

there will be a high level of variability in parasite success, due to the host population

being composed entirely of susceptible, or resistant, host genotypes, and an

intermediate level of mean parasite success (determined by the overall frequency of

resistant cf. susceptible populations).

B) High host x low parasite population genetic diversity (Table 1B): We predict that

there will be a low level of both mean parasite success and variability in parasite

success, due to the consistency of hosts to resist infection through a reduced

encounter rate with matching parasite genotypes.

C) Low host x high parasite population genetic diversity (Table 1C): We predict that there will be a high level of mean parasite success and a low level of variability in parasite success, due to the consistency of parasite transmission through an enhanced encounter rate with matching host genotypes.

D) High host x high parasite population genetic diversity (Table 1D): We predict that there will be an intermediate level of both mean parasite success and variability in parasite success, due to the diverging effects of host and parasite genetic diversity leading to an inconsistent encounter rate between matching host and parasite genotypes.

Collectively, these predictions form our 'diversity-uncertainty' model for predicting the mean and variability of parasite success across populations with different levels of host and parasite diversity. This builds on previous work (31), which focused on the relationship between population diversity and variability in parasite-induced host mortality and pathogen abundance for only three out of the four possible combinations in Table 1, without also acknowledging the influence of the genetic specificity for infection on these hypotheses.

To test our diversity-uncertainty model, we examine the relationship between host population genetic diversity, parasite population genetic diversity and variability in parasite success for different levels of a proxy for genetic specificity using meta-analysis. After confirming the results of previous studies (5,6), which found a significant difference in mean parasite success between various host populations with high versus low genetic diversity, we then extend their analysis to a study of variability using a suite of different moderator variables.

In particular, we compare the difference in the variability of parasite success between host populations with high versus low genetic diversity using a combination of host

range and parasite population genetic diversity variables. Since the requisite

population-level genetic specificity data does not exist for all of the combined studies,

and MAM dynamics are more likely to rely on coevolution between a single host and

parasite species compared to GFG, due to being driven by negative frequency-

dependent selection (32), we assume that there is a relatively higher level of genetic

specificity for parasites with a narrow host range (1 species) compared to parasites

with a broad host range (> 1 species). On the other hand, we do not make any

predictions about the mean level of, or level of variability in, parasite success for

systems with a low genetic specificity for infection.

Overall, we find that the relationship between host genetic diversity and both the mean

level of, and level of variability in, parasite success depends on a combination of host

range and parasite genetic diversity.

Table 1. Our 'diversity-uncertainty' model. This shows the expected effects of host and

parasite population genetic diversity on i) the mean (μ) and ii) the variability (σ) of

parasite success for highly specific host-parasite systems (e.g. matching-allele model

for infection genetics).

	Low diversity parasite	High diversity parasite
Low diversity host	A) Medium μ High σ	C) High μ Low σ
High diversity host	B) Low μ Low σ	D) Medium μ Medium σ

METHODS

Summary

We combined the data from two previous meta-analyses (5,6), which compared the

standardized mean difference (SMD) in parasite success between host populations

with qualitatively 'high' versus 'low' levels of genetic diversity, to calculate the log

coefficient of variation ratio (lnCVR). In other words, we calculate the mean difference

and ratio of variation in parasite success between host populations with different levels

of genetic diversity. Across all the studies we include these replicates are sometimes

experimental replicates in a lab, sometimes multiple natural populations, and

sometimes repeated measures on single populations along a time series.

Data collection

The data collection for each control versus treatment comparison, which was later used

in calculating effect sizes, involved five main steps (Figure 1):

1) First, we combined the list of studies from (5) and (6), removed any duplicate studies

and added the data used to calculate the effect size, SMD, and its sampling variance

in the original studies (mean, standard deviation, sample size), the metric of parasite

success and the unique study, experiment and replicate identifiers used to account for

the non-independence of separate effect sizes. We did not use the parasite success

data from (6) because the original data extracted from each study was missing from

their online supplementary material, so we would be unable to check the accuracy of

their data during validation (step 3), so we extracted the replicate or population

summaries from the data ourselves after step 3 and recalculated the mean, standard

deviation, sample size etc. ourselves.

In addition, the data used to calculate Fisher's z (an effect size for the difference

between two correlation coefficients) for the observational field studies from (5), was

not in the correct format to calculate either SMD or lnCVR. Therefore, we did not

include this information (from multiple populations with a continuous measure of

genetic diversity) at this stage and instead extracted the data ourselves and

recalculated it during steps 4 and 5. Also, we excluded any plant (wild or agricultural)

host data because a more detailed analysis of the plant literature would require a

separate review and we did not want to add effect size data for studies we already

knew were going to exclude during the next step.

2) Second, we amended the original study inclusion criteria (Table SX) and removed

any studies, experiment or comparisons which did not meet these criteria:

(i) 'Parasite success', which we define as the ability of a parasite to spread among

hosts (transmission rate, infection rate, prevalence), replicate on / within hosts (macro

/ microparasite load, disease severity) or kill hosts (virulence i.e. host survival /

mortality rate) was measured among replicate populations across time or space.

(ii) Parasite success data was collected from two or more host populations with a

difference in genetic diversity, such as the level of relatedness among individuals

(inbred versus outbred), genotypic diversity (high versus low) or heterozygosity.

(iii) Genetic diversity was quantified at the host population level and not community

diversity or individual-level genetic heterozygosity.

(iv) The study focused on an animal (or bacterial) host species.

(v) The study does not re-analyze the data from a previously published study.

(vi) The parasite success data was not replicated simply by using an alternate way of

measuring host population diversity.

(vii) An attempt to take the parasite success data from clearly illegible figures was not

made.

3) Third, we checked the accuracy of the data from the excel spreadsheets used to

calculate the summary of the parasite data for each control and treatment group (5).

There were 11 studies, 13 experiments and 27 comparisons which did not match the published raw data (available in the main text or online or in the supplementary material of each publication), 3 studies, 8 experiments and 32 comparisons which were calculated wrong and 4 studies, 5 experiments and 10 comparisons which had not been transferred from the excel spreadsheets into the final metadata file correctly. Therefore, we fixed any errors in how the parasite success data was calculated, annotated the data correctly with the corresponding information and removed any data which was incorrect. Admittedly, there was one study which we could not check, because the original data was sent by personal communication from (33) to (5), but we included it in our analysis anyway.

4) Fourth, we collected any data we had removed by going back to the original publications and extracting the data from the main text or supplementary files (we used PlotDigitizer (https://plotdigitizer.com/) for any figures). This included 26 studies which met our inclusion criteria, but were removed because they were either missing the replicate-level raw data (6) or they were observational field studies based on multiple populations with a continuous measure of genetic diversity (5), 11 studies, 13 experiments and 27 comparisons which we removed because they did not match the published data, 4 studies, 5 experiments and 10 comparisons which were removed because they were annotated incorrectly, and additional data for 3 comparisons (34–36) that were not made in the original analysis by (5).

We also collected information on 10 different moderators (for why, see Table 2) by standardizing or recoding existing moderator variables used by (5), including host range, parasite diversity, metric of parasite success, host species, parasite type, source of host genetic diversity, scale of host diversity, mode of host reproduction, whether the parasite induces host mortality and whether the study was performed in a laboratory environment. Note that parasite diversity was categorized as high if it was collected from a natural population for a lab study, part of an observational or

experimental field study, or if more than one genotype had been identified (but this

only applied to a small number of comparisons) or low if it was a laboratory strain of a

parasite or only one genotype had been identified (but again, this only applied to a

small number of comparisons). Where the information on these moderator variables

was not already available from the supplementary material of (5) and was not available

in the published article, we either performed a general search online, using Google, or

a more detailed search on the Web of Science database if this information was not

readily available.

5) Fifth, we calculated the mean, standard deviation and sample size for each

comparison of control and treatment groups from the data we extracted, as well as the

data calculated incorrectly in the previous analysis by (5). In total, this included 36

studies, 56 experiments and 130 comparisons.

For studies based on multiple populations with one or more continuous measures of

genetic diversity, (33,35,37–55), we chose the measure of genetic diversity which was

the best reflection of population-level genetic diversity (such as a measure based on a

Hardy-Weiberg equilibrium) and split the populations into two even groups of high

diversity and low diversity, before finally calculating a pooled mean, standard deviation

and sample size for each control and treatment group. Similarly, for studies with

multiple control groups (34,56), we calculated a pooled mean, standard deviation and

sample size. For studies which measured host survival, this was converted into host

mortality to better reflect parasite success.

After finishing all five steps of data collection, there was enough parasite success data

to calculate both the SMD and lnCVR for 211 non-independent comparisons of a group

of control versus treatment host populations.

Figure 1. How data was collected for each control versus treatment comparison of parasite success, which was later used in calculating effect sizes. n = number of studies / experiments / comparisons (the multi-level structure of the data make it appear that some sums are incorrect). Adapted from the preferred reporting items for systematic reviews and meta-analyses (PRIMSA) statement (57).

Calculation of effect sizes (SMD, lnCVR, lnRR, lnVR)

We calculated two main effect sizes (standardized mean difference [SMD] and log coefficient of variation ratio [lnCVR]) and two additional effects sizes (log response ratio [lnRR] and log variability ratio [lnVR]) to test the robustness of our results. Please note that all calculations of effect sizes and any subsequent calculations in this paper were performed in R v4.3.2 (58).

Main effect sizes - We chose SMD and lnCVR as our main effects sizes because they compare the difference in either the mean or variability of two groups whilst accounting for certain factors: (i) SMD measures the mean difference between two groups (high versus low diversity) in terms of standard deviations (59,60), so it can be used to compare different ways of measuring the same outcome (prevalence, load and virulence) (61), and uses a correction for small sample sizes, which is a common feature of ecological studies (62). (ii) lnCVR measures the ratio of variability between two groups adjusted for the size of the group means (63) and as a result, accounts for the possibility that the magnitude of the variability may scale with the mean, as is the case for many types of count data (such as parasite load) that follow a Poisson distribution.

Alternative effect sizes - We calculated lnRR and lnVR as alternatives to our main effect sizes, and although they did not account for all of the same factors, still measured the same difference in the mean and variability between two groups (63,64). Therefore, we were able to test the robustness of our results by comparing these two sets of effect sizes (65).

Before calculating our effect sizes, we added a small value (0.001) to the mean and standard deviation in parasite success for each pair of control and treatment groups to avoid any errors when taking logs of these data. For consistency, we calculated SMD and its sampling variance from the formula derived from the supplementary material of

(5), and calculated all variability effect sizes and their sampling variances using the

code from (63), as these were not used in the previous meta-analyses discussed

above (5,6), and instead are unique to our analysis. We calculated lnRR using the

escalc function from the metafor package v4.4.0 (66) because it was not clear how this

was calculated in (5). To account for comparisons based on shared controls, we

calculated the variance-covariance matrix for each effect size, using the

make_VCV_matrix function from the metaAidR package v0.0.0.9000 (67).

Publication bias

Before analyzing the data fully, we calculated the overall effect sizes for SMD and

lnCVR and tested for any potential publication bias using funnel plots and Egger's

regression (68).

Meta-analytic models were fitted to the data using the rma.mv function from the metafor

package v4.4.0 (66). We included fixed effects for each type of effect size, the

variance-covariance matrix of sampling errors, standard random effects for study and

host genus, and correlated random effects for comparisons taken from the same

experiment. Standard random effects for study and host genus were used to account

for the possibility of non-independence between experiments originating from the same

study and potential correlations between effects from closely related host species.

Similarly, correlated random effects were used to account for potential non-

independence of comparisons taken from the same experiment (multiple timepoints for

a single comparison of control and treatment groups, or effect sizes from the same

group of populations based on different measures of parasite success).

Funnel plots were used to identify whether published effect sizes were evenly

distributed around model means by examining how outcomes varied as a function of

their precision (standard error). This was achieved from a visual inspection of these

plots and statistical evaluation using Egger's regression.

Meta-analysis of overall data

To test whether there was a significant difference in mean parasite success (SMD) or

the variability in parasite success (lnCVR) between host populations with high versus

low genetic diversity, we fitted mixed effects meta-analytic models. All of the models

used in this paper were based on the same structure as those used for testing the

presence of publication bias.

Context dependence

Partial moderator analysis - To test if the overall effect of host population genetic

diversity on the mean and variability in parasite success depended on an interaction

between host range and parasite genetic diversity, we introduced an interaction term

for these two moderators in our original meta-analytic models. Therefore, we could

compare:

1) High versus low single-host parasite population genetic diversity.

2) High versus low multi-host parasite population genetic diversity.

We compared the significance level of each individual predictor within the model, as

well as the contrasts between them using the *glht* function from the *multcomp* package

v1.4.25 (69).

Full moderator analysis – To test our additional hypotheses (Table 2) for the eight

remaining moderator variables, we modelled each moderator separately with its own

individual mixed effects model. Before running the models, we removed redundant

moderator categories with a limited sample size, such as transmission or infection rate

(versus prevalence) and disease severity (versus load) for the metric of parasite

success, and prokaryotic (versus vertebrate or invertebrate) for host species.

We compared the significance level of each individual predictor within the model, as

well as the contrasts between them using ANOVA with a correction for multiple

comparisons (Holm's method).

Table 2. Hypotheses for additional moderators.

Moderator	Hypothesis
Metric of parasite success	Our study of 'parasite success' combined data of several types (eg prevalence, virulence, infection load). Using this moderator, we tested if the effects of host population genetic diversity differed between these different metrics.
Host species	The effect of host population genetic diversity may be influenced by the specificity of genetic interactions between host and parasite. These genetic interactions are thought to be more specific in invertebrates than in vertebrates (70), therefore we tested for inconsistency of the host population genetic diversity effect in these two groups.
Parasite type	Since microparasite infections tend to be short-lived, whereas macroparasites are usually long-lasting, it is generally believed that they can withstand the host immune response (71). This suggests that they might not be as tightly coevolved, and have a lower genetic specificity for infection, than microparasites. Therefore, we tested for inconsistency of the host population genetic diversity effect in these two groups.
Source of host genetic diversity	Studies typically investigate the impact of host genetic diversity by either (i) inbreeding lineages to create a comparison between inbred and outbred populations, (ii) selecting a suite of wildtype genotypes for controlled experiments with either low genetic diversity or high genetic diversity, or (iii) sampling organisms from the wild from populations that have been characterised as having different levels of genetic diversity. We used this moderator to test if these different sources of genetic diversity affected the influence of host population genetic diversity.
Scale of host diversity	Host populations were predetermined as having either high or low diversity (discrete) or we separated them into such categories as part of our data collection (Fig. 1, step 5) because they used multiple populations with a continuous measure of diversity. We used this moderator to test if this feature of how studies were designed had an effect on the influence of host population genetic diversity.
Mode of host reproduction	Host species reproduced sexually, asexually or using a combination of the two (i.e. facultatively sexual, such as *Daphnia*). We used this moderator to test if these different modes of host reproduction affected the influence of host population genetic diversity.
Host mortality?	The range of parasites studied can be further categorized by how virulent they are. Therefore, we used this moderator as a proxy for how virulent the parasites were, in terms of whether or not they kill the host, to test if

	differences in parasite virulence affected the influence of host population genetic diversity.
Laboratory?	We used this moderator to test if the difference in study setting (laboratory versus field) affected the influence of host population genetic diversity.

Sensitivity analysis of overall effects

To test the robustness of our results for the combined (overall) dataset, we performed

a series of 'leave-one-out' sensitivity analyses. This involved the iterative exclusion of

either one independent comparison (i.e. treatments with shared controls were

considered grouped together into a single comparison) or study at a time.

RESULTS

Absence of publication bias

Our dataset contained 211 estimates of the effect that changes in host population

genetic diversity have on parasite success; we assessed this effect on both mean

parasite success (SMD) and the variability in parasite success (lnCVR). Visual

inspection of the funnel plots for the effect of host population diversity on mean parasite

success (Fig. 2A) and its effect on the variability in parasite success (Fig. 2B), showed

no evidence for publication bias. More stringent evaluation showed that there was no

correlation between the size of the effects themselves and their standard error (Egger's

test for both SMD and lnlnCVR: $R = 0.06$, 95% CI [-0.24, 0.37], $p = 0.67$ and $R = -0.03$,

95% CI [-0.38, 0.32], $p = 0.86$ respectively).

Figure 2. Testing for publication bias: the distribution of published effect sizes for our meta-analysis as a function of their precision (standard error). The x-axis in both plots shows effects of an increase in host population genetic diversity (high vs low) for A) the mean difference in parasite success (SMD) and B) the ratio of variation in parasite success (lnCVR). Model means and their 95% confidence intervals are shown by the dashed black lines.

331
Evenly distributed animal host and parasite taxa

Our dataset included a diverse range of hosts and parasites, including 31 unique host
genera, 60 unique parasite genera and 71 (92) unique host-parasite genera (or
species) combinations.

The distribution of unique host taxa was biased towards animals (invertebrate and
vertebrate genera / species), with only two unique non-animal (prokaryotic) host
species (Table 3). However, there was an even distribution of the unique parasite taxa
across the combination of all unique host taxa.

Table 3. The number of unique host and parasite combinations and how evenly they
are distributed across different taxonomic groups. The number of unique combinations

of host and parasite genera and species is shown by the first two numbers separated

by a backslash (genera / species) and the number of studies they correspond to in

parentheses. The total number of studies is higher than 48 (the total number of studies

in our dataset), because there were some studies with multiple comparisons of unique

host and parasite combinations. The colour coding is based on the number of unique

combinations of host and parasite genera.

		Host taxon			
		Prokaryote	Invertebrate	Vertebrate	Total
Parasite taxon	Animal	0 / 0 (0)	12 / 14 (6)	10 / 13 (8)	22 / 27 (14)
	Bacteria	0 / 0 (0)	2 / 2 (2)	13 / 16 (5)	15 / 18 (7)
	Fungi	1 / 1 (1)	15 / 18 (16)	1 / 1 (1)	17 / 20 (18)
	Protozoa	0 / 0 (0)	7 / 12 (11)	3 / 3 (2)	10 / 15 (13)
	Virus	1 / 1 (1)	3 / 7 (2)	4 / 4 (4)	7 / 12 (7)
	Total	1 / 2 (2)	39 / 53 (37)	31 / 37 (20)	71 / 92 (59)

348
349
Overall effect of host population genetic diversity on mean parasite success is negative

There was a significant effect of host population genetic diversity on mean parasite

success (SMD = -0.29, 95% CI = [-0.57, -0.02], n = 211; Fig. 3A), which shows that

there is a negative relationship between host population genetic diversity and the mean

parasite success. However, the effect of host population genetic diversity on the

variability of parasite success was not significant (lnCVR = 0.02, 95% CI = [-0.30, 0.35],

n = 211; Fig. 3B), which suggests that there is no relationship between host population

genetic diversity and the variability in parasite success.

In these analyses the residual variation (heterogeneity) in the data for both the

difference in the mean and variability of parasite success was high (I^2 = 84.0% & 82.0%

respectively). Most of this variation was explained by the effect of study (84.0% &

80.7%) and only a small amount was explained by host genus (0.0% & and 3.3%).

Figure 3. Overall effect of host population genetic diversity on the mean and variability in parasite success. The x-axis in each plot shows the effect of increasing host population genetic diversity on either A) the difference in mean parasite success (SMD) or B) the difference in the variability in parasite success (lnCVR). The dashed line indicates an effect size of zero where there is no influence of host population genetic diversity on parasite success. Model means are shown with 95% confidence intervals (thick black lines) and prediction intervals (thin black lines). Individual effect sizes (circles) are scaled according to the inverse of their standard error. n = sample size of the data (the number of effect sizes : the number of studies). Model means that are significantly different from zero are indicated by an asterisk ($p < 0.05$). Forest plot alternatives are shown in the online supplementary material (Fig. S1).

Variability in the success of single-host parasites depends on the combined host and

parasite population genetic diversity

The overall conclusion that higher host population genetic diversity reduces mean

parasite success, but does not the affect the variability in parasite success, is relatively

simplified. In fact, there are significant effects of host population genetic diversity on

both the mean and variability in parasite success which depend on a combination of

parasite population genetic diversity and host range.

In contrast to the overall effect of host population genetic diversity on mean parasite

success, which was significantly negative (SMD = -0.29, 95% CI = [-0.57, -0.02], n =

211; Fig. 3A), separating the effects of host population genetic diversity by a

combination of parasite population genetic diversity and host range showed that the

effect of host population genetic diversity on mean parasite success was only

significant for single versus multi-host parasites.

In addition, although there was no overall effect of host population genetic diversity on

the variability in parasite success (lnCVR = 0.02, 95% CI = [-0.30, 0.35], n = 211; Fig.

3B), there was a significant difference in the effect of host population genetic diversity

on the variability in the success of singe-host parasites with low versus high population

genetic diversity (glht: p = 0.03; Fig. 4B). Specifically, increased host population

genetic diversity lead to either an increase or decrease in the variability of the success

of single-host parasites when their own population genetic diversity was either high or

low.

Figure 4. The influence of host range and parasite population genetic diversity on the effect of host population genetic diversity on the mean and variability in parasite success. The x-axis in each plot shows the effect of increasing host population genetic diversity on either A) the difference in mean parasite success (SMD) or B) the difference in the variability in parasite success (lnCVR). The dashed line indicates an effect size of zero where there is no influence of host population genetic diversity on parasite success. Model means are shown with 95% confidence intervals (thick black lines) and prediction intervals (thin black lines). Individual effect sizes (circles) are scaled according to the inverse of their standard error. n = sample size of the data (the number of effect sizes : the number of studies). The significance level of individual model means, as well as any pairwise contrasts, is indicated by one (p < 0.05) or three (p < 0.001) asterisks.

Context-dependent effect of host population genetic diversity

To evaluate our table of hypotheses (Table 2), we investigated how eight other aspects

of study design influenced the effect of host population genetic diversity on the mean

and variability in parasite success (Fig. 5)

We found that the effect of host population genetic diversity was significantly negative

on the mean success of microparasites (Fig. 5C), for inbred versus outbred hosts (Fig.

5D), for sexually reproducing hosts (SMD = -0.40, 95% CI = [-0.73, -0.07], p = 0.02,

Fig. 5F), parasites which caused host mortality (SMD = -0.34, 95% CI = [-0.68, -0.00],

p = 0.05, Fig. 5G) and non-lab based studies (SMD = -0.33, 95% CI = [-0.65, -0.01], p

= 0.04, Fig. 5H). For the effect of host population genetic diversity on the variability in

parasite success, we found that this was significantly negative for asexually

reproducing hosts (Fig. 5N).

In addition, comparisons between specific levels of these moderators showed a highly

significant difference in the mean difference in parasite success between micro- and

macroparasites (QM = 13.2, df = 1, p < 0.001, Fig. 5C) and also a significant difference

in the ratio of variability in parasite success between both sexual hosts and either

asexual (QM = 6.40, df = 1, p = 0.01, Fig. 5N) or facultatively sexual hosts (QM = 5.53

df = 1, p = 0.02, Fig. 5N).

Figure 5. The context-dependence of the effect of host population genetic diversity on the mean and variability in parasite success. The x-axis in each plot shows the effect of increasing host population genetic diversity on either A) the difference in mean parasite success (SMD) or B) the difference in the variability in parasite success (lnCVR). Model means are shown with 95% confidence intervals and are scaled according to the number of experiments. The dashed line indicates an effect size of zero. The significance level of individual model means, as well as any pairwise contrasts, is indicated by one (p < 0.05), two (p < 0.01) or three (p < 0.001) asterisks. The following abbreviations are used; Prev. (Prevalence), Vir. (Virulence), Vert. (Vertebrate), Invert.

(Invertebrate), Macro. (Macroparasite), Micro. (Microparasite), Source gen. var. (Source of host genetic diversity), Art. (Artificial), Nat. (Natural), Scale host div. (Scale of host diversity), Cont. (Continuous), Disc. (Discrete), Host mode repro. (Mode of host reproduction), Sex. (Sexual), Asex. (Asexual), Mixd (Mixed).

408
Our results are robust to leaving data out, but require the right 'mean' effect size

To test the robustness our of results, we performed a suite of 'leave-one-out' sensitivity

analyses and remodeled our parasite success data with an alternative set of effect

sizes.

The suite of sensitivity analyses showed that the results of our main effects were not

dependent on the inclusion of a particular study (Table 4., Fig. S2) or independent

comparison in our dataset (Table 4., Fig. S3). However, they were less robust to using

the log response ratio (lnRR) as an alternative effect size to the standardized mean

difference (SMD) to measure to effect of host population genetic diversity on mean

parasite success. Although both measures showed a negative effect of host population

genetic diversity on mean parasite success, the alternate way of measuring this was

not significant (lnRR = 0.93, 95% CI = [-0.40, 2.25], n = 211). In comparison, the

alternate variability measure, the log variability ratio (lnVR), supported the result of the

main effect size (the log coefficient of variation ratio, lnCVR) by showing that there was

no significant effect of host population genetic diversity on the variability in parasite

success.

Table 4. Results of the leave-one-out sensitivity analyses. To test the robustness of

the results using our main effect sizes, we re-modelled the data using an iterative

exclusion of either one study (Leave1studyout) or one independent comparison

(Leave1trtout) and calculated the mean model estimate and the mean p-value across

all the models. The following abbreviations are used; ES (effect size), SE (mean

standard error across all models), ci.lb and ci.ub (mean lower and upper bounds of

95% confidence intervals across all models respectively).

Method	ES	Estimate	SE	z-value	p-value	ci.lb	ci.ub
Leave1trtout	SMD	-0.29	0.14	-2.07	0.04	-0.56	-0.02
Leave1trtout	lnCVR	0.02	0.17	0.13	0.9	-0.3	0.35
Leave1studyout	SMD	-0.29	0.14	-2.05	0.04	-0.57	-0.01
Leave1studyout	lnCVR	0.02	0.17	0.13	0.87	-0.31	0.35

DISCUSSION

Building on the recent studies by (5,6), we show that the conventional theory, which

suggests that there is a negative effect of host population genetic diversity on mean

parasite success, is not always correct (1). Although there is a benefit to high host

population genetic diversity, which protects against mean parasite success, our

moderator analysis shows that it strongly depends on the nature of the host-parasite

association, such that the parasite success of host specialists, but not generalists, is

affected. We propose that this is driven by the amount of genetic specificity for infection

because specialist parasites could be more likely to have highly specific, matching-

allele-type interactions between host resistance and parasite infectivity alleles.

The benefit of host population genetic diversity in protecting against the mean success

of single-host, but not multi-host, parasites is an important distinction. It highlights the

susceptibility of vulnerable host populations (small, fragmented populations with low

genetic diversity) to infection by host-specific parasites, rather than infection by multi-

host parasites. This represents a change of perspective that could help to protect

vulnerable host populations by prioritizing how genetic diversity within these

populations is managed (72) or even by finding a way to alter the genetic specificity of

host and parasite associations to make them less specific, such as by introducing

alternative hosts for the parasite.

We also found that the effect of host population genetic diversity on parasite success

was highly context dependent. Using our partial moderator analysis, in which we

modelled the interaction between parasite population genetic diversity and host range,

we showed the importance of our proposed diversity-uncertainty model (Table 1).

Specifically, increased host population genetic diversity lead to either an increase or

decrease in the variability of the success of single-host parasites when their own

population genetic diversity was either high or low.

This has big implications for identifying disease threats, such as epidemics, to host

populations. Our results show that the combination of low host and low parasite

population genetic diversity can sometimes lead to an extremely high level of parasite

success. However, the most vulnerable host populations are those with a combination

of low genetic diversity themselves and high parasite population genetic diversity

because there is a consistently high level of parasite success. In addition, the

protective effect of host population genetic diversity diminishes with increasing parasite

population genetic diversity, so we might want to prioritize the genetic diversity of low

host-high parasite genetic diversity associations.

Similarly, there are also consequences for the repeatability of disease experiments and

the predictability of recurrent disease outbreaks. We might expect that combinations

of 1) low host and parasite population genetic diversity and 2) high host and parasite

genetic diversity have the biggest implications for study repeatability because lab

experiments are likely to be low x low and field experiments are likely to be high x high.

Due to their high levels of variability, these systems will also have similar issues in

terms of predictability of future disease occurrence and emergence.

In conjunction with our partial moderators, our full moderator analysis highlighted

several other significant influences on the effect of host population genetic diversity on

parasite success, including parasite type, the source of genetic diversity, the mode of

host reproduction, whether or not host mortality is affected and whether or not the study

was performed in a laboratory. In particular, the effect of host population genetic

diversity on the mean success of microparasites was much stronger (more negative)

than macroparasites. This confirmed our original hypothesis, which suggested that

macroparasites have a lower genetic specificity for infection due the longer-lasting

nature of their infections and thus being less tightly coevolved with their hosts.

One of the other key moderators was the source of host population genetic diversity.

This showed that there was a significant reduction in mean parasite success between

inbred versus outbred hosts, but not between other host comparisons which had been

established by selecting suites of unique genotypes or sampling organisms from wild

populations. This result makes sense and may have occurred due to combination of

an inbreeding effect, where inbred hosts are more susceptible to disease than outbred

hosts (24), and population-level genetic diversity effects, which was not necessarily

the case for the other categories of how host population genetic diversity was

assembled.

Despite using the same set of moderators, our results were inconsistent with previous

meta-analytic studies (5,6). They modelled each moderator separately (6) or using all

possible combinations of different moderators and calculated model averages (5) and

either found that most of their moderators had no significant influence, or a limited

influence, on the effect of host population genetic diversity on mean parasite success.

Specific differences between our moderator analysis to theirs include the use of a

larger number of effect sizes, which meant we did not suffer a limited sample size in

certain moderator categories as they did (e.g. single-host parasites and parasites with

one or more strains), and the interaction between host range and parasite population

genetic diversity, which was not included in one of these studies (6).

However, we temper the exciting nature of results with some considerations:

1) There is a large number of effect sizes (68) for single-host parasites with low

population genetic diversity, but most of these actually come from a prokaryotic

506   bacterial host study (73), rather than a vertebrate or invertebrate host, which is

507    the case for most of our data.

508  2) Host range may not be a reliable estimate of the genetic specificity for infection.

509    This is because we based this on the prediction that highly specific interactions

510    between host and parasite genotypes would be more likely for tightly

511    coevolving pathogens (MAM), as we might expect between a single host and

512    parasite species, compared to multiple hosts and a single parasite species with

513    a broad host range.

514  3) The results of our moderator analysis rely on particular ways of labelling the

515    data. For example, comparing mainly natural versus laboratory strains of

516    parasites could be a poor indication of the effect of parasite population genetic

517    diversity because the exact level of diversity was not actually quantified.

518    Similarly, our measure of host range is somewhat subjective and based on an

519    incomplete literature.

520  4) The suite of leave-one-out analyses showed that our overall effects, which did

521    not include any moderators, were robust to the potential influence of extreme

522    values from particular studies or data points, but the overall effect of host

523    population genetic diversity on mean parasite success effect size relies on our

524    use of the standardized mean difference versus the log response ratio (refer to

525    the methods for why this was selected as an alternative effect size).

526 Our final thought is that most of our data concentrates on the effect of host population

527 genetic diversity on both the mean and variability in parasite success for spatially

528 replicated groups of host populations (except for one study (74)). We expect that the

529 diversity-uncertainty model may also apply to the effect of host population genetic

530 diversity on both the mean and variability in parasite success across time, but with

531 some key differences. For example, repeated bouts of directional selection from

532 parasite outbreaks can reduce host and parasite population genetic diversity through

time (75,76), which could increase the mean and reduce the variability in parasite

success. However, the precise nature of selection and its resulting impact on host and

parasite population genetic diversity depends on the underlying host-parasite infection

genetics (i.e. MAM versus GFG (77)).

SUMMARY

In this study, we measured the difference in the mean and variability in parasite

success between host populations with high versus low genetic diversity. First, we

challenge so-called 'conventional wisdom' (*sensu* (1)) and propose a diversity-

uncertainty model to better understand the context in which the effect of host

population genetic diversity really matters for both the mean and variability in parasite

success. We find that host population genetic diversity affects the mean success of

host-specific parasites, but not host generalists, which could be a result of greater

genetic specificity for infection. We then demonstrate that our diversity-uncertainty

model is appropriate for our collection of studies, such that the effect of host population

diversity on the variability in parasite success depends on a combination of parasite

diversity and host range. Finally, we find that there are a number of other context

dependent effects of host population genetic diversity on both the mean and variability

in parasite success, such as parasite type. Overall, these findings represent a change

of perspective that could help to protect vulnerable host populations by prioritizing how

genetic diversity within these populations is managed. Future study of the diversity-

uncertainty hypothesis across a range of plant host-parasite systems would help

generalize these findings.

REFERENCES

1.   King KC, Lively CM. Does genetic diversity limit disease spread in natural host
populations? Heredity (Edinb) [Internet]. 2012 Oct 20 [cited 2018 Oct
11];109(4):199–203. Available from: http://www.nature.com/articles/hdy201233

2.   Hamilton WD. Kinship, recognition, disease, and intelligence: constraints of
social evolution. Tokyo: Japan Scientific Societies Press; 1987. 81–102 p.

3.   Sherman PW, Seeley TD, Reeve HK. Parasites, Pathogens, and Polyandry in
Social Hymenoptera [Internet]. Vol. 131, Source: The American Naturalist.
1988 [cited 2019 Jan 23]. Available from:
https://www.jstor.org/stable/pdf/2461747.pdf?refreqid=excelsior%3A10d16706
d07e226edd1f3ace6f23a6b4

4.   O'Brien SJ, Evermann JF. Interactive influence of infectious disease and
genetic diversity in natural populations. Vol. 3, Trends in Ecology and
Evolution. 1988.

5.   Gibson AK, Nguyen AE. Does genetic diversity protect host populations from
parasites? A meta-analysis across natural and agricultural systems. Evol Lett.
2020;5(1).

6.   Ekroth AKE, Rafaluk-Mohr C, King KC. Host genetic diversity limits parasite
success beyond agricultural systems: a meta-analysis. Proceedings of the
Royal Society B: Biological Sciences. 2019 Sep 25;286(1911):20191811.

7.   Alan Pounds J, Bustamante MR, Coloma LA, Consuegra JA, Fogden MPL,
Foster PN, et al. Widespread amphibian extinctions from epidemic disease
driven by global warming. Nature [Internet]. 2006 Jan [cited 2019 Sep
13];439(7073):161–7. Available from:
http://www.nature.com/articles/nature04246

8.   De Castro F, Bolker B. Mechanisms of disease-induced extinction. Ecol Lett
[Internet]. 2004 Dec 3 [cited 2019 Sep 13];8(1):117–26. Available from:
http://doi.wiley.com/10.1111/j.1461-0248.2004.00693.x

9.   Gibson AK. Genetic diversity and disease: The past, present, and future of an
old idea. Evolution (N Y) [Internet]. 2022 Feb 19;76(S1):20–36. Available from:
https://academic.oup.com/evolut/article/76/s1/20/6728439

10.   Altizer S, Dobson A, Hosseini P, Hudson P, Pascual M, Rohani P. Seasonality
and the dynamics of infectious diseases. Ecol Lett [Internet]. 2006 Mar 31
[cited 2019 Jun 29];9(4):467–84. Available from:
http://doi.wiley.com/10.1111/j.1461-0248.2005.00879.x

11.   Keesing F, Ostfeld RS. Dilution effects in disease ecology. Ecol Lett.
2021;24(11).

12.   Ostfeld RS, Keesing F. Effects of Host Diversity on Infectious Disease. Annu
Rev Ecol Evol Syst [Internet]. 2012 Dec 5 [cited 2019 Jan 21];43(1):157–82.
Available from: http://www.annualreviews.org/doi/10.1146/annurev-ecolsys-
102710-145022

13.   Johnson PTJ, Ostfeld RS, Keesing F. Frontiers in research on biodiversity and
disease. Ecol Lett. 2015;18(10):1119–33.

14.   Keesing F, Holt RD, Ostfeld RS. Effects of species diversity on disease risk.
Ecol Lett [Internet]. 2006 Mar 31 [cited 2019 Jan 10];9(4):485–98. Available
from: http://doi.wiley.com/10.1111/j.1461-0248.2006.00885.x

15.   Keesing F, Belden LK, Daszak P, Dobson A, Harvell CD, Holt RD, et al.
Impacts of biodiversity on the emergence and transmission of infectious
diseases. Nature. 2010.

16.   Altermatt F, Ebert D. Genetic diversity of Daphnia magna populations
enhances resistance to parasites. Ecol Lett. 2008;

17.   Mihaljevic JR, Joseph MB, Orlofske SA, Paull SH. The Scaling of Host Density
with Richness Affects the Direction, Shape, and Detectability of Diversity-
Disease Relationships. Fernandez-Reyes D, editor. PLoS One [Internet]. 2014
May 21;9(5):e97812. Available from:
https://dx.plos.org/10.1371/journal.pone.0097812

18.   Johnson PTJ, Preston DL, Hoverman JT, Henderson JS, Paull SH, Richgels
KLD, et al. Species diversity reduces parasite infection through cross-
generational effects on host abundance. Ecology [Internet]. 2012
Jan;93(1):56–64. Available from:
https://esajournals.onlinelibrary.wiley.com/doi/10.1890/11-0636.1

19.   Elton CS. The Ecology of Invasions by Animals and Plants. The Ecology of
Invasions by Animals and Plants. 1958.

20.   van der Plank JE. Plant diseases: epidemics and control. Academic Press;
1963.

21.   Browning JA, Frey KJ. Multiline Cultivars as a Means of Disease Control. Annu
Rev Phytopathol [Internet]. 1969 Sep;7(1):355–82. Available from:
https://www.annualreviews.org/doi/10.1146/annurev.py.07.090169.002035

22.   Leonard KJ. Factors affecting rates of stem rust increase in mixed plantings of
susceptible and resistant oat varieties cultivars. Phytopathology.
1969;59:1845–50.

23.   Garrett KA, Mundt CC. Epidemiology in Mixed Host Populations.
Phytopathology [Internet]. 1999;89(11):984–90. Available from:
http://apsjournals.apsnet.org/doi/10.1094/PHYTO.1999.89.11.984

24.   Coltman DW, Pilkington JG, Smith JA, Pemberton JM. PARASITE-MEDIATED
SELECTION AGAINST INBRED SOAY SHEEP IN A FREE-LIVING, ISLAND
POPULATION [Internet]. Vol. 53, Evolution. 1999 [cited 2019 Feb 12].
Available from: https://about.jstor.org/terms

25.   Anderson RM, May RM, Joysey K, Mollison D, Conway GR, Cartwell R, et al.
The invasion, persistence and spread of infectious diseases within animal and
plant communities. Philosophical Transactions of the Royal Society of London
B, Biological Sciences [Internet]. 1986 Dec 15 [cited 2019 Jan
9];314(1167):533–70. Available from:
http://rstb.royalsocietypublishing.org/cgi/doi/10.1098/rstb.1986.0072

26.   Schmid-Hempel P, Ebert D. On the evolutionary ecology of specific immune
defence. Trends Ecol Evol [Internet]. 2003 Jan;18(1):27–32. Available from:
https://linkinghub.elsevier.com/retrieve/pii/S0169534702000137

27.   Agrawal A, Lively CM. Infection genetics: Gene-for-gene versus matching-
alleles models and all points in between. Evol Ecol Res. 2002;

28. Boomsma JJ. Paternity in eusocial Hymenoptera. Philos Trans R Soc Lond B
Biol Sci [Internet]. 1996 Jul 29;351(1342):947–75. Available from:
https://royalsocietypublishing.org/doi/10.1098/rstb.1996.0087

29. Van Baalen M, Beekman M. The costs and benefits of genetic heterogeneity in
resistance against parasites in social insects. American Naturalist. 2006.

30. Ganz HH, Ebert D. Benefits of host genetic diversity for resistance to infection
depend on parasite diversity [Internet]. Vol. 91, Source: Ecology. 2010 [cited
2018 Oct 12]. Available from: https://www-jstor-
org.ezproxy.stir.ac.uk/stable/pdf/25661177.pdf?refreqid=excelsior%3A5c0f30d
81cb9568626bbc1823fa625af

31. Bensch HM, O'Connor EA, Cornwallis CK. Living with relatives offsets the
harm caused by pathogens in natural populations. Elife [Internet]. 2021 Jul
26;10. Available from: https://elifesciences.org/articles/66649

32. Koskella B, Lively CM. Evidence for negative frequency-dependent selection
during experimental coevolution of a freshwater snail and a sterilizing
trematode. Evolution (N Y). 2009;63(9):2213–21.

33. King KC, Jokela J, Lively CM. Parasites, sex, and clonal diversity in natural
snail populations. Evolution (N Y). 2011;65(5).

34. Agha R, Gross A, Rohrlack T, Wolinska J. Adaptation of a chytrid parasite to
its cyanobacterial host is hampered by host intraspecific diversity. Front
Microbiol. 2018;9(MAY):1–10.

35. Giese AR, Hedrick PW. Genetic variation and resistance to a bacterial
infection in the endangered Gila topminnow. Anim Conserv. 2003;6(4).

36. Baer B, Schmid-Hempel P. Unexpected consequences of polyandry for
parasitism and fitness in the bumblebee, Bombus terrestris. Evolution (N Y).
2001;

37. Puurtinen M, Hytönen M, Knott KE, Taskinen J, Nissinen K, Kaitala V. The
effects of mating system and genetic variability on susceptibility to trematode
parasites in a freshwater snail, Lymnaea stagnalis. Evolution (N Y).
2004;58(12).

38. Dagan Y, Liljeroos K, Jokela J, Ben-Ami F. Clonal diversity driven by
parasitism in a freshwater snail. J Evol Biol. 2013;26(11).

39. Dionne M, Miller KM, Dodson JJ, Bernatchez L. MHC standing genetic
variation and pathogen resistance in wild Atlantic salmon. Philosophical
Transactions of the Royal Society B: Biological Sciences. 2009;364(1523).

40. Ellison A, Cable J, Consuegra S. BEST OF BOTH WORLDS? ASSOCIATION
BETWEEN OUTCROSSING AND PARASITE LOADS IN A SELFING FISH.
Evolution (N Y) [Internet]. 2011 Oct 1 [cited 2019 Mar 14];65(10):3021–6.
Available from: http://doi.wiley.com/10.1111/j.1558-5646.2011.01354.x

41. Field SG, Lange M, Schulenburg H, Velavan TP, Michiels NK. Genetic
diversity and parasite defense in a fragmented urban metapopulation of
earthworms. Anim Conserv. 2007;10(2).

42. Kyle CJ, Rico Y, Castillo S, Srithayakumar V, Cullingham CI, White BN, et al.
Spatial patterns of neutral and functional genetic variations reveal patterns of
local adaptation in raccoon (Procyon lotor) populations exposed to raccoon
rabies. Mol Ecol. 2014;23(9).

43. Loiseau C, Zoorob R, Robert A, Chastel O, Julliard R, Sorci G. Plasmodium
relictum infection and MHC diversity in the house sparrow (Passer
domesticus). Proceedings of the Royal Society B: Biological Sciences.
2011;278(1709).

44. Meagher S. Genetic diversity and capillaria hepatica (nematoda) prevelance in
Michigan deer mouse populations. Evolution (N Y). 1999;53(4):1318–24.

45. Neumann P, Moritz RFA. Testing genetic variance hypotheses for the
evolution of polyandry in the honeybee (Apis mellifera L.). Insectes Soc.
2000;47(3).

46. Parsche S, Lattorff MHG. The relative contributions of host density and genetic
diversity on prevalence of a multi-host parasite in bumblebees. Biological
Journal of the Linnean Society. 2018;125(4).

47. Pierce AA, De Roode JC, Altizer S, Bartel RA. Extreme heterogeneity in
parasitism despite low population genetic structure among monarch butterflies
inhabiting the Hawaiian Islands. PLoS One. 2014;9(6).

48. Queirós J, Vicente J, Alves PC, de la Fuente J, Gortazar C. Tuberculosis,
genetic diversity and fitness in the red deer, Cervus elaphus. Infection,
Genetics and Evolution. 2016;43.

49. Rahn AK, Krassmann J, Tsobanidis K, MacColl ADC, Bakker TCM. Strong
neutral genetic differentiation in a host, but not in its parasite. Infection,
Genetics and Evolution. 2016;44.

50. Savage AE, Becker CG, Zamudio KR. Linking genetic and environmental
factors in amphibian disease risk. Evol Appl. 2015;8(6).

51. Trouvé S, Degen L, Renaud F, Goudet J. EVOLUTIONARY IMPLICATIONS
OF A HIGH SELFING RATE IN THE FRESHWATER SNAIL LYMNAEA
TRUNCATULA. Evolution (N Y). 2003;57(10).

52. Velavan TP, Weller S, Schulenburg H, Michiels NK. High genetic diversity and
heterogeneous parasite load in the earthworm Lumbricus terrestris on a
German meadow. Soil Biol Biochem. 2009;41(8).

53. Whitehorn PR, Tinsley MC, Brown MJF, Darvill B, Goulson D. Genetic
diversity and parasite prevalence in two species of bumblebee. J Insect
Conserv. 2014;18(4).

54. Whitehorn PR, Tinsley MC, Brown MJF, Darvill B, Goulson D. Genetic
diversity, parasite prevalence and immunity in wild bumblebees. Proceedings
of the Royal Society B: Biological Sciences [Internet]. 2011 Apr
22;278(1709):1195–202. Available from:
https://royalsocietypublishing.org/doi/10.1098/rspb.2010.1550

55. Whiteman NK, Matson KD, Bollmer JL, Parker PG. Disease ecology in the
Galápagos Hawk (Buteo galapagoensis): host genetic diversity, parasite load

and natural antibodies. Proceedings of the Royal Society B: Biological
Sciences [Internet]. 2006 Apr 7;273(1588):797–804. Available from:
https://royalsocietypublishing.org/doi/10.1098/rspb.2005.3396

56.   Schmidt AM, Linksvayer TA, Boomsma JJ, Pedersen JS. No benefit in
diversity? The effect of genetic variation on survival and disease resistance in
a polygynous social insect. Ecol Entomol. 2011;36(6).

57.   Page MJ, McKenzie JE, Bossuyt PM, Boutron I, Hoffmann TC, Mulrow CD, et
al. The PRISMA 2020 statement: an updated guideline for reporting systematic
reviews. BMJ [Internet]. 2021 Mar 29;n71. Available from:
https://www.bmj.com/lookup/doi/10.1136/bmj.n71

58.   R Core Team. _R: A Language and Environment for Statistical Computing_
[Internet]. Vienna, Austria: R Foundation for Statistical Computing; 2023.
Available from: https://www.r-project.org/

59.   Borenstein M, Hedges L V., Higgins JPT, Rothstein HR. Introduction to meta-
analysis. Introduction to Meta-Analysis. 2009.

60.   Field AP, Gillett R. How to do a meta-analysis. British Journal of Mathematical
and Statistical Psychology. 2010;63(3).

61.   Higgins JPT, Green S, editors. Cochrane Handbook for Systematic Reviews of
Interventions [Internet]. Version 5. The Cochrane Collaboration, 2011;
Available from: www.handbook.cochrane.org

62.   Jennions MD. A survey of the statistical power of research in behavioral
ecology and animal behavior. Behavioral Ecology [Internet]. 2003 May
1;14(3):438–45. Available from: https://academic.oup.com/beheco/article-
lookup/doi/10.1093/beheco/14.3.438

63.   Nakagawa S, Poulin R, Mengersen K, Reinhold K, Engqvist L, Lagisz M, et al.
Meta-analysis of variation: ecological and evolutionary applications and
beyond. O'Hara RB, editor. Methods Ecol Evol [Internet]. 2015 Feb
17;6(2):143–52. Available from:
https://onlinelibrary.wiley.com/doi/10.1111/2041-210X.12309

64.   Nakagawa S, Noble DWA, Lagisz M, Spake R, Viechtbauer W, Senior AM. A
robust and readily implementable method for the meta-analysis of response
ratios with and without missing standard deviations. Ecol Lett [Internet]. 2023
Feb 26;26(2):232–44. Available from:
https://onlinelibrary.wiley.com/doi/10.1111/ele.14144

65.   Koricheva J, Gurevitch J. Uses and misuses of meta-analysis in plant ecology.
Gómez-Aparicio L, editor. Journal of Ecology [Internet]. 2014 Jul
23;102(4):828–44. Available from:
https://besjournals.onlinelibrary.wiley.com/doi/10.1111/1365-2745.12224

66.   Viechtbauer W. Conducting Meta-Analyses in R with the metafor Package. J
Stat Softw [Internet]. 2010;36(3). Available from:
http://www.jstatsoft.org/v36/i03/

67.   Lagisz M, Senior A, Nakagawa S, Noble D. _metaAidR: Functions and Data
for Aiding Meta-analyses in Ecology and Evolution_. 2024.

68. Sutton AJ. Publication bias. In: The handbook of research synthesis and meta-
analysis. 2nd ed. Russell Sage Foundation; 2009. p. 435–452.

69. Hothorn T, Bretz F, Westfall P. Simultaneous Inference in General Parametric
Models. Biometrical Journal [Internet]. 2008 Jun 6;50(3):346–63. Available
from: https://onlinelibrary.wiley.com/doi/10.1002/bimj.200810425

70. Dybdahl MF, Jenkins CE, Nuismer SL. Identifying the Molecular Basis of Host-
Parasite Coevolution: Merging Models and Mechanisms. Am Nat [Internet].
2014 Jul;184(1):1–13. Available from:
https://www.journals.uchicago.edu/doi/10.1086/676591

71. Sorci G. Immunology of Parasitism. In: Reference Module in Biomedical
Sciences [Internet]. Elsevier; 2014. Available from:
https://linkinghub.elsevier.com/retrieve/pii/B9780128012383002531

72. Meuwissen THE, Sonesson AK, Gebregiwergis G, Woolliams JA.
Management of Genetic Diversity in the Era of Genomics. Front Genet
[Internet]. 2020 Aug 13;11. Available from:
https://www.frontiersin.org/article/10.3389/fgene.2020.00880/full

73. Van Houte S, Ekroth AKE, Broniewski JM, Chabas H, Ashby B, Bondy-
Denomy J, et al. The diversity-generating benefits of a prokaryotic adaptive
immune system. Nature. 2016;532(7599).

74. Hale KA, Briskie J V. Decreased immunocompetence in a severely
bottlenecked population of an endemic New Zealand bird. Anim Conserv.
2007;10(1).

75. Buckling A, Rainey PB. Antagonistic coevolution between a bacterium and a
bacteriophage. Proceedings of the Royal Society B: Biological Sciences
[Internet]. 2002 [cited 2018 Nov 13];269(1494):931–6. Available from:
https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1690980/pdf/12028776.pdf

76. Obbard DJ, Jiggins FM, Bradshaw NJ, Little TJ. Recent and recurrent
selective sweeps of the antiviral RNAi gene Argonaute-2 in three species of
Drosophila. Mol Biol Evol [Internet]. 2011 Feb [cited 2018 Oct 31];28(2):1043–
56. Available from: http://www.ncbi.nlm.nih.gov/pubmed/20978039

77. Boots M, White A, Best A, Bowers R. How specificity and epidemiology drive
the coevolution of static trait diversity in hosts and parasites. Evolution (N Y).
2014;68(6):1594–606.

78. Paplauskas S, Duthie B, Tinsley MC. Meta-analysis of host population-level
diversity and variation in 'parasite success' [Dataset]. Dryad.

DATA ACCESSIBILITY

Reviewer URL:

https://datadryad.org/stash/share/E2NqLZ8KL2oYaPQLSYatmNNIeUub3aeC4ExfjJv

E5Hw

Data available from the Dryad Digital Repository: doi:10.5061/dryad.2bvq83bzq (78).

AUTHOR CONTRIBUTIONS

All authors discussed the results and contributed to the final manuscript. S.P.

performed the data collection, analysed the data and wrote the manuscript. B.D. and

M.T. contributed to the final version of the manuscript and supervised the project.

COMPETING INTERESTS

The authors have no competing interests.

FUNDING

S.P. acknowledges funding from the Natural Environment Research Council through

the IAPETUS doctoral training programme (grant no. S154).

ACKNOWLEDGEMENTS

We are grateful to A. Strauss, C. Ameline, P. Thrall, J. Walsman and A.K. Gibson for

sharing their raw data and clarifying their methods with us. We also thank

L.Braunholtz, K. Raines, L.F. Bussière and K. Morrison for their general statistical

and meta-analytical advice.