Competition effects regulating the composition of the microRNA pool

Sofia B. Raak[1, *], Jonathan G. Hanley[1], Cian O'Donnell[2, 3, *]

1 School of Biochemistry, University of Bristol, Biomedical Sciences Building, University Walk, Clifton, Bristol, BS8 1TD, United Kingdom
2 School of Engineering Mathematics and Technology, University of Bristol, University Walk, Clifton, Bristol, BS8 1TD, United Kingdom
3 School of Computing, Engineering and Intelligent Systems, Ulster University, Magee Campus, Derry/Londonderry, BT48 7JL, United Kingdom

* Corresponding authors

Abstract

MicroRNAS (miRNAs) are short non-coding RNAs that can repress mRNA translation to regulate localised protein synthesis. MiRNAs are matured in two steps: first primary-miRNA is processed into pre-miRNA by Drosha in the nucleus, then, following export to the cytosol, pre-miRNA is matured to miRNA by Dicer. Several pre-miRNA attributes affect maturation efficiency, and multiple pre-miRNAs are likely to compete for a limited pool of Dicer. However it is unknown how this competition influences the relative expression of mature miRNAs. Here, we study this in a model of pre-miRNA maturation fitted to experimental data. We find that pre-miRNAs that readily associate with Dicer, and pre-miRNAs with efficient processing by Dicer, are preferentially matured in competitive conditions when Dicer is scarce. To test our model predictions, we re-analysed previously published experimental data with reduced *Dicer1* mRNA expression *in vivo*. We calculated a proxy measure for pre-miRNA affinity to TRBP (a protein which loads pre-miRNAs to Dicer). According to our model, proteins with high TRBP affinity should outcompete proteins with low TRBP affinity. Accordingly, we found that pre-miRNAs with stronger TRBP association are more efficiently matured, which validated our proxy measure. We also found that pre-miRNAs with strong TRBP association are over-represented in competition conditions, consistent with our model predictions. Finally, using further simulations, we discovered that pre-miRNAs with low maturation rates can regulate the composition of mature miRNA through competition among pre-miRNAs. Overall, this work presents evidence of pre-miRNA competition regulating the composition of mature miRNAs.

Introduction

MicroRNAs (miRNAs) are small non-coding RNAs that inhibit protein translation via the RNA-induced gene silencing complex (RISC). MiRNAs are synthesised in the nucleus by RNA poly-

merase II/III as primary-miRNAs (pri-miRNAs), which are then cleaved by Drosha/DGCR8 to form precursor miRNas (Results; Figure 1 A). Pre-miRNAs are exported into the cytosol via Exportin-5 and transported to sites of local inhibition of protein translation, such as neuronal dendrites, where they are loaded onto Dicer, which cleaves the characteristic hairpin-loop structure to produce mature miRNA. The double-stranded miRNA is then loaded onto Argonaut proteins (Ago), which finish the maturation by ejecting the passenger strand to leave a single-stranded miRNA bound to Ago. The Ago-bound miRNA can subsequently undergo complementary base-pairing with target mRNAs and trigger RISC assembly, leading to silencing of protein translation.

MiRNAs are of particular importance in regulating gene expression in dendrites due to the size and morphology of neurons. For example, in a cortical pyramidal neuron, the soma is typically around 20µm in length, whereas some dendrites can extend hundreds of micrometres [1] and when considering the entire dendritic arborisation, the total length of dendrite for a single neuron can reach tens of millimetres [1, 2]. Given these large distances, it seems likely that neurons must use local mechanisms to control the spatial pattern of protein expression, rather than orchestrating control completely from the nucleus. Control and maintenance of dendritic pools of mRNA transcripts offer an elegant solution to highly localised and highly specific translational control of the post-synaptic proteome via RNA-induced gene silencing by miRNAs [3]. MiRNAs therefore play an important role in synaptic function and plasticity in the brain.

Competition is a recurring theme at all levels of biology, from competition between species and individual organisms to the competition for resources on the molecular level within the cell [REF]. These competition effects can indirectly regulate population dynamics by *e.g.* regulating noise in a gene circuit [4]. In tissues with high pre-miRNA expression levels, such as the brain, where up to 70% of known miRNAs have been detected [5], it is reasonable to assume that a large number of pre-miRNAs with different Dicer affinities and maturation efficiencies are competing for a limited amount of available Dicer. If this is the case, pre-miRNA competition for Dicer may indirectly regulate the composition of the mature miRNA pool.

For competition to be meaningful, individual components of a system must display distinct and diverse attributes. Among pre-miRNAs, both sequence and structural characteristics have been linked to the efficiency of maturation. Tsutsumi *et al.* (2011) [6] showed that *Drosophila* Dicer1 is more efficient at cleaving pre-miRNAs with a large loop size *in vitro*. Work by Luo *et al.* in HEK293T cells expressing recombinant Dicer also showed a preference of Dicer towards pre-miRNAs with a large loop structure and strong binding in the stem region [7], suggesting that loop size may also play a role in regulating pre-miRNA association *in vivo*. More recently,

Lee *et al.* (2023) [8] identified a conserved sequence motif, the GYM motif, which is recognised by the human Dicer1 double-stranded RNA binding domain and is associated with highly efficient cleavage of specific pre-miRNAs. These are all provide different advantages for select pre-miRNAs in maturation and might drive an over-representation of specific mature miRNAs in conditions with reduced Dicer availability.

MiRNA competition has been studied before, but most studies have focused on the competition between mature miRNAs, or miRNAs and non-coding competing endogenous RNAs, for the same mRNA targets (see [9–11] for some examples). To our knowledge, competition between different pre-miRNA species for proteins in the miRNA maturation pathway has not been reported previously. Here, we present a simple computational model of pre-miRNA maturation and pre-miRNA competition for Dicer based on mass-action kinetics (Results; Figure 1B). The model predicts that pre-miRNAs with both a high rate of association with Dicer and efficient dicing rates have a competitive advantage over other pre-miRNAs in systems with both abundant and severely reduced Dicer. Based on our model predictions, we identify pre-miRNA competition for Dicer *in vivo* from previously published experimental data [8]. Our work highlights the non-specific effects of pre-miRNA competition for Dicer on the global miRNA pool.

Results

MiRNAs with a fast association rate to Dicer display robust maturation levels in competitive conditions.

We designed a minimal model of pre-miRNA maturation (Figure 1B) that could account for the dynamics of mature miRNA production in previously published *in vitro* time series data [6] (See Methods). To summarise the model briefly, a pool of pre-miRNA can reversibly bind with free Dicer, then go through a subsequent maturation step, resulting in the conversion of pre-miRNA to mature miRNA and the release of Dicer back to the free pool. Alternatively, pre-miRNA could irreversibly bind with Dicer forming a stalled complex. We initialised the model with pre-miRNA and Dicer only.

In neurons, pre-miRNA maturation can take place hundreds of μm from the soma where transcription and initial maturation from pri-miRNA to pre-miRNA takes place (Results; Figure 1A). Experimental data has shown that mRNA transcripts can be transported across dendrites in bursts of speeds from 0.5-5 μms^{-1} between short pauses of being stationary [12, 13]. Under the assumption that pre-miRNAs are transported at a similar rate in a similar fashion to mRNAs we decided to omit pre-miRNA replenishing from our model.

During a 60 minute simulation of the model (Figure 1C), the concentration of free Dicer and pre-miRNA drops over time while the concentration of pre-miRNA complexed with Dicer increases, along with mature miRNA and stalled pre-miRNA and Dicer. At the end of the simulation, no free pre-miRNA remains and the mature miRNA concentration has reached a plateau.

Fig. 1: **MicroRNA maturation in the neuron and model design**

A MiRNA maturation and function in the neuron. MiRNA is synthesised in the nucleus as pri-miRNA by RNApol II/III as a long single-stranded RNA molecule with a central hairpin loop. Pri-miRNA is then cleaved by the Drosha/DGCR8 complex to release the hairpin loop as pre-miRNA. Pre-miRNA is then transported out of the nucleus into the cytosol by the Exportin-5 complex. In the cytosol, pre-miRNA can be transported into dendrites where Dicer, assisted by e.g. TRBP, binds pre-miRNA. Dicer cleaves the pre-miRNA by the loop structure and double-stranded mature miRNA can then be loaded into Argonaute proteins, where the passenger strand is ejected and the remaining single-stranded miRNA can form complimentary base-pair binding with target mRNA for targeted repression of local protein translation. **B** Model diagram of pre-miRNA maturation. In the computational model, pre-miRNA (P) can associate with Dicer (D) to form a transient Dicer-pre-miRNA complex (PD) with rate k_a. The Dicer-pre-miRNA complex can either deteriorate back to free pre-miRNA and Dicer at rate k_b, or go through pre-miRNA maturation at rate k_d to form free Dicer and mature miRNA (M). Alternatively, free pre-miRNA and Dicer can associate to form a stalled complex of Dicer-pre-miRNA (S) at rate k_c, which permanently binds pre-miRNA and Dicer in the system. **C** Dynamics of species concentration in the model. As time increases, the concentration of free Dicer ([D]) and pre-miRNA ([P]) reduces while the concentration of mature miRNA ([M]) and stalled pre-miRNA and Dicer ([S]) increases. The transient pre-miRNA Dicer complex [PD] is highly unstable and does not accumulate in the system.

Next, we used this model in a set of simulation experiments to investigate how multiple different pre-miRNA types might compete for a shared pool of Dicer. In real mammalian cells there are over 1900 types of mature human miRNAs recorded in miRBase [14], the online repository of identified miRNAs. While miRNAs are highly localised, many different miRNA species may still compete for a shared Dicer pool. The expression levels of these miRNAs vary substantially [15], implying that they have heterogeneous Dicer affinities and maturation rates.

In our model, four distinct reaction rates can affect pre-miRNA maturation: association

to Dicer (k_a), dissociation from Dicer without maturing (k_b), irreversible association to Dicer leading to a stalled complex (k_c), and maturation through dicing (k_d). This allowed us to dissect the role of each stage in pre-miRNA maturation by varying each reaction rate independently. To achieve this, we designed 8 theoretical species of pre-miRNA (Figure 2A). We increased each parameter value in turn either 10-fold (high) or 20-fold (2x high) from the optimised values to investigate what characteristics can be expected to confer advantages in competitive environments. As the actual concentration of Dicer in a cellular environment is unknown, we ran a series of simulations with 1nM of each pre-miRNA species present and a range of 0.01 to 10 nM Dicer available (Figure 2B-G). Since there was 8 nM total pre-miRNA, we should expect to see competition effects when Dicer concentration is $\lesssim$ 8 nM. In contrast, when Dicer concentration is $\gtrsim$ 8 nM, there should be little competition as there is sufficient Dicer available to bind all miRNA.

In a competitive environment with a single Dicer pool, high rates of association (k_a) leads to a higher amount of mature miRNA (Figure 2B). This is also true in the absence of competition (Figure 2D), however in competitive regimes, when Dicer concentration is low, slightly higher levels of pre-miRNAs with fast association rates reach the mature state compared to pre-miRNAs with high rates of maturation (k_d). When investigating fold-change from the simulation with 10 nM Dicer, a high rate of association (k_a) is highly advantageous both in presence and absence of competition (Figure 2C and E). In contrast, pre-miRNAs with a high maturation rate (k_d) and high level of stalling (k_c) are almost equally sensitive to a drop in fold change than pre-miRNAs with high association rates (Figure 2C and G), despite a high maturation rate being highly advantageous in pre-miRNA maturation (Figure 2B-E). When Dicer is available in abundance and competition is negligible, a high association rate (k_a) also leads to a higher rate of pre-miRNA maturation than a high maturation rate (k_d, Figure 2F). This suggests that pre-miRNAs with features that promote Dicer association provide both a competitive advantage in environments with reduced Dicer availability and reach the highest level of maturation efficiency.

Fig. 2: **Competition effects between multiple pre-miRNA species**

A Competition diagram. Each reaction rate was increased 10-fold (high) or 20-fold (2x high) per pre-miRNA species. **B-C** Effects of varying Dicer availability on final mature miRNA concentration in the presence (B) and absence (C) of competition. **D-E** Effects of varying Dicer availability on miRNA maturation fold change, as calculated based on 10nM Dicer availability in the presence (D) and absence (E) of competition. **F** Final mature miRNA concentration at 10nM Dicer availability. **G** Minimum Dicer needed to reach 0.5 fold-change for each pre-miRNA species in the presence and absence of competition.

Signatures of pre-miRNA competition for Dicer in experimental data

To test whether any evidence of pre-miRNA competition effects can be detected in experimental data, we next investigated the characteristics of miRNA sequences from the YAC128 mouse model of Huntington's disease where *Dicer1* mRNA expression levels have been reported to be reduced by half (Figure 4D), while mRNA expression levels of proteins in the pri-miRNA

processing machinery or pre-miRNA export were unaffected [16]. We hypothesised that, since Dicer expression was reduced in the Huntington's model, there should be stronger competition between the pre-miRNAs in that scenario, compared to wild-type animals where Dicer was more abundant. Our strategy was as follows. Our simulation results above predicted that pre-miRNAs with stronger Dicer affinity should out-compete those with low Dicer affinity. Therefore, we aimed to identify some proxy measure that correlates with Dicer affinity, calculate that quantity for each pri-miRNA, and ask if it is predictive of the fold-change in mature miRNA expression in the Huntington model relative to wild-type. The specific prediction was that high-Dicer-affinity miRNAs should show a lower fold-drop in expression than low-Dicer-affinity miRNAs (red line in Figure 3C). In contrast, a lack of competition for Dicer should result in a flat fold-change in miRNA expression, independent of Dicer affinity (dashed blue line in Figure 3C).

We used the miRNA sequencing data from 12-month old mice produced in Lee *et al.* (2011) [16] to look for pre-miRNA features that might influence maturation. The RNA-loading complex (RLC) protein TRBP has been shown to promote efficient loading and processing of pre-miRNAs in crowded environments [17], as well as promoting cleavage of the hairpin loop at the correct site and strand selection during loading onto Ago proteins [18] (Figure 3A). Additionally Takahashi *et al.* (2018) [19] showed that TRBP preferentially binds pre-miRNAs with a strong base-pair binding probability (BPP) in the stem region, where the mature miRNA sequence is located. Therefore, we considered a strong TRBP association to promote pre-miRNA association to Dicer (parameter k_a) and therefore promote maturation efficiency (parameter k_d), while antagonising dissociation (parameter k_b) and stalling (parameter k_c, see Figure 3B). We hypothesised that if high BPP leads to high Dicer association, and competition effects are present in pre-miRNA maturation, we would see a positive correlation between the fold change of miRNA expression following Dicer reduction and BPP (Figure 3C), as predicted by the efficient maturation of fast associating pre-miRNAs in Figure 1B and C. Conversely, if high BPP did not increase Dicer association or no competition is present in pre-miRNA maturation *in vivo* we would expect no correlation between fold change levels and BPP (Figure 3C).

Fig. 3: **Strategy for identifying competition in experimental data**

A Role of TRBP in pre-miRNA maturation. TRBP forms a complex with Dicer where it aids in recognition and loading of pre-miRNA onto the catalytic site of Dicer (i) in crowded environments [20]. TRBP also aids in determining the appropriate cleavage site (ii) and influences strand selction by Ago (iii) [18]. **B** Proposed effects of TRBP on model parameters. TRBP association corresponds to boosted association rate k_a and cleavage rate k_d, while the dissociation rate k_b and stalling rate k_c is moderately reduced. **C** Predicted effects in data. If competition effects are present, we expect that a fast association rate (k_a) should have the biggest effect. A strong base-pair binding probability (BPP) in the stem region should be associated with stronger TRBP binding and therefore more efficient Dicer loading. Consequently, we expect a positive correlation between BPP and miRNA expression if pre-miRNAs with high BPP have a competitive advantage in miRNA maturation (i; red solid line). If no competition is present, or fast Dicer association is not providing a significant advantage in miRNA maturation, we expect no relationship between BPP and miRNA expression (ii; blue dotted line).

We decided to use BPP for each pre-miRNA in Lee *et al.* (2011) [16] as a measure of TRBP association and therefore indirect association to Dicer. To find the BPP, the miRNA registry miRBase [14] was automatically scraped for structure information. For each miRNA, the mature sequence was extracted along with the published base pair bonds. The 5' and 3' strands were then aligned using RNACofold [21] and the BPP for the published bonds were extracted (see Figure 4A for the processing pipeline). The BPP for each base was then plotted against base position to provide an estimate of the stem structure (Figure 4B-C). To investigate whether BPP had any relation to miRNA expression levels, we took the area under the curve (AUC) as a single measurement and used Pearson's correlation measure to investigate the relationship between AUC and the miRNA expression level of wild-type and YAC128 mice (Figure 4E-F).

In wild-type mice with normal *Dicer1* mRNA expression, there was a positive correlation with the log_{10} expression level of mature miRNAs (r=0.34283, p<0.01; Figure 4E). In the YAC128 mice with significantly reduced *Dicer1* mRNA expression (Figure 4D, [16]), the correlation was also positive (r=0.43718, p < 0.001; Figure 4F). The fact that miRNA expression was

positively correlated with BPP in these two independent datasets demonstrates the validity of our strategy for using BPP as a proxy measure for pre-miRNA affinity to Dicer.

To test if the relationship between the $\log_{10}$(Expression) and BPP was significantly steeper in the YAC128 than in the wild-type mice, we calculated the t-score for the regression slopes as follows:

$$t = \frac{\beta_{WT} - \beta_{YAC128}}{\sqrt{SE_{\beta_{WT}} + SE_{\beta_{YAC128}}}} \tag{1}$$

where β_{WT} and β_{YAC128} are the estimated regression slopes for wild-type and YAC128 mice respectively, and $SE_{\beta_{WT}}$ and $SE_{\beta_{YAC128}}$ are the relevant standard errors for the estimated slopes. For our model fit, the calculated slopes were 1.49±0.449 and 1.89±0.427 for wild-type and YAC128 mice respectively. These produced a t-score of t=-2.720971 with df=84, leading to a p-value of p=0.00791. Thus, the YAC128 mice had a significantly steeper positive association between $\log_{10}$(Expression) and BPP than the wild-type mice, is indicative of competition effects partly driving the shift in miRNA expression in the YAC128 mouse model of Huntington's disease.

Following Dicer reduction in our model, the fold-change for all miRNAs eventually decreased (Figure 2C). When we looked at the relationship between the fold-change of YAC128 and wild-type mice we saw an overall reduction for most miRNAs, consistent with our model predictions. However, a positive correlation between fold-change and high AUC persisted (r=0.2458, p<0.05; Figure 4G). Consequently, pre-miRNAs with high BPP in the stem region are not as strongly affected by reduction in *Dicer* levels, providing further evidence for competition effects between pre-miRNA for Dicer affecting the composition of the pool of mature miRNA.

Fig. 4: Analysis of pre-miRNA structures suggest maturation advantage of pre-miRNAs with strong associations to Dicer in an HD mouse model

A Bioinformatics approach. Pre-miRNA hairpin structures were obtained from mirBase [14] and the mature sequence with recorded base pair bonds extracted. Mature 5' and 3' sequences were then aligned using RNACofold v. 2.4.13 from the ViennaRNA 2.0 package [21]. The base pair binding probability (BPP) was then extracted for the recorded base pair bonds. Where no base pair bond was recorded the mean BPP for the i_{th} 5' nucleotide was used. **B** Subset of base pair binding probabilities. Black solid line represents mean base pair binding probabilities across analysed miRNAs, with shaded area representing standard error of the mean. Red dashed line represents an example miRNA (let-7g) with higher mean BPP, blue dotted line an example miRNA (miR-511) with lower than mean BPP. Position on x-axis denotes nucleotide position, with negative numbers referring to position on the 5' strand and positive numbers position on the 3' strand. The centre arbitrarily assigned 0 corresponds to the cleavage site. **C** Base pair binding probabilities of mature miRNA sequences. Graphical representation of subset of the mean base pair binding probabilities of all miRNAs (black) and an example miRNA with higher (let-7g, red) and lower (miR-511, blue) mean BPP. **D** *Dicer1* mRNA expression in wild-type and YAC128 mice at 5 months and 12 months of age as determined by qPCR. P**<0.01 as determined by Mann-Whitney U-test, error bars represent standard deviation. Figure adapted from [16]. **E-F** Pearson correlation of miRNA expression in 12 month old wild-type (**E**) and YAC128 (**F**) mice with BPP area under curve (AUC). **G** Pearson correlation of fold change between 12 month old YAC128 and wild-type miRNA expression with BPP AUC. In **E-G** solid line represents least squares linear regression, with shaded area calculated from the standard error of the intercept and gradient. In **G** outliers above fold change 4 were removed.

Effects of differential pre-miRNA expression on global miRNA composition

In the previous simulations (Figure 2) we studied the effects of pre-miRNA competition for Dicer by varying Dicer concentration from low, scarce regimes to high, abundant regimes. In those simulations all eight pre-miRNA species initially had equal concentration, and differed only in their reaction kinetics. However, in real cells different pre-miRNA types likely have different abundances. This heterogeneity may have knock-on effect on Dicer competition. For example if one pre-miRNA is highly upregulated, then it may sequester more Dicer, leaving less Dicer free for other pre-miRNA types.

To investigate whether and how differential pre-miRNA expression could effect the global mature miRNA pool via Dicer competition, we returned to the same computational model used previously and successively 'overexpressed' (Figure 5A-B) or 'knocked out' (Figure 5C-D) each of our eight simulated pre-miRNAs in turn, by changing the initial pre-miRNA concentration to either 5 nM (overexpression) or 0 nM (knockout). We then ran our model with 1.55 nM available Dicer, chosen as a condition with notable competition (Figure 2B-E, F), and calculated the fold change between the mature miRNA expression in conditions with increased or knocked-down pre-miRNAs compared to the same conditions where all pre-miRNAs were expressed equally at an initial concentration of 1nM.

Fig. 5: **Effects of pre-miRNA increase and removal on global miRNA population**

Effects on miRNA maturation in the absence (**B, D**) and presence (**A, C**) of competition following increase (5nM pre-miRNA, **A-B**) and removal (0nM, **C-D**) of specific pre-miRNAs in a system with 1.55nM available Dicer.

The simulation results are summarised in the four matrices shown in Figure 5. For a given matrix, each row corresponds to a single simulation run where a particular pre-miRNA concentration was either increased (panels A, B) or knocked out (panels C, D). The left two matrices (panels A, C) show results with 1.55 nM Dicer – encouraging competition – whereas the right two matrices (panels B, D) are the same simulations but with abundant Dicer and no competition, as a control comparison. The number and colour of each matrix element indicates the fold-change in mature miRNA expression for each of the eight species (matrix columns).

As expected, increasing the pre-miRNA concentration of a given species always increased its mature miRNA expression (red diagonals with values > 1 in Figure 5A, B). Similarly, knocking out a pre-miRNA species lead to zero mature miRNA expression (blue diagonals with values = 0 in Figure 5C, D).

Following increase of specific miRNAs, there was a broad trend of a reduction in mature

miRNA of all other species in the system in a competitive environment, while the increased pre-miRNAs were all upregulated (Figure 5A). The degree of upregulation was not homogeneous between the different miRNAs, and when compared with upregulated pre-miRNAs in the absence of competition (Figure 5B) it is evident that pre-miRNAs with a high k_b and k_c are more sensitive to competition even when available in abundance. Pre-miRNAs with a high kb also seem to be sensitive to changes in the amount of global pre-miRNA, while increasing the initial amount of pre-miRNAs with either high kb or k_c appears to have strong effects on most pre-miRNAs in the system. While these pre-miRNAs are not highly represented among mature miRNAs (Figure 2B), their broad effects on other miRNAs in Figure 5A suggest that they play an indirect role in regulating the composition of mature miRNAs through competition for Dicer. These effects are largely mirrored following removal of each initial pre-miRNA (Figure 5C). Overall, these results demonstrate that change in expression levels of one type of pre-miRNA species can have heterogeneous knock-on effects on the expression of other miRNAs, in regimes where Dicer is scarce.

Discussion

Modelling studies in the last few decades have discovered key principles underlying competition for resources during during gene expression and protein translation [22]. In contrast, competition's effect on miRNA maturation is less well understood. Here, we have presented a simple model of pre-miRNA competition for a limited Dicer availability during miRNA maturation. In models of competition between multiple hypothetical species of pre-miRNA, we found that high association rates provides a competitive advantage in conditions with reduced Dicer availability. We also found empirical evidence of competition effects partially regulating the composition of neuronal miRNAs in the YAC128 model of Huntington's disease. Finally, we found that certain pre-miRNAs are more sensitive to changes in Dicer availability in conditions with increased or removed pre-miRNAs.

We showed that a strong association with Dicer (high k_a) is beneficial both in increasing the miRNA maturation efficiency, but also provides a competitive advantage following reductions in Dicer availability (Figure 2). Based on the work of Takahashi *et al.* [19] we reasoned that pre-miRNAs with high BPP in the stem region would preferentially associate with TRBP and in turn more efficiently be loaded onto Dicer [17]. As the expression of Dicer, but not other components of the pre-miRNA maturation pathway, was significantly reduced at the mRNA level in the 12-month old YAC128 mice [16] we decided to use this as an existing model of Dicer competition *in vivo*.

We investigated the relationship between miRNA expression in the YAC128 Huntington's mouse model [16] and BPP in the stem region. We identified a weakly positive but statistically significant positive correlation between higher expressing miRNAs and BPP in both the wild-type and YAC128 mice; though the slope of the association was significantly stronger in the YAC128 mice (Figure 4). These results provide proof-of-principle for our assumption that TRBP BPP is a valid proxy measure for Dicer affinity. We also found a weakly positive correlation in the fold change between YAC128 and wild-type mice. This shows that pre-miRNAs with a high BPP, and therefore higher TRBP association (and consequently more efficient Dicer loading and maturation), are less affected by reduction in available Dicer. These results are important because they show that competition for Dicer in part regulate the composition of the global miRNA pool. They also suggest that competition effects might play a role for the disruption of the miRNA expression profile in Huntington's disease.

The degree of competition can not only be affected by Dicer availability, but also by differential expression of the various pre-miRNAs. We investigated this by either increasing or removing specific pre-miRNAs (mimicking up-regulation or knock-out experiments) and assessing the changes in the global mature miRNA pool in the presence or absence of competition (Figure 5). We found that pre-miRNAs with fast dissociation and and stalling rates were strongly affected when changes were made to any other pre-miRNA. These effects were mirrored following removal of pre-miRNAs from the system, but to a lesser degree. This can be explained by the relative ratio of increased or decreased available Dicer in the system. Following an increase in initial pre-miRNA availability, the Dicer availability drops by almost 40%. In contrast, when one pre-miRNA species is removed there is only a relative increase by 12% of Dicer availability to the system. Consequently, the effects of increasing specific pre-miRNAs are more pronounced than those following a removal.

What might be the functional benefit of pre-miRNA competition? In evolutionary terms, MiRNAs are phylogenetically stable once they emerge. Novel miRNAs are rarely lost in descendants after arising [23]. MiRNAs are also continuously emerging and undergo changes in sequence specificity and increase sequence diversification [23]. Taken together, these suggest that competition for pre-miRNA maturation is not detrimental and could even be positive. The selective effect of pre-miRNA expression following changes in availability of pre-miRNAs with either fast dissociation and stalling rates (Figure 5 indicate that, while these inefficiently matured pre-miRNAs are not highly represented among mature miRNAs (Figure 2B, D, F), they do play an important role in shaping the composition of mature miRNAs. Competition between pre-miRNAs for Dicer might therefore help stabilise and fine-tune the mature miRNA expression.

As with all models, there are limitations with our model. First, the permanently stalled pre-miRNA Dicer complex, which can neither dissociate or complete miRNA maturation, is not biologically plausible. While useful to account for the ceiling effect after around 40% of wild-type pre-miRNA are diced [6] (Figure 6A), there is no evidence of pre-miRNA and Dicer being removed together from the pre-miRNA maturation pathway *in vivo*. Nevertheless, our model is a single, well-mixed compartment observed for 1 hour. It is not unreasonable to consider the stalling a prolonged, but temporary, interruption to the pre-miRNA maturation process, for example by strong but misaligned association with a subset of pre-miRNAs within a species. In a more strongly biological model version, this term could be exchanged to *e.g.* dynamic pre-miRNA availability, spatial constraints, or including active and inactive Dicer states. Second, our model has a fixed initial concentration of pre-miRNA, and therefore does not take into account pre-miRNA production or transportation rates. However, we do not expect this to affect our conclusions, because these processes likely happen on a longer timescale of hours–days, whereas our simulations were over tens of minutes.

In conclusion, we have presented a simple model of pre-miRNA maturation capable of replicating experimental data. We used our model to dissect pre-miRNA characteristics that influence maturation, and based on our model predictions formed a testable hypothesis for identifying pre-miRNA competition for Dicer in experimental data. Through bioinformatic analysis we then found evidence of pre-miRNA competition *in vivo*. Finally, we uncovered a possible indirect regulatory role of lowly expressed miRNAs as directors of the global miRNA composition through pre-miRNA competition effects.

Methods

Software

All simulations and bioinformatics were performed in Python v3.9 if not otherwise indicated in the text. Ordinary differential equations (ODEs) were solved with the scipy integrate [24] library using the LSODA solver [25, 26], except for simulations in Figure **??**B and D where the Runge-Kutta method of order 5(4) [27, 28] was used instead to reduce numerical errors in solving. All code used in this paper is available from `https://github.com/SofiaRaak/Dicer-miRNA-dynamics-model`.

Model design

We designed our initial model (Results; Figure 1B) to replicate the *in vitro* test-conditions described in Tsutsumi *et al.* (2011) [6], where 1nM pre-miRNA was incubated with 5nM recombinant *Drosophila* Dicer1 for 60 minutes and pre-miRNA maturation tracked by gel shift to determine the fraction of diced miRNA (Figure 6). Based on this, we included a set amount of pre-miRNA and Dicer in the model at the start of each simulation (Figure 6A). The pre-miRNA was then allowed to associate with Dicer to form a pre-miRNA-Dicer complex, which could either dissociate or go through a maturation step to form mature miRNA and release the Dicer again to re-join the pool. Alternatively, the pre-miRNA could irreversibly form a complex with Dicer to form a stalled, permanently sequestered pre-miRNA-Dicer species. While not necessarily biologically intuitive, this term was added to the model to account for the dynamics of pre-miRNA maturation in the experimental data.

We mathematically described our model as a series of ordinary differential equations (ODEs), where the left hand side describes the change in concentration for each species in the system and the right hand side describes the interactions and reactions that increase or decrease the concentration for each species:

$$\frac{d[P_i]}{dt} = k_{b_i}[PD_i] - (k_{a_i} + k_{c_i})[P_i][D] \tag{2}$$

$$\frac{d[PD_i]}{dt} = k_{a_i}[P_i][D] - (k_{b_i} + k_{d_i})[PD_i] \tag{3}$$

$$\frac{d[D]}{dt} = (k_{b_i} + k_{d_i})[PD_i] - (k_{a_i} + k_{c_i})[P_i][D] \tag{4}$$

$$\frac{d[S_i]}{dt} = k_{c_i}[P_i][D] \tag{5}$$

$$\frac{d[M_i]}{dt} = k_{d_i}[PD_i] \tag{6}$$

where P_i represents the concentrations of free pre-miRNA of species i, D represents free Dicer, PD_i represents pre-miRNA bound to Dicer, S_i represents stalled pre-miRNA-Dicer complex and M_i represents mature miRNA concentration (all in nM). The reaction rates $k_{a_i-d_i}$ represent the reaction rates for pre-miRNA-Dicer association, dissociation, stalling and dicing respectively, see table 1. The model was then fitted to experimental data.

Fitting model to data

In order to fit the parameters of the model to experimental data initial parameter values were specified and the system of ODEs solved and output compared to the experimental data in Tsutsumi *et al.* (2011) [6]. The difference between the model output and experimental data was then quantified as the sum squared error (SSE) between the simulation time series and the experimental data time series and fed into the CMA-ES [29] optimiser, iteratively updating the parameter values and repeating the process until a good fit was achieved (see fig 6). The optimised parameter values for both wild-type (WT) and mutant (short-loop; SL) miRNA are shown in table 1. In all subsequent simulations, the WT parameters were used as a foundation for the different pre-miRNA species.

Tab. 1: **Parameter values used in model**

Symbol	Parameter	Value	Source
$k_{a_{WT}}$	Pre-miRNA association rate, wild-type	$0.2218\text{s}^{-1}\text{nM}^{-1}$	Optimisation
$k_{a_{SL}}$	Pre-miRNA association rate, short-loop	$6.4614\text{s}^{-1}\text{nM}^{-1}$	Optimisation
$k_{b_{WT}}$	Pre-miRNA dissociation rate, wild-type	5.6353s^{-1}	Optimisation, [6]
$k_{b_{SL}}$	Pre-miRNA dissociation rate, short-loop	954.35s^{-1}	Optimisation, [6]
$k_{c_{WT}}$	Pre-miRNA stalling rate, wild-type	$0.0123\text{s}^{-1}\text{nM}^{-1}$	Optimisation
$k_{c_{SL}}$	Pre-miRNA stalling rate, short-loop	$0.01995\text{s}^{-1}\text{nM}^{-1}$	Optimisation
$k_{d^{WT}}$	Pre-miRNA dicing rate, wild-type	4.0307s^{-1}	Optimisation
$k_{d_{SL}}$	Pre-miRNA dicing rate, short-loop	0.0343s^{-1}	Optimisation

Fig. 6: **Optimisation of reaction rates in computational model**

A Optimisation flow for fitting computational model to experimental data from [6]. The fraction of diced miRNA at specific time points in the model was compared to experimental data and the error calculated as the sum of squares (SSE) of the difference between the experimental data and data generated by the model. The error was then fed back into the optimisation algorithm CMA-ES [29] and reaction rates adjusted before output was compared again. The procedure was repeated for both wild-type (long-loop) and mutant (short-loop) pre-miRNAs until the error was maximally minimised. **B** Reproduced wild-type experimental data from model optimisation. **C** Reproduced short-loop experimental data from model optimisation.

Bioinformatics

All miRNA expression data used in bioinformatics are available in [16]. Briefly, miRNAs present in expression data from wild-type and YAC128 mice were identified in miRBase [14] and their pre-miRNA structure recorded. MiRNAs that were not found in miRBase or that did not have a pre-miRNA structure were excluded. From the pre-miRNA structure, the mature sequences were isolated and their base pair bonds recorded. The mature sequences were then aligned with RNACofold v. 2.4.13 [21] and the base pair binding probability (BPP) for the recorded base pair bonds were extracted. In the absence of a recorded base pair bond, the mean BPP was chosen (Results; Figure 4A). The BPP was then plotted against the nucleotide position to create a BPP curve (Results; Figure 4B). As a single measure of BPP, the area under the curve was taken and plotted either against the $\log_{10}$(miRNA Expression) or the fold-change in miRNA expression between wild-type and YAC128 mice (Results; Figure 4).

References

1. Gao L, Liu S, Wang Y, Wu Q, Gou L, Yan J. Single-neuron analysis of dendrites and axons reveals the network organization in mouse prefrontal cortex. Nature Neuroscience. 2023;26(6):1111–1126. doi:10.1038/s41593-023-01339-y.

2. Mohan H, Verhoog MB, Doreswamy KK, Eyal G, Aardse R, Lodder BN, et al. Dendritic and Axonal Architecture of Individual Pyramidal Neurons across Layers of Adult Human Neocortex. Cerebral Cortex. 2015;25(12):4839–4853. doi:10.1093/cercor/bhv188.

3. Hu Z, Li Z. miRNAs in synapse development and synaptic plasticity. Current Opinion in Neurobiology. 2017;45:24–31. doi:10.1016/j.conb.2017.02.014.

4. Wei L, Li S, Zhang P, Hu T, Zhang MQ, Xie Z, et al. Characterizing microRNA-mediated modulation of gene expression noise and its effect on synthetic gene circuits. Cell Reports. 2021;36(8):109573. doi:10.1016/j.celrep.2021.109573.

5. Cao X, Yeo G, Muotri AR, Kuwabara T, Gage FH. NONCODING RNAS IN THE MAMMALIAN CENTRAL NERVOUS SYSTEM. Annual Review of Neuroscience. 2006;29(1):77–103. doi:10.1146/annurev.neuro.29.051605.112839.

6. Tsutsumi A, Kawamata T, Izumi N, Seitz H, Tomari Y. Recognition of the pre-miRNA structure by Drosophila Dicer-1. Nature Structural & Molecular Biology. 2011;18(10):1153–1158. doi:10.1038/nsmb.2125.

7. Luo QJ, Zhang J, Li P, Wang Q, Zhang Y, Roy-Chaudhuri B, et al. RNA structure probing reveals the structural basis of Dicer binding and cleavage. Nature Communications. 2021;12(1). doi:10.1038/s41467-021-23607-w.

8. Lee YY, Kim H, Kim VN. Sequence determinant of small RNA production by DICER. Nature. 2023;615(7951):323–330. doi:10.1038/s41586-023-05722-4.

9. Nyayanit D, Gadgil CJ. Mathematical modeling of combinatorial regulation suggests that apparent positive regulation of targets by miRNA could be an artifact resulting from competition for mRNA. RNA. 2015;21(3):307–319. doi:10.1261/rna.046862.114.

10. Yuan Y, Liu B, Xie P, Zhang MQ, Li Y, Xie Z, et al. Model-guided quantitative analysis of microRNA-mediated regulation on competing endogenous RNAs using a synthetic gene circuit. Proceedings of the National Academy of Sciences. 2015;112(10):3158–3163. doi:10.1073/pnas.1413896112.

11. Yuan Y, Ren X, Xie Z, Wang X. A quantitative understanding of microRNA-mediated competing endogenous RNA regulation. Quantitative Biology. 2016;4(1):47–57. doi:10.1007/s40484-016-0062-5.

12. Yoon YJ, Wu B, Buxbaum AR, Das S, Tsai A, English BP, et al. Glutamate-induced RNA localization and translation in neurons. Proceedings of the National Academy of Sciences. 2016;113(44):E6877–E6886. doi:10.1073/pnas.1614267113.

13. Lionnet T, Czaplinski K, Darzacq X, Shav-Tal Y, Wells AL, Chao JA, et al. A transgenic mouse for in vivo detection of endogenous labeled mRNA. Nature Methods. 2011;8(2):165–170. doi:10.1038/nmeth.1551.

14. Kozomara A, Birgaoanu M, Griffiths-Jones S. miRBase: from microRNA sequences to function. Nucleic Acids Research. 2019;47(D1):D155–D162. doi:10.1093/nar/gky1141.

15. Keller A, Gröger L, Tschernig T, Solomon J, Laham O, Schaum N, et al. miRNATissueAtlas2: an update to the human miRNA tissue atlas. Nucleic Acids Research. 2022;50(D1):D211–D221. doi:10.1093/nar/gkab808.

16. Lee ST, Chu K, Im WS, Yoon HJ, Im JY, Park JE, et al. Altered microRNA regulation in Huntington's disease models. Experimental Neurology. 2011;227(1):172–179. doi:https://doi.org/10.1016/j.expneurol.2010.10.012.

17. Fareh M, Yeom KH, Haagsma AC, Chauhan S, Heo I, Joo C. TRBP ensures efficient Dicer processing of precursor microRNA in RNA-crowded environments. Nature Communications. 2016;7(1):13694. doi:10.1038/ncomms13694.

18. Ross, Tambe A, Mary, Cameron, Catherine, Jennifer. Dicer-TRBP Complex Formation Ensures Accurate Mammalian MicroRNA Biogenesis. Molecular Cell. 2015;57(3):397–407. doi:10.1016/j.molcel.2014.11.030.

19. Takahashi T, Nakano Y, Onomoto K, Murakami F, Komori C, Suzuki Y, et al. LGP2 virus sensor regulates gene expression network mediated by TRBP-bound microRNAs. Nucleic Acids Research. 2018;46(17):9134–9147. doi:10.1093/nar/gky575.

20. Fareh M, Yeom KH, Haagsma AC, Chauhan S, Heo I, Joo C. TRBP ensures efficient Dicer processing of precursor microRNA in RNA-crowded environments. Nature Communications. 2016;7(1):13694. doi:10.1038/ncomms13694.

21. Lorenz R, Bernhart SH, Höner Zu Siederdissen C, Tafer H, Flamm C, Stadler PF, et al. ViennaRNA Package 2.0. Algorithms for Molecular Biology. 2011;6(1):26. doi:10.1186/1748-7188-6-26.

22. Sabi R, Tuller T. Modelling and measuring intracellular competition for finite resources during gene expression. Journal of The Royal Society Interface. 2019;16(154):20180887. doi:10.1098/rsif.2018.0887.

23. Berezikov E. Evolution of microRNA diversity and regulation in animals. Nature Reviews Genetics. 2011;12(12):846–860. doi:10.1038/nrg3079.

24. Virtanen P, Gommers R, Oliphant TE, Haberland M, Reddy T, Cournapeau D, et al. SciPy 1.0: Fundamental Algorithms for Scientific Computing in Python. Nature Methods. 2020;17:261–272. doi:10.1038/s41592-019-0686-2.

25. Hindmarsh AC. ODEPACK, a systemized collection of ODE solvers. Scientific computing. 1983;.

26. Petzold L. Automatic Selection of Methods for Solving Stiff and Nonstiff Systems of Ordinary Differential Equations. SIAM Journal on Scientific and Statistical Computing. 1983;4(1):136–148. doi:10.1137/0904010.

27. Dormand JR, Prince PJ. A family of embedded Runge-Kutta formulae. Journal of Computational and Applied Mathematics. 1980;6(1):19–26. doi:https://doi.org/10.1016/0771-050X(80)90013-3.

28. Shampine LF. Some Practical Runge-Kutta Formulas. Mathematics of Computation. 1986;46(173):135. doi:10.2307/2008219.

29. Hansen N, Akimoto Y, Baudis P. CMA-ES/pycma on Github; 2019. Zenodo, DOI:10.5281/zenodo.2559634. Available from: `https://doi.org/10.5281/zenodo.2559634`.