Genetic background affects the strength of crossover interference in house mice

Andrew P Morgan[1] and Bret A Payseur[2]

[1]Department of Medicine, University of North Carolina, Chapel Hill, NC

[2]Laboratory of Genetics, University of Wisconsin, Madison, WI

Corresponding author:
Andrew P Morgan (`andrew.parker.morgan@gmail.com`)
Department of Medicine
University of North Carolina
101 Manning Drive #7085
Chapel Hill NC 27599-7085

Word count: 4509

Abstract

Meiotic recombination is required for faithful chromosome segregation in most sexually reproducing organisms and shapes the distribution of genetic variation in populations. Both the overall rate and the spatial distribution of crossovers vary within and between species. Adjacent crossovers on the same chromosome tend to be spaced more evenly than expected at random, a phenomenon known as crossover interference. Although interference has been observed in many taxa, the factors that influence the strength of interference are not well understood. We used house mice (*Mus musculus*), a well-established model system for understanding recombination, to study the effects of genetics and age on recombination rate and interference in the male germline. We analyzed crossover positions in 503 progeny from reciprocal F1 hybrids between inbred strains representing the three major subspecies of house mice. Consistent with previous studies, autosomal alleles from *M. m. musculus* tend to increase recombination rate, while inheriting a *M. m. musculus* X chromosome decreases recombination rate. Old males transmit an average of 0.6 more crossovers per meiosis (5.0%) than young males, though the effect varies across genetic backgrounds. We show that the strength of crossover interference depends on genotype, providing a rare demonstration that interference evolves over short timescales. Differences between reciprocal F1s suggest that X-linked factors modulate the strength of interference. Our findings motivate additional comparisons of interference among recently diverged species and further examination of the role of paternal age in determining the number and positioning of crossovers.

Introduction

Meiotic recombination is a potent cellular and evolutionary force. In most species that reproduce sexually, recombination is required for the segregation of chromosomes during meiosis, with too few or too many crossovers leading to aneuploid offspring that are usually either non-viable or phenotypically abnormal (reviewed in Hassold and Hunt (2001)). Recombination creates new allelic combinations and thereby shapes the trajectories of beneficial and deleterious mutations in populations (Fisher 1930; Hill and Robertson 1966; Felsenstein 1974; Barton 1995).

Although its functional and evolutionary roles impose constraints on recombination, the number and positioning of crossovers nonetheless vary among individuals, both within and between species (Smukowski and Noor 2011; Stapley *et al.* 2017; Brazier and Glémin 2022). Evidence for a heritable component to this variation comes from responses to artificial selection (Chinnici 1971; Kidwell and Kidwell 1976), resemblance among relatives (Kong *et al.* 2004; Coop *et al.* 2008; Brekke *et al.* 2022), and the identification of genetic variants associated with recombination rate differences in populations (Kong *et al.* 2008; Johnston *et al.* 2016; Hunter *et al.* 2016). Likewise, the fine-scale spatial distribution of recombination has been shown to vary within and between populations in several taxa (reviewed in Paigen and Petkov (2010)). In mammals, one determinant of this variation is the interaction between by nucleotide sequence at so-called "hotspots" and the protein PRDM9, which applies chromatin marks that designate the location of double-strand breaks that may mature into crossovers (Parvanov *et al.* 2010).

Despite ample evidence for genetic variation in the overall rate and fine-scale distribution of recombination, the causes of the relative positions of crossovers on the same chromosome at macroscopic scale remain poorly understood. Crossovers are spaced more evenly than expected if they are positioned at random with respect to each other, an observation made during the construction of the first linkage map (Sturtevant 1913, 1915). This pattern implies that the designation of a crossover in one chromosomal interval inhibits the formation of another crossover nearby, a phenomenon known as crossover interference. Crossover interference is observed across a wide variety of species (Berchowitz and Copenhaver 2010; Otto and Payseur 2019), with *Schizosaccharomyces pombe* (Munz 1994), *Aspergillus nidulans* (Strickland 1958), and *Aspergillus fumigatus* (Auxier *et al.* 2023) providing documented exceptions.

The magnitude of crossover interference is rarely compared across taxa, but distantly related species appear to differ (Otto and Payseur 2019). For example, interference in dogs is estimated to be 2-3 times as strong as interference in humans (Campbell *et al.* 2016). The main evidence for variation in interference landscapes within

species comes from contrasts between the sexes (Otto and Payseur 2019). Males show stronger interference in humans (Campbell *et al.* 2015), dogs (Campbell *et al.* 2016), mice (Peterson and Payseur 2021), and gray geese (Torgasheva and Borodin 2017), whereas females show stronger interference in cattle (Wang *et al.* 2016) and *Arabidopsis thaliana* (Basu-Roy *et al.* 2013). In *A. thaliana*, a lower proportion of crossovers escapes interference in females than in males (Basu-Roy *et al.* 2013).

The extent to which the magnitude of interference and the percentage of interfering crossovers vary among individuals of the same sex is virtually unknown. One study in cattle found an association between the strength of interference in males and variation in *Nek9*, a gene involved in meiotic spindle formation, implying the existence of heritable differences (Wang *et al.* 2016). A quantitative trait locus modulates a difference in interference between two wild-derived strains of house mice (Wang *et al.* 2019). The absence of information about variation in interference among individuals is striking given that this phenomenon is a key determinant of crossover positioning and exerts substantial influence on the distribution of relatedness in populations (Caballero *et al.* 2019; Veller *et al.* 2020).

House mice from the *Mus musculus* species complex provide a useful system for examining variation in crossover interference. Interference in mice is strong, with distances between double crossovers that amount to appreciable percentages of chromosomal length (Broman *et al.* 2002). For example, the minimal physical separation between double crossovers on chromosome 1 detected in 3026 male mouse meioses was 57 Mb (Petkov *et al.* 2007; Paigen *et al.* 2008). Variation in crossover interference has been proposed to explain sex differences in recombination rate in mice (Petkov *et al.* 2007). The molecular and cellular pathways that culminate in crossovers are well understood in mice (reviewed in Lam and Keeney (2014) and Gray and Cohen (2016)). Furthermore, wild-derived inbred strains capture natural variation in recombination rate (Peterson and Payseur 2021) and enable reproducible comparisons of interference. Here, we examine the transmission of crossovers in F1 hybrids generated from crosses between wild-derived strains of mice belonging to the three principal subspecies of house mice: *M. m. domesticus*, *M. m. musculus* and *M. m. castaneus*. We uncover heterogeneity in the strength of crossover interference, demonstrating that this important determinant of the recombination landscape evolves on short timescales.

Materials and methods

Mouse breeding

The breeding scheme used in this work has been described previously (Morgan *et al.* 2019) and is shown in **Figure 1**. Reciprocal F1 hybrid males were generated in all possible pairwise combinations between CAST/EiJ (*M. m. castaneus*), PWK/PhJ (*M. m. musculus*) and WSB/EiJ (*M. m. domesticus*). Founder individuals were

purchased from the Jackson Laboratory (Bar Harbor, ME) between 2008 and 2011. Shortly after reaching sexual maturity, F1 males were mated to multiple FVB/NJ females and liveborn N2 progeny were collected and sacrificed at birth by cervical dislocation. At least 2 litters were collected from each male by age 4 months ("young age"). F1 males were allowed to continue mating freely, and females were replaced if they stopped producing offspring. Once F1 males were approaching reproductive senescence (as evidenced by increasing interval between litters or declining litter size), at least 2 additional litters were collected at an age of at least 15 months ("old age"). Breeding was completed in June 2011. The list of N2 progeny used in this study is provided in **Table S1**. The study was approved by the Institutional Animal Care and Use Committee of the University of North Carolina at Chapel Hill, and all animal husbandry was conducted in the university's Association for Assessment and Accreditation of Laboratory Animal Care-accredited facility (#A3410-01; Animal Welfare Assurance) in accordance with institutional and federal regulations.

DNA extraction and genotyping

As previously described (Morgan *et al.* 2019), genomic DNA was extracted from whole tails or heads of N2 mice and from tail clips of adult representatives of parental inbred strains, using DNeasy Blood & Tissue Kits (catalog no. 69506; QIAGEN, Valencia, CA). Approximately 1.5μg of DNA per sample was shipped to Neogen (Lincoln, NE) for genotyping with one of two Mouse Universal Genotyping Arrays, dubbed MegaMUGA (approximately 77K probes) (Morgan *et al.* 2016) or MiniMUGA (approximately 11K probes) (Sigmon *et al.* 2020). Both are designed on the Illumina Infinium HD platform (Steemers *et al.* 2006) and SNP content is optimized for genetic mapping in populations descended from common laboratory and wild-derived mouse strains. The vendor provided raw genotype calls at all markers on each array. The number of N2 progeny genotyped on each array, from each genotype and paternal age, are shown in **Table 1**.

Filtering and harmonizing genotypes

Genotype data were processed in R v4.1.1 (R Foundation for Statistical Computing, http://www.r-project.org/) using the argyle package v0.2.2 (Morgan 2016). Physical positions (in the GRCm39 assembly) and genetic positions on the standard mouse genetic map (Cox *et al.* 2009) for array markers were assigned based on the annotations produced by Karl Broman and colleagues (https://github.com/kbroman/MUGAarrays). Initial filtering and quality control steps were performed separately for each combination of array and paternal genotype, since different subsets of markers will be informative depending on paternal genotype. Autosomal markers which met the following criteria were retained: (1) $< 10\%$ missing genotypes; (2) fixed for opposite alleles in the paternal grandparental strains; and (3) not subject to substantial segregation distortion (non-reference allele frequency $> 20\%$ and $< 80\%$). For all paternal genotypes besides PWK/PhJ×WSB/EiJ and

WSB/EiJ×PWK/PhJ, genotyping was split between the two array platforms. To mitigate biases arising from differential marker density, for each marker on the more sparse array (MiniMUGA), the nearest marker on the more dense array (MegaMUGA) was retained, and other markers were discarded. (This approach retains markers that are shared between the two arrays.) We also sought to mitigate technical biases arising from differences between array platform in the proportion of each chromosome spanned by informative markers, as follows. For each paternal genotype, let $(p_{\mathrm{mini}}, d_{\mathrm{mini}})$ and $(p_{\mathrm{mega}}, d_{\mathrm{mega}})$ be the most proximal and distal markers on MiniMUGA and MegaMUGA, respectively; and let $p_{\mathrm{inner}} = max\,(p_{\mathrm{mini}}, p_{\mathrm{mega}})$ and $d_{\mathrm{inner}} = min\,(d_{\mathrm{mini}}, d_{\mathrm{mega}})$. Any markers proximal to p_{inner} or distal to d_{inner} were discarded. The number of markers retained for each combination of paternal genotype and platform, and their physical span, are shown in **Table 2**.

Haplotype and crossover inference

Filtered genotypes were converted to `R/qtl2` format, recoded with respect to the inbred paternal grandparents. Although our breeding scheme is not actually a backcross, expected genotype frequencies are 0.5 in N2 offspring so genotype probabilities were modelled as a a backcross using `R/qtl2`'s hidden Markov model (HMM) (Broman *et al.* 2019). The most likely sequence of haplotype states at each marker was inferred using the Viterbi algorithm (`qtl2::viterbi()`), allowing 1% genotyping error. Uninterrupted runs of the same marker-wise haplotype state thus correspond to transmitted haplotype segments, and crossovers are implicitly defined as boundaries between haplotype segments. Inferred haplotype segments are provided in **Table S2**, and the number of autosomal crossovers per individual in **Table S3**.

Statistical analyses of recombination rate

Genetic and non-genetic effects on the autosome-wide ("global") recombination rate were modelled by treating our experiment as an incomplete diallel between WSB/EiJ, PWK/PhJ and CAST/EiJ. The phenotype of interest is the number of autosomal crossovers per meiosis, and each N2 offspring represents an independent observation. Additive, parent-of-origin, strain-pair-specific, and age effects were estimated using the `BayesDiallel` package v0.982 (Lenarcic *et al.* 2012). The hierarchical Bayesian model implemented in this package can also include terms for inbreeding and dominance effects, but without phenotypes measured in inbred parental strains, these cannot be estimated. The Gibbs sampler was run for $10\,000$ iterations (the first $1\,000$ of which were discarded as burn-in), in 3 replicate chains.

Statistical analyses of crossover interference

Crossover interference was analyzed under two well-studied models: the gamma model, in which the strength of interference is expressed by a single unitless parameter, ν (Broman and Weber 2000); and the interference-

escape model of Housworth and Stahl (2003), in which a proportion p crossovers are not subject to interference, and the remaining $1 - p$ are subject to interference under the gamma model with parameter ν. We sought genotype-specific estimates of the strength of crossover interference, and allowed this effect to vary with paternal age. First we fit the gamma and interference-escape models to each subgroup (defined by paternal genotype, or the combination of paternal genotype and paternal age cohort) separately, using the maximum-likelihood approach implemented in the `xoi` package v0.72 (Broman 2023). Uncertainty in parameter estimates was obtained by performing 1000 bootstrap replicates, in which individuals were sampled randomly with replacement from within each group. This approach implicitly assumes no relationship between group-specific parameter values. We reasoned that the strength of interference is likely biologically constrained and that treating group-specific parameter values as being drawn from a common distribution would provide regularization that might lead to more stable and precise estimates. The following hierarchical Bayesian model was therefore implemented. Let ν_i be the group-wise interference, and define it as follows:

$$\log(\nu_i) = \beta_0 + \beta_i + \alpha \tag{1}$$

and define the following terms:

$$
\begin{aligned}
\beta_0 &= \text{intercept term} \\
\beta_i &= \text{genotype-specific effect } (i = 1, \ldots, k) \\
\alpha &= \text{age-specific effect}
\end{aligned}
\tag{2}
$$

The prior distributions for these parameters were $\beta_0 \sim \mathcal{N}(2.5, \sigma^2)$, and the remaining $\beta_i \sim \mathcal{N}(0, \sigma^2)$. The prior mean and variance were set to encompass the range of plausible values in mice, based on prior work (Broman *et al.* 2002). We experimented with other hyperparameter values and they had little impact on results. Similarly:

$$
\alpha \sim \begin{cases} 0 & \text{for young males} \\ \mathcal{N}(0, \sigma^2) & \text{for old males} \end{cases}
\tag{3}
$$

We implemented a simple Metropolis-coupled Markov chain Monte Carlo (MCMC) sampler to do inference of these parameters. The proposal distribution was $\mathcal{N}(0, \tau^2)$, and we found that $\tau^2 = 0.005$ allowed adequate mixing. The likelihood of a parameter vector $\theta = (\beta_0, \ldots, \beta_k, \lambda)$ given the complete set of haplotype segments was calculated with the C routine `GammaS()`, which is exposed in the `xoi` package. The sampler was run for $25\,000$ iterations (the first $5\,000$ of which were discarded as burn-in), in 4 replicate chains. Time required to compute the likelihood grows approxiately linearly in the number of haplotype segments, so runtime of

each MCMC chain was about one day. Convergence and mixing were assessed by manual inspection of trace plots and by Gelman-Rubin diagnostics as implemented in the `coda` package (Plummer *et al.* 2006). Out-of-sample predictive performance was assessed using smoothed importance sampling as implemented in the `loo` package, and influential outliers were defined as those with Pareto k scores ≥ 0.5 (Vehtari *et al.* 2017). The UNC Research Computing cluster was used for most computation.

Results

Recombination rate varies by genotype and age

The study design is shown in **Figure 1**. A total of 503 offspring of 17 males representing 6 reciprocal F1 crosses were collected and genotyped with SNP arrays. After iterative filtering and pruning, marker maps cover at least 97.8% of the physical span of the autosomes in each paternal genotype. The overall genotyping rate was 99.8%; no individual had more than 1.2% genotypes missing. We identified a total $15\,834$ haplotype segments that define $6\,271$ autosomal crossovers. On average 12.5 crossovers (range $5-22$) were transmitted in each male meiosis. Consistent with many prior studies, crossovers were concentrated in the distal ends of chromosomes (**Figure S1**).

The distributions of crossover count by genotype and paternal age are shown in **Figure 2**. The effects of interest are the influence of genotype and paternal age on recombination rate. As a preliminary step, we analyzed crossover counts under a generalized linear model (with Poisson response and log link) with terms for genotype, age, and genotype-by-age interaction. There is significant heterogeneity in recombination rate across genotypes ($p = 6.1 \times 10^{-9}$, likelihood ratio test). Qualitatively, paternal age appears to have a modest effect that differs by genotype. In a model allowing an age-by-genotype interaction, neither the main effect of age nor its interaction term reach statistical significance (**Table 3**). However, treating observations as conditionally independent, given genotype, ignores correlation structure that may be present when genotypes share parents. We therefore re-analyzed our experiment as an incomplete diallel between WSB/EiJ, PWK/PhJ and CAST/EiJ. Additive, parent-of-origin and cross-specific genetic effects were included in the model. Results are shown in **Figure 3**. Alleles from the PWK/PhJ background tend to increase the recombination rate, but having a PWK/PhJ mother decreases recombination rate by 0.51 (95% highest posterior density interval [HPDI] $0.28 - 0.73$) crossovers per meiosis. In this analysis, older males transmit 0.59 (95% HPDI $0.06 - 1.11$) more crossovers than younger males. These effects are modest: the contributions of additive, maternal and age effects are 13.8%, 8.4% and 0.8% of trait variance, respectively.

As expected, increase in recombination rate is associated with an increase in the number of chromosomes with multiple crossovers; there does not appear to be a threshold effect (**Figure S2**). The distribution of inter-

crossover distances on chromosomes with multiple crossovers is summarized in **Figure 4**, and plotted against the expected distribution in the absence of interference in **Figure S3**. Qualitatively, the distribution of inter-crossover distances strongly suggests the presence of interference, and appears to differ between genotypes. We explore this more rigorously in the next section. Similar to previous work (Broman *et al.* 2002), we find that inter-crossover distances are longer – implying stronger interference – on shorter chromosomes (**Figure S4**).

Strength of crossover interference varies by genotype

Crossover interference across all autosomes was quantified under the gamma model (McPeek and Speed 1995; Broman and Weber 2000). The gamma model describes the strength of interference via a single unitless parameter, ν; the distribution of intercrossover distances in this model has mean 50 cM and standard deviation $50/\sqrt{\nu}$ cM. We used two approaches for inference: first, using maximum likelihood to estimate ν_i separately in each genotype (for each of $i = 0, \ldots k$ genotypes), with confidence bounds obtained by nonparametric bootstrap; and second, using a hierarchical Bayesian model to estimate all ν_i jointly. The maximum likelihood approach has been published previously (Broman *et al.* 2002); the Bayesian method is novel but uses the same code for computing the likelihood. The two methods provide very similar estimates (root mean square difference = 0.44, or 4.7% of the smallest value of ν_i), as shown in **Figure S5**.

We first evaluate the strength of evidence for heterogeneity of crossover interference across genotypes. This evaluation amounts to comparing the fit of a model with separate values of ν_i for each genotype (full model) to the fit of a model with with a single common ν for all individuals (null model). In the maximum likelihood framework, the full model is superior to the null model ($p = 0.038$ by likelihood-ratio test). The better fit of the full model comes at the expense of model complexity; using the Akaike information criterion (AIC), which penalizes the likelihood by the number of parameters (Akaike 1998), the improvement in fit is modest ($\Delta \mathrm{AIC} = 1.8$; see **Table 5**). In the Bayesian framework, a common approach to model selection is to assess out-of-sample predictive performance, which implicitly penalizes model complexity. The "expected log pointwise predictive density" for new observations (ELPD) provides one measure of predictive accuracy and can be estimated by cross-validation or by importance sampling (Vehtari *et al.* 2017). We take the latter approach as it is more computationally efficient. Results are presented in **Table 6**. The difference in estimated predictive accuracy of the full versus the null model is similar and well within their respective standard errors. However, out-of-sample predictive performance is sensitive to influential outliers, which are more prevalent under the full model than under the null model. Taking the Bayesian and frequentist analysis together, we conclude that our data is consistent with heterogeneity in strength of interference across genotypes, but the magnitude of these effects is challenging to estimate precisely with the current sample size.

Estimates of the strength of interference by genotype and paternal age are shown in **Figure 4**. Our es-

timates are close to the 11.3 obtained by Broman *et al.* (2002), the only other published application of the gamma model to house mice, although that study used an interspecies backcross between C57BL/6J and *Mus spretus*. The largest and smallest ν values are obtained from the reciprocal crosses PWK/PhJ×WSB/EiJ and WSB/EiJ×PWK/PhJ, respectively. Contrasts between genotypes were calculated on the log scale (since $\nu > 0$ by definition) and therefore represent ratios between values of ν when transformed back to the natural scale (**Figure S6**). Three comparisons reach conventional levels of statistical significance: PWK/PhJ×WSB/EiJ vs WSB/EiJ×PWK/PhJ, PWK/PhJ×CAST/EiJ vs WSB/EiJ×PWK/PhJ and CAST/EiJ×PWK/PhJ vs WSB/EiJ×PWK/PhJ. Given the evidence for parent-of-origin effects on recombination rate, we focused particular attention on differences in strength of interference in reciprocal pairs. Interference is 1.28 times (95% HPDI $1.02 - 1.59$) stronger in PWK/PhJ×WSB/EiJ than WSB/EiJ×PWK/PhJ; there is little difference between the other reciprocal pairs (**Figure S7**).

The effect of age on strength of interference is more modest: interference is 1.13 times (95% HPDI $0.91 - 1.13$) stronger in older versus younger males. The null value falls at the 14^{th} percentile of the posterior distribution.

Interference escape varies by genotype

Although the gamma model provides a succinct statistical description of intercrossover distances, it is known that some proportion of crossovers are not subject to interference (reviewed by **?**.) The Housworth-Stahl, or gamma-escape, model allows a proportion p of non-interfering crossovers, with the remaining $1 - p$ subject to the gamma model of interference with strength ν (Housworth and Stahl 2003).Intuitively we expect that a wide range of (p, ν) pairs might be compatible with a given set of haplotype intervals – if a large number of crossovers escape interference (and can therefore be placed close together), then the strength of interference inferred from the remaining intercrossover distances will be higher. We fit this model to our data by maximum likelihood, first to the entire dataset (null model) and then separately for each genotype (full model). (The computation time of the Bayesian method applied to this model was prohibitive.) The model fit with genotype-specific values of (p_i, ν_i) is superior to the model in which all genotypes share the same values ($p = 0.0023$ by likelihood-ratio test, ΔAIC $= 7.3$; see **Table 7**).

Inspection of parameter estimates from bootstrap replicates confirms our intuition that the strength of interference and escape fraction are challenging to estimate independently (**Figure S8**). This observation implies non-identifiability that undercuts the biological validity of any between-group comparisons of parameter values. With this limitation in mind, parameter estimates from the gamma and gamma-escape models are shown in **Figure 5**. The inferred proportion of non-interfering crossovers ranges from zero (in CAST/EiJ×PWK/PhJ) to 0.20 (in WSB/EiJ×CAST/EiJ). Estimates of the strength of interference are higher when escape is allowed, by a factor of up to 3.1 times the value without escape (in WSB/EiJ×CAST/EiJ).

Discussion

Our portrait of male recombination obtained from crosses between wild-derived strains representing three subspecies of house mice corroborates and extends previous work in this area. We find that alleles from the *M. m. musculus* strain PWK/PhJ tend to increase recombination, whereas alleles from the *M. m. castaneus* strain CAST/EiJ tend to decrease it. This finding is consistent with autosomal QTL detected in CAST/EiJ×C57BL/6J (Murdoch *et al.* 2010) and CAST/EiJ×PWD/PhJ (Dumont and Payseur 2011b) intercrosses – QTL at which CAST/EiJ alleles reduce the number of crossovers. (PWD/PhJ is another *M. m. musculus* wild-derived strain that is closely related to PWK/PhJ (Gregorová and Forejt 2000).) In addition, autosomal alleles from PWK/PhJ are associated with more recombination and those from CAST/EiJ are associated with lower recombination in the early generations of the Collaborative Cross (Liu *et al.* 2014). The decrease in recombination connected to PWK/PhJ mothers and the increase in recombination connected to CAST/EiJ mothers in this study are also consistent with previous inferences of X-linked QTL with effects in the same directions (Murdoch *et al.* 2010; Dumont and Payseur 2011b; Liu *et al.* 2014).

Our results support the conclusion that crossover interference is strong in mice (Broman *et al.* 2002; de Boer *et al.* 2006). The shape parameter ν in the gamma model determines whether the distribution of distances between crossovers is exponential ($\nu = 1$), corresponding to independence among crossovers, or humped ($\nu > 1$), indicating that crossovers are spaced more regularly along the chromosome. Although values of ν are continuous, $\nu - 1$ is equivalent to the number of recombination intermediates that are located between two crossovers in the counting model of interference (Foss *et al.* 1993; Otto and Payseur 2019). Hence, our estimates of ν suggest that on average, crossovers are separated by 10 or more recombination intermediates that are repaired as non-crossovers in three subspecies of mice. Furthermore, the superior fit of the gamma-escape model to our data is in line with the existence of both interfering and non-interfering crossovers in mice (de Boer *et al.* 2006) as well as several other species (Otto and Payseur 2019).

The primary source of novelty in our study is the demonstration that the level of crossover interference varies among genotypes. Given that crosses were conducted in a common environment, heterogeneity among F1s detected using the gamma and gamma-escape models implies heritable differences in the level of interference and possibly the proportion of crossovers that avoid interference. The largest contrast comes from reciprocal crosses involving *M. m. musculus* (PWK/PhJ) and *M. m. domesticus* (WSB/EiJ). Although other maternal effects might contribute, stronger interference in PWK/PhJ×WSB/EiJ F1s than in WSB/EiJ×PWK/PhJ F1s raises the possibility that the X chromosome modulates interference. This explanation is interesting in light of reports that PWK/PhJ and PWD/PhJ alleles on the X chromosome reduce recombination in other crosses (Dumont and Payseur 2011b; Liu *et al.* 2014), reductions that could be accomplished by amplifying interference. We also estimated the level of interference to be higher in PWK/PhJ×CAST/EiJ F1s than CAST/EiJ×PWK/PhJ F1s,

though the difference is small. Another informative pairwise comparison is WSB/EiJ×PWK/PhJ F1s (lower interference) and CAST/EiJ×PWK/PhJ F1s (higher interference). These F1s differ by one set of autosomes as well as the X chromosome. This last comparison suggests that alleles from CAST/EiJ increase interference, again directionally consistent with the lower rate of recombination in CAST/EiJ than in WSB/EiJ and PWD/PhJ (Dumont and Payseur 2011a; Peterson and Payseur 2021). However, at least among F1 genotypes, there is not a monotonic relationship between recombination rate and strength of interference (**Figure S9**).

Although parameter estimates associated with each cross should be treated with caution, our observation that the level of interference depends on genetic background provides evidence for evolution of this important determinant of crossover positioning. The three subspecies of mice to which WSB/EiJ, PWK/PhJ, and CAST/EiJ belong are estimated to have begun diverging about $350\,000 - 500\,000$ years ago (reviewed in Phifer-Rixey and Nachman (2015)). To the extent that the wild-derived strains we examined are representative of their subspecies, this divergence time suggests the evolution of interference can occur over relatively short timescales.

The cellular and molecular causes of the evolution of interference we document are unclear. Multiple mechanisms of interference have been proposed (Otto and Payseur 2019). Recent discoveries in *Arabidopsis* species suggest that interference is accomplished by diffusion-mediated growth of evenly spaced foci of the ubiquitin ligase HEI10 at the expense of closely spaced foci along the synaptonemal complex (Morgan *et al.* 2021; Durand *et al.* 2022; Fozard *et al.* 2023). Whether a similar "coarsening" mechanism operates in mice and contributes to the evolution of interference remains to be seen.

Another novel inference from our study is that older mouse males (at least 15 months of age) transmit an average of 0.59 more crossovers per meiosis than younger males (no more than 4 months of age). Although this effect is quite modest, it is consistent in magnitude and direction with a previous study in two classical inbred strains that found about 2 more interfering crossovers per spermatocyte in 1-year-old males versus males undergoing the first wave of spermatogenesis at 20 days postpartum (Vrooman *et al.* 2014). The effects of age on recombination rate in mammals are not well established in general. Studies reported no effect of paternal age in humans (Hussin *et al.* 2011) or sheep (Johnston *et al.* 2016). Maternal age has been associated with increased recombination in humans (Kong *et al.* 2004; Hussin *et al.* 2011; Campbell *et al.* 2015; Martin *et al.* 2015) and pigs (Lozada-Soto *et al.* 2021), decreased recombination in house mice (Polani and Jagiello 1976), increased and decreased recombination in cattle (Shen *et al.* 2021), and no change in recombination in sheep (Johnston *et al.* 2016). We also find that interference is about 9% stronger in older males, although the effect does not meet the usual nominal threshold for significance. Our results suggest that paternal age deserves further investigation as a modulator of crossover number and positioning.

We attach several caveats to our findings. First, distances between crossovers were computed by assum-

ing that the standard genetic map applies to all F1s, which could affect our characterization of interference. If the true recombination rate is higher in a given genotype than in the common map, we would expect more crossovers per interval than predicted by the map and downstream estimates of the strength of interference would be biased downward. As shown in **Figure S9**, we do not find a consistent relationship between interference and recombination rate across genotypes. Second, our analyses are based on transmitted genotypes in live progeny, and therefore reflect the recombination landscape in viable gametes only. Certain crosses between *M. m. musculus* mothers and *M. m. domesticus* fathers generate F1 hybrid males with signs of reduced fertility caused by incompatibilities that disrupt meiosis I (Forejt and Iványi 1974; Forejt 1996; White *et al.* 2012; Schwahn *et al.* 2018). If interference is systematically different in spermatocytes that are eliminated by meiotic checkpoints, or in gametes that do not mature into functional sperm, then our estimates of interference parameters will be biased. At this point there is no reason to suspect that the transmitted crossovers we surveyed comprise a non-random subset. Finally and most importantly, our study is restricted to males. Differences in interference between females and males in mice (Peterson and Payseur 2021) and other species (Basu-Roy *et al.* 2013; Campbell *et al.* 2016; Wang *et al.* 2016; Torgasheva and Borodin 2017) predict distinct evolutionary dynamics for interference in the two sexes.

Acknowledgements

This project was launched in the laboratory of Fernando Pardo-Manuel de Villena in 2012 as an outgrowth of existing studies of aging and male infertility. Support was provided by the following grants from the National Institutes of Health: F30 MH103925 (APM), K01 MH094406 (James J Crowley), R21 MH096261 (Fernando Pardo-Manuel de Villena), R01 HD065024 (Fernando Pardo-Manuel de Villena), P50 MH090338 (Fernando Pardo-Manuel de Villena), P50 HG006582 (Fernando Pardo-Manuel de Villena), U42 OD010924 (Terry Magnuson), and R35 GM139412 (Bret Payseur). We thank Timothy A Bell and James J Crowley for mouse husbandry. We thank Grace Clark, Justin Gooch, the late Mark Calaway, and Darla Miller for assistance with DNA preparation and submission of samples for genotyping. We benefited from fruitful correspondence with Fernando Pardo-Manuel de Villena, Beth Dumont, Karl Broman and several other members of the mouse genetics community in the course of this work.

References

Akaike, H., 1998 Information theory and an extension of the maximum likelihood principle. In *Selected Papers of Hirotugu Akaike*, edited by E. Parzen, K. Tanabe, and G. Kitagawa, pp. 199–213, Springer New York, New York, NY.

Auxier, B., A. J. M. Debets, F. A. Stanford, J. Rhodes, F. M. Becker, *et al.*, 2023 The human fungal pathogen aspergillus fumigatus can produce the highest known number of meiotic crossovers. PLoS Biol. **21**: e3002278.

Barton, N. H., 1995 A general model for the evolution of recombination. Genet. Res. **65**: 123–145.

Basu-Roy, S., F. Gauthier, L. Giraut, C. Mézard, M. Falque, *et al.*, 2013 Hot regions of noninterfering crossovers coexist with a nonuniformly interfering pathway in arabidopsis thaliana. Genetics **195**: 769–779.

Berchowitz, L. E. and G. P. Copenhaver, 2010 Genetic interference: don't stand so close to me. Curr. Genomics **11**: 91–102.

Brazier, T. and S. Glémin, 2022 Diversity and determinants of recombination landscapes in flowering plants. PLoS Genet. **18**: e1010141.

Brekke, C., P. Berg, A. B. Gjuvsland, and S. E. Johnston, 2022 Recombination rates in pigs differ between breeds, sexes and individuals, and are associated with the RNF212, SYCP2, PRDM7, MEI1 and MSH4 loci. Genet. Sel. Evol. **54**: 33.

Broman, K. W., 2023 *xoi: Tools for Analyzing Crossover Interference*.

Broman, K. W., D. M. Gatti, P. Simecek, N. A. Furlotte, P. Prins, *et al.*, 2019 R/qtl2: Software for mapping quantitative trait loci with High-Dimensional data and multiparent populations. Genetics **211**: 495–502.

Broman, K. W., L. B. Rowe, G. A. Churchill, and K. Paigen, 2002 Crossover interference in the mouse. Genetics **160**: 1123–1131.

Broman, K. W. and J. L. Weber, 2000 Characterization of human crossover interference. Am. J. Hum. Genet. **66**: 1911–1926.

Caballero, M., D. N. Seidman, Y. Qiao, J. Sannerud, T. D. Dyer, *et al.*, 2019 Crossover interference and sex-specific genetic maps shape identical by descent sharing in close relatives. PLoS Genet. **15**: e1007979.

Campbell, C. L., C. Bhérer, B. E. Morrow, A. R. Boyko, and A. Auton, 2016 A Pedigree-Based map of recombination in the domestic dog genome. G3 .

Campbell, C. L., N. A. Furlotte, N. Eriksson, D. Hinds, and A. Auton, 2015 Escape from crossover interference increases with maternal age. Nat. Commun. **6**: 6260.

Chinnici, J. P., 1971 Modification of recombination frequency in drosophila. i. selection for increased and decreased crossing over. Genetics **69**: 71–83.

Coop, G., X. Wen, C. Ober, J. K. Pritchard, and M. Przeworski, 2008 High-resolution mapping of crossovers reveals extensive variation in fine-scale recombination patterns among humans. Science **319**: 1395–1398.

Cox, A., C. L. Ackert-Bicknell, B. L. Dumont, Y. Ding, J. T. Bell, *et al.*, 2009 A new standard genetic map for the laboratory mouse. Genetics **182**: 1335–1344.

de Boer, E., P. Stam, A. J. J. Dietrich, A. Pastink, and C. Heyting, 2006 Two levels of interference in mouse meiotic recombination. Proc. Natl. Acad. Sci. U. S. A. **103**: 9607–9612.

Dumont, B. L. and B. A. Payseur, 2011a Evolution of the genomic recombination rate in murid rodents. Genetics **187**: 643–657.

Dumont, B. L. and B. A. Payseur, 2011b Genetic analysis of Genome-Scale recombination rate evolution in house mice. PLoS Genet. **7**: e1002116.

Durand, S., Q. Lian, J. Jing, M. Ernst, M. Grelon, *et al.*, 2022 Joint control of meiotic crossover patterning by the synaptonemal complex and HEI10 dosage. Nat. Commun. **13**: 5999.

Felsenstein, J., 1974 The evolutionary advantage of recombination. Genetics **78**: 737–756.

Fisher, R. A., 1930 *The genetical theory of natural selection*. Oxford, Clarendon Press.

Forejt, J., 1996 Hybrid sterility in the mouse. Trends Genet. **12**: 412–417.

Forejt, J. and P. Iványi, 1974 Genetic studies on male sterility of hybrids between laboratory and wild mice (mus musculus l.). Genet. Res. **24**: 189–206.

Foss, E., R. Lande, F. W. Stahl, and C. M. Steinberg, 1993 Chiasma interference as a function of genetic distance. Genetics **133**: 681–691.

Fozard, J. A., C. Morgan, and M. Howard, 2023 Coarsening dynamics can explain meiotic crossover patterning in both the presence and absence of the synaptonemal complex. Elife **12**.

Gray, S. and P. E. Cohen, 2016 Control of meiotic crossovers: From Double-Strand break formation to designation. Annu. Rev. Genet. **50**: 175–210.

Gregorová, S. and J. Forejt, 2000 PWD/Ph and PWK/Ph inbred mouse strains of mus m. musculus subspecies–a valuable resource of phenotypic variations and genomic polymorphisms. Folia Biol. **46**: 31–41.

Hassold, T. and P. Hunt, 2001 To err (meiotically) is human: the genesis of human aneuploidy. Nat. Rev. Genet. **2**: 280–291.

Hill, W. G. and A. Robertson, 1966 The effect of linkage on limits to artificial selection. Genet. Res. **8**: 269–294.

Housworth, E. A. and F. W. Stahl, 2003 Crossover interference in humans. Am. J. Hum. Genet. **73**: 188–197.

Hunter, C. M., W. Huang, T. F. C. Mackay, and N. D. Singh, 2016 The genetic architecture of natural variation in recombination rate in drosophila melanogaster. PLoS Genet. **12**: e1005951.

Hussin, J., M.-H. Roy-Gagnon, R. Gendron, G. Andelfinger, and P. Awadalla, 2011 Age-Dependent recombination rates in human pedigrees. PLoS Genet. **7**: e1002251.

Johnston, S. E., C. Bérénos, J. Slate, and J. M. Pemberton, 2016 Conserved genetic architecture underlying individual recombination rate variation in a wild population of soay sheep (ovis aries). Genetics **203**: 583–598.

Kidwell, M. G. and J. F. Kidwell, 1976 Selection for male recombination in drosophila melanogaster. Genetics **84**: 333–351.

Kong, A., J. Barnard, D. F. Gudbjartsson, G. Thorleifsson, G. Jonsdottir, *et al.*, 2004 Recombination rate and reproductive success in humans. Nat. Genet. **36**: 1203–1206.

Kong, A., G. Thorleifsson, H. Stefansson, G. Masson, A. Helgason, *et al.*, 2008 Sequence variants in the RNF212 gene associate with genome-wide recombination rate. Science **319**: 1398–1401.

Lam, I. and S. Keeney, 2014 Mechanism and regulation of meiotic recombination initiation. Cold Spring Harb. Perspect. Biol. **7**: a016634.

Lenarcic, A. B., K. L. Svenson, G. A. Churchill, and W. Valdar, 2012 A general bayesian approach to analyzing diallel crosses of inbred strains. Genetics **190**: 413–435.

Liu, E. Y., A. P. Morgan, E. J. Chesler, W. Wang, G. A. Churchill, *et al.*, 2014 High-Resolution Sex-Specific linkage maps of the mouse reveal polarized distribution of crossovers in male germline. Genetics **197**: 91–106.

Lozada-Soto, E. A., C. Maltecca, H. Wackel, W. Flowers, K. Gray, *et al.*, 2021 Evidence for recombination variability in purebred swine populations. J. Anim. Breed. Genet. **138**: 259–273.

Martin, H. C., R. Christ, J. G. Hussin, J. O'Connell, S. Gordon, *et al.*, 2015 Multicohort analysis of the maternal age effect on recombination. Nat. Commun. **6**: 7846.

McPeek, M. S. and T. P. Speed, 1995 Modeling interference in genetic recombination. Genetics **139**: 1031–1044.

Morgan, A. P., 2016 argyle : An R package for analysis of illumina genotyping arrays. G3 **6**: 281–286.

Morgan, A. P., T. A. Bell, J. J. Crowley, and F. Pardo-Manuel de Villena, 2019 Instability of the pseudoautosomal boundary in house mice. Genetics **212**: 469–487.

Morgan, A. P., C.-P. Fu, C.-Y. Kao, C. E. Welsh, J. P. Didion, *et al.*, 2016 The mouse universal genotyping array: From substrains to subspecies. G3 **6**: 263–279.

Morgan, C., M. A. White, F. C. H. Franklin, D. Zickler, N. Kleckner, *et al.*, 2021 Evolution of crossover interference enables stable autopolyploidy by ensuring pairwise partner connections in arabidopsis arenosa. Curr. Biol. **31**: 4713–4726.e4.

Munz, P., 1994 An analysis of interference in the fission yeast schizosaccharomyces pombe. Genetics **137**: 701–707.

Murdoch, B., N. Owen, S. Shirley, S. Crumb, K. W. Broman, *et al.*, 2010 Multiple loci contribute to genome-wide recombination levels in male mice. Mamm. Genome **21**: 550–555.

Otto, S. P. and B. A. Payseur, 2019 Crossover interference: Shedding light on the evolution of recombination. Annu. Rev. Genet. **53**: 19–44.

Paigen, K. and P. Petkov, 2010 Mammalian recombination hot spots: properties, control and evolution. Nat. Rev. Genet. **11**: 221–233.

Paigen, K., J. P. Szatkiewicz, K. Sawyer, N. Leahy, E. D. Parvanov, *et al.*, 2008 The recombinational anatomy of a mouse chromosome. PLoS Genet. **4**: e1000119.

Parvanov, E. D., P. M. Petkov, and K. Paigen, 2010 Prdm9 controls activation of mammalian recombination hotspots. Science **327**: 835–835.

Peterson, A. L. and B. A. Payseur, 2021 Sex-specific variation in the genome-wide recombination rate. Genetics **217**: 1–11.

Petkov, P. M., K. W. Broman, J. P. Szatkiewicz, and K. Paigen, 2007 Crossover interference underlies sex differences in recombination rates. Trends Genet. **23**: 539–542.

Phifer-Rixey, M. and M. W. Nachman, 2015 Insights into mammalian biology from the wild house mouse mus musculus. Elife **4**: e05959.

Plummer, M., N. Best, K. Cowles, and K. Vines, 2006 CODA: Convergence diagnosis and output analysis for MCMC.

Polani, P. E. and G. M. Jagiello, 1976 Chiasmata, meiotic univalents, and age in relation to aneuploid imbalance in mice. Cytogenet. Cell Genet. **16**: 505–529.

Schwahn, D. J., R. J. Wang, M. A. White, and B. A. Payseur, 2018 Genetic dissection of hybrid male sterility across stages of spermatogenesis. Genetics **210**: 1453–1465.

Shen, B., E. Freebern, J. Jiang, C. Maltecca, J. B. Cole, *et al.*, 2021 Effect of temperature and maternal age on recombination rate in cattle. Front. Genet. **12**: 682718.

Sigmon, J. S., M. W. Blanchard, R. S. Baric, T. A. Bell, J. Brennan, *et al.*, 2020 Content and performance of the MiniMUGA genotyping array: A new tool to improve rigor and reproducibility in mouse research. Genetics **216**: 905–930.

Smukowski, C. S. and M. A. F. Noor, 2011 Recombination rate variation in closely related species. Heredity **107**: 496–508.

Stapley, J., P. G. D. Feulner, S. E. Johnston, A. W. Santure, and C. M. Smadja, 2017 Variation in recombination frequency and distribution across eukaryotes: patterns and processes. Philos. Trans. R. Soc. Lond. B Biol. Sci. **372**.

Steemers, F. J., W. Chang, G. Lee, D. L. Barker, R. Shen, *et al.*, 2006 Whole-genome genotyping with the single-base extension assay. Nat. Methods **3**: 31–33.

Strickland, W. N., 1958 An analysis of interference in aspergillus nidulans. Proceedings of the Royal Society of London. Series B - Biological Sciences **149**: 82–101.

Sturtevant, A. H., 1913 The linear arrangement of six sex-linked factors in drosophila, as shown by their mode of association. J. Exp. Zool. **14**: 43–59.

Sturtevant, A. H., 1915 The behavior of the chromosomes as studied through linkage. Z. Indukt. Abstamm. Vererbungsl. **13**: 234–287.

Torgasheva, A. A. and P. M. Borodin, 2017 Immunocytological analysis of meiotic recombination in the gray goose (anser anser). Cytogenet. Genome Res. **151**: 27–35.

Vehtari, A., A. Gelman, and J. Gabry, 2017 Practical bayesian model evaluation using leave-one-out cross-validation and WAIC. Stat. Comput. **27**: 1413–1432.

Veller, C., N. B. Edelman, P. Muralidhar, and M. A. Nowak, 2020 Variation in genetic relatedness is determined by the aggregate recombination process. Genetics **216**: 985–994.

Vrooman, L. A., S. I. Nagaoka, T. J. Hassold, and P. A. Hunt, 2014 Evidence for paternal Age-Related alterations in meiotic chromosome dynamics in the mouse. Genetics **196**: 385–396.

Wang, R. J., B. L. Dumont, P. Jing, and B. A. Payseur, 2019 A first genetic portrait of synaptonemal complex variation. PLoS Genet. **15**: e1008337.

Wang, Z., B. Shen, J. Jiang, J. Li, and L. Ma, 2016 Effect of sex, age and genetics on crossover interference in cattle. Sci. Rep. **6**: 37698.

White, M. A., M. Stubbings, B. L. Dumont, and B. A. Payseur, 2012 Genetics and evolution of hybrid male sterility in house mice. Genetics **191**: 917–934.

Data availability

Sample metadata, raw and processed genotype data are available on Figshare: doi:10.6084/m9.figshare.25859950.

Analysis code is available on Github: `https://github.com/andrewparkermorgan/mouse_crossover_interference`.

Author contributions

APM designed and performed the analyses. APM and BAP and wrote the manuscript.

Competing interests

None of the authors have any competing financial interests in the work described in this manuscript.

Figure 1: Incomplete diallel design to study effects of age and genetics on recombination rate. (**A**) All reciprocal F1 genotypes between three parental strains representing the cardinal subspecies of *Mus musculus*. (**B**) F1 males mated at young age (< 4 months) and old age (>15 months) to highly fertile FVB/NJ females. (**C**) Number of progeny collected for each paternal genotype.

Figure 2: Recombination rate by genotype and age. (**A**) Number of transmitted crossovers per meiosis. Each point is an individual observation; solid dots show group means and bars show $2.5\% - 97.5\%$ (thin) and $25\% - 75\%$ (thick) quantile intervals. (**B**) Bootstrap estimates of the length of the autosomal recombination map in each genotype. Solid dots show group means; error bars defined as in panel **A**.

Figure 3: Diallel analysis of recombination rate. (**A**) Posterior distribution of estimated additive and maternal genotype effects (see main text) and age effect. Solid dots show group means and bars show $2.5\% - 97.5\%$ (thin) and $25\% - 75\%$ (thick) quantile intervals. (**B**) Posterior distribution of trait variance attributed to each class of genetic effects. Note square-root scale on horizontal axis. Quantile intervals indicated as in panel **A**.

Figure 4: Posterior distributions of interference parameter (ν) by genotype, averaged over age groups; and by age group, averaged over genotypes. Solid dots and text labels show posterior mean; bars show $2.5\% - 97.5\%$ (thin) and $25\% - 75\%$ (thick) quantile intervals. Dashed line shows previous estimate from Broman *et al.* (2002). Note that the horizontal axis is in log scale.

Figure 5: Interference escape may vary by genotype. (**A**) Estimates of strength of crossover interference by genotype, under the gamma and gamma-escape (Houseworth-Stahl) models. Error bars show $2.5\% - 97.5\%$ (thin) and $25\% - 75\%$ (thick) bootstrap confidence intervals. (**B**) Estimates of proportion of crossovers escaping interference by genotype. Error bars extend to the margin of the chart when confidence intervals include zero. (**C**) Apparent positive correlation between p and ν across genotypes.

F0 male	paternal age	platform	F0 female WSB/EiJ	PWK/PhJ	CAST/EiJ
WSB/EiJ	young	mega	–	14	0
		mini	–	0	8
	old	mega	–	12	0
		mini	–	0	52
PWK/PhJ	young	mega	48	–	9
		mini	0	–	0
	old	mega	47	–	7
		mini	0	–	45
CAST/EiJ	young	mega	0	105	–
		mini	31	0	–
	old	mega	0	75	–
		mini	11	39	–

Table 1: Counts of N2 progeny by genotype, paternal age, and genotyping platform ("mega" = MegaMUGA, "mini" = MiniMUGA).

genotype	platform	markers	Mb covered	n
CASTxPWK	mega	1940	2352	16
	mini	2059	2354	45
PWKxCAST	mega	1940	2352	180
	mini	2059	2354	39
CASTxWSB	mega	–	–	0
	mini	2712	2359	60
WSBxCAST	mega	–	–	0
	mini	2712	2359	42
PWKxWSB	mega	32010	2393	26
	mini	–	–	0
WSBxPWK	mega	32005	2393	95
	mini	–	–	0

Table 2: Density and coverage of pruned marker maps by genotype and platform ("mega" = MegaMUGA, "mini" = MiniMUGA, n = number of N2 progeny genotyped).

	df	deviance	residual df	residual deviance	p
genotype	5	46.84	497	318.00	6.1×10^{-9}
age	1	3.02	496	314.97	0.08
genotype $\times$ age	5	7.09	491	307.89	0.21
residuals	–	–	502	364.84	–

Table 3: Analysis of deviance for model of crossover counts. The p-values shown are from a likelihood-ratio test. df = degrees of freedom.

genotype	mean	median	2.5%ile	25%ile	75%ile	97.5%ile
CAST/EiJ×PWK/PhJ	51.3	50.0	26.6	40.2	62.5	79.3
PWK/PhJ×CAST/EiJ	54.2	53.3	28.7	44.6	64.6	81.8
CAST/EiJ×WSB/EiJ	54.4	55.3	27.3	46.2	62.8	82.5
WSB/EiJ×CAST/EiJ	60.3	61.6	34.7	52.1	69.5	80.2
PWK/PhJ×WSB/EiJ	51.9	52.0	33.9	43.7	60.4	73.7
WSB/EiJ×PWK/PhJ	53.4	53.5	24.1	44.9	63.3	81.9

Table 4: Distribution of inter-crossover distances (in cM) by cross.

model	df	LL	AIC	χ^2	p
full	6	-5888.4	11788.8	11.8	0.038
null	1	-5894.3	11790.5	–	–

Table 5: Comparison of gamma models fit by maximum likelihood, allowing strength of crossover interference to vary by genotype (full) or not (null). df = degrees of freedom, LL = log-likelihood, AIC = Akaike information criterion

model	ELPD	std error	% influential obs
full	-5864.3	56.8	8.7%
null	-5862.0	58.4	1.2%

Table 6: Comparison of gamma models fit by MCMC, allowing strength of crossover interference to vary by genotype (full) or not (null). ELPD = expected log pointwise posterior density

model	df	LL	AIC	χ^2	p
full	12	-5854.2	11732.4	27.3	0.0023
null	2	-5867.8	11739.6	–	–

Table 7: Comparison of gamma-escape models fit by maximum likelihood, allowing strength of crossover interference and escape fraction to vary by genotype (full) or not (null). df = degrees of freedom, LL = log-likelihood, AIC = Akaike information criterion

Supplementary material

Table S1. Sample metadata for $n = 503$ N2 progeny of F1 fathers.

Table S2. All inferred autosomal haplotype segments ($m = 15\,834$) for $n = 503$ mice.

Table S3. Number of autosomal crossovers inferred per individual for $n = 503$ mice.

Raw and processed genotype matrices are available from Figshare.

Figure S1: Recombination rate (in cM/Mb) in 5 Mb sliding windows across the autosomes, estimated separately for young and old males.

Figure S2: The number of multiply-recombinant chromosomes increases smoothly with recombination rate. (**A**) Distribution of chromosomes by number of crossovers in each genotype-by-age combination. (**B**) Count of chromosomes with multiple crossovers versus total number of crossovers in each genotype-by-age combination. Error bars are 95% confidence intervals from generalized linear model with genotype, age and their interaction as covariates. Solid grey line is a linear fit through the group means.

Figure S3: Empirical cumulative distribution of distance between crossovers (COs) on the same chromosome (blue curves), compared to expected distribution without crossover interference.

Figure S4: Distribution of inter-crossover distance by chromosome, for chromosomes on which multiply-recombinant progeny were obtained, plotted by chromosome number (**A**) and against total chromosome length (**B**). Solid dots are medians; bars show $2.5\% - 97.5\%$ (thin) and $25\% - 75\%$ (thick) quantile intervals. Solid grey line is a linear fit through the per-chromosome medians.

Figure S5: Comparison of group-wise estimates of the interference parameter ν obtained by maximum-likelihood versus joint estimates from Bayesian hierarchical model by MCMC. Solid grey line is a linear fit through the point estimates; dashed grey line has unit slope and passes through the origin.

Figure S6: All pairwise contrasts between per-genotype estimates of strength of interference. Dots show posterior means and bars show $2.5\% - 97.5\%$ (thin) and $25\% - 75\%$ (thick) quantile intervals. Open dots indicate comparisons that meet nominal statistical significance; filled dots nominally non-significant. Dashed line gives the null value of 1. Note that the horizontal axis is in log scale.

Figure S7: Contrasts between estimates of strength of interference in reciprocal genotypes. Text labels show posterior means. Dashed line gives the null value of 1. Note that the horizontal axis is in log scale.

Figure S8: Maximum-likelihood estimates for the strength of interference (ν) and proportion of non-interfering crossovers (p) for 1000 bootstrap replicates of crossover data from six genotypes. Note that horizontal axis is in log scale and vertical axis in square-root scale.

Figure S9: Relationship between length of the recombination map (in cM) and strength of crossover interference (ν). Error bars give 95% confidence bounds, calculated as described for the respective parameters elsewhere.