Alzheimer's disease-associated protective variant Plcg2-P522R modulates peripheral macrophage function in a sex-dimorphic manner

Hannah A. Staley[1,2,3], Janna Jernigan[1,2,3], MacKenzie L. Bolen[1,2,3], Ann M. Titus[1,2,3], Noelle Neighbarger[1,2,3], Cassandra Cole[1,2,3], Kelly B. Menees[1,2,3], Rebecca L. Wallings[1,2,3], Malú Gámez Tansey[1,2,3,4]*

[1] Department of Neuroscience, University of Florida College of Medicine, Gainesville, Florida, USA
[2] Center for Translational Research in Neurodegenerative Disease, University of Florida College of Medicine, Gainesville, Florida, USA
[3] McKnight Brain Institute, University of Florida, Gainesville, Florida, USA
[4] Norman Fixel Institute for Neurological Diseases, University of Florida Health, Gainesville, Florida, USA

*Corresponding Author
Malú Gámez Tansey, PhD
mgtansey@UFL.edu

ABSTRACT

Genome-wide association studies have identified a protective mutation in the phospholipase C

gamma 2 (*PLCG2*) gene which confers protection against Alzheimer's disease (AD)-associated

cognitive decline. Therefore, PLCG2, which is primarily expressed in immune cells, has become

a target of interest for potential therapeutic intervention. The protective allele, known as *P522R*,

has been shown to be hyper-morphic in microglia, increasing phagocytosis of amyloid-beta (Aβ),

and increasing the release of inflammatory cytokines. However, the effect of this protective

mutation on peripheral tissue-resident macrophages, and the extent to which sex modifies this

effect, has yet to be assessed. Herein, we show that peripheral macrophages carrying the *P522R*

mutation do indeed show functional differences compared to their wild-type (WT) counterparts,

however, these alterations occur in a sex-dependent manner. In macrophages from females, the

P522R mutation increases lysosomal protease activity, cytokine secretion, and gene expression

associated with cytokine secretion and apoptosis. In contrast, in macrophages from males, the

mutation causes decreased phagocytosis and lysosomal protease activity, modest increases in

cytokine secretion, and induction of gene expression associated with negative regulation of the

immune response. Taken together, these results suggest that the mutation may be conferring

different effects dependent on sex and cell type, and highlight the importance of considering sex

as a biological variable when assessing the effects of genetic variants and implications for

potential immune system-targeted therapies.

Keywords: Alzheimer's disease, macrophage, PLCG2, sex differences

BACKGROUND

Alzheimer's disease (AD) is a devastating neurodegenerative disorder that afflicts millions of people worldwide over two-thirds of which are women (1). Classically, AD is histologically characterized by the presence of amyloid-beta (Aβ) plaques extracellularly, and the formation of tau tangles within neurons. However, over recent years, the immune system has been implicated in the pathogenesis and progression of the disease. Of particular interest are cells of the innate immune system, which includes both peripheral monocytes and macrophages, and the brain-resident macrophages, microglia. These cells are considered to be the body's first line of defense, and are main propagators of inflammation, which has been shown to exacerbate neurodegeneration (2-4). Furthermore, these cells, having the ability to phagocytose, can take up these proteinaceous Aβ plaques as well as dying neurons. It has been noted for some time that although they possess the ability to do so, microglia and peripheral monocytes and macrophages fail to effectively clear these plaques. This has been hypothesized to occur with declining immune function with age, because as the disease progresses, the cells of the immune system become dysregulated, senescent, and associated with low-grade chronic inflammation (5). The lack of disease-modifying interventions for AD until recently has fueled the search for identification of dysregulated immune pathways as potential new targets to delay, arrest, and perhaps prevent disease development (6).

Genome-wide association studies identified a mutation within the phospholipase C gamma 2 (*PLCG2*) gene, called *P522R*, that protects against AD-related cognitive decline (7, 8). PLCG2 is highly expressed in immune cells including brain-resident microglia, peripheral monocytes, and tissue-resident macrophages (9). Determining how this mutation confers its protection in these cell populations will allow us to better understand how these cells play a role

in the progression of AD, and may open up new avenues for therapeutic development. PLCG2

has been shown to support numerous myeloid cell functions including phagocytosis, cytokine

secretion, survival, and lipid metabolism (10). To do so, it acts downstream of receptors such as

triggering receptor expressed on myeloid cells 2 (TREM2) and toll-like receptors (TLRs) to

cleave phosphatidylinositol (4,5) bisphosphate (PIP2) into diacylglycerol (DAG) and inositol

1,4,5 triphosphate (IP3). DAG and IP3 go on to both release intracellular calcium and activate

downstream pathways including nuclear factor light chain enhancer of activated B cells (NF-κB)

(10, 11). The *P522R* mutation has been shown to be hypermorphic, indicating an enhancement of

immune cell functions (12). As the primary brain cells that express PLCG2 are microglia, much

of the work aimed at investigating how its protective effect is conferred has focused on how it

alters microglial function. Studies have shown that the *P522R* mutation increases the release of

inflammatory cytokines by microglia, and has variable effects on phagocytosis depending on

cargo size (13). For example, studies showed that the *P522R* mutation reduced phagocytosis of

Escherichia coli but increased phagocytosis of Aβ (14, 15). Further *in vivo* work confirmed this

increased uptake of Aβ in a 5XFAD mouse model of amyloid-pathology (16).

However, the effect of this mutation in peripheral tissue-resident macrophages has yet to

be studied. As this protective pathway is of interest in the development of potential therapeutics,

it is important then to understand how those therapeutics may affect the immune system globally,

and not just within the brain. Furthermore, peripheral macrophages may not just be bystanders in

the progression of AD, as it has been shown that they are not only more efficient phagocytes of

Aβ than microglia, but they are also capable of traversing the blood-brain barrier during times of

injury and disease (17-21). There is further evidence to suggest this traversal of peripheral

myeloid cells to the brain may not even be necessary to impact AD progression. Indeed, studies

have shown that AD progression and pathology can be directly impacted by peripheral immune

cell functions. For example, Aβ efflux out of the brain and into peripheral reservoirs such as the

blood and tissues has been shown to occur (22, 23). The clearance of Aβ from these reservoirs

was associated with lower cerebral Aβ pathology (24). Furthermore, peripherally derived Aβ has

been shown to enter the brain and exacerbate neuropathology, further supporting the importance

of peripheral Aβ stores and clearance (25). Therefore, determining the effect of this protective

mutation on peripheral macrophage function is of vital importance.

Additionally, given that women are at higher risk for the development of AD, we posit

that sex-dependent differences in the immune system may be partially responsible for this

increased risk. For instance, it has been well established that females are at greater risk for

autoimmune conditions due to an overactive and inflammatory immune system (26). Given that

activated microglia and infiltrating monocytes are detectable in the early stages of AD (27), basal

over-activation of innate immune cells in females could be a potential link to the increased

prevalence of AD in women. Further studies have also shown that relative to healthy control

females, females with AD possess a higher prevalence of less phagocytic pro-inflammatory

macrophages (28). This could lead to lower Aβ uptake, and over-activation of surrounding

immune cells. With this in mind, we determined that sex as a biological variable should be

investigated in the present study, as sexually-dimorphic differences have been described for both

immune function and risk for AD.

The aims of our study were two-fold. First, we aimed to study the effect of the *P522R*

mutation on a peripheral macrophage population to determine how these cells may be

differentially affected by the mutation. To do so, we assessed changes in major macrophage

functions including phagocytosis and inflammatory cytokine secretion, as well as the

transcriptional changes that underlie these functions. We utilized peritoneal macrophages

(pMacs) which can be separated into two main populations, large peritoneal macrophages

(LPMs) and small peritoneal macrophages (SPMs). In broad terms, LPMs are known to be

resident to the peritoneal cavity, and SPMs are derived from blood-monocytes, which travel to

the peritoneal cavity upon infection or other inflammatory events (29). Recent studies have

shown that these recruited monocytes can, over time, differentiate into macrophages that closely

mimic the resident LPMs (30). This population was selected over bone-marrow derived

macrophages as peritoneal macrophages differentiate within the tissue, which allows them to

mold to their environment and experience hormonal changes throughout the animal's lifespan

(29, 31, 32). Additionally, this population of peripheral macrophages is well characterized,

lending to a more informed study that aims to assess tissue-resident macrophages specifically

(29, 31, 32). Furthermore, the immune cells of the peritoneum are of great interest in relation to

the gut, which has been identified as a peripheral reservoir for Aβ, and as a conduit of peripheral

messengers to the brain via the gut-brain-axis (22, 33). Not only that, but immune responses

occurring in this region have been shown to directly activate the brain's insular cortex, a region

known to be affected during the preclinical stages of AD (34), which can in turn store this

immune-related information and alter the peripheral immune response if reactivated, further

supporting a direct connection between the brain and peritoneum (35). Second, we aimed to

address how sex may influence the effector functions of this population of macrophages, both

with and without the *P522R* mutation, as knowing the baseline sex differences in wild-type (WT)

mice would more easily allow us to assess additional effects of the *P522R* mutation. We found

that the *P522R* mutation does indeed alter pMac function, supporting the idea that this mutation

may potentially have protective effects on the organism due to its expression in the periphery and

not just in brain-resident immune cells. However, we also found that the effects of this mutation

are largely sexually dimorphic, indicating that future studies should utilize both male and female

sexes to better inform potential drug development strategies for the clinic.

METHODS

Animals

Congenic mice expressing the *P522R* mutation in the *Plcg2* gene were purchased from the

Jackson Laboratory (strain #029598) and housed at the University of Florida McKnight Brain

Institute vivarium. Mice heterozygous for the *P522R* mutation were produced by breeding

heterozygous mice and C57BL/6 control littermates. They were kept at 22°C in 60-70%

humidity and on a 12-hour light-dark cycle and fed *ad libitum*. All animal procedures were

approved by the University of Florida Institutional Animal Care and Use Committee and were in

accordance with the National Institute of Health Guide for the Care and Use of Laboratory

Animals (NIH Publications No. 80-23) revised 1996. Both male and female mice were bred and

used in these experiments and were 3-5 months old at the time of sacrifice. Mice were sacrificed

via cervical dislocation, and all experiments included 5-6 biological replicates.

pMac isolation and culture

Three days prior to pMac extraction, mice were injected with 1mL of a 3% thioglycolate

(Brewer) solution. Thirty minutes prior to thioglycolate injection mice were injected with 50uL

of sustained-release Buprenorphine (ZooPharm) to alleviate pain. After three days, the mice were

sacrificed via cervical dislocation. The abdomen was sprayed with 70% ethanol and the skin was

split along the midline. Once the peritoneal cavity was exposed, a 10mL syringe filled with

10mL of cold RPMI (Gibco, 11875119) was injected into the peritoneal cavity using a 27G

needle, avoiding blood vessels. Once the 10mL of cold RPMI was injected, another empty 10mL

syringe with a 25-gauge needle was used to withdraw and place the peritoneal fluid into a 15mL

Falcon collection tube. Any remaining fluid was aspirated via transfer pipette and placed into the

collection tube. Once all mice were sacrificed and peritoneal fluid was collected into separate

collection tubes, the fluid was passed through a 70uM nylon filter pre-wet with 5mL of cold

HBSS-/- (Gibco, 14175103) into a 50mL conical tube. 5mL of cold HBSS-/- was then placed

into the collection tube to rinse any remaining cells and passed through the filter as well. The

filter was then washed with 5mL of HBSS-/- and the 50mL tube containing the fluid and HBSS-

/- were centrifuged at 400g for 5 minutes at 4°C to pellet the cells. If any blood cells were

present the pellet was resuspended in 1mL of ammonium-chloride-potassium lysis buffer for 1

minute and then centrifuged again at 400g for 5 minutes at 4°C. Once all cells were pelleted, the

supernatant was aspirated and each pellet was resuspended in 3mL of plating media (RPMI, 10%

fetal bovine serum, and 1% penicillin/streptomycin). Viable cells were identified and counted

using trypan-blue exclusion on an automated cell counter (Countess™). The volume of growth

media was adjusted so that cells were plated at a density appropriate for the intended assay

(detailed below). Cells were incubated at 37°C, 5% CO_2 for a minimum of 2-hours to allow

macrophages to adhere. Wells were washed with sterile DPBS (Gibco, 14190235) to remove any

non-adherent cells, and new, pre-warmed growth media was added.

Flow Cytometry

For flow cytometry analysis, cells were plated at a density of 3×10^5 cells/well in a 24-well plate

and left to incubate overnight. 1 hour prior to collection, DQ-BSA green (Invitrogen, D12050) was

added to each well at a concentration of 10ug/mL to measure protein degradation. After 1 hour,

the cells were washed 3 times with sterile phosphate buffered saline (PBS) (Gibco, 10010023),

detached from the plate via cell scraper, and then transferred to a 96-well v-bottom plate and

centrifuged at 300*g* for 5 minutes at 4°C. The cells were then resuspended in 50uL of PBS

containing fluorophore-conjugated antibodies for CD11b (1:100, Biolegend, 101216), MHCII

(1:100, Biolegend, 107628), and Live/Dead stain (1:2000, Invitrogen, L34965). The cells were

then incubated in the dark at 4°C for 20 minutes at which point they were centrifuged again at

300*g* for 5 minutes at 4°C. The cells were then washed with sterile PBS, and this process was

repeated twice. The cells were then fixed in 50uL per well of 1% paraformaldehyde in PBS for 30

minutes in the dark at 4°C. After 30 minutes the cells were centrifuged at 300*g* for 5 minutes at

4°C, resuspended in 200uL of FACs buffer (PBS, 0.5 mM EDTA, 0.1% sodium azide), and taken

to the Macs quant analyzer (Miltenyi) for flow cytometric analysis. A minimum of 100,000 events

were captured per sample and data were analyzed using FlowJo version 10.6.2 software (BD

Biosciences). When validating flow cytometry panels and antibodies, fluorescence minus one

controls (FMOCs) were used to set gates and isotype controls were used to ensure antibody-

specific binding.

Phagocytosis Assay

For phagocytosis studies, cells were plated at a density of $5x10^4$ cells/well in a 96-well black

wall plate (Cellvis, NC1648269) in triplicate. Cells were left to incubate for 6 hours at 37°C in a

5% CO_2 humidified cell culture incubator. 2mL of Live Cell imaging solution (Invitrogen,

A59688DJ) was added to each vial of pHrodo green *E. coli* bioparticles (Invitrogen, P35366),

equating to a final concentration of 1mg/mL, and these solutions were sonicated for 10 minutes.

After sonication, the media was removed from the cells in the plate and replaced with 100uL of

E. coli bioparticle solution, at which point the plate was placed into an onstage incubator

connected to an EVOS M7000 microscope to maintain the cells at 37°C and 5% CO_2 humidified

atmosphere throughout the duration of the live-cell imaging. GFP images were captured at 20X

magnification with 5 images per well every 30 minutes for 15 hours within each well, and the

EVOS Celleste (Version 5) software was used to count cells and determine the integrated optical

density of GFP fluorescence within each well over time. Brightfield images were used for

automated-focus throughout the time course. Treatment conditions were performed in triplicate

wells and averaged for analysis.

Multiplexed Immunoassays

For inflammatory cytokine secretion, isolated pMacs were plated at a density of $1x10^6$ cells/well

in a 6-well plate and treated with either vehicle (PBS) or LPS (O111:B4 Sigma, L4391,

Lot#0000185317) at a concentration of 10ng/mL. The cells were then placed in a humidified cell

culture incubator at 37°C and 5% CO_2. Conditioned media was collected at 2, 6, and 24 hours

post stimulation for assessment of specific protein concentrations of TNF, IL-6, IL-1β, KC/GRO,

IL-4, IL-10, IFN-γ, IL-5, IL-2, and IL-12p70 (MSD, K15048D-2) using multiplexed

immunoassays on the MesoScale Discovery (MSD) platform. Samples were plated in duplicate

at a 1:1 dilution with MSD diluent 41 and the rest of the experimental procedure was performed

according to the manufacturer's instructions. The plate was then analyzed on the MSD

Quickplex machine and MSD software (Discovery Workbench Version 4.0). Cytokines with

values below the lower limit of detection (LLOD listed below) were not quantified, these being

0.10pg/mL for TNF, 0.04pg/mL for IFN-γ, 0.11pg/mL for IL-1β, 0.22pg/mL for IL-2,

0.11pg/mL for IL-4, 0.06pg/mL for IL-5, 0.61pg/mL for IL-6, 0.94 pg/mL for IL-10, 9.95pg/mL

for IL-12p70, and 0.24pg/mL for KC/GRO.

RNA Isolation

For RNA extraction, cells were plated at $1x10^6$ cells/well in a 6-well plate and treated with either

vehicle or LPS at 10ng/mL. The cells were placed back in a humidified cell culture incubator at

37°C and 5% CO_2 for 6 hours. Buffer RLT from the RNeasy mini kit (Qiagen, 74104) was

combined with β-mercaptoethanol (Sigma, 63689) at a concentration of 20uL of BME per 1mL

of buffer RLT. Media was aspirated from the plate and then 350uL of RLT+BME solution was

then added to each well to lyse the cells. Each well was scraped with a mini cell scraper (Fisher,

NC0325221) and collected into a Qiashredder (Qiagen) column and centrifuged at 10,000g for 2

minutes. 350uL of 70% ethanol was added to the homogenized lysate and then transferred to a

RNeasy spin column. The columns were then centrifuged for 30 seconds at 10,000g and

afterwards the flow-through was discarded. 700uL of buffer RW1 was then added to each

column and the columns were centrifuged at 10,000g for 30 seconds and the flow-through was

discarded. 500ul of buffer RPE was added to each column and they were spun at 10,000g for 2

minutes and the flow-through was again discarded. This step was repeated twice, and the column

was then placed into a 1.5mL collection tube, 30uL of elution buffer was added to the column,

and then centrifuged for 1 minute at 10,000g to elute the RNA. Concentration of RNA was

analyzed on a spectrophotometer, and RNA quality was assessed using Bioanalyzer 2100

(Agilent, Denmark).

Gene expression analysis

For quantification of mRNA transcripts in pMacs, the NanoString nCounter platform and Mouse

Immune Exhaustion Panel RNA (CAT#PSTD-M-EXHAUST-12) containing 785 genes was

used. 25ng/ul of RNA was loaded and hybridized to the provided capture and reporter probes

overnight at 65°C according to the manufacturer's instructions. The samples were then added

into the provided nCounter chip via the use of the NanoString MAX/FLEX

robot. Unhybridized capture and reporter probes were removed, and the hybridized mRNAs were

immobilized for transcript count quantification on the nCounter digital analyzer. The data was

then imported into the nSolver analysis software v4.0 for quality check and analysis according to

the manufacturer's instructions. Raw counts were normalized to the geometric mean of the

positive and negative controls and count expression was calculated via nSolver Advanced

Analysis v2.0.134. Gene expression was considered significant if $p < 0.05$ after adjustment using

the Benjamini-Hochberg method for false discovery rate, and if the fold change was greater than

1.5. Pathway analysis was performed using Enrichr for GO biological process analysis (36-38).

Statistical Analysis

Statistical analysis was performed using Prism Graphpad version 10.2.2. Differences between

groups was determined using either T-test with Welch's correction or 2-way ANOVA with

multiple comparisons using Tukey's post-hoc correction. Results were considered statistically

significant if $p < 0.05$. Each graph depicts individual data points with error bars representing the

standard error of the mean (SEM).

RESULTS

The P522R mutation does not alter pMac sub-populations in male mice but increases antigen

presentation

Before addressing cellular functionality, it is beneficial to first understand and characterize the

macrophage populations present within our samples. As mentioned, pMacs are not a

homogenous population, with SPMs arising from blood monocytes and LPMs being resident to

the peritoneal cavity. To determine whether the *P522R* mutation has any effect on the

frequencies of these cell populations we isolated pMacs from the peritoneal cavity of male mice,

and then used flow cytometry to distinguish SPMs from LPMs on the basis of CD11b

expression, as has been previously described (Supplemental Figure 1A) (39). Prior to isolation,

we performed an intraperitoneal injection of a 3% thioglycolate solution. This step is crucial

prior to isolation, as it allows us to harvest both LPMs and SPMs, not just the resident LPMs.

However, we do allow a three-day rest period post injection and prior to cell isolation so that

acute inflammation and cell activation induced by the injection subside (Figure 1).

Figure 1: Schematic of pMac collection and the purpose of thioglycolate injections prior to cell isolation.
Diagram describing the process of pMac isolation (A) and visual representation of the purpose of starting the
process with a thioglycolate injection (B). Created with Biorender.com.

Additionally, after plating the cells collected from the peritoneum, we leave the cells to rest for

two hours and then rinse with sterile DPBS and replace the media, as this allows only the

adherent cells (pMacs) to remain and all other immune cell types to be washed away

(Supplemental Figure 1B). We further confirmed LPM vs SPM populations by assessing

canonical peritoneal macrophage markers of LPMs (CD80 and F4/80) and SPMs (Dectin1)

(Supplemental Figure 1C-E). We found that male mice carrying the *P522R* mutation showed no

difference in the frequency of either LPM or SPM sub-populations compared to WT male mice

(Figure 2A, B). This was confirmed via raw counts (Supplemental Figure 2A-D). Next, we used

major histocompatibility complex class II (MHCII) as a marker of antigen presentation. We

found that although LPM and SPM populations were not affected by the *P522R* mutation in male

mice, antigen presentation on LPMs was. A higher frequency of LPMs were MHCII+ in *P522R*

males compared to LPMs from WT males (Figure 2C).

Figure 2: The *P522R* mutation does not alter pMac sub-populations in male mice but increases antigen
presentation. To determine the prevalence of different sub-populations of macrophages in our *ex vivo* cultures
pMacs from male WT and *P522R* animals were isolated, plated, and non-adherent cells were washed away after a 2-
hour rest period and populations of (A) LPM (B) SPM (C) MHCII+ LPM and (D) MHCII+ SPM were quantified via
flow cytometry. Data was quantified via T-test with Welch's correction. Groups are considered non-significant if
p>0.05 and significantly different if *p*<0.05.

We did not find any significant differences between *P522R* and WT males in MHCII expression

on SPMs (Figure 2D). Representative plots for this analysis can be found in Supplemental Figure

2E and 2F. These data indicate that in males the *P522R* mutation does not affect the makeup of

peritoneal macrophage sub-populations, but does increase the amount of resident LPMs

displaying antigens.

The P522R mutation alters both sub-populations and antigen presentation in pMacs from

female mice

Based on the literature indicating distinct sex-differences in pMac populations, we aimed to

determine whether the *P522R* mutation would have similar effects on population frequencies and

antigen presentation in female mice as was seen in male mice (40). However, we found that the

P522R mutation had quite different effects in female mice compared to those seen in male mice.

Female mice carrying the *P522R* mutation exhibited significant increases in the frequency of

LPMs, and significantly lower frequencies of SPMs compared to WT females (Figure 3A, B).

After reviewing the counts, we determined that these effects were primarily driven by an

increase in the raw count of LPMs in females that carry the *P522R* mutation (Supplemental

Figure 3A-D). Furthermore, contrary to the effect seen in pMacs from males, the *P522R*

mutation's presence in females actually decreased the frequency of MHCII+ LPMs compared to

pMacs from WT females (Figure 3C). But similar to males, there were no observable differences

between the frequencies of MHCII+ SPMs between WT or *P522R* females (Figure 3D),

representative plots for which can be found in Supplemental Figure 3E and 3F.

Figure 3: The *P522R* mutation alters both sub-populations and antigen presentation in pMacs from female
mice. To determine the prevalence of different sub-populations of macrophages in our *ex vivo* cultures pMacs from
female WT and *P522R* animals were isolated, plated, and non-adherent cells were washed away after a 2-hour rest
period and populations of (A) LPM (B) SPM (C) MHCII+ LPM and (D) MHCII+ SPM were quantified via flow
cytometry. Data was quantified via T-test with Welch's correction. Groups are considered non-significant if p>0.05
and significantly different if *p*<0.05.
327
328   These data support a role for the *P522R* mutation in altering macrophage sub-populations within

329   the peritoneum of female mice. However, these effects are largely sex-specific, and even bi-

330   directional in the case of antigen presentation. These data begin to confirm a role for the *P522R*

331   mutation in peripheral tissue resident macrophages, and further support previously published

332   data indicating significant sex-differences in immune cell populations and function.

The P522R mutation reduces phagocytic capacity and lysosomal protease activity in pMacs

from males

After determining that the *P522R* mutation can alter pMac populations in female mice, and

antigen presentation on LPMs in a bidirectional manner in females and males, we aimed to

determine whether this mutation also has functional consequences in pMacs from each sex.

Macrophages are known to have many dynamic functions, yet phagocytosis and cytokine

secretion are among the most recognized, both of which PLCG2 is actively involved in. We

began with assessing phagocytosis capabilities in pMacs from male WT and *P522R* mice, as the

ability to uptake and degrade bacterial/viral particles is among the most important functions of

macrophages in the fight against infection. Furthermore, this function has been directly

associated with AD, as *E. coli* particles and LPS particles have been shown to associate directly

with pathology in the brains of AD patients (41). Additionally, AD progression has been

correlated with infection, emphasizing the importance of the pMacs ability to uptake and degrade

bacterial particles (42). To assess the phagocytic capabilities of these macrophages, we utilized

fluorescently labeled *E. coli* bioparticles which only fluoresce after being internalized and

exposed to the low pH associated with the lysosome, thereby reporting on the extent of bacterial

uptake by the cells. Interestingly, we found that pMacs from *P522R* male mice performed less

uptake of the fluorescent *E. coli* over the course of 15 hours than did pMacs from WT male mice

(Figure 4A-C). Up until approximately the 6 hour mark, the *P522R* and WT pMacs performed

very similar amounts of *E. coli* uptake, but the *P522R* pMacs were not able to keep up with the

WT pMacs, which seemingly performed better (Figure 4B). Although the macrophages were

able to take up the *E. coli* particles, we further assessed whether pMacs from males displayed

any differences in lysosomal function, or the ability to successfully break down that which has

been phagocytosed. To do so, we treated the cells with DQ-BSA, a substrate for proteases that

only fluoresces after it has been cleaved within the lysosome, and quantified via flow cytometry.

Figure 4: The *P522R* mutation reduces phagocytic capacity and lysosomal protease activity in pMacs from
males. pMacs from WT and *P522R* males were plated and incubated for 6 hours, at which point 200ug of *E. coli*
was added to each experimental well to assess *E. coli* uptake. (A) Representative brightfield images and GFP
fluorescence for each group are presented at 20X magnification. (B) Integrated optical density of GFP fluorescence
was plotted over 15 hours. (C) Groups were compared statistically via area under the curve. 1 hour prior to cell
collection DQ-BSA was added to the plate and then cells were collected for flow cytometry and (D) DQ-BSA MFI
is represented. Groups are considered non-significant if $p>0.05$ and significantly different if $p<0.05$.
366
Again, we found that pMacs from WT males performed better than pMacs from *P522R* males,

which exhibited lower lysosomal protease activity compared to pMacs from WT males (Figure

4D). These data indicate that the *P522R* mutation does indeed alter phagocytosis and lysosomal

activity in males, but these alterations may not be beneficial to pMac function.

The P522R mutation does not alter phagocytic capacity in pMacs from females, but increases

lysosomal protease activity

As the *P522R* mutation has been associated with protection, the decrease in phagocytic capacity

and lysosomal protease activity seen in pMacs from *P522R* males was quite surprising. We

repeated both the *E. coli* uptake assay, as well as the DQ-BSA lysosomal protease activity assay

in pMacs from WT and *P522R* females to determine whether similar patterns were seen in

females. Similar to what we found in our assessment of pMac populations, the *P522R* mutation

had different effects in pMacs from females compared to pMacs from males. Although there

appeared to be a visual difference in GFP fluorescence between pMacs from *P522R* and WT

females, quantification showed that uptake of *E. coli* in pMacs from females was largely

unaffected by the *P522R* mutation (Figure 5A-C). However, lysosomal protease activity

measured via fluorescence of the DQ-BSA probe was shown to be increased in pMacs from

P522R females (Figure 5D).

Figure 5: The *P522R* mutation does not alter phagocytic capacity in pMacs from females, but increases
lysosomal protease activity. pMacs from WT and *P522R* females were plated and incubated for 6 hours, at which
point 200ug of *E. coli* was added to each experimental well to assess *E. coli* uptake. (A) Representative brightfield
images and GFP fluorescence for each group are presented at 20X magnification. (B) Integrated optical density of
GFP fluorescence was plotted over 15 hours. (C) Groups were compared statistically via area under the curve. 1
hour prior to cell collection DQ-BSA was added to the plate and then cells were collected for flow cytometry and
(D) DQ-BSA MFI is represented. Groups are considered non-significant if *p*>0.05 and significantly different if
p<0.05.
393

These data indicate that the *P522R* mutation may confer functional advantages in lysosomal

activity in pMacs from females, but not in males.

The P522R mutation modestly increases inflammatory cytokine secretion by pMacs from

males

As mentioned, the other function of macrophages that has been linked to PLCG2 function is

cytokine secretion. We have already demonstrated that the *P522R* mutation has sex-dependent

effects on phagocytosis and lysosomal function, and we hypothesized that it would also alter

cytokine secretion by pMacs, likely also in a sex-dependent manner. To test this, we treated

pMacs from WT and *P522R* males with LPS and measured the protein concentration of

cytokines within the media at 2, 6, and 24 hours post-LPS treatment. We found that vehicle

treated cells produced very low concentrations of inflammatory cytokines, with no difference

between WT and *P522R* cells, further supporting that our *ex vivo* cultures are not activated or

inflammatory at baseline conditions (Figure 6A-H). For this reason, we treated with the immune

cell stimulant LPS to elicit an inflammatory response, as although there were no genotype

differences at baseline, the different genotypes may respond differently to inflammatory stimuli.

Figure 6: The *P522R* mutation modestly increases inflammatory cytokine secretion by pMacs from males.
pMacs from male WT and *P522R* mice were plated and treated with LPS, and their conditioned media was collected
at 2, 6, and 24 hours post LPS stimulation. Concentrations of (A) TNF (B) IL-6 (C) KC/GRO (D) IL-1β (E) IL-

413 12p70 (F) IL-2 (G) IL-10 and (H) IFN-γ were then assessed via MesoScale Discovery immunoassays. Data were
414 quantified via two-way ANOVA using Tukey's post-hoc correction for multiple comparisons. Groups are
415 considered non-significant if *p*>0.05 and significantly different if *p*<0.05.
416
Indeed, we found that upon LPS stimulation, pMacs from *P522R* males showed modest increases

in inflammatory cytokine secretion of TNF, IL-6, and IL-12p70 at 6 hours post-LPS stimulation

(Figure 6A, B and E), and IFN-γ at 24 hours post-LPS stimulation compared to pMacs from WT

males (Figure 6H). We found no differences between genotypes in secretion of KC/GRO, IL-1β,

IL-2, or IL-10 even after LPS stimulation (Figure 6C, D, F and G). These data support that the

P522R mutation has a modest effect on pMac cytokine secretion in males.

The P522R mutation increases inflammatory cytokine secretion by pMacs from females

To determine whether these same stimulation dependent effects were seen in females, we

repeated this same experiment but with pMacs from WT and *P522R* females. Similar to pMacs

from males, we found the pMacs from both WT and *P522R* females produce very low levels of

inflammatory cytokines when not stimulated with LPS (Figure 7A-H), supporting that, like our

male cultures, the female *ex vivo* cultures are not inflammatory or activated prior to stimulation.

Figure 7: The *P522R* mutation increases inflammatory cytokine secretion by pMacs from females. pMacs from
female WT and *P522R* mice were plated and treated with LPS, and their conditioned media was collected at 2, 6,
and 24 hours post-LPS stimulation. Concentrations of (A) TNF (B) IL-6 (C) KC/GRO (D) IL-1β (E) IL-12p70 (F)
IL-2 (G) IL-10 and (H) IFN-γ were then assessed via MesoScale Discovery immunoassays. Data were quantified via
two-way ANOVA using Tukey's post-hoc correction for multiple comparisons. Groups are considered non-
significant if *p*>0.05 and significantly different if *p*<0.05.

Similar to what we saw in our male data, LPS stimulation did reveal genotype-dependent

differences in cytokine secretion, however these differences were different and more prevalent

than in our male cultures. For example, at 6 hours post-LPS stimulation, pMacs from *P522R*

females were secreting more TNF, KC/GRO, IL-1β, and IL-12p70 than pMacs from WT females

(Figure 7A, C, D, and E). At 24 hours post-LPS stimulation, pMacs from *P522R* females were

secreting more TNF, IL-6, KC/GRO, IL-1β, and IL-2 compared to pMacs from WT females

(Figure 7A-D, and F). We found no genotype dependent differences in IL-10 or IFN-γ at any

time point (Figure 7G, H). So, although the *P522R* mutation caused increases in cytokine

secretion by pMacs from both males and females, the pMacs from females seemed to be more

affected by the presence of the mutation and produced significantly more inflammatory

cytokines over time.

Genes selectively upregulated in P522R males are associated with negative regulation of the

immune response

In our selected assays, the *P522R* mutation seemed to have a dampening effect on pMacs from

males, indicating a potential sex-specific role for the mutation. Furthermore, many of these

differences were more noticeable after treatment with some sort of immune cell stimulant, such

as *E. coli* or LPS, supporting the idea that the *P522R* mutation's effects may be activation-

dependent, and not as prevalent under baseline conditions. To attempt to identify molecular

pathways associated with the effects of stimulation, we performed a targeted transcriptomic

analysis using Nanostring nCounter to assess 785 genes associated with immune cell function in

pMacs from both WT and *P522R* males. We did so both at baseline and after treatment with LPS

to determine whether these effects were truly activation dependent. As expected, we did not find

any significant differences in gene expression between pMacs from WT and *P522R* males under

baseline conditions (Supplemental Figure 5A). However, after treatment with LPS, differences

between the genotypes did begin to arise.

Figure 8: Genes selectively upregulated in *P522R* males are associated with negative regulation of the immune
response. pMacs from male WT and *P522R* mice were plated and treated with either vehicle or LPS (10ng/mL) and
left to incubate for 6 hours. Cells were then collected and RNA was isolated for gene expression analysis. (A)
Volcano plot comparing gene expression of LPS treated pMacs from male WT mice to gene expression of vehicle
treated pMacs from male WT mice, horizontal grey dashed line indicating adjusted *p*<0.05, and vertical dashed lines
indicating fold change of greater than 1.5. (B) Volcano plot comparing gene expression of LPS treated pMacs from
male *P522R* mice to gene expression of vehicle treated pMacs from male *P522R* mice, horizontal grey dashed line
indicating adjusted *p*<0.05, and vertical dashed lines indicating fold change of greater than 1.5. (C) Venn diagram
comparing number of DEGs between pMacs from WT males and pMacs from *P522R* males after LPS stimulation.

(D) GO biological process pathways associated with DEGs specific to pMacs from WT males in response to LPS
stimulation. (D) GO biological process pathways associated with DEGs specific to pMacs from *P522R* males in
response to LPS stimulation.

Volcano plots representing each genotype's response to LPS stimulation are represented in

Figure 8A and 8B, with 8A showing pMacs from WT male's response to LPS, and 8B showing

pMacs from *P522R* male's response to LPS. We found that many of the differentially expressed

genes (DEGs) induced by LPS stimulation were shared between pMacs from WT and *P522R*

males, encompassing 264 shared DEGs (Figure 8C). However, there were a number of genes that

were genotype specific, including 13 DEGs specific to *P522R* males, and 21 specific to WT

males (Figure 8C). We took each of these sets of genotype specific genes and assessed what

pathways were associated with them. We found that the DEGs specific to pMacs from WT males

were associated with pathways including calcineurin-NFAT signaling cascade, regulation of

calcium ion transport across the plasma membrane, calcineurin-mediated signaling, and protein

modification/phosphorylation (Figure 8D). Many of these pathways involve calcium signaling,

and the calcineurin cascade which in macrophages has been shown to down-regulate TLR

mediated activation pathways (43). Contrarily, the DEGs specific to pMacs from *P522R* males

were associated with pathways including negative regulation of immune responses, regulation of

type II interferon production, regulation of NK T cell activation, and response to UV (Figure

8E). These results suggest that the *P522R* mutation induces gene expression that has a

dampening effect on pMac immune responses in males.

Genes selectively upregulated in P522R females are associated with cytokine signaling and

apoptosis

As so far we have seen sex-dependent differences in pMac responses to the *P522R* mutation, we

did targeted transcriptomics on our female samples as well to determine which pathways might

be differentially implicated in the female's varied responses. Similar to our male data, when

pMacs from females are not treated with LPS, there are no significant gene expression

differences between pMacs from WT females and *P522R* females (Supplemental Figure 5B).

Figure 9: Genes selectively upregulated in *P522R* females are associated with cytokine signaling and apoptosis. pMacs from female WT and *P522R* mice were plated and treated with either vehicle or LPS (10ng/mL) and left to incubate for 6 hours. Cells were then collected and RNA was isolated for gene expression analysis. (A) Volcano plot comparing gene expression of LPS treated pMacs from female WT mice to gene expression of vehicle treated pMacs from female WT mice, horizontal grey dashed line indicating adjusted *p*<0.05, and vertical dashed lines indicating fold change of greater than 1.5. (B) Volcano plot comparing gene expression of LPS treated pMacs from female *P522R* mice to gene expression of vehicle treated pMacs from female *P522R* mice, horizontal grey dashed line indicating adjusted *p*<0.05, and vertical dashed lines indicating fold change of greater than 1.5. (C) Venn diagram comparing number of DEGs between pMacs from WT females and pMacs from *P522R* females after LPS

509 stimulation. (D) GO biological process pathways associated with DEGs specific to pMacs from WT females in
510 response to LPS stimulation. (D) GO biological process pathways associated with DEGs specific to pMacs from
511 *P522R* females in response to LPS stimulation.
512
However, upon LPS treatment, different responses to this activating stimulus occur. DEGs are

represented in volcano plots for pMacs from WT females after LPS treatment (Figure 9A), and

pMacs from *P522R* females after LPS treatment (Figure 9B). Many of the DEGs implicated in

the cellular response to LPS are the same between pMacs from WT and *P522R* females,

comprising 190 shared DEGs (Figure 9C), but each genotype did again have specific DEGs

related to them, with 50 DEGs specific to WT females, and 12 DEGs specific to *P522R* females

in response to LPS. These distinct gene groups were once again analyzed via GO biological

process pathway analysis, and the results were, unsurprisingly, quite distinct from the male

groups. We found that DEGs specific to pMacs from WT females were associated with pathways

including regulation of defense response, regulation of immune response, regulation of

autophagy, and negative regulation of the inflammatory response (Figure 9D). DEGs specific to

pMacs from *P522R* females on the other hand were associated with pathways including cytokine

mediated signaling, apoptotic processing, extrinsic apoptotic signaling, and positive regulation of

peptidyl-tyrosine phosphorylation (Figure 9E). This suggests that the *P522R* mutation induces

genes related to the upregulation of immune responses in pMacs from females.

DISCUSSION

Numerous studies have implicated the immune system as a player in the progression of

neurodegenerative disorders, both within the brain and in the periphery. This includes evidence

that peripheral monocytes and tissue-resident macrophages participate in Aβ plaque uptake, both

around the brain and in peripheral tissue reservoirs (22-24). Furthermore, macrophages are

known to be one of the main producers of inflammatory cytokines, which, when chronic, can

exacerbate neurodegeneration (2-4), but when acute, act as messengers to brain immune cells

and signals to other local immune cells (44-46). Therefore, it is of critical importance to

investigate the nature of peripheral immune mechanisms to identify potential new targets for

therapeutic intervention. With this objective in mind, many groups have studied the effects of the

Plcg2-P522R protective mutation in myeloid cells, primarily microglia, to try to determine how

this protection is being conferred since the effects manifest themselves in improved cognitive

function. However, the effect of the *P522R* mutation on peripheral tissue-resident macrophages,

which have become of increasing importance in the context of AD, had yet to be determined.

Additionally, we assessed how the mutation's effects may differ between sexes, as it is well

known that both AD risk and immune system function is highly affected by sex. Herein, we

show that the *P522R* mutation does indeed confer functional alterations in the periphery, but

these changes seem to vary by sex.

When assessing pMac sub-populations, we found that the *P522R* mutation had different

effects that were dependent on sex. In WT male mice, we found that our *ex vivo* cultures were

predominantly comprised of LPMs, making up approximately 80% of the population (the other

portion being made up of SPMs), and we found this to be the same in cultures obtained from

P522R males, indicating that the mutation does not have an effect on pMac sub-populations in

male mice. In cultures obtained from female WT mice however, we found that only around 60-

65% of the population on average was comprised of LPMs. The addition of the *P522R* mutation

altered this significantly, bringing this percentage up to closer to 80% LPMs. These data indicate

that the mutation alters pMac sub-populations in females but not males, and interestingly seemed

to bring female sub-population composition up to meet that which was found in males. Although

the canonical sub-populations known as LPMs and SPMs are recognized as separate groups (29),

recent studies have shown that not only can SPMs continue to alter their functional and

transcriptional profiles to look more like LPMs, and even occupy that same niche over time, but

they actually do so more readily in males (30, 40). It might be possible then that the *P522R*

mutation somehow ablates this male-bias, making it possible for the infiltrating cells to become

more LPM-like in females over time as is seen in males. However, it could also stand to reason

that this increase in LPMs in *P522R* females may happen prior to/separate from our

interventions, potentially during the time of peritoneum colonization by the early resident

macrophages. Our data is not enough from which to draw substantiated conclusions on this topic,

however it can be concluded that the *P522R* mutation does alter sub-populations (particularly

LPMs) in female mice via a currently unknown mechanism. Based on our data and the

aforementioned studies assessing the dynamic properties of these macrophages to shift from one

sub-group to another at any particular time, we decided to perform the rest of the experiments on

the entire population of pMacs available within a well, instead of trying to isolate LPMs and

SPMs specifically for functional assays. Although possible, the ability of these cells to take on

characteristics of the other group make this unnecessary, as these sub-populations are not static

regardless. We do however, use our flow data pertaining to the makeup of our populations in

each sex to influence our interpretation of functional results, but we do not rely solely on this for

the aforementioned reasons.

The first functions we assessed were phagocytosis and lysosomal activity, these being the

ability to take up particles, and the ability to break them down, respectively. The literature

addressing the effect of the *P522R* mutation on phagocytosis is inconsistent, showing both

increased and decreased phagocytosis (14-16). This discrepancy was hypothesized to be due to

the size of the cargo, with studies showing lower phagocytosis of larger particles such as

zymosan and *E. coli*, and more phagocytosis of smaller-sized dextrans (14, 15). We show here

that this observed difference in phagocytosis caused by the *P522R* mutation could also be due to

the sex in which the testing was performed. In our cultures, we found that the *P522R* mutation

did alter phagocytosis and lysosomal activity in pMacs, but in opposing directions in males and

females. Our data showed that pMacs from *P522R* males exhibited less uptake of *E. coli*

particles over time compared to pMacs from WT males. In females, these same differences were

not found, and the *P522R* mutation seemingly had little effect on *E. coli* uptake over time. We

further observed changes in lysosomal activity, with the *P522R* mutation significantly reducing

lysosomal protease activity in males, and significantly increasing it in females, further supporting

the sex-dependent differences this mutation confers. It should be noted however, that differences

in phagocytosis in males were not observed until around the 6-7 hour mark, meaning that initial

uptake of *E. coli* was similar between groups, but the pMacs from *P522R* males may become

exhausted earlier or more rapidly. Taken together, these results suggested the potential of the

P522R mutation to be positive in females, but potentially negative in males via the reduction in

bacterial uptake and degradation.

Further functional assays surveyed inflammatory cytokine secretion by pMacs from

females and males, as previous studies showed that the *P522R* mutation was capable of

increasing cytokine secretion in microglia (13). Our data indicated that this effect is also present

in pMacs, but to different extents in males and females. We measured cytokine release both at

baseline and post-LPS to determine whether there may be differences prior to any sort of

external stimulation. In both male and female cultures, we saw that at baseline, regardless of

genotype, the pMacs were not producing inflammatory cytokines, supporting that our *ex vivo*

culture model allows for the cells to return to baseline non-inflammatory conditions. For this

reason, we chose to treat the cells with the immune cell stimulant LPS to elicit an inflammatory

response that could be compared between the groups. After LPS stimulation, pMacs from *P522R*

males showed increased release of TNF, IL-6, IL-12p70, and IFN-γ over 24 hours compared to

pMacs from WT males, while pMacs from *P522R* females showed increased release of TNF, IL-

6, KC/GRO, IL-1β, IL-12p70, and IL-2 over 24 hours compared to pMacs from WT females.

Interestingly, the *P522R* mutation had no effect on the release of IL-10, a canonical cytokine

involved in the resolution of inflammatory responses, in either females or males. Although it

could be assumed that these increases in inflammatory cytokines could be negative, our study

only looked at acute cytokine secretion, which is likely to have been an adaptive and beneficial

response.

In our assays assessing the effects of the *P522R* mutation on pMac function and

population dynamics we found different functional outcomes depending on sex. Furthermore, we

found that many of these differences were more/only prominent after some form of cell

activation either via *E. coli* or LPS treatment. This led us to believe that the *P522R* mutation may

exert stronger effects upon cell activation in females and males, and we wondered whether we

might be able to potentially determine some cellular pathways implicated in these differential

responses, and therefore we performed targeted transcriptomics to assess immune cell specific

gene expression changes. First, we compared gene expression in pMacs from *P522R* males to

pMacs from WT males and found no differences at baseline. From that point, we compared gene

expression after LPS stimulation between these two groups to determine what changes might be

occurring due to the *P522R* mutation in response to cellular activation. What we found was quite

fascinating. We saw that in response to LPS stimulation, the DEGs specific to *P522R* males were

related to pathways associated with negative regulation of the immune response. This lines up

with the functional data that revealed that in pMacs from males, the *P522R* mutation decreases

phagocytosis and lysosomal activity, and does not exert as strong of an effect on inflammatory

cytokine secretion as is seen in females. In females however, we found that the DEGs specific to

pMacs from *P522R* females were quite different, being related to cytokine mediated signaling

pathways, positive regulation of peptidyl-tyrosine phosphorylation, and apoptotic signaling.

Tyrosine phosphorylation has been characterized as a main event following LPS stimulation in

macrophages which aids in cellular activation (47), in line with the observed increase in

lysosomal activity and cytokine secretion in females. Furthermore, cytokine mediated pathways

and apoptotic signaling can go hand in hand, as cytokines are main promoters of apoptosis,

particularly TNF (48, 49). It appears based on our transcriptomic data that the *P522R* mutation

activates different intracellular signaling pathways and concurrent gene expression in response to

LPS leading to the different outcomes in response to activation seen between male and females.

At this time, we cannot say directly how these sex-dependent responses may contribute to

the neuroprotection associated with this mutation. However, we can say with certainty that

peripheral immune cells, particularly peritoneal macrophages, are affected by the presence of this

mutation, and show transcriptional, lysosomal, phagocytic, and inflammatory alterations due to

its presence, particularly after an inflammatory challenge. With recent studies indicating a role

for peripheral macrophages in the progression of AD, this study opens up new avenues of

interest for more in-depth studies of the contribution of peripheral macrophages to the *P522R*

mutation's protective effects. It should be noted however, that these responses and functional

differences may not solely be due to intrinsic changes in macrophages due to the *P522R*

mutation. As we isolated these cells from the peritoneum, a location filled with many immune

cell types (many of which would also carry the KI *P522R* mutation), some of these functional

changes could be due to the macrophages' direct interaction with/response to other mutated

immune cells which might leave lasting impacts on macrophage function. While we hypothesize

that the functional changes seen in this study are due to direct intrinsic changes in macrophages

brought about by the *P522R* mutation, which is supported by the studies showing intrinsic

mechanisms of the *P522R* mutation in similar cell type microglia (12, 13), we cannot completely

rule out the contribution of the peritoneal environment prior to our isolation protocol.

Additionally, we provide ample evidence to necessitate the use of both sexes in any future study

aiming to assess the protective effects of this mutation. Many drugs aiming to cure or mediate

disease have been pulled from the market due to unforeseen issues arising in one sex or another

(50, 51). Over recent years the issue of sex-differences in biomedical research has come to the

forefront, and we provide further evidence to support the advantages of utilizing both sexes in

pre-clinical studies whenever possible. Although this study does not provide an exact mechanism

by which these peripheral macrophages may contribute to the protective effects of this mutation,

it provides compelling reasons to support future inquiries into these questions, which we aim to

do moving forward using *in vivo* models of neurodegeneration to determine whether these same

functional alterations directly impact neuroinflammation and cognitive performance. It further

supports the importance of including sex as a biological variable in studies as well, showing that

not only can the magnitude of response to protective genetic mutations be different between

sexes, but also the directionality of response can vary by sex and will have implications for

future therapeutic interventions.

CONCLUSION

Inflammation and the cells of the peripheral immune system have been implicated in the

progression of AD. Therefore, identifying pathways that could be targeted in therapeutic

interventions will give the field the potential to modulate these dysregulated immune cell

functions. Others have demonstrated that the *P522R* mutation has the ability to enhance

important immune functions in microglia, such as phagocytosis and cytokine secretion. Herein

we provide the first evidence that the *P522R* mutation also modulates peripheral tissue-resident

macrophage functions, and that these effects are highly sex-dependent. Taken together, these

results reveal that peripheral immune cell changes should be taken into account when assessing

the mechanism(s) by which deleterious/beneficial genetic mutations confer their effects, as

microglia may not be the only cells contributing to the functional outcome of the genetic variant,

and that sex as a biological variable should be an integral part of any study looking to assess

peripheral inflammatory mechanisms that are likely to affect brain function.

LIST OF ABBREVIATIONS

PLCG2, phospholipase C gamma 2

AD, Alzheimer's disease

Aβ, amyloid-beta

TREM2, triggering receptor expressed on myeloid cells 2

TLR, toll-like receptor

PIP2, phosphatidylinositol (4,5) bisphosphate

DAG, diacylglycerol

IP3, inositol 1,4,5 triphosphate

LPM, large peritoneal macrophage

SPM, small peritoneal macrophage

WT, wild-type

PBS, phosphate buffered saline

FMOC, fluorescence minus one control

MSD, MesoScale Discovery

LLOD, lower limit of detection

SEM, standard error of the mean

MHCII, major histocompatibility complex II

LPS, lipopolysaccharide

DEG, differentially expressed gene

DECLARATIONS

Ethics approval and consent to participate

Mice were utilized in accordance with protocols approved by the Institutional Animal Care and

Use Committee of the University of Florida.

Consent for publication

Not applicable.

Availability of data and materials

All data are included in the present study or contained within the supplemental files. Raw data

can be provided upon request.

Competing Interests

The authors declare no conflicts of interest.

Funding

The funding for this project was provided by the McKnight Brain Institute and Norman Fixel

Institute for Neurological Diseases.

Authors Contributions

HAS and MGT conceived the study. HAS designed the experiments, performed them, and

analyzed and interpreted the data. MLB, RLW, JJ, KBM, CC, NN, and AMT assisted with

various experimental procedures. HAS designed all figures. HAS and MGT wrote and edited the

manuscript. All authors approved the final version of the manuscript and figures.

Acknowledgements

The authors thank the Tansey, Khoshbouei, and Chakrabarty labs for useful discussions.

REFERENCES
729
1. Scheyer O, Rahman A, Hristov H, et al. Female Sex and Alzheimer's Risk: The
Menopause Connection. J Prev Alzheimers Dis. 2018;5(4):225-230.
doi:10.14283/jpad.2018.34
2. Cunningham C, Campion S, Lunnon K, et al. Systemic inflammation induces acute
behavioral and cognitive changes and accelerates neurodegenerative disease. Biol
Psychiatry. 2009;65(4):304-312. doi:10.1016/j.biopsych.2008.07.024
3. Tejera D, Mercan D, Sanchez-Caro JM, et al. Systemic inflammation impairs microglial
Aβ clearance through NLRP3 inflammasome. EMBO J. 2019;38(17):e101064.
doi:10.15252/embj.2018101064
4. Marshall JS, Warrington R, Watson W, Kim HL. An introduction to immunology and
immunopathology. Allergy Asthma Clin Immunol. 2018;14(Suppl 2):49. Published 2018
Sep 12. doi:10.1186/s13223-018-0278-1
5. Cao W, Zheng H. Peripheral immune system in aging and Alzheimer's disease [published
correction appears in Mol Neurodegener. 2018 Oct 24;13(1):58]. Mol Neurodegener.
2018;13(1):51. Published 2018 Oct 3. doi:10.1186/s13024-018-0284-2
6. van Dyck CH, Swanson CJ, Aisen P, et al. Lecanemab in Early Alzheimer's Disease. *N
Engl J Med*. 2023;388(1):9-21. doi:10.1056/NEJMoa2212948
7. Kleineidam L, Chouraki V, Próchnicki T, et al. PLCG2 protective variant p.P522R
modulates tau pathology and disease progression in patients with mild cognitive
impairment. Acta Neuropathol. 2020;139(6):1025-1044. doi:10.1007/s00401-020-02138-
6
8. Sims R, van der Lee SJ, Naj AC, et al. Rare coding variants in PLCG2, ABI3, and
TREM2 implicate microglial-mediated innate immunity in Alzheimer's disease. Nat
Genet. 2017;49(9):1373-1384. doi:10.1038/ng.3916

9. Jackson JT, Mulazzani E, Nutt SL, Masters SL. The role of PLCγ2 in immunological disorders, cancer, and neurodegeneration. J Biol Chem. 2021;297(2):100905. doi:10.1016/j.jbc.2021.100905

10. Magno L, Bunney TD, Mead E, Svensson F, Bictash MN. TREM2/PLCγ2 signalling in immune cells: function, structural insight, and potential therapeutic modulation. Mol Neurodegener. 2021;16(1):22. Published 2021 Apr 6. doi:10.1186/s13024-021-00436-5

11. Andreone BJ, Przybyla L, Llapashtica C, et al. Alzheimer's-associated PLCγ2 is a signaling node required for both TREM2 function and the inflammatory response in human microglia. Nat Neurosci. 2020;23(8):927-938. doi:10.1038/s41593-020-0650-6

12. Magno L, Lessard CB, Martins M, et al. Alzheimer's disease phospholipase C-gamma-2 (PLCG2) protective variant is a functional hypermorph. Alzheimers Res Ther. 2019;11(1):16. Published 2019 Feb 2. doi:10.1186/s13195-019-0469-0

13. Takalo M, Wittrahm R, Wefers B, et al. The Alzheimer's disease-associated protective Plcγ2-P522R variant promotes immune functions. Mol Neurodegener. 2020;15(1):52. Published 2020 Sep 11. doi:10.1186/s13024-020-00402-7

14. Maguire E, Menzies GE, Phillips T, et al. PIP2 depletion and altered endocytosis caused by expression of Alzheimer's disease-protective variant PLCγ2 R522. EMBO J. 2021;40(17):e105603. doi:10.15252/embj.2020105603

15. Solomon S, Sampathkumar NK, Carre I, et al. Heterozygous expression of the Alzheimer's disease-protective PLCγ2 P522R variant enhances Aβ clearance while preserving synapses. Cell Mol Life Sci. 2022;79(8):453. Published 2022 Jul 27. doi:10.1007/s00018-022-04473-1

16. Tsai AP, Dong C, Lin PB, et al. Genetic variants of phospholipase C-γ2 alter the phenotype and function of microglia and confer differential risk for Alzheimer's disease. Immunity. 2023;56(9):2121-2136.e6. doi:10.1016/j.immuni.2023.08.008

17. Gate D, Rezai-Zadeh K, Jodry D, Rentsendorj A, Town T. Macrophages in Alzheimer's disease: the blood-borne identity. J Neural Transm (Vienna). 2010;117(8):961-970. doi:10.1007/s00702-010-0422-7

18. Town T, Laouar Y, Pittenger C, et al. Blocking TGF-beta-Smad2/3 innate immune signaling mitigates Alzheimer-like pathology. Nat Med. 2008;14(6):681-687. doi:10.1038/nm1781

19. Hawkes CA, McLaurin J. Selective targeting of perivascular macrophages for clearance of beta-amyloid in cerebral amyloid angiopathy. Proc Natl Acad Sci U S A. 2009;106(4):1261-1266. doi:10.1073/pnas.0805453106

20. Malm TM, Koistinaho M, Pärepalo M, et al. Bone-marrow-derived cells contribute to the recruitment of microglial cells in response to beta-amyloid deposition in APP/PS1 double transgenic Alzheimer mice. Neurobiol Dis. 2005;18(1):134-142. doi:10.1016/j.nbd.2004.09.009

21. Simard AR, Soulet D, Gowing G, Julien JP, Rivest S. Bone marrow-derived microglia play a critical role in restricting senile plaque formation in Alzheimer's disease. Neuron. 2006;49(4):489-502. doi:10.1016/j.neuron.2006.01.022

22. Wang J, Gu BJ, Masters CL, Wang YJ. A systemic view of Alzheimer disease — insights from amyloid-β metabolism beyond the brain. Nat Rev Neurol. 2017;13(10):612-623. doi:10.1038/nrneurol.2017.111

23. Qosa H, Abuasal BS, Romero IA, et al. Differences in amyloid-β clearance across mouse and human blood–brain barrier models: Kinetic analysis and mechanistic

modeling. Neuropharmacology. 2014;79:668-678.
doi:10.1016/j.neuropharm.2014.01.023

24. Xiang Y, Bu XL, Liu YH, et al. Physiological amyloid-beta clearance in the periphery
and its therapeutic potential for Alzheimer's disease. Acta Neuropathol.
2015;130(4):487-499. doi:10.1007/s00401-015-1477-1

25. Eisele YS, Obermüller U, Heilbronner G, et al. Peripherally applied aβ-containing
inoculates induce cerebral β-amyloidosis. Science. 2010;330(6006):980-982.
doi:10.1126/science.1194516

26. Klein SL, Flanagan KL. Sex differences in immune responses. Nat Rev Immunol.
2016;16(10):626-638. doi:10.1038/nri.2016.90

27. Muñoz-Castro C, Mejias-Ortega M, Sanchez-Mejias E, et al. Monocyte-derived cells
invade brain parenchyma and amyloid plaques in human Alzheimer's disease
hippocampus. Acta Neuropathol Commun. 2023;11(1):31. Published 2023 Feb 28.
doi:10.1186/s40478-023-01530-z

28. Paranjpe MD, Belonwu S, Wang JK, et al. Sex-Specific Cross Tissue Meta-Analysis
Identifies Immune Dysregulation in Women With Alzheimer's Disease. Front Aging
Neurosci. 2021;13:735611. Published 2021 Sep 30. doi:10.3389/fnagi.2021.735611

29. Cassado Ados A, D'Império Lima MR, Bortoluci KR. Revisiting mouse peritoneal
macrophages: heterogeneity, development, and function. Front Immunol. 2015;6:225.
Published 2015 May 19. doi:10.3389/fimmu.2015.00225

30. Bain CC, Hawley CA, Garner H, et al. Long-lived self-renewing bone marrow-derived
macrophages displace embryo-derived cells to inhabit adult serous cavities. Nat
Commun. 2016;7:ncomms11852. Published 2016 Jun 13. doi:10.1038/ncomms11852

31. Zajd CM, Ziemba AM, Miralles GM, et al. Bone Marrow-Derived and elicited
peritoneal macrophages are not created equal: the questions asked dictate the cell type
used. Frontiers in Immunology. 2020;11. doi:10.3389/fimmu.2020.00269

32. Wang C, Yu X, Cao Q, et al. Characterization of murine macrophages from bone
marrow, spleen and peritoneum. BMC Immunology. 2013;14(1). doi:10.1186/1471-
2172-14-6

33. Kowalski K, Mulak A. Brain-gut-microbiota axis in alzheimer's disease. J
Neurogastroenterol Motil. 2019;25(1):48-60. doi:10.5056/jnm18087

34. Kitamura J, Nagai M, Ueno H, et al. The insular cortex, Alzheimer disease pathology,
and their effects on blood pressure variability. Alzheimer Disease & Associated
Disorders. 2020;34(3):282-291. doi:10.1097/wad.0000000000000340

35. Koren T, Yifa R, Amer M, et al. Insular cortex neurons encode and retrieve specific
immune responses. Cell. 2021;184(24):5902-5915.e17. doi:10.1016/j.cell.2021.10.013

36. Chen EY, Tan CM, Kou Y, et al. Enrichr: interactive and collaborative HTML5 gene list
enrichment analysis tool. BMC Bioinformatics. 2013;14(1):128. doi:10.1186/1471-
2105-14-128

37. Kuleshov MV, Jones MR, Rouillard AD, et al. Enrichr: a comprehensive gene set
enrichment analysis web server 2016 update. Nucleic Acids Res. 2016;44(W1):W90-
W97. doi:10.1093/nar/gkw377

38. Xie Z, Bailey A, Kuleshov MV, et al. Gene set knowledge discovery with enrichr.
Current Protocols. 2021;1(3):e90. doi:10.1002/cpz1.90

39. Ghosn EE, Cassado AA, Govoni GR, et al. Two physically, functionally, and developmentally distinct peritoneal macrophage subsets. Proc Natl Acad Sci U S A. 2010;107(6):2568-2573. doi:10.1073/pnas.0915000107

40. Bain CC, Gibson DA, Steers NJ, et al. Rate of replenishment and microenvironment contribute to the sexually dimorphic phenotype and function of peritoneal macrophages. Sci Immunol. 2020;5(48):eabc4466. doi:10.1126/sciimmunol.abc4466

41. Zhan X, Stamova B, Jin LW, DeCarli C, Phinney B, Sharp FR. Gram-negative bacterial molecules associate with Alzheimer disease pathology. Neurology. 2016;87(22):2324-2332. doi:10.1212/WNL.0000000000003391

42. Maheshwari P, Eslick GD. Bacterial infection and alzheimer's disease: a meta-analysis. JAD. 2014;43(3):957-966. doi:10.3233/JAD-140621

43. Kang YJ, Kusler B, Otsuka M, et al. Calcineurin negatively regulates tlr-mediated activation pathways. The Journal of Immunology. 2007;179(7):4598-4607. doi:10.4049/jimmunol.179.7.4598

44. Dantzer R. Neuroimmune interactions: from the brain to the immune system and vice versa. *Physiological Reviews*. 2018;98(1):477-504. doi:10.1152/physrev.00039.2016

45. Jin H, Li M, Jeong E, Castro-Martinez F, Zuker CS. A body–brain circuit that regulates body inflammatory responses. Nature. 2024;630(8017):695-703. doi:10.1038/s41586-024-07469-y

46. Zang X, Chen S, Zhu J, Ma J, Zhai Y. The emerging role of central and peripheral immune systems in neurodegenerative diseases. Front Aging Neurosci. 2022;14:872134. doi:10.3389/fnagi.2022.872134

47. Weinstein SL, Gold MR, DeFranco AL. Bacterial lipopolysaccharide stimulates protein tyrosine phosphorylation in macrophages. Proc Natl Acad Sci USA. 1991;88(10):4148-4152. doi:10.1073/pnas.88.10.4148

48. Behar SM, Martin CJ, Booty MG, et al. Apoptosis is an innate defense function of macrophages against Mycobacterium tuberculosis. Mucosal Immunology. 2011;4(3):279-287. doi:10.1038/mi.2011.3

49. Xaus J, Comalada M, Valledor AF, et al. LPS induces apoptosis in macrophages mostly through the autocrine production of TNF-α. *Blood*. 2000;95(12):3823-3831. doi:10.1182/blood.V95.12.3823

50. Soldin OP, Mattison DR. Sex differences in pharmacokinetics and pharmacodynamics: Clinical Pharmacokinetics. 2009;48(3):143-157. doi:10.2165/00003088-200948030-00001

51. U.S. Government Accountability Office. *Drug Safety: Most Drugs Withdrawn in Recent Years Had Greater Health Risks for Women*. 2001. GAO-01-286R. Accessed August 22, 2024.