Title

Direct lineage conversion of postnatal mouse cortical astrocytes to oligodendrocyte lineage cells.

Authors

Justine Bajohr[1-2], Erica Y. Scott[1,3-5], Arman Olfat[1], Mehrshad Sadria[5-6], Kevin Lee[1], Maria Fahim[1], Hiba T. Taha[1,2], Daniela Lozano Casasbuenas[1,7], Ann Derham[1], Scott A. Yuzwa[7], Gary D. Bader[5,8], Maryam Faiz[1,2,7 # $]

Affiliations

1. Department of Surgery, Temerty Faculty of Medicine; University of Toronto; Toronto, Ontario, M5S 1A8, Canada
2. Institute of Medical Science, Temerty Faculty of Medicine; University of Toronto; Toronto, Ontario, M5S 1A8, Canada
3. Institute of Biomedical Engineering, Faculty of Applied Science and Engineering; University of Toronto; Toronto, Ontario, M5S 3G9, Canada
4. Department of Chemistry, Faculty of Arts and Science; University of Toronto; Toronto, Ontario, M5S 3G3, Canada
5. Donnelly Centre for Cellular and Biomolecular Research, Temerty Faculty of Medicine; University of Toronto; Toronto, Ontario, M5S 3E, Canada
6. Department of Applied Mathematics, University of Waterloo, Waterloo, Ontario N2L 3G1, Canada.
7. Department of Laboratory Medicine and Pathobiology, Temerty Faculty of Medicine; University of Toronto; Toronto, Ontario, M5S 1A8, Canada
8. Department of Molecular Genetics, Temerty Faculty of Medicine; University of Toronto; Toronto, Ontario, M5S 1A8
#Correspondence: maryam.faiz@utoronto.ca
$Lead contact: maryam.faiz@utoronto.ca

SUMMARY

Oligodendrocyte lineage cells (OLCs) are lost in many CNS diseases. Here, we investigate the generation of new OLCs via ectopic expression of *Sox10*, *Olig2* or *Nkx6.2* in mouse postnatal astrocytes. Using stringent analyses including, Aldh1l1-astrocyte fate mapping and live cell imaging we confirm that *Sox10* and *Olig2*, but not *Nkx6.2*, convert Aldh1l1[pos] astrocytes to induced OLCs (iOLCs). With single cell RNA sequencing (scRNA-seq) we uncover the molecular signatures of iOLCs. Transcriptomic analysis of *Sox10-* and control cultures over time reveals a clear trajectory from astrocytes to iOLCs. Finally, perturbation models CellOracle and Fatecode support the idea that *Sox10* drives cells towards a terminal iOLC fate. Altogether, this multidimensional analysis shows bonafide conversion of astrocytes to iOLCs using *Sox10* or *Olig2* and provides a foundation for astrocyte DLR strategies to promote OLC repair.

KEYWORDS

Astrocyte, oligodendrocyte, reprogramming, deep learning, single cell RNA sequencing, fate mapping

INTRODUCTION
Oligodendrocytes (OLs) are best known as the myelinating cells of the central nervous
system (CNS). OLs ensheath neuronal axons to enable fast propagation of action potentials.
Consequently, the loss or dysfunction of OLs results in impaired neurological function that
is characteristic of many types of CNS disease and injury. Multiple sclerosis (MS),
Alzheimer's Disease, spinal cord injury, white matter stroke and cerebral palsy are all
characterized by oligodendrocyte failure. Thus, therapeutic strategies aimed at replacing
OLs are of significant clinical interest (1).
9
Direct lineage reprogramming (DLR) aims to generate new target cells lost to disease via the
forced conversion of donor cells. Typically, DLR is performed by the overexpression of
transcription factors (TFs). Early pioneering work from the Tesar Lab demonstrated fibroblast
to induced oligodendrocyte progenitor cell (iOPC) conversion by ectopic expression
of *Olig2*, *Sox10*, and *Nkx6.2*; determinants of OL cell fate in the embryonic brain (2). The
combination of *Sox10* and *Olig2* was also later used to generate iOPCs from pericytes (3).
Therapeutically, however, these newly generated iOPCs still require transplantation into the
brain. Therefore, methods to reprogram endogenous CNS cells would be advantageous.
18
In parallel, astrocytes, CNS-resident cells, have emerged as donor cells in DLR strategies
aimed at generating new neurons (4–7). Astrocytes are an attractive donor cell type for OL
conversion given their shared neural origin (8,9). Astrocytes may already have relevant
epigenetic marks and active TFs that could make DLR faster or more efficient (10–13). In
addition, closely related cells may require fewer TFs for conversion. Indeed, conversion of
astrocytes to an induced oligodendrocyte-like cell was suggested using *Sox10* alone (14).
Therefore, we reasoned that single 'Tesar' factors, *Olig2*, *Sox10*, or *Nkx6.2*, could be used to
force astrocyte conversion to new, induced oligodendrocyte lineage cells (iOLCs).
27
The OL lineage is comprised of oligodendrocyte progenitor cells (OPCs) that give rise to
committed oligodendrocyte progenitors (COPs). These COPs differentiate into mature OLs
(mOLs), comprising at least 6 different states (15). Of interest, during OL development,
Olig2, Sox10, and Nkx6.2 show different temporal expression and play different roles in OL
fate specification. *Olig2,* long considered an OL lineage fate determinant (16), is also
expressed in astrocytes (17,18), which suggests a broad role in early glial commitment
(19,20). *Sox10* is important throughout OL development; *Sox10* promotes early OL lineage
specification by re-inducing *Olig2* (21) and by inhibiting *Sufu (22)*, but is also required for OL
survival following myelination (23,24). In contrast, *Nkx6.2* is expressed late in OL
development, with myelin genes *Mbp* and *Mog,* and plays a role in regulating myelination
(25,26). Given the unique roles of each of these TFs in development, we further hypothesized
that each of these single factors would have varying efficiency in generating iOLCs.
40

Recent controversy in the field of astrocyte to neuron DLR *in vivo (27)* has highlighted the
need for rigorous reporting of DLR outcomes. In a landmark study using astrocyte fate
mapping strategies, the authors suggested that off-target transduction of endogenous cells
was misrepresented as DLR (27). Therefore, it is important that new DLR paradigms utilize
fate mapping, and stringent, multi-faceted analysis to determine the origin of the newly
generated cells.

Here, we used lentiviral delivery of *Olig2*, *Sox10*, or *Nkx6.2* to investigate single
TF conversion of postnatal (P0-P5) GFAP+ cortical astrocytes to iOLCs. Lineage
tracing experiments using Aldh1l1-CreERT2;Ai14 mice confirmed
that *Sox10* and *Olig2* directly convert Aldh1l1+ astrocytes to iOLCs. Moreover, live cell
imaging, single cell RNA sequencing (scRNA seq) and deep learning methods support the
findings that iOLCs can be generated from astrocytes following TF delivery. Altogether, these
findings show bonafide astrocyte to iOLC DLR and lay the groundwork for future studies
utilizing DLR for diseases involving OLC dysfunction and loss.

METHODS
Animals
All experiments were performed in accordance with approved Animal Use Protocols (AUP
20012151, 25-0389H) from the Division of Comparative Medicine at the University of
Toronto. P0-P5 Ai14 **(**B6;129S6- *Gt(ROSA)26Sor^{tm14(CAG-tdTomato)Hze}*/J, RRID:IMSR_JAX:007908)
and Ai14;Ald1l1-CreERT2 mice (Ai14 crossed to B6N.FVB-Tg(Aldh1l1-
cre/ERT2)1Khakh/J [RRID:IMSR_JAX:031008]) were used to generate postnatal astrocyte
cultures.

Cell Culture
Cortical astrocytes were isolated from male and female P0-P5 mice as previously described
(28). Briefly, mice were decapitated, followed by the removal of the skull and meninges.
Cortices were dissected, pooled, and mechanically dissociated in astrocyte media [DMEM
(Gibco Catalogue No. 10569-010), 10% fetal bovine serum (FBS) (Gibco Catalogue No.
10082147) and 1% penicillin/streptomycin (Gibco Catalogue No. 15140122)]. Cells were
cultured in flasks pre-coated with 10 µg/ml poly-d-lysine (Sigma Catalogue No. P6407), and
incubated at 37°C, 5% CO_2. Media was changed the day following isolation and every other
day thereafter. Once the cells reached 80% confluency, typically after 6 days, flasks were
placed on an orbital shaker for 30 minutes to remove contaminating microglia. Astrocyte
media was replaced. For Ai14;Aldh1l1-CreERT2 cultures, 1uM 4-OHT was added to the
astrocyte media at this step. Flasks were returned to the orbital shaker overnight, followed
by vigorous shaking for one minute to remove contaminating OLCs. Media was removed and
astrocytes were incubated in TrypLE Express Enzyme (Gibco Catalogue No. 12604013) for 5
minutes at 37°C, 5% CO_2 to lift off the astrocytes. To inactivate the enzyme, astrocyte media
was added at a 3:1 ratio (media:TrypLE). Cell suspension was collected and centrifuged at
300 x g for 5 minutes. Following removal of the supernatant, the pellet was resuspended in
astrocyte media. Cells were then plated on poly-l-ornithine/laminin coated coverslips at
either 50,000 or 70,000 cells/well in 24 well plates and incubated at 37°C, 5% CO_2. For live

cell analysis, cells were plated at 10,000 cells/ well in 96 well plates with poly-l-
ornithine/laminin coating. For poly-l-ornithine/laminin coating, 0.1mg/ml poly-l-ornithine
(Sigma Catalogue No. P4957) was added to dishes overnight at 37˚C, washed 3 times with
1X phosphate buffered saline (PBS), and then dishes were incubated for two hours at 37˚C
with 10 µg/ml laminin (Sigma Catalogue No. L2020).
90
Reprogramming
Lentiviral particles were purchased from VectorBuilder. For Ai14 reprogramming, LV-
hGFAP::*Sox10*-P2A-Cre, LV-hGFAP::*Olig2*-P2A-Cre, LV-hGFAP::*Nkx6.2*-P2A-Cre, and control
LV-hGFAP::BFP-T2A-Cre were used. For Ai14;Aldh1l1-CreERT2 reprogramming, LV-
hGFAP::*Sox10*-P2A-zsGreen,   LV-hGFAP::*Olig2*-P2A-zsGreen,   LV-hGFAP::*Nkx6.2*-P2A-
zsGreen and control LV-hGFAP::zsGreen were used. A multiplicity of infection (MOI) of 100
was used for all experiments. Virus-containing astrocyte media was placed on the cells and
left overnight. Viral media was replaced with fresh astrocyte media one day post
transduction (DPT). Three DPT, cells were switched to OPC differentiation media
(2) [DMEM/F12/Glutamax (Gibco Catalogue No. 11330032), 1X N2 (Gibco Catalogue No.
17502048), 1X B27 without vitamin A (Gibco Catalogue No. 17504044), 200 ng/ml SHH (R&D
Systems Catalogue No. 464-SH), 20 ng/ml FGF (R&D Systems Catalogue No. 3139-FB), 4
ng/ml PDGF (Sigma Catalogue No. SRP3228)]. At 10 DPT, the cells were switched to OL
differentiation media (2) [DMEM/F12/Glutamax, 1X N2, 1X B27 without vitamin A, 40 ng/ml
T3   (T2877   Catalogue   No.   Sigma),   200 ng/ml   SHH,
100 ng/ml Noggin (R&D Systems Catalogue No. 1967-NG), 10 µM cAMP (Sigma Catalogue
No. A9501), 100 ng/ml IGF (R&D Systems Catalogue No. 791-MG), 10 ng/ml NT3 (Sigma
Catalogue No. SRP6007)].
109
110
Live Cell Analysis
Astrocytes isolated from Ai14;Aldh1l1-CreERT2 mice were plated in 96 well plates,
transduced and imaged every hour from 7 to 12DPT using the Apotome live cell system
(Zeiss). 25 z-stack tiled images per well were captured with brightfield as well as the 488nm
and 568nm fluorescent wavelength. Images were stitched to create a continuous video for
each well. At 12DPT, cells were fixed and stained for OLC markers to confirm fate. Each well
was then re-imaged at this final timepoint and OLC+ reprogrammed cells were matched to
the live cell video and retrospectively analyzed for starting cell morphology and fluorescent
expression. To characterize starting cell morphology, an analysis was first performed on 70
Aldh1l1+tdTomato+ astrocytes in tamoxifen-labelled Aldh1l1-CreERT2;Ai14 cultures and 70
PDGFRa+ OPCs from iOLC cultures (Table S1). Astrocyte and OPC paramenters were
established for cell size, nucleus size, number of branches and branch thickness
(Supplemental Figure 1). These parameters were then used to characterize starting cells as
astrocytes or OPCs.
125
Immunocytochemistry
Cells were fixed in 4% paraformaldehyde (PFA) (Sigma Catalogue No. P6148) for 20 minutes
followed by three washes with 1X PBS. Cell membranes were permeabilized with 0.1%

Triton-X-100 (Sigma Catalogue No. X100) for 10 minutes at room temperature, followed by
three washes with 1X PBS, and then blocked with 5% milk for one hour at room temperature.
Cells were incubated with primary antibodies in 1X PBS overnight at 4°C, washed three times
with 1X PBS, and then incubated with secondary antibodies and DAPI (Sigma Catalogue No.
D9542) in 1X PBS at room temperature for one hour. Following three final 1X PBS washes,
coverslips were mounted on glass slides (Fisher Scientific Catalogue No. 125523)
with Mowiol mounting solution (Sigma Catalogue No. 81381). For staining of membrane
bound proteins (O4, PDGFRα), no permeabilization step with Triton-X-100 was performed.
Primary antibodies: mouse anti-SOX10 (RRID:AB_10844002, 1:250), rabbit anti-PDGFRα
(RRID:AB_2892065, 1:500), mouse anti-O4 (RRID:AB_357617, 1:1000) and rat anti-
MBP RRID:AB_305869, 1:50).
Secondary antibodies: anti-mouse IgG 488 (Invitrogen Catalogue No. A32723) and 647
(Invitrogen Catalogue No. A32728), anti-mouse IgM heavy chain 488 (Invitrogen Catalogue
No. A21042) and 647 (Invitrogen Catalogue No. A21238), anti-rabbit 488 IgG (Invitrogen
Catalogue No. A11034) and 647 (Invitrogen Catalogue No. A32733), anti-rat IgG 488
(Invitrogen Catalogue No. A21208) and 647 (Invitrogen Catalogue No. A21247) all at 1:1000.
145

Microscopy and Image Analysis
Fluorescent images for quantification were taken on an LSM 880 Elyra Superresolution and
LSM 900 (Zeiss) using a 20x objective and Zen Blue software (Zeiss). Post-acquisition linear
adjustments of brightness for all channels were made to micrographs using the Zen Blue
software in Figure 1C, Figure 2E-G and Supplemental Figure 5D. For quantification, ten
regions of interest were selected at random for each well. Images were analyzed using
ImageJ software (National Institutes of Health, RRID:SCR_003070). Reprogramming
efficiency was calculated as a measure of total OLC marker$^+$reporter$^+$DAPI$^+$ cells over
total reporter$^+$DAPI$^+$cells.
155

scRNA-seq capture and processing
At 14DPT, LV-hGFAP::*Sox10*-P2A-Cre, LV-hGFAP::*Olig2*-P2A-Cre, LV-hGFAP::*Nkx6.2*-P2A-
Cre, and control LV-hGFAP::BFP-T2A-Cre cultures were processed using the BD Rhapsody
System (BD Biosciences) and then sequenced. For single-cell isolation, an average
of 9813.75 viable cells were captured in wells at cell load (Table S2). The BD Rhapsody
scanner reported an average multiplet rate of 10.13% and an average number of wells with
viable cells and a bead of 7081.5 (Table S2). Detailed metrics for each sample can be found
in Table S2. Samples were down-sampled to 2500 cells and carried through and converted
to cDNA using the BD Rhapsody WTA Reagent Kit (Becton Dickinson Canada, Catalogue No.
633802). Each cell was sequenced at approximately 100 million reads per cell (at least
2x150 bp paired-end reads) on a Novaseq (Donnelly Sequencing Centre, University of
Toronto).
In addition, LV-GFAP::*Sox10* and control LV-GFAP::Cre cultures were collected prior to
transduction, at 3DPT and 8DPT, and processed using the BD Rhapsody System (BD
Biosciences) and then sequenced. For single-cell isolation, an average of 10263.8 viable
cells were captured in wells at cell load (Table S3). The BD Rhapsody scanner reported an
average multiplet rate of 6.86% and an average number of wells with viable cells and a bead

of 7785 (Table S3). Detailed metrics for each sample can be found in Table S3. Samples were
down-sampled to 2500 cells and carried through and converted to cDNA using the BD
Rhapsody WTA Reagent Kit (Becton Dickinson Canada, Catalogue No. 633802). Each cell
was sequenced at approximately 100 million reads per cell (at least 2x150 bp paired-end
reads) on a Novaseq (Donnelly Sequencing Centre, University of Toronto).
178
scRNA-seq analysis
Fastq files were first demultiplexed with Kallisto (29) (RRID:SCR_016582)(v0.48.0) and
Bustools (30) (RRID:SCR_018210)(V 0.41.0) using supplied whitelists (Data S1) with the -
BDWTA option and aligning to GRCm38.96 with Cre sequence appended to the end.
Bustools (30) was then used to generate gene count tables. Cells were plotted based upon
UMI counts per barcode, thresholds were selected based on inflection point of UMI count
per barcode plots. These thresholds produced read and gene count distributions that were
comparable between all treatment groups (Supplemental Figure 2A, Supplemental
Figure 6A). Gene count tables were made into S4 objects, scaled, normalized and
dimensions reduced (PCA then UMAP) using the Seurat (31) package (RRID:SCR_016341)
(v4.1). Clusters identified as microglia and as vascular and leptomeningeal cells
(VLMCs) (Supplemental Figure 2B-C [cluster 8], Supplemental Figure 6B-C [cluster 6 and 8])
were removed prior to further analysis.
192
Gene markers for oligodendrocyte lineage cells were adopted from studies observing *in*
vivo mouse oligodendrocyte lineage cells across several areas in young and mature CNS
tissues (15). These markers were converted into percent expression of each UMAP cluster
using the PercentageFeatureSet function from Seurat (31), with further gene resolution
displayed by heatmaps created using ComplexHeatmap (32)(RRID:SCR_017270) (v12.13.1).
An additional set of gene markers, demonstrating similar, but less resolved, conclusions was
also used from an *in vitro* rat study looking at OLCs from the cortex (33). Stacked violin
cluster dot plots were made with scCustomize (34)(RRID:SCR_024675)(v2.1.2).
For pseudotime and trajectory analysis Slingshot (35) (RRID:SCR_017012)(v2.3.1) and
Monocle3 (36–40) (RRID:SCR_018685) were used. For *in silico Sox10* perturbation,
CellOracle (41) (v0.14.0) was used. Plots were produced using ggplot2
(42) (RRID:SCR_014601)(v3.3.5) and figure generating scripts were run in R studio
(43) (v4.2.0), with demultiplexing using Kallisto (29) and Bustools (30) run on a Compute
Canada HPC cluster. All scripts used for processing of scRNA-seq data and for figure
generation can be found at github.com/eyscott/Bajor_BDRhapsodyData_DLR.
208
Deep-learning analysis
Fatecode (44) was used to identify key genes for cellular transition. To optimize the
autoencoder and subsequent classifier configuration, a grid search was conducted to
systematically evaluate various combinations of hyperparameters. These included: latent
layer size, number of nodes in the first and second layers of the autoencoder, classifier
architecture, and type of activation function. The grid search aimed to identify the
hyperparameters that minimized a combined reconstruction and classification loss
function, signifying the optimal performance for our specific dataset. Perturbations on the

latent space were performed and cell classifications as well as genes associated with each
perturbation were obtained.
219

Statistical Analysis
Percentage values were transformed using the arcsine square root transformation and
assessed for normal distribution using the Shapiro-Wilks test. When distribution and
variance were equal, a matched pairs one-way ANOVA ([Ai14:D8, D10, D12 O4, D12 MBP],
[Ai14;Aldh1l1-CreERT2: D12 MBP zsGreen+, D12 MBP tdTomato+) or one-way ANOVA
([Ai14:D14], [Ai14;Aldh1l1-CreERT2: D12 MBP zsGreen+,tdTomato+]) , or paired t-test
([Ai14;Aldh1l1-CreERT2: D12 PDGFRα zsGreen+tdTomato+, D12 PDGFRα tdTomato+,
O4 zsGreen+, O4 tdTomato+]) was performed to compare reprogramming efficiency of TF
groups to a control group (Ai14: LV-GFAP::Cre, Ai14;Aldh1l1-CreERT2: LV-GFAP::zsGreen).
When transformed values did not follow a Gaussian distribution, a Kruskal-Wallis test (Ai14:
D12 PDGFRα, D14 PDGFRα) or Wilcoxon test (Ai14;Aldh1l1-CreERT2: PDGFRα zsGreen+,
O4 zsGreen+tdTomato+) was performed to compare reprogramming efficiency to a control
group (Ai14: LV-GFAP::Cre, Ai14;Aldh1l1-CreERT2: LV-GFAP::zsGreen). In both cases,
Dunnett's post-hoc testing was performed to correct for multiple comparisons. Differences
were considered significant at $p < 0.05$. Values are presented as mean ± SEM. The statistical
software used for transformation, distribution, variance, ANOVA, Kruskal-Wallis, t-test,
Wilcoxon test and Dunnett's analysis was GraphPad Prism version 9.0.1
(RRID:SCR_002798).
238

RESULTS
iOLCs are generated following expression of *Olig2*, *Sox10* or *Nkx6.2*
To investigate astrocyte to OLC conversion, we established postnatal, cortical astrocyte
cultures from Ai14 mice. To understand the purity of our cultures, we quantified the numbers
of contaminating OLCs. Quantification of SOX10+, O4+ and MBP+ OLCs showed less
than 2.5% in our cultures (Supplemental Figure 3A).
245

To examine the reprogramming potential of *Olig2, Sox10,* and *Nkx6.2*, we transduced Ai14
astrocytes with either LV-GFAP::*Olig2,* LV-GFAP::*Sox10,* LV- GFAP::*Nkx6.2* or a control LV-
GFAP::*Cre* (Figure 1A). When we quantified the numbers of tdTomato+ cells that co-
expressed OPC (PDGFRa), COP (O4) or OL (MBP) markers at 8 and 10 DPT, no differences
were seen in any of the TF-transduced cultures compared to controls (Figure 1B). By 12
DPT, *Olig2*- cultures showed an increase in the percentage of tdTomato$^+$PDGFRα$^+$ OPCs (p =
0.0309, H-statistic: 7.514), whereas *Sox10*- and *Nkx6.2*- cultures showed an increase
in tdTomato$^+$MBP$^+$ OLs (*Sox10:* p = 0.0004, *Nkx6.2:* p = 0.0108, F statistic = 13.28, degrees of
freedom =3) (Figure 1B-C). No differences were seen in the percentage of
tdTomato$^+$O4$^+$ COPs in any condition at 12DPT (Figure 1B). However, by 14DPT we observed
an increase in the percentage of tdTomato$^+$O4$^+$ COPs in *Sox10*- cultures when compared to
controls (p = 0.0213, F statistic = 3.512, degrees of freedom = 3) (Figure 1B-C). Finally, to
understand whether a longer time in culture could increase the number of MBP+ OLs, we
cultured cells for an additional 8 days and analyzed the percent of tdTomato$^+$MBP$^+$ cells at
22 DPT. No differences were seen in tdTomato$^+$MBP$^+$ OLs in any condition compared to

controls   (Supplemental   Figure 3B).   Altogether,   these   findings   suggest
that *Olig2*, *Sox10* and *Nkx6.2* increase the percentage of iOLCs relative to controls in Ai14
astrocyte cultures.
264
Canonical OLC cluster found in TF-treated cells following scRNA-seq
To further characterize *Olig2-, Sox10-, Nkx6.2-* and control cultures, we performed scRNA-
seq at 14DPT (Figure 1A). Clustering analysis showed the appearance of nine clusters (Figure
1D). We then used proportion analysis to identify clusters that were unique to TF-treated
samples. Clusters 3 and 8 were predominantly comprised of cells from TF-induced
cultures (Figure 1E-F). The top 10 genes marking each cluster in Figure 1D are highlighted in
Figure 1G. Canonical OLC genes such as *Mbp, Plp1, Bcas1* and *Cldn11* were expressed in
clusters 3 and 8 (Figure 1G), further suggesting that the increase in iOLCs is a result of TF
delivery.
274
Next, we compared the molecular profiles of clusters 3 and 8 to established OL lineage
datasets. First, we used OLC specific annotations of mouse fetal and adult OLCs derived
from Marques *et al* (15) to bin the data. We found that cluster 3 was characterized by
COP signatures, while cluster 8 was characterized by COP, newly formed oligodendrocyte
(nfOL) and myelin forming oligodendrocyte (mfOL) signatures (Supplemental Figure 4A). We
also used OLC annotations of *in vitro* OLCs derived from Dugas *et al* (33) to bin the data.
Again, we saw that cluster 8 was characterized by OL signatures, but cluster 3 was not.
(Supplemental Figure 4B). Altogether, these findings further support the observation of
increased OLC generation in TF-induced Ai14 cultures (Figure 1B).
284
Bonafide astrocyte to OLC conversion is confirmed with astrocyte fate mapping.
We had previously observed contaminating OLCs in our cultures and the presence
of tdTomato⁺OLC marker⁺ cells in our Cre controls. Therefore, to confirm the origin of these
newly generated OLCs, we performed astrocyte fate mapping using Ai14;Aldh1l1-CreERT2
cultures, the current gold standard in the field (27). Prior to transduction, post-natal
cortical Ai14;Aldh1l1-CreERT2 astrocyte cultures were treated with 1uM 4-OHT to
permanently label all *Aldh1l1* expressing cells with tdTomato (Figure 2A). Cultures were
transduced with LV-GFAP::*Sox10*-zsGreen, LV-GFAP::*Olig2*-zsGreen, LV-GFAP::*Nkx6.2*-
zsGreen or a LV-GFAP::zsGreen control (Figure 2A). Any OLCs that were directly converted
by   a   TF   from   these *Aldh1l1* expressing   astrocytes   would   therefore
express zsGreen and tdTomato (Figure 2A). To understand whether the OPCs generated
from *Olig2* overexpression were the product of astrocyte reprogramming, we quantified the
percentage of PDGFRα+zsGreen⁺tdTomato⁺ cells at 12DPT. We observed an increase in
PDGFRα⁺zsGreen⁺tdTomato⁺ OPCs compared to controls, confirming the astrocytic origin
of iOPCs (Figure 2B,E). To assess the origin of *Sox10*-induced COPs we quantified the
percentage of O4⁺zsGreen⁺tdTomato⁺ cells at 14DPT. No increase was observed
in *Sox10* cultures compared to controls (Figure 2C). Finally, to determine the origin of OLs
generated   by *Sox10* or *Nkx6.2,* we   quantified   the   percentage   of
MBP⁺zsGreen⁺tdTomato⁺ OLs   at   12DPT.   We   observed   an   increase   of
MBP⁺zsGreen⁺tdTomato⁺ OLs generated by *Sox10,* but not from *Nkx6.2* (Figure 2D,F). This

suggests that *Olig2* and *Sox10* reprogram *Aldh1l1* astrocytes to iOLCs, while *Nkx6.2*-derived
OLs and *Sox10*-derived COPs previously observed (Figure 1B), are not the result of *Aldh1l1*
astrocyte conversion. To further confirm these findings, we performed live cell imaging
of Ai14;Aldh1l1-CreERT2 *Sox10*-treated and control cultures from 7DPT to 12DPT. At 12DPT,
we observed tdTomato$^+$zsGreen$^+$ cells that co-expressed OLC markers (PDGFRa/O4/MBP+)
and showed OPC-like morphologies (small nucleus, multiple thin branches) (Figure 2G,
Supplemental Figure 1). When we tracked these cells back to 7DPT, we found that they
showed a characteristic astrocyte morphology (large cell body and nucleus, with minimal
branching) (Figure 2G, Supplemental Figure 1, Supplemental Video 1). Taken together, these
findings further demonstrate bonafide astrocyte to iOLC conversion.
315
We also examined the numbers of Aldh1l1$^+$ non-transduced (tdTomato$^+$only) cells, as well
as Aldh1l1^{neg} virally transduced (zsGreen$^+$ only) cells (Supplemental Figure 5). No difference
in the percentage of OLC marker$^+$tdTomato$^+$zsGreen^{neg} was seen between TF-treated and
control cultures (Supplemental Figure 5). However, when we examined
zsGreen$^+$tdTomato^{neg} cells, we observed an interesting increase in the percentage of
MBP$^+$zsGreen$^+$tdTomato^{neg} OLs at 12DPT in *Sox10*-treated cells compared to controls
(Supplemental Figure 5C). When we analyzed our live cell imaging experiment to understand
the origin of these cells, we found examples of tdTomato^{neg}zsGreen$^+$OLCmarker$^+$ cells that
arose from a tdTomato^{neg} cell (Supplemental Figure 5D). These tdTomato^{neg} cells did not have
OPC morphology 7DPT (Supplemental Figure 1,5D, Supplemental Video 2). Rather, they
showed morphologies typical of astrocytes (Supplemental Figure 1,5D, Supplemental Video
2), suggesting the conversion of an astrocyte- like Aldh1l1^{neg} cell.
328
Characterization of *Sox10*-mediated DLR using scRNA-seq
Given that *Sox10* increased the number of mOLs from *Aldh1l1* astrocytes (Figure 2D,F), we
next probed the mechanisms underlying *Sox10*-mediated reprogramming. We
performed scRNA-sequencing at three additional timepoints (D0 prior to transduction,
3DPT, and 8DPT) on Cre-control and *Sox10*-treated cultures. For analysis, we combined
cells from D0, 3DPT, 8DPT and 14DPT (Supplemental Figure 6B). We excluded cells with
microglia and VLMC markers (Supplemental Figure 6C). Clustering analysis of the remaining
cells showed the appearance of 10 distinct clusters (Figure 3A). We characterized these
clusters based on expression of genes associated with astrocytes, oligodendrocytes, and
NG2 cells, as well as proliferation markers (Figure 3B-C).
339
First, we examined the molecular identities of cells in our starting cultures. D0 cells were
found in clusters 1, 3, 5 and 9 and showed expression of canonical astrocyte markers,
including *Slc1a3, Aqp4, Atp1b2, Glul* and *Gfap* (Figure 3B,D). Of interest, clusters 1, 3 and 5
showed expression of *Aldh1l1*, but this gene was absent in cluster 9 (Figure 3B), which could
explain the conversion of tdTomato^{neg} cells observed in our fate mapping experiments
(Supplemental Figure 5). Altogether, this suggests that early postnatal cultures are
heterogeneous and comprised of distinct astrocyte-like populations (clusters).
347

To understand the cell fate transitions that occur over the reprogramming timecourse, we
performed trajectory analysis with Slingshot (35) (Figure 3E) and Monocle3 (36–40) (Figure
3F). Slingshot and Monocle3 both reconstructed trajectories with clusters 6 and 4 as
terminal branches (Figure 3E-F). We then examined the origin of cells comprising each
cluster. As expected, the clusters formulating earlier roots of the Monocle3 branches
(clusters 0, 1, 3, 9) were primarily comprised of D0, 3DPT, and 8DPT timepoints (Figure 3G).
Clusters 6 and 4, were predominantly comprised of 14DPT samples, and in particular, cells
from *Sox10*-treated samples (D14_S) (Figure 3G). Analysis of the differentially expressed
genes (DEGs) at these branches showed that cluster 6 and branch 14 of the Monocle3
trajectory were enriched in the OLC genes *Sox10* (21–24) (p- and q-value=0), *Bcas1* (*45)* (p-
and q-value=0), and *Omg* (46,47) (p- and q-value=0 (Supplemental Figure 7A). In contrast,
cluster 4 and the Monocle trajectory branch 11 showed expression of the genes *Lcn2* (p- and
q-value=0) and *Igfbp5* (p- and q-value=0), (Supplemental Figure 7B) previously shown to be
expressed in reactive astrocytes (48–50). All Monocle3-derived differentially expressed
genes can be found in Data S2. Altogether, these analyses suggest that the trajectory
to branch 14/cluster 6 represents the path of astrocyte to OLC conversion.

To determine how *Sox10* would influence the gene regulatory network of cells in clusters 6
and 4, we performed *in silico* perturbation of *Sox10* using CellOracle (41). *in silico* knock out
(KO) of *Sox10* predicted a large shift away from a cluster 6 identity and little change away
from the identity of cluster 4 (Figure 3H). In agreement, *in silico* knock in (KI)
of *Sox10* showed a large shift towards a cluster 6 identify and little change towards a cluster
4 identity (Figure 3I). This suggests that leaf 14 and cluster 6 represent cells with a gene
regulatory network most affected by *Sox10*.

As cluster 6 was comprised of both *Sox10*-treated and Cre-control cells, we then performed
a differential gene expression analysis of 14DPT Cre-control and *Sox10*-treated cells within
this cluster. 175 DEGs were found, with the top genes (expressed in at least 50% of Cluster
6, *Sox10*-treated cells and with a difference of at least 0.5 pct between Cre-control
and *Sox10*-treated cultures) clustered and provided in a dotplot (Figure 3J). This analysis
highlighted that genes including *Omg*, *Bcas1*, *Pdgfra*, *Cldn11 (51)*, and *Sox10* are enriched
in *Sox10*-treated cells compared to Cre-control cells, suggesting that *Sox10*-treated cells
are more representative of OLCs than the Cre-control cells exposed to OPC and OL media
alone.

Understanding the genetic drivers of astrocyte to OLC DLR
Our analysis showed that *Sox10* was important for determining the OLC identity of cluster 6.
We specifically chose *Sox10* based on its role in OLC fate specification and development
(21–24). To understand whether there might be other [better] candidate genes that would
promote an OLC identity, we used an unbiased, deep learning perturbation model, Fatecode
(44), to predict genes that would allow cells to shift from a cluster 4 identity (astrocytes at
14DPT, not fully reprogrammed) to a cluster 6 identity (OLCs at 14DPT, end state of
DLR). Following training of the Fatecode model on our *Sox10* and Cre-control treated DLR
dataset, we identified perturbation on node 16 of the latent layer as one which increased the

number of cells in cluster 6 whilst simultaneously reducing the number of cells in clusters
0, 2, 4, 5 and 7. (Figure 3K). Given that the total number of cells in the dataset does not
change, this suggests that the perturbation on node 16 of the latent layer was pushing cells
that did not reprogram (astrocytes at 8 and 14DPT) towards a reprogrammed (OLCs at
14DPT) fate. To identify the genes driving this shift,we ranked the absolute value of gene
expression changes to obtain a list of 3000 genes involved in this perturbation, with the top
40 genes listed in Table S4. Strikingly, we observed that *Sox10* was ranked as the ninth most
correlated gene and the top TF driving this shift (Table S4). This supports our previous
findings that suggest *Sox10* drives astrocyte to iOLC DLR. Additionally, we observed genes
important for myelination (*Plp1, Mbp*) (52), as well as other TFs previously identified with
OLC differentiation and myelination (*Sox3* (53), *Klf9* (54)) as integral to this shift (Table S4).
Taken together, these findings further support the importance
of *Sox10* in reprogramming astrocytes to iOLCs and identify additional genes and TFs that
may be ideal candidates for astrocyte to iOLC reprogramming.
406
DISCUSSION
Here, we show astrocyte to iOLC conversion with *Sox10* or *Olig2* using a battery of
experimental tools, including astrocyte fate mapping, live cell imaging, scRNA-
seq timecourse and unbiased deep learning. While previous studies demonstrated the
generation of iOLCs from different types of somatic cells using combinations
of *Sox10*, *Olig2* and *Nkx6.2* (2,3,55), ours is the first to compare the individual
reprogramming ability of each these TFs in astrocytes. Using *Aldh1l1*-based fate-mapping,
we found that that *Sox10* and *Olig2* convert Aldh1l1$^+$ astrocytes to iOLCs.
415
In contrast to *Sox10* and *Olig2*, *Nkx6.2* was unable to convert *Aldh1l1$^+$*astrocytes to new
iOLCs. Unlike *Sox10* and *Olig2*, which define oligodendrocyte identity and continue to shape
their gene regulatory network throughout life (56), *Nkx6.2* may be unable to direct a stable
iOLC identity. In support of this, previous work investigating pericyte to OL reprogramming
found that inclusion of *Nkx6.2* in their reprogramming cocktail was refractory to
reprogramming (3). In addition to observing the activation of genes unrelated
to oligodendrogenesis, genes required for OPC identity were downregulated when cells were
transduced with *Nkx6.2* (3).
424
In a previous study, Khanghahi *et al*. investigated *Sox10*-mediated astrocyte to OLC
conversion. Ectopic expression of *Sox10* in astrocytes *in vitro* led to an increase in OPCs at
21DPT (14), a different OLC type and longer time to conversion than we observed in our
study. Although similar experimental designs were used in both studies (cortical P3-P5
astrocytes cultured in OPC media in Khanghahi *et al*. versus P0-P5 cortical astrocytes
cultured in OPC followed by OL media in our study), these discrepancies may be due to the
use of different viral delivery strategies and metrics of reprogramming. In our study, we used
lentiviral delivery of *Sox10*-P2A-zsGreen under the control of the long
(2178bp) hGFAP promoter (57). In addition, we used Ai14;Aldh1l1-CreERT2 mice and
quantified only the iOLCs with an astrocyte origin (based
on tdTomato$^+$OLC marker$^+$expression). In contrast, in the Khanghahi *et al*. study, the authors

used a SFFV promoter to deliver *Sox10*-IRES-GFP and reported the number of GFP+ iOLCs in
their *Sox10*-transduced cultures. Without an astrocyte specific promoter and stringent
lineage tracking, it is difficult to conclude that iOPCs were the result of astrocyte conversion.
A non-specific delivery strategy could instead hit a contaminating, perhaps more distantly
related cell that would need more time to convert.
441
In this regard, a recent study suggested that the conversion reported in studies of *in*
vivo astrocyte to neuron DLR (6,58,59) was not true reprogramming, but rather the result of
erroneous labelling of endogenous neurons due to technical confounds (27). As a result, the
DLR community has advocated for the stringent validation of DLR paradigms. In this study,
we first examined astrocyte to iOLC conversion in Ai14 cells. This enabled the permanent
labeling of transduced cells and therefore, the tracking of those cells through the
DLR timecourse. To then validate our findings, we used Aldh1l1-astrocyte fate mapping,
which highlighted a lack of conversion with *Nkx6.2*.
450
When performing our lineage tracing experiments, we also discovered a subpopulation of
cells (*Aldh1l1*-tdTomato^{neg}zsGreen^{+}OLCmarker^{+}) that was converted by *Sox10* at a high
reprogramming efficiency (Supplemental Figure 5C). scRNA-seq analysis of cells prior to
conversion also showed a cluster with low *Aldh1l1* expression compared to the other
astrocyte clusters, but with similar levels of *Gfap* expression (Figure 3B), which could
explain its transduction by the LV-GFAP::*Sox10*. These *Aldh1l1*^{lo}*Gfap*^{hi} cells were
predominantly characterized by mature astrocyte markers but also showed expression of
genes found in NG2 glia (*Cspg, Pdgfra*) (Figure 3B-C). Curiously, this population did not show
expression of proliferation marker *Mki67* (Figure 3C), in contrast to previous studies showing
that astrocyte-like NG2 glia are proliferative (60,61). Of interest, one study reported high co-
expression of *Gfap*, *Cspg4* and *Pdgfra* in early postnatal astrocytes (62). This leads to the
intriguing idea that these cells represent a population of *Aldh1l1*^{lo}expressing astrocytes. If
so, *Aldh1l1*-fate-mapping would underestimate reprogramming efficiencies. The presence
of these *Aldh11l*^{lo} cells shows that astrocyte cultures are heterogeneous and that different
types of astrocytes or astrocyte-like cells may be suitable targets for DLR. Further studies
using scRNA barcoding and CellOracle to profile the different types of astrocytes that are
amenable to reprogramming will be beneficial in designing more specific and tailored DLR
strategies.
469
Conversion of *Aldh1l1*^{pos} astrocytes to PDGFRa^{+}tdTomato^{+}zsGreen^{+} iOPCs using *Olig2*
occurred at a relatively low rate, with an average conversion efficiency of 16.75% cells. This
conversion was even lower for the generation of mature MBP^{+}tdTomato^{+}zsGreen^{+}OLs by
Sox10, with an average conversion efficiency of 2.83%. This may suggest that
although *Sox10* may generate more mature OLCs, it can only do this in a select number of
'elite' donor cells. Alternatively, the absence of a substrate to myelinate in our cultures may
preclude the true reprogramming ability of *Sox10*. OL survival *in vitro* and *in vivo* has been
shown to be dependent on the presence of axons (63). Furthermore, previous studies have
shown that mRNA expression of myelin genes is increased when OLs are cultured in the
presence of neurons (64), and differentiation of OPCs can be induced with bead or nanofiber

scaffolding (65,66). However, additional factors, or the discovery of "better" TFs
may also allow for increased reprogramming efficiency. It remains to be determined whether
the genes predicted by Fatecode in this study will improve the efficiencies of OLC
generation. Nonetheless, it is important to note that the extent of iOLC generation must be
balanced with the extent of astrocyte loss as many studies have shown that widespread loss
of astrocytes can be detrimental (67,68).
486
Investigation of the reprogramming trajectory of *Sox10* and Cre-control treated cells showed
a bifurcation, where the cells either converted to iOLCs or remained as astrocytes (Figure
3F). It remains to be determined if this is a genetic hurdle or a metabolic hurdle, similar to
studies of astrocyte to neuron conversion (69). Future studies investigating the genes that
characterize the 'tipping point' of conversion will be useful in identifying the mechanisms of
astrocyte to iOLC reprogramming. A clearer understanding of how different astrocyte types
convert to iOLCs and the mechanisms that underly the reprogramming processes will help
us identify the best DLR paradigm for therapeutic applications.
495

ACKNOWLEDGMENTS
The authors are grateful to Dr. Lindsey Fiddes at the Microscope Imaging Facility for
assistance with live cell imaging. Next-generation sequencing was performed at the
Donnelly
Sequencing Facility. J.B. was supported by the Baden Havard Endowment Fund, the Ontario
Graduate Scholarship and the MITACs Accelerate Fellowship. This work was supported by
grants to M.F. from CIHR (PJT-175254), Medicine by Design (Cycle 2
Funding), Temerty Foundation (Pathway Grant), Ontario Institute of Regenerative Medicine
(Kickstart Innovation Investment Program), the Stem Cell Network (Early Career Researcher
Jump Start Award), the Connaught Foundation (Innovation Grant), CFA-JELF program for
microscopy infrastructure, and to S.A.Y. from the CFI-JELF program for scRNA-seq related
infrastructure.

DATA AVAILABILITY
RNA sequence data is deposited in GEO (accession number
GSE263185). Genetables are provided as a Cyvers link through the Github page (see
methods) housing the scripts used in the manuscript.
513

AUTHOR CONTRIBUTIONS
Conceptualization, M.F.; Methodology, J.B., G.D.B., M.F.; Software, E.Y.S., M.S.; Formal
Analysis, J.B., E.Y.S., M.S., S.A.Y., M.F.; Investigation, J.B., E.Y.S., A.O., M.S., K.L., M. Fahim,
H.T.T., D.L.C., A.D., and S.A.Y.; Writing – Original Draft, J.B. and E.Y.S.; Writing – Review and
Editing, M.F.; Visualization, J.B., E.Y.S., M.S.; Supervision, G.D.B., M.F.; Funding
Acquisition, S.A.Y., M.F.
520

521 **COMPETING INTERESTS**

FIGURE LEGENDS

Figure 1. *Sox10*, *Olig2* and *Nkx6.2* convert GFAP+ cells to oligodendrocyte lineage
cells. (A) Experimental design and timeline. **(B)** Quantification of PDGFRα⁺tdTomato⁺iOPCs,
O4⁺tdTomato⁺ iCOPs and MBP⁺tdTomato⁺ iOLs at 8 (n=5), 10 (n=4), 12 (n=5) and 14 (n=7)
days post transduction (DPT). Data are presented as mean ± SEM, each data point
represents one individual cell culture experiment; at each time point and for each cell type
marker, a matched pairs one-way ANOVA or Kruskal-Wallis (d8, d10, d12) or one- way ANOVA
(d14) was performed with Dunnet's post testing (*= p<0.05, *** = p<
0.001). **(C)** Representative images of PDGFRα⁺tdTomato⁺,O4⁺tdTomato⁺ and
MBP⁺tdTomato⁺ cells at 12DPT. Single channel images are shown of the boxed cells (arrows
indicate double positive cells). Scale bar = 50um. **(D)** UMAP clustering of *Olig2*-, *Sox10*-
, *Nkx6.2*- and control (Cre) transduced cells at 14DPT **(E)** UMAP of (D) overlaid with
treatment (*Olig2*-, *Sox10*-, *Nkx6.2*- and control (Cre)) **(F)** Proportion analysis of clusters
found in *Olig2*-, *Sox10*-, *Nkx6.2*-, and control (Cre) transduced cells. **(G)** Heatmap of top
upregulated genes from each cluster in (D).

Figure 2. Lineage tracing confirms true conversion of astrocytes to oligodendrocyte
lineage cells.
(A) Experimental design, timeline and outcomes. **(B)** Quantification
of tdTomato⁺zsGreen⁺PDGFRα⁺ iOPCs at 12DPT (n= 3). Data are presented as mean ± SEM,
each data point represents one individual cell culture, a paired t-test was performed (** =
p<0.01). **(C)** Quantification of tdTomato⁺zsGreen⁺O4⁺ iCOPs at 14DPT (n = 4). Data
are presented as mean ± SEM, each data point represents one individual cell
culture, a Wilcoxon test was performed (ns). **(D)** Quantification
of tdTomato⁺zsGreen⁺MBP⁺ iOLs at 12DPT (n= 4 for *Sox10, Nkx6.2*, n= 3 for *Cre*). Data are
presented as mean ± SEM, each data point represents one individual cell culture, a one-way
ANOVA with Dunnet's post testing was performed (* = p<0.05). **(E)** Representative image of
PDGFRα⁺tdTomato⁺zsGreen⁺ cells 12DPT. Single channel images are shown of the boxed
cells (arrows indicate triple positive cells, scale bar = 50um (merge) and 20um (single
channel)). **(F)** Representative image of MBP⁺tdTomato⁺zsGreen⁺ cells 12DPT. Single channel
images are shown of the boxed cells (arrows indicate triple positive cells, scale bar = 50um
(merge) and 20um (single channel)). **(G)** Representative tdTomato⁺zsGreen⁺ cell with
astrocyte-like morphology at onset (7DPT) and OLC expression at the end (12DPT) of live cell
tracking (arrow indicates tracked cell, scale bar = 50um).

Figure 3. Characterization of DLR using scRNA-seq shows terminal oligodendrocyte
cluster of cells at day 14 driven by *Sox10*. (A) UMAP clustering
of *Sox10* and *Cre* control treated cells from prior to transduction, as well as 3, 8 and
14DPT **(B)** Canonical astrocyte gene expression (log-normalized, y-axis) separated by

cluster (x-axis) and coloured by timepoint and treatment group. **(C)** Canonical gene expression (log-normalized, y-axis) of NG2 glia, VLMC, OLC, microglia and proliferation, separated by cluster (x-axis) and coloured by timepoint and treatment group. **(D)** UMAP clustering from (A) overlayed with timepoint and treatment group **(E)** Slingshot lineage analysis of *Sox10* and *Cre* control treated clusters overlaid with UMAP embeddings. **(F)** Monocle3 lineage analysis of *Sox10* and *Cre* control treated clusters overlaid on UMAP plot and coloured by pseudotime predictions. **(G)** The number of cells (y-axis) per cluster (x-axis) originating from each of the coloured timepoint+treatmentgroups. **(H)** CellOracle modeling of *in silico Sox10* knock out (KO) overlaid onto UMAP plot. Arrows indicate trajectory prediction with *Sox10* KO. **(I)** CellOracle modeling of *in silico Sox10* knock in (KI) overlaid onto UMAP plot. Arrows indicate trajectory prediction with *Sox10* KI **(J)** Clustered, top differentially expressed genes (dots, y-axis) between 14DPT *Sox10 (D14_S,* Beige, left*)* and *Cre (D14_C,* Blue, right*)* control treated cells from cluster 6. Size of dot is scaled to percent of cells in the cluster expressing that gene, colour of dot represents the average scaled expression of the gene across cells. **(K)** Predicted cell types following Fatecode perturbation on node 16 of the latent layer in the *Sox10* and control treated dataset. Blue represents the number of cells prior to perturbation. Orange (overlayed) represents the number of cells following perturbation.

REFERENCES

1. Bajohr J, Faiz M. Direct Lineage Reprogramming in the CNS. In: Turksen K, editor. Cell Biology and Translational Medicine, Volume 6: Stem Cells: Their Heterogeneity, Niche and Regenerative Potential [Internet]. Cham: Springer International Publishing; 2020 [cited 2024 Mar 15]. p. 31–48. Available from: https://doi.org/10.1007/5584_2019_374

2. Najm FJ, Lager AM, Zaremba A, Wyatt K, Caprariello AV, Factor DC, et al. Transcription factor–mediated reprogramming of fibroblasts to expandable, myelinogenic oligodendrocyte progenitor cells. Nat Biotechnol. 2013 May;31(5):426–33.

3. Kim KP, Li C, Bunina D, Jeong HW, Ghelman J, Yoon J, et al. Donor cell memory confers a metastable state of directly converted cells. Cell Stem Cell. 2021 Jul 1;28(7):1291-1306.e10.

4. Heinrich C, Blum R, Gascón S, Masserdotti G, Tripathi P, Sánchez R, et al. Directing astroglia from the cerebral cortex into subtype specific functional neurons. PLoS Biol. 2010 May 18;8(5):e1000373.

5. Heins N, Malatesta P, Cecconi F, Nakafuku M, Tucker KL, Hack MA, et al. Glial cells generate neurons: the role of the transcription factor Pax6. Nat Neurosci. 2002 Apr;5(4):308–15.

6.   Guo Z, Zhang L, Wu Z, Chen Y, Wang F, Chen G. In vivo direct reprogramming of reactive
glial cells into functional neurons after brain injury and in an Alzheimer's disease
model. Cell Stem Cell. 2014 Feb 6;14(2):188–202.

7.   Liu Y, Miao Q, Yuan J, Han S, Zhang P, Li S, et al. Ascl1 Converts Dorsal Midbrain
Astrocytes into Functional Neurons In Vivo. J Neurosci. 2015 Jun 24;35(25):9336–55.

8.   Levison SW, Goldman JE. Both oligodendrocytes and astrocytes develop from
progenitors in the subventricular zone of postnatal rat forebrain. Neuron. 1993
Feb;10(2):201–12.

9.   Kriegstein A, Alvarez-Buylla A. The glial nature of embryonic and adult neural stem
cells. Annu Rev Neurosci. 2009;32:149–84.

10. Masserdotti G, Gascón S, Götz M. Direct neuronal reprogramming: learning from and
for development. Development. 2016 Jul 15;143(14):2494–510.

11. Hörmanseder E. Epigenetic memory in reprogramming. Current Opinion in Genetics &
Development. 2021 Oct 1;70:24–31.

12. Bussmann LH, Schubert A, Manh TPV, Andres LD, Desbordes SC, Parra M, et al. A
Robust and Highly Efficient Immune Cell Reprogramming System. Cell Stem Cell. 2009
Nov 6;5(5):554–66.

13. Xie H, Ye M, Feng R, Graf T. Stepwise reprogramming of B cells into macrophages. Cell.
2004 May 28;117(5):663–76.

14. Mokhtarzadeh Khanghahi A, Satarian L, Deng W, Baharvand H, Javan M. In vivo
conversion of astrocytes into oligodendrocyte lineage cells with transcription factor
Sox10; Promise for myelin repair in multiple sclerosis. PLoS One. 2018 Sep
13;13(9):e0203785.

15. Marques S, Zeisel A, Codeluppi S, van Bruggen D, Mendanha Falcão A, Xiao L, et al.
Oligodendrocyte heterogeneity in the mouse juvenile and adult central nervous system.
Science. 2016 Jun 10;352(6291):1326–9.

16. Zhou Q, Wang S, Anderson DJ. Identification of a novel family of oligodendrocyte
lineage-specific basic helix-loop-helix transcription factors. Neuron. 2000
Feb;25(2):331–43.

17. Wang H, Xu L, Lai C, Hou K, Chen J, Guo Y, et al. Region-specific distribution of Olig2-
expressing astrocytes in adult mouse brain and spinal cord. Molecular Brain. 2021 Feb
17;14(1):36.

18. Buffo A, Vosko MR, Ertürk D, Hamann GF, Jucker M, Rowitch D, et al. Expression pattern
of the transcription factor Olig2 in response to brain injuries: Implications for neuronal
repair. Proc Natl Acad Sci U S A. 2005 Dec 13;102(50):18183–8.

19. Marshall CAG, Novitch BG, Goldman JE. Olig2 directs astrocyte and oligodendrocyte
formation in postnatal subventricular zone cells. J Neurosci. 2005 Aug 10;25(32):7289–
98.

20. Tatsumi K, Isonishi A, Yamasaki M, Kawabe Y, Morita-Takemura S, Nakahara K, et al.
Olig2-Lineage Astrocytes: A Distinct Subtype of Astrocytes That Differs from GFAP
Astrocytes. Front Neuroanat. 2018;12:8.

21. Liu Z, Hu X, Cai J, Liu B, Peng X, Wegner M, et al. Induction of oligodendrocyte
differentiation by Olig2 and Sox10: evidence for reciprocal interactions and dosage-
dependent mechanisms. Dev Biol. 2007 Feb 15;302(2):683–93.

22. Pozniak CD, Langseth AJ, Dijkgraaf GJP, Choe Y, Werb Z, Pleasure SJ. Sox10 directs
neural stem cells toward the oligodendrocyte lineage by decreasing Suppressor of
Fused expression. Proc Natl Acad Sci U S A. 2010 Dec 14;107(50):21795–800.

23. Takada N, Kucenas S, Appel B. Sox10 is necessary for oligodendrocyte survival
following axon wrapping. Glia. 2010 Jun;58(8):996–1006.

24. Stolt CC, Rehberg S, Ader M, Lommes P, Riethmacher D, Schachner M, et al. Terminal
differentiation of myelin-forming oligodendrocytes depends on the transcription factor
Sox10. Genes Dev. 2002 Jan 15;16(2):165–70.

25. Awatramani R, Scherer S, Grinspan J, Collarini E, Skoff R, O'Hagan D, et al. Evidence
That the Homeodomain Protein Gtx Is Involved in the Regulation of Oligodendrocyte
Myelination. J Neurosci. 1997 Sep 1;17(17):6657–68.

26. Cai J, Zhu Q, Zheng K, Li H, Qi Y, Cao Q, et al. Co-localization of Nkx6.2 and Nkx2.2
Homeodomain Proteins in Differentiated Myelinating Oligodendrocytes. Glia. 2010
Mar;58(4):458–68.

27. Wang LL, Serrano C, Zhong X, Ma S, Zou Y, Zhang CL. Revisiting astrocyte to neuron
conversion with lineage tracing in vivo. Cell. 2021 Oct 14;184(21):5465-5481.e16.

28. McCarthy KD, de Vellis J. Preparation of separate astroglial and oligodendroglial cell
cultures from rat cerebral tissue. J Cell Biol. 1980 Jun;85(3):890–902.

29. Bray NL, Pimentel H, Melsted P, Pachter L. Near-optimal probabilistic RNA-seq
quantification. Nat Biotechnol. 2016 May;34(5):525–7.

30. Melsted P, Booeshaghi AS, Liu L, Gao F, Lu L, Min KH (Joseph), et al. Modular, efficient
and constant-memory single-cell RNA-seq preprocessing. Nat Biotechnol. 2021
Jul;39(7):813–8.

31. Hao Y, Hao S, Andersen-Nissen E, Mauck WM, Zheng S, Butler A, et al. Integrated
analysis of multimodal single-cell data. Cell. 2021 Jun 24;184(13):3573-3587.e29.

32. Gu Z, Eils R, Schlesner M. Complex heatmaps reveal patterns and correlations in
multidimensional genomic data. Bioinformatics. 2016 Sep 15;32(18):2847–9.

33. Dugas JC, Tai YC, Speed TP, Ngai J, Barres BA. Functional genomic analysis of
oligodendrocyte differentiation. J Neurosci. 2006 Oct 25;26(43):10967–83.

34. Marsh S. scCustomize:Custom Visualizations & Functions for Streamlined Analyses of
Single Cell Sequencing. [Internet]. 2021. Available from:
https://doi.org/10.5281/zenodo.5706430

35. Street K, Risso D, Fletcher RB, Das D, Ngai J, Yosef N, et al. Slingshot: cell lineage and
pseudotime inference for single-cell transcriptomics. BMC Genomics. 2018 Jun
19;19(1):477.

36. Trapnell C, Cacchiarelli D, Grimsby J, Pokharel P, Li S, Morse M, et al. The dynamics and
regulators of cell fate decisions are revealed by pseudotemporal ordering of single
cells. Nat Biotechnol. 2014 Apr;32(4):381–6.

37. Qiu X, Mao Q, Tang Y, Wang L, Chawla R, Pliner HA, et al. Reversed graph embedding
resolves complex single-cell trajectories. Nat Methods. 2017 Oct;14(10):979–82.

38. Cao J, Spielmann M, Qiu X, Huang X, Ibrahim DM, Hill AJ, et al. The single cell
transcriptional landscape of mammalian organogenesis. Nature. 2019
Feb;566(7745):496–502.

39. Traag VA, Waltman L, van Eck NJ. From Louvain to Leiden: guaranteeing well-connected
communities. Sci Rep. 2019 Mar 26;9(1):5233.

40. Levine JH, Simonds EF, Bendall SC, Davis KL, Amir E ad D, Tadmor M, et al. Data-driven
phenotypic dissection of AML reveals progenitor-like cells that correlate with prognosis.
Cell. 2015 Jul 2;162(1):184–97.

41. Kamimoto K, Stringa B, Hoffmann CM, Jindal K, Solnica-Krezel L, Morris SA. Dissecting
cell identity via network inference and in silico gene perturbation. Nature. 2023
Feb;614(7949):742–51.

42. Wickham H. ggplot2: Elegant Graphics for Data Analysis [Internet]. New York, NY:
Springer; 2009 [cited 2024 Mar 16]. Available from:
https://link.springer.com/10.1007/978-0-387-98141-3

43. RStudio: Integrated Development for R [Internet]. 2020. Available from:
http://www.rstudio.com/

44. Sadria M, Layton A, Goyal S, Bader GD. Fatecode: Cell fate regulator prediction using
classification autoencoder perturbation [Internet]. bioRxiv; 2022 [cited 2024 Mar 15]. p.
2022.12.16.520772. Available from:
https://www.biorxiv.org/content/10.1101/2022.12.16.520772v1

45. Fard MK, van der Meer F, Sánchez P, Cantuti-Castelvetri L, Mandad S, Jäkel S, et al.
BCAS1 expression defines a population of early myelinating oligodendrocytes in
multiple sclerosis lesions. Sci Transl Med. 2017 Dec 6;9(419):eaam7816.

46. Vourc'h P, Andres C. Oligodendrocyte myelin glycoprotein (OMgp): evolution, structure
and function. Brain Res Brain Res Rev. 2004 May;45(2):115–24.

47. Mikol DD, Stefansson K. A phosphatidylinositol-linked peanut agglutinin-binding
glycoprotein in central nervous system myelin and on oligodendrocytes. J Cell Biol.
1988 Apr 1;106(4):1273–9.

48. Bi F, Huang C, Tong J, Qiu G, Huang B, Wu Q, et al. Reactive astrocytes secrete lcn2 to
promote neuron death. Proceedings of the National Academy of Sciences. 2013 Mar
5;110(10):4069–74.

49. Liddelow SA, Guttenplan KA, Clarke LE, Bennett FC, Bohlen CJ, Schirmer L, et al.
Neurotoxic reactive astrocytes are induced by activated microglia. Nature. 2017 Jan
26;541(7638):481–7.

50. Lee WH, Wang GM, Seaman LB, Vannucci SJ. Coordinate IGF-I and IGFBP5 Gene
Expression in Perinatal Rat Brain after Hypoxia-Ischemia. J Cereb Blood Flow Metab.
1996 Mar 1;16(2):227–36.

51. Morita K, Sasaki H, Fujimoto K, Furuse M, Tsukita S. Claudin-11/OSP-based Tight
Junctions of Myelin Sheaths in Brain and Sertoli Cells in Testis. J Cell Biol. 1999 May
3;145(3):579–88.

52. Kister A, Kister I. Overview of myelin, major myelin lipids, and myelin-associated
proteins. Front Chem [Internet]. 2023 Feb 21 [cited 2024 Mar 17];10. Available from:
https://www.frontiersin.org/articles/10.3389/fchem.2022.1041961

53. Hoffmann SA, Hos D, Küspert M, Lang RA, Lovell-Badge R, Wegner M, et al. Stem cell
factor Sox2 and its close relative Sox3 have differentiation functions in
oligodendrocytes. Development. 2014 Jan 1;141(1):39–50.

54. Dugas JC, Ibrahim A, Barres BA. The T3-induced gene KLF9 regulates oligodendrocyte
differentiation and myelin regeneration. Mol Cell Neurosci. 2012 May;50(1):45–57.

55. Yang N, Zuchero JB, Ahlenius H, Marro S, Ng YH, Vierbuchen T, et al. Generation of
oligodendroglial cells by direct lineage conversion. Nat Biotechnol. 2013
May;31(5):434–9.

56. Sock E, Wegner M. Using the lineage determinants Olig2 and Sox10 to explore
transcriptional regulation of oligodendrocyte development. Dev Neurobiol. 2021
Oct;81(7):892–901.

57. Lee Y, Messing A, Su M, Brenner M. GFAP promoter elements required for region-
specific and astrocyte-specific expression. Glia. 2008;56(5):481–93.

58. Chen YC, Ma NX, Pei ZF, Wu Z, Do-Monte FH, Keefe S, et al. A NeuroD1 AAV-Based Gene
Therapy for Functional Brain Repair after Ischemic Injury through In Vivo Astrocyte-to-
Neuron Conversion. Molecular Therapy. 2020 Jan 8;28(1):217–34.

59. Qian H, Kang X, Hu J, Zhang D, Liang Z, Meng F, et al. Reversing a model of Parkinson's
disease with in situ converted nigral neurons. Nature. 2020 Jun;582(7813):550–6.

60. Janeckova L, Knotek T, Kriska J, Hermanova Z, Kirdajova D, Kubovciak J, et al. Astrocyte-
like subpopulation of NG2 glia in the adult mouse cortex exhibits characteristics of
neural progenitor cells. Glia. 2024;72(2):245–73.

61. Kirdajova D, Valihrach L, Valny M, Kriska J, Krocianova D, Benesova S, et al. Transient
astrocyte-like NG2 glia subpopulation emerges solely following permanent brain
ischemia. Glia. 2021 Nov;69(11):2658–81.

62. Rusnakova V, Honsa P, Dzamba D, Ståhlberg A, Kubista M, Anderova M. Heterogeneity
of Astrocytes: From Development to Injury – Single Cell Gene Expression. PLOS ONE.
2013 Aug 5;8(8):e69734.

63. Barres BA, Jacobson MD, Schmid R, Sendtner M, Raff MC. Does oligodendrocyte
survival depend on axons? Current Biology. 1993 Aug;3(8):489–97.

64. Macklin W b., Weill C l., Deininger P l. Expression of myelin proteolipid and basic
protein mRNAS in cultured cells. Journal of Neuroscience Research. 1986;16(1):203–
17.

65. Rosenberg SS, Kelland EE, Tokar E, De La Torre AR, Chan JR. The geometric and spatial
constraints of the microenvironment induce oligodendrocyte differentiation. Proc Natl
Acad Sci U S A. 2008 Sep 23;105(38):14662–7.

66. Lee S, Leach MK, Redmond SA, Chong SYC, Mellon SH, Tuck SJ, et al. A culture system
to study oligodendrocyte myelination-processes using engineered nanofibers. Nat
Methods. 2012 Sep;9(9):917–22.

67. Bush TG, Puvanachandra N, Horner CH, Polito A, Ostenfeld T, Svendsen CN, et al.
Leukocyte infiltration, neuronal degeneration, and neurite outgrowth after ablation of
scar-forming, reactive astrocytes in adult transgenic mice. Neuron. 1999 Jun;23(2):297–
308.

68. Davis N, Mota BC, Stead L, Palmer EOC, Lombardero L, Rodríguez-Puertas R, et al.
Pharmacological ablation of astrocytes reduces Aβ degradation and synaptic
connectivity in an ex vivo model of Alzheimer's disease. J Neuroinflammation. 2021 Mar
17;18:73.

69. Gascón S, Masserdotti G, Russo GL, Götz M. Direct Neuronal Reprogramming:
Achievements, Hurdles, and New Roads to Success. Cell Stem Cell. 2017 Jul
6;21(1):18–34.

780
781
782
783 **SUPPLEMENTARY INFORMATION**
784
Supplemental Figure 1. Morphological characterization of astrocytes and
oligodendrocyte progenitor cells. Quantification of the **(A)** cell length, **(B)** nucleus size, **(C)**
number of branches, and **(D)** branch thickness in Aldh1l1+tdTomato+ and PDGFRα+ OPCs
(n= 70 per cell, data are presented as mean ± SEM, each data point represents one individual
cell , significance was calculated using Welch's t-test (A), Mann Whitney test (B,C) and
Kruskal Wallis one-way ANOVA (D), ** = p<0.01, **** = p<0.0001)
791
792 **Supplemental Figure 2. 14DPT scRNA-seq processing for initial UMAP generation.**
793 **(A)** Distribution of genes (nFeature), reads (nCount) and percent of mitochondrial reads
794 (percent.mt) per sample. **(B)** UMAP clustering prior to removal of microglia cluster. **(C)**Top
795 upregulated genes characterizing each cluster observed in (B).
796
797 **Supplemental Figure 3. Characterization of DLR cultures. (A)** Quantification of
798 contaminating iOLCs in post-natal astrocyte cultures 3DPT. **(B)** Quantification of MBP
799 expression in tdTomato$^+$ transduced cells 22DPT (n=3, data are presented as mean ± SEM,
800 each data point represents one individual cell culture, a matched pairs one way ANOVA was
801 performed (ns)).
802
Supplemental Figure 4. Comparison of DLR clusters 14DPT to established datasets of
OLCs. (A) Violin plots of cell types annotated by Marques *et al*. 2016 within each cluster
(OPC = oligodendrocyte progenitor cells, COPs = committed oligodendrocyte progenitors,
NFOL = newly formed oligodendrocyte, MFOL = myelin forming oligodendrocyte, mOL =
mature oligodendrocyte).**(B)** Violin plots of cell types annotated by Dugas *et al*. 2006 within
each cluster (OL = oligodendrocyte, OPC = oligodendrocyte progenitor cells).
809

Supplemental Figure 5. Characterization of *Aldh1l1^{neg}* and zsGreen^{neg} iOLCs
(A) Quantification of (i) tdTomato$^+$ and (ii) zsGreen$^+$ PDGFRα$^+$iOPCs at 12DPT (n= 3). Data are
presented as mean ± SEM, each data point represents one individual cell culture, a paired t-
test for (i) and a Wilcoxon test for (ii) was
performed. **(B)** Quantification of (i) tdTomato$^+$ and (ii) zsGreen$^+$ O4$^+$iCOPs at 14DPT (n = 4).
Data are presented as mean ± SEM, each data point represents one individual cell culture, a
paired t-test was used. **(C)** Quantification of (i) tdTomato$^+$ (ii) zsGreen$^+$ MBP$^+$ OLs at 12DPT
(n= 3). Data are presented as mean ± SEM, each data point represents one individual cell
culture, a matched pairs one way ANOVA with Geisser –Greenhouse correction
and Dunnet's post testing was used (* = p<0.05). **(D)** Representative tdTomato^{neg}zsGreen$^+$
cell with astrocyte-like morphology at onset (7DPT) and OLC expression at the end (12DPT)
of live cell tracking (arrow indicated tracked cell, scale bar = 50um).
822
823 **Supplemental Figure 6. DLR timecourse scRNA-seq processing for initial UMAP**
824 **generation. (A)** Distribution of genes (nFeature), reads (nCount) and percent of
825 mitochondrial reads (percent.mt) per sample. **(B)** UMAP clustering prior to removal of
826 microglia and VLMC clusters. **(C)** Top upregulated genes characterizing each cluster
827 observed in (B).
828
829 **Supplemental Figure 7. Differential gene expression of clusters in the Monocle3**
830 **trajectory. (A)** Genes enriched in leaf 14 of Monocle3 trajectory. Cells in UMAP colored
831 according to log-normalized gene expression values. **(B)** Genes enriched in leaf 11 of
832 Monocle3 trajectory. Cells in UMAP colored according to log-normalized gene expression
833 values.
834
Video S1. Live cell conversion of tdTomato$^+$zsGreen$^+$ astrocyte to iOLC
Video S2. Live cell conversion of tdTomato^{neg}zsGreen$^+$ astrocyte to iOLC
Table S1. Astrocyte and oligodendrocyte progenitor morphology characterization
Table S2. 14DPT scRNA seq cell collection metrics
Table S3. Pre-transduction, 3 and 8DPT scRNA seq cell collection metrics
Table S4. Top 40 genes associated with Fatecode perturbation on node 16 of the latent
layer
Data S1. BD Rhapsody Whole Transcriptome Analysis whitelist
Data S2. Monocle3-derived differentially expressed genes
844
845