TITLE PAGE

Title: Abscisic acid and GIGANTEA signalling converge to regulate the recruitment of

CONSTANS to the *FT* promoter and activate floral transition

Running title: ABA and GIGANTEA control CO recruitment to the *FT* promoter

Alice Robustelli Test[1]*, Giorgio Perrella[1]*, Beatrice Landoni[1], Sara Colanero[1], Aldo Sutti[1],

Paolo Korwin Krukowski[1], Tianyuan Xu[2], Elisa Vellutini[2], Giulia Castorina[3], Massimo

Galbiati[4], Damiano Martignago[1], Eirini Kaiserli[2], Chiara Tonelli[1], and Lucio Conti[1]

Affiliations

1 = Dipartimento di Bioscienze, Università degli Studi di Milano, Italy

2 = School of Molecular Biosciences, College, of Medical, Veterinary and Life Sciences,

University of Glasgow, UK

3 = Dipartimento di Scienze agrarie e ambientali, Università degli Studi di Milano, Italy

4 = Istituto di Biologia e Biotecnologia Agraria – IBBA, CNR – Milano, Italy

* = equal contribution

Corresponding author: L. Conti; Dipartimento di Bioscienze, Università degli studi di

Milano/ Via Celoria 26 /Milano / 20133/ Italy/ 0039 0250315012/ Lucio.Conti@unimi.it

Highlight:

22 ABA and GIGANTEA signalling promote *FLOWERING LOCUS T* (*FT*) transcriptional

23 activation by regulating the binding of the transcription factor CONSTANS to the proximal

24 *FT* promoter.

Abstract

Plants align flowering with optimal seasonal conditions to increase reproductive success. This process depends on modulating signalling pathways that respond to diverse environmental and hormonal inputs, thereby regulating the transition to flowering at the shoot apical meristem. In *Arabidopsis*, long-day photoperiods (LDs) stimulate the transcription of *FLOWERING LOCUS T* (*FT*), encoding the main florigenic signal. *FT* activation is mediated by the transcriptional regulator CONSTANS (CO), which binds to the CO responsive elements (COREs) located in the proximal *FT* promoter region. The phytohormone abscisic acid also (ABA) contributes to *FT* activation together with GIGANTEA (GI) to regulate drought escape (DE). Whether CO is a target of ABA and GI actions for the regulation of *FT* is, however, unknown. Here, we report that ABA and its signalling components promote CO recruitment to the CORE1/2, without causing reductions in the diel pattern of CO protein accumulation. ChIPseq analyses show that ABA broadly shapes the CO DNA binding landscape, which is enriched at the promoters of genes involved in the response to abiotic stress. We also found that GI promotes CO recruitment to the CORE1/2 region, and that CO recruitment is required for the accumulation of RNAPol II at the transcription start site of *FT*. Finally, we show that GI and ABA signalling pathways are largely epistatic in the control of flowering time, suggesting their involvement in the same molecular process. Taken together, these observations suggest that varying water deficit conditions modulate CO recruitment and *FT* expression, thus dictating DE strategies in *Arabidopsis*.

Keywords: flowering time regulation, environmental and stress responses, florigen signalling, regulation of gene expression

INTRODUCTION

During floral transition plants shift from vegetative to reproductive growth. This complex process is orchestrated by a combination of internal (e.g., age) and external (environmental) signals that ultimately cause transcriptional reprogramming and changes of cell fates in young primordia located at the shoot apical meristem. At the molecular level, these signals are relayed through a multitude of molecular pathways, hormonal signalling cascades and transcriptional events that unfold at different temporal and tissue scales. Variations in daylengths (photoperiod) play a key role in flowering time regulation and its measurement occurs in the vascular tissue of leaves (GARNER And ALLARD, 1920; Wigge et al., 2005; Song et al., 2015; Gendron and Staiger, 2023). Following perception of an inductive photoperiod condition, a specialised protein signal referred to as florigen is produced and exported systemically to activate floral transition at the shoot apical meristem (Kardailsky et al., 1999; Kobayashi et al., 1999). In *Arabidopsis* the photoperiodic cascade is activated under long day conditions (LDs) and requires the activity of three key genes including *GIGANTEA* (*GI*), *CONSTANS* (*CO*) and *FLOWERING LOCUS T* (*FT*) (Song et al., 2015). GI is a key activator of the network and encodes a plant-specific protein with unclear molecular functions. GI forms an oscillatory (diel regulated) complex with blue light receptors to set the stage for the transcriptional activation of *CO* during late afternoon in LDs (Suárez-López et al., 2001; Imaizumi et al., 2005; Sawa et al., 2007; Fornara et al., 2009; Ito et al., 2012). Light-stabilized CO protein promotes the transcriptional activation of *FT*, encoding the main florigen of *Arabidopsis*, so that peak *FT* transcript accumulation occurs at dusk under standard growth conditions (Kardailsky et al., 1999; Kobayashi et al., 1999; Valverde et al., 2004; Corbesier et al., 2007; Song et al., 2018).

The molecular mechanisms that lead to the stabilization of CO, its post-translational modifications, binding to specific DNA cis-elements, and interactions with a diverse array of

co-regulators are collectively important to finely tune the expression of *FT* (Takagi *et al.,*

2023). CO binds to *FT* regulatory regions at the CO Response Elements (*CORE*s) and *P1/2*

sites. These are 4-bp TGTG core motifs located in the proximal promoter of *FT* (-285bp from

ATG), broadly corresponding to Block A regulatory region (Adrian *et al.,* 2010; Tiwari *et al.,*

2010; Hayama *et al.,* 2017; Lv *et al.,* 2021). CO binding to DNA occurs through the positively

charged CCT domain, located at the C terminus, which is evolutionarily related to the DNA

binding domain observed in NUCLEAR FACTOR-Y subunit A (NF–YA) encoding genes

(Wenkel *et al.,* 2006). CO forms a trimeric complex with NF–YB and NF–YC subunits to bind

to DNA (Ben-Naim *et al.,* 2006; Tiwari *et al.,* 2010; Gnesutta *et al.,* 2017). At the same time,

through the N terminal B-BOX domain CO can form multimers, each one being able to form

NF–YB/YC complexes via the CTT for high affinity binding to the TGTG core motifs (Lv *et*

al., 2021; Zeng *et al.,* 2022). Besides NF-YB/C, other mechanisms regulate CO recruitment

to chromatin (Bu *et al.,* 2014; Wang *et al.,* 2016). All in all, these factors together with CO

participate in forming local chromatin looping and initiate *FT* transcription (Adrian *et al.,* 2010;

Tiwari *et al.,* 2010; Farrona *et al.,* 2011; Cao *et al.,* 2014; Tian *et al.,* 2021).

Several environmental and hormonal pathways influence CO function and, ultimately *FT*

transcriptional activation (Pin and Nilsson, 2012; Conti, 2017). The phytohormone abscisic

acid (ABA) is emerging regulator of flowering time, influencing the expression multiple floral

targets (Martignago *et al.,* 2020). The ABA biosynthesis pathway has been well characterised

(Seo and Marion-Poll, 2019) with key steps occurring in the plastids catalysed by the

zeaxanthin epoxidase (ABA1) and the nine-cis-epoxycarotenoid dioxygenases (NCEDs)

and the cytosol, operated by ABA2 and other enzymes. The ABA signalling cascade

comprises a family of soluble receptors known as PYR/PYL/RCARs (Cutler *et al.,* 2010) that

can bind to ABA. Upon binding the receptor can interact with a clade of protein

phosphatases, the PP2C (Fujii *et al.,* 2009). This interaction inhibits the phosphatase activity

of PP2Cs, allowing their substrate, protein kinases of the SNF1-related protein kinase 2 (SnRK2) group, to auto phosphorylate. Active SnRK2s can in turn phosphorylate downstream components of ABA signalling including the ABA-regulated bZIPs (ABRE binding factors, ABFs) which mediate transcriptional reprogramming (Furihata *et al.,* 2006; YOSHIDA *et al.,* 2015).

Much of our knowledge about the impact of ABA in flowering time derives from the study of the drought escape response (DE), an acceleration of flowering triggered by water deficit. Under these conditions, water deficit trigger the expression of *FT* and the flowering time regulator *SUPPRESSOR OF OVEREXPRESSION OF CONSTANS 1* (*SOC1*) in leaves, in an ABA-dependent manner and likely through different mechanisms (Riboni *et al.,* 2013; Hwang *et al.,* 2019). The ABFs in association with NF-YC subunits activate *SOC1* (Hwang *et al.,* 2019), whereas little is known about the regulatory role of ABA on CO upstream of *FT*. Recently, the SNF1-related protein kinase 2 (SnRK2) substrate 1 (SNS1) was shown to promote DE by inducing *FT*. SNS1 activity is regulated by phosphorylation and subsequent stabilization by ABA stimulated SnRK2 protein kinases (Katagiri *et al.,* 2024). Impairing ABA signalling causes a reduction in *FT* expression with relatively weaker changes in *CO* transcript accumulation suggesting that ABA signalling may act at least in part downstream of *CO* transcriptional activation (Riboni *et al.,* 2016). In this process, GI function is key for the ABA - dependent activation of *FT* and DE (Riboni *et al.,* 2016). GI might act as a general regulator of ABA signals for its involvement in modulation of physiological traits related to gas exchange (which are also ABA regulated) as well as ABA production (Ando *et al.,* 2013; Baek *et al.,* 2020; Simon *et al.,* 2020). However, *gi* plants also display a general hyper activation of ABA signalling responses, suggesting that the mode of action of GI in DE regulation may be separate (and more general) from its role in regulating ABA signals (Siemiatkowska *et al.,* 2022).

In this study tests were made to understand how changes in ABA accumulation could alter

FT expression through CO. We reveal that CO recruitment to the *FT* promoter is ABA

regulated which impacts the recruitment of the RNAPol II. Surprisingly GI is also required

for proper CO recruitment to the *FT* promoter, indicating that GI and ABA may operate in

the same genetic pathway. Overall, we hypothesise that GI could be an important factor

bridging photoperiodic stimuli with ABA signalling to relay water deficit information and

regulate flowering time.

MATERIAL AND METHODS

Plant material, genotyping, and growing conditions

Arabidopsis thaliana plants used in this study are of ecotype Columbia (Col-0) or Landsberg

erecta (L.*er*). Mutant and transgenic lines of *aba1-6* (Niyogi *et al.,* 1998)*, aba2-1* (Leon-

Kloosterziel *et al.,* 1996)*, nced3/5* (Frey *et al.,* 2012)*, abi1-1* (Bertauche *et al.,* 1996)*,

SUC2::HA:CO* (Jang *et al.,* 2009)*, co-10* (Laubinger *et al.,* 2006)*, gi-2* (Rédei, 1962)*, gi-100*

(Huq *et al.,* 2000) were previously described. Seeds were stratified and plants grown under

long day conditions (LDs, 16 h light / 8 h dark), under controlled-environment cabinets, as

previously described (Riboni *et al.,* 2016). Temperature range in the growth chamber

fluctuated between 24 °C during the day and 21 °C in the night and air humidity was about

60%. For *in vitro Arabidopsis* growth Murashige and Skoog (MS) medium was prepared by

dissolving the MS salt mix (Duchefa) in distilled water. After adjusting the pH solution to 5.8,

agar (Duchefa) was added to a final concentration of 0.8% w/v. Sterilized seeds (70% v/v

ethanol and 1% w/v Sodium Dodecyl Sulphate, SDS for 10 minutes) were spread onto

solidified agar plates, stratified for 2-3 days (4°C and dark) and then moved to the growth

chamber. ABA application experiments on soil were performed following the procedure

detailed previously (Riboni *et al.,* 2016). A mock solution (0.025% v/v ethanol) was used as

a mock control. Mutant combinations were selected via PCR of genomic DNA with primers

described previously (Kaiserli *et al.,* 2015; Riboni *et al.,* 2016) or in Supplementary Table 1

and by exploiting markers encoded by the different transgenes; *gi-100, SUC2::CO:CITRINE*

(Basta resistance), *SUC2::HA:CO* (GFP expression in seeds).

Molecular cloning

Cloning was done with Gateway and Multi-Site Gateway (three-fragment vector) cloning

technology (Invitrogen). To generate entry clones (pDONR207, Invitrogen), the *AttB1/AttB2-*

containing sequences were fused to the *CONSTANS* specific primers (Supplementary Table

2). The Phusion High Fidelity DNA polymerase (New England Biolabs) was used for all the

PCR reactions. The 5' and 3' elements and destination vectors were previously described;

SUC2 promoter / pDONR221 P4-P1r (Marquès-Bueno *et al.,* 2016), *mCITRINE / pDONR221*

P2r-P3 (Jaillais *et al.,* 2011) and *pB7m34GW* (Karimi *et al.,* 2005). All the recombinant

destination vectors were transformed into *Agrobacterium* cells, strain GV3101 (Koncz and

Schell, 1986), for *Arabidopsis* stable transformation.

The reporter plasmid used for monitoring luciferase (Luc) activity carries a 5' *FT* regulatory

region (1666 bp upstream of ATG). The sequence was custom synthetized (Officinae Bio,

Italy) and cloned in the reporter plasmid *pGreenII 0800-LUC* (Hellens *et al.,* 2005). This

plasmid also contains the *35S:Renilla* (Ren) cassette. The effector construct (*35S:CO*) was

built by recombining a CO entry vector (which includes the stop codon) into the Gateway

destination vector *p2GW7* (Karimi *et al.,* 2002).

Plants transformation and BASTA selection

Destination vectors *SUC2::CO:CITRINE* and *pSUC2::GW* (here used as a control for Basta

resistance) (An *et al.,* 2004), were introduced into *Agrobacterium* by electroporation.

Transformed *Agrobacterium* were used to generate *Arabidopsis* transgenic plants via the

floral dip technique (Clough and Bent, 1998). Transgenic plants were selected on soil after

continuous Basta applications (12 mg/l dilution, Agrevo) and single locus insertion events

were selected based on a Mendellian 3:1 ratio in T_2 generation. T_3 homozygous lines were

selected on soil or MS+Basta plates, according to the absence of Basta resistance

segregation. Transgenic *SUC2* plants propagated up to the 5th generation by selfing did not

show obvious signs of silencing.

Protoplast Isolation and transactivation Assays

184   Leaf protoplasts were isolated from 3-week-old Arabidopsis plants as previously described

185   (Iacopino *et al.*, 2019). Luc and Ren activities were measured by providing the specific

186   substrates (Dual-Luciferase reporter assay system, Promega) and luminescence was

187   measured with the EnSight plate reader (Perkin Elmer). In each assay, data were expressed

188   as Luc activity relative to Ren.

Flowering time measurement

Flowering time was measured by scoring the number of rosette leaves, excluding

cotyledons. When indicated, flowering was expressed as days to bolting (days between

sowing and a visible emergence of the primary inflorescence). Data regarding cauline leaf

number (I1 phase) were also recorded for each genotype. To avoid flowering time alterations

associated to soil transfer from plates, Basta resistance selection in T1 generation was

performed directly on soil.

RNA extraction and Real-Time qPCR

Total RNA was extracted with QIAzol reagent (Invitrogen) and suspended in RNase-free

milliQ dH_2O. RNA concentration was measured with a UV spectrophotometer and 1000 ng

aliquots were used for cDNA synthesis with the Maxima Reverse Transcriptase kit

(Thermofisher). Quantitative Real-Time PCR (qPCR) was performed as previously detailed

including primers to amplify *CO*, *FT*, *ACT2* and *IPP2* were described (Kaiserli *et al.,* 2015;

Riboni *et al.,* 2016). Primers to amplify *CITRINE* are listed in Supplementary Table 1.

Nuclei isolation and CO protein detection in *Arabidopsis*

Approximately 100 mg of *Arabidopsis* seedlings were harvested and immediately frozen in liquid nitrogen. Tissue samples were ground by shaking with glass beads in a TissueLyser II (Qiagen, 2 pulses of 30 s each at 28Hz shaking). Leaf powder was suspended in 1.2 ml of cold nuclear isolation buffer (20 mM Tris-HCl, pH 8.8, 25 mM NaCl, 5 mM MgCl$_2$, 30% (v/v) glycerol, 5% (w/v) sucrose, 0.5% (v/v) Triton X-100, 0.08% (v/v) β-mercaptoethanol, 0.2% (v/v) SIGMA plant protease inhibitor, 1mM DTT, 1.3 mM PMSF) (Hayama *et al.,* 2017). The samples were filtrated trough two layers of Miracloth (Millipore) and centrifuged at *5,000 x g*, at 4 °C, for 10 minutes. The pellet was washed four times with 1 ml of nuclear isolation buffer and, after each wash, pelleted at 4 °C at decreasing speed: *5,000 x g, 2,700 x g, 2,200 x g* and *2,200 x g*, 8 minutes each time. Nuclei were suspended in 30 µl of nuclear isolation buffer, mixed with 10 µl of Laemmli Buffer and heated 10 minutes at 95 °C. Samples were centrifuged for 1 minute at *3,000 x g* to pellet all nuclear membranes, and 20 µl of the supernatants (enriched in soluble nuclear proteins) were loaded onto an 10% SDS-PAGE gel. For total protein detection, tissue from 50 mg seedlings was harvested in liquid Nitrogen and lysed in pre-cooled TissueLyser II (Qiagen) for 3 min at the frequency 30/s. Proteins were isolated by homogenisation in 450 µL sample buffer (100 mM Tris-HCl pH 6.8, 20% (v/v) glycerol, 5% (w/v) SDS, 20 mM DTT, 1mM bromophenol blue, 1× proteinase inhibitor cocktail (Sigma) and 80 µM MG132), boiled at 95 °C for 5 min, centrifugated for 1 min at *12000* x g and loaded onto an 4%-12% Bis-Tris Plus SDS Mini Protein Gel run in 1× Bolt MOPS SDS buffer (Thermo Fisher). Proteins were transferred to a nitrocellulose membrane and incubated with anti-HA-Peroxidase, High Affinity antibody (Roche), anti-Histone H3 (anti H3) or anti-Actin (Agrisera), used for loading control. Chemiluminescent signals were detected through a ChemiDoc Touch Imaging System (Biorad) and quantified with the Image Lab software (Biorad).

ChIP experiments and sequencing

Chromatin immunoprecipitation assays were performed as described previously with minor modification, starting from 1-2 grams (FW) of seedlings (Perrella *et al.,* 2018*,* 2023). A bioruptor sonicator (Diagenode, B01020001) was used to shear the chromatin using the following settings: 40 Cycles, 30 sec ON, 30 sec OFF at high power. With these conditions the size of sheared DNA is below 100 bp. An anti-RNAPol II antibody (AbCam Ab5408) or GFP-Trap magnetic nanobodies (Chromotek) were used. qPCR was performed at the following cycles: 95°C x 3 min; 95°C x 10 sec, 59.5°C x 30 sec (45 cycles); 65°C x 0.05 sec and 95°C x 30 sec (Melting curve). Relative enrichment was calculated as described previously (Kaiserli *et al.,* 2015; Perrella *et al.,* 2018). Reactions were performed on four technical replicates and two independent biological experiments. Amplicons of the *CBS*, *CORE1/2*, *CORE3*, *CORE4* and *CORE5* regions of *FT* were obtained with primers previously described (Hayama *et al.,* 2017), whereas the *TSS* regions of *FT*, *YUCCA8* and *ACTIN2* were amplified with primers lcm217/lcm218, *YUC8 FW*/*RV* and *ACT2 FW*/*RV*, respectively (Supplementary Table 1).

For ChIP-sequencing 5 grams (FW) of seedlings were used for each genotype/replicate and processed as described above. Inputs and ChIP samples (two biological replicates) were quantified with Qubit dsDNA Quantification High sensitivity Assay Kit (Invitrogen). For library preparation, the VAHTS Universal Plus DNA Library Prep Kit for Illumina (ND617-02, Vazyme, Nanjing, China) was used, following the manufacturer's recommendations without fragmentation. Library quality control was performed using the QSep-400 (BiOptic, Taipei, China) and Qubit 3.0 (Thermo Fisher Scientific, Wilmington, USA). The qualified library was sequenced on the Illumina Novaseq X platform (Illumina, San Diego, USA) with paired-end 150 bp (PE150) mode. Library construction and sequencing were performed at Biomarker Technologies (BMKGENE) GmbH. Mapping of raw reads and peak calling followed the greenscreen pipeline (Klasfeld *et al.,* 2022) with slight modifications. The adapted code is

available at https://github.com/beaLando/chipseq_aba.git. In brief, fastp (Chen *et al.,* 2018)

was used to trim and filter reads after quality checks in fastqc (Andrews S., 2010). Mapping

of filtered reads was then executed with Bowtie2 with TAIR10.1 as reference genome. Peak

calling was done in macs2 (Gaspar, 2018) for each replicate separately using input samples

as background. Peaks found for each replicate were then processed with irreproducible

discovery rate (IDR) to retain reproducible peaks and with a global IDR (FDR corrected)

greater or equal to -log10(0.1). This list was then further filtered to retain only peaks falling

outside greenscreen's blacklisted regions. Further quality checks were executed with

qualimap (García-Alcalde *et al.,* 2012) and deeptools (Ramírez *et al.,* 2014). Finally,

ChIPseeker (Yu *et al.,* 2015) was used to estimate how frequently peaks would fall within

gene promoters or other genomic regions.

Because the two replicates of *SUC2::CO:CIT aba1-6* resulted in very low mapping rates

(between 3-15%; Table S1), we simulated a peak calling analysis on *SUC2::CO:CIT*, by

subsampling mapped reads until 3-15% mapping rate was achieved. The logic for

subsampling *SUC2::CO:CIT* was the following: if a sizeable number of peaks could still be

called for *SUC2::CO:CIT* at low mapping rates, the low number of called peaks in the

SUC2::CO:CIT aba1-6 mutant could be due to a real biological effects leading to low binding.

The ChIPseq data from this study is available under the GEO accession number

GSE291256.

Metanalysis studies of RNA-seq and ChIP-seq datasets

We retrieved CO ChIPseq peaks data (de los Reyes *et al.,* 2024) - GEO accession number

GSE222657 - and gene expression data of upregulated DEGs in *SUC2::CO* and *35S::CO*

genotypes relative to the WT (same study). Using ChIPseeker, CO peaks were then

annotated with gene IDs based on the genes' distance from the peak. Using GeneOverlap

(Shen, 2024) we assessed the overlap in Click or tap here to enter text.peaks, or genes in

proximity of peaks, and their significance, for the two datasets - our study and (de los Reyes

et al., 2024) using the number of protein-coding genes in *A. thaliana* as background. We

then obtained the intersection between genes falling next to *SUC2::CO:CIT* peaks and

upregulated *SUC2::CO* DEGs compared to WT. Based on this intersection, we ran an

enrichment analysis for biological process terms using the union of all *SUC2::CO*

upregulated DEGs and genes falling next to *SUC2::CO:CIT* peaks as background. The

same procedure was followed for *35S::CO* peaks and DEGs. All plots were produced with

ggplot2 (Wickham, 2011) and Inkscape.

Statistical analysis

Analysis of variance (ANOVA) was conducted with linear models where flowering time/gene

expression was predicted by genotype according to the formula: flowering time (gene

expression) ~ genotype. Here, "flowering time" was expressed as either one of: leaf number

at bolting, days to bolting, or number of I1 nodes. "genotype" refers to the allelic status at

ABA1/2 (WT, *aba1-6*, *aba2-1*) or *GI* (WT, *gi-2*, *gi-100*) loci, and different isogenic lines for

the various mutants. When *aba1* mutants were in different backgrounds (WT,

SUC2::CO:CIT, *SUC2::HA:CO*), the same model was run separately for the different

backgrounds and the models were qualitatively compared. In Fig. 2A, the model used was

gene expression ~ genotype*timepoint. If the interaction was not statistically significant, only

the additive model (genotype + timepoint) was considered for post-hoc tests.

Before running the ANOVAs, the condition of homogeneity of variances between groups

was checked with a Bartlett's test in base R (R Core Team 2023). If group variances were

not homogeneous, a log transformation was applied to the data. All ANOVA models were

conducted with the R function lm (R Core Team 2023) using sum contrasts and significance

of predictors was assessed with the ANOVA function from the car package (Fox and

Weisberg, 2018). Post-hoc Tukey tests to assess differences between each pair of

genotypes were conducted with the R packages multcomp (Hothorn *et al.,* 2008) and

emmeans (Lenth, 2023). Even for models including timepoint as predictor, post-hoc tests

were conducted exclusively to assess differences between genotypes within timepoints.

RESULTS

CO function is sensitive to ABA signalling status in the phloem companion cells

To evaluate the contribution of ABA accumulation in regulating CO function and *FT* expression we generated transgenic plants constitutively expressing *CO* in different ABA-impaired/defective backgrounds. Since CO regulates *FT* in the phloem companion cells (An *et al.,* 2004) we engineered a *CO:CITRINE* (*CO:CIT*) translational fusion gene under the control of the phloem companion cell-specific *SUCROSE-PROTON SYMPORTER 2* (*SUC2*) promoter (Truernit and Sauer, 1995). Basta resistant T1 plants of wild-type (Col-0) and two different ABA-deficient backgrounds (*aba1-6* and *aba2-1*, impaired at different steps of the ABA biosynthetic pathway) (Nambara and Marion-Poll, 2005) displayed a clear early flowering phenotype compared with the corresponding genotypes transformed with an empty vector (Fig. 1A). *SUC2::CO:CIT aba1-6* and *aba2-1* plants showed a mild but significant delay in flowering (p = 6.1E-07 and 1.2E-14, respectively) compared with the wild type transformed with *SUC2::CO:CIT*. Ruling out an indirect effect of reduced ABA production on *SUC2* promoter activity, we detected similar levels of *CIT* transcript accumulation across randomly chosen T1 plants (Supplementary Figure S1). This suggests a role for ABA production downstream of *CO* transcriptional activation. In support of this view, we observed similarly reduced level of *FT* accumulation in *SUC2::CO:CIT* in either *aba1-6 or aba2-1* independent T1 transgenic events compared with *SUC2::CO:CIT* transgenics events in the wild-type background (p = 0.047 and 0.026, respectively, Supplementary Figure S1). *FT* transcript levels detected in *SUC2::CO:CIT aba2-1/aba1-6* plants were, however, slightly increased compared with the wild type transformed with an empty vector, consistent with their earlier flowering.

To confirm the observed flowering time phenotypes we crossed two homozygous T3 *SUC2::CO:CIT* plants with *aba1-6* to generate isogenic lines (#5 and #18). We also performed genetic crosses between *SUC2::HA:CO* plants (Jang *et al.,* 2009) and *aba1-6* to

obtain *SUC2::HA:CO aba1-6*. *SUC2::HA:CO* plants were extremely early flowering, similar

to *SUC2::CO:CIT* plants. However, in the *aba1-6* background, the early flowering phenotype

conferred by *SUC2::HA:CO* or *SUC2::CO:CIT* was significantly alleviated ($p < 4.6E-06$, Fig.

1B and 1C). Collectively, our findings suggest that ABA may promote CO function and *FT*

transcript accumulation to activate flowering.

Reduced ABA levels impair *FT* transcript accumulation

We next aimed to evaluate potential modes of regulations of CO function controlled by ABA.

Using stable transgenic isogenic lines expressing *SUC2::HA:CO* or *SUC2::CO:CIT #5* in the

wild-type or *aba1-6* backgrounds, we confirmed that *FT* transcript accumulation was

reduced in *aba1-6* plants compared with the corresponding ABA producing WT background

($p < 3.33E-03$, Fig. 2A) irrespective of the time point analysed (ZT12 and 16 under LDs). In

contrast, *CO* levels remained not affected in *SUC2::HA:CO aba1-6* plants or even increased

($p = 1.82E-03$) in *SUC2::CO:CIT#5 aba1-6* compared with the corresponding *SUC2::CO:CIT*

#5. Interestingly, we also detected reduced levels of *FT* transcript accumulation in *aba1-6*

plants compared with the wild type ($p = 1.92E-04$) at ZT16 when we also observed a

marginal reduction ($p = 0.06$) in *CO* transcript levels. To investigate further the role of ABA

production in *FT* transcript accumulation we analysed 9-day-old seedlings of wild type and

strong ABA deficient mutant backgrounds, including *aba1-6*, *aba2-1* and double mutants of

9-cis-epoxycarotenoid dioxygenase (*nced*) *3* and *5* (Frey *et al.,* 2012). Compared with the

wild type, at ZT16 we observed reduced *FT* accumulation in all ABA deficient mutants ($p <$

$6.09E-06$) and correspondingly reduced levels of *CO* transcript ($p < 0.01$) (Fig. 2B).

Collectively, our data indicate a role for ABA in promoting *FT* accumulation with ABA

affecting both *CO* expression and its function.

To gain further insights into the potential mode of ABA-dependent stimulation of CO we

analysed HA:CO protein levels derived from *SUC2::HA:CO* and *SUC2::HA:CO aba1-6*

plants. HA:CO protein accumulation was broadly similar between the two genotypes at ZT12

and ZT16, suggesting that reduced ABA accumulation impairs HA:CO protein function but

not HA:CO accumulation (Fig. 2C). We next extended this analysis to morning timepoints

including ZT 0.5 (i.e., within 30 min after the switch on of the lights), ZT4 till midday-dusk

(ZT8 and ZT12). At ZT0.5 we observed a much stronger increase in HA:CO accumulation

in *SUC2::HA:CO aba1-6* plants compared to *SUC2::HA:CO* (Supplementary Figure S2).

This peak of accumulation was, however, transient and might suggest a role of ABA in

repressing this phase of CO stabilization in this early morning timepoint (Valverde *et al.,*

2004; Song *et al.,* 2012; Hayama *et al.,* 2017). HA:CO protein was still clearly detectable in all

the other time points in *SUC2::HA:CO aba1-6* plants further pointing to defective CO function

with respect to *FT* and consequent flowering activation.

To independently test for potential effects of ABA on HA:CO protein abundance we

exogenously applied three different concentrations of ABA (0.25 -2.5 -25 μM) on a daily

basis for 12 days (Riboni *et al.,* 2016). Tissue from ABA-treated plants was collected at ZT4

and ZT12 and HA:CO protein was detected from nuclear extracts. We did not observe

obvious changes in HA:CO protein levels at any of the concentrations and time points

assayed (Supplementary Figure S3). In addition, we did not observe obvious changes in the

pattern of HA:CO protein gel mobility in the ABA deficient background *aba1-6*

(Supplementary Figure S3), which are determined by its phosphorylation status (Sarid-Krebs

et al., 2015; Chen *et al.,* 2020). In summary, our data indicate that CO function might be

stimulated by ABA in the cells where *FT* is transcriptionally activated without clear changes

at the post translational level.

ABA stimulates the recruiting of CO to the *FT* promoter region

We next investigated what other aspects of CO function could be influenced by ABA status.

We first set up a transient reporter expression assay in *Arabidopsis* mesophyll protoplasts,

by fusing a 1666 bp promoter region of *FT* upstream of the *LUCIFERASE*-encoding sequence (hereafter *FT~pro~::LUC*). This promoter region encompasses several cis elements that are critical for the *FT* transcriptional activation, including the well characterised CO responsive element (*CORE)1/2* (Tiwari *et al.,* 2010). Co-transformation of *FT~pro~::LUC* with *35S::CO* vectors resulted in a strong activation of LUC activity, indicating that the fused promoter can respond to CO binding (Supplementary Figure S4). Co-incubation of *FT~pro~::LUC/35S::CO* transformed protoplasts in presence of ABA (10 and 100 nM), however, did not produce significant changes in LUC activity suggesting that exogenous ABA does not directly influence CO-mediated stimulation of the *FT* promoter, at least in the absence of a native chromatin context.

We therefore evaluated CO recruitment to the *FT* promoter through a Chromatin Immunoprecipitation (ChIP) assay by comparing CO binding in isogenic transgenic seedlings of *SUC2::CO:CIT* #5 (wild type and *aba1-6* backgrounds) under LDs. Consistent with previous reports (Cao *et al.,* 2014; Song *et al.,* 2016; Hayama *et al.,* 2017; de los Reyes *et al.,* 2024), at ZT12 we observed a strong enrichment of the CO:CIT protein at the *CORE1/2 FT* promoter regions (Fig. 3 and Supplementary Figure S5). In contrast, such CO:CIT enrichment was much decreased (approximately 4X) in the *aba1-6* background, suggesting that ABA promotes CO recruitment to the *CORE1/2* region. A much weaker CO:CIT binding was also observed at the *CORE3* position (Hayama *et al.,* 2017), located in the second intron of *FT* and the *CBS* (*CAAT Binding Site*) region, located at 1.8kb upstream of the ATG start codon - Block B enhancer - (Adrian *et al.,* 2010). These enrichment signals were also reduced in in the *aba1-6* background.

To gain a better resolution and coverage of the CO:CIT DNA binding landscape we carried out a ChIP experiment followed by sequencing (ChIPseq). We sequenced the immunoprecipitated DNA derived from *SUC2::CO:CIT*#5 (wild type and *aba1-6* backgrounds) at ZT12 in two biological replicates. We initially retrieved around 20000 peaks

per replicate in the *SUC2::CO:CIT* genotype, but only 7312 were retained after

irreproducible discovery rate (IDR) filtering (Figure 4A, Table S2 and S3). These peaks were

mostly located in proximity of promoter (ca. 60%) and distal-intergenic (ca 32%) regions

(Figure 4B) which resulted in 5755 unique genes being recovered in proximity of peaks after

annotation (Table S3). The same procedure resulted in around 150-250 peaks called and 1

peak retained in the *SUC2::CO:CIT aba1-6* line, although very low mapping rates (3-15%)

were achieved in this background (Figure 4A, Table S2 and S3). We thus performed

subsampling of mapped reads in the *SUC2::CO:CIT* genotype and called peaks again at

lower mapping rates. Even at mapping rates close to 3-15% (corresponding to 949907

mapped reads per replicate), it was possible to retrieve around 1300 peaks per replicate for

SUC2::CO:CIT, that is over 1000 more peaks than in the mutant *SUC2::CO:CIT aba1-6*

(Table S2). From a qualitative perspective, the mapping density around transcription start

sites for all immunoprecipitated replicates showed no enrichment of reads in *SUC2::CO:CIT*

aba1-6, while enrichment was visible for *SUC2::CO:CIT* (Figure 4A). This suggests that,

despite low DNA yields likely affect our results for the *SUC2::CO:CIT aba1-6* mutant, the

almost total absence of reproducible peaks could be partly attributed to low binding by

CO:CIT in *aba1-6*.

The overlap of peaks detected in *SUC2::CO:CIT* and those found in *35S::CO* (de los Reyes

et al., 2024) was not statistically significant (Fisher's exact test $p > 0.05$, overlap = 252, total

= 10272), but the overlap in genes closest to peaks was ($p < 0.05$, overlap = 648, total =

7823). This might be due to differences in how peaks were called and filtered, as well as

differences in the promoters used in the two different studies. The comparison of

SUC2::CO:CIT peaks associated with corresponding upregulated DEGs found in *SUC2::CO*

(de los Reyes *et al.,* 2024) further disclosed an enrichment for biological process terms related

environmental stimuli or stress (abiotic and biotic) responses, including response to salt and

water deficit (Fig 4C). A similar over representation of these terms was independently

confirmed when comparing CO peaks derived from *35S::CO* and DEGs of *35S::CO* (de los

Reyes *et al.,* 2024) (Figure S6).

As expected, *FT* was found within the *SUC2::CO:CIT* ChIP peaks - *SUC2::CO* DEGs and

35S::CO ChIP peaks-DEGs intersections. We found overlapping CO ChIP peaks in the *FT*

CORE1/2 region in the two ChIP datasets indicating that the CORE1/2 region represents

the main CO binding site *in vivo* (Figure 4D). The ChIPseq analyses also identified potential

peaks in regions corresponding to the CBS but not the CORE3. The CBS peak was not

significant in our study, but present in *35S::CO* ChIP dataset, indicating weaker binding

whereas no binding could be confirmed around the CORE3. Notably, the CORE1/2 peak

was absent in *SUC2::CO:CIT aba1-6*, further supporting a role for ABA in promoting CO

chromatin binding.

ABA promotes RNAPol II recruitment to the *FT* promoter

To define the role of ABA in regulating the early transcriptional events at the *FT* promoter

we monitored the RNA Polymerase II (RNAPol II) recruitment. We assayed the Ser5

phosphorylated pool of RNAPol II, which in Arabidopsis is distributed over the transcribed

region of regulated genes (Obermeyer *et al.,* 2023). In the wild type at ZT12 we could detect

a peak of RNAPol II enrichment primarily around the *FT* transcription start site (TSS) (Fig.

5A and Supplementary Figure S7). In the *co* and the *gi* mutants, RNAPol II occupancy was

much reduced compared with the wild type. This data indicates that the assay is a sensitive

readout for CO-stimulated recruitment of RNAPol II to the *FT* promoter (although it does not

imply a physical contact between CO and the RNAPol II) (Fig. 5A). Remarkably, we

observed a similar strong reduction in RNAPol II occupancy at the TSS in ABA deficient

mutants *aba1-6* and *aba2-1* (Fig. 5A). Ruling out a general defect in RNAPol II recruitment

at actively transcribed genes in ABA deficient or other mutants utilised in these assays, we

observed no clear changes in RNAPol II occupancy at the TSS of *ACTIN2* and *YUCCA8*

genes in the same panel of genotypes (Supplementary Figure S8). Therefore, the reduced

occupancy of RNAPol II could depend on the lower levels of CO recruitment to the *CORE1/2*

region observed in ABA deficient mutants. ABA signals promote the transcriptional

activation of *CO* in *Arabidopsis* and other *Brassicaceae* (Yoshida *et al.,* 2010; Koops *et al.,*

2011; Zhang *et al.,* 2022) and we observed reduced *CO* transcript levels in ABA deficient

backgrounds, albeit only at the peak of *FT* accumulation (ZT16) (Fig. 2B). To verify further

the post-transcriptional control of CO mediated by ABA and its impact on RNAPol II

recruitment, we examined the same *FT* regions in *abscisic acid insensitive 1-1* (*abi1-1*)

mutants. *abi1-1* encodes a dominant allele of the ABI1 (encoding a PP2C) which severely

impairs ABA signalling (Bertauche *et al.,* 1996). *abi1-1* plants show similar levels of *CO*

transcript accumulation during the day, but diminished levels of *FT* accumulation (Riboni *et*

al., 2016). Consistent with this observation, RNAPol II occupancy was reduced in *abi1-1*

mutants compared with the wild type (Fig. 5B and Supplementary Figure S9). Interestingly,

RNAPol II recruitment in L.*er* (unlike Col-0) was reproducibly enriched primarily towards the

CORE1/2 regions. These results suggest that impairing ABA signalling may also cause

similar reductions in CO occupancy at the *FT* promoter which impacts correct RNAPol II

recruitment, despite RNAPol II positioning is subject to unknown ecotype-specific controls.

Finally, we assessed the role of ABA on RNAPol II recruitment to the *FT* promoter in the

presence of constitutive CO accumulation, thus independently of ABA-mediated

transcriptional effects on *CO*. We observed a clear decrease in RNAPol II enrichment at the

TSS of *FT* in *SUC2::HA:CO aba1-6* compared to *SUC2::HA:CO*, which correlates with the

reduced levels of CO occupancy at the *FT* promoter detected in previous ChIP assays (Fig.

5C and Supplementary Figure S7). Notably, RNAPol II was more strongly enriched in

SUC2::HA:CO plants compared with the wild type, further supporting the limiting role of CO

accumulation in RNAPol II recruitment.

ABA and GI largely act in the same pathway to control CO recruitment to the *FT* promoter and promote floral transition

Previously we reported the key role for GI in mediating flowering under water deficit conditions (Riboni *et al.,* 2016). *gi* mutants are defective in *FT* upregulation under control (Song *et al.,* 2014) or water deficit conditions (Riboni *et al.,* 2016) even in genetic backgrounds overexpressing *CO.* Consistent with these findings, we observed a strong reduction in RNAPol II recruitment to the TSS of *FT* in *SUC2::HA:CO gi-100* lines, suggesting that GI might also be important for CO recruitment (Fig. 5C and Supplementary Figure S7). A ChIP assay using *SUC2::CO:CIT gi-2* #5 further revealed reduced levels of CO:CIT recruitment to the *CORE1/2* region compared to *SUC2::CO:CIT* #5 (Fig 5D). However, flowering time of *SUC2::CO:CIT gi-2* #5 plants was similar to *SUC2::CO:CIT* #5 indicating that an excess of transgene-derived CO:CIT expression could ultimately compensate for its reduced recruitment observed at the *FT* promoter in *gi* mutants (Supplementary Figure S10).

Because both ABA and GI mediate CO recruitment to the *FT* promoter, we tested whether they acted in the same genetic pathway. We thus compared flowering time (bolting time and leaves counts) of *SUC2::HA:CO aba1-6* and *SUC2::HA:CO gi-100* with triple mutants of *SUC2::HA:CO aba1-6 gi-100* under LDs. As expected, flowering time of *SUC2::HA:CO aba1-6* was significantly delayed compared to *SUC2::HA:CO* plants both in terms of number of rosette leaves (p = .01) or bolting time (p = 1.78E-11) (Fig. 6A and Supplementary Figure S11). We did not detect a delay of flowering in any *SUC2::HA:CO gi* mutants combination (i.e., *gi-2, gi-100*) compared to *SUC2::HA:CO* in terms of leaf numbers, but we observed a significant delay in days to flowering (p = 1.95E-05 for *SUC2::HA:CO gi-2*). Finally, we compared the impact of combined *aba1-6* and *gi-100* mutant alleles in the *SUC2::HA:CO* background. This showed a minor but significant (p < 0.01) aggravation of flowering time of *SUC2::HA:CO aba1-6 gi-100* compared with *SUC2::HA:CO aba1-6* in terms of rosette leaves but no consistent effects in days to flowering (Fig. 6A and Supplementary Figure

S11), suggesting a partially epistatic relationship between GI and ABA signalling in this process. We also scored the number of I1 coflorescences (i.e., nodes presenting side shoots derived from the main inflorescence, Supplementary Figure S11). I1 nodes were similarly reduced in *SUC2::HA:CO* plants and *SUC2::HA:CO gi* backgrounds. We observed a significant increase in I1 nodes in *SUC2::HA:CO aba1-6* (p = 0.002) but no further aggravation of this phenotype in *SUC2::HA:CO aba1-6 gi-100* mutants lines. Therefore, GI and ABA signalling action is largely non additive to regulate flowering time and during inflorescence development in genetic background that overexpress *CO*.

DISCUSSION

Our study describes a yet uncharacterised role for ABA on CO function. Since ABA biosynthesis and signalling is enriched in the phloem companion cells of the leaf (Mustilli *et al.,* 2002; Kuromori *et al.,* 2014), endogenous CO function might be subject to varying ABA accumulation, which is usually related to water deficit status. Our experiments do not imply a direct role of ABA in controlling CO function as this could be mediated by one or more CO-interacting protein(s) or factors that are not present in ABA-deficient plants. However, they establish a role for ABA and its signalling components in influencing *FT* promoter accessibility to CO, particularly the *CORE1/2* region, subsequently leading to RNAPol II engagement at the *FT 5' UTR*.

Recent evidence supports the role of GI as a potential target of ABA action during DE. First, we previously highlighted the key role of GI in enhancing *FT* expression in response to ABA, and that GI function was sensitive to the ABA signalling status (Riboni *et al.,* 2016). Second, the epistatic interaction between *gi* and *aba1* mutants in the regulation of flowering time (bolting time in particular) of *SUC2::HA:CO* plants aligns with a model where ABA and GI could act in the same pathway to regulate CO recruitment to the *FT* promoter, although the regulatory role of ABA on GI function remains unknown. GI was shown to stimulate ABA

production which might suggest an indirect effect on CO recruitment exerted by GI (Baek *et*

al., 2020). However, we reported an increased levels of ABA sensitivity in *gi* plants under

controlled water deficit conditions (Siemiatkowska *et al.,* 2022), suggesting that GI might be

crucial for relaying ABA signalling upstream of *FT*. GI exerts a direct influence on *FT*

activation by associating to its promoter (Sawa and Kay, 2011). To further define the sites

of GI binding onto the *FT* regulatory region, we analysed GI ChIPseq data describing the

genome-wide occupancy of GI (Nohales *et al.,* 2019). GI binding regions on the *FT* promoter

(Fig. 6B) closely overlap with two DNAse accessible chromatin locations (Zhang *et al.,* 2012).

These regions were functionally defined as block C and E enhancers (Adrian *et al.,* 2010;

Liu *et al.,* 2014; Zicola *et al.,* 2019). Block C contains NF-Y binding sites, which facilitate

looping with the *CORE* region and consequent *FT* transcriptional activation (Cao *et al.,* 2014;

Siriwardana *et al.,* 2016). Block E sequence exhibits binding sites for different transcriptional

regulators, including the *FT* repressor SCHLAFMUTZE (SMZ), which has been shown to

bind to GI (Mathieu *et al.,* 2009; Sawa and Kay, 2011). While GI is unlikely to bind DNA and

despite its molecular function is still unclear, it is required to control transcriptional events

through direct interaction with specific transcriptional regulators (Sawa and Kay, 2011; Song

et al., 2014; Kubota *et al.,* 2017), possibly by acting as a general chaperone-like protein (Cha

et al., 2017). These observations could point to a model whereby GI controls CO recruitment

through binding to Block E and C enhancers, perhaps by affecting local chromatin

conformation or CO protein complex accessibility (Fig.6B).

Different NF-Y encoding genes act as regulators of ABA transcriptional responses

(Kumimoto *et al.,* 2013; Song *et al.,* 2016) and *nf-yb2* mutants display an exaggerated delay

in flowering time compared to wild-type controls under osmotic stress conditions (Chen *et al.,*

2007). Considering the mild flowering time defects of *nf-yb2* mutants under normal growth

conditions (Kumimoto *et al.,* 2008), this observation may point to a specific requirement for

NF-YB2 subunit in promoting flowering under osmotic stress, which is ABA regulated. A similar observation about the requirement for NF-Y in mediating ABA-dependent flowering derives from the lack of DE of *nf-yc3 yc4 yc9* mutants (Hwang *et al.,* 2019). Because NF-YB/C can form complexes with both CO and NF-YA subunits, their contribution in relaying ABA signals may depend on either CO, NF-YA or both functions. Moreover, NF-YC subunits interact with the ABFs bZIP to promote the transcriptional activation of *SOC1* (Hwang *et al.,* 2019). Whether this model of interplay between ABFs and NF-YA/B/C can be applied to the *FT* promoter regulation is unclear mainly because the ABFs do not appear to bind to the *FT* chromatin region (Song *et al.,* 2016; Hwang *et al.,* 2019). Intriguingly, recent data support the physical interaction between CO and ABF1/2/3/4 and their antagonistic roles at salinity-responsive genes promoters (Du *et al.,* 2023). Our ChIPseq analyses revealed an over-representation of genes related to biotic and abiotic responses (particularly salt and water stress response) among *bona fide* CO direct targets. This points to a broader role for CO in influencing processes unrelated to flowering time regulation in Arabidopsis and other species (Du *et al.,* 2023; Biancucci *et al.,* 2024, Preprint). It might also suggest a potential positive feedback mechanism, in which ABA mediates between abiotic stress levels and CO occupancy at regulated promoters. Previous reports have described direct links between ABA signalling and function of chromatin regulatory complexes that affect *FT* expression (Farrona *et al.,* 2004; Peirats-Llobet *et al.,* 2016; Katagiri *et al.,* 2024). Here, genetic, and molecular evidence suggests that ABA might also act through GI-mediated recruitment of CO to regulate *FT* expression. Therefore, our data and experimental approaches offer new tools for investigating the ABA-dependent level of regulation of photoperiodic flowering.

FIGURES LEGENDS

593
594
595

596
Figure 1. ABA promotes CO function in the phloem companion cells
(A) Boxplot of flowering time (right Y axis, rosette leaves number) of Basta resistant transgenic T1
plants of the indicated genotypes. Whiskers represent the range, min to max values. The left Y axis
shows the number of plants scored for each genotype, and their phenotype distribution according to
rosette leaves number (X axis). An ANOVA test to assess the impact of *aba1* and *2* mutations was
run separately for *SUC2::CO:CIT* and empty vector - transformed plants (vertical bar). For both
groups (empty vector or *SUC2::HA:CO* transformed), the genotype was a statistically significant
predictor of leaf number at bolting ($p < 0.001$). Letters at the top of boxplots indicate statistically
significant differences ($p < 0.05$) according to a Tukey post-hoc test. (B) Boxplot of flowering time of
the indicated genotypes and isogenic lines derived from the introgression of different *SUC2::CO*
transgenes (wild type Col-0) into *aba1-6*. An ANOVA test to assess the impact of *aba1-6* mutation
was run separately for transgenic (*SUC2::CO:CIT/SUC2::HA:CO*) and non-transgenic controls
(vertical bar). For both groups, the *aba1-6* genotype was a statistically significant predictor of leaf
number at bolting ($p < 0.001$). Letters at the top of boxplots indicate if genotypes showed statistically
significant differences ($p < 0.05$) according to a Tukey post-hoc test. The number of samples
analysed for each genotype is shown at the bottom of the graph (C) Representative plants of the
indicated genotypes pictured 27 days after sowing.

616
617
618
619
Figure 2. ABA regulates CO function without altering its accumulation in *Arabidopsis*
(A) Real-Time qPCR of *CO* (upper panels) and *FT* (lower panels) transcripts accumulation in
seedlings of the indicated genotypes grown on plates for 13 days and harvested at the indicated
Zeitgeber times (ZT, h) in the diurnal phase. Horizontal Lines represent the mean fold change of
IPP2-normalised *CO/FT* accumulation relative to the wild type (the three biological replicates are
shown as dots). Each replicate comprises approximately 50 mg of seedlings. Different letters denote
differences ($p < 0.05$) between genotypes within each ZT according to post-hoc tests. (B) Same as
A, from seedlings grown for 9 days under LDs and harvested at ZT16. Different letters denote
different ($p < 0.05$) groups according to post-hoc tests. In (A) and (B), the plot background colours
denote the ABA producing vs the ABA deficient genotype (C) Immunoblot detection of HA:CO protein
from nuclear fractions of seedlings grown on plates for 12 days under LDs and harvested at the
indicated ZT points. WT, non-transgenic Col-0 plants were used as negative controls whereas
asterisks denote non-specific bands detected by the antibody and used as loading controls. Two
exposures of the same blots are shown. Numbers refer to molecular mass according to the migration
of a protein ladder.

635
636

637
638
639
640
Figure 3. ABA promotes CO recruitment to the *FT* promoter
Schematic representation of the *FT* locus (upper panel). Numbers indicate the genomic coordinates
(Mbp) of the *FT* region. The diagram illustrates the *FT* transcript, including the 5' and 3' untranslated
regions (UTRs) (small rectangles), exons (large rectangles), and introns (lines). The location of the
sequences amplified by Real-Time qPCR - the CO responsive elements (CORE 1,2,3,4,5), the
transcription start site (TSS) and the CAAT binding site (CBS) - are shown below as rectangles. The
bottom panel shows the relative enrichment of CO:CIT on the *FT* promoter in isogenic lines
SUC2::CO:CIT #5 (WT and *aba1-6* mutant backgrounds) grown on plates and harvested at ZT12.
Non transgenic, WT Col-0 was used as negative control. Bars are the mean ± SE (n = 4 technical
replicates). An independent experiment is shown as supplementary material.

666
667

668
669
670
Figure 4. The genome wide CO binding is promoted by ABA
(A) Density of mapped reads around the TSS of known genes following ChIPseq of *SUC2::CO:CIT#5*
(WT and *aba1-6* mutant backgrounds) plants (two replicates each) grown on plates and harvested
at ZT12. (B) proportion of ChIPseq peaks detected in *SUC2::CO:CIT #5* plants (after filtering by IDR
and greenscreen blacklisted regions) falling in proximity of various genomic features. (C) intersection
between genes falling next to *SUC2::CO:CIT #5* ChIPseq peaks and *SUC2::CO* upregulated genes,
and enrichment of biological process terms for the intersection, where also *FT* is found. (D)
Distribution of ChIPseq peaks detected in this study across the *FT* genomic region. The bar charts
represent the ChIPseq averaged profiles for *SUC2::CO:CIT #5* (upper panel) and *SUC2::CO:CIT #5*

aba1-6 (lower panel) superimposed to their respective inputs. Significant peaks, including those
found in the ChIPseq of *35S::CO* plants (de los Reyes *et al.,* 2024), are highlighted as colour-coded
rectangles.

Figure 5. ABA production and signalling are required for proper RNA Pol II recruitment at the TSS of *FT*

(A to C) RNAPol II deposition on the *FT* locus in the indicated genotypes expressed as % of input. Each data point represents the mean ± SE (n = 4 technical replicates). Seedlings were grown for 12 days under LDs and harvested at ZT12. In (B), no antibody denotes a negative control for the ChIP assay. In (A) and (C) the WT Col-0 data are the same as samples derived from the same experiment. Iindependent experiments are shown as supplementary material. (D) Relative enrichment of CO:CIT on the *FT* promoter in isogenic lines *SUC2::CO:CIT #5* (WT and *gi-2* mutant backgrounds) at ZT12. Non transgenic, WT Col-0 was used as negative control. Bars are the mean ± SE (n = 4 technical replicates). An independent experiment is shown as supplementary material.

Figure 6. ABA signalling and *GI* act together to promoter CO recruitment to the *FT* promoter and regulate flowering time.
(A) Boxplot of flowering time of the indicated genotypes and isogenic lines derived from the introgression of *SUC2::HA:CO* transgenes into *aba1-6* or *gi* mutant backgrounds. Note that for the *SUC2::HA:CO aba1-6 gi-100* combination, two sibling lines were analysed. An ANOVA test to assess the impact of mutations at the *ABA1* and *GI* loci on leaf number at bolting was run separately for transgenic (*SUC2::HA:CO*) and control plants (vertical bar). For both groups, genotype was a statistically significant predictor of leaf number at bolting ($p < 0.001$). Letters at the top of boxplots indicate if genotypes showed statistically significant differences ($p < 0.05$) according to a Tukey post-hoc test. The number of samples analysed for each genotype is shown at the bottom of the graph.
(B) GI ChIPseq peaks on the *FT* region coincide with DNAse accessible chromatin (referred to as enhancer blocks C and E). CO recruitment to the CORE region is regulated by ABA and GI (thick arrows). ABA and GI might act independently to promote CO binding to the CORE region. ABA might also influence the accessibility of GI to Block C/E (dotted arrows), thus indirectly controlling CO recruitment.

Acknowledgments

We thank personnel at Orto Botanico Città Studi and Viktoriia Kyrychenko for plant care, Fabio Fornara and Federico Zambelli (University of Milan) for insightful comments and guidance on ChIPseq analysis, respectively. This study has been supported by a grant from the Italian Ministerial of Research PRIN 2022 "Light and drought signals integration driving development transitions and adaptations in plants – LIDS" Ref: 2022T2737Y and by a research grant from the HFSP Ref: RGP0011/2019. DM is supported by a research fellowship co-funded by the European Union - ESF, REACT-EU, PON Ricerca e Innovazione 2014-2020. The funders had no role in study design, data collection and analysis, decision to publish, or preparation of the manuscript.

Author contribution

ART, GP, LC: conceptualization and investigation; AS, SC, PKK, GC, TX, EV, DM: investigation; BL: formal analysis; GP, ART, LC: writing - original draft; EK, MG, CT, LC: writing - review & editing; LC: supervision; LC: funding acquisition

No conflict of interest declared

REFERENCES

Adrian J, Farrona S, Reimer JJ, Albani MC, Coupland G, Turck F. 2010. *cis* -Regulatory Elements and Chromatin State Coordinately Control Temporal and Spatial Expression of *FLOWERING LOCUS T* in *Arabidopsis* . The Plant Cell **22**, 1425–1440.

An H, Roussot C, Suárez-López P, *et al.* 2004. CONSTANS acts in the phloem to regulate a systemic signal that induces photoperiodic flowering of *Arabidopsis*. Development **131**, 3615–3626.

Ando E, Ohnishi M, Wang Y, Matsushita T, Watanabe A, Hayashi Y, Fujii M, Ma JF, Inoue S, Kinoshita T. 2013. TWIN SISTER OF FT, GIGANTEA, and CONSTANS have a positive but indirect effect on blue light-induced stomatal opening in Arabidopsis. Plant physiology **162**, 1529–38.

Andrews S. 2010. FastQC: a quality control tool for high throughput sequence data.

Baek D, Kim WY, Cha JY, *et al.* 2020. The GIGANTEA-ENHANCED em LEVEL Complex Enhances Drought Tolerance via Regulation of Abscisic Acid Synthesis1[OPEN]. Plant Physiology **184**, 443–458.

Ben-Naim O, Eshed R, Parnis A, Teper-Bamnolker P, Shalit A, Coupland G, Samach A, Lifschitz E. 2006. The CCAAT binding factor can mediate interactions between CONSTANS-like proteins and DNA. The Plant Journal **46**, 462–476.

Bertauche N, Leung J, Giraudat J. 1996. Protein Phosphatase Activity of Abscisic Acid Insensitive 1 (ABII) Protein from Arabidopsis Thaliana. European Journal of Biochemistry **241**, 193–200.

Biancucci M, Chirivì D, Baldini A, *et al.* 2024. Impact of mutations in the *HEADING DATE 1* gene on transcription and cell wall composition of rice. doi: 10.1101/2024.10.16.618758. [Preprint].

Bu Z, Yu Y, Li Z, Liu Y, Jiang W, Huang Y, Dong A-W. 2014. Regulation of Arabidopsis Flowering by the Histone Mark Readers MRG1/2 via Interaction with CONSTANS to Modulate FT Expression. (B Holt, Ed.). PLoS Genetics **10**, e1004617.

Cao S, Kumimoto RW, Gnesutta N, Calogero AM, Mantovani R, Holt BF. 2014. A Distal *CCAAT* /NUCLEAR FACTOR Y Complex Promotes Chromatin Looping at the *FLOWERING LOCUS T* Promoter and Regulates the Timing of Flowering in *Arabidopsis*. The Plant Cell **26**, 1009–1017.

Cha J-Y, Kim J, Kim T-S, Zeng Q, Wang L, Lee SY, Kim W-Y, Somers DE. 2017. GIGANTEA is a co-chaperone which facilitates maturation of ZEITLUPE in the Arabidopsis circadian clock. Nature Communications **8**, 3.

Chen Y, Song S, Gan Y, Jiang L, Yu H, Shen L. 2020. SHAGGY-like kinase 12 regulates flowering through mediating CONSTANS stability in *Arabidopsis*. Science Advances **6**.

Chen N-Z, Zhang X-Q, Wei P-C, Chen Q-J, Ren F, Chen J, Wang X-C. 2007. AtHAP3b Plays a Crucial Role in the Regulation of Flowering Time in Arabidopsis during Osmotic Stress. BMB Reports **40**, 1083–1089.

Chen S, Zhou Y, Chen Y, Gu J. 2018. fastp: an ultra-fast all-in-one FASTQ preprocessor. Bioinformatics **34**, i884–i890.

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. The Plant journal **16**, 735–743.

Conti L. 2017. Hormonal control of the floral transition: Can one catch them all? Developmental Biology **430**, 288–301.

Corbesier L, Vincent C, Jang S, *et al.* 2007. FT Protein Movement Contributes to Long-Distance Signaling in Floral Induction of *Arabidopsis*. Science **316**, 1030–1033.

Cutler SR, Rodriguez PL, Finkelstein RR, Abrams SR. 2010. Abscisic Acid: Emergence of a Core Signaling Network. Annual Review of Plant Biology **61**, 651–679.

Du J, Zhu X, He K, Kui M, Zhang J, Han X, Fu Q, Jiang Y, Hu Y. 2023. CONSTANS interacts with and antagonizes ABF transcription factors during salt stress under long-day conditions. Plant Physiology doi: 10.1093/plphys/kiad370.

Farrona S, Hurtado L, Bowman JL, Reyes JC. 2004. The *Arabidopsis thaliana* SNF2 homolog AtBRM controls shoot development and flowering. Development **131**, 4965–4975.

Farrona S, Thorpe FL, Engelhorn J, Adrian J, Dong X, Sarid-Krebs L, Goodrich J, Turck F. 2011. Tissue-Specific Expression of *FLOWERING LOCUS T* in *Arabidopsis* Is Maintained Independently of Polycomb Group Protein Repression . The Plant Cell **23**, 3204–3214.

Fornara F, Panigrahi KCS, Gissot L, Sauerbrunn N, Rühl M, Jarillo JA, Coupland G. 2009. Arabidopsis DOF Transcription Factors Act Redundantly to Reduce CONSTANS Expression and Are Essential for a Photoperiodic Flowering Response. Developmental Cell **17**, 75–86.

Fox J, Weisberg S. 2018. *An R Companion to Applied Regression*. Sage.

Frey A, Effroy D, Lefebvre V, Seo M, Perreau F, Berger A, Sechet J, To A, North HM, Marion-Poll A. 2012. Epoxycarotenoid cleavage by NCED5 fine-tunes ABA accumulation and affects seed dormancy and drought tolerance with other NCED family members. The Plant Journal **70**, 501–512.

Fujii H, Chinnusamy V, Rodrigues A, Rubio S, Antoni R, Park S-Y, Cutler SR, Sheen J, Rodriguez PL, Zhu J-K. 2009. In vitro reconstitution of an abscisic acid signalling pathway. Nature **462**, 660–664.

Furihata T, Maruyama K, Fujita Y, Umezawa T, Yoshida R, Shinozaki K, Yamaguchi-Shinozaki K. 2006. Abscisic acid-dependent multisite phosphorylation regulates the activity of a transcription activator AREB1. Proceedings of the National Academy of Sciences **103**, 1988–1993.

García-Alcalde F, Okonechnikov K, Carbonell J, Cruz LM, Götz S, Tarazona S, Dopazo J, Meyer TF, Conesa A. 2012. Qualimap: evaluating next-generation sequencing alignment data. Bioinformatics **28**, 2678–2679.

Garner WW, Allard HA. 1920. Effect of the relative length of day and night and other factors of the environment on growth and reproduction in plants. Monthly Weather Review **48**, 415–415.

Gaspar JM. 2018. Improved peak-calling with MACS2. doi: 10.1101/496521. [Preprint].

Gendron JM, Staiger D. 2023. New Horizons in Plant Photoperiodism. Annual Review of Plant Biology **74**, 481–509.

Gnesutta N, Kumimoto RW, Swain S, Chiara M, Siriwardana C, Horner DS, Holt BF, Mantovani R. 2017. CONSTANS Imparts DNA Sequence Specificity to the Histone Fold NF-YB/NF-YC Dimer. The Plant Cell **29**, 1516–1532.

Hayama R, Sarid-Krebs L, Richter R, Fernández V, Jang S, Coupland G. 2017. PSEUDO RESPONSE REGULATORs stabilize CONSTANS protein to promote flowering in response to day length. The EMBO Journal **36**, 904–918.

Hellens RP, Allan AC, Friel EN, Bolitho K, Grafton K, Templeton MD, Karunairetnam S, Gleave AP, Laing WA. 2005. Transient expression vectors for functional genomics, quantification of promoter activity and RNA silencing in plants. Plant Methods **1**, 13.

Hothorn T, Bretz F, Westfall P. 2008. Simultaneous Inference in General Parametric Models. Biometrical Journal **50**, 346–363.

Huq E, Tepperman JM, Quail PH. 2000. GIGANTEA is a nuclear protein involved in phytochrome signaling in Arabidopsis. Proceedings of the National Academy of Sciences of the United States of America **97**, 9789–9794.

Hwang K, Susila H, Nasim Z, Jung J-Y, Ahn JH. 2019. Arabidopsis ABF3 and ABF4 Transcription Factors Act with the NF-YC Complex to Regulate SOC1 Expression and Mediate Drought-Accelerated Flowering. Molecular Plant **12**, 489–505.

Iacopino S, Jurinovich S, Cupellini L, Piccinini L, Cardarelli F, Perata P, Mennucci B, Giuntoli B, Licausi F. 2019. A Synthetic Oxygen Sensor for Plants Based on Animal Hypoxia Signaling. Plant Physiology **179**, 986–1000.

Imaizumi T, Schultz TF, Harmon FG, Ho LA, Kay SA. 2005. FKF1 F-Box Protein Mediates Cyclic Degradation of a Repressor of *CONSTANS* in *Arabidopsis*. Science **309**, 293–297.

Ito S, Song YH, Josephson-Day AR, Miller RJ, Breton G, Olmstead RG, Imaizumi T. 2012. FLOWERING BHLH transcriptional activators control expression of the photoperiodic flowering regulator *CONSTANS* in *Arabidopsis*. Proceedings of the National Academy of Sciences **109**, 3582–3587.

Jaillais Y, Hothorn M, Belkhadir Y, Dabi T, Nimchuk ZL, Meyerowitz EM, Chory J. 2011. Tyrosine phosphorylation controls brassinosteroid receptor activation by triggering membrane release of its kinase inhibitor. Genes & Development **25**, 232–237.

Jang S, Torti S, Coupland G. 2009. Genetic and spatial interactions between *FT* , *TSF* and *SVP* during the early stages of floral induction in Arabidopsis. The Plant Journal **60**, 614–625.

Kaiserli E, Páldi K, O'Donnell L, Batalov O, Pedmale UV, Nusinow DA, Kay SA, Chory J. 2015. Integration of Light and Photoperiodic Signaling in Transcriptional Nuclear Foci. Developmental Cell **35**, 311–321.

Kardailsky I, Shukla VK, Ahn JH, Dagenais N, Christensen SK, Nguyen JT, Chory J, Harrison MJ, Weigel D. 1999. Activation Tagging of the Floral Inducer *FT*. Science **286**, 1962–1965.

Karimi M, Inzé D, Depicker A. 2002. GATEWAY vectors for Agrobacterium-mediated plant transformation. Trends in Plant Science **7**, 193–195.

Karimi M, De Meyer B, Hilson P. 2005. Modular cloning in plant cells. Trends in Plant Science **10**, 103–105.

Katagiri S, Kamiyama Y, Yamashita K, Iizumi S, Suzuki R, Aoi Y, Takahashi F, Kasahara H, Kinoshita T, Umezawa T. 2024. Accumulation of Phosphorylated SnRK2 Substrate 1 Promotes Drought Escape in *Arabidopsis*. Plant And Cell Physiology **65**, 259–268.

Klasfeld S, Roulé T, Wagner D. 2022. Greenscreen: A simple method to remove artifactual signals and enrich for true peaks in genomic datasets including ChIP-seq data. The Plant Cell **34**, 4795–4815.

Kobayashi Y, Kaya H, Goto K, Iwabuchi M, Araki T. 1999. A Pair of Related Genes with Antagonistic Roles in Mediating Flowering Signals. Science **286**, 1960–1962.

Koncz C, Schell J. 1986. The promoter of TL-DNA gene 5 controls the tissue-specific expression of chimaeric genes carried by a novel type of Agrobacterium binary vector. MGG Molecular & General Genetics **204**, 383–396.

Koops P, Pelser S, Ignatz M, Klose C, Marrocco-Selden K, Kretsch T. 2011. EDL3 is an F-box protein involved in the regulation of abscisic acid signalling in Arabidopsis thaliana. Journal of experimental botany **62**, 5547–5560.

Kubota A, Ito S, Shim JS, *et al.* 2017. TCP4-dependent induction of CONSTANS transcription requires GIGANTEA in photoperiodic flowering in Arabidopsis. (S Hake, Ed.). PLoS Genetics **13**, e1006856.

Kumimoto RW, Adam L, Hymus GJ, Repetti PP, Reuber TL, Marion CM, Hempel FD, Ratcliffe OJ. 2008. The Nuclear Factor Y subunits NF-YB2 and NF-YB3 play additive roles in the promotion of flowering by inductive long-day photoperiods in Arabidopsis. Planta **228**, 709–723.

Kumimoto RW, Siriwardana CL, Gayler KK, Risinger JR, Siefers N, Holt BF. 2013. NUCLEAR FACTOR Y Transcription Factors Have Both Opposing and Additive Roles in ABA-Mediated Seed Germination. (E Huq, Ed.). PLoS ONE **8**, e59481.

Kuromori T, Sugimoto E, Shinozaki K. 2014. Intertissue Signal Transfer of Abscisic Acid from Vascular Cells to Guard Cells. Plant Physiology **164**, 1587–1592.

Laubinger S, Marchal V, Gentillhomme J, Wenkel S, Adrian J, Jang S, Kulajta C, Braun H, Coupland G, Hoecker U. 2006. Arabidopsis SPA proteins regulate photoperiodic flowering and interact with the floral inducer CONSTANS to regulate its stability. Development **133**, 3213–3222.

Lenth R V. 2023. *emmeans: Estimated Marginal Means, aka Least-Squares Means.* .

Leon-Kloosterziel KM, Gil MA, Ruijs GJ, Jacobsen SE, Olszewski NE, Schwartz SH, Zeevaart JAD, Koornneef M. 1996. Isolation and characterization of abscisic acid-deficient Arabidopsis mutants at two new loci. The Plant Journal **10**, 655–661.

Liu L, Adrian J, Pankin A, Hu J, Dong X, von Korff M, Turck F. 2014. Induced and natural variation of promoter length modulates the photoperiodic response of FLOWERING LOCUS T. Nature Communications **5**, 4558.

de los Reyes P, Serrano-Bueno G, Romero-Campero FJ, Gao H, Romero JM, Valverde F. 2024. CONSTANS alters the circadian clock in Arabidopsis thaliana. Molecular Plant **17**, 1204–1220.

Lv X, Zeng X, Hu H, *et al.* 2021. Structural insights into the multivalent binding of the Arabidopsis *FLOWERING LOCUS T* promoter by the CO–NF–Y master transcription factor complex. The Plant Cell **33**, 1182–1195.

Marquès-Bueno MM, Morao AK, Cayrel A, Platre MP, Barberon M, Caillieux E, Colot V, Jaillais Y, Roudier F, Vert G. 2016. A versatile Multisite Gateway-compatible promoter and transgenic line collection for cell type-specific functional genomics in Arabidopsis. The Plant Journal **85**, 320–333.

Martignago D, Siemiatkowska B, Lombardi A, Conti L. 2020. Abscisic Acid and Flowering Regulation: Many Targets, Different Places. International Journal of Molecular Sciences **21**, 9700.

Mathieu J, Yant LJ, Mürdter F, Küttner F, Schmid M. 2009. Repression of Flowering by the miR172 Target SMZ. (C Dean, Ed.). PLoS Biology **7**, e1000148.

Mustilli A-C, Merlot S, Vavasseur A, Fenzi F, Giraudat J. 2002. Arabidopsis OST1 Protein Kinase Mediates the Regulation of Stomatal Aperture by Abscisic Acid and Acts Upstream of Reactive Oxygen Species Production. The Plant Cell **14**, 3089–3099.

Nambara E, Marion-Poll A. 2005. ABSCISIC ACID BIOSYNTHESIS AND CATABOLISM. Annual Review of Plant Biology **56**, 165–185.

Niyogi KK, Grossman AR, Björkman O. 1998. Arabidopsis Mutants Define a Central Role for the Xanthophyll Cycle in the Regulation of Photosynthetic Energy Conversion. The Plant Cell **10**, 1121–1134.

Nohales MA, Liu W, Duffy T, Nozue K, Sawa M, Pruneda-Paz JL, Maloof JN, Jacobsen SE, Kay SA. 2019. Multi-level Modulation of Light Signaling by GIGANTEA Regulates Both the Output and Pace of the Circadian Clock. Developmental Cell **49**, 840-851.e8.

Obermeyer S, Kapoor H, Markusch H, Grasser KD. 2023. Transcript elongation by <scp>RNA</scp> polymerase <scp>II</scp> in plants: factors, regulation and impact on gene expression. The Plant Journal doi: 10.1111/tpj.16115.

Peirats-Llobet M, Han S-K, Gonzalez-Guzman M, Jeong CW, Rodriguez L, Belda-Palazon B, Wagner D, Rodriguez PL. 2016. A Direct Link between Abscisic Acid Sensing and the Chromatin-Remodeling ATPase BRAHMA via Core ABA Signaling Pathway Components. Molecular Plant **9**, 136–147.

Perrella G, Davidson MLH, O'Donnell L, Nastase A-M, Herzyk P, Breton G, Pruneda-Paz JL, Kay SA, Chory J, Kaiserli E. 2018. ZINC-FINGER interactions mediate transcriptional regulation of hypocotyl growth in *Arabidopsis*. Proceedings of the National Academy of Sciences **115**.

Perrella G, Fasano C, Donald NA, Daddiego L, Fang W, Martignago D, Carr C, Conti L, Herzyk P, Amtmann A. 2023. Histone Deacetylase Complex 1 and histone 1 epigenetically

moderate stress responsiveness of *Arabidopsis thaliana* seedlings. New Phytologist doi: 10.1111/nph.19165.

Pin Pa, Nilsson O. 2012. The multifaceted roles of FLOWERING LOCUS T in plant development. Plant, Cell & Environment **35**, 1742–1755.

Ramírez F, Dündar F, Diehl S, Grüning BA, Manke T. 2014. deepTools: a flexible platform for exploring deep-sequencing data. Nucleic Acids Research **42**, W187–W191.

Rédei GP. 1962. Supervital Mutants of Arabidopsis. Genetics **47**, 443–460.

Riboni M, Galbiati M, Tonelli C, Conti L. 2013. GIGANTEA enables drought escape response via abscisic acid-dependent activation of the florigens and SUPPRESSOR of OVEREXPRESSION of CONSTANS11[c][w]. Plant Physiology **162**, 1706–1719.

Riboni M, Robustelli Test A, Galbiati M, Tonelli C, Conti L. 2016. ABA-dependent control of *GIGANTEA* signalling enables drought escape via up-regulation of *FLOWERING LOCUS T* in *Arabidopsis thaliana*. Journal of Experimental Botany **67**, 6309–6322.

Sarid-Krebs L, Panigrahi KCS, Fornara F, Takahashi Y, Hayama R, Jang S, Tilmes V, Valverde F, Coupland G. 2015. Phosphorylation of <scp>CONSTANS</scp> and its <scp>COP</scp> 1-dependent degradation during photoperiodic flowering of Arabidopsis. The Plant Journal **84**, 451–463.

Sawa M, Kay SA. 2011. GIGANTEA directly activates *Flowering Locus T* in *Arabidopsis thaliana*. Proceedings of the National Academy of Sciences **108**, 11698–11703.

Sawa M, Nusinow DA, Kay SA, Imaizumi T. 2007. FKF1 and GIGANTEA Complex Formation Is Required for Day-Length Measurement in *Arabidopsis*. Science **318**, 261–265.

Seo M, Marion-Poll A. 2019. Abscisic acid metabolism and transport. 1–49.

Shen L. 2024. GeneOverlap: Test and visualize gene overlaps. R package version 1.42.0.

Siemiatkowska B, Chiara M, Badiger BG, *et al.* 2022. GIGANTEA Is a Negative Regulator of Abscisic Acid Transcriptional Responses and Sensitivity in Arabidopsis. Plant and Cell Physiology **63**, 1285–1297.

Simon NML, Graham CA, Comben NE, Hetherington AM, Dodd AN. 2020. The Circadian Clock Influences the Long-Term Water Use Efficiency of Arabidopsis. Plant Physiology **183**, 317–330.

Siriwardana CL, Gnesutta N, Kumimoto RW, Jones DS, Myers ZA, Mantovani R, Holt BF. 2016. NUCLEAR FACTOR Y, Subunit A (NF-YA) Proteins Positively Regulate Flowering and Act Through FLOWERING LOCUS T. (GK Muday, Ed.). PLOS Genetics **12**, e1006496.

Song YH, Estrada DA, Johnson RS, Kim SK, Lee SY, MacCoss MJ, Imaizumi T. 2014. Distinct roles of FKF1, GIGANTEA, and ZEITLUPE proteins in the regulation of CONSTANS stability in *Arabidopsis* photoperiodic flowering. Proceedings of the National Academy of Sciences **111**, 17672–17677.

Song L, Huang SC, Wise A, Castanon R, Nery JR, Chen H, Watanabe M, Thomas J, Bar-Joseph Z, Ecker JR. 2016. A transcription factor hierarchy defines an environmental stress response network. Science **354**, aag1550–aag1550.

Song YH, Kubota A, Kwon MS, *et al.* 2018. Molecular basis of flowering under natural long-day conditions in Arabidopsis. Nature Plants **4**, 824–835.

Song YH, Shim JS, Kinmonth-Schultz HA, Imaizumi T. 2015. Photoperiodic Flowering: Time Measurement Mechanisms in Leaves. Annual Review of Plant Biology **66**, 441–464.

Song YH, Smith RW, To BJ, Millar AJ, Imaizumi T. 2012. FKF1 Conveys Timing Information for CONSTANS Stabilization in Photoperiodic Flowering. Science **336**, 1045–1049.

Suárez-López P, Wheatley K, Robson F, Onouchi H, Valverde F, Coupland G. 2001. CONSTANS mediates between the circadian clock and the control of flowering in Arabidopsis. Nature **410**, 1116–1120.

Takagi H, Hempton AK, Imaizumi T. 2023. Photoperiodic flowering in Arabidopsis: Multilayered regulatory mechanisms of CONSTANS and the florigen FLOWERING LOCUS T. Plant Communications, 100552.

Tian H, Li Y, Wang C, *et al.* 2021. Photoperiod-responsive changes in chromatin accessibility in phloem companion and epidermis cells of Arabidopsis leaves. The Plant Cell **33**, 475–491.

Tiwari SB, Shen Y, Chang H, *et al.* 2010. The flowering time regulator CONSTANS is recruited to the *FLOWERING LOCUS T* promoter via a unique *cis* -element. New Phytologist **187**, 57–66.

Truernit E, Sauer N. 1995. The promoter of the Arabidopsis thaliana SUC2 sucrose-H+ symporter gene directs expression of beta-glucuronidase to the phloem: Evidence for phloem loading and unloading by SUC2. Planta **196**.

Valverde F, Mouradov A, Soppe W, Ravenscroft D, Samach A, Coupland G. 2004. Photoreceptor Regulation of CONSTANS Protein in Photoperiodic Flowering. Science **303**, 1003–1006.

Wang H, Pan J, Li Y, Lou D, Hu Y, Yu D. 2016. The DELLA-CONSTANS Transcription Factor Cascade Integrates Gibberellic Acid and Photoperiod Signaling to Regulate Flowering. Plant Physiology **172**, 479–488.

Wenkel S, Turck F, Singer K, Gissot L, Le Gourrierec J, Samach A, Coupland G. 2006. CONSTANS and the CCAAT Box Binding Complex Share a Functionally Important Domain and Interact to Regulate Flowering of *Arabidopsis*. The Plant Cell **18**, 2971–2984.

Wickham H. 2011. ggplot2. WIREs Computational Statistics **3**, 180–185.

Wigge PA, Kim MC, Jaeger KE, Busch W, Schmid M, Lohmann JU, Weigel D. 2005. Integration of spatial and temporal information during floral induction in Arabidopsis. Science (New York, N.Y.) **309**, 1056–1059.

YOSHIDA T, FUJITA Y, MARUYAMA K, MOGAMI J, TODAKA D, SHINOZAKI K, YAMAGUCHI-SHINOZAKI K. 2015. Four *Arabidopsis* AREB/ABF transcription factors function predominantly in gene expression downstream of SnRK2 kinases in abscisic acid signalling in response to osmotic stress. Plant, Cell & Environment **38**, 35–49.

Yoshida T, Fujita Y, Sayama H, Kidokoro S, Maruyama K, Mizoi J, Shinozaki K, Yamaguchi-Shinozaki K. 2010. AREB1, AREB2, and ABF3 are master transcription factors that cooperatively regulate ABRE-dependent ABA signaling involved in drought stress tolerance and require ABA for full activation. The Plant Journal **61**, 672–685.

Yu G, Wang L-G, He Q-Y. 2015. ChIPseeker: an R/Bioconductor package for ChIP peak annotation, comparison and visualization. Bioinformatics **31**, 2382–2383.

Zeng X, Lv X, Liu R, He H, Liang S, Chen L, Zhang F, Chen L, He Y, Du J. 2022. Molecular basis of CONSTANS oligomerization in *FLOWERING LOCUS T* activation. Journal of Integrative Plant Biology **64**, 731–740.

Zhang W, Zhang T, Wu Y, Jiang J. 2012. Genome-Wide Identification of Regulatory DNA Elements and Protein-Binding Footprints Using Signatures of Open Chromatin in *Arabidopsis*. The Plant Cell **24**, 2719–2731.

Zhang C, Zhou Q, Liu W, Wu X, Li Z, Xu Y, Li Y, Imaizumi T, Hou X, Liu T. 2022. *BrABF3* promotes flowering through the direct activation of *CONSTANS* transcription in pak choi. The Plant Journal **111**, 134–148.

Zicola J, Liu L, Tänzler P, Turck F. 2019. Targeted DNA methylation represses two enhancers of FLOWERING LOCUS T in Arabidopsis thaliana. Nature Plants **5**, 300–307.