The theory of massively repeated evolution and full identifications of Cancer Driving Nucleotides (CDNs)

Lingjie Zhang[1], Tong Deng[1], Zhongqi Liufu[1,2], Xueyu Liu[1], Bingjie Chen[1,3], Zheng Hu[4], Chenli Liu[4], Xuemei Lu[2], Haijun Wen[1,*], Chung-I Wu[1,5*].

[1] State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, Guangdong, China.

[2] State Key Laboratory of Genetic Resources and Evolution/Yunnan Key Laboratory of Biodiversity Information, Kunming Institute of Zoology, The Chinese Academy of Sciences, Kunming 650223, China.

[3] GMU-GIBH Joint School of Life Sciences, Guangzhou Medical University, Guangzhou 511436, China.

[4] CAS Key Laboratory of Quantitative Engineering Biology, Shenzhen Institute of Synthetic Biology, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China.

[5] Department of Ecology and Evolution, University of Chicago, Chicago, Illinois 60637, USA.

* Corresponding authors: wenhj5@mail.sysu.edu.cn; ciwu@uchicago.edu.

ABSTRACT

Tumorigenesis, like most complex genetic traits, is driven by the joint actions of many mutations. At the nucleotide level, such mutations are Cancer Driving Nucleotides (CDNs). The full sets of CDNs are necessary, and perhaps even sufficient, for the understanding and treatment of each cancer patient. Currently, only a small fraction of CDNs is known as most mutations accrued in tumors are not drivers. We now develop the theory of CDNs on the basis that cancer evolution is massively repeated in millions of individuals. Hence, any advantageous mutation should recur frequently and, conversely, any mutation that does not is either a passenger or deleterious mutation. In the TCGA cancer database (sample size $\mathbf{n}$ = 300 - 1000), point mutations may recur in i out of $\mathbf{n}$ patients. This study explores a wide range of mutation characteristics to determine the limit of recurrences (i^*) driven solely by neutral evolution. Since no neutral mutation can reach i^* = 3, all mutations recurring at $i \geq 3$ are CDNs. The theory shows the feasibility of identifying almost all CDNs if $\mathbf{n}$ increases to 100,000 for each cancer type. At present, only < 10% of CDNs have been identified. When the full sets of CDNs are identified, the evolutionary mechanism of tumorigenesis in each case can be known and, importantly, gene targeted therapy will be far more effective in treatment and robust against drug resistance.

INTRODUCTION

Cancers are complex genetic traits with multiple mutations that interact to yield the ensemble of tumor phenotypes. The ensemble has been characterized as "cancer hallmarks" that include sustaining growth signaling, evading growth suppression, resisting apoptosis, achieving immortality, executing metastasis and so on (Hanahan and Weinberg 2000; Hanahan and Weinberg 2011; Hanahan 2022). It seems likely that each of the 6-10 cancer hallmarks is governed by a set of mutations. Most, if not all, of these mutations are jointly needed to drive the tumorigenesis.

In the genetic sense, cancers do not differ fundamentally from other complex traits whereby multiple mutations are simultaneously needed to execute the program (Wu and Ting 2004; Stratton et al. 2009; Yates and Campbell 2012; Ortmann et al. 2015; Marques et al. 2019; Campbell et al. 2020). A recent example is SARS-CoV-2. The early onset of COVID-19 requires all four mutations of the D614G group and the later Delta strain has 31 mutations accrued in three batches (Ruan, Hou, et al. 2022; Ruan, Wen, et al. 2022). While cancer research has often proceeded one mutation at a time, each of the

mutations has been shown to be insufficient for tumorigenesis until many (Ortmann et al. 2015; Takeda 2021; Hodis et al. 2022) are co-introduced.

We now aim for the identification of all (or at least most) of the driver mutations in each patient. Both functional tests and treatments demand such identifications. The number of key drivers has been variously estimated to be 6 – 10 (Martincorena et al. 2017; Anandakrishnan et al. 2019; Campbell et al. 2020). Although cancer driving "point mutations", referred to as Cancer Driving Nucleotides (or CDNs), are not the only drivers, they are indeed abundant (Campbell et al. 2020). Furthermore, CDNs, being easily quantifiable, may be the only type of drivers that can be fully identified (see below). Here, we will focus on the clonal mutations present in all cells of the tumor without considering within-tumor heterogeneity for now (Ling et al. 2015; Turajlic et al. 2019; Black and McGranahan 2021).

Since somatic evolution proceeds in parallel in millions of humans, point mutations can recur multiple times as shown in **Fig. 1**. The recurrences should permit the detection of advantageous mutations with unprecedented power. The converse should also be true that mutations that do not recur frequently are unlikely to be advantageous. **Fig. 1** depicts organismal evolution and cancer evolution. While both panels **A** and **B** show 7 mutations, there can be two patterns for cancer evolution - the pattern in **(C)** where all mutations are at different sites is similar to the results of organismal evolution whereas the pattern in **(D)** is unique in cancer evolution. The hotspot of recurrences shown in **(D)** holds the key to finding all CDNs in cancers.

Fig. 1

In the literature, hotspots of recurrent mutations have been commonly reported (Gartner et al. 2013; Chang et al. 2016; Cannataro et al. 2018; Buisson et al. 2019; Hess et al. 2019; Stobbe et al. 2019; Juul et al. 2021; Nesta et al. 2021; Zhao et al. 2021; Bergstrom et al. 2022; Sherman et al. 2022; Wong et al. 2022; Zeng and Bromberg 2022). A hotspot, however, could be either a mutational or functional hotspot. Mutational hotspots are the properties of the mutation machinery that would include nucleotide composition, local chromatin structure, timing of replication, etc. (Stamatoyannopoulos et al. 2009; Pleasance et al. 2010; Makova and Hardison 2015; Polak et al. 2015; Martincorena et al. 2017). In contrast, functional hotspots are CDNs under positive selection. CDN evolution is akin to "convergent evolution" that repeats itself in different taxa (He, Xu, and Shi 2020; He, Xu, Zhang, et al. 2020; Wu et al. 2020) and is generally considered the most convincing proof of positive selection.

While many studies conclude that sites of mutation recurrence are largely mutational hotspots (Buisson et al. 2019; Hess et al. 2019; Stobbe et al. 2019; Nesta et al. 2021; Bergstrom et al. 2022), others deem them functional hotspots, driven by positive selection (Gartner et al. 2013; Chang et al. 2016; Bailey et al. 2018; Cannataro et al. 2018; Juul et al. 2021; Zhao et al. 2021; Zeng and Bromberg 2022). In the attempt to distinguish between these two hypotheses, studies make assumptions, often implicitly, about the relative importance of the two mechanisms in their estimation procedures. The conclusions naturally manifest the assumptions made when extracting information on mutation and selection from the same data (Elliott and Larsson 2021).

This study consists of three parts. First, the mutational characteristics of sequences surrounding CDNs are analyzed. Second, a rigorous probability model is developed to compute the recurrence level at any sample size. Above a threshold of recurrence, all mutations are CDNs. Third, we determine the necessary sample sizes that will yield most, if not all, CDNs. In the companion study (supplement File

95 S3), the current cancer genomic data are analyzed for the characteristics of CDNs that have already
96 been discovered. Together, these two studies show how full functional tests and precise target therapy
97 can be done on each cancer patient.
98
RESULTS
In **PART I**, we search for the general mutation characteristics at and near high recurrence sites by both
machine learning and extensive sequence comparisons. In **PART II**, we develop the mathematical
theory for the maximal level of recurrences of neutral mutations (designated i^*). CDNs are thus defined
as mutations with $\geq i^*$ recurrences in **n** cancer samples. We then expand in **PART III** the theory to very
large sample sizes ($n \geq 10^5$), thus making it possible to identify all CDNs.
105
To carry out the analyses, we first compile from the TCGA database the statistics of multiple hit sites (i
hits in **n** samples) in 12 cancer types. This study focuses on the mutational characteristics pertaining
to recurrence sites. We often present the three cancer types with the largest sample sizes (lung, breast
and CNS cancers), while many analyses are based on pan-cancer data. Analyses of every of the 12
cancer types individually will be done in the companion paper. The TCGA database is used as it is well
established and covers the entire coding region (Weinstein et al. 2013). Other larger databases
(Cerami et al. 2012; Tate et al. 2019; de Bruijn et al. 2023) are employed when the whole exon
analyses are not crucial.
114
The compilation of multi-hit sites across all genes in the genome
Throughout the study, S_i denotes the number of synonymous sites where the same nucleotide
mutation occurs in i samples among **n** patients. A_i is the equivalent of S_i for non-synonymous (amino
acid altering) sites. **Table 1** presents the numbers from lung cancer for demonstration. It also shows
the A_i and S_i numbers with CpG sites filtered out. There are 22.7 million nonsynonymous sites, among
which ~ 0.2 million sites have one hit (A_1 = 195,958) in 1035 patients. The number then decreases
sharply as i increases. Thus, A_2 = 2946 (number of 2-hit sites), A_3 = 99, and $A_4 + A_5 + ... = 79$. We also
note that the A_i/S_i ratio increases from 2.89, 2.82, 3.04 to 4.71 and so on.
123
124   **Table 1**
125   **Fig. 2**
126
Fig. 2 shows the average of A_i and S_i among the 12 cancer types (see **Methods**). The salient features
are shown by differences between the solid and dotted lines. As will be detailed in **PART II**, the dotted
lines, extending linearly from i = 0 to i = 1 in logarithmic scale, should be the expected values of A_i and
S_i, if mutation rate is the sole driving force. In the actual data, A_1 and S_1 decrease to ~ 0.002 of A_0 and
S_0, the step being least affected by selection (see **PART II** later). For A_2 and S_2, the decrease is only
~ 0.01 of A_1 and S_1. The decrease from i to i+1 becomes smaller and smaller as i increases, suggesting
that the process in not entirely neutral. Furthermore, the lower panel of **Fig. 2** shows that A_i/S_i
continues to rise as i increases. These patterns again suggest a stronger positive selection at higher i
values. The extrapolation lines shown in **Fig. 2** roughly define i = 3 as a cutoff where the expected A_3
falls below 1 (see **PART II** for details). The precise model of **PART II** will define high recurrence sites (i
≥ 3) as CDNs.
138
139
140
141

PART I - The mutational characteristic of high recurrence sites

In this part, the analyses are done in two different ways. The sequence-feature approach is to examine the mutation characteristics of sequence features (say, 3 mers, 5 mers, etc.) across patients. The patient-feature approach is to examine patients for their mutation signatures and mutation loads.

1. The sequence-feature approach

The simplest and best-known sequence feature associated with high mutation rate is CpG sites. In mammals, methylation and de-amination would enhance the mutation rate from <u>C</u>pG to <u>T</u>pG or Cp<u>A</u> by 5~10-fold (Hodgkinson and Eyre-Walker 2011; Ségurel et al. 2014). As the CpG site mutagenesis has been extensively reported, we only present the confirmation in the Supplement (**Fig. S1**). Indeed, CpG sites account for ~6.5% of the coding sequences but contributing ~22% among the mutated sites in **Fig. 2**. Hence, the 7-fold increase in the CpG mutation rate should contribute more to A_i and S_i as i increases. **Table 1** has shown the effects of filtering out CpG sites in the counts of recurrences. Clearly, CpG sites do contribute disproportionately to the recurrences but, even when they are separately analyzed, the conclusion is unchanged. As shown later in **PART II**, every increment of i should decrease the site number by ~0.002 in the TCGA database. Thus, even with a 10-fold increase in mutation rate, the decrease rate would still be 0.02. In the theory sections, CpG site mutations are incorporated into the model.

In this section, we aim to find out how extreme the mutation mechanisms must be to yield the observed recurrences. If these mechanisms seem implausible, we may reject the mutational-hotspot hypothesis and proceed to test the functional hotspot hypothesis.

The analyses of mutability variation by Artificial Intelligence (AI)

The variation of mutation rate at site level could be shaped by multiple mutational characteristics. Epigenomic features, such as chromatin structure and accessibility, could affect regional mutation rate at kilobase or even megabase scale (Stamatoyannopoulos et al. 2009; Makova and Hardison 2015), while nucleotide biases by mutational processes typically span only a few base pairs around the mutated site (Roberts et al. 2013; Haradhvala et al. 2018; Herzog et al. 2021). AI-powered multi-modal integration offers a new tool to quantify the joint effect of various factors on mutation rate variability (Luo et al. 2019; Sherman et al. 2022; Song et al. 2023). Here we explore the association between the mutation recurrence (i) and site-level mutation rate predicted by AI.

Fig. 3

Fig. 3A shows the mutation rate landscape across all recurrence sites in breast, CNS and lung cancer using the deep learning framework *Dig* (Sherman et al. 2022). In this approach, the mutability of a focus site is calculated based on both local stretch of DNA and broader scale of epigenetic features. The X-axis shows all mutated sites with $i > 0$ scanned by *Dig*. While the mutation rate fluctuates around the average level, we detect no significant difference in mutation rate as a function of i (**Methods**). In CNS, two sites exhibited exceptional mutability at $i = 6$, surpassing the average by 10-fold. Unsurprisingly, these two are CpG sites and correspond to amino acid change of V774M and R222C in *EGFR*, which are canonical actionable driver mutations in glioma target therapy. In other words, the two sites called by AI for possible high mutability appear to be selection driven.

In **Fig. 3B**, we take a closer look at how CDN is situated against the background of mutation rate variation, using the example of *PAX3* (Paired Box Homeotic Gene 3) (Wang et al. 2008; Li et al. 2019).

In this typical example, *Dig* predicts site mutability to vary from site to site. In the lower panel is an expanded look at a stretch of 200 bps. In this stretch, about 8% of sites are 5 to 10-fold more mutable than the average. But none of them are mutated in the data ($i = 0$). There are indeed three sites with $i >$ 0 in this DNA segment including a CDN site C1271T (marked by the red star). This CDN site is estimated to have a 2-fold elevation in mutability, which is less than 1/50 of the necessary mutability to reach $i \geq 3$. The other two mutated sites, marked by the green star, are also indicated.

Other AI methods have also been used in the mutability analysis (Fang et al. 2022), reaching nearly the same conclusion. Overall, while AI often suggests sequence context to influence the local variation in mutation rate, the reported variation does not correspond to the distribution of CDNs. In the next subsection, we further explore the local contexts for potential biases in mutability.

The conventional analyses of local contexts - from 3-mers to 100-mers
Since the AI analyses suggest the dominant role of local sequence context in mutability, we carry out such conventional analyses in depth. Other than the CpG sites, local features such as the T<u>C</u>W (W = A or T) motif recognized by APOBEC family of cytidine deaminases (Burns et al. 2013; Roberts et al. 2013), would have impacts as well. We first calculate the mutation rate for motifs of 3-mer, 5-mer and 7-mer, respectively, with 64, 1024, 16384 in number, (see **Methods**). The pan-cancer analyses across the 12 cancer types are shown in **Fig. 4A**. We use **α** to designate the fold change between the most mutable motif and the average. Since the number of motifs increases 16-fold between each length class, the **α** value increases from 4.7 to 8.8 and then to 11.5. Nevertheless, even the most mutable 7-mer, TAA<u>C</u>GCG, which has a CpG site at the center, is only 11.52-fold higher than the average. This spread is insufficient to account for the high recurrences, which decrease to ~0.002 for each increment of i (see **PART II** below).

Fig. 4

A more direct approach is given in **Fig. 4B** explained in the legends. The absence of a trend shows that the high recurrence sites are not associated with the mutability of the motif. In **Fig. 4C**, the analysis extends to longer motifs of 21 bp, 41 bp, 61 bp, 81 bp and 101 bp surrounding the high-recurrence sites. For example, the motif of 101 bp may be (10, 90), (20, 80) and so on either side of a recurrence site. We then compute the pairwise differences in sequences of the motifs among recurrence sites. The logic is that, if certain motifs dictate high mutation rates, we may observe unusually high sequence similarity in the pairwise comparisons. As can be seen in all 5 panels of **Fig. 4C**, there are no outliers in the tail of the distribution. In other words, the sequences surrounding the high-recurrence sites appear rather random.

In conclusion, the analyses by the sequence approach do not find any association between high recurrence sites ($i \geq 3$) and the mutability of the local sequences.

2. The patient-feature approach
In this second approach, we examine the mutation characteristics among patients across sequence features. The first question is whether high recurrence sites tend to happen in patients with higher mutation loads. **Fig. 5A** depicts the distribution of mutation loads among patients harboring a CDN of recurrence i. Hence, a patient's load may appear several times in the plot, each appearance corresponding to one CDN in the patient's data. For the comparison across i values, the mutation load is normalized by a z-score within each cancer population to equalize the 3 cancer types. The overall

236 trend shows consistently that patients with recurrence sites do not bias toward high mutation loads.
237 The presence of recurrence sites in patients with low mutation loads suggests that overall mutation
238 burden is not a determining factor of recurrence.
239
240 <center>**Fig. 5**</center>
241
With the results of **Fig. 5A**, we then ask a related question: whether these high recurrence mutations
are driven by factors that affect mutation characteristics. Such influences have been captured by the
analyses of "mutational signatures" (Alexandrov et al. 2013; Alexandrov et al. 2020). Each signature
represents a distinct mutation pattern (e.g., high rate of TCT -> TAT and other tri-nucleotide changes)
associated with a known factor, such as smoking or an aberrant mutator (aristolochic acid, for
example). Each patient's mutation profile can then be summarized by the composite of multiple
mutational signatures.
249
The issue is thus whether a patient's CDNs can be explained by the patient's composite mutational
signatures. **Fig. 5B** reveals that in lung cancer, the signature compositions among patients with
different recurrence cutoffs are statistically indistinguishable (**Methods**). Smoking (signature SBS4)
consistently emerges as the predominant mutational process across all levels of recurrences. In breast
cancer, while SBS2 and SBS13 exhibit some differences in the bins of $i^* = 2$ and $i^* = 3$, the profiles
remain rather constant for all bins of $i^* \geq 3$. The two lowest bins, not unexpectedly, are also different
from the rest in the total mutation load (see **Fig. 5A**). In **table S1**, we provide a comprehensive review
of supporting literature on genes with recurrence sites of $i \geq 3$ for breast cancer. In CNS cancer, SBS11
appears significantly different across bins, in particular, $i \geq 20$. This is a signature associated with
Temozolomide treatment and should be considered a secondary effect. In short, while there are
occasional differences in mutational signatures across i^* bins, none of such differences can account for
the recurrences (see **PART II**).
262
263 To conclude **PART I**, the high-recurrence sites do not stand out for their mutation characteristics.
264 Therefore, the variation in mutation rate across the whole genome can reasonably be approximated
265 mathematically by a continuous distribution, as will be done below.
266
PART II - The theory of Cancer Driving Nucleotides (CDNs)
We now develop the theory for S_i and A_i under neutrality where i is the recurrence of the mutation at
each site. We investigate the maximal level of neutral mutation recurrences (i^*), above which the
expected values of S_i and A_i are both < 0. Since no neutral mutations are expected to reach i^*, every
mutation with the recurrence of i^* or larger should be non-neutral. Importantly, given that the
expected S_i and A_i is a function of U^i where $U = \mathbf{n}E(\mathbf{u})$ is in the order of 10^{-2} and 10^{-3}, i^* is insensitive to
a wide range of mutation scenarios. For that reason, the conclusion is robust.
274
The mutation rate of each nucleotide ($\boldsymbol{u}$) follows a Gamma distribution with a scale parameter θ and a
shape parameter k. Gamma distribution is often used for its flexibility and, in this context, models the
waiting time required to accumulate k mutations. Its mean (=$k\theta$) and variance (=$k\theta^2$) are determined
by both parameters but the shape (skewness and kurtosis) is determined only by k. In particular, the
Gamma distribution has a long tail suited to modeling a small number of sites with very high mutation
rate.
281
282 We now use synonymous (S_i) mutations as the proxy for neutrality. Hence, in **n** samples,

283
$$S_i = \sum_{l=1}^{L_S} C_n^i u(l)^i [1 - u(l)]^{n-i} \sim C_n^i L_S E(u^i)$$

284   Eq. 1

285 where L_S is the total number of synonymous sites and $u(l)$ is the mutation rate of site l. In the equation

286 above, the term $[1 - u(l)]^{n-i}$ is dropped. We note that $[1 - u(l)]^{n-i} \sim e^{[-u(l)(n-i)]} \sim 1$ as u is in the

287 order of 10^{-6} and n is in the order of 10^2 from the TCGA data. With the gamma distribution of u

288 whereby the ith moment is given by

289
$$E(u^i) = \frac{\Gamma(k+i)}{\Gamma(k)} \theta^i = \frac{\Gamma(k+i)}{\Gamma(k) \cdot k^i} E(u)^i$$

290 we obtain:

291
$$S_i = L_S \cdot g(i,k) [nE(u)]^i$$

292 Where:

293
$$g(i,k) = C_{k+i-1}^i \cdot \frac{1}{k^i}$$

294 In a condensed form,

295
$$S_i = G \cdot [nE(u)]^i$$

296   Eq. 2

297 where $G = L_S \cdot g(i,k)$. Similarly, if nonsynonymous mutations are assumed neutral, then,

298
$$A_i = L_A \cdot g(i,k) [nE(u)]^i \sim 2.3 \cdot S_i$$

299   Eq. 3

The number of 2.3 is roughly the ratio of the total number of nonsynonymous over that of synonymous

sites (Hartl and Clark 1989; Li 1997; Chen et al. 2019). This number would vary moderately among

cancers depending on their nucleotide substitution patterns.

$E(u)$ of Eqs. 2-3 is generally $(1 \sim 5) \times 10^{-6}$ per site in cancer genomic data and n is generally between

300 and 1000. Hence, $nE(u)$ is the total mutation rate summed over all n patients and is generally

between 0.001 and 0.005 in the TCGA data. Given $nE(u)$ is in the order of 10^{-3}, S_i and A_i would both

decrease by $2 \sim 3$ orders of magnitude with each increment of i by 1. We note that the total number of

synonymous sites, L_S, is $\sim 0.9 \times 10^7$ and L_A is ~ 2.3 times larger. Therefore, $S_3 < 1$ and $A_3 < 1$. When i

reaches 4, $S_{i \geq 4}$ and $A_{i \geq 4}$ would both be $\ll 1$ when averaged over cancer types.

For each cancer type, the conclusion of $S_3 < 1$ and $A_3 < 1$ is valid with the actual value of S_3 and A_3

ranging between 0.01 and 1. In other words, with $n < 1000$, neutral mutations are unlikely to recur 3

times or more in the TCGA data ($i^* = 3$). While $S_{i \geq 3}$ and $A_{i \geq 3}$ sites are high-confidence CDNs, the value of

i^* is a function of n. At $n \leq 1000$ for the TCGA data, i^* should be 3 but, when n reaches 10,000, i^* will be

6. The benefits of large n's will be explored in **PART III**.

Possible outliers to the distribution of mutation rate

Although we have explored extensively the sequence contexts, other features beyond DNA sequences

could still lead to outliers to mathematically distributions. These features may include DNA stem loops

(Buisson et al. 2019) or unusual epigenetic features (Zheng et al. 2014; Makova and Hardison 2015;

Supek and Lehner 2015). We therefore expand the model by assuming a small fraction of sites (p) to

be hyper-mutable that is α fold more mutable than the genomic mean. Most likely, α and p are the

inverse of each other. For the bulk of sites ($1-p$) of the genome, we assume that their mutations follow

the Gamma distribution. (Nevertheless, the bulk can be assumed to have a fixed mutation rate of $E(u)$

without affecting the conclusion qualitatively.)

We let p range from 10^{-2} to 10^{-5} and α up to 1000. As no stretches of DNA show such unusually high

mutation rate (1000-fold higher than the average), such sites are assumed to be scattered across the

genome and are rare. With the parameter space of defined above, we choose the (p, α) pairs that agree

with the observed values of S_1 to S_3 which are sufficiently large for estimations. **Table 2** presents the

value range and standard deviation for p and α across the 6 cancer types that have >500 patient

samples. Among the 6 cancer types, the lung cancer data do not conform to the constraints and we set

$p = 0$. With observed values for $S_{1\sim3}$ as constraints, S_4 rarely exceeds 1. Hence, even with the purely

conjectured existence of outliers in mutation rate, $i^* = 4$ is already too high a cutoff.

Table 2

Table 2 suggests that p has to be smaller than 10^{-5} and $\alpha > 1000$ to yield $S_4 > 1$. Since the coding

region has 3×10^7 sites, $p < 10^{-5}$ would mean that the outliers are at most in the low hundreds. In other

words, the number of high recurrence sites projected by the theory is close to the observed numbers.

Therefore, there are really no unknown outlier sites of high mutation rate. Positive selection would be

a more straightforward explanation, explained below.

The influence of selection on mutation occurrences

We now show that, although the mutational bias alone cannot account for the high occurrences,

selection can easily do so. We assume a fraction, f, of A_i's to be under positive selection. The fraction

should be small, probably $\ll 0.01$, and will be labeled A_i^*. The rest, labeled A_i is considered neutral.

Hence, A_i is proportional to $[nE(u)]^i$ and $A_i/S_i = L_A/L_S \sim 2.3$. Like $S_{i\geq4}$, $A_{i\geq4} \ll 1$. In contrast,

$$A_i^* = G^*[w \cdot nE(u)]^i$$

350 Eq.4

$$\frac{A_i^*}{S_i} = \frac{G^*}{G}w^i$$

Eq.5

where G^* is also a constant but its value depends on f. The crucial parameter, w (= $2Ns$), is the selective

advantage (s) scaled by the population size of progenitor cancer cell (N). Since w can easily be >10,

even at $i = 3$, A_3^*/S_3 would be >100 as large as A_1^*/S_1. In other words, observed mutation recurrences

at $i \geq 3$ for advantageous mutation should not be uncommon. Eq. 4 also shows that w and $E(u)$ jointly

affect the recurrence; therefore, CpG sites (many of which fall in functional sites) are expected to be

strongly represented among high recurrence sites.

PART III. The theory of large samples (n > 10^5) and identification of all CDN's

Using the theory of CDN developed above, the companion paper shows that each sequenced cancer

genome in the current databases, on average, harbors only $1\sim2$ CDNs. The number varies in this range

depending on the cancer type (supplement File S3). For comparison, tumorigenesis may require at

least $5\sim10$ driver mutations as estimated by various criteria (Armitage and Doll 1954; Hanahan and

Weinberg 2011; Belikov 2017; Martincorena et al. 2017; Anandakrishnan et al. 2019; Campbell et al.

2020). The results show that there are many more CDNs that have not been discovered. This is not

unexpected since most CDNs are found in <1% of patients. If each CDN is observed in 1% of patients

and each patient has 5 CDNs, then there should be at least 500 CDNs for each cancer type.

In the companion study that uses the A/S ratios of **Fig. 1**, the estimated number of CDNs ranges from

500 to 2000, whereas the current estimates based on $A_{i\geq3}$ sites is only $50\sim100$ (supplement File S3).

Where, then, are the undiscovered CDNs and how to find them? Since all $A_{i\geq3}$ sites are concluded to be
CDNs, the bulk of CDNs must be among A_1 and A_2 sites. The best way to identify the CDNs hidden in A_1
and A_2 is to increase the sample size, **n**, dramatically.
375

376   We hence extend Eq. 1 for S_i and A_i to large **n**'s. Note that Eq. 1 drops the term of $(1-u)^{n-i} \sim e^{-u(n-i)}$
377   as it is ~ 1, when **nE(u)** $\ll 1$. With a large **n** when $e^{-u(n-i)}$ is not near 1, the recurrence of mutations
378   would follow a Poisson distribution with the expected value of **nE(u)**. Assuming that u follows a
379   gamma distribution with a shape parameter of k, the probability of observing i mutations would
380   follow the negative binomial distribution as shown below:

$$f\big(i|k,n,E(u)\big) = \frac{\Gamma(i+k)}{\Gamma(i+1)\Gamma(k)} k^k [nE(u)]^i [k+nE(u)]^{-k-i}$$

381   Eq. 6
383   The cumulation density function for Eq. 6 is then:

$$F(i \leq i^*) = \sum_{i=1}^{i^*} f\big(i|k,n,E(u)\big)$$

385   Eq. 7
386   Then, by definition, $A_{i\geq i^*}$ should be $\ll 1$ so that mutations with recurrences $i > i^*$ could be defined as
387   CDN. Thus:

$$F(i \leq i^*) = 1 - \frac{\varepsilon}{L_A}$$

389   Eq. 8

390   Where $\varepsilon = A_{i\geq i^*}$ denotes the number of sites with mutation recurrence $\geq i^*$ under the sole influence of

391   mutational force. $\frac{\varepsilon}{L_A}$ could then be regarded as significance of i^* since it controls the overall false

392   positive rate of CDNs.
393

394   Specifically, with $k = 1$, the probability function of mutation recurrence of a given site would transform
395   to a geometric distribution with $p = 1 / (1 + $**nE(u)**$)$, the cumulative density function (CDF) is then:

$$F(i \leq i^*) = 1 - \left(1 - \frac{1}{1+nE(u)}\right)^{i^*}$$

397   Eq. 9
398   Combined with Eq. 5, the mutation recurrence cutoff i^* of being a CDN could be expressed as:

$$i^* = \frac{log\left(\frac{\varepsilon}{L_A}\right)}{log\left(\frac{nE(u)}{1+nE(u)}\right)}$$

Eq. 10
For very large **n**, $1 / $**nE(u)** is small and $i^*/$**n** can be approximated as
$$i^*/n = log(L_A) \cdot E(u) \sim 5 \times 10^{-5}$$
Eq. 11
Eq. 11 shows that $i^*/$**n** would approach $log(L_A) \cdot E(u)$ asymptotically as **n** increases. This asymptotic
value is attained when n reaches $\sim 10^6$.
406

407   **Fig. 6**

Fig. 6 shows the range of i^* for **n** up to 10^6. As expected, i^* increases by small increments while **n** increases in 10-fold jumps. For example, when **n** increases by 3 orders of magnitude, from 100 to 100,000, i^* only doubles from 3 to 12. The disproportional increment between i^* and **n** explains why we use the actual number of i^* for the cutoff, instead of the ratio $i^*/\mathbf{n}$. As shown in the inset of **Fig. 2**, the ratio of $i^*/\mathbf{n}$ would approach the asymptote at $\mathbf{n}\sim10^6$, where an advantageous mutation only needs to rise to 0.00006 to be detected. With **n** reaching this level, we shall be able to separate most CDNs apart from the mutation background.

When **n** approaches 10^5, the number of CDNs will likely increase more than 10-fold as conjectured in **Fig. 7A**. In that case, every patient would have, say, 5 CDNs that can be subjected to gene targeting. (The companion study shows that, at present, an average cancer patient would have fewer than one targetable CDN.) Before the project is realized, it is nevertheless possible to test some aspects of it using the GENIE data of targeted sequencing. Such screening for mutations in the roughly 700 canonical genes serves the purpose of diagnosis with **n** ranging between 10~17 thousands for the breast, lung and CNS cancers. Clearly, GENIE efforts did not engage in discovering new mutations although they would discover additional CDNs in the canonical genes.

Fig. 7A assumes that the prevalence, $i/\mathbf{n}$, should not be much affected by **n**, but the cutoff for CDNs, $i^*/\mathbf{n}$, would decrease rapidly as **n** increases. **Fig. 7B** shows that the $i/\mathbf{n}$ ratios indeed correspond well between TCGA and GENIE, which differ by 10 to 20-fold in sample size. Importantly, as predicted in **Fig. 6**, the number of CDNs increases by 3 to 5-fold (**Fig. 7C - E**). Many of these newly discovered CDNs from GENIE are found in the A_1 and A_2 classes of TCGA while many more are found in the A_0 class in TCGA. In conclusion, increasing **n** by one to two orders of magnitude would be the simplest means of finding all CDNs.

DISCUSSION
The nature of high-recurrence mutations has been controversial. Many authors have argued for mutational hotspots (Hess et al. 2019; Stobbe et al. 2019; Nesta et al. 2021; Bergstrom et al. 2022; Wong et al. 2022) but just as many have contended that they are CDNs driven by selection (Gartner et al. 2013; Chang et al. 2016; Bailey et al. 2018; Cannataro et al. 2018; Juul et al. 2021; Zhao et al. 2021; Zeng and Bromberg 2022). While the two views co-exist, they are in fact incompatible. If the mutational hotspot hypothesis is correct, the selection hypothesis, and the determination of CDNs, would not be needed.

We believe that this study is the first to comprehensively test of the null hypothesis of mutational hotspots. In **PART I**, the mutational characteristics near all putative CDNs are examined and **PART II** presents the probability theory based on the analyses. The conclusion is that it is possible to reject the null hypothesis for recurrences as low as 3 in the TCGA data. The main reason for the high sensitivity is shown in Eqs. 2 and 3 where A_i and S_i is proportional to $[\mathbf{n}E(\mathbf{u})]^i$ or, roughly 0.002^i. We recognized that the conclusion is based on what we currently know about mutation mechanisms. In a sense, the theory developed here can help the search for such unknown mutation mechanisms, if they do exist.

Finally, the theory developed would permit the explorations in several new fronts when the sample size, **n**, expands to 10^5. The first front is to identify (nearly) all CDNs. When **n** reaches 10^5, any point mutation with a prevalence higher than 12/100,000 would be a CDN, which is 25-fold more sensitive

than in the TCGA data (3/1000). The companion analysis suggests that CDNs with lower prevalence, say 12/100,000 may still be highly tumorigenic in patients with the said mutation. If prevalence and potence are indeed poorly correlated, the search for lower prevalence CDNs by increasing $\mathbf{n}$ to 10^5 is equivalent to searching for less common but still potent cancer driving mutations.

The second one is functional tests in patient-derived cell lines. When we have all CDNs identified, a patient can be expected to have multiple (≥5; supplement File S3) CDNs. These mutations will be the basis of in vitro test, as well as in animal model experiments, by gene editing. As shown recently (Hodis et al. 2022), targeting multiple mutations simultaneously is necessary and may even be sufficient.

The third front, and arguably the most important one, is cancer treatment by targeted therapy. When multiple CDN mutations in the same patient can be simultaneously targeted, the efficacy should be high. No less crucial, resistance to treatment should be diminished since it would be harder on cancer cells to evolve multiple escape routes to evade multiple drugs.

There will be other fronts to explore with the full set of CDNs. For example, cancer screening can be more precise at the nucleotide level than at the gene level. A large database will also facilitate the detection of negative selection which has eluded detection. The failure to pinpoint deleterious mutations has led to the curious view that, in somatic evolution, deleterious mutations are rare. Chen et al. have addressed this curious phenomenon, which they term "quasi-neutral evolution" (Chen et al. 2019). In addition, it will be possible to study the evolution of mutation mechanisms in cancer cells based on such large samples (Jackson and Loeb 1998; Ruan et al. 2020). This last topic is addressed in the Supplement. As cancer genomics is increasingly adopted in cancer treatment, these benefits should become apparent when $\mathbf{n}$ reaches 10^5 for most cancer types.

References

Alexandrov LB, Kim J, Haradhvala NJ, Huang MN, Tian Ng AW, Wu Y, Boot A, Covington KR, Gordenin DA, Bergstrom EN, et al. 2020. The repertoire of mutational signatures in human cancer. *Nature* 578:94–101.

Alexandrov LB, Nik-Zainal S, Wedge DC, Aparicio SAJR, Behjati S, Biankin AV, Bignell GR, Bolli N, Borg A, Børresen-Dale A-L, et al. 2013. Signatures of mutational processes in human cancer. *Nature* 500:415–421.

Anandakrishnan R, Varghese RT, Kinney NA, Garner HR. 2019. Estimating the number of genetic mutations (hits) required for carcinogenesis based on the distribution of somatic mutations. *PLOS Computational Biology* 15:e1006881.

Armitage P, Doll R. 1954. The Age Distribution of Cancer and a Multi-stage Theory of Carcinogenesis. *Br J Cancer* 8:1–12.

Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, Colaprico A, Wendl MC, Kim J, Reardon B, et al. 2018. Comprehensive Characterization of Cancer Driver Genes and Mutations. *Cell* 173:371-385.e18.

Belikov AV. 2017. The number of key carcinogenic events can be predicted from cancer incidence. *Sci*
Rep 7:12170.

Bergstrom EN, Luebeck J, Petljak M, Khandekar A, Barnes M, Zhang T, Steele CD, Pillay N, Landi MT,
Bafna V, et al. 2022. Mapping clustered mutations in cancer reveals APOBEC3 mutagenesis of
ecDNA. *Nature* 602:510–517.

Black JRM, McGranahan N. 2021. Genetic and non-genetic clonal diversity in cancer evolution. *Nat Rev*
Cancer 21:379–392.

Blough MD, Beauchamp DC, Westgate MR, Kelly JJ, Cairncross JG. 2011. Effect of aberrant p53 function
on temozolomide sensitivity of glioma cell lines and brain tumor initiating cells from
glioblastoma. *J Neurooncol* 102:1–7.

de Bruijn I, Kundra R, Mastrogiacomo B, Tran TN, Sikina L, Mazor T, Li X, Ochoa A, Zhao G, Lai B, et al.
2023. Analysis and Visualization of Longitudinal Genomic and Clinical Data from the AACR
Project GENIE Biopharma Collaborative in cBioPortal. *Cancer Res* 83:3861–3867.

Buisson R, Langenbucher A, Bowen D, Kwan EE, Benes CH, Zou L, Lawrence MS. 2019. Passenger
hotspot mutations in cancer driven by APOBEC3A and mesoscale genomic features. *Science*
364:eaaw2872.

Burns MB, Temiz NA, Harris RS. 2013. Evidence for APOBEC3B mutagenesis in multiple human
cancers. *Nat Genet* 45:977–983.

Campbell PJ, Getz G, Korbel JO, Stuart JM, Jennings JL, Stein LD, Perry MD, Nahal-Bose HK, Ouellette
BFF, Li CH, et al. 2020. Pan-cancer analysis of whole genomes. *Nature* 578:82–93.

Cannataro VL, Gaffney SG, Townsend JP. 2018. Effect Sizes of Somatic Mutations in Cancer. *JNCI: Journal*
of the National Cancer Institute 110:1171–1177.

Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E,
et al. 2012. The cBio cancer genomics portal: an open platform for exploring multidimensional
cancer genomics data. *Cancer Discov* 2:401–404.

Chang MT, Asthana S, Gao SP, Lee BH, Chapman JS, Kandoth C, Gao J, Socci ND, Solit DB, Olshen AB, et
al. 2016. Identifying recurrent mutations in cancer reveals widespread lineage diversity and
mutational specificity. *Nat Biotechnol* 34:155–163.

Chen B, Shi Z, Chen Q, Shen X, Shibata D, Wen H, Wu C-I. 2019. Tumorigenesis as the Paradigm of
Quasi-neutral Molecular Evolution. *Mol Biol Evol* 36:1430–1441.

Elliott K, Larsson E. 2021. Non-coding driver mutations in human cancer. *Nat Rev Cancer* 21:500–509.

Fang Y, Deng S, Li C. 2022. A generalizable deep learning framework for inferring fine-scale germline
mutation rate maps. *Nat Mach Intell* 4:1209–1223.

Gartner JJ, Parker SCJ, Prickett TD, Dutton-Regester K, Stitzel ML, Lin JC, Davis S, Simhadri VL, Jha S,
Katagiri N, et al. 2013. Whole-genome sequencing identifies a recurrent functional synonymous
mutation in melanoma. *Proceedings of the National Academy of Sciences* 110:13481–13486.

Gojobori T, Li W-H, Graur D. 1982. Patterns of nucleotide substitution in pseudogenes and functional
genes. *J Mol Evol* 18:360–369.

Hanahan D. 2022. Hallmarks of Cancer: New Dimensions. *Cancer Discovery* 12:31–46.

Hanahan D, Weinberg RA. 2000. The Hallmarks of Cancer. *Cell* 100:57–70.

Hanahan D, Weinberg RA. 2011. Hallmarks of Cancer: The Next Generation. *Cell* 144:646–674.

Haradhvala NJ, Kim J, Maruvka YE, Polak P, Rosebrock D, Livitz D, Hess JM, Leshchiner I, Kamburov A,
Mouw KW, et al. 2018. Distinct mutational signatures characterize concurrent loss of
polymerase proofreading and mismatch repair. *Nat Commun* 9:1746.

Hartl DL, Clark AG. 1989. Principles of population genetics. Sunderland, Mass: Sinauer

He Z, Xu S, Shi S. 2020. Adaptive convergence at the genomic level—prevalent, uncommon or very
rare? *National Science Review* 7:947–951.

He Z, Xu S, Zhang Z, Guo W, Lyu H, Zhong C, Boufford DE, Duke NC, The International Mangrove
Consortium, Shi S. 2020. Convergent adaptation of the genomes of woody plants at the land–
sea interface. *National Science Review* 7:978–993.

Herzog M, Alonso-Perez E, Salguero I, Warringer J, Adams DJ, Jackson SP, Puddu F. 2021. Mutagenic
mechanisms of cancer-associated DNA polymerase ε alleles. *Nucleic Acids Research* 49:3919–
3931.

Hess JM, Bernards A, Kim J, Miller M, Taylor-Weiner A, Haradhvala NJ, Lawrence MS, Getz G. 2019.
Passenger Hotspot Mutations in Cancer. *Cancer Cell* 36:288-301.e14.

Hodgkinson A, Eyre-Walker A. 2011. Variation in the mutation rate across mammalian genomes. *Nat
Rev Genet* 12:756–766.

Hodis E, Triglia ET, Kwon JYH, Biancalani T, Zakka LR, Parkar S, Hütter J-C, Buffoni L, Delorey TM,
Phillips D, et al. 2022. Stepwise-edited, human melanoma models reveal mutations' effect on
tumor and microenvironment. *Science* 376:eabi8175.

Jackson AL, Loeb LA. 1998. The Mutation Rate and Cancer. *Genetics* 148:1483–1490.

Juul RI, Nielsen MM, Juul M, Feuerbach L, Pedersen JS. 2021. The landscape and driver potential of site-
specific hotspots across cancer genomes. *npj Genom. Med.* 6:1–15.

Kim J, Mouw KW, Polak P, Braunstein LZ, Kamburov A, Tiao G, Kwiatkowski DJ, Rosenberg JE, Van Allen
EM, D'Andrea AD, et al. 2016. Somatic ERCC2 mutations are associated with a distinct genomic
signature in urothelial tumors. *Nat Genet* 48:600–606.

Li M, Chen Wei, Sun X, Wang Zhipeng, Zou X, Wei H, Wang Zhan, Chen Wansheng. 2019. Metastatic
colorectal cancer and severe hypocalcemia following irinotecan administration in a patient
with X-linked agammaglobulinemia: a case report. *BMC Med Genet* 20:157.

Li W-H ,. 1997. Molecular evolution. Sunderland, Mass: Sinauer Associates

Lin L, Cai J, Tan Z, Meng X, Li R, Li Y, Jiang C. 2021. Mutant IDH1 Enhances Temozolomide Sensitivity via Regulation of the ATM/CHK2 Pathway in Glioma. *Cancer Res Treat* 53:367–377.

Ling S, Hu Z, Yang Z, Yang F, Li Y, Lin P, Chen K, Dong L, Cao L, Tao Y, et al. 2015. Extremely high genetic diversity in a single tumor points to prevalence of non-Darwinian cell evolution. *Proc Natl Acad Sci USA* 112:E6496–E6505.

Luo P, Ding Y, Lei X, Wu F-X. 2019. deepDriver: Predicting Cancer Driver Genes Based on Somatic Mutations Using Deep Convolutional Neural Networks. *Frontiers in Genetics* [Internet] 10. Available from: https://www.frontiersin.org/articles/10.3389/fgene.2019.00013

Makova KD, Hardison RC. 2015. The effects of chromatin organization on variation in mutation rates in the genome. *Nat Rev Genet* 16:213–223.

Marques DA, Meier JI, Seehausen O. 2019. A Combinatorial View on Speciation and Adaptive Radiation. *Trends in Ecology & Evolution* 34:531–544.

Martincorena I, Raine KM, Gerstung M, Dawson KJ, Haase K, Van Loo P, Davies H, Stratton MR, Campbell PJ. 2017. Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* 171:1029-1041.e21.

Nesta AV, Tafur D, Beck CR. 2021. Hotspots of Human Mutation. *Trends in Genetics* 37:717–729.

Noeuveglise A, Sarafan-Vasseur N, Beaussire L, Marguet F, Modzelewski R, Hanzen C, Alexandru C, Magne N, Langlois O, Di Fiore F, et al. 2023. Impact of EGFRA289T/V mutation on relapse pattern in glioblastoma. *ESMO Open* 8:100740.

Ortmann CA, Kent DG, Nangalia J, Silber Y, Wedge DC, Grinfeld J, Baxter EJ, Massie CE, Papaemmanuil E, Menon S, et al. 2015. Effect of Mutation Order on Myeloproliferative Neoplasms. *N Engl J Med* 372:601–612.

Pleasance ED, Cheetham RK, Stephens PJ, McBride DJ, Humphray SJ, Greenman CD, Varela I, Lin M-L, Ordóñez GR, Bignell GR, et al. 2010. A comprehensive catalogue of somatic mutations from a human cancer genome. *Nature* 463:191–196.

Polak P, Karlić R, Koren A, Thurman R, Sandstrom R, Lawrence MS, Reynolds A, Rynes E, Vlahoviček K, Stamatoyannopoulos JA, et al. 2015. Cell-of-origin chromatin organization shapes the mutational landscape of cancer. *Nature* 518:360–364.

Roberts SA, Lawrence MS, Klimczak LJ, Grimm SA, Fargo D, Stojanov P, Kiezun A, Kryukov GV, Carter SL, Saksena G, et al. 2013. An APOBEC cytidine deaminase mutagenesis pattern is widespread in human cancers. *Nat Genet* 45:970–976.

Ruan Y, Hou M, Tang X, He X, Lu X, Lu J, Wu C-I, Wen H. 2022. The Runaway Evolution of SARS-CoV-2 Leading to the Highly Evolved Delta Strain. *Molecular Biology and Evolution* 39:msac046.

Ruan Y, Wang H, Chen B, Wen H, Wu C-I. 2020. Mutations Beget More Mutations—Rapid Evolution of Mutation Rate in Response to the Risk of Runaway Accumulation. *Mol Biol Evol* 37:1007–1019.

Ruan Y, Wen H, Hou M, He Z, Lu X, Xue Y, He X, Zhang Y-P, Wu C-I. 2022. The twin-beginnings of COVID-19 in Asia and Europe—one prevails quickly. *National Science Review* 9:nwab223.

Ségurel L, Wyman MJ, Przeworski M. 2014. Determinants of Mutation Rate Variation in the Human Germline. *Annual Review of Genomics and Human Genetics* 15:47–70.

Sherman MA, Yaari AU, Priebe O, Dietlein F, Loh P-R, Berger B. 2022. Genome-wide mapping of somatic mutation rates uncovers drivers of cancer. *Nat Biotechnol*:1–10.

Song Q, Li M, Li Q, Lu Xun, Song K, Zhang Z, Wei Jiale, Zhang L, Wei Jiacheng, Ye Y, et al. 2023. DeepAlloDriver: a deep learning-based strategy to predict cancer driver mutations. *Nucleic Acids Research* 51:W129–W133.

Stamatoyannopoulos JA, Adzhubei I, Thurman RE, Kryukov GV, Mirkin SM, Sunyaev SR. 2009. Human mutation rate associated with DNA replication timing. *Nat Genet* 41:393–395.

Stobbe MD, Thun GA, Diéguez-Docampo A, Oliva M, Whalley JP, Raineri E, Gut IG. 2019. Recurrent somatic mutations reveal new insights into consequences of mutagenic processes in cancer. *PLOS Computational Biology* 15:e1007496.

Stratton MR, Campbell PJ, Futreal PA. 2009. The cancer genome. *Nature* 458:719–724.

Supek F, Lehner B. 2015. Differential DNA mismatch repair underlies mutation rate variation across the human genome. *Nature* 521:81–84.

Takeda H. 2021. A Platform for Validating Colorectal Cancer Driver Genes Using Mouse Organoids. *Front Genet* 12:698771.

Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, Boutselakis H, Cole CG, Creatore C, Dawson E, et al. 2019. COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Research* 47:D941–D947.

Turajlic S, Sottoriva A, Graham T, Swanton C. 2019. Resolving genetic heterogeneity in cancer. *Nat Rev Genet* 20:404–416.

Wang Q, Fang W-H, Krupinski J, Kumar S, Slevin M, Kumar P. 2008. Pax genes in embryogenesis and oncogenesis. *Journal of Cellular and Molecular Medicine* 12:2281–2294.

Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, Shmulevich I, Sander C, Stuart JM. 2013. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet* 45:1113–1120.

Wong JKL, Aichmüller C, Schulze M, Hlevnjak M, Elgaafary S, Lichter P, Zapatka M. 2022. Association of mutation signature effectuating processes with mutation hotspots in driver genes and non-coding regions. *Nat Commun* 13:178.

Wu C-I, Maeda N. 1987. Inequality in mutation rates of the two strands of DNA. *Nature* 327:169–170.

Wu C-I, Ting C-T. 2004. Genes and speciation. *Nat Rev Genet* 5:114–122.

Wu C-I, Wang G-D, Xu S. 2020. Convergent adaptive evolution—how common, or how rare? *National*
Science Review 7:945–946.

Yates LR, Campbell PJ. 2012. Evolution of the cancer genome. *Nat Rev Genet* 13:795–806.

Zeng Z, Bromberg Y. 2022. Inferring Potential Cancer Driving Synonymous Variants. *Genes* 13:778.

Zhao W, Yang J, Wu Jingcheng, Cai G, Zhang Y, Haltom J, Su W, Dong MJ, Chen S, Wu Jian, et al. 2021.
CanDriS: posterior profiling of cancer-driving sites based on two-component evolutionary
model. *Briefings in Bioinformatics* 22:bbab131.

Zheng CL, Wang NJ, Chung J, Moslehi H, Sanborn JZ, Hur JS, Collisson EA, Vemula SS, Naujokas A, Chiotti
KE, et al. 2014. Transcription Restores DNA Repair to Heterochromatin, Determining Regional
Mutation Rates in Cancer Genomes. *Cell Reports* 9:1228–1234.

646
Methods
Data Collection
Single nucleotide variation (SNV) data for the TCGA cohort was downloaded from the GDC Data Portal
(https://portal.gdc.cancer.gov/). Mutations were further filtered based on their population frequency
recorded in the Genome Aggregation Database (*gnomAD*), with an upper threshold of 1‰. We focused
on coding region mutations of missense, nonsense, and synonymous types on autosomes. The
mutation load for each patient was defined as the sum of these three types of mutations. Patients with
a mutation load exceeding 3000 were identified as having a mutator phenotype and were excluded
from our analysis. In total, 7369 samples representing 12 cancer types were included for mutation
analysis.
657
Additional mutation data was acquired from the AACR Project GENIE Consortium via cBioPortal. Due
to the prevalence of targeted sequencing within the dataset, filtering was implemented to ensure the
inclusion of samples with sequencing assays encompassing all exonic regions of target genes.
Furthermore, sample-level filtering was performed to guarantee a unique sequencing sample per
patient. Germline filtering was applied to the resulting point mutations, removing mutations with SNP
frequencies exceeding 0.0005 in any subpopulation annotated by gnomAD. To exclude patients
exhibiting hypermutator phenotypes, mutation loads were scaled to the whole exon level with $\widetilde{n_L} =$
$n_l \cdot \frac{L}{l}$, where n_l and l represent the mutation load and genomic length of target sequencing region,
respectively. L denotes the genome-wide whole coding region length. Patients with $\widetilde{n_L} > 3000$ were
subsequently excluded, consistent with the threshold employed for the TCGA dataset.
668
Calculation of missense and synonymous site number (L_A **and** L_S)
The idea of missense or synonymous sites originates from the question that how many missense or
synonymous mutations would be expected if each site of the genome were mutated once given
background mutation patterns. Here, the background mutation patterns refer to intrinsic biases in the
mutational process, such as the over-representation of C>T (or G>A) mutations at CpG sites due to
spontaneous deamination of 5-methylcytosine. In coding regions, four-fold degenerate sites are
generally considered neutral, as any mutation path would not alter the encoded amino acid sequence.
This analysis follows established methods at the single-base level to infer the expected number of

missense and synonymous sites across the genome (Gojobori et al. 1982; Wu and Maeda 1987; Hartl
and Clark 1989; Martincorena et al. 2017).
679

To illustrate the calculation process, we provide an example of synonymous site estimation. At four-
fold degenerate sites throughout the genome, we tally the number of mutations from base m to base v
as $n_{m>v}$ ($m, v \in \{A, C, G, T\}$, $m \neq v$). The likelihood of observing a mutation from reference base m to
variant base v will be $r_{m>v} = \frac{n_{m>v}}{N_m}$, where N_m represents the number of four-fold degenerate site
with reference base m. We then normalize all likelihoods by $R_{m>v} = \frac{r_{m>v}}{\sum r}$, where $\sum r$ represents the
sum of likelihoods across 12 possible mutation paths, $R_{m>v}$ thus describes the relative probability of
an occurred mutation to be $m > v$ at any site. For a given genomic coding region of length L, the
synonymous site number will be:

688
$$L_S = \sum_L \delta_{m>v}^{syn} R_{m>v}$$

689   Eq. S1
690   with $\delta_{m>v}^{syn}$ being a Kronecker delta function where:
691

692
$$\delta_{m>v}^{syn} = 1 \text{ if } m > v \text{ is synonymous}$$

694
$$\delta_{m>v}^{syn} = 0 \text{ otherwise}$$

696   Similarly, the expected number of missense sites L_A, is calculated as follows:

697
$$L_A = \sum_L \delta_{m>v}^{mis} R_{m>v}$$

698   Eq. S2
699

Calculation of A_i and S_i
For each mutated site, we track its number of recurrent mutations i ($i > 0$) across two mutation
categories: missense and synonymous. Subsequently, we aggregate across entire coding region to
count the number of sites harboring i missense mutations (A_i) and synonymous mutations (S_i). For $i =$
0, we define A_0 and S_0 as the estimated number of potential missense and synonymous sites that
remain unmutated within the current sample size. $A_0 = L_A - \sum_{i>0} A_i$, $S_0 = L_S - \sum_{i>0} S_i$.
706

AI-based mutation rate analysis
To capture the complex interplay of genomic and epigenomic factors influencing mutation
susceptibility, we employed pre-trained artificial intelligence (AI) models from *Dig*, an aggregated tool
combining deep learning and probabilistic models (Sherman et al. 2022). Downloaded from the *Dig*
data portal (http://cb.csail.mit.edu/cb/DIG/downloads/), these models leverage a rich set of features
encompassing both kilobase-scale epigenomic context (replicating timing, chromatin accessibility,
etc.) and fine-grained base-pair level information (such as sequence context biases) to predict the site
level mutation rate. For each cancer type, we re-fitted the pre-trained models with mutations analyzed
in our study. The mutation rate for each site, scaled by population size, was obtained via the
elementDriver function within Dig, and was represented by *EXP_SNV* from the final results. For a closer
look at mutation rate landscape of *PAX3*, we re-fitted the AI-model with point mutations from large
intestine cancer. The mutation rates were generated site-by-site for the coding regions of *PAX3*.
719

Given the scarcity of mutated sites with recurrence $i \geq 3$ (comprising only 0.15% of all mutated sites),
a rigorous statistical approach was adopted to assess the significance of mutability differences
between these high-hit groups and low-hit groups. We implemented the procedure as follows:
1). Raw significance level: For each recurrence group i (containing A_i sites), a one-sided Kolmogorov-
Smirnov (K-S) test was employed to calculate a raw significance level (denoted as p_0) against the low-
hit group.
2). Resampling for Significance Pool: we resample A_i sites from the entire pool of mutated sites with
missense mutations. The significance p_j from one-sided K-S test is calculated against the low hit
group. The resampling process was repeated 100,000 times, generating a distribution of resampled
significance levels, denoted as $\{p_j, j = 1,2,\ldots,100{,}000\}$.

3). Adjusted Significance Level: the raw significance p_0 was then compared to the resampled
significance pool $\{p_j, j = 1,2,\ldots,100{,}000\}$. The proportion of $p_0 \leq p_j$ was then calculated as the
adjusted significance level that accounted for potential sampling effects.
733
Motif based mutability
For a given nucleotide base, we extended the sequence to each side by 1, 2, and 3 base pairs, producing
sets of 3-mer, 5-mer, and 7-mer motifs, respectively. We then pooled point mutations from 12 cancer
types to create a comprehensive dataset. For 3-mer and 5-mer motifs, we utilized synonymous
mutations as the reference for mutation rate calculations. For 7-mer motifs, the vast number of
possible sequence combinations ($4^7 = 16{,}384$) posed a challenge, as synonymous mutations alone
might not adequately cover all potential contexts. To address this, we employed all singleton
mutations (1-hit mutations) from both missense and synonymous categories for 7-mer motif analysis.
This decision was based on the assumption that singleton mutations are less affected by selective
pressures, supported by the genome-wide observation that the ratio of missense to synonymous
singletons (A_1/S_1) approximates the ratio of unmutated missense to synonymous sites (A_0/S_0).
The site number for 3-mer and 5-mer motifs of a given context c was calculated as follows:

746
$$L_{c,S} = \sum_L \delta_{c,m>v}^{syn} \cdot R_{m>v}$$

747   Which is an extension of Eq. S1, with $\delta_{c,m>v}^{syn}$ being a Kronecker delta function where:

748        $\delta_{c,m>v}^{syn} = 1$ if base change of m to v ($m > v$) is synonymous under sequence context c.

749        $\delta_{c,m>v}^{syn} = 0$ otherwise.

750   The mutation rate then could be expressed as:

751
$$\mu_c = \frac{n_{c,m>v}^{syn}}{L_{c,S}}$$

752   Eq. S3
753   for 7-mer motifs, the calculation is:

754
$$L_c = L_{c,S} + L_{c,A} = \sum_L \delta_{c,m>v}^{syn} \cdot R_{m>v} + \sum_L \delta_{c,m>v}^{mis} \cdot R_{m>v}$$

755
$$\mu_c = \frac{n_{c,m>v}^{syn} + n_{c,m>v}^{mis}}{L_c}$$

756   Eq. S4

757 Where for Eqs. S3 and S4, $n_{c,m>v}^{syn}$ and $n_{c,m>v}^{mis}$ represent the mutation numbers with **$m > v$** being

758 synonymous and missense under sequence context c, respectively. The mutation rate is then scaled as
759 the expected mutation number per 10^5 corresponding sequence motifs for better presentation.
760

761 Significance for motif enrichment (**Fig. 4B**) mirrored the AI analysis. For each $i \geq 3$ site, we calculated
762 raw K-S p-values against motif mutabilities (denoted as p_0). These were then compared to a

763 resampled significance pool $\{p_j, j = 1,2,\dots,100,000\}$, with the proportion of $p_0 \leq p_j$ employed as the

764 final p-value, depicting enrichment significance for highly mutable motifs in recurrence group i against
765 low hits.
766

Consensus length comparison
To explore potential sequence motifs associated with recurrent mutations ($i \geq 3$), we employ a sliding
window of 10 bp stride to extract the local context from reference genome (**Fig. S5**). We examined
diverse window sizes (21, 41, 61, 81, and 101 bp) to capture potential motifs of varying lengths and
distances to the mutated site. Consensus length of local contexts was measured by Hamming distance
in pairwise comparisons of aligned windows (with same stride) between mutated sites.
773

To prioritize sequence similarities likely driven by mutational mechanisms rather than functional
constraints or gene structure, we restricted consensus comparisons to non-homologous genes. This
approach effectively mitigated potential biases arising from homologous genes (e.g., *KRAS* and *NRAS*)
or repeated domains within a single gene (e.g., *FBXW7*). The statistical significance of observed
consensus lengths was assessed using the K-S test, which compared the empirical distribution of
consensus lengths against a Poisson distribution, with mean of λ set to one-quarter of the window
size, which reflects the expected distribution under random scenarios.
781

Mutational signature analysis
The mutation load of each patient could be further decomposed to several known mutational
processes, which is represented by mutational signatures. In general, each mutational signature
embodies the relative mutabilities across distinct mutational contexts. Leveraging single base
substitution (SBS) signatures from COSMIC (v3.3), we employed the *SignatureAnalyzer* tool to quantify
the contribution of each signature to individual mutational loads (Kim et al. 2016). For composition
analysis in **Fig. 5B**, we focused on signatures contributing at least 2% to the total mutations within a
given cancer type, given that there are 79 mutational signatures in use for deconvolution.
790

791 To assess signature contribution changes across recurrence cutoffs (i), we grouped patients with
792 mutations of recurrence $i \geq i^*$ and scaled signature contributions to 1 to cancel out the population size
793 effect. Pairwise K-S test between different i^*s is employed to determine whether signature
794 contributions are significantly different under each i^*.
795

796 **Outlier model**
797 The purpose of the outlier model is to investigate if high-hit sites could be explained by a fraction (p)
798 of highly mutable sites (with mutability α-fold higher). By definition, we have:

$$S_i = (1 - p) \cdot L_S \cdot [nE(u)]^i + p \cdot L_S \cdot [\alpha \cdot nE(u)]^i$$

Eq. S5
With **n** represents the population size and **E(u)** denotes the average mutation rate per site per patient.

We let p range from 10^{-5} to 10^{-2} and α from 1.1 to 1000. For each (p, α) pair, we solve Eq. S5 based on
observed S_1 ($i = 1$) to obtain $E(u)$. Then, we calculate expected S_i with $i = 2, 3$. The (p, α) pairs are
filtered by imposing constraints grounded in observed S_2 and S_3 values. Specifically, we retained only
those pairs whose expected S_2 and S_3 values resided within the 95% quantile range of a Poisson
distribution with λ set to the observed values. This filtering process yielded biologically plausible (p,
α) pairs that were then used to derive S_4 and S_5. Finally, we computed the mean and standard
deviation for p, α, S_4, and S_5 across all filtered pairs to capture their central tendencies and variability.

CDN analysis in GENIE
To circumvent potential biases in $E(u)$ estimation stemming from the varying target gene coverage
across sequencing panels within the GENIE dataset, we leveraged $E(u)$ values derived from the
corresponding cancer types in the TCGA dataset. Specifically, we focused on 1-hit synonymous
mutations within coding regions, as these are generally considered to be the least influenced by
selective pressures in coding regions. Based on Eq. 1 from the main text, we have:

$$S_1 = L_S \cdot nE(u)e^{-(n-1)E(u)}$$

Eq. S6

Where $e^{-(n-1)E(u)}$ comes from approximation of $[1 - E(u)]^{n-1}$ from binomial distribution, and $\mathbf{n}$
is the population size in TCGA. The calculation for the threshold i^*, based on Eq. 10 from the main
text, is:

$$i^* = \frac{log\left(\frac{\varepsilon}{L_A}\right)}{log\left(\frac{n_e E(u)}{1 + n_e E(u)}\right)}$$

Eq. S7

The only difference in Eq. S7 is that we use $\mathbf{n_e}$ to represent the number of patients sequenced for a
target gene in GENIE, considering the overlapping between different assays in use. In essence, we will
have for each gene a CDN threshold i^*.

The comparison of CDNs between TCGA and GENIE are restricted to genes sequenced by GENIE
panels. For CDNs identified in GENIE using Eq. S7, we investigated their hit information and CDN
identity within TCGA dataset. The CDN flow proportion depicted in **Fig. 7 C-E** represents the ratio of
sites identified as CDN in GENIE to A_i^* ($i = 0, 1, 2$) of TCGA. A_i^* mirrors A_i but specifically considers
the coding region sequenced by the GENIE panel. For CDNs identified in TCGA, the flow ratio is just the
proportion of sites being identified as CDNs in GENIE. Notably, for CDNs identified in GENIE but
lacking mutations in TCGA ($i = 0$), A_0^* is obtained using Eq. S2 with L being the length of coding region
sequenced in GENIE.

ACKNOWLEDGEMENTS
The authors gratefully acknowledge the following for their support in the initiation of the Cancer
Driving Nucleotide (CDN) project: the First Affiliated Hospital, the Seventh Affiliated Hospital of Sun
Yat-sen University; Cancer Center of Clifford Hospital, Jinan University; Cancer Hospital Chinese
Academy of Medical Sciences, Shenzhen Center; Guangdong Academy of Medical Sciences, Guangdong
Provincial People's Hospital. We thank the Kunming Institute of Zoology for valuable discussions on
the CDN concept. We are also grateful to Weiwei Zhai, Qianfei Wang, and Weini Huang for their
insightful comments and suggestions. Finally, we acknowledge the American Association for Cancer
Research (AACR) and The Cancer Genome Atlas (TCGA) project for providing invaluable datasets and

resources that have significantly advanced our understanding of cancer biology and improved patient
outcomes. This work was supported by the National Natural Science Foundation of China
(32293193/32293190 and 32150006 to C.I.W.), the National Key Research and Development Projects
of the Ministry of Science and Technology of China (2021YFC2301300), National Key R&D Program of
China (2021YFC0863400), and the Innovation Group Project of Southern Marine Science and
Engineering Guangdong Laboratory (Zhuhai; No. 311021006).
852

853 **DECLARATION OF INTERESTS**
854 The authors declare no competing interests.

TABLES AND FIGURES

856   **Table 1**: An example of A_i and S_i (from lung cancer, n = 1035).

i	All sites			CpG sites removed		
	A_i	S_i	A_i / S_i	A_i	S_i	A_i / S_i
0	22540623	7804281	2.89	21375384	7014012	3.04
1	195958	69393	2.82	168371	56821	2.96
2	2946	969	3.04	2188	643	3.4
3	99	21	4.71	68	16	4.25
4	23	1	23	17	1	17
5	16	0	16 : 0	9	0	9 : 0
6	10	0	10 : 0	6	0	6 : 0
7	5	0	5 : 0	5	0	5 : 0
8	8	0	8 : 0	6	0	6 : 0
9	4	0	4 : 0	3	0	3 : 0
≥3	178	22	8.09	122	17	7.18
≥4	79	1	79	54	1	54
[10-20]	7	1	7	4	0	4 : 0
≥20	6	0	6 : 0	4	0	4 : 0

857   Note – The ratio of A_i / S_i is provided as a measure of selection strength.

860 **Table 2**: Summary for modeling outlier sites in 6 cancer types

Cancer Type	S_3	p	α	S_4	S_5
Lung*	--	*0.0*	--	--	--
breast	0.12	8.75E-04 (8.21E-04)	88.6 (32.0)	0.102 (0.068)	0.004 (0.004)
CNS	0.02	2.73E-04 (1.09E-04)	295.1 (57.0)	0.448 (0.173)	0.026 (0.015)
kidney	0.03	3.03E-05 (2.98E-05)	304.1 (108.0)	0.067 (0.056)	0.005 (0.006)
Upper-AD tract	0.47	0.002 (0.001)	48.9 (10.7)	0.174 (0.078)	0.005 (0.003)
Large intestine	1.03	0.009 (0.001)	51.6 (1.4)	0.998 (0.087)	0.026 (0.003)

861 Note – For each cancer type, p stands for the proportion of highly mutable sites, with mutation rate
862 being α-fold of the average. S_3 gives the expected number without mutable outliers ($p = 0$). S_4 and S_5
863 denote the expected number with the best (p, α) pairs with the standard deviation in parentheses. For
864 lung cancer, S_2 and S_3 do not fit the outlier model (supplement File S1); therefore, we set $p = 0$.
865
866
867

Figure 1: Two modes of DNA sequence evolution. **(A)** A hypothetical example of DNA sequences in organismal evolution. **(B)** Cancer evolution that experiences the same number of mutations as in **(A)** but with many short branches. **(C)** A common pattern of sequence variation in cancer evolution. **(D)** In cancer evolution, the same mutation at the same site may occasionally be seen in multiple sequences. The recurrent sites could be either mutational or functional hotspots, their distinction being the main objective of this study.

878
879 **Figure 2**: The average A_i and S_i values across different i ranges (X-axis). **Top**: The average of A_i and S_i
880 in the log scale. Color lines - full data; gray lines - CpG sites removed. The dash lines are linear
881 extrapolations. **Bottom: The A_i / S_i** ratio as a function of i.
882
883

884   A)

886   B)

Figure 3: Site-level mutation rate variation obtained from *Dig* (Sherman et al. 2022), a published AI
tool. **(A)** Each dot represents the expected SNVs (Y-axis) at a site where missense mutations occurred *i*
times in the corresponding cancer population. The boxplot shows the overall distribution of mutability
at *i*, with the red dashed line denoting the average. There is no observable trend that sites of higher *i*
are more mutable. **(B)** A detailed look at the coding region of *PAX3* gene in colon cancer. The expected
mutability of sites in the 200 bp window is plotted. The three mutated sites in this window, marked by
green and red (a CDN site) stars, are not particularly mutable. Overall, the mutation rate varies by
about 10-fold as is generally known for CpG sites.

896
897

Figure 4: Conventional analyses of local contexts at recurrence sites. **(A)** From top panel down - For the 64 (4^3) 3-mer motifs, their mutational rates are shown on the X-axis. The most mutable motif over the average mutability (α) is 4.69. For the 1024 (=4^5) 5-mer and 16,384 (4^7) 7-mer motifs, the α values are, respectively, 8.79 and 11.52. The most mutable motifs, as expected, are dominated by CpG's. **(B)** Each dot represents the motif surrounding a high-recurrence site. The recurrence number is shown on the X-axis and the mutability of the associated motif's mutability (mutations per 0.1M) is shown on the Y-axis. The average mutation rate across all motifs of given length category is indicated by a red horizontal dashed line. The absence of a trend indicates that the high recurrence sites are not associated with the mutability of the motif. **(C)** The analysis is extended to longer motifs surrounding each CDN (21, 41, 61, 81, and 101 bp). For each length group, all pairwise comparisons are enumerated. The observed distributions (black bars and points) are compared to the expected Poisson distributions (red bars and curves) and no difference is observed. Thus, local sequences of CDNs do not show higher-than-expected similarity.

A)

B)

Figure 5: Patient level analysis for mutation load and mutational signatures. **(A)** Boxplot depicting the
distribution of mutation load among patients with recurrent mutations. The X-axis denotes the count
of recurrent mutations, while the Y-axis depicts the normalized z-score of mutation load (see
Methods). The green dashed line indicates the mean mutation load. In short, the mutation load does
not influence the mutation recurrence among patients. **(B)** Signature analysis in patients with
mutations of recurrences ≥ i^* (X-axis). For lung cancer (left), the upper panel presents the number of
patients for each group, while the lower panel depicts the relative contribution of mutational
signatures. For breast cancer, APOBEC-related signatures (SBS2 and SBS13) are notably elevated in all
groups of patients with i^* ≥ 3, while patients with mutations of recurrence ≥ 20 in CNS cancer exhibit
an increased exposure to SBS11 (Blough et al. 2011; Lin et al. 2021; Noeuveglise et al. 2023). Again,
patients with higher mutation recurrences do not differ in their mutation signatures.

Figure 6: The Y axis presents the i^* values under different sample sizes (**n** of the X-axis) in log scale.
Five shape parameters (k) of the gamma-Poisson model are used. In the literatures on the evolution of
mutation rate, k is usually greater than 1. The inset figure illustrates how $i^*/$ **n** (prevalence) would
decrease with increasing sample sizes. The prevalence would approach the asymptotic line of
$[log(L_A) \cdot E(u)]$ when **n** reaches 10^6. In short, more CDNs (those with lower prevalence) will be
discovered as **n** increases. Beyond **n** = 10^6, there will be no gain.
941
942

943
Figure 7: **(A)** The two vertical lines show the cutoffs of i^*/n at (3/1000) vs. (12/100,000). The Y axis
shows that the potential number of site would decrease with i^*/n, which is a function of selective
advantage. The area between the two cutoffs below the line represent the new CDNs to be discovered
when **n** reaches 100,000. The power of **n** = 100,000 is even larger if the distribution follows the blue
dashed line. **(B)** The prevalence (i/n) of sites is well correlated between datasets of different **n** (TCGA
with **n** < 1000 and GENIE with **n** generally 10-fold higher), as it should be. Sites are displayed by color.
"1-hit": CDNs identified in GENIE but remain in singleton in TCGA, "2-hit": CDNs identified in GENIE
but present in doubleton in TCGA. "CDN both": CDNs identified in both databases. **(C-E)** CDNs
discovered in GENIE (**n** > 9000) but absent in TCGA (**n** < 1000). The newly discovered CDNs may fall in
TCGA as 0 - 2 hit sites. The numbers in the middle column show the percentage of lower recurrence
(non-CDN) sites in TCGA that are detected as CDNs in the GENIE database, which has much larger **n**'s.