On the discovered Cancer Driving Nucleotides (CDNs) –Distributions across genes, cancer types and patients

Lingjie Zhang[1], Tong Deng[1], Zhongqi Liufu[1,2], Xiangnyu Chen[1], Shijie Wu[1], Xueyu Liu[1], Changhao Shi[1], Bingjie Chen[1,3], Zheng Hu[4], Qichun Cai[5], Chenli Liu[4], Mengfeng Li[6], Miles E. Tracy[1], Xuemei Lu[2], Chung-I Wu[1,7,*] and Haijun Wen[1,*]

[1] State Key Laboratory of Biocontrol, School of Life Sciences, Sun Yat-sen University, Guangzhou 510275, Guangdong, China.
[2] Center for Excellence in Animal Evolution and Genetics, The Chinese Academy of Sciences, Kunming 650223, China.
[3] GMU-GIBH Joint School of Life Sciences, Guangzhou Medical University, Guangzhou 511436, China.
[4] CAS Key Laboratory of Quantitative Engineering Biology, Shenzhen Institute of Synthetic Biology, Shenzhen Institute of Advanced Technology, Chinese Academy of Sciences, Shenzhen, China.
[5] Cancer Center, Clifford Hospital, Jinan University, Guangzhou 511495, China.
[6] Cancer Research Institute, School of Basic Medical Sciences, Southern Medical University, Guangzhou 510515, China.
[7] Department of Ecology and Evolution, University of Chicago, Chicago, Illinois 60637, USA.
* Corresponding authors: ciwu@uchicago.edu; wenhj5@mail.sysu.edu.cn.

ABSTRACT

A central goal of cancer genomics is to identify, in each patient, all the cancer driving mutations. Among them, point mutations are referred to as Cancer Driving Nucleotides (CDNs), which recur in cancers. The companion study shows that the probability of i recurrent hits in **n** patients would decrease exponentially with i; hence, any mutation with $i \geq 3$ hits in the TCGA database is a high-probability CDN. This study characterizes the 50~150 CDNs identifiable for each cancer type of TCGA (while anticipating 10 times more undiscovered ones) as follows: **i**) CDNs tend to code for amino acids of divergent chemical properties. **ii**) At the genic level, far more CDNs (>5-fold) fall on non-canonical than canonical cancer driving genes (CDGs). Most undiscovered CDNs are expected to be on unknown CDGs. **iii**) CDNs tend to be more widely shared among cancer types than canonical CDGs, mainly because of the higher resolution at the nucleotide than the whole-gene level. **iv**) Most important, among the 50~100 coding region mutations carried by a cancer patient, 5~8 CDNs are expected but only 0~2 CDNs have been identified at present. This low level of identification has hampered functional test and gene targeted therapy. We show that, by expanding the sample size to 10^5, most CDNs can be identified. Full CDN identification will then facilitate the design of patient-specific targeting against multiple CDN-harboring genes.

INTRODUCTION

Tumorigenesis in each patient is driven by mutations in the patient's genome. Hence, a central goal of cancer genomics is to identify *all* driving mutations in each patient. This task is particularly challenging because each driving mutation is present in only a small fraction of patients. As the number of driver mutations in each patient has been estimated to be >5 (Armitage and Doll 1954; Bozic et al. 2010; Hanahan and Weinberg 2011; Belikov 2017; Anandakrishnan et al. 2019), the total number of driver mutations summed over all patients must be quite high.

This study, together with the companion paper (Zhang et al. 2024), are based on one simple premise: In the massively repeated evolution of cancers, any advantageous cancer-driving mutation should recur frequently, say i times in **n** patients. The converse that non-recurrent mutations are not advantageous is part of the same premise. We focus on point mutations, referred to as Cancer Driving Nucleotides (CDNs), and formulate the maximum of i (denoted i^*) in **n** patients if mutations are not advantageous. For example, in the TCGA database with **n** generally in the range of $500 \sim 1000$, $i^* = 3$. Hence, any point mutation with $i \geq 3$ is a CDN. At present, a CDN would have a prevalence of 0.3% among cancer patients. If the sample size approaches 10^6, a CDN only needs to be prevalent at 5×10^{-5}, the theoretical limit (Zhang et al. 2024).

Although there are many other driver mutations (e.g., fusion genes, chromosomal aberrations, epigenetic changes, etc.), CDNs should be sufficiently numerous and quantifiable to lead to innovations in functional tests and treatment strategies. Given the current sample sizes of various databases (Cerami et al. 2012; Weinstein et al. 2013; Tate et al. 2019; de Bruijn et al. 2023), each cancer type has yielded $50 \sim 150$ CDNs while the CDNs to be discovered should be at least 10 times more numerous. The number of CDNs currently observed in each patient is $0 \sim 2$ for most cancer types. This low-level of discovery has limited functional studies and hampered treatment strategies.

While we are proposing the scale-up of sample size to discover most CDNs, we now characterize CDNs that have been discovered. The main issues are the distributions of CDNs among genes, across cancer types and, most important, among patients. In this context, cancer driver genes (CDGs) would be a generic term. We shall use "canonical CDGs" (or conventional CDGs) for the driver genes in the union set of three commonly used lists (Bailey et al. 2018; Sondka et al. 2018; Martínez-Jiménez et al. 2020). In parallel, CDN-harboring genes, referred to "CDN genes", constitute a new and expanded class of CDGs.

The first issue is that CDNs are not evenly distributed among genes. The canonical cancer drivers such as *TP53*, *KRAS* and *EGFR* tend to have many CDNs. However, the majority of CDNs, especially those yet-to-be-identified ones, may be rather evenly distributed with each gene harboring only $1 \sim 2$ CDNs. Hence, the number of genes with tumorigenic potential may be far larger than realized so far. The second issue is the distribution of CDNs and CDGs among cancer types. It is generally understood that the canonical CDGs are not widely shared among cancer types. However, much (but not all) of the presumed cancer-type specificity may be due to low statistical resolution at the genic level.

The third issue concerns the distribution of CDNs among patients. Clearly, the CDN load of a patient is crucial in diagnosis and treatment. However, the conventional diagnosis at the gene level may have two potential problems. One is that many CDNs do not fall in canonical CDGs as signals from one or two CDNs get diluted. Second, a canonical CDG, when mutated, may be mutated at a non-CDN site. In those patients, the said CDG does not drive tumorigenesis. We shall clarify the relationships between CDN mutations and genes that may or may not harbor them.

The characterizations of discovered CDNs are informative and offer a road map for expanding the CDN list. A complete CDN list for each cancer type will be most useful in functional test, diagnosis and treatment. A full list of mutations that drive the evolution of complex traits is at the center of evolutionary genetics. Such phenomena as complex human diseases (e.g., diabetes.) (Vujkovic et al.

2020; Lagou et al. 2023; Xue et al. 2023; Suzuki et al. 2024), the genetics of speciation (Q. Chen et al. 2022; Wang et al. 2022; Wu 2022) and the evolution of viruses in epidemics (Deng et al. 2022; Ruan et al. 2022; Cao et al. 2023; Ruan et al. 2023) are all prime examples in need of a full list). Thanks to their massively repeated evolution, cancers could be the first complex systems well resolved at the genic level.

RESULTS

In molecular evolution, a gene under positive selection is recognized by its elevated evolutionary rate (**Fig. 1A and 1C**). There have been numerous methods for determining the extent of rate elevation (Li et al. 1985; Nei and Gojobori 1986; Yang and Swanson 2002; Lawrence et al. 2013; Martincorena et al. 2017; Pan et al. 2022; Sherman et al. 2022; Wang et al. 2022; Ruan et al. 2023) and cancer evolution studies have adopted many of them. However, no model has been developed to take advantage of the massively repeated evolution of cancers (**Fig. 1B**), which happens in tens of millions of people at any time.

Fig. 1

In the whole-gene analysis, **Fig. 1C-E** are identical, each with $A : S = 10 : 1$ where A and S denote nonsynonymous and synonymous mutations, respectively. However, the presence of a 4-hit site in **Fig. 1E** is far less likely to be neutral than **Fig. 1C and 1D**. Although ratio in **Fig. 1F**, $A : S = 4 : 1$, is statistically indistinguishable from the neutral ratio of about 2.5 : 1, **Fig. 1F** in fact has much more power to reject the neutral ratio than **Fig. 1C** and **1D**. After all, the probability that multiple hits are at the same site in a big genome is obviously very small.

1. The analyses of CDNs across the whole genome

For the entire coding regions in the cancer genome data, we define A_i (or S_i) as the number of nonsynonymous (or synonymous) sites that harbor a mutation with i recurrences. **Table 1** presents the distribution of A_i and S_i across the 12 cancer types with $n > 300$ (Weinstein et al. 2013).

For neutral mutations, we define i^* as the threshold above which the expected numbers of A_i would be <1, i.e. $E[A_{i \geq i^*}] < 1$, The corollary is that all $A_{i \geq i^*}$ sites are advantageous CDNs. (Since S_i is $\sim A_i / 2.3$, the same i^* would apply to S_i as well: $E[S_{i \geq i^*}] < 1$) As i^* is a function of the number of patients (n), it is shown mathematically in the companion study (Zhang et al. 2024) that $i^* = 3$ for $n < 1000$. Interestingly, while the $E[A_{i \geq 3}]$ is < 1, the expected $E[A_{i \geq 4}]$ is ⊠ 1, in the order of 0.001. Hence, $i^* = 4$ may be considered unnecessarily stringent.

We should note that this study is constrained by $n < 1000$ in the TCGA databases. (Databases with larger n's are also used where the actual n's are often uncertain.) At $i^* = 3$, we could detect only a fraction (<10%; see below) of CDNs. Many more tumorigenic mutations may be found in the $i = 1$ or 2 classes although not every one of them is a CDN. Since these two classes of mutations are far more numerous, they should account for the bulk of CDNs to be discovered. Indeed, **Table 1** shows 76 $A_{i \geq 3}$ CDN mutations per cancer type but 681 A_2 and 56,648 A_1 mutations in the lower recurrence groups. If n reaches $10^{5\sim6}$, most of the undiscovered CDNs in the A_1 and A_2 classes should be identified (Zhang et al. 2024).

Table 1

Table 2

In Table 2, we estimate the proportion of the A_1 and A_2 mutations that are possible CDNs. The relationships of $A_3/S_3 > A_2/S_2$, $A_2/S_2 > A_1/S_1$, and $A_1/S_1 > A_0/S_0$ are almost always observed in **Table 1** with 32 *(3 ×8 + 2 ×4)* out of 36 such relationships. The use of A/S ratios may still under-estimate the selective advantages of $A_{1\sim3}$ mutations because $S_{1\sim3}$ may have slight advantages as well (Zhang et al. 2024). Assuming S_1 is truly neutral, we use S_0 to S_1 as the basis to calculate the excess of $A_{1\sim3}$ in **Table 2** where 35 of the 36 $Obs(A_i) > Exp(A_i)$ relationships can be observed. The implication is that hundreds and, likely low thousands, of A_1's and A_2' should be CDNs whereas we have only confidently identified ~76 strong CDNs, on average, for a cancer type. (Note that A_1 excesses are less reliable since a 1% error in the calculation would mean 566 CDNs.)

2. CDNs and the amino acids affected

We now ask whether the amino acid changes associated with CDNs bear the signatures of positive selection. Amino acids that have divergent physico-chemical properties have been shown to be under strong selection, both positive and negative (Chen, He, et al. 2019; Chen, Lan, et al. 2019; Q. Chen et al. 2022). We note that, in almost all cases in cancer evolution, when a codon is altered, only one nucleotide of the triplet codon is changed. Among the 190 amino acid (AA, 20×19/2) pairs, only 75 of the pairs differ by one bp (Tang et al. 2004). For example, Pro (CCN) and Ala (GCN) may differ by only one bp but Pro and Gly (GGN) must differ by at least 2 bp. These 75 AA changes, referred to as the elementary AA changes (Grantham 1974; Li et al. 1985; Yang et al. 2003; Meyer et al. 2021), account for almost all AA substitutions in somatic evolution.

In a series of studies (Tang et al. 2004; Chen, He, et al. 2019; Chen, Lan, et al. 2019), we have defined the physico-chemical distances between AAs of the 75 elementary pairs as ΔUi, where i = 1 to 75. ΔUi reflects 47 measures of AA differences including hydrophobicity, size, charge etc. and ranges between 0 and 1. The most similar pair, Ser and Thr, has ΔUi = 0 and the most dissimilar pair is Asp and Try with ΔUi = 1. These studies show that ΔUi is a strong determinant of the evolutionary rates of DNA sequences and that large-step changes (i.e., large ΔUi's) are more acutely "recognized" by natural selection. These large-step changes are either highly deleterious or highly advantageous. Most strikingly, advantageous mutations are enriched with AA pairs of ΔUi > 0.8 (Chen, He, et al. 2019).

Fig. 2

To analyze the properties of CDNs, we choose 6 cancer types from **Table 1** that have the largest sample sizes (**n** > 500) but leap over kidney since kidney cancers have unusually low CDN counts. In **Fig. 2**, we divide the CDNs into groups according to the number of recurrences, *i*. CDNs of similar *i*'s are merged into the same group in the descending order of *i*, until there are at least 10 CDNs in the group. The 6 cancer types show two clear trends: 1) the proportion of CDNs with ΔUi > 0.8 (red color segments) increases in groups with higher recurrences; 2) in contrast, the proportion of CDNs with ΔUi < 0.4 (green segments) decreases as recurrences increase. These two trends would mean that highly recurrent CDNs tend to involve larger AA distances (ΔUi > 0.8) and similar AAs tend not to

manifest strong fitness increases. In general, CDNs alter amino acids in ways that expose the changes to strong selection.

3. CDNs in relation to the genes harboring them

We shall use the term "CDN genes" for genes having at least one CDN site. Since CDN genes contribute to tumorigenesis when harboring a CDN mutation, they should be considered cancer drivers as well. CDN genes have two desirable qualities for recognition as driver genes. First, CDNs are straightforward and unambiguous to define (e.g., $i \geq 3$ for $\mathbf{n} < 1000$). In the literature, there have been multiple definitions of cancer driver genes (Reimand and Bader 2013; Porta-Pardo and Godzik 2014; Mularoni et al. 2016; Arnedo-Pac et al. 2019), resulting in only modest overlaps among cancer gene lists (see **Fig. S3**). Second, the evolutionary fitness of CDN, and hence the tumorigenic potentials of CDN genes, can be computed (supplement File S1).

Table 3

We now present the analyses of CDN genes, using the same 6 cancers of **Fig. 2.** Two types of CDN genes are shown in **Table 3**. Type I genes fulfill the conventional criterion of fast evolution with the whole-gene Ka/Ks (or dN/dS) significantly larger than 1 (Martincorena et al. 2017). Averaged across cancer types, Type I overlaps by 95.7% with the canonical CDG list, which is the union of three popular lists (Bailey et al. 2018; Sondka et al. 2018; Martínez-Jiménez et al. 2020). Type I genes are mostly well-known canonical CDGs (e.g., *TP53*, *PIK3CA*, and *EGFR*).

Type II (CDN genes) is the new class of cancer driver genes. These genes have CDNs but do not meet the conventional criteria of whole-gene analysis. Obviously, if a gene has only one or two CDNs plus some sporadic hits, the whole-gene Ka/Ks would not be significantly greater than 1. As shown in **Table 3**, over 80% of CDN genes have only 1~2 CDN sites. The salient result is that Type II genes outnumber Type I genes by a ratio of 5 : 1 (229 : 45, column 8, **Table 3**). Furthermore, Type II genes overlap with the canonical CDG list by only 23%.

Type II genes represent a new class of cancer drivers that concentrate their tumorigenic strength on a small number CDN sites. They have been missed by the conventional whole-gene definition of cancer drivers. One such example is the *FGFR3* gene in lung cancer. This gene of 809 codons has only 8 hits, among which one is a CDN ($i = 3$) in lung cancer. It is noticed solely for this CDN. In the supplemental text, we briefly annotate these new cancer driver genes for comparisons with the canonical driver genes (supplement File S1). Possible functional tests in the future can be found in Discussion.

We now briefly discuss the driver genes listed in previous studies as shown at the lower part of **Table 3** (Bailey et al. 2018; Sondka et al. 2018; Martínez-Jiménez et al. 2020). From the total number of CDGs listed, it is clear that the overlaps are limited. As analyzed before (Wu et al. 2016), conventional gene lists overlap mainly by a core set of high Ka/Ks genes. This core set has not changed much as various criteria such the replication timing, expression profiles, epigenetic features are introduced. These criteria are the reasons for the many CDGs recognized by only a small subset of CDG lists. CDN genes, in contrast, can be objectively defined as CDN mutations (i recurrences in $\mathbf{n}$ samples) themselves are unambiguous.

Variation in CDN number and tumorigenic contribution among genes

By and large, the distribution of CDNs among genes is very uneven. **Fig. 3A** shows 10 genes with at least 6 CDNs whereas 87 genes have only one CDN. Two genes stand out for the number of CDNs they harbor, *TP53* and *PIK3CA*, which also happen to be the only genes mutated in >15% of all cancer patients surveyed (Kandoth et al. 2013). Clearly, the prevalence of mutations in a gene is a function of the number of strong CDNs it harbors.

Fig. 3

Although a small number of genes have unusually high number of CDNs, these genes may not drive the tumorigenesis in proportion to their CDN numbers in individual patients. **Fig. 3B** shows the number of CDN mutations on *TP53* that occur in any single patient. Usually, only one CDN change is observed in a patient whereas 2 or 3 CDN mutations are expected. It thus appears that CDNs on the same genes are redundant in their tumorigenic effects such that the second hit may not yield additional advantages. This pattern of disproportionally lower contribution by CDN-rich genes is true in other genes such as *EGFR* and *KRAS*. Consequently, the large number of genes with only 1 or 2 CDN sites are disproportionately important in driving the tumorigenesis of individual patients.

4. CDNs in relation to the cancer types - The pan-cancer properties
In the current literature, cancer driver genes (however they are defined) generally meet the statistical criteria for driver genes in only one or a few cancer types. However, genes may in fact contribute to tumorigenesis but are insufficiently prevalent to meet the statistical requirements for CDGs. Many genes are indeed marginally qualified as drivers in some tissues and barely miss the statistical cutoff in others. To see if genes that drive tumorigenesis in multiple tissues are more common than currently understood, we need to raise the sensitivity of cancer driver detection. Thus, CDNs may provide the resolution.

To test the pan-cancer driving capacity of CDNs, we define i_{max} as the largest i values among the 12 cancer types for each CDN. The number of cancer types where the said mutation can be detected (i.e., i > 0) is designated NC12. **Fig. 4** presents the relationship between the observed NC12 of each CDN against i_{max} of that CDN. Clearly, many CDNs are observed in multiple cancer types (NC12 > 3), even though they do not qualify as a driver gene in all but a single cancer type. It happens frequently when a mutation has i > 3 in one cancer type but has i < 3 in others. One extreme example is C394 and G395 in *IDH1*. In CNS, both sites show i ≥ 3, while in 6 other cancer types (lung, breast, large intestine, prostate, urinary tract, liver), their hits are i < 3 but > 0. Conditional on a specific site informed by a cancer type, a mutation in another cancer type should be very unlikely if the mutation is not tumorigenic in multiple tissues. Hence, the pattern in **Fig. 4** is interpreted to be drivers in multiple cancer types, but with varying statistical strength.

Fig. 4

Examining **Fig. 4** more carefully, we could see that CDNs with a larger i_{max} in one cancer type are more likely to be identified as CDNs in multiple cancer types (red dots, r = 0.97, p = 9.23×10^{-5}, Pearson's correlation test). Of 22 sites with i_{max} > 20, 15 are identified as CDNs (i ≥ 3) in multiple cancer types, with a median NC12 of 9. On the opposite end, 2 CDNs with i_{max} > 20 are observed in only one cancer

type (*EGFR*: T2573 in lung and *FGFR2*: C755 in endometrium cancer). The bimodal pattern suggests that a few cancer driver mutations are tissue specific whereas most others appear to have pan-cancer driving potentials.

To conclude, when a driver is observed in more than one cancer type, it is often a cancer driver in many others, but insufficiently powerful to meet the statistical criteria for driver mutations. This pan-cancer property can be seen at the higher resolution of CDN, but is often missed at the whole-gene level. Cancers of the same tissue in different patients, often reported to have divergent mutation profiles (Nik-Zainal et al. 2012; Roberts and Gordenin 2014), should be a good test of this hypothesis.

5. CDNs in relation to individual patients and therapeutic strategies
In previous sections, the focus is on the population of cancer patients; for example, how many in the patient population have certain mutations. We now direct the attention to individual patients. It would be necessary to pinpoint the CDN mutations in each patient in order to delineate the specific evolutionary path and to devise the treatment strategy. We shall first address the cancer driving power of CDN vs. non-CDN mutations in the same gene.

1) Efficacy of targeted therapy against CDNs vs non-CDNs
In general, a patient would have many point mutations, only a few of which are strong CDNs. We may ask whether most mutations on the canonical genes, such as *EGFR*, are CDNs. Presumably, synonymous, and likely many nonsynonymous, mutations on canonical genes may not be CDNs. It would be logical to hypothesize that patients whose *EGFR* has a CDN mutation (Group1 patients) should benefit from the gene-targeted therapy more than patients with a non-CDN mutation on the same gene (Group2 patients). In the second group, *EFGR* may be a non-driver of tumorigenesis.

Fig. 5

Published data (AACR Project GENIE Consortium 2017; Choudhury et al. 2023) are re-analyzed as shown in **Fig. 5**. The hypothesis that patients of Group2 would not benefit as much as those of Group1 is supported by the analysis. This pattern further strengthens the underlying assumption that non-CDN mutations, even on canonical genes, are not cancer drivers.

2) Number of CDNs in each patient
We postulate that a full set of CDNs should be able to inform about the cause of each cancer as well as the design of gene-targeted therapy. In **Table 4**, the known CDNs based on TCGA are tallied. Note that only a few CDNs fall on the canonical driver genes whereas most CDNs fall on the non-conventional ones.

Table 4

In most cancer types, 10%~30% of patients, shown in the n_0 row of **Table 4**, have no known CDNs (and >50% among breast cancer patients). Hence, the current practice is to rely on missense mutations, regardless of CDNs or non-CDNs, on the canonical genes. The CDN column vs. the gene column in **Table 4** address this issue. For example, the CDN column suggests that 33% of lung cancer patients (the n_0 row) would not respond well to gene-targeted therapy whereas the gene column

show only 5.3%. The difference is due to a higher, and likely inflated, detection rate of candidate drivers in the gene column. We suggest that patients who have a non-CDN mutation on a driver gene would not respond to the targeted therapy against that gene, as demonstrated in **Fig. 5**. In the above example, 27.7% (33%~5.3%) of patients may be subjected to the targeted treatment but may not respond well.

3) Prevalence vs. potency of CDN-bearing genes in driving tumorigenesis

The last question is the relationship between mutation prevalence and tumorigenic strength (or potency) among CDN-bearing genes. For example, when a patient is diagnosed to have 5 CDNs in 5 genes, what may be their relative contributions to the tumorigenesis? Are they equally valid candidates for targeted therapy? It would seem logical that canonical CDGs with many CDNs should be the targets. However, because these genes would contribute at most one CDN to the tumorigenesis (**Fig. 3B**), targeting a high prevalence gene may not yield more benefits to the patients than targeting a low prevalence gene that has a CDN.

The implication is that prevalence and potency of CDNs may not be strongly correlated. Some genes may be prevalently mutated in the patient population but, in each affected patient, these genes may not be more potent than the less prevalent genes with a CDN mutation. Potency can be tested in vitro by gene editing or in vivo by targeting treatment. In this interpretation, targeting a CDN of low prevalence (say, i = 3) may be as effective in treatment as targeting a high prevalence CDN with i = 20. The model and **Table 5** present this hypothesis based on cancer hallmarks.

Table 5

The hallmarks of cancer were first proposed by (Hanahan and Weinberg 2000) with several updates (Hanahan and Weinberg 2011; Hanahan 2022). Each hallmark is a cancer phenotype shown in **Table 5** that lists the number of genes involved in each particular hallmark (see **Methods**). While each hallmark may be associated with a number of genes, many genes are also involved in multiple hallmarks. As even the highly prevalent genes would usually have at most one mutation in each patient, we assume that each gene is associated with one hallmark in each patient.

Suppose that tumorigenesis requires a mutation in most (but perhaps not all) of the hallmarks, then the number of mutation combinations would be the product of all numbers in the corresponding column. For breast cancer, it would be $8 \times 12 \times 4 .... \times 11 \times 2 \sim 1.7 \times 10^{11}$. In other words, the possible mutation combinations that can drive breast cancer is over a billion. Hence, two breast cancers are unlikely to have the same set of CDGs or CDNs. In this view, the prevalence of a gene would be inversely proportional to the hallmark gene number. For example, genes of "*invasion and metastasis*" in breast cancer would have a prevalence of < 1/52. In contrast, the potency in tumorigenesis should depend on the hallmark phenotype itself, and independent of gene number for that hallmark. In this example, each gene of "*invasion and metastasis*" may be lowly prevalent, but could also be highly potent in each patient.

In short, the prevalence and potency of CDNs may be poorly correlated. The hypothesis can be functionally tested (by gene-editing in vitro or targeting treatment in vivo) in conjunction with the data on the attraction (i.e., co-occurrences) vs repulsion (lack of co-occurrences) of CDNs.

DISCUSSION

The companion study presents the theory that computes the limit of recurrences (*i*/**n**, *i* times in **n** patients) of reachable by neutral mutations. Above the cutoff (e.g., 3/1000), a recurrent mutation is deemed an advantageous CDN (Zhang et al. 2024). At present, the power of CDN analysis is hampered by the still small sample sizes, generally between 300 and 3000. We show that, when **n** reaches 10^5, a mutation only has to recur 12 times to be shown as a CDN, i.e., 25 times more sensitive than 3/1000. In short, nearly all CDNs should be discovered with **n** $\geq 10^5$.

In this study, we apply the theory on existing data to characterize the discovered CDNs. Based on the TCGA data, this study concludes that each cancer patient carries only 1~2 CDNs, whereas 6~10 drivers are usually hypothesized to be present in each cancer genome (Hanahan and Weinberg 2011; Vogelstein et al. 2013; Campbell et al. 2020). This deficit signifies the current incomplete understanding of cancer driving potentials. Across patients of the same cancer type, about 50 to 150 CDNs have been discovered for each cancer type, representing perhaps only 10% of all possible CDNs. Given a complete set of CDNs, it should be possible to delineate the path of tumor evolution for each individual patient.

Direct functional test of CDNs would be to introduce putative cancer-driving mutations and observe the evolution of tumors. Such a task of introducing multiple mutations that are collectively needed to drive tumorigenesis has been done only recently, and only for the best-known cancer driving mutations (Ortmann et al. 2015; Takeda et al. 2015; Hodis et al. 2022). In most tumors, the correct combination of mutations needed is not known. Clearly, CDNs, with their strong tumorigenic strength, are suitable candidates.

Many CDNs in a patient may not fall on conventional CDGs, whereas these conventional CDGs may have passenger or weak mutations. Therefore, the efforts in gene-targeting therapy may well be shifted to the CDN-harboring genes. Given a complete set of CDNs, many more driver genes can be identified. Since many driver genes cannot be targeted for biological or technical reasons (Dang et al. 2017; Danesi et al. 2021; Waarts et al. 2022), a large set of CDGs will be desirable. The goal is that each cancer patient would have multiple targetable CDGs, all driven by CDNs they carry. In that case, the probability that resistance mutations eluding multiple targeting drugs should be diminished (B. Chen et al. 2022; Zhai et al. 2022; Bian et al. 2023; Lin et al. 2023; Zhu et al. 2023).

In this context, we should comment on the feasibility of targeting CDNs that may occur in either oncogenes (ONCs) or tumor suppressor genes (TSGs). It is generally accepted that ONCs drive tumorigenesis thanks to the gain-of-function (GOF) mutations whereas TSGs derive their tumorigenic powers by loss-of-function (LOF) mutations. Nevertheless, since LOF mutations are likely to be widespread on TSGs, they are less likely to recur as CDNs. The even distributions of non-sense mutations along the length of many TSGs provide such evidence. Importantly, as gene targeting aims to diminish gene functions, GOF mutations are perceived to be targetable whereas LOF mutations are not. By extension, ONCs should be targetable but TSGs are not, an assertion we address below.

The data suggest that mis-sense mutations on TSGs may often be of the GOF kind. If mis-sense mutations are far more prevalent than nonsense mutations in tumors, the mis-sense mutations cannot possibly be LOF mutations. (After all, it is not possible to lose *more* functions than nonsense mutations.) In a separate study (Deng et al. in prep.), we compare mis-sense and nonsense mutations (referred to as the escape-route analysis). For example, AAA to AAC (K to Q) is a mis-sense mutation while the same AAA codon to AAT (K to stop) is a non-sense mutation. We found many cases where the mis-sense mutations on TSGs are more prevalent (> 10X) than nonsense mutations. We interpret these mis-sense mutations to be of the GOF kind because they could not possibly "lose" more functions than the nonsense mutations

Another interesting pattern may be the distributions of CDNs across different cancer types. Cancer evolution in different tissues represents parallel evolution driven by similar selection for cell proliferation but under different ecological conditions. **Fig. 4** suggests that CDNs previously identified to be cancer-specific may have pan-cancer effects. In different cancer types, the same CDNs may drive the tumorigenesis but the strength may not be sufficient to raise the data above the statistical threshold.

The CDN approach has two additional applications. First, it can be used to find CDNs in non-coding regions. Although the number of whole genome sequences at present is still insufficient for systematic CDN detection, the preliminary analysis suggests that the density of CDNs in non-coding regions is orders of magnitude lower than in coding regions. Second, CDNs can also be used in cancer screening with the advantage of efficiency as the targeted mutations are fewer. For the same reason, the false negative rate should be much lower too. Indeed, the false positive rate should be far lower than the gene-based screen which often shows a false positive rate of >50% (supplement File S1).

Cancer evolution falls within the realm of ultra-microevolution (Wu et al. 2016). The repeated evolution addresses the single most severe criticism of evolutionary studies, namely all evolutionary events have a sample size of one. Such repeated evolution offers the opportunity to uncover the full list of mutations underling complex traits that is at the heart of molecular evolution. The genetics of speciation (Wu and Ting 2004; Pan et al. 2022; Wang et al. 2022; Wu 2023) and the emergence of major viral strains (such as COVID-19) (Deng et al. 2022; Ruan et al. 2022; Cao et al. 2023; Ruan et al. 2023) are both phenomena of complex gene interactions. The two companion studies may thus unite evolutionary biology and cancer medicine.

References

AACR Project GENIE Consortium. 2017. AACR Project GENIE: Powering Precision Medicine through an International Consortium. *Cancer Discov* 7:818–831.

Anandakrishnan R, Varghese RT, Kinney NA, Garner HR. 2019. Estimating the number of genetic mutations (hits) required for carcinogenesis based on the distribution of somatic mutations. *PLOS Computational Biology* 15:e1006881.

Armitage P, Doll R. 1954. The Age Distribution of Cancer and a Multi-stage Theory of Carcinogenesis. *Br J Cancer* 8:1–12.

Arnedo-Pac C, Mularoni L, Muiños F, Gonzalez-Perez A, Lopez-Bigas N. 2019. OncodriveCLUSTL: a sequence-based clustering method to identify cancer drivers. *Bioinformatics* 35:4788–4790.

Bailey MH, Tokheim C, Porta-Pardo E, Sengupta S, Bertrand D, Weerasinghe A, Colaprico A, Wendl MC, Kim J, Reardon B, et al. 2018. Comprehensive Characterization of Cancer Driver Genes and Mutations. *Cell* 173:371-385.e18.

Belikov AV. 2017. The number of key carcinogenic events can be predicted from cancer incidence. *Sci Rep* 7:12170.

Bian S, Wang Y, Zhou Y, Wang W, Guo L, Wen L, Fu W, Zhou X, Tang F. 2023. Integrative single-cell multiomics analyses dissect molecular signatures of intratumoral heterogeneities and differentiation states of human gastric cancer. *National Science Review* 10:nwad094.

Bozic I, Antal T, Ohtsuki H, Carter H, Kim D, Chen S, Karchin R, Kinzler KW, Vogelstein B, Nowak MA. 2010. Accumulation of driver and passenger mutations during tumor progression. *Proceedings of the National Academy of Sciences* 107:18545–18550.

de Bruijn I, Kundra R, Mastrogiacomo B, Tran TN, Sikina L, Mazor T, Li X, Ochoa A, Zhao G, Lai B, et al. 2023. Analysis and Visualization of Longitudinal Genomic and Clinical Data from the AACR Project GENIE Biopharma Collaborative in cBioPortal. *Cancer Res* 83:3861–3867.

Campbell PJ, Getz G, Korbel JO, Stuart JM, Jennings JL, Stein LD, Perry MD, Nahal-Bose HK, Ouellette BFF, Li CH, et al. 2020. Pan-cancer analysis of whole genomes. *Nature* 578:82–93.

Cao Y, Chen L, Chen H, Cun Y, Dai X, Du H, Gao F, Guo F, Guo Y, Hao P, et al. 2023. Was Wuhan the early epicenter of the COVID-19 pandemic?—A critique. *National Science Review* 10:nwac287.

Cerami E, Gao J, Dogrusoz U, Gross BE, Sumer SO, Aksoy BA, Jacobsen A, Byrne CJ, Heuer ML, Larsson E, et al. 2012. The cBio cancer genomics portal: an open platform for exploring multidimensional cancer genomics data. *Cancer Discov* 2:401–404.

Chen B, Wu X, Ruan Y, Zhang Y, Cai Q, Zapata L, Wu C-I, Lan P, Wen H. 2022. Very large hidden genetic diversity in one single tumor: evidence for tumors-in-tumor. *Natl Sci Rev* 9:nwac250.

Chen Q, He Z, Lan A, Shen X, Wen H, Wu C-I. 2019. Molecular Evolution in Large Steps—Codon Substitutions under Positive Selection. *Molecular Biology and Evolution* 36:1862–1873.

Chen Q, Lan A, Shen X, Wu C-I. 2019. Molecular Evolution in Small Steps under Prevailing Negative Selection: A Nearly Universal Rule of Codon Substitution. *Genome Biology and Evolution* 11:2702–2712.

Chen Q, Yang H, Feng X, Chen Qingjian, Shi S, Wu C-I, He Z. 2022. Two decades of suspect evidence for adaptive molecular evolution—negative selection confounding positive-selection signals. *National Science Review* 9:nwab217.

Choudhury NJ, Lavery JA, Brown S, de Bruijn I, Jee J, Tran TN, Rizvi H, Arbour KC, Whiting K, Shen R, et al. 2023. The GENIE BPC NSCLC Cohort: A Real-World Repository Integrating Standardized

Clinical and Genomic Data for 1,846 Patients with Non–Small Cell Lung Cancer. *Clin Cancer Res* 29:3418–3428.

Danesi R, Fogli S, Indraccolo S, Del Re M, Dei Tos AP, Leoncini L, Antonuzzo L, Bonanno L, Guarneri V, Pierini A, et al. 2021. Druggable targets meet oncogenic drivers: opportunities and limitations of target-based classification of tumors and the role of Molecular Tumor Boards. *ESMO Open* 6:100040.

Dang CV, Reddy EP, Shokat KM, Soucek L. 2017. Drugging the "undruggable" cancer targets. *Nat Rev Cancer* 17:502–508.

Deng S, Xing K, He X. 2022. Mutation signatures inform the natural host of SARS-CoV-2. *National Science Review* 9:nwab220.

Grantham R. 1974. Amino Acid Difference Formula to Help Explain Protein Evolution. *Science* 185:862–864.

Hanahan D. 2022. Hallmarks of Cancer: New Dimensions. *Cancer Discovery* 12:31–46.

Hanahan D, Weinberg RA. 2000. The Hallmarks of Cancer. *Cell* 100:57–70.

Hanahan D, Weinberg RA. 2011. Hallmarks of Cancer: The Next Generation. *Cell* 144:646–674.

Hodis E, Triglia ET, Kwon JYH, Biancalani T, Zakka LR, Parkar S, Hütter J-C, Buffoni L, Delorey TM, Phillips D, et al. 2022. Stepwise-edited, human melanoma models reveal mutations' effect on tumor and microenvironment. *Science* 376:eabi8175.

Kandoth C, McLellan MD, Vandin F, Ye K, Niu B, Lu C, Xie M, Zhang Q, McMichael JF, Wyczalkowski MA, et al. 2013. Mutational landscape and significance across 12 major cancer types. *Nature* 502:333–339.

Lagou V, Jiang L, Ulrich A, Zudina L, González KSG, Balkhiyarova Z, Faggian A, Maina JG, Chen S, Todorov PV, et al. 2023. GWAS of random glucose in 476,326 individuals provide insights into diabetes pathophysiology, complications and treatment stratification. *Nat Genet* 55:1448–1461.

Lawrence MS, Stojanov P, Polak P, Kryukov GV, Cibulskis K, Sivachenko A, Carter SL, Stewart C, Mermel CH, Roberts SA, et al. 2013. Mutational heterogeneity in cancer and the search for new cancer-associated genes. *Nature* 499:214–218.

Li WH, Wu CI, Luo CC. 1985. A new method for estimating synonymous and nonsynonymous rates of nucleotide substitution considering the relative likelihood of nucleotide and codon changes. *Molecular Biology and Evolution* 2:150–174.

Lin J, Zhan G, Liu J, Maimaitiyiming Y, Deng Z, Li B, Su K, Chen J, Sun S, Zheng W, et al. 2023. YTHDF2-mediated regulations bifurcate BHPF-induced programmed cell deaths. *National Science Review* 10:nwad227.

Martincorena I, Raine KM, Gerstung M, Dawson KJ, Haase K, Van Loo P, Davies H, Stratton MR, Campbell PJ. 2017. Universal Patterns of Selection in Cancer and Somatic Tissues. *Cell* 171:1029-1041.e21.

Martínez-Jiménez F, Muiños F, Sentís I, Deu-Pons J, Reyes-Salazar I, Arnedo-Pac C, Mularoni L, Pich O, Bonet J, Kranas H, et al. 2020. A compendium of mutational cancer driver genes. *Nat Rev Cancer* 20:555–572.

Meyer D, Kames J, Bar H, Komar AA, Alexaki A, Ibla J, Hunt RC, Santana-Quintero LV, Golikov A, DiCuccio M, et al. 2021. Distinct signatures of codon and codon pair usage in 32 primary tumor types in the novel database CancerCoCoPUTs for cancer-specific codon usage. *Genome Med* 13:122.

Mularoni L, Sabarinathan R, Deu-Pons J, Gonzalez-Perez A, López-Bigas N. 2016. OncodriveFML: a general framework to identify coding and non-coding regions with cancer driver mutations. *Genome Biology* 17:128.

Nei M, Gojobori T. 1986. Simple methods for estimating the numbers of synonymous and nonsynonymous nucleotide substitutions. *Molecular Biology and Evolution* 3:418–426.

Nik-Zainal S, Alexandrov LB, Wedge DC, Van Loo P, Greenman CD, Raine K, Jones D, Hinton J, Marshall J, Stebbings LA, et al. 2012. Mutational Processes Molding the Genomes of 21 Breast Cancers. *Cell* 149:979–993.

Ortmann CA, Kent DG, Nangalia J, Silber Y, Wedge DC, Grinfeld J, Baxter EJ, Massie CE, Papaemmanuil E, Menon S, et al. 2015. Effect of Mutation Order on Myeloproliferative Neoplasms. *N Engl J Med* 372:601–612.

Pan Y, Zhang C, Lu Y, Ning Z, Lu D, Gao Y, Zhao X, Yang Y, Guan Y, Mamatyusupu D, et al. 2022. Genomic diversity and post-admixture adaptation in the Uyghurs. *National Science Review* 9:nwab124.

Passaro A, Leighl N, Blackhall F, Popat S, Kerr K, Ahn MJ, Arcila ME, Arrieta O, Planchard D, De Marinis F, et al. 2022. ESMO expert consensus statements on the management of EGFR mutant non-small-cell lung cancer. *Annals of Oncology* 33:466–487.

Porta-Pardo E, Godzik A. 2014. e-Driver: a novel method to identify protein regions driving cancer. *Bioinformatics* 30:3109–3114.

Reimand J, Bader GD. 2013. Systematic analysis of somatic mutations in phosphorylation signaling predicts novel cancer drivers. *Molecular Systems Biology* 9:637.

Roberts SA, Gordenin DA. 2014. Hypermutation in human cancer genomes: footprints and mechanisms. *Nat Rev Cancer* 14:786–800.

Ruan Y, Wen H, Hou M, He Z, Lu X, Xue Y, He X, Zhang Y-P, Wu C-I. 2022. The twin-beginnings of COVID-19 in Asia and Europe—one prevails quickly. *National Science Review* 9:nwab223.

Ruan Y, Wen H, Hou M, Zhai W, Xu S, Lu X. 2023. On the epicenter of COVID-19 and the origin of the pandemic strain. *National Science Review* 10:nwac286.

Sherman MA, Yaari AU, Priebe O, Dietlein F, Loh P-R, Berger B. 2022. Genome-wide mapping of somatic mutation rates uncovers drivers of cancer. *Nat Biotechnol*:1–10.

Sondka Z, Bamford S, Cole CG, Ward SA, Dunham I, Forbes SA. 2018. The COSMIC Cancer Gene Census: describing genetic dysfunction across all human cancers. *Nat Rev Cancer* 18:696–705.

Sun S, Wang Y, Maslov AY, Dong X, Vijg J. 2022. SomaMutDB: a database of somatic mutations in normal human tissues. *Nucleic Acids Research* 50:D1100–D1108.

Suzuki K, Hatzikotoulas K, Southam L, Taylor HJ, Yin X, Lorenz KM, Mandla R, Huerta-Chagoya A, Melloni GEM, Kanoni S, et al. 2024. Genetic drivers of heterogeneity in type 2 diabetes pathophysiology. *Nature* 627:347–357.

Takeda H, Wei Z, Koso H, Rust AG, Yew CCK, Mann MB, Ward JM, Adams DJ, Copeland NG, Jenkins NA. 2015. Transposon mutagenesis identifies genes and evolutionary forces driving gastrointestinal tract tumor progression. *Nat Genet* 47:142–150.

Tang H, Wyckoff GJ, Lu J, Wu C-I. 2004. A universal evolutionary index for amino acid changes. *Mol Biol Evol* 21:1548–1556.

Tate JG, Bamford S, Jubb HC, Sondka Z, Beare DM, Bindal N, Boutselakis H, Cole CG, Creatore C, Dawson E, et al. 2019. COSMIC: the Catalogue Of Somatic Mutations In Cancer. *Nucleic Acids Research* 47:D941–D947.

Vogelstein B, Papadopoulos N, Velculescu VE, Zhou S, Diaz LA, Kinzler KW. 2013. Cancer Genome Landscapes. *Science* 339:1546–1558.

Vujkovic M, Keaton JM, Lynch JA, Miller DR, Zhou J, Tcheandjieu C, Huffman JE, Assimes TL, Lorenz K, Zhu X, et al. 2020. Discovery of 318 new risk loci for type 2 diabetes and related vascular outcomes among 1.4 million participants in a multi-ancestry meta-analysis. *Nat Genet* 52:680–691.

Waarts MR, Stonestrom AJ, Park YC, Levine RL. 2022. Targeting mutations in cancer. *J Clin Invest* 132:e154943.

Wang X, He Z, Guo Z, Yang M, Xu S, Chen Q, Shao S, Li S, Zhong C, Duke NC, et al. 2022. Extensive gene flow in secondary sympatry after allopatric speciation. *National Science Review* 9:nwac280.

Weinstein JN, Collisson EA, Mills GB, Shaw KRM, Ozenberger BA, Ellrott K, Shmulevich I, Sander C, Stuart JM. 2013. The Cancer Genome Atlas Pan-Cancer analysis project. *Nat Genet* 45:1113–1120.

Wu C-I ed. 2022. What are species and how are they formed? *National Science Review* 9:nwad017.

Wu C-I. 2023. The genetics of race differentiation—should it be studied? *National Science Review* 10:nwad068.

Wu C-I, Ting C-T. 2004. Genes and speciation. *Nat Rev Genet* 5:114–122.

Wu C-I, Wang H-Y, Ling S, Lu X. 2016. The Ecology and Evolution of Cancer: The Ultra-Microevolutionary Process. *Annu. Rev. Genet.* 50:347–369.

Xue D, Narisu N, Taylor DL, Zhang M, Grenko C, Taylor HJ, Yan T, Tang X, Sinha N, Zhu J, et al. 2023. Functional interrogation of twenty type 2 diabetes-associated genes using isogenic human embryonic stem cell-derived β-like cells. *Cell Metabolism* 35:1897-1914.e11.

Yang Z, Ro S, Rannala B. 2003. Likelihood Models of Somatic Mutation and Codon Substitution in Cancer Genes. *Genetics* 165:695–705.

Yang Z, Swanson WJ. 2002. Codon-Substitution Models to Detect Adaptive Evolution that Account for Heterogeneous Selective Pressures Among Site Classes. *Molecular Biology and Evolution* 19:49–57.

Zhai W, Lai H, Kaya NA, Chen J, Yang H, Lu B, Lim JQ, Ma S, Chew SC, Chua KP, et al. 2022. Dynamic phenotypic heterogeneity and the evolution of multiple RNA subtypes in hepatocellular carcinoma: the PLANET study. *National Science Review* 9:nwab192.

Zhang L, Deng T, Liufu Z, Liu X, Chen B, Hu Z, Liu C, Lu X, Wen H, Wu C-I. 2024. The theory of massively repeated evolution and full identifications of Cancer Driving Nucleotides (CDNs). *eLife* [Internet] 13. Available from: https://elifesciences.org/reviewed-preprints/99340

Zhu H, Lin Y, Lu D, Wang S, Liu Y, Dong L, Meng Q, Gao J, Wang Y, Song N, et al. 2023. Proteomics of adjacent-to-tumor samples uncovers clinically relevant biological events in hepatocellular carcinoma. *National Science Review* 10:nwad167.

Methods

Data preparation

Single-nucleotide variant (SNV) data for TCGA patients were downloaded from the GDC Data Portal (https://portal.gdc.cancer.gov/, data version 2022-02-28), with mutations identified by at least two pipelines were included in this study. Mutations exceeding a 1‰ frequency in the Genome Aggregation Database (*gnomAD*, version v2.1.1) were excluded to minimize potential false positives arising from germline variants. Patients with more than 3000 coding region point mutations were filtered out as potential hypermutator phenotypes. This filtering process yielded a final analysis set encompassing 7369 patients across 12 diverse cancer types for subsequent analysis. The calculation of A_i and S_i follows the same method as described in the companion paper (https://elifesciences.org/reviewed-preprints/99340).

For CDN analysis in non-cancerous tissues, mutation profiles for normal tissues were retrieved from *SomaMutDB* (Sun et al. 2022). Mutations from different samples originating from the same individual were consolidated. Donners above the age of 80 were excluded from our dataset. The mutation processing followed the same pipeline as previously described. In total, we have mutation profiles from 487 donners serving as a negative control.

The canonical lists of cancer driver genes were obtained from three distinct data sources. The CGC Tier 1 genes, encompassing genes with the highest confidence of driver status, were retrieved from the COSMIC Cancer Gene Census (https://cancer.sanger.ac.uk/census) (Sondka et al. 2018). The IntOGen driver gene list, which employs an integrated pipeline for gene discovery, was downloaded from https://www.intogen.org/download (Martínez-Jiménez et al. 2020). Bailey's driver gene list comprises 299 cancer driver genes identified through a *PanSoftware* strategy, with further experimental validation confirming their role in driving cell lines (Bailey et al. 2018). The consistency of cancer types across all studies was manually verified using *oncotree* (https://oncotree.mskcc.org/#/home). For the analysis of driver gene overlap, only drivers from the same cancer type were compared.

The hallmark annotation of genes is downloaded from COSMIC (https://cancer.sanger.ac.uk/cosmic/download), encompassing 331 genes with annotated dysregulated biological processes. It is important to note that these hallmarks are manually annotated as part of an ongoing effort to characterize the role of genes in cancer based on literature evidence. The actual scale of hallmark genes may be substantially larger than the current version.

For gene-level selection analysis, we utilized the R package '*dndscv*' to quantify selection signals for missense and nonsense mutations in a given gene (Martincorena et al. 2017). Specifically, the package calculates the Ka/Ks ratio, denoted as 'w' in the final results, for a given mutation impact (missense or nonsense). The significance of selection is presented as q values after Benjamini-Hochberg (BH) adjustment. Genes with $w > 1$ and $q < 0.1$ were identified as being significantly under positive selection.

We employ $i^* = 3$ as a cutoff for identifying Cancer Driving Nucleotides (CDNs) across various cancer types. The specific value of i^* is detailed in Eq. 10 of the companion paper (Zhang et al. 2024). Here, $i^* = 3$ is chosen consistently across all cancer types, taking into account the abundance of sites under positive selection given $i = 3$ in **Table 2**. Throughout our analysis, emphasis is placed on CDNs of the missense category, where missense mutations with a recurrence ≥3 are identified as CDNs. For ΔUi analysis, the reference table for 75 single-step amino acid changes was obtained from (Chen, He, et al. 2019), and the ΔUi for each CDN is derived by mapping the amino acid change to the reference table.

Calculation of A_i_e
We employ Eq. 9 from the companion paper to calculate the expected value for A_i under neutrality. For a given site, the cumulative probability for recurrence $x \le i - 1$ could be expressed as:

$$F(x \le i - 1) = 1 - \left(1 - \frac{1}{1 + nE(u)}\right)^{i-1}$$

Eq. S1

where **n** is the population size of a given cancer type, and **E(u)** is the mutation rate per site per patient derived from singleton synonymous mutations:

$$S_1 = L_S \cdot nE(u)e^{-(n-1)E(u)}$$

Eq. S2

Then by expectation, site number of recurrence i ($A_{x \ge i}$) could be represented by:

$$A_{x \ge i} = L_A - L_A \cdot F(x \le i - 1)$$

Following the same logic, we'll have $\boldsymbol{A_{x \geq i+1}}$ as:

$$A_{x \geq i+1} = L_A - L_A \cdot F(x \leq i)$$

Then the expected value for $\boldsymbol{A_i\_e}$ is then:

$$A_i\_e = A_{x \geq i} - A_{x \geq i+1} = L_A \cdot [F(x \leq i) - F(x \leq i-1)]$$

Eq. S3

L_A and L_S are missense and synonymous sites, respectively. The calculation procedure is described in methods of the companion paper (Zhang et al. 2024).

With Eqs. S1~S3, we could solve for the expected number of sites with missense mutation recurrence i.

Survival analysis of *EGFR*-targeted therapy

The mutation and clinical profiles of 23,253 patients were retrieved from the GENIE project (Cerami et al. 2012; de Bruijn et al. 2023), with 7,216 patients harboring *EGFR* mutations. Survivor months were calculated as the time elapsed between the date of sequencing and the date of the last contact (or day of death). In cases where patients had multiple sequencing reports, the earliest one was selected. For CDN calling, we applied Eq. 10 from the companion paper (Zhang et al. 2024). With $\varepsilon = 0.01$, we set the CDN cutoff $i^* = 14$. To mitigate potential biases from other common drivers in lung cancer, patients with indels in exon 19 and 20 of *EGFR*, G12/13 mutations in *KRAS*, V600 mutations in *BRAF*, exon 20 insertions in *HER2*, fusions in *MET, ALK, ROS1, RET, NTRK*, and *MET* were filtered out. The final survival analysis was conducted using *GraphPad Prism 8*.

Annotation for non-canonical CDN genes

We conducted functional annotation and enrichment analysis for newly identified non-canonical CDN genes using four independent databases (Gene Ontology, KEGG, Disease Ontology, and Reactome) with R packages (*clusterProfiler, DOSE, ReactomePA*). For each analysis, we set a *p*-value cutoff of 0.05 and a *q*-value cutoff of 0.2, with *p* value adjustment method set to "BH". To explore the connections between non-canonical CDN genes and canonical cancer driver genes (CDGs), enrichment analyses were performed alongside cancer drivers from IntOGen. Specifically, for enrichment annotations related to cancer hallmarks, the corresponding genes were subjected to manual confirmation using CancerGeneNET (https://signor.uniroma2.it/CancerGeneNet/).

Data Availability

The scripts for generating the key results of this study and the accompanying paper (https://doi.org/10.7554/eLife.99340.1) are available at https://gitlab.com/ultramicroevo/cdn_v1. Example files for breast cancer analysis have also been included. The complete set of CDNs can be found in Supplementary File S2 of the accompanying paper (https://elifesciences.org/reviewed-preprints/99340).

ACKNOWLEDGEMENTS

We wish to acknowledge the supports from the First Affiliated Hospital, the Seventh Affiliated Hospital of Sun Yat-sen University, Cancer Center of Clifford Hospital, Jinan University, Cancer Hospital Chinese Academy of Medical Sciences, Shenzhen Center, and Guangdong Academy of Medical Sciences, Guangdong Provincial People's Hospital on the startup of the Cancer Driving Nucleotide (CDN) project. We would like to acknowledge Kunming Institute of Zoology for discussing the ideas of CDN. We thank Weiwei Zhai, Qianfei Wang, and Weini Huang for insightful comments and suggestions. We would also

like to acknowledge the American Association for Cancer Research (AACR) and The Cancer Genome Atlas (TCGA) project, which have provided invaluable datasets and resources that have significantly enriched our understanding of cancer biology and improved patient outcomes. This work was supported by the National Natural Science Foundation of China (32150006, 32293193/32293190 to C.I.W., 82341092 to HJ Wen, and 32200493 to Y.R.), the National Key Research and Development Projects of the Ministry of Science and Technology of China (2021YFC2301300, 2021YFC0863400), and Guangdong Key Research and Development Program (No. 2022B1111030001).

DECLARATION OF INTERESTS

The authors declare no competing interests.

TABLES AND FIGURES

Table 1- Mutation recurrences (A_i's and S_i's) in 12 cancer types.

	Lung	Breast	CNS	Kidney	Upper-AD tract	Colon	Endo-met rium	Prostate	Stomach	Urinary tract	Ovary	Liver	Average
patients #	*1035*	*963*	*873*	*711*	*688*	*571*	*465*	*465*	*423*	*404*	*404*	*367*	*614*
A_0	22540623	21683136	20783835	22247653	21580444	20601026	20766001	21300810	20892755	21628918	22278124	22618059	21576782
S_0	7804281	9388418	10298911	8781483	9333283	10428913	10375596	9754331	10243634	9426888	8746002	8255268	9403084
A/S_0	2.89	2.31	2.02	2.53	2.31	1.98	2.00	2.18	2.04	2.29	2.55	2.74	2.29
A_1	195958	44696	25122	25669	66924	94634	78870	9583	78834	66153	21138	25731	61109
S_1	69393	16732	10182	9317	26151	38606	31982	3613	32538	26546	7227	9398	23474
A/S_1	2.82	2.67	2.47	2.76	2.56	2.45	2.47	2.65	2.42	2.49	2.92	2.74	2.60
A_2	2946	233	287	56	489	1662	1052	29	1176	816	51	46	737
S_2	969	62	75	11	159	736	386	9	489	308	9	12	249
A/S_2	3.04	3.76	3.83	5.09	3.08	2.26	2.73	3.22	2.40	2.65	5.67	3.83	2.74
A_3	99	18	42	14	28	91	52	6	79	60	9	9	42.3
S_3	21	2	6	1	5	28	11	0	14	9	0	0	8.08
A/S_3	4.71	9	7	14	5.6	3.25	4.73	6 : 0	5.64	6.67	9 : 0	9 : 0	5.23
$A_{i≥3}$	178	51	84	18	77	148	142	14	124	100	26	23	82.1
$A_{i≥4}$	79	33	42	4	49	57	90	8	45	40	17	14	39.8
A_4	23	10	8	2	14	23	21	3	23	11	4	3	11.1
A_5	16	6	10	2	10	6	20	2	9	9	3	5	8.2
A_{6-9}	27	10	10	0	13	9	32	2	7	12	6	2	10.8
$A_{[10,20)}$	7	3	10	0	9	11	9	1	6	5	4	4	5.75
$A_{≥20}$	6	4	4	0	3	8	8	0	0	3	0	0	3
Total	202828	45669	26596	25841	68387	98931	81898	9706	81678	68297	21387	25944	63097
SiteNbr	22739705	21728116	20809328	22273396	21647934	20697470	20846065	21310436	20972889	21695987	22299339	22643859	21638710
nE(u)	9.07E-03	1.79E-03	1.00E-03	1.06E-03	2.83E-03	3.84E-03	3.15E-03	3.72E-04	3.27E-03	2.88E-03	8.28E-04	1.14E-03	2.6E-03

a – A_i and S_i are as defined in the text.

b – "Total" represents the total number of missense mutations, or $\sum i \cdot A_i$. "Site number" refers to the count of missense sites. **nE(u)** is calculated based on synonymous mutations, representing the expected number of neutral mutations per site in a population of size **n**.

c – See methods for the calculations of A_0 and S_0.

Table 2- Excess of A_i's of each i class.

Recurrences	Lung	Breast	CNS	Kidney	Upper-AD tract	Colon	Endo-met rium	prostate	Stomach	Urinary tract	Ovary	liver
A_1_o	195958	44696	25122	25669	66924	94634	78870	9583	78834	66153	21138	25731
A_1_e	198627	38586	20532	23582	60316	76049	63860	7888	66194	60751	18396	25720
Excess	-2669	6110	4590	2087	6608	18585	15010	1695	12640	5402	2742	11
Ratio	**-1.36%**	**13.67%**	**18.27%**	**8.13%**	**9.87%**	**19.64%**	**19.03%**	**17.69%**	**16.03%**	**8.17%**	**12.97%**	**0.04%**
A_2_o	2946	233	287	56	489	1662	1052	29	1176	816	51	46
A_2_e	1750	69	20	25	169	280	196	3	210	171	15	29
Excess	1195.61	164.36	266.72	31.01	320.48	1381.54	855.77	26.08	966.42	645.41	35.81	16.75
Ratio	**40.58%**	**70.54%**	**92.93%**	**55.37%**	**65.54%**	**83.13%**	**81.35%**	**89.93%**	**82.18%**	**79.09%**	**70.22%**	**36.42%**
A_3_o	99	18	42	14	28	91	52	6	79	60	9	9
A_3_e	15.43	0.12	0.02	0.03	0.47	1.03	0.60	0.00	0.66	0.48	0.01	0.03
Excess	83.57	17.88	41.98	13.97	27.53	89.97	51.40	6.00	78.34	59.52	8.99	8.97
Ratio	**84.42%**	**99.32%**	**99.95%**	**99.81%**	**98.32%**	**98.86%**	**98.84%**	**99.98%**	**99.16%**	**99.20%**	**99.86%**	**99.63%**
A_4_o	23	10	8	2	14	23	21	3	23	11	4	3
A_4_e	0.13593	0.00022	1.98E-05	2.81E-05	0.00132	0.00381	0.00185	4.00E-07	0.00210	0.00135	1.04E-05	3.78E-05
Excess	22.8641	9.99978	7.99998	1.99997	13.9987	22.9962	20.9981	3	22.9979	10.9987	3.99999	2.99999
Ratio	**99.41%**	**100 %**	**100 %**	**100 %**	**99.99%**	**99.98%**	**99.99%**	**100.00%**	**99.99%**	**99.99%**	**100 %**	**100.00%**

a - The notation of "**o**" and "**e**" following A_i's represents the observed A_i and expected A_i.
b - See methods for the calculation of expected A_i's.
c - Ratio is the proportion of observed sites in excess, i.e., the proportion of putative CDNs in the observation.

Table 3- Distribution of CDNs among genes.

CDN calls based on $i^* = 3$	Lung	Breast	CNS	Upper-AD tract	Colon	Endo-metrium	Mean	Total[1]	Overlap with the conventional set	Criteria of classification
# of patients (n)	1035	963	873	688	571	465	-	-	-	
CDN count	178	50	83	77	148	142	113.3	495	-	
# CDN-carrying genes (Type I fulfills the convention of Ka/Ks > 1**; Type II does not. ** means statistical significance)										
Type I (Ka/Ks > 1 **)	10	8	12	13	10	21	12.33	45	95.7%	Conventional
Type II (Ka/Ks ∼ 1)	79	9	12	19	86	35	40	229	26.1%	This study only
All CDN genes	89	17	24	32	96	56	52.33	258	47%	Both types
Genes with 1-2 CDNs (% all CDN genes)	80 (89.9 %)	14 (82.4 %)	19 (79.2 %)	27 (84.4 %)	90 (93.8 %)	45 (80.4 %)	45.8 (85 %)	250 (96.9%)		A subset of both types
Number of driver genes in 3 major CDG lists										
Other criteria*:									---	Variable (see Legends)
IntOGen	118	100	100	106	86	72	97	321		
Bailey et al.	36	29	32	38	20	55	35	134		
CGC Tier1	30	32	32	24	44	23	30.83	118		

Note –

* *intOGen*, *Bailey et al.*, and *CGC Tier1* are the three major CDG lists adopted here for comparison (Bailey et al. 2018; Sondka et al. 2018; Martínez-Jiménez et al. 2020).

Table 4 - Numbers of patients with CDNs vs. number of patients with any non-synonymous mutations in the same genes.

	Lung		Breast		CNS		Upper-AD tract		Colon		Endometrium	
	CDN[1,3] (178)	gene[2,3] (89)	CDN (50)	Gene (17)	CDN (83)	Gene (24)	CDN (77)	Gene (32)	CDN (148}	Gene (96)	CDN (142)	Gene (56)
n_0	342 (33%)[4]	53 (5.3%)	492 (51.1%)	415 (43.1%)	235 (26.9%)	163 (18.7%)	268 (39%)	140 (20.3%)	102 (17.9%)	42 (7.4%)	42 (9%)	14 (3%)
n_1	411 (39.7%)	70 (6.8%)	379 (39.4%)	395 (41%)	359 (41.1%)	306 (35.1%)	268 (39%)	229 (33.3%)	159 (27.8%)	79 (13.8%)	108 (23.2%)	59 (12.7%)
n_2	192 (18.6%)	84 (8.1%)	73 (7.6%)	114 (11.8%)	225 (25.8%)	293 (33.6%)	101 (14.7%)	171 (24.9%)	140 (24.5%)	93 (16.3%)	169 (36.3%)	101 (21.7%)
$n_{>2}$	90 (8.7%)	826 (79.8%)	18 (1.9%)	38 (3.9%)	53 (6.1%)	110 (12.6%)	50 (7.3%)	147 (21.4%)	170 (29.8%)	357 (62.5%)	146 (31.4%)	291 (62.6%)
Total **n**	1035	1035	963	963	873	873	688	688	571	571	465	465
Mean #	1.06	7.19	0.61	0.78	1.12	1.44	0.93	1.63	1.96	4.6	2.17	3.7

Note –

[1] - n_i designates the number of patients with i CDN mutations.

[2] - In this column, n_i designates the number of patients with any nonsynonymous mutation in the same gene as the CDN column.

[3] - The number in the parentheses is the total number of CDNs or genes.

[4] - There are 684 CDNs summed over all cancer types. The percentage is n_i/Total **n**.

Table 5 - Gene numbers for different cancer hallmarks.

hallmark	gene number		
	all records	breast	colon
angiogenesis	78	8	6
cell division control	107	12	10
cell replicative immortality	44	4	3
change of cellular energetics	70	10	4
escaping immune response to cancer	51	1	1
escaping programmed cell death	202	32	20
genome instability and mutations	106	10	7
invasion and metastasis	206	52	27
proliferative signaling	176	40	20
senescence	48	3	5
suppression of growth	130	11	12
tumor promoting inflammation	54	2	3

Note - data downloaded from COSMIC (https://cancer.sanger.ac.uk/cosmic/download), see methods.

Figure 1: Mutations in organismal evolution vs. cancer evolution. **(A, B)** A hypothetical example of DNA sequence evolution in organism vs. in cancer with the same number of mutations. **(C)** Mutation distribution in two species in the organismal evolution of **A**. **(D** and **E)** Mutation distribution in cancer evolution among 10 sequences may have D and E patterns. **(F)** Another pattern of mutation distribution in cancer evolution with a recurrent site but shows too few total mutations. Mutations of **(F)** are CDNs missed in the conventional screens.

Figure 2: analysis across 6 cancer types. , ranging between 0 and 1 (Tang et al. 2004; Chen, He, et al. 2019), is a measure of physico-chem differences among the 20 amino acids (see the text). The most similar amino acids have near 0 and the most dissimilar ones have near 1 Each panel corresponds to one cancer type, with horizontal bar represents distribution of each recurrence group. The numbers on the left of panel are *i* values and on the right are the number of sites. Note that the proportion of dark red segments increases as *i* increases. This figure shows that mutations at high recurrence sites (larger *i*'s) code for amino acids that are chemically very different from the wild type.

A)

B)

Figure 3: Distribution of CDNs among genes. (**A**) Out of 119 CDN-carrying genes (red bars), 87 have only one CDN. For the rest, *TP53* possesses the most CDNs with three others having more than 10 CDNs. (**B**) CDN number in *TP53* among patients. The dark bar represents the observed patient number with corresponding CDNs of the X-axis. The grey bar shows the expected patient distribution. Clearly, *TP53* only needs to contribute one CDN to drive tumorigenesis. Hence, *TP53* (and other canonical driver genes; see text), while prevalent, does not contribute disproportionately to the tumorigenesis of each patient.

	1	**2**	**3**	**4**	**5**	**6**	**7**	**8**	**9**	**10–20**	**>20**
Sites	702008	8514	390	95	61	37	18	11	14	38	22
∑Sites	711208	9200	686	296	201	140	103	85	74	60	22
Genes	17815	5069	313	67	38	25	13	9	11	20	13
∑Genes	17816	5186	393	119	72	50	37	33	31	26	13

Figure 4: Sharing of CDNs across cancer types. The X-axis shows i_{max}, which is the largest i a CDN reaches among the 12 cancer types. The Y axis shows the number of cancer types whre the mutation also occurs. Each dot is a CDN and the number of dots in the cloud is given. The blue and red dots denote, respectively, mutations classified as a CDN in one or multiple cancer types. Grey dots are non-CDNs. The table in the lower panel summarizes the number of sites and the number of genes harboring these sites.

Figure 5. Survival analysis of non-small cell lung cancer (NSCLC) patients based on EGFR mutation status. Patient data were retrieved from the GENIE database (https://genie.cbioportal.org/) and stratified into three groups based on *EGFR* mutation profiles: Group1 comprises patients with *EGFR* CDN mutations; Group2 includes patients with nonsynonymous mutations in *EGFR* that are not CDNs; The *EGFR^{WT}* group consists of patients with no *EGFR* mutations (see **methods**). Patients of Group1 and Group2 received *EGFR*-targeted therapies in accordance with the guidelines for managing *EGFR* mutant NSCLC (Passaro et al. 2022; Choudhury et al. 2023). Survival analysis using the Kaplan-Meier method revealed a significantly higher survival rate for Group1 patients compared to Group2 and the *EGFR^{WT}* group (p < 0.001).