TITLE

Systematic genetic characterization of the human PKR kinase domain highlights its functional

malleability to escape a poxvirus substrate mimic

AUTHORS

Michael J. Chambers (1,2), Sophia B. Scobell (1), Meru J. Sadhu (1)

AFFILIATIONS

1.   Center for Genomics and Data Science Research, National Human Genome Research

Institute, National Institutes of Health, Bethesda, Maryland, USA

2.   Department of Microbiology & Immunology, Georgetown University, Washington DC, USA

ABSTRACT

Evolutionary arms races can arise at the contact surfaces between host and viral

proteins, producing dynamic spaces in which genetic variants are continually pursued.

However, the sampling of genetic variation must be balanced with the need to maintain protein

function. A striking case is given by protein kinase R (PKR), a member of the mammalian innate

immune system. PKR detects viral replication within the host cell and halts protein synthesis to

prevent viral replication by phosphorylating eIF2α, a component of the translation initiation

machinery. PKR is targeted by many viral antagonists, including poxvirus pseudosubstrate

antagonists that mimic the natural substrate, eIF2α, and inhibit PKR activity. Remarkably, PKR

has several rapidly evolving residues at this interface, suggesting it is engaging in an

evolutionary arms race, despite the surface's critical role in phosphorylating eIF2α. To

systematically explore the evolutionary opportunities available at this dynamic interface, we

generated and characterized a library of 426 SNP-accessible nonsynonymous variants of

human PKR for their ability to escape inhibition by the model pseudosubstrate inhibitor K3,

encoded by the vaccinia virus gene *K3L*. We identified key sites in the PKR kinase domain that

harbor K3-resistant variants, as well as critical sites where variation leads to loss of function. We

find K3-resistant variants are readily available throughout the interface and are enriched at sites

under positive selection. Moreover, variants beneficial against K3 were also beneficial against

an enhanced variant of K3, indicating resilience to viral adaptation. Overall, we find that the

eIF2α-binding surface of PKR is highly malleable, potentiating its evolutionary ability to combat

viral inhibition.

INTRODUCTION

Molecular interfaces between interacting host and pathogen proteins are atomic battlegrounds that give rise to evolutionary arms races (1). Successful infection typically involves critical interactions between host and pathogen proteins, such as binding to cell-surface receptors or inhibition of cellular defense mechanisms. In such scenarios, the host benefits from genetic mutations that disrupt the interaction while the pathogen benefits from mutations that maintain the interaction. The pursuit of genetic mutations by the host and pathogen can develop into an evolutionary arms race where the accumulation of mutations leads to abnormally high sequence diversity, which is used to identify sites experiencing positive selection.

A key member of the antiviral innate immune repertoire in vertebrates is protein kinase R (PKR) (2–4). PKR contains two N-terminal double-stranded RNA (dsRNA) binding domains and a C-terminal kinase domain, separated by a long linker (5). The presence of dsRNA in the cytoplasm is a hallmark of viral infection (6). PKR binds dsRNA, initiating PKR homodimerization and autophosphorylation, which activates the kinase (7–10). Active PKR binds and phosphorylates eIF2α on Ser51 (Figure 1A), a member of the translation initiation machinery. Phosphorylation of eIF2α leads to a halt in protein synthesis, thereby preventing viral replication (8,11,12).

Vertebrate PKR homologs have many sites under positive selection (13), including residues at its binding surface with eIF2α (Figure 1B). This is remarkable given the critical nature of this interface and that eIF2α is an unvarying protein under purifying selection (14). Many viruses encode PKR inhibitors to evade its antiviral activity, some of which are known to interact with PKR at its eIF2α-binding interface (15–17). One example is vaccinia virus, whose well-characterized pseudosubstrate inhibitor protein K3 is encoded by the *K3L* gene (Figure 1C) (18–20). Vaccinia K3 mimics the structure and sequence of eIF2α to inhibit PKR and prevent the halt of protein synthesis (19,21–23). Thus, despite evolutionary pressure on PKR to

recognize the conserved eIF2α substrate, variation at the same interface would be evolutionarily

beneficial if it allowed evasion of viral pseudosubstrate antagonists. Indeed, previous studies

have identified some single-residue variants at sites under positive selection in the PKR kinase

domain that maintain recognition of eIF2α yet resist vaccinia K3 (13,14,24). However, without

comprehensively exploring the genetic landscape at this interface, it is unclear how readily PKR

can mutate to evade viral antagonists while preserving its kinase function.

Here we use a prospective, systematic, and unbiased approach to map SNP-accessible

PKR variants in the kinase domain that are resistant or susceptible to the vaccinia wild-type K3,

nonfunctional K3, and the enhanced mutant allele K3-H47R. We found many novel K3-resistant

variants, especially at sites under positive selection, and note a strong correlation between

variants that resist wild-type K3 and the enhanced allele. We also identified sites in the PKR

kinase domain where variants increased susceptibility to vaccinia K3. Essential PKR residues

that did not tolerate variation were infrequent despite the critical nature of this interface; they

were primarily found surrounding the ATP-binding site. These results paint a portrait of PKR

balancing the tension between conserved function and evasion of viral inhibitors through

mutational resilience in its substrate interface.

RESULTS

Generation of nonsynonymous PKR variant library

We systematically made genetic variants in four windows of interest in the kinase

domain of PKR (Figure 1D). The four windows were selected based on having sites within five

angstroms of eIF2α or K3 and sites under positive selection across vertebrate PKR homologs

as identified by Rothenburg et al. (13). Our windows of interest include helix αD and αG, as well

as the glycine-rich loop and activation loop surrounding the ATP-binding site. In these windows,

we focused on testing 426 variants that could be acquired with a single nucleotide change from

the predominant human PKR allele (Genbank M85294.1) (Supplemental Figure 1A), as those

are the most likely to be evolutionarily sampled.

Overexpression of PKR in the budding yeast *Saccharomyces cerevisiae* is toxic, as PKR

auto-activates and shuts down protein synthesis by phosphorylating yeast eIF2α, which is highly

conserved between yeast and humans (25,26). This toxicity can be partially counteracted by co-

expression of vaccinia K3 (23). Thus, yeast provides a useful experimental system to test the

functionality of PKR variants: variants that maintain the ability to phosphorylate eIF2α while

escaping K3 inhibition will further arrest growth, while variants that lose eIF2α kinase activity or

become more susceptible to K3 will improve growth.

We sought to leverage this assay in a high-throughput manner to determine the

functionality of PKR variants by tracking their abundances in a pool over time. We generated the

426 single-residue variants of PKR on a single-copy plasmid with a galactose-inducible

promoter in a plasmid pool (Figure 1E). Each unique plasmid in the pool carried a DNA barcode

randomly generated during cloning. We used long-read circular consensus sequencing to pair

DNA barcodes with PKR variants, identifying an average of 43 barcodes per variant

(Supplemental Figure 1B). We then created three libraries by combining the PKR variant library

with wild-type K3, nonfunctional K3, or the enhanced mutant allele K3-H47R (23,27). The K3

alleles were expressed from a constitutive *TDH3* promoter. We transformed the three PKR-K3

paired libraries in duplicate into yeast grown in glucose media, such that PKR expression was

repressed. We then inoculated the libraries in liquid galactose media to induce PKR expression,

and sampled cells at 0, 12, 16, and 20 hours post-inoculation. We quantified the abundance of

the PKR variants at each timepoint using high-throughput short-read sequencing of the DNA

barcodes. For each PKR variant we calculated a functional score against each K3: PKR variants

that evade K3 inhibition and reduce yeast growth have high functional scores, while PKR

variants that are non-functional or are inhibited by vaccinia K3, and thus are associated with

good yeast growth, have low functional scores (Supplemental Figure 2). The correlation of

functional scores between replicates 1 and 2 was very strong (Pearson correlation coefficient >

0.98 for each K3 allele, Supplemental Figure 3), so we combined the reads from replicates 1

and 2 to generate PKR functional scores for subsequent analyses.

Many PKR variants evade K3 while maintaining kinase function

We first examined how each PKR variant performed against wild-type K3. We were

encouraged to find that variants previously characterized as resistant to K3-H47R (24) were

resistant to wild-type K3 in our screen (Figure 2A). We found many additional K3-resistant

variants across all four tested windows and validated select PKR variants using serial dilution to

measure yeast growth (Supplemental Figure 4). Notably, every variant made at Glu375

increased evasion of K3, with the exception of Glu375Asp. In the AlphaFold2-predicted

complex, Glu375 forms an attractive electrostatic interaction with Lys45 of K3 (Figure 2B).

Consistent with our findings, this predicted interaction would be disrupted by all tested variants

at site 375 aside from the negatively charged aspartic acid. Interestingly, Glu375 is the only site

under positive selection in vertebrates that was proximal to K3 but not eIF2α in the AlphaFold2

models, making it a prime candidate for disrupting K3 inhibition without affecting eIF2α

phosphorylation.

We found a large number of K3-resistant variants in the vicinity of PKR's helix αG

(Figure 2C,D). Compared to other protein kinase structures, eIF2α kinases have a noncanonical

helix αG that is extended by one full turn and rotated 40° counterclockwise with a 5 angstrom

translation relative to its C-terminus, owing to a reduced αF-αG linker (5). The noncanonical

orientation of helix αG contributes to both eIF2α binding specificity and Ser51 presentation to

PKR (5). It has been proposed that mutations causing movement in helix αG might be especially

capable of disrupting K3 binding due to the rigid structure of K3 between its contact points on

helix αG and the PKR active site. In contrast, PKR mutations that move helix αG could be

tolerated in the context of eIF2α binding because eIF2α contacts PKR's active site using a

flexible loop (5). We observe a gradient of PKR functional scores along the helix αG region,

starting with highly resistant or susceptible variants (positions 480-492) and shifting to neutral

wild-type-like scores (493-500) (Figure 2A). We identified four sites within or near the N-

terminus of helix αG for which many variants evade K3: Glu480, Val484, Glu490, and Thr491

(Figure 2C).

We looked at what features of PKR were associated with sites of K3-resistant variants

and were excited to find that sites under positive selection in vertebrates were strongly enriched

for K3-resistant variants relative to sites conserved across vertebrate PKRs (Figure 2E).

Comparing K3 contact and non-contact sites, we find significantly lower functional scores at K3

contact sites (Figure 2F). However, functional scores at K3 contact sites form a bimodal

distribution (Hartigan's dip test statistic = 0.105, p < 0.0001). Considering only functional

variants (PKR functional score > -2) we find that variants at K3 contact sites have significantly

higher functional scores than those at non-contact sites (Supplemental Figure 5). Thus, variants

at K3 contact sites tend towards large effects either improving or diminishing PKR function

versus K3.

Sampled sites in the PKR kinase domain are largely resilient to genetic variation

To understand the constraint placed on the PKR kinase domain, we examined variants

that failed to maintain PKR function. We screened our PKR variant library against a

nonfunctional K3 (hereafter referred to as K3Δ58) in which the first 173 bases were deleted,

removing the first 58 residues, including the initial methionine. When paired with K3Δ58, most

PKR variants were comparable to wild-type PKR (Figure 3A), suggesting the tested regions

were not highly functionally constrained for kinase activity despite their proximity to eIF2α during

its phosphorylation. The few highly constrained sites include Gly276 and Gly277, which

compose the glycine-rich loop above the ATP-binding site; Gly450 and Thr451 in the activation

loop; and Thr487 in helix αG, which stabilizes helix αG (5) and induces the conformational

change in eIF2α required for its phosphorylation (Figure 3B) (28). We also found proline variants

to generally disrupt PKR function (Figure 3C). We found nonfunctional variants of varying

degrees in all four windows of interest, but with very few in helix αD and the C-terminus of helix

αG (Figure 3C), consistent with these regions having pliable secondary structures in which

variants can evade K3 while not disrupting eIF2α recognition.

More PKR variants had low functional scores in the presence of functional K3 (Figure

2A) than in its absence (Figure 3C). We plotted the functional scores against each other (Figure

4A), and found that variants with low PKR functional scores versus K3 separated into two

classes: some switched to high functional scores in the absence of K3, indicating they caused

enhanced susceptibility to K3 but otherwise retained PKR function, while others had low

functional scores in both conditions, indicating they had diminished eIF2α kinase activity

irrespective of K3 antagonism. We quantified this effect by calculating residuals from the line

connecting PKR nonsense variants to PKR-WT, which highlights variants whose functional

scores diverged between the K3 conditions. We find that the activation loop and the start of

helix αG, which were less tolerant of mutation when paired with wild-type K3 (Figure 2D),

contain a mix of variants that lose function outright or that increase susceptibility to K3 (Figures

3B-C, 4B-C). Variants at five sites were especially susceptible to K3 inhibition: Phe278, Thr373,

Arg453, Phe489, and Ser492 (Figure 4B). None appear under positive selection across

vertebrates, and Phe278 is conserved across vertebrate PKR homologs. Interestingly, Phe278

is predicted to form a pi interaction with His47 of K3, which aligns to Arg52 of eIF2α

(Supplemental Figure 6) (23). Mutation of K3-H47 to arginine, matching eIF2α, is known to

improve K3's antagonism of PKR (23,27), suggesting PKR can discriminate against K3 based

on its difference from eIF2α at this position. This is consistent with our observation of the

importance of Phe278 specifically in the presence of K3.

K3-resistant variants were largely resistant to enhanced K3-H47R

We also characterized PKR variants in the presence of an enhanced mutant allele of K3,

K3-H47R (Figure 5). This mutant allele was first identified in a genetic screen of K3 mutants as

an enhanced antagonist of PKR (23) and independently identified in a directed evolution screen

(27). We sought to understand the resilience of improved PKR variants to alternative K3 alleles

by examining the concordance between beneficial PKR variants against wild-type K3 and K3-

H47R. We found strong correlation between PKR functional scores versus K3 and K3-H47R

(Spearman correlation = 0.832, Figure 5A). Indeed, many of the K3-resistant variants are also

resistant to K3-H47R (Figure 5B-D), suggesting resilience of PKR to allelic variation in K3. A

general change is the struggle for PKR to overcome the enhanced inhibition of K3-H47R,

compressing the functional score range between wild-type-like and nonfunctional scores. This

nonlinear relationship between the functional scores in the two K3 conditions (Supplemental

Figure 7) led to some differences in the patterns of resistance to the two K3 alleles. For

instance, positively selected PKR sites did not have significantly higher PKR functional scores

versus K3-H47R than conserved sites (Figure 5E), a sharp contrast to the PKR functional

scores versus wild-type K3 (Figure 2E). This result appears to be caused by the many wild-type-

like scores at positively selected sites now being similar to the many nonsense-like scores at

conserved sites. We also noted that variants at sites proximal to K3 had slightly higher

functional scores than those at non-contact sites (Figure 5F), the opposite of what we observed

against wild-type K3 (Figure 2F). This could result from wild-type-like PKR variants being

enriched at non-contact sites. Overall, the strong correlation of functional scores versus K3 and

K3-H47R suggests K3-resistant variants would often be resilient against alternate K3 alleles.

Previous work identified twelve PKR variants with enhanced resistance to K3-H47R (24),

nine of which were included in our experiment (Supplemental Figure 8). We examined the other

nonsynonymous variants introduced at these sites and found that most sites contained

additional novel resistant variants, indicating the previously uncovered individual mutants reflect

sites of general opportunity for pseudosubstrate evasion.

DISCUSSION

Here we characterize the local evolutionary space available to the human innate

immunity protein PKR to maintain functionality while evading the well-studied vaccinia virus

antagonist, K3. One strategy viruses leverage against PKR is pseudosubstrate inhibition, in

which proteins like K3 bind PKR by structurally mimicking its natural substrate eIF2α, inhibiting

PKR kinase activity. Interestingly, this binding interface contains many residues under positive

selection. However, positively selected genetic variants across the binding interface would need

to maintain binding with PKR's natural substrate while evading viral pseudosubstrate

antagonists. In this study, we quantified the abundance of such "tightrope-walking" genetic

variants relative to the abundance of functionally deleterious variants, which highlighted

particular sites of opportunity and constraint within the kinase domain.

We identified many sites containing variants that evade vaccinia K3 without losing eIF2α

targeting. Resistant variants clustered around helices αD and αG, secondary structures with

sites under positive selection and where resistant variants have been previously characterized

(13,24). Overall, we found few genetic variants that impair the kinase function of PKR (Figure 3),

underscoring the genetic resilience of PKR, with functionally constrained sites mostly localized

to the vicinity of the ATP-binding site. Finally, we found a strong correlation between PKR

variants that are resistant against both K3 and the enhanced antagonist K3-H47R, suggesting

the K3-resistant variants are resilient to K3 allelic variation.

We identified sites across the PKR kinase domain that were K3-resistant, K3-

susceptible, or essential to eIF2α phosphorylation (Figure 6A). Sites enriched with genetic

variants that improve evasion of K3 were classified as K3-resistant sites, such as Glu375 and

Glu480. We find clusters of these sites located across helices αD and αG, at the binding surface

of K3 and eIF2α. Positive selection sites are enriched for K3-resistant variants. There were sites

under positive selection where we did not observe much K3 resistance, such as sites 269-272,

which may suggest this region is antagonized by a completely different viral antagonist. Still, we

found these sites to be generally able to adopt variants that maintain PKR function (Figure 3),

making them prime "tightrope-walking" sites for thwarting any alternate viral inhibitors that

interact at those sites. We also identified some K3-resistant sites that are not under positive

selection, such as Glu480, which is in fact conserved across vertebrate PKR homologs and the

four human eIF2α kinases (9,13). Perhaps it has an important role in a function not tested in our

assay, such as avoiding off-target phosphorylation of other human proteins.

247   Our analysis does not account for variation in PKR expression that may result from the

248   introduced variation. Previous work demonstrated that K3-resistant PKR variants were not

249   enhanced due to increased expression (13). We note that we did not observe much variation in

250   PKR functional scores in the absence of K3 inhibition (Figure 3A), which suggests PKR variants

251   did not often substantially modulate inherent properties of PKR, like its expression.

252   We found PKR variants that were not tolerated, though perhaps fewer than we would

expect given the kinase domain is responsible for carrying out its critical function. These came

in two flavors: variants that increased susceptibility to K3, and variants that were deleterious for

PKR function. We were surprised to find sites enriched for K3-susceptible variants, such as

Phe278, Arg453, and Ser492. These could be sites where the wild-type residues help

discriminate between eIF2α and K3. These sites do not cluster together; however, when

projected onto the surface of PKR they appear to be intermingled with essential residues

(Figure 6A). This could suggest that while they are not needed to recognize eIF2α in normal

conditions, they are well-positioned to discriminate between eIF2α and a pseudosubstrate

inhibitor. Finally, the correlation of PKR functional scores between K3 and K3-H47R suggests

resistant variants are resilient. This agrees with prior work screening a few PKR variants against

vaccinia K3-H47R and the K3 homolog from variola virus (13). We would be curious to know if

this pattern holds for more highly diverged K3 orthologs (29) and other pseudosubstrate

inhibitors, such as Ranavirus vIF2α (30).

The genetic pliability observed in the PKR kinase domain is reminiscent of that seen with

another innate immunity factor, TRIM5α, which is locked in an evolutionary arms race with

retroviral capsids (31,32). Both PKR and TRIM5α have access to "rolling hills" of resistant

variants (Figure 6B), in opposition to sharp cliffs that are highly optimized (32). Notably, unlike

PKR, TRIM5α does not have to balance pathogen conflict with enzymatic function. We also note

the structural context in which we identified K3-resistant sites, with many being found within

secondary structures (e.g. Glu375, Il378, and Glu379 in helix αD, and Glu490, Lys493, and

Thr491 in helix αG), whereas in TRIM5α, as well as the immune proteins MxA and NLRP1

(33,34), positively selected sites and improved variants have been described in structurally

flexible loops and linkers. We also considered contrasts between PKR and ACE2, a host

enzyme with sites under positive selection likely due to being targeted by coronaviruses for cell

entry (35–37). However, the ACE2 sites under positive selection are not located near the

catalytic domain of the enzyme, in contrast to PKR (38–40). Ultimately, it appears that the

resolution of this conundrum is that PKR maintains pliability in this region despite its criticality.

This pliability is likely made possible by the flexibility of eIF2α's Ser51 loop in which the

phosphorylation site resides; indeed, the loop is unstructured in the co-crystal of PKR and eIF2α

(5). As such, while we still don't have a clear picture of what phosphorylation of eIF2α looks like,

we can appreciate the flexible nature of eIF2α during its interaction with PKR (5,14,24,28).

Further biochemical and structural investigation will be necessary to interrogate the ability of

variants in PKR to disrupt interactions with K3 while maintaining eIF2α targeting. We note that

while we were able to identify many sites of interest across the PKR kinase domain, variation at

many other sites in PKR remains to be explored and may harbor allosteric effects on

pseudosubstrate resistance.

Barring possible variation in under-studied populations, there is little allelic diversity of

PKR in the human population, with all nonsynonymous variation listed as rare in gnomAD (41).

PKR does exhibit sequence diversity in comparisons across species, including primates (14,42).

We expect viruses could exert selective pressure on rare heterozygous variation in PKR, as we

would expect beneficial variants in PKR that evade pseudosubstrate inhibitors to be dominant,

given that a small amount of eIF2α phosphorylation is capable of halting protein synthesis (43).

We found an enrichment for vaccinia K3-resistant variants at sites previously found to be

under positive selection (13,14). In effect, these are two independent approaches that both

pinpoint these sites as genetically pliable: positive selection analysis retrospectively considers

existing variation in nature across multiple homologs, whereas deep mutational scanning

prospectively tests novel variants in a single homolog. We and others (32,33) view deep

mutational scanning experiments as a complementary approach to positive selection analysis.

One strength of deep mutational scanning is its potential to illuminate sites of opportunity

against pathogens that have not exerted historical evolutionary pressure. It can also allow for

the testing of insertions or deletions (44) which are generally discarded in positive selection

analysis. On the other hand, a limitation of the deep mutational scanning approach is that it only

tests variant function in a limited set of conditions, when the evolutionary selection could be a

symphonic signal made up of a myriad of unknown pressures. Deep mutational scanning also

requires the function of the protein to be testable in a selectable assay. While proteins with cell-

extrinsic or whole-organism phenotypes would be harder to test, recent advances in high-

throughput genetic experimentation may allow deep mutational scans of those proteins as well

(45,46).

IDEAS AND SPECULATION

One might naively expect a PKR pseudosubstrate inhibitor to be a closer mimic of the

natural PKR substrate, eIF2α, which would make it more difficult for PKR to distinguish it from

eIF2α. One possibility is that K3 pseudosubstrate divergence from eIF2α has been selected to

allow the inhibitor to bind PKR with even higher affinity than eIF2α does. But mutations that

improve binding to one species' PKR homolog might decrease binding to the divergent PKR

homologs from other species (47–49). Thus, pseudosubstrate inhibitors of generalist poxviruses

may need to evolutionarily balance improving PKR binding for one host against maintaining

PKR binding of multiple hosts. Interestingly, the structural component most diverged between

K3 and eIF2α, a rigid helix in K3 in place of eIF2α's flexible phospho-acceptor site, primarily

interacts with the residues around the activation loop of PKR (Figure 6C), which we found to be

highly functionally constrained. In contrast, amino acids 72-83 of K3 are highly identical to eIF2α

and interact primarily with PKR's divergent helix αG (Supplemental Figure 6). Thus, vaccinia K3

may well have been evolutionarily constrained to avoid limiting host range. While the host range

of vaccinia is not fully understood (50), current evidence suggests vaccinia is a generalist

(29,48). Poxviruses as a whole infect a very wide range of species, which suggests the

ancestral K3 needed to be compatible with a wide host range.

Despite its pliability, PKR helix αG is believed to play a critical role in eIF2α binding and

phosphorylation, as it induces a conformational shift in eIF2α that moves the Ser51 phospho-

acceptor site into PKR's ATP-binding site. Based on our findings and positive selection

analyses, this conformational shift is apparently amenable to variation in the PKR kinase

domain. Perhaps the resilience of the binding surface between PKR and eIF2α could be

informative for the budding field of de novo protein design, as resilient design is a critical

consideration for vaccines and therapeutics faced with rapidly evolving infections agents.

MATERIALS AND METHODS

PKR and K3 plasmid construction

We generated the plasmid MSp508_MCS as the base plasmid for all experiments by inserting a multiple-cloning site (MCS) into the single-copy plasmid YCp50, a gift from Mark Rose (51). The MCS contained seven unique restriction enzyme cut sites: AgeI, BsiWI, BstEII, NotI, MluI, PvuII, and SacI. YCp50 was digested using restriction enzymes EcoRI-HF (NEB Cat#R3101S) and BspDI (NEB Cat#R0557S) with rCutSmart buffer, followed by purification and size selection on a 1% agarose gel with 0.6 µg/mL ethidium bromide (BioRad Cat#1610433), selecting for the 7,961 bp band. The extracted band was purified using a QIAquick Gel Extraction kit (Qiagen Cat#28706) and eluted with 30 µL of water. For Gibson Assembly the purified band was combined with a single-stranded DNA oligonucleotide containing the MSC and homology arms spanning the digest site, named Oligo 1 (Supplemental File), in a 1:5 molar ratio in a total volume of 5 µL, along with an additional 5 µL 2x Gibson Assembly Master Mix (NEB Cat#E2611L). The reaction mixture was incubated at 50°C for 1 hour. One microliter of the Gibson Assembly reaction was transformed into 10 µL 5-alpha competent E. coli (NEB Cat#C2987I) and plated onto 10-cm LB-AMP plates (IPM Scientific Cat#11006-016) and incubated overnight at 37°C. Colonies were selected for overnight outgrowth in 3 mL LB-AMP (IPM Scientific Cat#11006-004), followed by a QIAprep spin miniprep (Qiagen Cat#27106) and Sanger sequencing to validate the plasmid sequence using the primer Oligo 2. This produced the plasmid MSp508_MCS.

We cloned wild-type PKR, wild-type K3, and K3-H47R into the MCS of MSp508_MCS to produce the MSp509_PKR-WT, MSp510_K3-WT, and MSp511_K3-H47R plasmids, respectively. The plasmid p1419 (23) contains the *PKR* gene (standard name *EIF2AK2)* under the control of the galactose-inducible pGAL10/CYC1 promoter for yeast expression. We PCR-amplified the PKR expression construct using primers (Oligos 3 and 4) to place pGAL10/CYC1-*PKR* on MSp508_MCS between AgeI and BstEII. Oligo 4 incorporated a synthetic terminator for

PKR, T$_{synth1}$ (52), a 28-bp spacer sequence, and a 26-bp barcode

(CGCTTAATATGCAATGAAATTGCTTA); the primers added 20-bp homology arms for cloning

onto the MSp508_MCS plasmid. PCR was performed using the polymerase PfuUltraII (Agilent

Technologies Cat#600674) for 30 cycles with an annealing temperature of 55°C, followed by gel

purification. MSp508_MCS was digested with restriction enzymes AgeI-HF (NEB Cat#R3552S)

and BstEII-HF (NEB Cat#R3162S), followed by gel purification. The digested MSp508_MCS

and pGAL10/CYC1-PKR fragments were joined via Gibson Assembly in a 1:5 molar ratio,

followed by transformation into 5-alpha competent *E. coli* and plating onto LB-AMP. A resultant

clone, validated by Sanger sequencing, was designated MSp509_PKR-WT.

To produce MSp510_K3-WT, the *K3L* coding sequence was amplified from the plasmid

pC140 (23) using primers Oligo 5 and Oligo 6. Similarly, to produce MSp511_K3-H47R, the

K3L-H47R coding sequence was amplified from the plasmid pC407 using primers Oligo 6 and

Oligo 7. pC140 and pC407 were gifts from Thomas Dever. We note that the *K3L* alleles on

these plasmids carry a Val2Leu mutation relative to the VACV-WR *K3L* (Genbank

AAO89313.1), which was introduced during the cloning of *K3L* on the original plasmid pTM1

(19). Oligos 5 and 7 incorporated a synthetic terminator (T$_{synth8}$) for the K3 allele (52), a 28-bp

nucleotide spacer sequence, a 26-bp barcode (ATCGTAATAGGTTTCCGGCTTGTTCG and

ACAGGAAATGCTTTCGGGGTTGTATT, respectively), and a 20-bp homology arm to

MSp508_MCS. The pTDH3 promoter, also known as pGPD, was amplified from genomic DNA

of the yeast strain BY4742 (53) using primers Oligo 8 and Oligo 9). K3-WT, K3-H47R, and

pTDH3 were amplified using PfuUltraII, followed by gel purification. MSp508_MCS was digested

with restriction enzymes NotI-HF (NEB Cat#R3189S) and PvuII-HF (NEB Cat#R3151S). Each

K3 allele (K3-WT and K3-H47R) was joined with a pTDH3 promoter onto the digested

MSp508_MCS via Gibson Assembly in a 1:5:5 molar ratio, followed by transformation into 5-

alpha competent *E. coli* and plating onto LB-AMP. Resultant clones, validated by Sanger

sequencing, were designated as MSp510_K3-WT and MSp511_K3-H47R.

MSp512_K3Δ58 was made by removing nucleotides 1-173 of the *K3L* sequence of

YCp50_K3, leaving codons 59-88 to generate a nonfunctional *K3L* allele (denoted K3Δ58)

under the pTDH3 promoter. The K3Δ58 sequence was amplified from the MSp510_K3-WT

plasmid using primers Oligo 10 and Oligo 11. MSp510_K3-WT was digested with restriction

enzymes NotI-HF and MluI-HF (NEB Cat#R3198S), followed by gel purification and size

selection for the 8,684 bp band. The K3Δ58 allele was joined with the MSp508_MCS digested

fragment via Gibson Assembly in a 1:5 molar ratio, followed by transformation into 5-alpha

competent E. coli and plating onto LB-AMP. A resultant clone was validated by Sanger

sequencing and designated as MSp512_K3Δ58.

Generation of the *PKR* variant library

We made the *PKR* variant library by 15 Gibson Assembly reactions using pairs of PCR

amplicons targeting 4 windows of PKR protein sequence: 255-278 (5 pairs of reactions), 371-

385 (3 pairs of reactions), 448-455 (2 pairs of reactions), and 480-506 (5 pairs of reactions). In

each pair, one of the PCR reactions, PCR-1, utilized a pool of forward primers, termed a variant

primer tile set, containing single-base deviations from the *PKR* sequence to generate all SNP-

accessible variants in a stretch of X to Y consecutive codons (Supplemental Figure 9). Each

forward primer in a tile set was composed of a 20-bp homology arm, the variant region, and a

priming region with a melting temperature of 50°C. The reverse primer of all PCR-1 reactions

had a constant 20-bp homology arm, a barcode region with 20 fully mixed bases, and a

constant PCR-priming region homologous to the end of *PKR* with a melting temperature of 55°C

(Supplemental Figure 10). Mixed bases in the barcodes were limited to 5-nucleotide stretches to

prevent the unintended generation of restriction sites, spacing these stretches with "AA" or "TT"

sequences (54). The forward PCR-1 primer tile sets are listed in the Supplemental File as

Oligos 12-26, and the reverse PCR-1 barcode primer is listed as Oligo 27. The paired PCR-2

reaction was an inverse PCR that generated the remainder of the *PKR* gene and plasmid

backbone, with overlaps to the homology arms of PCR-1; the forward primer for PCR-2

reactions was Oligo 28, and the reverse primers were Oligos 29-43. Thus, Gibson Assembly of

these PCR products would generate sets of *PKR* variants in defined segments of *PKR* along

with random barcodes at the end of the gene (Supplemental Figure 11).

We used custom Python (v3.8.18) scripts to design each variant-introducing primer,

utilizing IUPAC degenerate base symbols (e.g. "D" represents a mix of "A", "G", and "T")

(55,56). The introduced variant amino acids were all within a single base change from the

canonical human *PKR* coding sequence (Genbank M85294.1:31-1686); we note the *PKR*

sequence encoded on MSp509_PKR-WT differs somewhat from the canonical sequence

through synonymous substitutions. Primers were synthesized by Integrated DNA Technologies,

then were pooled manually into variant primer tiles sets.

PCR-1 reactions were amplified from the MSp509_PKR-WT plasmid using PfuUltraII

polymerase with an annealing temperature of 45°C. Each reaction generated a pool of *PKR*

gene fragments containing a single nonsynonymous variant paired with a unique barcode

sequence. In parallel, PCR-2 reactions amplified vector fragments from MSp509_PKR-WT

plasmid using Herculase II polymerase (Agilent Technologies Cat#600677) using cycling

conditions for vector targets >10 kb. PCR-1 and PCR-2 amplicons were digested with DpnI

(NEB Cat#R0176S) for 2 hours at 37°C to remove the MSp509_PKR-WT template DNA,

followed by gel purification. Each PCR-1 amplicon was paired with its corresponding PCR-2

amplicon for Gibson Assembly in a 1:5 molar ratio.

Two μL of each Gibson Assembly was transformed into 50 μL of 5-alpha competent *E.*

coli and all cells were plated on LB-AMP and incubated overnight at 37°C. Colonies were

counted on each plate and plates were bottlenecked at approximately 30 colonies per selected

nonsynonymous variant (i.e. each nonsynonymous variant would be linked to approximately 30

unique barcode sequences). Plates were washed with 15 mL LB-AMP, cell densities were

measured as the optical density at 600 nm (OD_{600}), and an equal number of cells (approximately

5×10^9 cells) from each reaction were pooled to form a single *PKR* variant library. A 200 mL LB-

AMP outgrowth was grown to an OD_{600} measurement of approximately 3, followed by a Qiagen

MAXI plasmid prep (Qiagen Cat#12963).

PacBio sequencing of barcoded variant libraries

The *PKR* variant library was sequenced using the PacBio Sequel II instrument to identify

the barcodes that were linked to each nonsynonymous variant. Sixteen micrograms of the *PKR*

variant library were digested with restriction enzymes AgeI-HF and NotI-HF to create linear

fragments of approximately 2,100 bp for sequencing. The digest was incubated for 2 hours at

37°C then purified using a QiaQuick PCR purification kit (Qiagen Cat#28106). Fragments of

2,100 bp were size-selected using a SageELF instrument (Sage Science) before PacBio circular

consensus sequencing (CCS) HiFi sequencing on a Sequel II instrument (Pacific Biosciences).

From the CCS reads we generated a table of *PKR* barcodes paired with genetic variants using

alignparse v0.2.6 and custom Python scripts (57).

Combining *PKR* variant library with K3 alleles

We next cloned the three K3 alleles (K3, K3-H47R, and K3Δ58) onto the PKR variant

library plasmids. The PKR variant library was digested with NotI-HF and PvuII-HF to generate a

10,270 bp receiver DNA fragment, while the K3 alleles were separately digested with BstEII-HF

and SacI-HF (NEB Cat#R3156S) to generate a 1,130 bp insert DNA fragment with 20-bp

homology arms with homology to the receiver fragment for Gibson Assembly. For each of the

reactions, 10 µg of plasmid DNA was digested with 5 µL of each restriction enzyme in a 50 µL

reaction to generate a linear fragment, followed by gel purification. The PKR and K3 fragments

were combined in three separate Gibson Assembly reactions in a 1:5 molar ratio with 100 ng of

the PKR vector fragment in a reaction volume of 20 µL and incubated at 50°C for 1 hour. A

standard ethanol precipitation was performed on each reaction, adding 100 µL of 100% ETOH

and incubating overnight at -20°C before resuspending the pellets in 2 µL of water (58).

We transformed each of the three concentrated reactions into *E. cloni* 10G SUPREME

electrocompetent cells (Lucigen Cat#60080-2) using a MicroPulser Electroporator (BioRad),

with 2 µL of the concentrated reaction combined with 25 µL of competent cells. Following the

electroporation and recovery, cells were plated onto two 15-cm LB-AMP plates. A 1:1000

dilution was plated with 100 µL of water on a 10-cm plate to estimate the number of transformed

colonies across the two 15-cm plates, which produced an estimate of approximately 2 million

colonies per electroporation. Plates were washed with 25 mL LB-AMP, combining the two plates

for each of the reactions. Each library was expanded in 200 mL LB-AMP to an OD_{600}

measurement of approximately 3, followed by a Qiagen MAXI plasmid prep.

Screening the PKR-K3 variant pairs in a yeast growth assay

All yeast growth was at 30°C in CSM-Ura (Sunrise Science Products Cat#1004-010),

and liquid cultures were shaken at 200 revolutions per minute. We transformed each of the

three paired libraries (PKR + K3, PKR + K3Δ58, and PKR + K3-H47R) into yeast in duplicate.

The paired plasmid libraries were transformed into the yeast strain BY4742 (*MATα ura3Δ0*

leu2Δ0 his3Δ1 lys2Δ0; strain name MSY2) (53) using a standard large-scale high efficiency

lithium acetate transformation protocol (59) and plated onto 15-cm plates with 2% dextrose,

using 1:1,000 and 1:10,000 dilutions on 10-cm plates to estimate colony counts, which

produced approximately 100,000 colonies. Colonies were washed off plates with 25 mL media

with 2% dextrose.

To start the yeast growth assay, six cultures were seeded at low density (OD_{600}

measurement of 0.01) in 40 mL media with 2% dextrose and grown overnight for 16 hours. The

following morning cultures were moved from 30°C to 4°C to pause growth, then restarted at an

OD_{600} of 0.25 in 40 mL media with 2% dextrose 3 hours prior to GAL induction. After 3 hours,

once all cultures were in a log growth phase, approximately 1×10^8 cells were taken from each

culture as the starting timepoint sample (designated as "0 hours"). All timepoint samples were

spun down in a 1.5 mL Eppendorf tube at 5,000 rpm for 1 minute, after which supernatant was

removed and the pellet was frozen at -80°C. Approximately 2×10^8 cells from the remaining

cultures were pelleted in 50 mL conical tubes at 3,000 rpm, supernatant was removed, and cells

were resuspended in 80 mL CSM-Ura media with 2% galactose (OD_{600} measurement of 0.125

OD_{600}) to induce PKR expression. Cultures were incubated overnight at 30°C, 200 rpm.

Additional samples were harvested at 12, 16, and 20 hours post-PKR induction, with

approximately 1×10^8 cells taken at each timepoint. The culture density was monitored and

back diluted to maintain a log growth phase (OD_{600} measurement less than 1). With three K3

allele conditions (K3, K3Δ58, and K3-H47R), four timepoints (0, 12, 16, and 20 hours) and two

replicates, a total of 24 samples were taken across the yeast growth assay.

Plasmid extraction and barcode amplification

To quantify changes in barcode abundance between timepoints in the yeast growth

assay, we harvested plasmids from the four timepoints for each K3 allele, then amplified and

sequenced the adjacent PKR and K3 barcodes in each plasmid. Plasmids were harvested from

the yeast samples using a modified QIAprep spin miniprep protocol (60). Sampled cell pellets

were first thawed at room temperature, followed by the addition of 250 μL QIAprep P1 buffer

and 2 μL zymolyase (2.5 units per μL), and incubation at 37°C for 30 minutes, followed by

adding 250 μL P2 buffer and following the manufacturer instructions for the remainder of the

protocol.

To identify and quantify the abundance of each PKR variant paired with each of the

three K3 alleles we performed Illumina sequencing of the PKR barcodes from the plasmids. We

used a two-step PCR protocol, the first to amplify the PKR barcode from the plasmids and the

second to attach Illumina adapters and Nextera indices for downstream sequencing and sample

demultiplexing. For the first PCR reaction, we designed forward and reverse primers as a pool

of five primers each, that had 0 to 4 "N" bases between the Nextera transposase adapter

sequence and the priming sequence to stagger the base signal per cycle and maintain

sequence diversity across the flow cell (61). The forward primers (Oligos 44-48) with a melting

temperature of 60°C were pooled together and used for all PCR-1 reactions, annealing

immediately upstream of the PKR barcode locus. Five reverse primers were designed and

pooled for amplification from each of the three K3 alleles that annealed within the three distinct

K3 barcode sequences (Oligos 49-53 for K3, Oligos 54-58 for K3Δ58, and Oligos 59-63 for K3-

H47R). One microliter of each primer pool was combined with 50 ng of sample plasmid DNA

and 25 µL Kapa Hifi Hotstart ReadyMix (Kapa Biosystems Cat# KK2602), and topped off to a

total volume of 50 µL with water. PCR cycling was performed as follows: (1) 95°C for 3 minutes,

(2) 98°C for 20 seconds, (3) 65°C for 15 seconds, (4) 72°C for 1 minute, repeat steps 2-4 for a

total of 18 cycles, (5) 72°C for 1 minute, (6) 12°C hold. PCR products were purified using a

MinElute PCR purification kit (Qiagen Cat#28006) using 11 µL water for the final elution, then

quantified using a Qubit dsDNA High Sensitivity (HS) kit (ThermoFisher Scientific Cat#Q32851).

For the second PCR reaction, 10 ng of amplicon DNA from the first PCR reaction was

combined with 10 µL of Nextera adapter index primers (Illumina Cat#20027213) and 25 µL

Kapa HiFi Hotstart ReadyMix, and topped off to a total volume of 50 µL with water. PCR cycling

was performed as follows: (1) 100°C for 45 seconds, (2) 100°C for 15 seconds, (3) 60°C for 30

seconds, (4) 72°C for 30 seconds, repeat steps 2-4 for a total of 8 cycles, (5) 72°C for 1 minute,

(5) 12°C hold. PCR products were purified using a MinElute PCR purification kit using 11 µL of

water for the final elution, then quantified using a Qubit dsDNA Broad Range (BR) kit

(ThermoFisher Scientific Cat#Q32850). 500 ng of the second PCR reaction amplicons were gel

purified on a Size Select II E-Gel (ThermoFisher Scientific Cat#G661012) for approximately 13

minutes to extract the approximately 250-bp amplicon band, followed by quantification with a

Qubit dsDNA HS kit. Amplicon samples were diluted to 4 nM before being pooled together,

followed by manufacture denature and dilution protocols (Illumina Document#15039740v10)

before sequencing on an Illumina NextSeq 2000 instrument.

PKR functional scores and screening analysis

Next, we extracted PKR barcode sequences from the Illumina reads and mapped the

barcodes back to their corresponding select nonsynonymous variants using the table of PKR

barcodes paired with genetic variants generated from the PacBio CCS HiFi reads. Paired reads

were assembled into contiguous sequences using PEAR v0.9.11 (62), followed by Bartender

v1.1 (63) to extract and cluster PKR barcodes using the barcode search pattern "CAAGG[25-

27]GGTGA".

We wrote Python v3.8.18 scripts to tally the PKR barcodes and map them back to

genetic variants using the table generated from the PacBio CCS HiFi reads. Barcode counts

were normalized to the total read count for each of the four timepoints across each of the K3

alleles, followed by a fold change calculation for each PKR barcode across timepoints 0, 12, 16,

and 20 hours:

558
$$FC_{TP} = -log_2(\frac{Normalized\ Reads_{TP}}{Normalized\ Reads_0})$$

where FC_{TP} is the fold change in normalized reads for a PKR barcode from 0 hours to a

subsequent timepoint (TP), $Normalized\ Reads_{TP}$ is the read count for a given barcode divided

by the total number of reads at the given timepoint, and $Normalized\ Reads_0$ is the read count for

a given barcode divided by the total number of reads at the 0-hour timepoint. Of note, the log_2 of

the fold change is inverted such that functional PKR variants that inhibit yeast growth and

decreased in abundance were assigned positive fold change value. We then calculated PKR

functional scores for each PKR barcode in each of the K3 alleles by calculating the area under

the curve using the composite trapezoidal rule. We calculated the PKR functional score for each

variant for each K3 allele by averaging the PKR functional score across all representative

barcodes for a given variant. As we found a strong correlation between replicates (Pearson

correlation coefficient > 0.98 for each K3 allele, Supplemental Figure 3), we proceeded by

combining all reads from the two replicate experiments for each K3 allele and recalculating the

fold changes and PKR functional scores for each barcode as described above.

Predicted PKR complexes and substrate contacts

All molecular graphics and analyses were performed using USCF ChimeraX v1.5 and

PyMol v2.5.4. To define PKR residues contacting K3 and eIF2α we used the AlphaFold2

structure prediction tool (ColabFold v1.5.5) in UCSF ChimeraX v1.5 (64,65) (Supplemental

Figure 12A). We aligned these predictions to existing crystal structures (PDB 2A1A and 1LUZ)

and found the AlphaFold2 predictions to largely represent the crystal structures (RMSD < 1

angstrom for both alignments, Supplemental Figure 12B-D) (5,22). We note the low confidence

score (pLDDT < 50) of the eIF2α flexible loop containing Ser51, which adopts a compact helical

conformation in the predicted model and closely matches the conformation observed in the

crystal structure of unbound yeast eIF2α (PDB 1Q46) (66). As Ser51 of eIF2α would need to

travel approximately 17 Å to reach PKR's active site (5), this predicted structure is likely

reflective of a pre-phosphorylation complex, and thus would not capture contacts between PKR

and eIF2α made during phosphorylation. The corresponding loop is unresolved in the crystal

structure of human PKR in complex with eIF2α (PDB 2A1A). Note that we opted not to use the

existing structure for defining contact residues, both for consistency with the K3 contact

definitions, as there is no PKR-K3 co-crystal, as well as for the potential to capture interactions

mediated by residues unresolved in the crystal structure (PDB 2A1A, PKR Asp338-Asn350 and

eIF2α Glu49-Lys60).

To identify potential PKR contact sites from the predicted complex structures, we used

PyMol v2.5.4 to select all PKR residue branches within 5 Å of K3 or eIF2α using the command:

"sele contacts, br. /{pdb file name}//A within 5 of /{pdb file name}//B" with chain "A" being PKR

and chain "B" being either K3 or eIF2α. Predicted PKR sites that contact K3 were: 275, 276,

278, 304, 339, 343, 345, 375, 379, 382, 414, 416, 435, 448, 449, 450, 451, 452, 453, 455, 460,

485, 486, 487, 488, 489, 490, 492, 493, and 496. Predicted PKR sites that contact eIF2α were:

274, 275, 276, 279, 335, 337, 338, 339, 340, 341, 342, 379, 382, 451, 452, 453, 483, 486, 487,

488, 489, 490, 491, 492, and 493.

PKR sites in the kinase domain under positive selection across vertebrate species were

identified by Rothenburg et al. (13): 261, 269-272, 307, 314, 322, 360, 368, 375, 378, 379, 382,

385, 389, 394, 405, 428, 448, 449, 462, 471, 483, 486, 488, 491, 493, 500, 502, 504, 505, 514,

520, and 524. Sites conserved across vertebrate PKR homologs were also identified by

Rothenburg et al. (13) which were used for supplementary analysis: 263, 267, 276-279, 281,

283, 296, 298, 308, 309, 312, 315, 317, 319, 320, 323, 327, 362, 364-367, 369, 374, 377, 397,

401, 404, 406, 407, 410-417, 419, 420, 429-433, 437, 444-446, 450, 451, 454, 455, 457-459,

465, 469, 470, 474-477, 480, 490, 511, 519, 523, and 526.

Data access

Supplemental data and code are available under the DOI: 10.5281/zenodo.13788100

ACKNOWLEDGEMENTS

We thank Thomas Dever, Stefan Rothenburg, Adam Phillippy, Bernard Moss, Stephanie

Jaquet, Darach Miller, and members of the Sadhu lab for helpful discussions. We thank Thomas

Dever and Mark Rose for strains and plasmids. Next-generation sequencing was performed by

both the NIH Intramural Sequencing Center (NISC) and the Microarrays and Single-Cell

Genomics Core of the National Human Genome Research Institute. This work utilized the

computational resources provided by the NIH HPC Biowulf Cluster (http://hpc.nih.gov). This

work was supported by the Intramural Research Program of the National Human Genome

Research Institute, NIH (1ZIAHG200401).

AUTHOR CONTRIBUTIONS

Designed research/conceptualization: M.J.C. and M.J.S. designed research;

Performed research: M.J.C. and S.B.S. performed research;

Analyzed data: M.J.C., S.B.S., and M.J.S. analyzed data;

Wrote paper: M.J.C. and M.J.S. wrote the paper.

All authors read and revised the paper.

AUTHOR DETAILS

Michael J Chambers

• Center for Genomics and Data Science Research, National Human Genome Research

Institute, National Institutes of Health, Bethesda, Maryland, USA

• Department of Microbiology & Immunology, Georgetown University, Washington DC,

USA

Contribution: Conceptualization, Investigation, Visualization, Writing - original draft

Competing interests: No competing interests declared

ORCID: 0000-0002-0658-6984

Sophia B Scobell

Center for Genomics and Data Science Research, National Human Genome Research Institute,

National Institutes of Health, Bethesda, Maryland, USA

Contribution: Investigation, Analysis, Writing - review and editing

Competing interests: No competing interests declared

ORCID: 0000-0003-4630-0167

Meru J Sadhu

Center for Genomics and Data Science Research, National Human Genome Research Institute,

National Institutes of Health, Bethesda, Maryland, USA

Contribution: Conceptualization, Supervision, Analysis, Writing - original draft

For correspondence: meru.sadhu@nih.gov

Competing interests: No competing interests declared

ORCID: 0000-0002-8636-9102

FIGURES

Figure 1. Exploring genetic variants of human PKR against the pseudosubstrate antagonist vaccinia K3. (A) (Left) AlphaFold2 model of the PKR kinase domain (green) bound to its target, eIF2α (gray). (Right) Rotated 90°, the eIF2α binding surface of PKR is shaded gray and outlined in white dashed line. (B) PKR kinase domain with vertebrate positive selection sites highlighted in red (13), with the K3 binding surface delineated with white dashed line. (C) AlphaFold2 model of the PKR kinase domain bound to vaccinia K3 (purple). Rotated 90°, the K3 binding surface of PKR is shaded purple and outlined in white dashed line. (D) (Left) The PKR kinase domain with four windows highlighted in which we designed 426 nonsynonymous variants. Secondary structures occurring in these windows (alpha helices α0, αD, and αG, beta strand β1, and the kinase activation loop) are marked (5). The positions of conserved sites and sites under positive selection (13) are denoted above with black and red triangles, respectively, and the positions of sites within 5 angstroms of either K3 or eIF2α in an Alphafold2 binding model are denoted with purple and gray triangles, respectively. Note that the modeled interaction between PKR and eIF2α resembles the pre-phosphorylation state in which the phosphoacceptor loop of eIF2α has not entered the PKR active site, and thus there are likely additional sites in PKR that contact eIF2α. (Right) Mutated windows are highlighted on the surface of the PKR kinase domain, with the K3 binding surface delineated with a white dashed line, and in cartoon form with secondary structures marked. (E) Methodological approach to explore the effects of variants of human PKR in the presence of different K3 alleles. 426 nonsynonymous variants of PKR were generated and paired with wild-type K3, K3Δ58, and K3-H47R. Variant effects were characterized using a high-throughput yeast growth assay and massively parallel sequencing.

675
Figure 2. PKR variants that evade K3 and maintain kinase function are enriched at
positive selection sites and helices αD and αG. (A-B) PKR functional scores versus K3 are
colored ranging from susceptible (red) to WT-like (white) to resistant (blue). (A) Heatmap of
PKR variants with cells colored by the PKR functional score versus K3 for SNP-accessible
variants. Wild-type PKR residues and untested variants are denoted with black circles and gray
squares, respectively. Previously characterized K3-H47R-resistant variants are noted with an
asterisk (24). (B) Attractive electrostatic interaction between PKR-Glu375 (blue) and K3-Lys45
(yellow) in the AlphaFold2 model of PKR bound to K3. (C) Surface structure of the PKR kinase
domain with sites colored by the mean PKR functional score versus K3 for missense variants.
The K3 binding surface is delineated with black dashed line. (Inset) Location of K3-resistant
sites cited in the text. (D-F) Strip plots of PKR functional score versus K3 for variants, with green
and red dashed lines representing mean scores for WT PKR and nonsense variants,
respectively. Variants are partitioned by nearest secondary structural element (D), level of
conservation in vertebrates (E), or predicted contact with K3 (F). Points in D are ordered left to
right by their position along the kinase domain, whereas they are randomly jittered along the x-
axis in E and F. (* p < .05, **** p < .0001, Tukey's HSD (D) and two-sample t-test (E, F).)

692
Figure 3. Few nonfunctional PKR variants identified in the absence of K3 inhibition. (A)
Scatter plot showing each variant's PKR functional score versus wild-type K3 plotted against its
PKR functional score versus K3Δ58. The data point for PKR-WT is colored green and a data
point representing the average of the four nonsense variants (PKRΔ) is colored red. (B-C) PKR
functional scores versus K3Δ58 are colored ranging from nonfunctional (red) to WT-like (white).
(B) Surface structure of the PKR kinase domain with sites colored by the mean PKR functional
score versus K3Δ58 for missense variants. The K3 binding surface is delineated with a black
dashed line. (Inset) Location of highly constrained sites cited in the text. (C) Heatmap of PKR
variants with cells colored by the PKR functional score versus K3Δ58 for each variant.

702
Figure 4. Identification of PKR sites highly susceptible to K3 inhibition. (A) A line was
drawn connecting the data point for PKR-WT to the position of the average data point for the
four nonsense variants (PKRΔ) from the data in the scatter plot from Figure 3A, with all other
data points colored by their residual from that line (K3 / K3Δ58 Residuals), ranging from K3-
susceptible (purple) to K3-indifferent (green) to K3-resistant (yellow). This color scheme is used
in panels B and C. (B) Surface structure of the PKR kinase domain with sites colored by the
mean K3 / K3Δ58 residuals for missense variants. The K3 contact site is delineated with a black
dashed line. (Inset) Location of K3-susceptible sites cited in the text. (C) Heatmap of PKR
variants with cells colored by the K3 / K3Δ58 residuals for each variant.

712
Figure 5. PKR variants that are K3-resistant are also largely K3-H47R-resistant. (A) Scatter
plot of PKR functional scores versus wild-type K3 plotted against PKR functional scores versus
K3-H47R. The data point for PKR-WT is colored green and a datapoint representing the
average of the four nonsense variants (PKRΔ) is colored red. (B-C) PKR functional score versus
K3-H47R ranging from susceptible (red) to WT-like (white) to resistant (blue). (B) Surface
structure of the PKR kinase domain with sites colored by the mean PKR functional score versus
K3-H47R for missense variants. The K3 binding surface is delineated with a black dashed line.
(Inset) Location of K3-H47R resistant sites cited in the text. (C) Heatmap of PKR variants with
cells colored by the PKR functional score versus K3-H47R for each variant. (D-F) Strip plots of
PKR functional scores versus K3-H47R for variants, as plotted in Figure 2D-F. Variants are
partitioned by nearest secondary structural element (D), level of conservation in vertebrates (E),
or predicted contact with K3 (F). Points in D are ordered left to right by their position along the
kinase domain, whereas they are randomly jittered along the x-axis in E and F. (* p < .05, ** p <
.01, **** p < .0001, Tukey's HSD (D) and two-sample t-test (E, F).)

727
Figure 6. Unified spatial view of highlighted PKR sites across the K3-binding interface.
Positions of sites classified as essential residues (black), K3-resistant (blue), and K3-
susceptible (red) based upon the above analyses. All other tested sites are marked white. (Left)
The vaccinia K3 binding surface of PKR is outlined with a black dashed line. (B) PKR functional
scores versus K3Δ58 (Left), K3 (Center), and K3-H47R (Right), with each vertical line
representing a PKR variant. Variants are sorted by functional score from high (left) to low (right).
Gray and black lines indicate the standard deviation and standard error, respectively, in PKR
functional scores across barcodes associated with the given variant. (C) (Left) AlphaFold2
model of Vaccinia K3 with surface residues colored as different from eIF2α (purple) or identical
to eIF2α (gray) based on sequence alignment (Supplemental Figure 6). (Right) Rotated 90°, K3
residues that contact the PKR activation loop or helix αD or αG are outlined in pink and dashed
orange and burgundy, respectively.

SUPPLEMENTAL FIGURES

Supplemental Figure 1. Composition of the PKR variant library. (A) Barchart depicts the number of variants made at each of the four PKR windows categorized by amino acid properties: Positive Charge = H, K, R; Negative Charge = D, E; Polar-Neutral = C, N, Q, S, T; Non-Polar = A, I, L, M, V; Aromatic = F, W, Y; Unique = G, P; Stop = *. A total of 140 variants were made in α0/β1, 86 in αD, 43 in the Activation Loop, and 161 in αG for a total of 430 variants, which includes the 4 nonsense variants made in each window. (B) Histogram depicts the number of unique barcodes linked to each PKR variant, with a mean of 42.83.

749
Supplemental Figure 2. Calculation of PKR functional scores from yeast growth assay.
(A) Line plot of mock barcode read count data over time for cells expressing nonfunctional (red)
or functional (blue) PKR across four sampled timepoints. As PKR activity is toxic to yeast, the
number of cells in the pool expressing the functional PKR will decrease over time, and thus the
associated barcode read count will also decrease, while the read count for the nonfunctional
PKR will increase. (B) Line plot depicting the fold changes from timepoint 0 of barcode
abundance derived from Panel A, with a -log2 transformation to assign positive values to
functional PKR and negative values to nonfunctional PKR, from which the area under the curve
(AUC) is calculated to produce a PKR functional score.

759
Supplemental Figure 3. Replication of PKR variants paired with K3 alleles. (A) Scatter plot
of PKR functional scores for variants paired with K3-WT for two biological replicate experiments.
Functional scores were calculated across four timepoints as the area under the curve (see
Materials and Methods). Thus, variants with increased evasion of K3 would have higher PKR
functional scores, while those with increased susceptibility to K3 or loss of eIF2α kinase activity
would have lower scores.

766
Supplemental Figure 4. Experimental validation of the K3 resistance phenotypes of PKR variants. Select PKR variants were generated and screened against K3Δ58, WT, and H47R alleles using a yeast growth assay visualized through serial dilution (23). PKR is under the inducible pGAL10/CYC1 hybrid promoter, and thus is repressed in dextrose media (left) and expressed in galactose media (right). We selected one variant that exhibited loss of function in the high-throughput screen (H483P, Figure 3C), one variant whose phenotype was similar to PKR WT (E375D, Figures 2A, 3C, and 5C), and five variants whose K3 resistance was improved (E375K, L452S, T487S, T491R, and T496P, Figures 2A and 5C).

775
Supplemental Figure 5. Bimodal distribution of PKR functional scores at K3 contact
sites. Strip plot of PKR functional scores versus K3 partitioned by predicted contact with K3.
Black dashed line denotes the threshold at which variants were separated as nonfunctional-like
or functional. Green and red dashed lines represent mean scores for WT PKR and nonsense
variants. (** p < 0.01, two-sample t-test, comparing just the variants classified as functional in
the two classes.)

```
                                                                    #
          0        10        20        30        40        50
Human eIF2α   MPGLSCRFYQHKFPEVEDVVMVNVRSIAEMGAYVSLLEYNNIEGMILLSELSR--RRIRS
              1        10        20          30        40        50
Vaccinia K3   -----MLAFCYSLPNAGDVIKGRVYE-KDYALYIYLFDYPHFEAILAESVKMHMDRYVEY
                  : :..:*:. **:  .* .  : . *: *::* ::*.::  *   .  * :

          60        70        80        90       100       110
Human eIF2α   INKLIRIGRNECVVVIRVDKEKGYIDLSKRRVSPEEAIKCEDKFTKSKTVYSILRHVAEV
              55      60        70        80
Vaccinia K3   RDKL--VGKTVKVKVIRVDYTKGYIDVNYKRMCRHQ------------------------
                  :**  :*.. * ***** *****:. .*:.  :
```

Supplemental Figure 6. Sequence similarity between human eIF2α and vaccinia K3.
Alignment of human eIF2α (RefSeq Accession NP_004085.1, residues 0-117) to vaccinia K3
(RefSeq Accession YP_232916.1). The Ser51 site of phosphorylation in eIF2α is indicated by #.
Sequences were aligned using Muscle 3.8 and displayed in Clustal format, * = fully conserved, :
= strong group conservation, . = weak group conservation. Sites 41-58 of K3 correspond to the
rigid helix insert, which is proximal to PKR's ATP-binding site in the Alphafold2-predicted
complex, while sites 72-83 are proximal to PKR's helix αG.

790
Supplemental Figure 7. Differing patterns of resistance between wild-type K3 and K3-
H47R. (A) A nonlinear exponential curve (black line) was fitted to the data in the scatter plot
from Figure 5A. Points are colored by their residuals from that curve (K3 / K3-H47R Residuals),
ranging from enhanced (purple) to decreased (yellow) resistance to K3-H47R relative to the
expectation from K3 resistance. This color scheme is used in panels B and C. (B) Surface
structure of PKR kinase domain with sites colored by the mean K3 / K3-H47R residuals for
missense variants. The K3 contact site is delineated with a black dashed line. (Inset) Location of
K3-H47R-resistant sites where variants deviated most strongly between the wild-type K3 and
K3-H47R conditions. (C) Heatmap of PKR variants with cells colored by the K3 / K3-H47R
residuals for each variant. (D) Two PKR variants that did not follow the general trend of
enhanced susceptibility to K3-H47R relative to K3 WT were selected for validation. The variants

were generated and screened against K3Δ58, WT, and H47R alleles using a yeast growth
assay visualized through serial dilution as in Supplemental Figure 4.

804
Supplemental Figure 8. Additional variants at the sites of previously identified K3-H47R-
resistant variants often also conferred resistance to K3. Strip plots of PKR functional scores
of variants paired with K3-H47R (top) and K3-WT (bottom). Each plot highlights PKR variants
made at sites where an improved PKR variant was previously identified (24). Blue markers
denote previously identified variants; black markers are additional variants made at the same
site. Green and red dashed lines represent mean scores for WT PKR and nonsense variants,
respectively. Histograms (right) show PKR functional scores versus K3-H47R (top) and K3
(bottom) for all variants. * = stop codon.

Supplemental Figure 9. Systematic generation of PKR variants using mixed-base primer
tile sets. (A) Each variant primer is composed of a homology arm, variant region, and priming
region. (B) Nonsynonymous SNP-accessible variants are generated by altering the codon in the
variant region of the primer. This example depicts the codon "AAT" encoding Asn. The first
nucleotide in the codon, "A", is underlined in red, with three codons above having changes to
"C", "G", and "T" underlined in red, which generate the nonsynonymous variants His, Asp, and
Tyr. (C) Variant primers were designed across all three nucleotides in each codon, as
underlined in red. (D) Variant primer tile sets, represented in dark green, were made by pooling
variant primers that modify adjacent codons. Primers included in a given variant primer tile sets
have differing variant regions but share homology arms and priming region sequences.

Supplemental Figure 10. Multiple unique barcode sequences were attached to each PKR
variant. (A) The barcode primer is composed of a homology arm, barcode region, and priming
region. The barcode primer is used as the reverse primer in the variant-generating PCR
reactions to attach a unique nucleotide sequence after the PKR variant sequence. (B) The
barcode region of the primer is composed of 20 "N", representing an equal mix of the
nucleotides "A", "C", "G", and "T". A barcode primer with 20 random nucleotides can take on 4^{20}
unique nucleotide sequences. Dinucleotide sequences "TT" and "AA" are interspersed
throughout the barcode region to avoid making unintended restriction enzyme cut sites.

833
Supplemental Figure 11. Assembly of PKR variant library using variant tile sets and
barcode primers. (A) 15 variant primer tile sets were designed to generate variants across four
windows of interest in PKR. The full-length PKR sequence is denoted in green, with Windows 1-
4 overlaid in yellow, orange, magenta, and burgundy, respectively. Variant primer tile sets were
used to generate 426 PKR missense variants, for a total of 15 PCR-1 reactions. (B) Two
separate PCR reactions were used to generate complementary insert and vector fragments.
PCR-1 primers (light gray box) include a single variant primer tile set (dark green, see
Supplemental Figure 9) and a single doped barcode primer (light blue, see Supplemental Figure
10) that amplified from WT PKR (green) and made the PCR-1 insert fragment containing a
select nonsynonymous variants (dark green) and a unique barcodes (blue). PCR-2 primers
included a single forward and reverse primer that amplified from WT PKR and made a larger
vector fragment with 20-bp homology arms that complement the homology arms of the PCR-1
insert fragment. The two fragments were combined via Gibson Assembly to form a pool of
complete vectors, each vector containing a single, nonsynonymous variant with a unique
barcode.

849
Supplemental Figure 12. Alphafold2 Multimer predictions used to identify PKR sites
proximal to eIF2α and K3. (A) AlphaFold2 Multimer prediction of PKR in complex with eIF2α
(Left) and K3 (Right). Residues are colored by AlphaFold2 pLDDT confidence scores per
residue. In both predicted models, PKR is on the left and its binding partner is on the right, as in
Figure 1A,B. (B) Crystal structure of PKR (green, PDB 2A1A) in complex with eIF2α (gray, PDB
2A1A) with the crystal structure of vaccinia K3 (purple, PDB 1LUZ) aligned to eIF2α (root-mean-
square deviation (RMSD) = 0.819 Å). (C) Overlay of the PKR-eIF2a from PDB 2A1A (green and
gray) and AlphaFold2 (blue), RMSD = 0.970 Å. (D) Overlay of the PKR-Vaccinia K3 complex
from PDB 2A1A and 1LUZ (green and purple) to AlphaFold2 (blue), RMSD = 0.944 Å.

SUPPLEMENTAL DATA
860
Supplementary File 1 - 1_Oligo-Table.xlsx
Supplementary File 2 - 2_Variant-Primers.xlsx
Supplementary File 3 - 3_Plasmids.xlsx
Supplementary File 4 - 4_PKR-Functional-Scores.csv
Supplementary File 5 - 5_Structure-Alignment-RMSD-Values.xlsx
Supplementary File 6 - 6_AF2_PKR-KD_eIF2a.pdb
Supplementary File 7 - 7_AF2_PKR-KD_VACV-K3.pdb

868 REFERENCES

1. Daugherty MD, Malik HS. Rules of Engagement: Molecular Insights from Host-Virus Arms
Races. Annual Review of Genetics. 2012;46(1):677–700.

2. Metz DH, Esteban M. Interferon inhibits Viral Protein Synthesis in L Cells infected with Vaccinia
Virus. Nature. 1972 Aug;238(5364):385–8.

3. Clemens MJ. PKR—A protein kinase regulated by double-stranded RNA. The International
Journal of Biochemistry & Cell Biology. 1997 Jul 1;29(7):945–9.

4. Balachandran S, Roberts PC, Brown LE, Truong H, Pattnaik AK, Archer DR, et al. Essential
Role for the dsRNA-Dependent Protein Kinase PKR in Innate Immunity to Viral Infection.
Immunity. 2000 Jul 1;13(1):129–41.

5. Dar AC, Dever TE, Sicheri F. Higher-Order Substrate Recognition of eIF2α by the RNA-
Dependent Protein Kinase PKR. Cell. 2005 Sep 23;122(6):887–900.

6. Weber F, Wagner V, Rasmussen SB, Hartmann R, Paludan SR. Double-Stranded RNA Is
Produced by Positive-Strand RNA Viruses and DNA Viruses but Not in Detectable Amounts by
Negative-Strand RNA Viruses. Journal of Virology. 2006 May 15;80(10):5059–64.

7. Meurs E, Chong K, Galabru J, Thomas NSB, Kerr IM, Williams BRG, et al. Molecular cloning
and characterization of the human double-stranded RNA-activated protein kinase induced by
interferon. Cell. 1990 Jul 27;62(2):379–90.

8. Kaufman RJ. Double-stranded RNA-activated protein kinase PKR. Cold Spring Harbor
Monograph Series. 2000;39:503–28.

9. Dey M, Cao C, Dar AC, Tamura T, Ozato K, Sicheri F, et al. Mechanistic Link between PKR
Dimerization, Autophosphorylation, and eIF2α Substrate Recognition. Cell. 2005 Sep
23;122(6):901–13.

10. Lemaire PA, Anderson E, Lary J, Cole JL. Mechanism of PKR Activation by dsRNA. Journal of
molecular biology. 2008;381(2):351–60.

11. Hershey JWB. Translational Control in Mammalian Cells. Annu Rev Biochem. 1991
Jun;60(1):717–55.

12. Sudhakar A, Ramachandran A, Ghosh S, Hasnain SE, Kaufman RJ, Ramaiah KVA.
Phosphorylation of Serine 51 in Initiation Factor 2α (eIF2α) Promotes Complex Formation
between eIF2α(P) and eIF2B and Causes Inhibition in the Guanine Nucleotide Exchange
Activity of eIF2B. Biochemistry. 2000 Oct 1;39(42):12929–38.

13. Rothenburg S, Seo EJ, Gibbs JS, Dever TE, Dittmar K. Rapid evolution of protein kinase PKR
alters sensitivity to viral inhibitors. Nature Structural & Molecular Biology. 2009 Jan;16(1):63–70.

14. Elde NC, Child SJ, Geballe AP, Malik HS. Protein kinase R reveals an evolutionary model for
defeating viral mimicry. Nature. 2009 Jan;457(7228):485–9.

15. Langland JO, Cameron JM, Heck MC, Jancovich JK, Jacobs BL. Inhibition of PKR by RNA and
DNA viruses. Virus research. 2006;119(1):100–10.

16. Nallagatla SR, Toroney R, Bevilacqua PC. Regulation of innate immunity through RNA structure
and the protein kinase PKR. Curr Opin Struct Biol. 2011 Feb;21(1):119–27.

17. Hull CM, Bevilacqua PC. Discriminating Self and Non-Self by RNA: Roles for RNA Structure,
Misfolding, and Modification in Regulating the Innate Immune Sensor PKR. Acc Chem Res.
2016 Jun 21;49(6):1242–9.

18. Beattie E, Tartaglia J, Paoletti E. Vaccinia virus-encoded eIF-2α homolog abrogates the antiviral
effect of interferon. Virology. 1991;183(1):419–22.

19. Carroll K, Elroy-Stein O, Moss B, Jagus R. Recombinant vaccinia virus K3L gene product
prevents activation of double-stranded RNA-dependent, initiation factor 2 alpha-specific protein
kinase. Journal of Biological Chemistry. 1993 Jun 15;268(17):12837–42.

20. Davies MV, Chang HW, Jacobs BL, Kaufman RJ. The E3L and K3L vaccinia virus gene
products stimulate translation through inhibition of the double-stranded RNA-dependent protein
kinase by different mechanisms. Journal of Virology. 1993 Mar;67(3):1688–92.

21. Sharp TV, Witzel JE, Jagus R. Homologous Regions of the α Subunit of Eukaryotic Translational Initiation Factor 2 (eIF2α) and the Vaccinia Virus K3L Gene Product Interact with the Same Domain within the dsRNA-Activated Protein Kinase (PKR). European Journal of Biochemistry. 1997;250(1):85–91.

22. Dar AC, Sicheri F. X-Ray Crystal Structure and Functional Analysis of Vaccinia Virus K3L Reveals Molecular Determinants for PKR Subversion and Substrate Recognition. Molecular Cell. 2002 Aug 1;10(2):295–305.

23. Kawagishi-Kobayashi M, Silverman JB, Ung TL, Dever TE. Regulation of the protein kinase PKR by the vaccinia virus pseudosubstrate inhibitor K3L is dependent on residues conserved between the K3L protein and the PKR substrate eIF2alpha. Molecular and Cellular Biology. 1997 Jul 1;17(7):4146–58.

24. Seo EJ, Liu F, Kawagishi-Kobayashi M, Ung TL, Cao C, Dar AC, et al. Protein kinase PKR mutants resistant to the poxvirus pseudosubstrate K3L protein. Proceedings of the National Academy of Sciences. 2008 Nov 4;105(44):16894–9.

25. Chong K l., Feng L, Schappert K, Meurs E, Donahue T f., Friesen J d., et al. Human p68 kinase exhibits growth suppression in yeast and homology to the translational regulator GCN2. The EMBO Journal. 1992 Apr 1;11(4):1553–62.

26. Dever TE, Chen JJ, Barber GN, Cigan AM, Feng L, Donahue TF, et al. Mammalian eukaryotic initiation factor 2 alpha kinases functionally substitute for GCN2 protein kinase in the GCN4 translational control mechanism of yeast. PNAS. 1993 May 15;90(10):4616–20.

27. Elde NC, Child SJ, Eickbush MT, Kitzman JO, Rogers KS, Shendure J, et al. Poxviruses Deploy Genomic Accordions to Adapt Rapidly against Host Antiviral Defenses. Cell. 2012 Aug 17;150(4):831–41.

28. Dey M, Velyvis A, Li JJ, Chiu E, Chiovitti D, Kay LE, et al. Requirement for kinase-induced conformational change in eukaryotic initiation factor 2α (eIF2α) restricts phosphorylation of Ser51. Proceedings of the National Academy of Sciences. 2011 Mar 15;108(11):4316–21.

29. Haller SL, Peng C, McFadden G, Rothenburg S. Poxviruses and the evolution of host range and virulence. Infection, Genetics and Evolution. 2014 Jan 1;21:15–40.

30. Rothenburg S, Chinchar VG, Dever TE. Characterization of a ranavirus inhibitor of the antiviral protein kinase PKR. BMC Microbiology. 2011 Mar 18;11(1):56.

31. Sawyer SL, Wu LI, Emerman M, Malik HS. Positive selection of primate TRIM5α identifies a critical species-specific retroviral restriction domain. Proceedings of the National Academy of Sciences. 2005 Feb 22;102(8):2832–7.

32. Tenthorey JL, Young C, Sodeinde A, Emerman M, Malik HS. Mutational resilience of antiviral restriction favors primate TRIM5α in host-virus evolutionary arms races. Schoggins JW, Weigel D, Berthoux L, editors. eLife. 2020 Sep 15;9:e59988.

33. Colón-Thillet R, Hsieh E, Graf L, McLaughlin RN, Young JM, Kochs G, et al. Combinatorial mutagenesis of rapidly evolving residues yields super-restrictor antiviral proteins. PLoS Biol. 2019;17(10):e3000181.

34. Chavarría-Smith J, Mitchell PS, Ho AM, Daugherty MD, Vance RE. Functional and Evolutionary Analyses Identify Proteolysis as a General Mechanism for NLRP1 Inflammasome Activation. PLOS Pathogens. 2016 Dec 7;12(12):e1006052.

35. Demogines A, Farzan M, Sawyer SL. Evidence for ACE2-Utilizing Coronaviruses (CoVs) Related to Severe Acute Respiratory Syndrome CoV in Bats. Journal of Virology. 2012 Jun 1;86(11):6350–3.

36. Damas J, Hughes GM, Keough KC, Painter CA, Persky NS, Corbo M, et al. Broad host range of SARS-CoV-2 predicted by comparative and structural analysis of ACE2 in vertebrates. PNAS. 2020 Sep 8;117(36):22311–22.

37. Frank HK, Enard D, Boyd SD. Exceptional diversity and selection pressure on coronavirus host receptors in bats compared to other mammals. Proc Biol Sci. 289(1979):20220193.

38. Lan J, Ge J, Yu J, Shan S, Zhou H, Fan S, et al. Structure of the SARS-CoV-2 spike receptor-
binding domain bound to the ACE2 receptor. Nature. 2020 May;581(7807):215–20.

39. Chan KK, Dorosky D, Sharma P, Abbasi SA, Dye JM, Kranz DM, et al. Engineering human
ACE2 to optimize binding to the spike protein of SARS coronavirus 2. Science. 2020 Sep
4;369(6508):1261–5.

40. Heinzelman P, Romero PA. Discovery of human ACE2 variants with altered recognition by the
SARS-CoV-2 spike protein. PLOS ONE. 2021 May 12;16(5):e0251585.

41. Chen S, Francioli LC, Goodrich JK, Collins RL, Kanai M, Wang Q, et al. A genomic mutational
constraint map using variation in 76,156 human genomes. Nature. 2024 Jan;625(7993):92–100.

42. Jacquet S, Culbertson M, Zhang C, El Filali A, De La Myre Mory C, Pons JB, et al. Adaptive
duplication and genetic diversification of protein kinase R contribute to the specificity of bat-virus
interactions. Science Advances. 2022 Nov 23;8(47):eadd7540.

43. Siekierka J, Manne V, Ochoa S. Mechanism of translational control by partial phosphorylation of
the alpha subunit of eukaryotic initiation factor 2. Proceedings of the National Academy of
Sciences. 1984 Jan;81(2):352–6.

44. Ogden PJ, Kelsic ED, Sinai S, Church GM. Comprehensive AAV capsid fitness landscape
reveals a viral gene and enables machine-guided design. Science. 2019 Nov
29;366(6469):1139–43.

45. Parvez S, Herdman C, Beerens M, Chakraborti K, Harmer ZP, Yeh JRJ, et al. MIC-Drop: A
platform for large-scale in vivo CRISPR screens. Science. 2021 Sep 3;373(6559):1146–51.

46. Popp NA, Powell RL, Wheelock MK, Zapp BD, Holmes KJ, Sheldon KM, et al. Multiplex,
multimodal mapping of variant effects in secreted proteins [Internet]. bioRxiv; 2024 [cited 2024
Apr 30]. p. 2024.04.01.587474. Available from:
https://www.biorxiv.org/content/10.1101/2024.04.01.587474v1

47. Park C, Peng C, Brennan G, Rothenburg S. Species-specific inhibition of antiviral protein kinase
R by capripoxviruses and vaccinia virus. Annals of the New York Academy of Sciences.
2019;1438(1):18–29.

48. Park C, Peng C, Rahman MJ, Haller SL, Tazi L, Brennan G, et al. Orthopoxvirus K3 orthologs
show virus- and host-specific inhibition of the antiviral protein kinase PKR. PLOS Pathogens.
2021 Jan 14;17(1):e1009183.

49. Peng C, Haller SL, Rahman MM, McFadden G, Rothenburg S. Myxoma virus M156 is a specific
inhibitor of rabbit PKR but contains a loss-of-function mutation in Australian virus isolates.
PNAS. 2016 Apr 5;113(14):3855–60.

50. Smith GL. Genus Orthopoxvirus: Vaccinia virus. In: Mercer AA, Schmidt A, Weber O, editors.
Poxviruses [Internet]. Basel: Birkhäuser; 2007 [cited 2024 Sep 18]. p. 1–45. Available from:
https://doi.org/10.1007/978-3-7643-7557-7_1

51. Rose MD, Novick P, Thomas JH, Botstein D, Fink GR. A *Saccharomyces cerevisiae* genomic
plasmid bank based on a centromere-containing shuttle vector. Gene. 1987 Jan 1;60(2):237–
43.

52. Curran KA, Morse NJ, Markham KA, Wagman AM, Gupta A, Alper HS. Short Synthetic
Terminators for Improved Heterologous Gene Expression in Yeast. ACS Synth Biol. 2015 Jul
17;4(7):824–32.

53. Brachmann CB, Davies A, Cost GJ, Caputo E, Li J, Hieter P, et al. Designer deletion strains
derived from Saccharomyces cerevisiae S288C: A useful set of strains and plasmids for PCR-
mediated gene disruption and other applications. Yeast. 1998;14(2):115–32.

54. Liu X, Liu Z, Dziulko AK, Li F, Miller D, Morabito RD, et al. iSeq 2.0: A Modular and
Interchangeable Toolkit for Interaction Screening in Yeast. Cell Systems. 2019 Apr 24;8(4):338-
344.e8.

55. Cornish-Bowden A. Nomenclature for incompletely specified bases in nucleic acid sequences:
recommendations 1984. Nucleic acids research. 1985;13(9):3021.

56. Guido VR, Drake Jr FL. Python 3 reference manual. Scotts Valley: CreateSpace. 2009;

57. Crawford KH d, Bloom JD. alignparse: A Python package for parsing complex features from high-throughput long-read sequencing. Journal of Open Source Software. 2019 Dec 11;4(44):1915.

58. Sambrook J, Russell DW. Standard Ethanol Precipitation of DNA in Microcentrifuge Tubes. Cold Spring Harb Protoc. 2006 Jun 1;2006(1):pdb.prot4456.

59. Gietz RD, Schiestl RH. Large-scale high-efficiency yeast transformation using the LiAc/SS carrier DNA/PEG method. Nat Protoc. 2007;2(1):38–41.

60. Singh MV, Anthony Weil P. A method for plasmid purification directly from yeast. Analytical Biochemistry. 2002 Aug 1;307(1):13–7.

61. Wu L, Wen C, Qin Y, Yin H, Tu Q, Van Nostrand JD, et al. Phasing amplicon sequencing on Illumina Miseq for robust environmental microbial community analysis. BMC Microbiol. 2015 Dec;15(1):1–12.

62. Zhang J, Kobert K, Flouri T, Stamatakis A. PEAR: a fast and accurate Illumina Paired-End reAd mergeR. Bioinformatics. 2014 Mar 1;30(5):614–20.

63. Zhao L, Liu Z, Levy SF, Wu S. Bartender: a fast and accurate clustering algorithm to count barcode reads. Bioinformatics. 2018 Mar 1;34(5):739–47.

64. Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, et al. UCSF ChimeraX: Tools for structure building and analysis. Protein Science. 2023;32(11):e4792.

65. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. ColabFold: making protein folding accessible to all. Nat Methods. 2022 Jun;19(6):679–82.

66. Dhaliwal S, Hoffman DW. The Crystal Structure of the N-terminal Region of the Alpha Subunit of Translation Initiation Factor 2 (eIF2α) from *Saccharomyces cerevisiae* Provides a View of the Loop Containing Serine 51, the Target of the eIF2α-specific Kinases. Journal of Molecular Biology. 2003 Nov 21;334(2):187–95.