1

Multimodal single-cell profiling reveals neuronal vulnerability and pathological cell

states in focal cortical dysplasia

4
5

Isabella C. Galvão[1,2], Manuela Lemoine[1,2], Lauana A. Messias[1,2], Patrícia A.O.R.A. Araújo[1,2],

Jaqueline C. Geraldis[1,2], Clarissa L. Yasuda[2,3], Marina K. M. Alvim[2,3], Enrico Ghizoni[2,3],

Helder Tedeschi[2,3], Fernando Cendes[2,3], Fabio Rogerio[2,4], Iscia Lopes-Cendes[1,2], and Diogo F.

T. Veiga[1,2,*]

[1]Department of Translational Medicine, School of Medical Sciences, University of Campinas,

Campinas (UNICAMP), Brazil

[2]The Brazilian Institute of Neuroscience and Neurotechnology (BRAINN), Campinas, Brazil

[3]Department of Neurology, School of Medical Sciences, University of Campinas (UNICAMP),

Campinas, SP, Brazil

[4]Department of Pathology, School of Medical Sciences, University of Campinas (UNICAMP),

Campinas, Brazil

[*]Corresponding author: dfveiga@unicamp.br

20
21
22

ABSTRACT

Focal Cortical Dysplasia (FCD) is a neurodevelopmental condition characterized by

malformations of the cerebral cortex that often cause drug-resistant epilepsy. In this study, we

performed multi-omics single-cell profiling to map the chromatin accessibility and

transcriptome landscapes of FCD type II, generating a comprehensive multimodal single-cell

dataset comprising 61,525 cells from 11 clinical samples of lesions and controls. Our findings

revealed profound chromatin, transcriptomic, and cellular alterations affecting neuronal and

glial cells in FCD lesions, including the selective loss of upper-layer excitatory neurons,

significant expansion of oligodendrocytes and immature astrocytic populations, and a unique

neuronal subpopulation harboring dysmorphic neurons. Furthermore, we uncovered activated

microglia subsets, particularly in FCD IIb cases. This comprehensive study unveils neuronal

and glial cell states driving FCD development and epileptogenicity, enhancing our

understanding of FCD and offering new directions for targeted therapy development.

36
37
38
39

40   **Keywords:** focal cortical dysplasia; cortical malformations; single-cell sequencing; multi-

41   omics; gene expression; chromatin accessibility; microglia; reactive astrocytes

INTRODUCTION

Focal Cortical Dysplasia (FCD) is a malformation of the cerebral cortex and a major cause of drug-resistant epilepsy in children. Treatment of this neurological condition with current anti-seizure medications is often ineffective, frequently requiring surgical removal of the affected brain tissue. FCD is stratified based on neuroimaging, histopathology, and genetic characteristics[1,2]. FCD type II, the most common presentation, is characterized by cytoarchitectural cortical abnormalities including loss of cortical lamination, blurring of the gray-white matter junction, and abnormal cellular development[2]. FCD II cortical lesions can be further divided based on cytological features: FCD IIa is characterized by the presence of dysmorphic neurons, while FCD IIb presents with both dysmorphic neurons and balloon cells. Dysmorphic neurons are abnormal cells featuring enlarged soma and nuclei[1,2], along with expression of neurofilament markers typical of immature neurons in the developing cortex. On the other hand, balloon cells are characterized by large cell bodies with vitreous, opalescent, and eosinophilic cytoplasm[1,2]. These cells express markers commonly associated with neuroglial progenitors, including nestin and vimentin, indicating their immature nature[3]. High expression of other progenitor cell markers such as c-Myc, SOX2, and Oct-4 were also observed throughout FCD II lesions[4,5].

At the molecular level, FCD II cases display activation of the mammalian target of rapamycin complex 1 (mTORC1) signaling pathway, which is implicated in cell growth, increased synaptic transmission and plasticity, as well as regulation of immune system responses[6–8]. Hyperactivation of mTOR signaling, which operates through the mTORC1 and mTORC2 complexes, is particularly observed in abnormal cells such as dysmorphic neurons and balloon cells[9]. Increased phosphorylation of the mTORC1 substrates S6K1 and S6 proteins can be used as a biomarker to distinguish FCD type II lesions[4]. Genetic alterations such as

somatic gain-of-function mutations in mTOR activating genes and germline loss-of-function

mutations of mTOR inhibitors are also found in FCD II[9–11].

The cellular and epileptogenic mechanisms of FCD are not yet fully understood.

Dysmorphic neurons appear to play a pivotal role in generating epileptic discharges[11], while

balloon cells are considered electrically quiescent and may even have an anti-seizure effect[3].

A recent study showed that dysmorphic neurons derive from excitatory neurons while balloon

cells have astrocytic lineage[12]. We and other groups have shown that FCD IIb lesions are also

characterized by increased microglia and astrocyte reactivity when compared to FCD IIa or

healthy controls[13–15]. However, little is known regarding the cellular landscape of FCD lesions

and the role of the diverse cortical cell types in the disease.

Single-cell RNA and transposase-accessible chromatin (ATAC) sequencing have been

applied to resolve the cellular diversity of the human brain in healthy and disease contexts[16–18]

and to discover cell states associated with neurological conditions[19–21]. Recent multimodal

single-cell approaches can obtain multiple measurements from the same cells simultaneously,

enabling the integration of epigenetic and transcriptomic modalities to identify cell types

involved in diseases[22,23].

In this study, we applied multi-omics single-cell profiling to characterize the chromatin

accessibility and gene expression landscapes in FCD type II, generating a comprehensive

multimodal single-cell dataset comprising 61,525 cells from 11 clinical samples of FCD lesions

and controls obtained from epilepsy surgeries. Integrative analyses of this dataset unraveled

cellular unbalances and cell states underlying FCD pathogenesis, and highlighted neuronal

populations susceptible in the disease.

RESULTS

High-resolution characterization of the chromatin accessibility and transcriptome landscapes of FCD type II

We performed multi-omics single-cell sequencing of cortical brain tissue from FCD IIa (n=3) and IIb (n=6) lesions, as well as from non-lesion tissues selected as controls (n=2; Figure 1A). Specimens were collected during epilepsy surgery to treat drug-resistant FCD. Internal controls correspond to histologically normal cortical tissues where neither abnormal cells (dysmorphic neurons and balloon cells) nor architectural changes were observed. Dissociated nuclei were profiled using the 10X Genomics Multiome ATAC + Gene Expression assay for simultaneous measurement of chromatin accessibility and gene expression in individual nuclei. Quality control was performed to remove low-quality nuclei and to filter out potential doublets, and Harmony[24] was applied to ATAC and RNA datasets to integrate samples from different sequencing batches (Figure 1B). The Weighted Nearest Neighbor (WNN) approach[25] from Seurat was used to create an integrated multimodal (ATAC + RNA) visualization (Figure 1B). The resulting multimodal single-cell compedium comprised 61,525 nuclei with paired chromatin and gene expression measurements. Clustering analysis identified a total of 40 populations (Figure S1A), which were annotated using Azimuth[25] followed by manual validation using canonical marker genes to define consensus cell types. Azimuth was able to assign labels with high confidence to the majority of clusters (Figure S1B), and none of the clusters were exclusive to a unique sample (Figure S1C). Consensus cell type annotation was performed using a broad cell types (Figure 1C) as well as a higher-resolution taxonomy including neuronal subtypes and cortical layers (Figure 1D).

Overall, our annotation identified the various cell types expected in the human cortex such as excitatory and inhibitory neurons, astrocytes, oligodendrocytes, oligodendrocytes progenitor cells (OPC), microglia, endothelial cells, and vascular and leptomeningeal cells

(VLMC, Figure 1C). We also identified the various subtypes of excitatory neurons (EN, Figure

1D) including upper layers EN L2/3 (termed EN Upper), and deep layer subtypes EN L5, EN

L5/6, EN L6 (EN Deep), while inhibitory neurons (IN) were further classified into Vip, Sst,

Pvalb, Lamp5 and Sncg/Lamp5 subtypes (Figure 1D). Based on the histological assessment,

most of the sequenced nuclei derived from IIb lesions, followed by internal controls and IIa

lesions (Figure 1E). Gene expression and open chromatin accessibility of marker genes were

inspected to confirm cell type identity (Figures 1F-G). For instance, *AQP4* (astrocytes),

SLC17A7 (EN), *GAD2* (IN), *PLP1* (oligodendrocytes), *CD86* (microglia), and *GPR17* (OPC)

were marker genes with concordant expression and chromatin accessibility in their

corresponding cell types.

Using signature enrichment analysis, mTORC1 signaling was found to be elevated in

dysplastic tissue across all cell types except VLMC and endothelia (Figure S2A), and cells with

the highest mTORC1 activation were identified in FCD IIb (Figure S2B). Thus, multimodal

single-cell profiling uncovered the heterogenous composition of FCD type II lesions and

recapitulated key aspects of their pathological state.

Unraveling cellular changes in FCD lesions

Next, we determined cell type abundance changes between FCD and control samples with

Speckle[26], using an approach that accounts for variability in cellular frequencies among

patients. The analysis uncovered alterations affecting several broad cell types, including

decreased frequency of the EN upper population and increased frequency of oligodendrocytes

in both FCD IIa and IIb lesions (Figure 1H). In addition, we found an expansion of the

microglia population specifically in IIb donors.

We also leveraged the hierarchical nature of the Azimuth annotation, which classifies

excitatory and inhibitory neurons into 116 subclasses based on their expression of markers, to

find out how these neuronal subclasses are affected in the disease. The Exc L2

LINC00507/*GLRA3*, the most common EN subclass identified in the cortex of FCD samples,

was also the most affected EN displaying both the highest number of transcriptional changes

and frequency reduction in lesions (Figure 2A,B). Among IN, FCD IIa displayed an increased

frequency of Parvalbumin-expressing (Pvalb) interneurons Inh L3 Pvalb/*SAMD13* (Figure

S3C); yet, this was not the subclass with most transcriptional changes (Figure 2D). On the other

hand, somatostatin-expressing (Sst) subclass Inh L1-2 Sst/PRRT4 and PAX6-expressing

subclass Inh L1 PAX6/*MIR101-1* were affected only in FCD IIb when compared to controls

(Figure 2C). Altogether, this data shows that cellular abundance changes affect both neuronal

and non-neuronal populations in FCD pathogenesis, and revealed the vulnerability of

excitatory neurons in the disease.

Identification of a disease-specific population containing dysmorphic neurons

During annotation, we found a heterogeneous cluster containing both excitatory neurons and

oligodendrocytes that we initially termed Unclassified since Azimuth could not confidently

assign a cell type (Figure 1C). This cluster was highly disease-specific as it contained 1,308

cells, being that 1,285 cells (98%) originated from FCD (Figure 3A). Despite clustering next

to oligodendrocytes after data integration, 54% of nuclei were classified as EN and 29% as

oligodendrocytes. The cluster was detected in all 9 FCD samples, denoting its reproducibility

across donors (cluster #15, Figure S1C). To further characterize this disease-specific (DS)

population, we identified its markers and performed pathway enrichment analysis. Notably, all

top enriched pathways were related to neurodegenerative diseases including Parkinson's,

amyotrophic lateral sclerosis, and Alzheimer's (Figure 3B). Also, the majority of marker genes

were associated with neuronal processes including oxidative phosphorylation, Ca^{2+} transport,

neuron development, and neurofilament subunits (Figure 3C).

In particular, we found two neurofilament subunits *NEFL* and *NEFM* overexpressed in the DS cluster (Figure 3C). *NEFM* was exclusively expressed in the DS cluster and not detected in any other cell type in either lesions or non-lesion tissues (Figure 3D). The chromatin accessibility in the *NEFM* and *NEFL* loci corroborates with their active gene expression in the DS cluster (Figure 3E).

Antibodies against neurofilaments are often used to identify dysmorphic neurons in dysplastic tissue during histopathological diagnosis of FCD[2,11]. Based on this observation, we used a signature enrichment analysis to inspect whether the DS cluster harbor dysmorphic neurons. Specifically, we analyzed a 30-gene signature expressed in a purified dysmorphic neuron population selected using laser capture microdissection in FCD tissue[27]. We found that nuclei with the highest expression of this signature were overlapping the DS cluster (Figure 1F). Finally, to find out whether the DS cluster could be detected in healthy individuals, we integrated our FCD dataset with single-cell RNA-seq from cortical samples obtained from age-matched autopsy controls[17]. This analysis confirmed that the DS cluster was not present in the normal healthy cortex (Figure 1G). Taken together, these data demonstrate that the DS population is uniquely present in FCD lesions and comprises dysmorphic neurons expressing genes associated with neurodegenerative conditions.

Pathological microglia cell states in FCD IIb

To further characterize microglia states in FCD, we performed subclustering and re-annotation of the broad microglial compartment. Subclustering and marker detection with Seurat uncovered 7 subpopulations with clearly distinct gene signatures (Figure S3; full list of subcluster markers provided in Supplemental Table 2). Diverse microglia signatures identified by single-cell sequencing have been previously reported in neurological diseases[20,21,28]. We used signatures of microglia subsets derived from healthy and epileptic temporal cortex[21] to

annotate subclusters in FCD. Using this approach, we identified three distinct microglia states

in FCD lesions (Figure 4A): a homeostatic subset (the largest population corresponding to

subcluster 0); a subset expressing high levels of *CD74* (subcluster 1, *CD74+,*), and a subset

expressing high levels of *CD83* (subclusters 2 and 5, *CD83+*). We also identified small

subclusters of lymphocyte/T cells, B cells, and perivascular macrophages that were annotated

using classical lymphocyte markers (Figure 4A). As shown in Figure 4B, *CD74+* microglia

expressed marker genes related to antigen processing and presentation/MHC class II genes

(*HLA-DRA*, *HLA-DPA1*, *HLA-DPB1*, *HLA-DRB1*, *HLA-DQA1*), adaptive immune system

response and microglial cell activation (*CD74*, *SC1N*, *C1QA*, *C1QB*, *C1QC*), and interferon

signaling (*IFNGR1*, *JAK2*, *B2M*). On the other hand, *CD83+* microglia expressed high levels

of pro-inflammatory cytokines including *IL1B*, *CCL2*, *CCL3*, *CCL4*, and *TNFSF18*.

Interestingly, *CD83* and *CD74* expression are largely non-overlapping, indicating that these

surface markers can be effectively used to distinguish microglia states in FCD (Figure 4C).

Of note, homeostatic microglia and activated states are expanded only in FCD IIb

tissues (Figure 4D). Compared to controls, FCD IIb had a higher frequency of *CD74+* and

homeostatic subclusters. Compared to FCD IIa, the expansion included *CD74+* and *CD83+*

subpopulations as well as lymphoid/T cells (Figure 4D).

Next, we performed differential expression analysis and identified 133 upregulated

genes in FCD IIb microglia compared to controls (Figure 4E). GO terms associated with

upregulated genes included antigen processing and presentation, MHC class II protein

complex, and positive regulation of cell activation (Figure 4F), in line with the expansion of

CD74+ microglia in FCD IIb. Motif activity analysis using scATAC-seq uncovered that top

upregulated motifs in FCD IIb microglia included IRF8 and STAT1/STAT2 (Figure 4G),

which are regulators related to microglia activation. Cells with the highest IRF8 and

STAT1/STAT2 motif activity overlapped the *CD74+* microglia (Figure 4H). Taken together,

these data highlight the emergence of pathological microglial states is specific to FCD IIb, and

indicated that microglia activation in FCD IIb is mediated by the *CD74+* subpopulation.

Chromatin changes and cell-type specific gene regulation in FCD

Next, we aimed to find open chromatin regions associated with gene regulation in FCD. First,

we found the set of differentially accessible chromatin regions (DACRs), which are peaks with

gain or loss of chromatin accessibility, by comparing chromatin profiles across cell types in

FCD against controls. Overall, FCD IIa had a higher number of DACRs, especially in upper

and deep excitatory neurons (Figure 5A). Most of the DACRs in FCD IIa displayed a gain of

accessibility in the lesion (opening peaks), while DACRs in FCD IIb were characterized by a

loss of chromatin accessibility (Figure 5A). Further analysis using Encode annotation found

that most DACRs were located within enhancer regions, with 34% overlapping proximal

enhancers near a transcriptional start site and 44% overlapping distal enhancers (Figure 5B).

We then sought to understand whether these DACRs were driving gene regulation. To

achieve this, we implemented a machine-learning approach based on random forests to learn

regulatory links using the paired single-cell chromatin and gene expression readouts in the

dataset (see Methods, Figure 5C). The method revealed open chromatin regions most likely to

predict gene expression in each cell type. For instance, we identified regulatory links for *FKBP*

prolyl isomerase 5 (*FKBP5*), a gene within the mTORC1 pathway upregulated in FCD

astrocytes (Figure 5D). The regulatory links involved DACRs with gain of chromatin

accessibility in FCD (regions highlighted in Figure 5D), suggesting that the increased *FKBP5*

expression is accompanied by chromatin opening. These peaks are also putative *FKBP5* cis-

regulatory elements since they are located in regions with enhancer activity according to

Encode. Similarly, we also found opening chromatin peaks located in enhancers linked to

C1QA and *C1QB*, which were genes involved in microglia cell activation increased in FCD IIb

(Figure S4). Taken together, these analyses demonstrated that chromatin changes affecting

enhancer regions are likely driving cell type-specific gene regulation in FCD.

Immature astrocyte cell states and balloon cells in FCD

We explored whether the astrocytic population is affected in FCD by employing an

unsupervised trajectory analysis. The trajectory organized astrocytes according to their

differentiation status, with immature and mature astrocytes on opposite ends of the trajectory

(Figure 6A). The root of the trajectory was chosen based on the expression of immature

astrocyte markers (Figure 6B). Remarkably, the quantification of FCD and control cells along

the trajectory revealed an expansion of immature astrocytes accompanied by a decrease of

mature astrocytes in FCD tissues (Figure 6C). On the other hand, fully mature astrocytes were

highly represented in controls (Figure 6C). We also analyzed the expression of gene signatures

corresponding to balloon cells, which are known to originate from the astrocytic lineage[27] as

well as reactive astrocytes often associated to neurodegenerative diseases[19]. Scores for both

balloon cells and reactive astrocytes signatures were more elevated in astrocytes from FCD

lesions and overlapped with immature states at the beginning of the trajectory (Figure 6D, top).

Of note, immature astrocytes identified as balloon cells were mostly exclusive to FCD IIb

cases, while reactive astrocytes were present in both FCD subtypes (Figure 6D, bottom). Genes

associated with balloon cells and reactive astrocytes including *CHI3L1*, *HSPB1*, *SERPINA3*,

GFAP, and *VIM* were up-regulated in FCD IIb astrocytes compared to FCD IIa or internal

controls (Figure 6E). Taken together, these analyses showed that immature astrocyte cell states,

including balloon cells and reactive astrocytes, are expanded in the disease, indicating that

normal astrocyte differentiation is impaired in FCD.

DISCUSSION

FCD is a leading cause of drug-resistant epilepsy in children and adolescents, accounting for up to 50% of epilepsy surgeries in this age group[10]. This neurological condition was found to be the most common diagnosis among children with drug-resistant focal epilepsy requiring surgical intervention[29]. Despite advances in the genetic characterization of FCD, brain surgery remains the only treatment option to reduce seizures in refractory cases. Our limited understanding of the molecular mechanisms and the role of diverse brain cell types in FCD epileptogenicity is a major obstacle to developing targeted treatments. To address this need, we employed a cutting-edge multi-omics single-cell profiling approach to create a high-resolution cellular census of the human cortex affected by FCD. We leveraged this unique dataset to uncover cellular changes and coordinated chromatin and transcriptional alterations in both neuronal and glial populations in response to the disease. Overall, we uncovered neuronal, microglial and astrocytic pathological cell states involved the disease.

First, our data showed that both excitatory and inhibitory neurons undergo changes in cellular abundance and transcriptional profiles during FCD development, yet neuronal loss affects mostly the EN L2/3 subtype. Interestingly, EN in upper cortical layers were also found to be reduced in cortical samples of patients with multiple sclerosis[30] and temporal lobe epilepsy[31], highlighting their vulnerability not only in FCD but in other neurological conditions.

In addition, we characterized a neuronal population specific to FCD lesions expressing a dysmorphic neuron-signature and neurofilament genes. This neuronal population was notably absent in non-lesion controls and the healthy cortex from autopsy cases. Also, the expression of *NEFM,* encoding the medium polypeptide subunit of the neurofilament protein, was found to be restricted to this subpopulation. Neurofilaments detected in the blood have emerged as reliable diagnostic biomarkers in several neurological diseases[33]. The remarkable specificity of

NEFM expression in dysplastic cells suggests its potential as a blood biomarker for FCD,

prompting the need for additional research to explore its utility as a molecular diagnostic tool

in the disease.

Furthermore, we described the expansion of microglia and the emergence of

pathological cell states specifically in FCD IIb lesions. We identified two FCD-associated

microglial subpopulations with distinct signatures: a *CD74+* subset overexpressing antigen

presentation molecules (MHC class-II components), and a *CD83+* subset with increased

expression of pro-inflammatory cytokines. Previous scRNA-seq studies have identified

microglial populations in neurological conditions[20,21,28]. For instance, using scRNA-seq

followed by in situ immunohistochemistry, Olah et al[21] described the existence of nine

microglial subpopulations in the human cortex (prefrontal cortex of autopsy samples and

temporal cortex of epilepsy), including the *CD83*[high] and *CD74*[high] subsets. Mathys et al[34] also

identified a microglia subset with increased expression of *CD74* and MHCII-related genes in

patients with Alzheimer's disease. A *CD74* subset was also described in the pathology of

multiple sclerosis[35], and *CD74* has been proposed as a marker for reactive microglia due to its

increased expression levels in disease[36]. Although there is no consensus in the field on how to

annotate microglia cell states, we adopted the terminology from Olah et al since the *CD83* and

CD74 molecules were identified as markers of microglia subclusters in FCD. Of note, we also

found higher lymphoid/T-cells infiltration is specific to FCD IIb, in agreement with previous

experimental findings[14].

There is strong emerging evidence of dysregulated neuroinflammation mediated by

microglia in FCD and epilepsy. Zimmer et al[14] showed that FCD IIb elicits a stronger

inflammatory response characterized by increased expression of the *HLA-II* and *CCL2* in

microglia near balloon cells. Kumar et al[37] described immune cell infiltration and increased

expression of pro-inflammatory genes (e.g. *IL1B*, *CCL4*, *CCL2*) in microglia from epilepsy

samples. Therefore, we postulate that the CD74[+] subset (expressing antigen presentation

molecules) and the CD83[+] subset (expressing pro-inflammatory cytokines) identified in this

study are the microglial populations mediating immune system activation and

neuroinflammation in FCD IIb.

Finally, we showed that astrocyte differentiation is impaired in FCD, and immature

astrocytic cell states are expanded in lesions. This shift from mature to immature astrocytes

can be explained in part by the appearance of reactive astrocytes and balloon cells, which are

abnormal cell populations typically found in the lesion microenvironment, that we

characterized as immature astrocytic states. Specifically, pseudotime trajectory analysis

revealed that reactive astrocytes resemble immature-like cells that emerge in FCD lesions,

supporting the view that astrogliosis induces a partial reversion of astrocyte maturation[38].

Similarly, we showed that balloon cells also represent immature astrocytic cell states. Balloon

cells were identified exclusively in FCD IIb, consistent with the histological characterization

of the clinical samples. Moving forward, it will be important to understand the contribution of

these immature astrocytes to FCD. Mature astrocytes are known to maintain neuronal

homeostasis and prevent excessive neuronal activity through neurotransmitter reuptake[38], and

their dysregulation can lead to epileptogenesis[39]. Thus, these immature astrocytic populations

might add to the intrinsic epileptogenicity in FCD.

Paired multimodal single-cell experiments enable exploration of cell-type specific gene

regulation by learning the relationship between chromatin accessibility and transcription in the

same cell[40]. We explored the integration of these paired readouts by predicting cell-type

specific regulatory links and identifying enhancer regions controlling key genes in the

mTORC1 pathway and microglia activation. Indeed, we anticipate that the high-resolution

multimodal dataset produced herein will be useful in future studies integrating chromatin and

transcriptomic data, including unraveling the impact of non-coding variants in open chromatin

regions in gene expression or identifying chromatin enhancers driving gene dysregulation in

FCD.

The findings presented in this study offer promising new directions for the development

of targeted drug treatments for FCD. By elucidating the underlying molecular and cellular

mechanisms common and specific to FCD IIa and IIb at single-cell resolution, this research

provides valuable insights that could inform the development of tailored therapeutic strategies

for each disease subtype. For instance, mTORC1 signaling was found to be elevated in FCD

in most cell types. Thus, mTOR inhibitors or antisense oligonucleotides targeting the mTOR

pathway hyperactivation may represent promising strategies for targeted treatment of FCD[11].

Gene therapies targeting the disease-specific population containing dysmorphic neurons can

also be explored since dysmorphic neurons are believed to play a crucial role in seizure

generation in FCD[3]. Further investigation into marker genes associated with this population

may lead to the identification of pharmacological interventions aimed at improving seizure

control in the disease.

Additionally, several rare neurological conditions lead to epilepsy[41]. These orphan

diseases pose significant challenges to research due to their low prevalence and unresolved

pathology[41]. However, the integration of single-cell datasets may provide a powerful approach

to uncover pathological cell states shared across various forms of rare focal epilepsies. By

comparing cellular profiles, it may be possible to identify common abnormal cell states,

molecular signatures and pathways, offering insights into disease mechanisms and potential

therapeutic targets applicable to multiple epilepsy subtypes[32].

In conclusion, our study uncovered diverse cellular abundance changes in FCD and

identified key neuronal, microglial and astrocytic subpopulations involved in the disease.

These findings improve our understanding of the condition and suggest novel avenues for

developing targeted treatments.

STAR METHODS

Clinical samples and neuropathological diagnosis

Fresh brain samples were collected from patients who underwent surgery for intractable seizures after clinical diagnosis of FCD at Hospital de Clínicas, University of Campinas. Surgical specimens were (i) formalin-fixed paraffin-embedded (FFPE) or (ii) snap-frozen in liquid nitrogen and stored at −80°C. FFPE samples were submitted to diagnostic routine in serial 4 μm-sections stained with hematoxylin and eosin (H&E) and submitted to immunohistochemical reactions. For the latter protocol, the sections were exposed to antibodies against NeuN (neuronal marker; 1:1,000, clone A60, Merck Millipore, cat#MAB377, Temecula, CA, USA), MAP2 (neuronal marker; 1:1,000, clone M13, Thermo Fisher, cat#13-1500, Waltham, MA, USA), SMI 32 (neuronal marker; 1:2,500, clone SMI 32, Biolegend, cat#SMI-32R, San Diego, CA, USA), GFAP (astrocytic marker, 1:100, clone 6F2, Dako/Agilent, cat#M0761, Santa Clara, CA, USA), vimentin (1:100, clone V9, e-Bioscience/Thermo Fisher, cat#14-9897-82, Waltham, MA, USA), CD34 (1:50, clone QBEnd-10, Dako, cat#M7165, Glostrup, Denmark) and CNPase (myeloarchitecture marker; 1:500, clone 11-5 B, Millipore, cat#MAB326, Darmstadt, Germany), for 18 h at 4°C. Then, a detection solution containing the secondary antibody and peroxidase (AdvanceTMHRP®, Dako, cat#K4068, Glostrup, Denmark; or EnvisionTM Flex+, Dako, cat#K8002, Glostrup, Denmark) was added for 30 min at 37°C. 3,3-diaminobenzidine (DAB) was used as a chromogenic substrate and counterstaining was performed with hematoxylin. Negative controls (without primary antibody) were run concurrently with all immunohistochemical reactions.

Samples were classified as FCD IIa or IIb according to the guidelines of the International League Against Epilepsy (ILAE)[2]. Specifically, specimens exhibiting cortical dyslamination, hypertrophic and dysmorphic neurons [disoriented neurons with anomalous

cytoplasmic distribution of Nissl substance and accumulation of non-phosphorylated

neurofilament (SMI 32-positive)] without or with balloon cells (large cells with opaque

eosinophilic cytoplasm, vesicular nucleus and immunopositivity for vimentin) were classified

as FCD ILAE type IIa or IIb, respectively. Internal controls were derived from histologically

normal tissue where neither abnormal cells nor architectural changes were observed. Detailed

patient information is provided in the Supplemental Table 1. All procedures were approved by

the University of Campinas's Research Ethics Board, and written informed consent was

obtained from all patients before surgery.

Nuclei isolation and multiome library construction

The nuclei extraction protocol for human brain tissue was adapted from Jessa et al[42]. Briefly,

frozen tissue was dounce homogenized and incubated for 2 minutes in lysis buffer containing

0.1% NP-40. Then, suspension was filtered through a 30 μm cell strainer (MACS), and washed

in buffer containing 10% BSA to remove debris, followed by a centrifugation at 500g and 4°C

for 5 minutes. Next, the suspension was carefully layered on double 29% and 50% iodixanol

(Sigma) cushion layers, and centrifuged at 10,000g and 4°C for 30 minutes for additional

removal of debris and cell aggregates. Retrieved nuclei were resuspended and counted. Nuclei

were stained with Trypan blue and counted using a hemocytometer. Nuclei integrity was

evaluated under a microscope using 40X or 60X magnifications, and nuclei were considered

viable if they were roundish and with an intact membrane. Next, we proceeded to the steps of

the library construction, including open chromatin transposition and droplet (GEM) formation

using a Chromium Controller equipment (10X Genomics). Libraries were prepared using the

Multiome ATAC + Gene Expression kit (10X Genomics) following the manufacturer's

protocol CG000338 revision E. The quality of purified ATAC and RNA libraries was assessed

using a TapeStation equipment (Agilent Technologies).

Sequencing and raw data processing

Multiome libraries were sequenced on either Illumina HiSeqX Ten or Illumina NovaSeq 6000 instruments according to the sequencing depth and the read length recommended in the Multiome kit protocol (CG000338). Demultiplexing, genome alignment, gene quantification, and peak accessibility in single nuclei were performed with the Cell Ranger ARC pipeline[43]. Specifically, fastq files were generated using cellranger mkfastq, then processed using the cellranger-arc count v.2.0.2 and the human genome GRCh38 to generate the count matrices for the RNA and ATAC modalities. Next, we performed data processing, normalization, and batch integration on both assays independently, using Seurat v.5.0.1[44] and Signac v.1.12.0[45] in the R environment v.4.1.2.

scRNA-seq analysis

Samples were preprocessed in Seurat to remove low-quality nuclei using the following criteria: $400 < \text{nFeatures} < 7,000$, $500 < \text{UMI counts} < 50,000$, and percentage of mitochondrial reads (percent.mt) < 15. Identification of doublets was performed per sample using scDblFinder v.1.8.0[46]. Next, standard Seurat data processing and normalization steps were performed: NormalizeData, FindVariableFeatures, ScaleData, RunPCA, RunUMAP, FindNeighbors, and FindClusters. For each sample, the filtered counts matrix was log-normalized with regression of sequencing mitochondrial gene percentages. The RunUMAP function was used with the first 40 PCs identified in the elbow plot to perform dimensional reduction. Clustering was performed using FindNeighbors and FindClusters functions using the smart local moving (SLM) algorithm, with a resolution of 1.2.

To correct batch effect among samples, we used Harmony v.1.2[24] to integrate sample-level PCA projections using automatic hyperparameter optimization and default parameters.

Harmony implements an algorithm that projects cells into a common space where they cluster

by cell type rather than other sample attributes such as sequencing batch.

Finally, we used SCTransform to create a normalized expression matrix containing all samples,

and with regression of sequencing batch, brain region, and mitochondrial gene percentages.

DEGs between FCD and controls were calculated using MAST or the Wilcox test implemented

in Seurat as indicated. For MAST, DEGs were selected using an adjusted P value of less than

0.05. The Wilcox test was applied using the SCT normalized data, using a log2 fold-change

threshold of 0.25 and an adjusted P value of less than 0.05.

scATAC-seq analysis

Samples were preprocessed in Signac to remove low-quality nuclei using the following criteria:

$1{,}000 < \text{ATAC fragments} < \alpha(\text{ATAC fragments})$, $400 < \text{ATAC fragments in peaks} < \alpha$ (ATAC

fragments in peaks), percent.mt < 15, nucleosome signal < 2, TSS enrichment < 2. Upper limits

(α) of ATAC fragments per sample were computed as mean plus 2 standard deviations. Peaks

within each sample were identified using MACS2[47]. Standard processing and normalization

steps were performed as follows: RunTFIDF, FindTopFeatures, RunSVD, and FindClusters.

For each sample, the peak counts matrix was normalized using the RunTFIDF function, to

correct for differences in nuclei sequencing depth. Dimensional reduction was performed with

latent semantic indexing (LSI) using the RunSVD function. UMAP projection was performed

utilizing LSI components 2-30, as the first LSI component was confirmed to be a technical

variation. Before sample integration, we created a common peak set across samples using

MACS2 and quantified peaks in each sample using the FeatureMatrix function. Peaks in non-

standard chromosomes and blacklisted regions were removed from downstream analyses. We

then re-computed RunTFIDF, FindTopFeatures, and RunSVD on the merged object and

applied Harmony to integrate sample-level low-dimensional cell embeddings (LSI components

2-30) using the same parameters defined in the RNA modality. Differentially accessible

chromatin regions (DACRs) between FCD tissues and controls were calculated using

FindMarkers with the LR test, with a min.pct of 0.05, an adjusted P value less than 0.05, and

included the number of ATAC fragments as a latent variable.

Multimodal data integration

To integrate gene expression data with chromatin accessibility data, we performed Weighted

Nearest Neighbor (WNN) analysis[25], an unsupervised approach that learns the relative utility

of each modality and builds a joint graph representing a weighted combination of RNA and

ATAC modalities. WNN was performed using the FindMultiModalNeighbors function with

20 neighbors (k.nn), based on the UMAP reductions obtained after Harmony integration. The

WNN graph was used for creating a joint UMAP visualization and clustering based on the

WNN graph was performed with the FindClusters function using the smart local moving

(SLM) algorithm. WNN clusters with fewer than 50 nuclei, a high percentage of doublets, or

enriched in mitochondrial markers were removed from downstream analyses.

Cluster annotation

We used the Azimuth R package[25] to perform cell type annotation in the FCD dataset, using as

a reference the Allen atlas of the human cortex[48]. Azimuth performed label-transferring by

assigning to each cell in the FCD dataset the most similar cell type in the reference based on

gene expression similarity. The assigned cell type chosen for each cluster was defined as the

cell type with the highest mean frequency within the cluster. The Azimuth annotation was

manually verified using the marker genes shown in Figure 1C. Cluster and subcluster markers

were calculated using the function FindAllMarkers based on the SCT normalized data, with

the Wilcox test using a log2 fold-change threshold of 0.25 and an adjusted P value of less than

0.05.

Differential cellular abundance

495 We used the function propeller of the Speckle R package[26] to detect changes in cellular

496 composition between FCD tissues and controls. The propeller function calculated cell type

497 proportions in biological replicates, and fitted a linear model for each cell type with the tissue

498 condition and brain cortical region as covariates.

Integration with scRNA-seq from normal cortex

We obtained normal cortex scRNA-seq data from three autopsy donors with no history of

neurological disease from Silleti et al[17]. To reduce computational costs, we randomly sampled

10,000 neuronal and 10,000 non-neuronal cortical nuclei from these donors, and processed this

data along with our FCD dataset following the procedures described in the "scRNA-seq

analysis" section. We used Harmony v.1.2 to integrate all samples with automatic

hyperparameter optimization and default parameters. Clustering was performed with

FindNeighbors and FindClusters functions using the smart local moving (SLM) algorithm and

a resolution of 2.0.

Gene signature analysis

We used the AddModuleScore function from Seurat and the AUCell[49] R package to calculate

gene set enrichment at the nuclei level. The AddModuleScore function from Seurat was

calculated based on the SCT-normalized RNA data. AUCell computes the gene set enrichment

scores based on ranked gene lists (highest to lowest gene expression) in each cell. AUCell

scores were computed using the calcAUC function using the top 5% of genes. The obtained

threshold was used to classify cells with active status for the given signature.

Motif enrichment analysis

We performed motif analysis using chromVAR v. 3.3.2[50] and Signac. Motif activity at the

nuclei level was calculated using the RunChromVAR function in Signac after matching the set

of background peaks. Differential motif activity was computed using FindAllMarkers and

FindMarkers (pairwise comparisons) using the LR test, with a min.pct of 0.05, an adjusted P

value less than 0.05, and included the number of ATAC fragments as a latent variable. We also

filtered context-relevant motifs using the regulator expression levels. The regulator associated

with the motif needed to be expressed in at least 10% of the cells in the cluster.

Trajectory Inference Analysis

Pseudotime trajectory analysis was conducted using Monocle v3[51,52]. Briefly, RNA counts

were preprocessed using the preprocess_cds function with the PCA method. To correct batch

effects, we used the align_cds function[53] with the sample as an alignment group and a residual

formula including the sequencing batch, brain region, and mitochondrial genes percentage as

covariates. Nuclei were then projected onto the UMAP space[54] and divided into clusters and

partitions with the cluster_cells function[55]. Subsequently, an unsupervised trajectory was

identified using the SimplePPT (Probabilistic Pseudotime Trajectory) algorithm implemented

in the learn_graph function[56], with the root set according to the expression of immature cellular

markers. Pseudotime values were automatically assigned to individual cells based on their

distance along the trajectory relative to the root node. Genes with differential expression along

the trajectory were calculated using the graph_test function, with adjusted P-value less than

0.05 and arranged according to Moran's I statistic. For each of these genes, differential

expression between tissue conditions was assessed using the fit_models and coefficient_table

functions, using a regression test with an adjusted P value less than 0.05, minimum fold-change

of 1.5, and with the number of counts, sequencing batch, brain region and mitochondrial gene

percentages as covariates.

545 Inference of regulatory links

We used a regression approach based on random forests to find open chromatin regions

associated with gene regulation. The goal was to predict the expression levels of the target gene

based on the accessibility of ATAC peaks (i.e. cis-regulatory elements) around the gene

promoter. Considering that a target gene g has n chromatin peaks, the linear regression was

defined as:

$$x_{gi} = \sum_{i=1}^{n} w_j a_{ji} + \epsilon_i$$

where x_{gi} is the expression of g in cell i, a_{ji} is the accessibility of a given peak j in cell i, w_j is

the weight and ϵ_i is the noise term. For the regression, we tested all peaks located in a 500kb

window around the gene's transcriptional start site (TSS), to include both proximal and distal

regulatory elements. The optimal weights, which were associated with the importance of the

cis-regulatory regions as predictors of gene expression, were determined using random forests.

Specifically, the regression was implemented using the GENIE3 R package[49], setting the

number of trees to 200, and peaks with weights in the top 5% (95% percentile) for each gene

were selected. We used the Encode annotation[57] downloaded from https:

//screen.encodeproject.org to identify open chromatin regions with potential enhancer activity.

Encode enhancers are classified based on their relative location to the TSS as proximal

enhancer-like sequences (pELS) if located within 200bp $<$ TSS $\leq$ 2kb, or distal enhancer-like

sequence (dELS) when located more than 2kb from a TSS. We used both pELS and dELS

datasets for annotation.

REFERENCES

1. Blümcke, I. *et al.* The clinicopathologic spectrum of focal cortical dysplasias: A consensus classification proposed by an ad hoc Task Force of the ILAE Diagnostic Methods Commission. *Epilepsia* **52**, 158–174 (2011).

2. Najm, I. *et al.* The ILAE consensus classification of focal cortical dysplasia: An update proposed by an ad hoc task force of the ILAE diagnostic methods commission. *Epilepsia* **63**, 1899–1919 (2022).

3. Ii, P. H. I. & Crino, P. B. Focal Cortical Dysplasia: Gene Mutations, Cell Signaling, and Therapeutic Implications. *Annual Review of Pathology: Mechanisms of Disease* **12**, 547–571 (2017).

4. Orlova, K. A. *et al.* Early Progenitor Cell Marker Expression Distinguishes Type II From Type I Focal Cortical Dysplasias. *J Neuropathol Exp Neurol* **69**, 850–863 (2010).

5. Hanai, S. *et al.* Abnormal maturation of non-dysmorphic neurons in focal cortical dysplasia: Immunohistochemical considerations. *Seizure* **19**, 274–279 (2010).

6. Lasarge, C. L. & Danzer, S. C. Mechanisms regulating neuronal excitability and seizure development following mTOR pathway hyperactivation. *Frontiers in Molecular Neuroscience* **7**, (2014).

7. Powell, J. D., Pollizzi, K. N., Heikamp, E. B. & Horton, M. R. Regulation of Immune Responses by mTOR. *Annual Review of Immunology* **30**, 39–68 (2012).

8. Weichhart, T., Hengstschläger, M. & Linke, M. Regulation of innate immune cell function by mTOR. *Nat Rev Immunol* **15**, 599–614 (2015).

9. Lin, Y. *et al.* Similar PDK1–AKT–mTOR pathway activation in balloon cells and dysmorphic neurons of type II focal cortical dysplasia with refractory epilepsy. *Epilepsy Research* **112**, 137–149 (2015).

10. Lim, J. S. *et al.* Brain somatic mutations in MTOR cause focal cortical dysplasia type II

leading to intractable epilepsy. *Nat Med* **21**, 395–400 (2015).

11. Gerasimenko, A., Baldassari, S. & Baulac, S. mTOR pathway: Insights into an established

pathway for brain mosaicism in epilepsy. *Neurobiology of Disease* **182**, 106144 (2023).

12. Baulac, S. *et al. Single-Cell Genotyping and Transcriptomic Profiling in Focal Cortical*

Dysplasia. https://www.researchsquare.com/article/rs-4014535/v1 (2024)

doi:10.21203/rs.3.rs-4014535/v1.

13. Boer, K. *et al.* Evidence of activated microglia in focal cortical dysplasia. *Journal of*

Neuroimmunology **173**, 188–195 (2006).

14. Zimmer, T. S. *et al.* Balloon cells promote immune system activation in focal cortical

dysplasia type 2b. *Neuropathol Appl Neurobiol* **47**, 826–839 (2021).

15. Galvão, I. C. *et al.* Identifying cellular markers of focal cortical dysplasia type II with cell-

type deconvolution and single-cell signatures. *Sci Rep* **13**, 13321 (2023).

16. Bakken, T. E. *et al.* Comparative cellular analysis of motor cortex in human, marmoset

and mouse. *Nature* **598**, 111–119 (2021).

17. Siletti, K. *et al.* Transcriptomic diversity of cell types across the adult human brain.

Science **382**, eadd7046 (2023).

18. Walter, T. J., Suter, R. K. & Ayad, N. G. An overview of human single-cell RNA sequencing

studies in neurobiological disease. *Neurobiology of Disease* **184**, 106201 (2023).

19. Liddelow, S. A. *et al.* Neurotoxic reactive astrocytes are induced by activated microglia.

Nature **541**, 481–487 (2017).

20. Keren-Shaul, H. *et al.* A Unique Microglia Type Associated with Restricting Development

of Alzheimer's Disease. *Cell* **169**, 1276-1290.e17 (2017).

21. Olah, M. *et al.* Single cell RNA sequencing of human microglia uncovers a subset

associated with Alzheimer's disease. *Nat Commun* **11**, 6129 (2020).

22. Flynn, E., Almonte-Loya, A. & Fragiadakis, G. K. Single-Cell Multiomics. *Annual Review of*

Biomedical Data Science **6**, 313–337 (2023).

23. Armand, E. J., Li, J., Xie, F., Luo, C. & Mukamel, E. A. Single-Cell Sequencing of Brain Cell

Transcriptomes and Epigenomes. *Neuron* vol. 109 11–26 (2021).

24. Korsunsky, I. *et al.* Fast, sensitive and accurate integration of single-cell data with

Harmony. *Nat Methods* **16**, 1289–1296 (2019).

25. Hao, Y. *et al.* Integrated analysis of multimodal single-cell data. *Cell* **184**, 3573-3587.e29

(2021).

26. Phipson, B. *et al.* propeller: testing for differences in cell type proportions in single cell

data. *Bioinformatics* **38**, 4720–4726 (2022).

27. Baulac, S. *et al.* Single-cell genotyping and transcriptomic profiling in focal cortical

dysplasia. Preprint at https://doi.org/10.21203/rs.3.rs-4014535/v1 (2024).

28. Paolicelli, R. C. *et al.* Microglia states and nomenclature: A field at its crossroads. *Neuron*

110, 3458–3483 (2022).

29. Blumcke, I. *et al.* Histopathological Findings in Brain Tissue Obtained during Epilepsy

Surgery. *N Engl J Med* **377**, 1648–1656 (2017).

30. Schirmer, L. *et al.* Neuronal vulnerability and multilineage diversity in multiple sclerosis.

Nature **573**, 75–82 (2019).

31. Pfisterer, U. *et al.* Identification of epilepsy-associated neuronal subtypes and gene

expression underlying epileptogenesis. *Nat Commun* **11**, 5038 (2020).

32. Sizemore, G. *et al.* Temporal Lobe Epilepsy, Stroke, and Traumatic Brain Injury:

Mechanisms of Hyperpolarized, Depolarized, and Flow-Through Ion Channels Utilized as

Tri-Coordinate Biomarkers of Electrophysiologic Dysfunction. *OBM Neurobiol* **2**, 009

(2018).

33. Khalil, M. *et al.* Neurofilaments as biomarkers in neurological disorders — towards

clinical application. *Nat Rev Neurol* **20**, 269–287 (2024).

34. Mathys, H. *et al.* Single-cell transcriptomic analysis of Alzheimer's disease. *Nature* **570**,

332–337 (2019).

35. Masuda, T. *et al.* Spatial and temporal heterogeneity of mouse and human microglia at

single-cell resolution. *Nature* **566**, 388–392 (2019).

36. Potru, P. S. & Spittau, B. CD74: a prospective marker for reactive microglia? *Neural*

Regen Res **18**, 2673–2674 (2023).

37. Kumar, P. *et al.* Single-cell transcriptomics and surface epitope detection in human brain

epileptic lesions identifies pro-inflammatory signaling. *Nat Neurosci* **25**, 956–966 (2022).

38. Lattke, M. & Guillemot, F. Understanding astrocyte differentiation: Clinical relevance,

technical challenges, and new opportunities in the omics era. *WIREs Mech Dis* **14**, e1557

(2022).

39. Verhoog, Q. P., Holtman, L., Aronica, E. & van Vliet, E. A. Astrocytes as Guardians of

Neuronal Excitability: Mechanisms Underlying Epileptogenesis. *Front. Neurol.* **11**, (2020).

40. Baysoy, A., Bai, Z., Satija, R. & Fan, R. The technological landscape and applications of

single-cell multi-omics. *Nat Rev Mol Cell Biol* **24**, 695–713 (2023).

41. Kioutchoukova, I. P. *et al.* Neurologic orphan diseases: Emerging innovations and role

for genetic treatments. *World J Exp Med* **13**, 59–74 (2023).

42. Jessa, S. *et al.* Stalled developmental programs at the root of pediatric brain tumors. *Nat*

Genet **51**, 1702–1713 (2019).

43. Zheng, G. X. Y. *et al.* Massively parallel digital transcriptional profiling of single cells. *Nat*

Commun **8**, 14049 (2017).

44. Dictionary learning for integrative, multimodal and scalable single-cell analysis | Nature

Biotechnology. https://www.nature.com/articles/s41587-023-01767-y.

45. Stuart, T., Srivastava, A., Madad, S., Lareau, C. A. & Satija, R. Single-cell chromatin state

analysis with Signac. *Nat Methods* **18**, 1333–1341 (2021).

46. Germain, P.-L., Lun, A., Garcia Meixide, C., Macnair, W. & Robinson, M. D. Doublet

identification in single-cell sequencing data using scDblFinder. *F1000Res* **10**, 979 (2021).

47. Zhang, Y. *et al.* Model-based Analysis of ChIP-Seq (MACS). *Genome Biology* **9**, R137

(2008).

48. Comparative cellular analysis of motor cortex in human, marmoset and mouse | Nature.

https://www.nature.com/articles/s41586-021-03465-8.

49. Aibar, S. *et al.* SCENIC: single-cell regulatory network inference and clustering. *Nat*

Methods **14**, 1083–1086 (2017).

50. Schep, A. N., Wu, B., Buenrostro, J. D. & Greenleaf, W. J. chromVAR: inferring

transcription-factor-associated accessibility from single-cell epigenomic data. *Nature*

Methods **14**, 975–978 (2017).

51. Trapnell, C. *et al.* The dynamics and regulators of cell fate decisions are revealed by

pseudotemporal ordering of single cells. *Nat Biotechnol* **32**, 381–386 (2014).

52. Cao, J. *et al.* The single-cell transcriptional landscape of mammalian organogenesis.

Nature **566**, 496–502 (2019).

53. Haghverdi, L., Lun, A. T. L., Morgan, M. D. & Marioni, J. C. Batch effects in single-cell

RNA-sequencing data are corrected by matching mutual nearest neighbors. *Nat*

Biotechnol **36**, 421–427 (2018).

54. McInnes, L., Healy, J. & Melville, J. UMAP: Uniform Manifold Approximation and

Projection for Dimension Reduction. Preprint at http://arxiv.org/abs/1802.03426 (2020).

55. Traag, V. A., Waltman, L. & van Eck, N. J. From Louvain to Leiden: guaranteeing well-

connected communities. *Sci Rep* **9**, 5233 (2019).

56. Qiu, X. *et al.* Reversed graph embedding resolves complex single-cell trajectories. *Nat*

Methods **14**, 979–982 (2017).

57. Moore, J. E. *et al.* Expanded encyclopaedias of DNA elements in the human and mouse

genomes. *Nature* **583**, 699–710 (2020).

Acknowledgments

We thank the patients and their families who participated in this study. We thank Claudia Kleimann, Steven Hebert, Bhavyaa Chandarana and Tomas V. Waichman for advice on single-cell analysis; Damien Faury for sharing the nuclei extraction protocol; Danielle C. Bruno for discussions regarding the tissue dissociation. This work was supported by the following grants: Fundação de Amparo à Pesquisa do Estado de São Paulo (FAPESP) Young Investigator grant 2019/07382-2 (D.F.T.V); the Chan Zuckerberg Initiative DAF, an advised fund of the Silicon Valley Community Foundation grant DAF2021-237598 (D.F.T.V.); FAPESP doctoral fellowship 2022/01530-2 (I.C.G); Coordenação de Aperfeiçoamento de Pessoal de Nível Superior (CAPES) master fellowship (L.A.M.); Conselho Nacional de Pesquisa (CNPq) grant 311923/2019-4 (I.L.-C.); FAPESP doctoral fellowship 2020/06168-4 (J.C.G.), FAPESP regular grant 2019/08259-0 (F.R.), and FAPESP Research Innovation and Dissemination Center grant 2013/07559-3 (F.C.).

Author contributions

I.C.G. and D.F.T.V. designed the study. L.A.M. and M.L. performed single-cell assays. I.C.G. and D.F.T.V. performed data analyses. P. A., J.C.G., C.L.Y., E.G., H.T., M.K.M.A., J.C.G, F.C., and I.L-C contributed to clinical sample collection. F.R. performed histopathological analyses. I.C.G. and D.F.T.V. wrote the manuscript. All authors reviewed and approved the final version of the manuscript.

Competing interests

The authors declare no competing interests.

 FIGURES

Figure 1. Overview of the multimodal single-cell sequencing in FCD type II.
727

(A) Schematic of FCD samples and study design. Cortical tissues from lesions and adjacent non-lesion areas in individuals with FCD IIa and IIb were profiled by multiome single-cell ATAC and RNA sequencing. Created with Biorender.com.

(B) Uniform manifold approximation and projection (UMAP) visualization of joint modalities (ATAC + RNA) after sample integration. Nuclei are colored by sequencing batch.

(C) UMAP joint visualization of nuclei colored by broad cell type annotation.

(D) UMAP joint visualization colored using a high-resolution annotation with classification of neuronal subtypes.

(E) UMAP plots depicting nuclei by tissue condition.

(F) Dot plot showing gene expression of known marker genes for major cortical cell types. Dot size corresponds to the fraction of cells expressing the marker, and color corresponds to the expression levels.

(G) Coverage plots of marker genes depicting chromatin accessibility across cell types. The normalized ATAC signal is depicted in a region of +/- 1kb around gene coordinates.

(H) Box plots showing cell type proportion by tissue condition. In box plots, the median is indicated by the center line; box limits represent upper and lower quartiles; and whiskers extend to 1.5 times the interquartile range. Overlay dots represent cell type proportions in individual samples. Significant changes in cell types were detected using Speckle. ˙$P < 0.1$, *$P < 0.5$, **$P < 0.01$, ***$P < 0.001$.

Figure 2. Excitatory and inhibitory neuron subtypes affected in FCD.

(A) Box plots showing excitatory neurons (EN) subtype frequency in FCD and controls. In box plots, the median is indicated by the center line; box limits represent upper and lower quartiles; and whiskers extend to 1.5 times the interquartile range. Overlay dots represent proportions in individual samples. Significant changes were detected using Speckle. ˙P < 0.1, *P < 0.5, **P < 0.01, ***P < 0.001.

(B) Bar graphs depicting the number of differentially expressed genes (DEGs) in EN subtypes. DEGs between FCD and controls were computed using MAST with adj. P < 0.05. Only the five subtypes with the most transcriptional changes are shown.

(C) Box plots showing inhibitory neurons (IN) subtype frequency in FCD and controls. Box plots legends and significance levels are defined as in (A).

(D) Bar graphs depicting DEGs in IN subtypes. DEGs between FCD and controls were computed using MAST with adj. P < 0.05. Only the five subtypes with the most transcriptional changes are shown.

Figure 3. Characterization of a disease-specific cluster containing dysmorphic neurons.

(A) UMAP highlighting the disease-specific (DS) cluster. The pie chart indicates the percentage of nuclei from FCD IIa, IIb, and controls within the cluster.

(B) Pathway enrichment of DS cluster markers. Marker genes were identified using Seurat´s FindMarkers and KEGG pathway enrichment was performed using g:Profiler. The bars represent the significance score of pathway enrichment, and the color indicates the number of genes in each pathway.

(C) Dot plot denoting expression of marker genes associated with neurons in the DS cluster. Neuronal genes are grouped by biological function. Dot size and color correspond to expression frequency and level in the cluster, respectively.

(D) Violin plots denoting gene expression of *NEFM* and *NEFL* neurofilaments in the various cell types.

(E) Coverage plots of neuronal genes expressed in the DS cluster. The normalized ATAC signal is depicted in a region of +/- 1kb around gene coordinates.

798 **(F)** UMAP showing the average expression of a dysmorphic neuron (DN) signature obtained
799 from Baldassari et al. Signature expression was calculated using the AddModuleScore function
800 from Seurat.
801
802 **(G)** UMAP RNA plots showing the DS cluster population in FCD IIa and IIb lesions,
803 histologically normal tissue from FCD individuals (internal controls), and healthy cortex
804 (autopsy controls). The leftmost UMAP depicts annotated cell types in the integrated dataset,
805 which was created by integrating scRNA-seq data from this study with Siletti et al using
806 Harmony.
807
808
809

810
Figure 4 – Microglia pathological cell states and activation in FCD type IIb.

(A) UMAP visualization of microglia and lymphoid subclusters sequenced in FCD lesions.

(B) Feature plots depicting expression of key genes distinguishing microglia subsets *CD83*⁺ (*CD83, CCL2, CCL3*) and *CD74*⁺ (*CD74, HLA-DRA, HLA-DPA1*).

(C) Dot plot denoting the expression of key marker genes in microglia and lymphoid subclusters.

(D) Box plots denoting subcluster frequency by tissue condition. Significant changes were detected using Speckle. Box plots and significance legends are defined in Fig. 1H.

(E) Volcano plot depicting differentially expressed genes (DEGs) in FCD IIb microglia. DEGs were obtained using a Wilcox test after SCT transformation with correction for sequencing batch, mitochondrial gene percentage, and brain region. The *x*-axis is the fold-change (log2) expression in FCD IIb *vs* controls, and the *y*-axis depicts the significance of the change (-log10 of the adjusted P-value). The color indicates the gene status.

(F) Dot plots showing top enriched GO terms in up- and down-regulated genes from (E). Dot size and color correspond to the number of genes associated with the GO term and enrichment P-value, respectively.

(G) Top five differential motifs in microglia from FCD IIb versus controls, as measured by chromVAR (see Methods).

(H) Feature plots depicting IRF8 and STAT1/STAT2 motif activity in microglia open chromatin regions from FCD IIb and controls. Cells are colored by the chromVAR score.

840
841 **Figure 5. Open chromatin changes and cell-type specific regulatory links in FCD.**
842

843 **(A)** Barplots depicting differentially accessible chromatin regions (DACRs) between FCD
844 tissues and controls. The number of chromatin regions with opening or closing status (gain or
845 loss of chromatin accessibility, respectively) is indicated for each cell type.
846

847 **(B)** Enhancer types overlapping DACRs. The number of opening and closing peaks associated
848 with distal or proximal enhancers from Encode are indicated.
849

850 **(C)** Diagram depicting the inference of regulatory links identifying neighboring open
851 chromatin regions (ATAC peaks) associated with gene expression using linear regression (see
852 Methods).
853

854 **(D)** Regulatory links predicted for *FKBP5* in astrocytes. ATAC coverage tracks and
855 corresponding expression levels in controls, FCD IIb, and IIa tissues are depicted at the top.
856 Regions highlighted in gray indicate opening peaks in lesions and red asterisks indicate
857 Encode-annotated enhancers. The gene track shows the location of the *FKBP5* locus in
858 chromosome 6, and the bottom track shows all peaks contained in the chromosomal region.
859

860
861 **Figure 6. Impaired Astrocyte Differentiation in FCD.**
862

863 **(A)** UMAP visualization of astrocytes temporally ordered on a trajectory as a function of
864 pseudotime built by Monocle3.
865

866 **(B)** Feature plots depicting gene expression of immature astrocyte markers *CD44*, *TNC*, *VCAN*,
867 and *GFAP* (top), and differentiated astrocyte markers *SLC1A2*, *SLC1A3*, and *GPC5* (bottom).
868

869 **(C)** UMAP visualization and density quantification along the trajectory. Density plots indicate
870 the normalized density of cells along the pseudotime according to tissue condition.
871
872

873   **(D)** Feature plots depicting expression of balloon cells and reactive astrocytes gene signatures
874   computed by AddModuleScore from Seurat (top). Bar graphs (bottom) quantify the proportion
875   of active cells for these signatures along the pseudotime trajectory. Active cells were defined
876   using the AUCell method and are colored by tissue condition.
877
878   **(E)** Expression levels of genes associated with balloon cells and reactive astrocytes *CHI3L1*,
879   *HSPB1*, *SERPINA3*, *GFAP*, and *VIM* in the trajectory and by tissue condition.