1

A refined human linear B cell epitope map of Outer surface protein C (OspC) from the Lyme disease spirochete, *Borreliella burgdorferi*

Grace Freeman-Gallant[1,#], Kathleen McCarthy[2, #], Jennifer Yates[1,2], Karen Kulas[1], Michael J. Rudolph[3], David J Vance[1,2], and Nicholas J Mantis[1,2,*]

[1] Division of Infectious Diseases, Wadsworth Center, New York State Department of Health, Albany, NY; [2] Department of Biomedical Sciences, University at Albany, Albany, NY;[3] New York Structural Biology Center, New York, NY

[#], these authors contributed equally to this study with last names listed alphabetically

Running title: Linear B cell epitopes on OspC

*, address correspondence to Dr. Nicholas Mantis, Division of Infectious Diseases, Wadsworth Center, New York State Dept Health, 120 New Scotland Ave, Albany, NY 12208; Tel: 518-473-7487; Email:nicholas.mantis@health.ny.gov

ABSTRACT

A detailed understanding of the human antibody response to Outer surface protein C (OspC) of *Borrelellia burgdorferi* has important implications for Lyme disease diagnostics and vaccines. In this report, a total of 13 peptides encompassing eight reported OspC linear B cell epitopes from OspC types A, B and K, including the conserved C-terminus (residues 193-210: peptide C10), were evaluated by multiplex immunoassay (MIA) for IgG reactivity with ~700 human serum samples confirmed positive in a two-tiered Lyme disease diagnostic assay and ~160 post-treatment Lyme disease (PTLD) serum samples. The VlsE C6-17 peptide was included as a positive control. Diagnostic serum IgG reacted with 11 of the 13 OspC-derived peptides, significantly more than controls, with the C10 peptide being the most reactive. In the PTLD serum samples, two OspC peptides including C10 were significantly more reactive than controls. Spearman's rank correlation matrices and hierarchical clustering indicated a strong correlation between C10 and VlsE C6-17 peptide reactivity but little demonstrable association between C10 and the other OspC peptides or recombinant OspC. OspC peptide reactivities (excluding C10) were strongly correlated with each other and were disproportionately influenced by a subset of pan-reactive samples. In the PTLD cohort, C10 clustered with the other OspC-derived peptides and was distinct from OspC and VlsE C6-17. The asynchronous serologic response to OspC, C10, and the OspC-derived peptides reveals the complexity of B cell responses to *B. burgdorferi* and confounds simple interpretation of antibody profiles associated with Lyme disease.

IMPORTANCE

Lyme disease is an emerging tick-borne infection caused by the spirochete, *Borreliella burgdorferi*. In humans, antibodies against spirochetal outer surface lipoproteins are proposed to play a role in disease resolution and in protection against reinfection. Some of those same antibodies also serve as diagnostic indicators of an active or history of Lyme disease. In this study, we sought to validate reported antibody binding sites on Outer surface protein C (OspC), a known target of both protective and diagnostic antibodies.

KEYWORDS: Lyme disease, antibody, epitope, vaccine, *Borrelia burgdorferi*

INTRODUCTION

Lyme (borreliosis) disease is the most common vector-borne infection in the United States, with an estimated 450,000 cases per year (1). The primary etiologic agent of Lyme disease is the spirochetal bacterium, *Borreliella burgdorferi*. In North America, the spirochete is transmitted to humans by black legged ticks, *Ixodes scapularis* and *Ixodes pacificus*, during the course of a blood meal. The spirochete proliferates at the site of the tick bite, typically resulting in an expanding skin lesion commonly referred to as a bull's eye rash or erythema migrans (2-4). In the absence of antibiotic intervention, *B. burgdorferi* can disseminate to peripheral tissues, organs, large joints, and the central nervous system, potentially resulting in severe complications including neuroborreliosis, carditis and/or Lyme arthritis weeks, months or even years later (2, 5). Moreover, a fraction of Lyme disease patients who receive a full regimen of antibiotics will report persistent health issues (e.g., fatigue, cognitive issues, musculoskeletal pain), a syndrome termed post-treatment Lyme disease (PTLD) (6-9). Developing rapid and accurate diagnostic tests capable of detecting and discriminating between LD, PTLD syndrome and *B. burgdorferi* reinfection are much needed (2l, 10).

The humoral response to *B. burgdorferi* is robust, resulting in detectable spirochete-specific B cells and serum IgM and IgG days and weeks following an infectious tick bite (11-14). In Lyme disease patients, antibodies are primarily directed against *B. burgdorferi*'s panoply of outer surface lipoproteins (15). Among these, Outer surface protein C (OspC; also known as BB_B19, P23 and P25) stands out. OspC is a ~21 kDa helical homodimeric lipoprotein expressed by *B. burgdorferi* during tick transmission and in the early stages of infection (16). OspC is implicated in facilitating spirochete egress from the tick during the course of a blood meal, enabling survival in the early stages of mammalian skin infection, and modulating transmigration across vascular walls (16-23). The highly immunogenic nature of OspC has lent itself to applications in Lyme disease serology-based diagnostics, including widely used commercial tests (24, 25). For example, the peptide derived from the conserved C-terminal 10 amino acids of OspC (peptide "C10") is a component of the FDA approved diagnostic test *Borrelia* VlsE1/pepC10 from Zeus Scientific (Branchburg, NJ) (26-30). In addition to their diagnostic utility, linear B cell epitopes from different OspC types are a component of a widely used canine Lyme disease vaccine and being considered for human use (see references within (31)).

While multiple linear human B cell epitopes on OspC have been reported over the past

three decades, differences in serologic assays, detection methodologies, and sample sizes from

varying Lyme disease cohorts, makes it difficult to draw conclusions about relative reactivities of

one peptide over another. This is problematic, because determining the relationships between

OspC-derived peptides (including C10) and total OspC antibodies has practical implications for

interpreting serologic assays. Moreover, there are >26 allelic variants or types of OspC reported

in the United States, with amino acid sequence identities that range from 60-90% (32, 33). As

different OspC types are associated with varying degrees of virulence and invasiveness, tracking

epitope-specific responses to those particular types is of paramount importance (34, 35). In New

York state, for example, OspC types A, B and K, which are associated with more invasive

disease, represent ~70% of all isolates (35-37).

With that in mind, we sought to validate human antibody reactivities to reported linear epitopes

associated with OspC to enable broad comparability of epitope usage across *B. burgdorferi* for

diagnostic and vaccine development purposes.

METHODS

Chemicals and biological reagents. Chemicals and reagents were obtained from Thermo Fisher

Scientific (Waltham, MA), unless noted otherwise. Buffers were prepared by the Wadsworth

Center's Cell and Tissue Culture core facility.

Recombinant OspC and OspC-derived peptides. Recombinant *B. burgdorferi* $OspC_A$

(residues 38 to 201; PDB ID 1GGQ; UniProt ID Q07337) (38), $OspC_B$ (residues 38 to 202; *B.*

burgdorferi strain ZS7; PDB ID 7UJ2) (39) and $OspC_K$ (residues 38 to 202; *B. burgdorferi* strain

297; PDB ID 7UJ6) (40) were expressed in *E. coli* strain BL21 (DE3) and purified by nickel-

affinity and size-exclusion chromatography, as described (41). Recombinant $OspC_A$ with the C10

sequence (residues 201-210; PVVAESPKKP; "$OspC_A$+C10") was expressed and purified as

above. Linear epitope prediction was done using Discotope 2.0 (42). OspC peptides (>80%

purity) were synthesized by Genemed Synthesis (San Antonio, TX) with a C-terminal GGGSK-

biotin extension. We also synthesized the VlsE-derived C6 B31-17 peptide

[MKKDDQIAAAIALRGMA] with the GGGSK-biotin linker (43).

Commercial and clinical LD serum samples. Commercial Lyme disease seronegative (Lot

10500586) and seropositive (Lot 10510438) pooled samples were used strictly as intra-assay and

bead coupling confirmation controls throughout this study (ACCURUN products 810 and 130,

respectively; SeraCare, Milford, MA). The Lyme disease seronegative samples (referred to as

"controls" throughout the manuscript) consist of a commercial panel of 81 serum samples

collected in 2017-2018 (Access Biologicals, Vista, CA). Five of the 81 serum samples were

classified as extreme outliers for VlsE C6-17 reactivity by interquartile range (IQR) calculated

using Microsoft Excel and therefore removed from the control panel. In the end, a total of 76

control samples were used as comparators with the diagnostic sample set and 75 samples were

used as comparators for the PTLD sample set as a result of sample availability.

     B. burgdorferi seropositive serum samples (n = 696) were obtained from the New York

State Department of Health Wadsworth Center's Diagnostic Immunology Laboratory. The

samples had been submitted for Lyme disease serology and subjected to two-tiered testing

consisting of (Tier 1) a C6 peptide screen (Immunetics®; C6 Lyme ELISA™) or Enzyme

Linked Fluorescent Assay (ELFA; BioMerieux, VIDAS® Lyme IgG II and Lyme IgM II;

Durham, NC), followed by (Tier 2) IgM and IgG detection by Western blot (MarDX®; Trinity

Biotech, Carlsbad, CA). *B. burgdorferi*-specific IgM reactivity was defined as $\geq$2 positive bands,

with IgG reactivity defined as $\geq$5 positive bands. Serum samples were aliquoted, de-identified,

and classified as IgM-positive/IgG-negative, IgM-positive/IgG-positive, or IgM-negative/IgG-

positive, based on the Western blot results. Post-treatment Lyme disease (PTLD) serum samples

(n=158) were kindly provided by the Lyme Disease Biobank at Nuvance Health® (Danbury, CT).

PTLD was defined as described by Aucott (8).

Multiplexed OspC and OspC peptide microsphere immunoassays (MIA). Recombinant

OspC$_A$, OspC$_B$, or OspC$_K$ (5 µg) were coupled to Magplex-C microspheres (1 x 10^6) using sulfo-

NHS (N-hydroxysulfosuccinimide) and EDC [1-ethyl-3-(3-dimethylaminopropyl) carbodiimide

hydrochloride], as recommended by the manufacture (Luminex Corp., Austin, TX). Coupled

beads were diluted in xMap® AbC Wash Buffer to a concentration of 5 x 10^6 beads/ml. Biotin-

labeled peptides were complexed to Magplex®-avidin microspheres following protocols provided

by the manufacturer (Luminex Corp.). Microspheres were resuspended in 250 µL of assay buffer

(1 x PBS, 2% BSA, pH 7.4) then subjected to vortexing and sonication. A total of 1×10^6 beads

in assay buffer was mixed with biotin-conjugated peptides (~5 μg) and incubated for 30 min at

room temperature. The microsphere suspensions were then washed three times using wash buffer

(1 x PBS, 2% BSA, 0.02% TWEEN-20, 0.05% Sodium azide, pH 7.4) and a magnetic separator,

resuspended in 500 μL of assay buffer, and stored at 4°C until use. For experimental use, assay

buffer was used to dilute bead stocks (1:50) and human serum (1:100). The bead dilution (50 μL)

and diluted serum (50 μL) were combined in black, clear-bottomed, non-binding, chimney 96-

well plates (Greiner Bio-One, Monroe, North Carolina) and allowed to incubate for 60 min in a

tabletop shaker (600 rpm) at room temperature. Plates were washed three times using a magnetic

separator and wash buffer. Secondary antibody goat anti-Human IgG Fc, eBioscience

(Invitrogen, Carlsbad, California) was diluted 1:500 in assay buffer and added (100 μL) to each

well. The secondary antibody was allowed to incubate for 30 min in a tabletop shaker (600 rpm)

at room temperature. Plates were washed as stated above and beads were resuspended in 100 μL

of wash buffer. Samples were analyzed using a FlexMap 3D (Luminex Corp.).

Statistical analysis. Statistical analyses were performed using R 4.3.0 with R packages readxl

and tidyverse (44-46). MFIs (log_{10}) were first subjected to the Shapiro-Wilk's test to assess

normality, then either Levene's or Fligner-Killeen tests to compare variances. Statistical

significance was determined by Mann-Whitney U tests with an alpha level of 0.05 using the R

package rstatix [https://CRAN.R-project.org/package=rstatix]. R package corrplot (47) was used

to generate the Spearman's Rank correlation matrix. R package pheatmap was used to create the

hierarchically clustered heatmaps (48). R packages ggthemes

[https://github.com/jrnold/ggthemes], RColorBrewer [https://cran.r-

project.org/web/packages/RcolorBrewer/index.html], and ggpubr [https://CRAN.R-

project.org/package=ggpubr] were used in formatting.

Molecular modeling. PyMol (DeLano Scientific LLC, Palo Alto, CA) was used for epitope

modeling using OspC PDB ID: **1GGQ** (strain B31, OspC$_A$).

RESULTS

We sought to validate the reactivity of OspC linear B cell epitopes, including those

already accessioned in the Immune Epitope Database (iedb.org), that have been directly or

indirectly implicated in Lyme disease diagnostics and/or immunity to *B. burgdorferi*. Eight

different epitopes, including the C10 peptide, were chosen for analysis **(Table 1)**. When localized

onto the structure of OspC$_A$ using PyMol, the epitopes represent ~30% of the surface area of the

molecule (**Figure 1A**). An alignment of the primary amino acid sequences of the three OspC

types (A, B, K) associated with invasive disease in northeast United States revealed

polymorphisms in a number of these epitopes (**Table 1**; **Figure 1B**) (49, 50). For this reason, a

total of 13 different peptides were generated to encompass the specific amino acid sequences for

each epitope within OspC$_A$, OspC$_B$, and OspC$_K$ (**Table 1**). The peptides were synthesized with a

C-terminal biotin tag and coupled to streptavidin-coated microspheres for multiplex

immunoassays (MIA) by Luminex. As a positive control, we also coupled microspheres with the

VlsE C6 B31-17 peptide from *B. burgdorferi* strain B31(43).

We employed two *B. burgdorferi* seropositive serum sample collections to gauge the

relative reactivity of recombinant OspC, OspC-derived peptides, and the VlsE C6-17 peptide. A

"diagnostic" collection consisted of 696 de-identified human serum samples from the Wadsworth

Center's Diagnostic Immunology laboratory classified as *B. burgdorferi* seropositive and

categorized based on Western blot banding profiles, as IgM$^+$/IgG$^-$ (+/-), IgM$^+$/IgG$^+$ (+/+), or IgM$^-$

/IgG$^+$ (-/+). A second collection provided by the Lyme Disease Biobank at Nuvance Health

consisted of 158 serum samples from PTLD patients. These two collections were compared to a

commercial panel consisting of ~75 serum samples confirmed negative for VlsE C6-17 peptide

reactivity. Reactivity was limited to IgG analysis, since preliminary IgM reactivity profiles were

low or inconclusive for the two sample sets (data not shown).

We found that the diagnostic immunology samples were significantly more reactive with

recombinant OspC than the control samples, although reactivity was skewed towards OspC$_A$ and

OspC$_B$ more than OspC$_K$ **(Figure 2A)**. We can only speculate that this bias reflects actual

exposure to *B. burgdorferi* expressing these two OspC types. In terms of peptides, the diagnostic

serum samples were on average significantly more reactive than control serums samples with all

the OspC-derived peptides tested, except for peptides 71-86$_A$ and 178-187$_B$ (**Figure 2A**). The

average serum IgG reactivity was highest against C10 (193-210$_A$) and VlsE C6-17 (p<0.0001)

with the remainder of the peptides being low to moderately reactive relative to the control.

To apply a more rigorous test of peptide reactivity, MFI values were subjected to a cutoff

value of ≥ 6 SD above the mean of the control MFI, $\log_{10}$ transformed, then plotted to reveal both

the number of serum samples above the cutoff (% positive) as well as the magnitude of samples

over the cutoff (fold increase). We also derived box and whisker plots to illustrate the upper and

lower quartiles as well as the median reactivity to each antigen/peptide with superimposed violin

plots to outline sample distribution. Using these metrics, reactivity with $OspC_A$, $OspC_B$ and

$OspC_K$ ranged from 24-42% of the diagnostic samples with the fold increase ranging from 2.8 to

4.7 (**Table 2**; **Figure 3A**). In terms of peptides, VlsE C6-17 was the most reactive (71% positive;

fold increase 5.8), followed by peptide C10 (26% positive; fold increase 4.4) (**Table 3; Figure**

3B). The percent of samples >6SD for the other 12 peptides ranged from 0.1 to 10% with the

fold increase between 1.2 and 2.5 (**Table 3; Figure 3B**). Thus, while overall MFI values

indicating that OspC peptides are widely recognized by *B. burgdorferi* immune sera, applying a

more stringent threshold to the analysis suggests a different story. Namely, that OspC-derived

peptides (apart from C10) are recognized by only a fraction of the total diagnostic serum

samples. While the profile suggests limited utility of peptides other than C10 for diagnostic

applications, it does raise questions about whether reactivity with particular linear epitopes on

OspC influence disease outcome, a point that we will return to in the Discussion.

The same overall serologic analysis was performed with a panel of PTLD samples. The

PTLD samples were significantly more reactive with $OspC_A$, $OspC_B$ and $OspC_K$ than the control

serum samples (**Figure 2B**). For $OspC_A$ and $OspC_K$, 16 and 18% of the samples were above the

≥ 6SD cutoff with the fold increase between 2.8-5 (**Table 2; Figure 3C**). In the case of $OspC_B$,

only 3.2% were above the cutoff with a fold increase of 2.1 (**Table 2; Figure 3C**). In terms of

linear epitopes, reactivity significantly above controls was limited to VlsE C6-17 (40% positive;

fold increase 5.6), C10 (14% positive; fold increase 3.3), and 178-187$_K$ (9.5%; positive; fold

increase 3.0) (**Figure 2B**, **Table 3, Figure 3D**). Comparatively, PTLD samples had significantly

less reactivity to peptide 9-33$_{ABK}$ relative to controls (**Figure 2B**). Only a small fraction of the

PTLD serum samples were found to be positive for the remaining 10 peptides (range 1.3-9.5%)

and the MFIs did not achieve statistical significance (**Table 3, Figure 2B**). PTLD samples were

plotted to visualize OspC and peptide reactivities relative to the cutoff (**Figure 3C-D**). There is

notably less reactivity to both C10 and C6-17 in the PTLD cohort in comparison to the

diagnostic population (**Figure 3B, 3D**).

Understanding the relationships between OspC and OspC-derived peptide epitopes has

implications for Lyme disease diagnostics and possibly immunity to *B. burgdorferi*. We expected

that antibody reactivity with the OspC-derived peptides would strongly correlate with overall

OspC antibody levels in any given sample, suggesting that the two populations track with each

other and serve as redundant indicators of a previous *B. burgdorferi* infection. To investigate

these relationships, we generated correlation matrices (Spearman's Rank) showing the

correlation between OspC and the each of the different OspC-derived peptides, including C10

and C6-17 (**Figure 4**). While all correlations were determined to be positive, in the diagnostic

samples, C10 reactivity was only weakly correlated with $OspC_A$, $OspC_B$, or $OspC_K$ [r_s = 0.2-0.4]

(**Figure 4A**). Rather, C10 reactivity was moderately to strongly correlated with other OspC-

derived peptides, including $132\text{-}146_A$ (r_s = 0.63), $183\text{-}190_{AB}$ (r_s = 0.58), and $40\text{-}55_{ABK}$ (r_s = 0.58).

Indeed, inter-peptide correlations were generally strong across the board (e.g., $132\text{-}146_B$ versus

$177\text{-}186_A$, r_s >0.9). A similar pattern was observed in the PTLD samples: OspC-derived peptide

reactivities were strongly correlated with each other (r_s = 0.63-0.98), but weakly correlated with

$OspC_A$, $OspC_B$, and $OspC_K$ (r_s = 0.2-0.4) (**Figure 4B**). In the case of VlsE C6-17, reactivity was

weakly correlated with OspC and OspC-derived peptides in both the diagnostic and PTLD

samples, except for moderate correlativity with C10 in the PTLD cohort (r_s = 0.56). These results

reveal a significant degree of discordance (asynchrony) among circulating antibody levels

against OspC, OspC-derived peptides and the VlsE C6-17 peptide in both diagnostic and PTLD

cohorts.

The absence of a strong correlation between antibody levels to OspC and OspC-derived

peptides led us to subject the diagnostic and PTLD datasets to hierarchical clustering (HC) as

means to identify possible relationships in immunoreactivity profiles. In the case of the

diagnostic samples, the horizontal (top) dendrogram revealed three distinct branches consisting

of (1) $OspC_A$, $OspC_B$, and $OspC_K$, (2) C6-17 and C10, and (3) the other OspC-derived peptides

(**Figure 5A**). In cluster 1 ($OspC_A$, $OspC_B$, $OspC_K$), high MFI values were distributed throughout

the column with a concentration towards the lower quartile. There was no obvious clustering on

the vertical dendrogram to indicate a relationship between OspC reactivity and seropositivity

categories (i.e., $IgM^{+/-}$, $IgG^{+/-}$).

In branch 2, high VlsE C6-17 reactivity was present throughout the column, albeit, with

three hot (orange and red shading) zones and one cold (deep blue shading) zone. In the C10

column, there was generally an even distribution of MFI intensities only occasionally punctuated

with hot spots that often mirror C10 reactivity. Samples that were "cold" for C6-17 and C10

tended to be within the IgM+/IgG- pool, possibly indicative of an early response to *B.*

burgdorferi infection.

There were several interesting patterns within branch 3, which consisted of OspC-derived

peptides: uniformly <u>low reactivity</u> with peptide 9-33$_{ABK}$, as noted in Figures 2 and 3; a

subpopulation of samples that were uniformly <u>nonreactive</u> with all the OspC peptides, as

evidenced by a blue band across the top of the figure, which in some instances spilled over to

C10 and VlsE C6-17 columns; and, finally, a subpopulation of samples with <u>moderate to high</u>

pan-OspC peptide reactivity, as evidenced by a horizontal band of orange/red shading situated in

the lower third of the figure. Collectively, the HC analysis reveals a remarkable degree of

discordance in serum antibody profiles associated with *B. burgdorferi* seropositivity within the

diagnostic samples.

The HC plot of the PTLD serum samples was clearly distinct from the diagnostic samples

(**Figure 5B**). On the horizontal dendrogram, for example, there were just two major branches not

three: one encompassing OspC$_{ABK}$ and C6-17, and the other containing the OspC-derived

peptides, including C10. On the vertical dendrogram, there were two branches: the top branch

corresponding to low peptide reactivity with or without concomitant C6-17 and OspC

reactivities; and the bottom branch corresponding to high peptide reactivity that were more

frequently than not associated with high C6-17 reactivity. Within the bottom branch, there was a

notable subbranch corresponding to 4 samples with strikingly high reactivity with C10, C6-17,

and most of the OspC-derived peptides (including 9-33$_{ABK}$). We would tentatively classify these

handful of samples as "polyreactive" (51).

Contribution of C10 to overall OspC antibody reactivity. There is some debate as to

what proportion C10-specific antibodies constitute relative to total OspC antibodies, with some

studies reporting that C10-specific antibodies make up an overwhelming proportion (52), while

others suggesting the opposite (53). As our prior analysis included truncated derivatives of OspC

that lacked C10, we expressed a second version of OspC in which residues 202-210

(VVAESPKKP) were included. We then revisited Luminex analysis in which we compared MFI

to OspC$_A$ versus OspC$_A$+C10. MFIs for OspC$_A$+C10 were significantly greater than OspC$_A$ in

both diagnostic and PTLD cohorts, demonstrating that antibody reactivity to the conserved C-

terminal residues constitute an important component of the overall antibody response (p <0.05;

Mann-Whitney U test) (**Figure 6A, Figure 6D**). We then investigated the relationship between

C10 reactivity and $OspC_A$ vs $OspC_A$+C10 using a linear regression model for diagnostic samples

(**Figure 6B, Figure 6C**) and PTLD samples (**Figure 6E, Figure 6F**). There is a slight increase

(0.0702) in the adjusted R-squared value between the C10 peptide MFI and $OspC_A$+C10, as

compared to $OspC_A$ for the DI cohort (**Figure 6B, Figure 6C**). A similar increase (0.0848) in the

adjusted R-squared value was observed between the C10 peptide and $OspC_A$+C10 in comparison

to $OspC_A$ for the PTLD population. However, the overall adjusted R-squared value was

negligible, thereby demonstrating that C10-specific antibodies constitute only a small fraction of

the total OspC reactivity.

DISCUSSION

Defining the breadth and the frequency of linear B cell epitope reactivity on OspC has

important implications for both Lyme disease diagnostics and vaccines. In this report, a panel of

peptides representing eight previously reported and predicted linear epitopes on OspC types A, B

and K, were examined by multiplex immunoassay for serum IgG reactivity in diagnostic (n~700)

and PTLD (n~150) *B. burgdorferi* seropositive sample sets. The results revealed that seven of the

eight peptides are indeed antibody targets even though reactivity to any given peptide may be

confined to a fraction (2-10%) of the total samples tested. The exception being peptide C10,

which was 26% positive (4.4 positive fold increase value) in the diagnostic sample set and 14%

positive (3.3 positive fold increase value) in the PLTDs sample set. Interestingly, in the

diagnostic sample set, C10 antibody titers did not correlate with either reactivity to OspC itself or

any of the other OspC-derived peptides. In the PTLD sample set, C10 antibody levels did

correlate with the other OspC-derived peptides but were distinct from OspC. These results have

clinical implications, as they demonstrate that the human B cell response to linear epitopes on

OspC is both variable and asynchronous.

The C10 peptide corresponds to the stretch of ~10 highly conserved amino acids at C-

terminus of OspC, first recognized as an immunodominant epitope in neuroborreliosis patients

(52). Since then, the C10 peptide and derivatives have been integrated into numerous diagnostic

assays, including the Zeus VlseE1/C10 ELISA (25, 27, 28). Dolange and colleagues recently

proposed the use of high affinity C10-specific monoclonal antibodies as tools to detect

circulating levels of *B. burgdorferi* antigens (54). While there is little debate as to whether C10

constitutes a human B cell epitope, the issue of whether antibodies to C10 contribute to

borreliacidal activity remains contentious. Lovrich and colleagues reported that OspC-specific

borreliacidal activity (in a handful of human serum samples) was almost entirely directed against

C10 (55, 56), whereas Izac and colleagues argued that, in rodents, canines and non-human

primates, C10-specific antibodies contribute little if any borreliacidal activity (57). Although our

results do not resolve this issue per se, they do raise questions about the underlying B cell

biology associated with antibodies to C10 and OspC. The fact that C10 and OspC antibody titers

did not correlate with each other in the two serum sample sets examined suggests differences in

the kinetics of antibody induction, circulating half-life, and/or plasma cell persistence. Others

have suggested, at least in the case of VlsE, there is a distinct progression of epitope reactivity

over the course of *B. burgdorferi* infection with readily accessible epitopes targeted first and

more obstructed, membrane proximal epitopes following (58). Resolving detailed questions

about the nature of the human B cell response to OspC is becoming more feasible as single cell

V_H and V_L repertoires are catalogued over the course of Lyme disease (11, 59).

B cell epitope prediction software like Discotope 3.0 identifies residues 133-147 as

having a high immunogenic propensity (60). That prediction was borne out in our study, as

evidenced by statistically significant reactivity of serum samples with the three peptides

corresponding to residues $132\text{-}146_{AB}$ and $133\text{-}147_K$. Others have reported similar findings in

mice and in a few human samples. Namely, Earnhart recognized residues 136-150 (which they

refer to as "loop 5") as a target of antibodies in *B. burgdorferi*-infected mice (36). In a follow-up

study, the same investigators identified two serum samples from early Lyme disease patients that

reacted with peptides spanning 136-144 (61). A subsequent report by Arnaboldi and colleagues

did not detect binding to three different 15-mer $OspC_K$-derived peptides spanning residues 131-

155, possibly because their sample size was limited (27). In our collection of almost 700

diagnostic samples, the frequency of peptide positivity was admittedly relatively low (3-7%).

Nonetheless, antibodies targeting this epitopic region may be consequential in terms of

promoting spirochete clearance (36). For example, the potent borreliacidal mouse monoclonal

antibody 16.22 recognizes $OspC_A$ residues 133-147 with the key contact points postulated to be

residues 139-141 ("KXK" motif) (62). Thus, a more detailed examination of B cell epitopes

within this region of OspC in human Lyme disease patients is warranted.

It has been postulated that serum antibody profiling at the antigen and possibly even

epitope level may be able to resolve different stages of Lyme disease (e.g., acute, acute resolved,

PTLD) (10, 58). While the antibody profiling in our study was limited to serum IgG reactivity to

OspC, OspC-derived peptides, and VlsE C6-17, the results revealed distinct patterns between the

diagnostic and PTLD sample sets when subjected to HC. Most notable was the classification of

the C10 peptide: in the diagnostic sample set C10 clustered with C6-17, whereas in the PTLD

samples it clustered with the other OspC-derived peptides. Thus, it is tempting to speculate that

the ratio of C10 to C6-17 serum IgG reactivity may have some utility in defining disease

progression. By the same token, as virtually all samples tested were positive for either C10

and/or C6-17, establishing a combined C10 + C6-17 threshold value might serve to alleviate

false positives in Lyme disease diagnostics based solely on serology (63, 64). That said we

recognize that any claims to this effect are premature, considering that the serum samples

employed in our study are not associated with any clinical information beyond seropositivity.

Moreover, we lack a "post-Lyme healthy" cohort that Chandra and colleagues so effectively used

in their study to compare with their PTLD cohort (10).

One interesting pattern that emerged from the PTLD cohort was a cluster of four

individuals who demonstrated pan-reactivity with the OspC-derived peptides and VlsE C6-17

peptide. In follow-up analysis, those same four samples in question also reacted with other *B.*

burgdorferi-derived peptides, as well as control non-*B. burgdorferi*-derived peptides, but not

necessarily with the corresponding recombinant proteins (G. Freeman-Gallant, unpublished

results). Thus, the serum antibodies from those particular individuals seemingly have a

propensity for peptide antigens, possibly due to a particular bias in germline chain usage and

corresponding paratopes conducive to accommodating peptides (65). Whether or not there is a

germline bias in antibodies associated with PTLD is worthy of investigation, especially in light

of finding related to COVID-19. Moreover, the availability of bulk plasmablast BCR sequences,

as well as single cell paired BCR sequences from LD patients opens the door to detailed antibody

reactivity profiles and links to disease manifestations (66). We are also exploring the possibility

that these particular samples may have profiles consistent with autoantibodies (67) or have

characteristics associated with polyreactivity that might promote ligation to spirochete surface

antigens (68, 69).

Acknowledgements

We thank Dr. William Lee and other members of the Diagnostic Immunology Laboratory at the
Wadsworth Center for providing residual diagnostic serum samples. We gratefully acknowledge
Dr. John Martignetti and Lisa Arrigo of the Nuvance Health Lyme disease Biobank for
generously providing PTLD serum samples. We thank Elizabeth Cavosie (Wadsworth Center)
for administrative assistance and Greta Van Slyke (Wadsworth Center) for establishing Luminex
protocols. This work was supported by the National Institute of Allergy and Infectious Diseases
(NIAID), National Institutes of Health, Department of Health and Human Services, Contract No.
75N93019C00040 (PI/PD Mantis). The content is solely the responsibility of the authors and
does not necessarily represent the official views of the National Institutes of Health.

Conflicts of Interest

The authors have no conflicts to declare.

References Cited

1. Kugeler KJ, Schwartz AM, Delorey MJ, Mead PS, Hinckley AF. 2021. Estimating the
Frequency of Lyme Disease Diagnoses, United States, 2010-2018. Emerg Infect Dis
27:616-619.
2. Bobe JR, Jutras BL, Horn EJ, Embers ME, Bailey A, Moritz RL, Zhang Y, Soloski MJ,
Ostfeld RS, Marconi RT, Aucott J, Ma'ayan A, Keesing F, Lewis K, Ben Mamoun C,
Rebman AW, McClune ME, Breitschwerdt EB, Reddy PJ, Maggi R, Yang F, Nemser B,
Ozcan A, Garner O, Di Carlo D, Ballard Z, Joung HA, Garcia-Romeu A, Griffiths RR,
Baumgarth N, Fallon BA. 2021. Recent Progress in Lyme Disease and Remaining
Challenges. Front Med (Lausanne) 8:666554.
3. Radolf JD, Strle K, Lemieux JE, Strle F. 2021. Lyme Disease in Humans. Curr Issues Mol
Biol 42:333-384.
4. Steere AC, Strle F, Wormser GP, Hu LT, Branda JA, Hovius JW, Li X, Mead PS. 2016.
Lyme borreliosis. Nat Rev Dis Primers 2:16090.
5. Lochhead RB, Strle K, Arvikar SL, Weis JJ, Steere AC. 2021. Lyme arthritis: linking
infection, inflammation and autoimmunity. Nat Rev Rheumatol 17:449-461.

6.   Klempner MS, Hu LT, Evans J, Schmid CH, Johnson GM, Trevino RP, Norton D, Levy
L, Wall D, McCall J, Kosinski M, Weinstein A. 2001. Two controlled trials of antibiotic
treatment in patients with persistent symptoms and a history of Lyme disease. N Engl J
Med 345:85-92.

7.   Marques A. 2008. Chronic Lyme disease: a review. Infect Dis Clin North Am 22:341-60,
vii-viii.

8.   Aucott JN, Crowder LA, Kortte KB. 2013. Development of a foundation for a case
definition of post-treatment Lyme disease syndrome. Int J Infect Dis 17:e443-9.

9.   Aucott JN, Yang T, Yoon I, Powell D, Geller SA, Rebman AW. 2022. Risk of post-
treatment Lyme disease in patients with ideally-treated early Lyme disease: A prospective
cohort study. Int J Infect Dis 116:230-237.

10.   Chandra A, Wormser GP, Marques AR, Latov N, Alaedini A. 2011. Anti-Borrelia
burgdorferi antibody profile in post-Lyme disease syndrome. Clin Vaccine Immunol
18:767-71.

11.   Blum LK, Adamska JZ, Martin DS, Rebman AW, Elliott SE, Cao RRL, Embers ME,
Aucott JN, Soloski MJ, Robinson WH. 2018. Robust B Cell Responses Predict Rapid
Resolution of Lyme Disease. Front Immunol 9:1634.

12.   Bockenstedt LK, Wooten RM, Baumgarth N. 2021. Immune Response to Borrelia: Lessons
from Lyme Disease Spirochetes. Curr Issues Mol Biol 42:145-190.

13.   Embers ME, Hasenkampf NR, Jacobs MB, Philipp MT. 2012. Dynamic longitudinal
antibody responses during Borrelia burgdorferi infection and antibiotic treatment of rhesus
macaques. Clin Vaccine Immunol 19:1218-26.

14.   Jiang R, Meng H, Raddassi K, Fleming I, Hoehn KB, Dardick KR, Belperron AA,
Montgomery RR, Shalek AK, Hafler DA, Kleinstein SH, Bockenstedt LK. 2021. Single-
cell immunophenotyping of the skin lesion erythema migrans identifies IgM memory B
cells. JCI Insight 6.

15.   Barbour AG, Jasinskas A, Kayala MA, Davies DH, Steere AC, Baldi P, Felgner PL. 2008.
A genome-wide proteome array reveals a limited set of immunogens in natural infections
of humans and white-footed mice with Borrelia burgdorferi. Infect Immun 76:3374-89.

16. Tilly K, Krum JG, Bestor A, Jewett MW, Grimm D, Bueschel D, Byram R, Dorward D,
Vanraden MJ, Stewart P, Rosa P. 2006. Borrelia burgdorferi OspC protein required
exclusively in a crucial early stage of mammalian infection. Infect Immun 74:3554-64.

17. Pal U, Yang X, Chen M, Bockenstedt LK, Anderson JF, Flavell RA, Norgard MV, Fikrig
E. 2004. OspC facilitates Borrelia burgdorferi invasion of Ixodes scapularis salivary
glands. J Clin Invest 113:220-30.

18. Tilly K, Bestor A, Jewett MW, Rosa P. 2007. Rapid clearance of Lyme disease spirochetes
lacking OspC from skin. Infect Immun 75:1517-9.

19. Earnhart CG, Leblanc DV, Alix KE, Desrosiers DC, Radolf JD, Marconi RT. 2010.
Identification of residues within ligand-binding domain 1 (LBD1) of the Borrelia
burgdorferi OspC protein required for function in the mammalian environment. Mol
Microbiol 76:393-408.

20. Caine JA, Coburn J. 2016. Multifunctional and Redundant Roles of Borrelia burgdorferi
Outer Surface Proteins in Tissue Adhesion, Colonization, and Complement Evasion. Front
Immunol 7:442.

21. Caine JA, Lin YP, Kessler JR, Sato H, Leong JM, Coburn J. 2017. Borrelia burgdorferi
outer surface protein C (OspC) binds complement component C4b and confers bloodstream
survival. Cell Microbiol 19.

22. Lin YP, Tan X, Caine JA, Castellanos M, Chaconas G, Coburn J, Leong JM. 2020. Strain-
specific joint invasion and colonization by Lyme disease spirochetes is promoted by outer
surface protein C. PLoS Pathog 16:e1008516.

23. Tan X, Castellanos M, Chaconas G. 2023. Choreography of Lyme Disease Spirochete
Adhesins To Promote Vascular Escape. Microbiol Spectr 11:e0125423.

24. Branda JA, Steere AC. 2021. Laboratory Diagnosis of Lyme Borreliosis. Clin Microbiol
Rev 34.

25. Hahm JB, Breneman JWt, Liu J, Rabkina S, Zheng W, Zhou S, Walker RP, Kaul R. 2020.
A Fully Automated Multiplex Assay for Diagnosis of Lyme Disease with High Specificity
and Improved Early Sensitivity. J Clin Microbiol 58.

26. Bacon RM, Biggerstaff BJ, Schriefer ME, Gilmore RD, Jr., Philipp MT, Steere AC,
Wormser GP, Marques AR, Johnson BJ. 2003. Serodiagnosis of Lyme disease by kinetic
enzyme-linked immunosorbent assay using recombinant VlsE1 or peptide antigens of

Borrelia burgdorferi compared with 2-tiered testing using whole-cell lysates. J Infect Dis
187:1187-99.

27.   Arnaboldi PM, Seedarnee R, Sambir M, Callister SM, Imparato JA, Dattwyler RJ. 2013.
Outer surface protein C peptide derived from Borrelia burgdorferi sensu stricto as a target
for serodiagnosis of early lyme disease. Clin Vaccine Immunol 20:474-81.

28.   Baarsma ME, Vrijlandt A, Ursinus J, Zaaijer HL, Jurriaans S, van Dam AP, Hovius JW.
2022. Diagnostic performance of the ZEUS Borrelia VlsE1/pepC10 assay in European LB
patients: a case-control study. Eur J Clin Microbiol Infect Dis 41:387-393.

29.   Coleman AS, Rossmann E, Yang X, Song H, Lamichhane CM, Iyer R, Schwartz I, Pal U.
2011. BBK07 immunodominant peptides as serodiagnostic markers of Lyme disease. Clin
Vaccine Immunol 18:406-13.

30.   Mathiesen MJ, Christiansen M, Hansen K, Holm A, Asbrink E, Theisen M. 1998. Peptide-
based OspC enzyme-linked immunosorbent assay for serodiagnosis of Lyme borreliosis. J
Clin Microbiol 36:3474-9.

31.   O'Bier NS, Hatke AL, Camire AC, Marconi RT. 2021. Human and Veterinary Vaccines
for Lyme Disease. Curr Issues Mol Biol 42:191-222.

32.   Wilske B, Preac-Mursic V, Jauris S, Hofmann A, Pradel I, Soutschek E, Schwab E, Will
G, Wanner G. 1993. Immunological and molecular polymorphisms of OspC, an
immunodominant major outer surface protein of Borrelia burgdorferi. Infect Immun
61:2182-91.

33.   Barbour AG, Travinsky B. 2010. Evolution and distribution of the ospC Gene, a
transferable serotype determinant of Borrelia burgdorferi. MBio 1.

34.   Cerar T, Strle F, Stupica D, Ruzic-Sabljic E, McHugh G, Steere AC, Strle K. 2016.
Differences in Genotype, Clinical Features, and Inflammatory Potential of Borrelia
burgdorferi sensu stricto Strains from Europe and the United States. Emerg Infect Dis
22:818-27.

35.   Seinost G, Dykhuizen DE, Dattwyler RJ, Golde WT, Dunn JJ, Wang IN, Wormser GP,
Schriefer ME, Luft BJ. 1999. Four clones of Borrelia burgdorferi sensu stricto cause
invasive infection in humans. Infect Immun 67:3518-24.

36.   Earnhart CG, Buckles EL, Dumler JS, Marconi RT. 2005. Demonstration of OspC type
diversity in invasive human lyme disease isolates and identification of previously

uncharacterized epitopes that define the specificity of the OspC murine antibody response. Infect Immun 73:7869-77.

37.   Hanincova K, Mukherjee P, Ogden NH, Margos G, Wormser GP, Reed KD, Meece JK,

Vandermause MF, Schwartz I. 2013. Multilocus sequence typing of Borrelia burgdorferi

suggests existence of lineages with differential pathogenic properties in humans. PLoS One

8:e73066.

38.   Fraser CM, Casjens S, Huang WM, Sutton GG, Clayton R, Lathigra R, White O, Ketchum

KA, Dodson R, Hickey EK, Gwinn M, Dougherty B, Tomb JF, Fleischmann RD,

Richardson D, Peterson J, Kerlavage AR, Quackenbush J, Salzberg S, Hanson M, van Vugt

R, Palmer N, Adams MD, Gocayne J, Weidman J, Utterback T, Watthey L, McDonald L,

Artiach P, Bowman C, Garland S, Fuji C, Cotton MD, Horst K, Roberts K, Hatch B, Smith

HO, Venter JC. 1997. Genomic sequence of a Lyme disease spirochaete, Borrelia

burgdorferi. Nature 390:580-6.

39.   Anderson JF, Flavell RA, Magnarelli LA, Barthold SW, Kantor FS, Wallich R, Persing

DH, Mathiesen D, Fikrig E. 1996. Novel Borrelia burgdorferi isolates from Ixodes

scapularis and Ixodes dentatus ticks feeding on humans. J Clin Microbiol 34:524-9.

40.   Steere AC, Grodzicki RL, Kornblatt AN, Craft JE, Barbour AG, Burgdorfer W, Schmid

GP, Johnson E, Malawista SE. 1983. The spirochetal etiology of Lyme disease. N Engl J

Med 308:733-40.

41.   Rudolph MJ, Davis SA, Haque HME, Weis DD, Vance DJ, Piazza CL, Ejemel M, Cavacini

L, Wang Y, Mbow ML, Gilmore RD, Mantis NJ. 2023. Structural Elucidation of a

Protective B Cell Epitope on Outer Surface Protein C (OspC) of the Lyme Disease

Spirochete, Borreliella burgdorferi. mBio 14:e0298122.

42.   Kringelum JV, Lundegaard C, Lund O, Nielsen M. 2012. Reliable B cell epitope

predictions: impacts of method development and improved benchmarking. PLoS Comput

Biol 8:e1002829.

43.   Gomes-Solecki MJ, Meirelles L, Glass J, Dattwyler RJ. 2007. Epitope length, genospecies

dependency, and serum panel effect in the IR6 enzyme-linked immunosorbent assay for

detection of antibodies to Borrelia burgdorferi. Clin Vaccine Immunol 14:875-9.

44.   R core Team. 2023. R: A language and environment for statistical computing. R

Foundation for Statistical Computing,.

45.   Wickham H, Bryan J. 2023. Read Excel Files, v1.4.3. CRAN. https://readxl.tidyverse.org,
550   https://github.com/tidyverse/readxl.

551   46.   Wickham H. 2019. Welcome to the Tidyverse. Journal of Open Source Software 4:1686.

552   47.   Wei T, Simko V, Levy M, Xie Y, Jin Y, Zemla J. 2017. Package 'corrplot'. Statistician
553   56:e24.

554   48.   Kolde R. 2018-05-19 2018. pheatmap: Pretty Heatmaps. https://cran.r-
555   project.org/web/packages/pheatmap/index.html. Accessed

556   49.   Lemieux JE, Huang W, Hill N, Cerar T, Freimark L, Hernandez S, Luban M, Maraspin V,
557   Bogovic P, Ogrinc K, Ruzic-Sabljic E, Lapierre P, Lasek-Nesselquist E, Singh N, Iyer R,
558   Liveris D, Reed KD, Leong JM, Branda JA, Steere AC, Wormser GP, Strle F, Sabeti PC,
559   Schwartz I, Strle K. 2023. Whole genome sequencing of human Borrelia burgdorferi
560   isolates reveals linked blocks of accessory genome elements located on plasmids and
561   associated with human dissemination. PLoS Pathog 19:e1011243.

562   50.   Wormser GP, Brisson D, Liveris D, Hanincova K, Sandigursky S, Nowakowski J,
563   Nadelman RB, Ludin S, Schwartz I. 2008. Borrelia burgdorferi genotype predicts the
564   capacity for hematogenous dissemination during early Lyme disease. J Infect Dis
565   198:1358-64.

566   51.   Dimitrov JD, Planchais C, Roumenina LT, Vassilev TL, Kaveri SV, Lacroix-Desmazes S.
567   2013. Antibody polyreactivity in health and disease: statu variabilis. J Immunol 191:993-
568   9.

569   52.   Mathiesen MJ, Holm A, Christiansen M, Blom J, Hansen K, Ostergaard S, Theisen M.
570   1998. The dominant epitope of Borrelia garinii outer surface protein C recognized by sera
571   from patients with neuroborreliosis has a surface-exposed conserved structural motif.
572   Infect Immun 66:4073-9.

573   53.   Earnhart CG, Rhodes DV, Smith AA, Yang X, Tegels B, Carlyon JA, Pal U, Marconi RT.
574   2014. Assessment of the potential contribution of the highly conserved C-terminal motif
575   (C10) of Borrelia burgdorferi outer surface protein C in transmission and infectivity.
576   Pathog Dis 70:176-84.

577   54.   Dolange V, Simon S, Morel N. 2021. Detection of Borrelia burgdorferi antigens in tissues
578   and plasma during early infection in a mouse model. Sci Rep 11:17368.

55. Jobe DA, Lovrich SD, Schell RF, Callister SM. 2003. C-terminal region of outer surface
protein C binds borreliacidal antibodies in sera from patients with Lyme disease. Clin
Diagn Lab Immunol 10:573-8.

56. Lovrich SD, Jobe DA, Schell RF, Callister SM. 2005. Borreliacidal OspC antibodies
specific for a highly conserved epitope are immunodominant in human lyme disease and
do not occur in mice or hamsters. Clin Diagn Lab Immunol 12:746-51.

57. Izac JR, Camire AC, Earnhart CG, Embers ME, Funk RA, Breitschwerdt EB, Marconi RT.
2019. Analysis of the antigenic determinants of the OspC protein of the Lyme disease
spirochetes: Evidence that the C10 motif is not immunodominant or required to elicit
bactericidal antibody responses. Vaccine 37:2401-2407.

58. Jacek E, Tang KS, Komorowski L, Ajamian M, Probst C, Stevenson B, Wormser GP,
Marques AR, Alaedini A. 2016. Epitope-Specific Evolution of Human B Cell Responses
to Borrelia burgdorferi VlsE Protein from Early to Late Stages of Lyme Disease. J Immunol
196:1036-43.

59. Kirpach J, Colone A, Burckert JP, Faison WJ, Dubois A, Sinner R, Reye AL, Muller CP.
2019. Detection of a Low Level and Heterogeneous B Cell Immune Response in Peripheral
Blood of Acute Borreliosis Patients With High Throughput Sequencing. Front Immunol
10:1105.

60. Hoie MH, Gade FS, Johansen JM, Wurtzen C, Winther O, Nielsen M, Marcatili P. 2024.
DiscoTope-3.0: improved B-cell epitope prediction using inverse folding latent
representations. Front Immunol 15:1322712.

61. Buckles EL, Earnhart CG, Marconi RT. 2006. Analysis of antibody response in humans to
the type A OspC loop 5 domain and assessment of the potential utility of the loop 5 epitope
in Lyme disease vaccine development. Clin Vaccine Immunol 13:1162-5.

62. Yang X, Li Y, Dunn JJ, Luft BJ. 2006. Characterization of a unique borreliacidal epitope
on the outer surface protein C of Borrelia burgdorferi. FEMS Immunol Med Microbiol
48:64-74.

63. Rebman AW, Crowder LA, Kirkpatrick A, Aucott JN. 2015. Characteristics of
seroconversion and implications for diagnosis of post-treatment Lyme disease syndrome:
acute and convalescent serology among a prospective cohort of early Lyme disease
patients. Clin Rheumatol 34:585-9.

64.   Reifert J, Kamath K, Bozekowski J, Lis E, Horn EJ, Granger D, Theel ES, Shon J, Sawyer
JR, Daugherty PS. 2021. Serum Epitope Repertoire Analysis Enables Early Detection of
Lyme Disease with Improved Sensitivity in an Expandable Multiplex Format. J Clin
Microbiol 59.

65.   Cobaugh CW, Almagro JC, Pogson M, Iverson B, Georgiou G. 2008. Synthetic antibody
libraries focused towards peptide ligands. J Mol Biol 378:622-33.

66.   Blum JS, Fiani ML, Stahl PD. 1991. Proteolytic cleavage of ricin A chain in endosomal
vesicles. Evidence for the action of endosomal proteases at both neutral and acidic pH. J
Biol Chem 266:22091-5.

67.   Trier NH, Houen G. 2023. Antibody Cross-Reactivity in Auto-Immune Diseases. Int J Mol
Sci 24.

68.   Boughter CT, Borowska MT, Guthmiller JJ, Bendelac A, Wilson PC, Roux B, Adams EJ.
2020. Biochemical patterns of antibody polyreactivity revealed through a bioinformatics-
based analysis of CDR loops. Elife 9.

69.   Mouquet H, Scheid JF, Zoller MJ, Krogsgaard M, Ott RG, Shukair S, Artyomov MN,
Pietzsch J, Connors M, Pereyra F, Walker BD, Ho DD, Wilson PC, Seaman MS, Eisen
HN, Chakraborty AK, Hope TJ, Ravetch JV, Wardemann H, Nussenzweig MC. 2010.
Polyreactivity increases the apparent affinity of anti-HIV antibodies by heteroligation.
Nature 467:591-5.

70.   Ikushima M, Yamada F, Kawahashi S, Okuyama Y, Matsui K. 1999. Antibody response
to OspC-I synthetic peptide derived from outer surface protein C of Borrelia burgdorferi in
sera from Japanese forestry workers. Epidemiol Infect 122:429-33.

71.   Pulzova L, Flachbartova Z, Bencurova E, Potocnakova L, Comor L, Schreterova E, Bhide
M. 2016. Identification of B-cell epitopes of Borrelia burgdorferi outer surface protein C
by screening a phage-displayed gene fragment library. Microbiol Immunol 60:669-677.

72.   Tokarz R, Mishra N, Tagliafierro T, Sameroff S, Caciula A, Chauhan L, Patel J, Sullivan
E, Gucwa A, Fallon B, Golightly M, Molins C, Schriefer M, Marques A, Briese T, Lipkin
WI. 2018. A multiplex serologic platform for diagnosis of tick-borne diseases. Sci Rep
8:3158.

73.   Yu Z, Carter JM, Sigal LH, Stein S. 1996. Multi-well ELISA based on independent peptide

640   antigens for antibody capture. Application to Lyme disease serodiagnosis. J Immunol

641   Methods 198:25-33.

642   74.   Baum E, Randall AZ, Zeller M, Barbour AG. 2013. Inferring epitopes of a polymorphic

643   antigen amidst broadly cross-reactive antibodies using protein microarrays: a study of

644   OspC proteins of Borrelia burgdorferi. PLoS One 8:e67445.

FIGURE LEGENDS

Figure 1. Relative location of linear epitopes on homdimeric OspC$_A$. (**A**) Surface
representation of homodimeric OspC$_A$ (residues 38-201; PDB 1GGQ] with one monomer
(OspC) colored gray and the other, denoted with an apostrophe (OspC'), colored in charcoal. The
linear epitopes examined in this study are coded and numbered according to **Table 1**. Epitope 1
is represented as a bracket (]) on OspC', as its corresponding residues were truncated in the
recombinant version of OspC$_A$ used for X-ray crystal structure analysis. Similarly, only residues
193-201 of epitope 8 (residues 193-210) are shown because the C-terminus of OspC$_A$ was
truncated at position 201. Epitopes 6 and 7 overlap with the overlap colored as epitope 6. (**B**)
The same image of homodimeric OspC$_A$ as Panel A except that amino acid differences between
OspC types A, B and K within the eight epitopes are colored red to illustrate degree of
polymorphisms.

Figure 2. Serum IgG reactivity of diagnostic and PTLD samples with OspC and OspC-derived peptides. (**A**) DI and (**B**) PTLD serum samples (diluted 1:100) were subject to MIA
using microspheres coated with recombinant dimeric OspC types A, B and K (top rows), OspC-peptides as described in Table 1, and C6-17 peptide. Panels are labeled by corresponding residue
numbers for each OspC type. MFI values were $\log_{10}$ transformed and compared to the control
sample set. Significance was determined by the Mann-Whitney U-test (*, p< 0.05).

**Figure 3. Indexed (>6SD) IgG reactivity of DI and PTLD serum samples with OspC and
OspC-derived peptides.** MFI values for OspC and OspC- and VlsE-derived peptides were
indexed using a cutoff of 6SD above mean control reactivity. The resulting values were then
$\log_{10}$ transformed with each point representing a single sample. OspC subtypes are listed along
the x-axis, residues are indicated along the top of the plot (**B**, **D**), and each antigen is represented
by a different color. In each of the graphs, the 6 SD cutoff corresponds to a value of 0 on the y-axis and is depicted with horizontal red line. All samples falling above y = 0 were found to be
positive for the given antigen and were used to calculate percent positivity and fold increase as
reported in Tables 2 and 3. Box and whisker plots illustrate the upper and lower quartiles as well

as the median reactivity to each antigen (thick center line). Sample points that fall above or

below the whiskers are considered outliers. Violin plots outline the distribution of data.

Figure 4. Correlation matrices of rOspC and OspC-derived peptide reactivity in diagnostic

and PTLD serum. Diagnostic (**A**) and PTLD (**B**) serum sample MFIs (indexed and $\log_{10}$

transformed) were subjected to Spearman's Rank correlation. Only significant correlation

coefficients (r_s) are displayed (confidence level = 0.95). The degree and direction of correlation is

indicated by the color scale. Interpretations of the strength of each correlation was conducted as

such: >0.7 indicates a strong correlation, 0.5 – 0.7 is moderate, 0.3 – 0.5 is weak, and 0.0 – 0.3 is

negligible.

Figure 5. Hierarchical clustering of rOspC and OspC-derived reactivity in diagnostic and

PTLD serum. Heatmaps were generated using diagnostic (**A**) and PTLD (**B**) serum sample MFIs

(indexed and $\log_{10}$ transformed). Rows represent individual samples and columns represent the

panel of antigens. The color scales of **A** and B are identical to one another (allowing for direct

comparison), with 0 representing the positivity cutoff (as seen in **Figure 3**). Annotation of the

diagnostic samples (**A**) with Western blot profile data (previously described in the Materials and

Methods) is designated by the legend. The columns were subjected to hierarchical clustering by

correlation and the rows were clustered by Euclidean distance to better visualize patterns in

reactivity profiles.

Figure 6: Correlation between OspC$_A$ reactivity and C10 region. Diagnostic (n=686) and

PTLD (n=158) samples were subjected to Luminex with OspC$_A$, OspC$_A$+C10, and 193-210$_A$ as

described in the Materials and Methods. OspC$_A$ MFIs were evaluated against OspC$_A$+C10 for

diagnostic **(A)** and PTLD **(D)** samples. Significance was determined by Mann-Whitney U test. *,

$p < 0.05$. To examine the contribution of the C10 region to OspC$_A$ reactivity, MFIs for peptide

193-210$_A$ were plotted (on the y-axis) versus MFIs from OspC$_A$ and OspC$_A$+C10 (on the x-axis)

for diagnostic samples **(B, C)** and PTLD samples **(E, F)**. Best fit lines and confidence intervals

(level = 0.95) were added to each plot using a linear regression model. Adjusted R-squared (r^2)

values were calculated in R.

 TABLES

Table 1. OspC and VlsE-derived peptides used in this study

#[a]	residues	OspC type sequence	IEDB ID [b]	References
1	9-33	ILMTLFLFISCNNSGKDGNTSANSA	181204 745120	(70) (27) (71) (72)
2	40-55	A/B–PNLTEISKKITDSNAV K–PNLTEISKKITESNAV	559957	(71)
3	71-86	A–EIAAKAIGKKIHQNNG	12383	(73)
4	132-146 132-146 133-147	A–ETFTNKLKEKHTDLG B–EEFSTKLKDNHAQLG K–EDFTKKLEGEHAQLG	63756 ,745097 14380, 18025	(62)
5	155-169	A–AKEAILKTNGTKTKG	558968 181187	(27) (71)
6	177-186 178-187 178-187	A–FESVEVLSKA B–SGSLESLSKA K–FKAVENLAKA	NA	(74)
7	183-190[c]	A/B–LSKAAKEM	560173	(71)
8	193-210	A–NSVKELTSPVVAESPKKP	745122	(25, 52, 71, 72)
C6	VlsE C6-17	MKKDDQIAAAIALRGMA	41843	(43)

[a], Numbers correspond to peptides highlighted in Figure 1, except C6. [b], Immune Epitope Database (IEDB.org) identifier. [c], Residues 184-191 in Type B.

Table 2: Serum reactivity greater than 6SD to OspC$_{A/B/K}$ in diagnostic and PTLD samples

| OspC | Diagnostic | | PTLD | |
	n = (%)[a]	Fold increase (SD)[b]	n = (%)[a]	Fold increase (SD)[b]
A	167 (24.0)	2.8 (1.9)	29 (18.4)	5.5 (5.6)
B	287 (41.2)	3.7 (3.4)	5 (3.2)	2.1 (0.6)
K	195 (28.0)	4.7 (4.3)	26 (16.5)	2.8 (2.2)

[a], The number of samples (n =) and percent reactivity ("%") to each antigen in the diagnostic (n=696) and PTLD (n=158) sample sets were calculated using a cutoff of >6SD above the mean of controls samples; [b], The fold increase (magnitude) of positive sample antibody reactivity over controls such than an index value of 1 is 6SD above the control mean, while 2.8 corresponds to ~16SD greater than the mean control.

Table 3: Serum reactivity greater than 6SD to OspC peptides in diagnostic and PTLD serum samples

Peptide	Diagnostic		PTLD	
	n = (%)[a]	Fold increase (SD)[b]	n = (%)[a]	Fold increase (SD)[b]
$9\text{-}33_{ABK}$	1 (0.1)	2.4 (N/A)	0 (0)	N/A (N/A)
$40\text{-}55_{AB}$	17 (2.4)	2.1 (1.4)	4 (2.5)	1.6 (0.5)
$40\text{-}55_{K}$	7 (1.0)	1.5 (0.5)	2 (1.3)	1.3 (0.2)
$71\text{-}86_{A}$	28 (4.0)	1.7 (0.8)	1 (0.6)	1.4 (NA)
$132\text{-}146_{A}$	18 (2.6)	2.4 (1.5)	5 (3.2)	1.7 (0.6)
$132\text{-}146_{B}$	48 (6.9)	2.0 (1.3)	11 (7.0)	3.0 (1.9)
$133\text{-}147_{K}$	36 (5.2)	2.4 (2.6)	8 (5.1)	2.5 (1.2)
$155\text{-}169_{A}$	30 (4.3)	2.5 (2.5)	5 (3.2)	2.5 (1.5)
$177\text{-}186_{A}$	38 (5.5)	1.6 (0.7)	13 (8.2)	2.6 (2.0)
$178\text{-}187_{K}$	70 (10.1)	1.5 (0.7)	15 (9.5)	3.0 (2.8)
$178\text{-}187_{B}$	7 (1.0)	1.2 (0.1)	12 (7.6)	2.7 (1.9)
$183\text{-}190_{AB}$[c]	53 (7.6)	1.9 (1.0)	14 (8.9)	3.0 (3.1)
$193\text{-}210_{A}$ (C10)	185 (26.6)	4.4 (4.5)	22 (14.0)	3.3 (3.0)
VlsE C6-17	496 (71.3)	5.8 (2.9)	64 (40.5)	5.6 (5.0)

[a], The number of samples (n =) and percent reactivity ("%") to each antigen in the diagnostic (n=696) and PTLD (n=158) sample sets were calculated using a cutoff of >6SD above the mean of controls samples; [b], The fold increase (magnitude) of positive sample antibody reactivity over controls such than an index value of 1 is 6SD above the control mean as described in Table 2. [c], Residues 184-191 in Type B.

A. Putative epitopes on OspC$_A$

B. Polymorphisms (red) on OspC$_{A, B, K}$

A.

B.

Figure 3

A. rOspC reactivity with DI (n=696)

B. OspC peptide reactivity with DI (n=696)

C. rOspC reactivity with PTLD (n=158)

D. OspC peptide reactivity with PTLD (n=158)

Figure 4

A.

B.