mTOR inhibition in Q175 Huntington's disease model mice facilitates neuronal autophagy and mutant huntingtin clearance

Philip Stavrides[1], Chris N. Goulbourne[1], James Peddy[1], Chunfeng Huo[1], Mala Rao[1, 2], Vinod Khetarpal[5], Deanna M. Marchionini[5], Ralph A. Nixon[1, 2, 3, 4, #], Dun-Sheng Yang[1, 2, @]

[1]Center for Dementia Research, Nathan S. Kline Institute, Orangeburg, NY 10962 USA; Departments of [2]Psychiatry and [3]Cell Biology, and [4]NYU Neuroscience Institute, New York University Grossman School of Medicine, New York, NY 10016 USA; [5]CHDI Management/CHDI Foundation, New York, NY 10001 USA.

[@] Correspondence: Dun-Sheng.Yang@NKI.RFMH.ORG

[#] **Lead contact:**

Ralph A. Nixon, MD, PhD

Center for Dementia Research

Nathan S. Kline Institute

140 Old Orangeburg Road

Orangeburg, NY 10962

Phone: (845) 398-5423

Fax: (845) 398-5422

Email: Ralph.Nixon@NKI.RFMH.ORG

ABSTRACT

Huntington's disease (HD) is caused by expansion of the polyglutamine stretch in huntingtin

protein (HTT) resulting in hallmark aggresomes/inclusion bodies (IBs) composed of mutant

huntingtin protein (mHTT) and its fragments. Stimulating autophagy to enhance mHTT clearance

is considered a potential therapeutic strategy for HD. Our recent evaluation of the autophagic-

lysosomal pathway (ALP) in human HD brain reveals upregulated lysosomal biogenesis and

relatively normal autophagy flux in early Vonsattel grade brains, but impaired autolysosome

clearance in late grade brains, suggesting that autophagy stimulation could have therapeutic

benefits as an earlier clinical intervention. Here, we tested this hypothesis by crossing the Q175

HD knock-in model with our autophagy reporter mouse TRGL (**T**hy-1-**R**FP-**G**FP-**L**C3) to

investigate *in vivo* neuronal ALP dynamics. In the Q175 and/or TRGL/Q175 mice, mHTT was

detected in autophagic vacuoles and also exhibited high level colocalization with autophagy

receptors p62/SQSTM1 and ubiquitin in the IBs. Compared to the robust lysosomal pathology in

late-stage human HD striatum, ALP alterations in Q175 models are also late-onset but milder that

included a lowered phospho-p70S6K level, lysosome depletion and autolysosome elevation

including more poorly acidified autolysosomes and larger-sized lipofuscin granules, reflecting

impaired autophagic flux. Administration of a mTOR inhibitor to 6-mo-old TRGL/Q175

normalized lysosome number, ameliorated aggresome pathology while reducing mHTT-, p62- and

ubiquitin-immunoreactivities, suggesting beneficial potential of autophagy modulation at early

stages of disease progression.

INTRODUCTION

Huntington's disease (HD) is an autosomal dominant disorder caused by a mutation in the gene encoding the huntingtin protein (HTT) resulting in expansion of the polyglutamine (polyQ) stretch on its amino-terminus (The-Huntington's-Disease-Collaborative-Research-Group 1993, Bates, Dorsey et al. 2015, Jiang, Handley et al. 2023, Franklin, Teive et al. 2024). HD pathogenesis advances in a spatiotemporal pattern, which is used to stage disease pathology severity as Grade 0-4 (HD0, HD1, HD2, etc.)(Vonsattel, Myers et al. 1985). GABA-containing medium spiny projection neurons in the striatum are most susceptible to cell death (Vonsattel 2008), and the cerebral cortex, particularly layer 5a, also shows cell loss (Sotrel, Paskevich et al. 1991, Pressl, Mätlik et al. 2024). Neuronal intranuclear inclusions (NIIs) and neuropil inclusions are present in HD brains and are positive for mutant huntingtin (mHTT) and ubiquitin (Ub) (DiFiglia, Sapp et al. 1997, Gutekunst, Li et al. 1999).

Recently, we completed a comprehensive evaluation of the autophagic-lysosomal pathway (ALP) in the HD brain at progressive disease stages, focusing on the most affected brain region, the striatum, in comparison to a less affected region, the neocortex (Berg, Veeranna et al. 2024 bioRxiv). Double fluorescence immunolabeling and immuno-electron microscopy (IEM) revealed colocalization of HTT/mHTT with the autophagy-related adaptor proteins, p62/SQSTM1 and ubiquitin, and cathepsin D (CTSD) within aggresome inclusions and autophagic compartments, documenting the involvement of ALP in HTT/mHTT turnover and the disease-related impairment of this process in late-stage disease. The temporal evolution of ALP alterations generally revealed minimal detectable impairment of upstream autophagy steps [e.g., autophagosome (AP) induction, formation, and fusion with lysosomes (LY)] and, in striatum, elevated levels of LAMP1 and LAMP2 markers suggesting modestly upregulated LY biogenesis. At late disease stages, mainly HD4, neuronal ALP dysfunction exhibited enlarged/clumped CTSD-immunoreactive autolysosomes (AL)/LY and ultrastructural evidence of autophagic vacuole (AV) fusion and transition to lipofuscin granule formation. These findings collectively suggest that a relatively competent autophagy machinery is maintained during the disease progression with a compensatory upregulation in lysosomal biogenesis, which together prevents against mHTT accumulation. This situation is failing at the late disease stages when AL clearance is impeded, substrates, including mHTT and its metabolites, accumulate in AL and aggresome inclusions increase.

A possible implication of the foregoing findings is that pharmacologic enhancement of

autophagy applied at a symptomatic but early stage of disease, when the ALP clearance machinery

is fully competent, may be therapeutic in clearing mHTT protein in affected neurons. By contrast,

in Alzheimer's Disease (AD) lysosomal clearance deficits develop at the earliest disease stages,

suggesting enhanced autophagy induction is counterproductive.

Modulation of autophagy as a therapeutic strategy for HD has been investigated in various

cell and animal models of HD (Sarkar and Rubinsztein 2008, Boland, Yu et al. 2018, Kim, Lalonde

et al. 2021, Jiang, Handley et al. 2023, Yang and Zhang 2023). Of particular relevance to our

present study are those involving autophagy induction with mTOR-dependent or -independent

autophagy-enhancing approaches in HD mouse models [e.g. Trehalose (Tanaka, Machida et al.

2004); Rapamycin analog Temsirolimus (CCI-779) (Ravikumar, Vacher et al. 2004); Rilmenidine

(Rose, Menzies et al. 2010); Rhes manipulations (Baiamonte, Lee et al. 2013, Lee, Tecedor et al.

2015)]. These studies have generally demonstrated ameliorative effects on outcome measures such

as mHTT lowering and behavioral/motor function assays. However, in many cases it is unclear

whether autophagy mechanisms are directly engaged in the brain or are critical to rescue.

Thus, our study aimed to comprehensively characterize the autophagy response with a range

of autophagy markers to interrogate the competence of the entire autophagy process in relationship

to mHTT in the well-characterized zQ175 Knock-In HD mouse model (Q175). Towards this goal,

we evaluated autophagy in neurons of Q175 after introducing by neuron-specific transgenesis the

dual fluorescence-tagged autophagy probe, **t**andem **f**luorescent mRFP-eGFP-**LC3** (tfLC3), an

autophagy adaptor protein associated with AP and metabolized via autophagy (Lee, Rao et al.

2019). We thereby generated a new Q175 cross, namely TRGL (**T**hy-1-**R**FP-**G**FP-**L**C3)/Q175.

tfLC3 expression, driven postnatally by the neuron-specific Thy1-promoter, allows for selective

monitoring of neuronal autophagy without the confounding influence of glial cells. Resolution and

sensitivity in reporting tfLC3 signal is high compared to conventional immunofluorescence

staining of LC3. The ability to ratiometrically report pH-dependent changes in fluorescence

wavelength (hue angle) enables neutral pH AP to be distinguished from acidic AL, that

progressively acidify intraluminally upon fusion with LY. We are further able to differentiate AL

subgroups differing in their extent of acidification after LY fusion. Distinguishing properly

acidified AL from those poorly acidified because of delayed or defective acidification is assisted

by immunolabeling AL with a marker, such as cathepsin D, which is then detected with a third

fluorophore. Together this triple fluorescence paradigm objectively quantified by computer-
assisted deconvolution of the proportions of each label within the analyzed neuron, a reflection of
their relative pH (and fusion with LY), which allows ALP organelle subtypes including LC3-
nagative LY to be identified and quantified for their numbers, sizes and spatial distributions in
intact brain sections. The collective data provides reports on the completion (or lack thereof) of
autophagy flux (ALP dynamics) and any blockage at particular steps in the ALP pathway,
including the normal acidification and further maturation of AL through their successful
elimination of fluorescence-tagged LC3 (Lee, Rao et al. 2019, Lie, Yang et al. 2021, Lee, Yang et
al. 2022, Lie, Yoo et al. 2022).

Crossing TRGL mice with a model of a neurodegenerative disease, the Q175 mouse model of
HD, has enabled us to assess disease-related autophagy alterations in the Q175 and TRGL/Q175
models and their response to a pharmacological inhibition of mTOR, (mTORi) INK-128 (hereafter
INK). Our data demonstrate target engagement and positive effects of the compound on rescuing
Q175 phenotypes including reversal on AL/LY subtypes as reported by the tfLC3 probe and
parallel reductions of mHTT-, p62- and Ub-immunoreactivity (IR), suggesting that the compound
targeted the ALP to degrade mHTT.

122
123

RESULTS

Identification of inclusions and HTT molecular species in Q175 mice

Immunohistochemistry (IHC) experiments with the antibody mEM48, which preferentially
recognizes aggregated mHTT (Gutekunst, Li et al. 1999), revealed age-dependent development of
mHTT-positive profiles in the striatum of Q175KI mice. Brain sections from 2.5-mo-old Q175
(not shown) only exhibited faint and diffuse nuclear mHTT staining without identifiable
aggresomes/inclusion bodies (IBs), while at 6 and 10 months of age, mHTT-positive IBs were
readily detected progressively with age (Fig. 1A2-A3). There were no similar mHTT
immunoreactive puncta in WT striatum (Fig. 1A1). These observations are consistent with
previous findings in Q175 and other HD mouse models (Li, Li et al. 1999, Menalled, Sison et al.
2003, Carty, Berson et al. 2015). To further determine the locations of mHTT IBs in Q175,
mEM48-immunostained sections were counterstained with cresyl violet (Fig. 1A4) to distinguish
nuclear IBs (Fig. 1A4, arrowheads) from extranuclear IBs, which were localized predominantly in

the neuropil (Fig. 1A4, arrows) and also detected rarely in the cytoplasmic portion of the

perikaryon. Thus, our results demonstrate age-dependent increase in mHTT aggresomes in the

Q175 model.

To reveal the ultrastructural locations and features of the mHTT aggregates, immunogold EM

(IEM) with antibody mEM48 was performed. EM images (Fig. 1B) demonstrated that the IEM

with this antibody was highly specific in detecting mHTT IBs which were localized in the neuronal

nuclei, dendrites and axons. Ultrastructurally, most IBs were cotton-ball shaped and composed of

fine fibrous or granular elements, somewhat similar to the unbundled short fibrils/protofibrils

found *in vitro* with recombinant mHTT protein fragments (Ko, Isas et al. 2018, Kolla, Gopinath et

al. 2021, Mario Isas, Pandey et al. 2021), and similar to the structure of NII type inclusions we

found in the human HD brain (Berg, Veeranna et al. 2024). Notably, however, human neuritic IBs

often displayed a more heterogeneous composition, such as fine fibrils mixed with AVs or bundles

composed of microtubule-like filaments, which were not found in the mouse neuritic IBs. Thus,

the result suggests a homogenous aggresome pathology in the mouse model, possible reflecting a

more rapid formation rate than in human.

Previous immunoblotting studies have observed fragmentation of mHTT molecules in the

human brain (Kim, Yi et al. 2001, Mende-Mueller, Toneff et al. 2001), including our own study

which detects mHTT fragments of 45-48 kDa, which predominantly exist in HD striatum (Berg,

Veeranna et al. 2024). To assess HTT molecular species in the Q175 mouse brain, we employed

multiple antibodies that preferentially detected mHTT over wild type HTT, including MAB1574

(Clone 1C2) (Fig. 1C1, C2), its epitope containing a 38-glns stretch (Trottier, Lutz et al. 1995),

and mAb PHP2 (Fig. 1C1), reacting with the peptide sequence QAQPLLPQP within the proline

rich domain of HTT (Ko, Isas et al. 2018). Both detected full-length mHTT and a ~120 kDa

fragment in the Q175 model (Figs. 1C1; 1C2 left and 1C3 top two graphs). By contrast, MAB5490

(Fig. 1C2 right), reacting with aa115-129 of HTT (C-terminal to the region of polyQ stretch-

containing exon 1), detected both wild type and mutant forms of HTT (Fig. 1C3, bottom right

graph). A ~48 kDa HTT fragment may correspond to the 45-48 kDa fragment seen in human brain

(Berg, Veeranna et al. 2024), and was detected by both MAB1574 and MAB5490 antibodies in

some samples (Fig. 1C2) but its levels in the Q175 models and the control TRGL were not

statistically significantly different (Fig. 1C3, bottom left graph). Thus, the result suggests that HTT

fragmentation is not obvious in the Q175 brains, unlike the far more prevalent occurrence of this
phenomenon in human HD striatum.

mHTT colocalizes with p62, Ub and CTSD in Q175 striatum

Our earlier study in human HD brains (Berg, Veeranna et al. 2024) found a high degree of
colocalization of mHTT/Ub or Ub/p62 colocalized signals in IBs, suggesting a relationship
between mHTT and the autophagy machinery since p62 and Ub are adaptor proteins mediating
autophagic cargo sequestration. Similarly, in Q175 mice, mHTT/Ub or Ub/p62 signals were highly
colocalized in IBs, particularly in NIIs, in striatal neurons of Q175 mice (Fig. S1). Additional triple
IF labeling experiments (mHTT/Ub/p62) with brain sections from 6- and 10-mo-old Q175 mice
identified IBs positive for mHTT, p62 and Ub within or outside the nuclei (Fig. 2A, arrows and
arrowheads, respectively). The 10-mo-old Q175 mice exhibited more and larger mHTT-positive
IBs than 6-mo mutants (Fig. 2A, 1st column), consistent with Fig. 1A and the literature (Carty,
Berson et al. 2015, Deng, Wang et al. 2021). Thus, our data demonstrate a close spatial relationship
among mHTT and autophagy receptor protein p62 and Ub in the Q175 model, which is consistent
with our observations in human HD brain (Berg, Veeranna et al. 2024).

Double IF labeling of mHTT with a LY marker cathepsin D (CTSD) (Fig. 2B1) detected small
punctate mHTT signal in CTSD-positive vesicles, suggesting a pool of mHTT within the ALP.
Consistent with this light microscopic (LM) finding, IEM with antibody MAB1574 directed
against mHTT clearly demonstrated that the AVs in either cell bodies, dendrites and axons were
labeled with concentrated silver-enhanced gold particles (Fig. 2B2), and the specificity of the IEM
labeling was very high, as reflected by a much higher number of silver-enhanced gold particles
associated with AVs versus the minimal number of silver-enhanced gold particles with
mitochondria that represent the background labeling (Fig. 2B3). Together, the LM and IEM
findings suggest that mHTT molecules exist within ALP vesicles in the absence of definable
aggresome/IB structures, similar to the IEM finding from a recent report (Zhou, Peskett et al.
2021).

Mild late-onset alterations in the ALP revealed in 17-mo-old Q175 mice

We crossed TRGL mice (Lee, Rao et al. 2019) with Q175 to generate TRGL/Q175 and
assessed AV/LY subtypes in striatal neurons with a hue angle-based analysis method (Lee, Rao et

al. 2019, Lee, Yang et al. 2022). Our pilot studies in young TRGL/Q175 mice (2.5 to 6-mo of age)

hardly detected autophagy alterations, compared to the control TRGL mice (not shown). Previous

studies have reported that autophagy impairment was hardly detected in 6-mo Q175/GFP-LC3

mice (Wold, Lim et al. 2016), but alterations in a limited number of autophagy markers (e.g.,

protein levels of p62, LC3, p-Beclin-1) were found in brain homogenates of 12-15-mo Q175 mice

(Wold, Lim et al. 2016, Abd-Elrahman, Hamilton et al. 2017, Heikkinen, Bragge et al. 2021).

Therefore, to further validate the status of the ALP and to discover potential additional autophagy

alterations, we expanded our study to old mice at 17-mo of age, where we found the following

ALP alterations.

(1) Alterations in AV/LY subtypes. Confocal images from 17-mo-old TRGL/Q175 and control

TRGL brain sections, triple-fluorescent labeled via an additional immunostaining with an anti-

CTSD antibody and far-red emitting fluorophore, showed enhanced mRFP- and eGFP-LC3 signals

in the striatal neurons of TRGL/Q175 compared to TRGL (Fig. 3A1). Hue angle-based analysis

confirmed the above observations by revealing significant increases in the numbers of AL and

poorly-acidified AL (pa-AL, as explained in Lee et al., 2019; 2022). It also detected a reduction in

the number of LY (Fig. 3A2). Thus, these results indicate that the tfLC3 reporter can reveal

alterations in the proportions of AV/LY subtypes in Q175, consistent with delayed and/or deficient

acidification of AL causing deficits in the reformation of LY to replenish the LY pool.

(2) Mild alterations in the levels of autophagy marker proteins detected by immunoblotting. To

further assess autophagy phenotypes in Q175 models, we conducted western blotting (WB) using

hemibrain homogenates from 17-mo-old mice for protein markers of individual steps in the ALP

(i.e., autophagy induction, membrane nucleation/AP formation, autophagy adaptor proteins and

AL formation/substrate degradation) (Fig. S2). At the stage of autophagy induction signaling, we

did not see alterations in the levels of mTOR and p-mTOR forms including its auto-phospho-form

at S2481, but we did observe a decreased level of a mTOR substrate, p-p70S6K (T389), in

TRGL/Q175 compared to the control TRGL (Fig. S2A), implying reduced mTOR activity, similar

to the finding from a previous study (Ravikumar, Vacher et al. 2004). Another mTOR substrate,

p-ULK1 (S757), did not exhibit changes in TRGL/Q175. However, there was an increase in the

level of total ULK1, resulting in a reduced ratio of p-ULK1 (S757)/total ULK1 in TRGL/Q175

compared to TRGL (Fig. S2A). The data collectively imply a down-regulated mTOR activity, as expected in response to accumulated aggregate-prone protein in Q175 mice, or as a result of sequestration of mTOR by mHTT IBs as suggested previously (Ravikumar, Vacher et al. 2004).

Except for these alterations, we did not detect statistically significant differences between TRGL/Q175 and the control TRGL in all other tested marker proteins in the downstream phases of the ALP (Fig. S2B-D). Of special interest is the ATG14-containing VPS34/Beclin-1 complex implicated in HD pathogenesis (Park, Jung et al. 2016, Wold, Lim et al. 2016) for which we did not detect statistically significant alterations in the levels of Beclin-1, VPS34 and ATG14 and their corresponding phosphor-forms. Notably, even if some TRGL/Q175 clearly exhibited diminished signals for p-ATG14 (S29) (Fig. S2B), no statistical significance could be established due to large variations among samples (See Discussion). Another notable point is that we did not find alterations in levels of full-length proteins or fragments (Norman, Cohen et al. 2010, Jamilloux, Lagrange et al. 2018, Sanchez-Garrido, Sancho-Shimizu et al. 2018, Valionyte, Yang et al. 2022) of autophagy adaptor protein such as p62, TRAF6 (Fig. S2C), in contrast to our immunoblot analysis of human HD brains (Berg, Veeranna et al. 2024). Together, our data suggest that, in general, autophagy alterations at the protein level, as can be detected by immunoblotting, are mild in TRGL/Q175 even at 17-mo-old (See Discussion).

(3) Increased numbers of larger lipofuscin granules. To assess possible ultrastructural pathologies, we conducted EM study for Q175 vs WT at 17-mo of age. As predicted, NIIs were not observed in WT neurons (Fig. 3B, left panel) but were detected (Fig. 3B, right panel, red circle) in ~10% striatal neurons of Q175 mice. Different from the substantial accumulation of AVs including lipofuscin granules in late stage human HD brain (e.g., HD4) (Berg, Veeranna et al. 2024), we did not observe gross accumulation of AVs in Q175 at this old age, except that quantitative analysis of EM images revealed that larger sized (>1 μm) lipofuscin granules (Fig. 3B, right panel, arrow) modestly increased in number in Q175 compared to WT [WT: 113 lipofuscin/420 neurons (n = 6 mice), i.e., 27 lipofuscin/100 neurons; Q175: 228 lipofuscin/586 neurons (n = 9 mice), i.e., 39 lipofuscin/100 neurons; Unpaired t-test, Two-tailed P value < 0.01; Bar graph not shown]. Thus, these EM findings suggest that ultrastructural autophagy alterations in the ALP were mild in Q175 even at an older age.

DARPP-32-IR decreases in striatal neurons of 17-mo-old TRGL/Q175 in the absence neuronal loss

To investigate potential neurodegeneration in our model, brain sections from 17-mo-old mice were double-labeled with the medium spiny neuron marker DARPP-32 (dopamine- and cAMP-regulated phosphoprotein, 32 kDa)(Ouimet, Langley-Gullion et al. 1998) and a general neuronal marker NeuN. We found that the intensity of DARPP-32-IR significantly diminished in TRGL/Q175 compared to TRGL (Fig. 4A, B). However, there were no alterations in NeuN-IR, including Area Covered and the Number of NeuN-positive neurons (Fig. 4A, B), and there were clear examples of neurons showing strong NeuN signal but faint or no DARPP-32 signal (Fig. 4A, arrows). Such a finding is consistent with other studies in Q175 and Q140 mice where neuronal loss was not found even though a reduction in DARPP-32 was seen in the same study, and that neuronal loss only occurred quite late, e.g., around 2 years of age (Hickey, Kosmalska et al. 2008, Rothe, Deliano et al. 2015, Peng, Wu et al. 2016, Deng, Wang et al. 2021). Thus, our data support the notion that changes in DARPP-32 may indicate alterations in its protein level and a loss of phenotype pattern, which is a potentially common prelude to neuronal loss in other neuronal cell types, e.g., basal forebrain cholinergic neurons (Jiang, Alam et al. 2022).

Oral mTORi INK engages mTOR target and induces downstream responses in the ALP in 7-mo-old TRGL/Q175

As a prelude to investigate effects of pharmacological autophagy stimulation with a mTORi, INK, on TRGL/Q175 mice, we first performed a safety, pharmacokinetic and pharmacodynamic evaluation of INK in 6-mo-old WT mice, administered via oral gavage. The WB results (Fig. S3A) indicated that all doses tested, from 2.5 mg/kg (mpk) to 10-mpk, exhibited target engagement as revealed by the reductions in the protein levels of mTOR targets, i.e., p-ULK1 (S757, phosphorylated by mTOR), p-p70S6K (not shown) or phosphorylated S6 ribosomal protein (p-S6, at S240/244), implying high BBB permeability of this compound and target engagement. Consistently, measurements for the levels of INK in cerebellar homogenates of WT mice treated with 1, 3 and 10-mpk INK revealed dose-dependent brain levels of INK (Fig. S3B).

We then decided an oral gavage treatment regimen for INK as 4-mpk, daily, for 3 weeks, to be administered in 6-mo-old TRGL/Q175. The brains from the mice (7-mo of age after the 3-week treatment duration) were then analyzed by multiple experimental approaches. By immunoblotting,

target engagement was verified, as indicated by decreased levels of p-mTOR (S2481,
autophosphorylation site), p-ULK1 (S757) and p-S6 (S240) in brain homogenates from INK-
treated TRGL/Q175 compared to Veh-treated TRGL/Q175 (Fig. 5A, B). Additionally, there were
also INK induced changes in marker proteins located downstream of autophagy induction,
including, for example, increased levels of p-ATG14 in INK-treated samples (Fig. 5A, B). It is
interesting that although INK did not induce any alterations in the transgene product tfLC3-I or -
II, there was a trend of decreased endogenous LC3-I, leading to a statistically significant increase
in the ratio of LC3-II/I, and such a result in LC3-I reduction was reproduced in a repeated
experiment (Fig. 5A, B). Thus, the pilot study in WT mice and the actual study in TRGL/Q175
mice together established mTORi INK's BBB permeability, target engagement and ability to
modify molecular events in the ALP.

mTOR inhibition alters AV/LY subtypes

We took advantage of the tfLC3 construct reporting *in vivo* autophagy flux (Lee, Rao et al.
2019) to assess INK effects on AV/LY subtypes in TRGL/Q175. Confocal images from INK- or
Veh-treated TRGL/Q175 and Veh-treated TRGL brain sections of 7-mo-old mice, immuno-
stained with anti-CTSD antibody and detected with a third fluorophore, were collected (Fig. 6A)
and processed for hue-angle based quantitative analysis using the protocol described previously
(Lee, Rao et al. 2019). The analysis in striatal neurons (Fig. 6B) revealed reversal of the two main
ALP alterations. First, the pre-existing abnormally lowered LY number in TRGL/Q175-Veh (Fig.
6B, LY group, red vs green), was reversed by INK treatment (Fig. 6B, LY group, red vs blue).
Second, INK reduced the number of AL in TRGL/Q175-Veh compared to TRGL-Veh (Fig. 6B,
AL group, red vs blue), implying improved AL clearance and likely providing the basis for
restoration of normal LY number. Together, the data suggest that autophagy phenotype in 7-mo
Q175 is measurable but milder than that in the 17-mo-old mice shown in Fig. 3A and that INK has
the potential to modulate the autophagy pathway and restore normal function at this mild disease
stage.

mTOR inhibition mitigates mHTT-related aggresome pathology

320   Importantly, we found that HD-like pathology such as mHTT IBs and associated protein
321   aggregation of adaptor proteins, p62 and Ub, was ameliorated by INK treatment. Confocal images

demonstrated that mHTT positive particles/inclusions were substantially reduced in INK-treated 7-mo-old TRGL/Q175, compared to Veh-treated TRGL/Q175 mice (Fig. 7A). Moreover, p62-IR or Ub-IR also changed in a similar trend (Fig. 7A), consistent with the high degree of colocalization of mHTT/p62/Ub signals shown in Fig. 2A. The corresponding quantitative analysis, shown as Area Covered by either mHTT-, p62- or Ub-IR on a per cell basis, confirmed statistically significant decreases in the Total Area Covered by mHTT-, p62-signals (Fig. 7B1). With the assistance of the endogenous tfLC3 signal (particularly the mRFP signal) in the TRGL to identify the association status of the mHTT-, p62- or Ub-IR with AVs, the images were additionally quantified to generate the separated "Area Covered by the AV-associated Form" (Fig. 7B2) and "Area Covered by the AV-unassociated Form" (Fig. 7B3) of mHTT-, p62- or Ub- signals. The results collectively suggest decreased mHTT-, p62- or Ub-IR in treated compared to untreated TRGL/Q175, implying that INK treatment was able to promote the clearance of mHTT and related receptor proteins p62 and/or Ub.

mTOR inhibition does not reverse preexisting DARPP-32-IR reduction in TRGL/Q175

Finally, we evaluated neurodegenerative changes by employing DARPP-32 IHC on brain sections of Veh-treated or INK-treated TRGL/Q175. Similar to what was found in 17-mo-old mice (Fig. 4), the group of TRGL/Q175-Veh, even at this young age (7-mo), already exhibited a diminished DARPP-32-IR signal compared to that in TRGL-Veh (Fig. 8A, B). However, no differences between the untreated and INK-treated TRGL/Q175 groups were established by quantitative analysis (Fig. 8A, B). Thus, the result suggests a lack of effectiveness of INK in reversing this pre-existing phonotype, implying that earlier intervention may be necessary (Marchionini, Liu et al. 2022).

DISCUSSION

Previous studies in cell and animal models have documented roles of HTT and mHTT in autophagy in relation to HD (Martin, Ladha et al. 2015, for reviews, see Croce and Yamamoto 2019, Klionsky, Petroni et al. 2021). WT HTT participates in normal autophagy by releasing ULK1 via mTOR inhibition and serving as a scaffold to facilitate cargo sequestration by enhancing p62 interaction with LC3 and ubiquitinated cargos (Ochaba, Lukacsovich et al. 2014, Rui, Xu et

al. 2015). mHTT in HD may activate autophagy by sequestering and thus inhibiting mTOR

(Ravikumar, Vacher et al. 2004). Most mHTT effects in cellular and mouse HD models appear to

be inhibitory for autophagy induction steps, from initiation signaling and phagophore nucleation

to cargo recognition/AP formation (Martinez-Vicente, Talloczy et al. 2010, Pryor, Biagioli et al.

2014, Rui, Xu et al. 2015, Wold, Lim et al. 2016, Ashkenazi, Bento et al. 2017). In addition, mHTT

may disturb endolysosomal homeostasis by reducing exocytosis and promoting AL accumulation

(Zhou, Peskett et al. 2021). In the current study using Q175, we have found that mHTT is an ALP

substrate and that the ALP is relatively intact, with some late-onset minor alterations, such as

insufficient mTOR activity reflected by lowered phospho-p70S6K level and ratio of phospho-

ULK1 (S757)/total ULK1, lysosome depletion and AL/pa-AL elevation. This mild late-onset

nature of ALP pathogenesis allowed us to modulate ALP with a mTOR inhibitor in 6-mo-old

TRGL/Q175 which resulted in positive effects on rescuing Q175 phenotypes including

normalization of AL/LY subtypes and reduction in aggresome pathology.

367   ALP impairments in Q175 are mild and late-onset

In the current study, we assessed ALP alterations in the Q175 model by analyzing protein

markers in all phases of the ALP including autophagy induction signaling, membrane

nucleation/AP formation, autophagy adaptor proteins and AL formation/substrate degradation. By

immunoblot analysis, alterations in the tested marker proteins even at 17 months of age in

TRGL/Q175 mice were limited to a reduction in p-P70S6K (T389), similar to that reported

previously (Ravikumar, Vacher et al. 2004), and a reduced ratio of p-ULK1 (S757)/total ULK1

as a result of increased total ULK1. While these two alterations together would suggest a lower

mTOR activity and autophagy activation, consequential alterations of downstream markers,

particularly proteins in the ATG14-containing Beclin-1/VPS34 complex, were not detected.

Although 4 out of the 10 tested TRGL/Q175 mice exhibited very low levels of p-ATG14, implying

a deficit in this complex as previously reported (Wold, Lim et al. 2016), variation among animals

was too great to draw statistical conclusions. In this regard, mice used in the current study were

heterozygous, which results in less autophagy alteration than that reported in homozygous mice

using certain autophagy markers (Abd-Elrahman, Hamilton et al. 2017). It may also be noted in

our experience that immunoblot analysis of brain tissue averages contributions from varied cell

types that may respond to disease states in opposite directions and mask changes in a given cell

population. Thus, we also applied a neuron-specific analysis using our neuron-specific tfLC3 reporter and hue-angle based image quantification approach, which yielded more sensitive detection and findings.

In brain sections from 17-mo-old TRGL/Q175 mice, we observed increased numbers of AL including the population of pa-AL and reduced numbers of LY. Such changes in AL/pa-AL were not readily observed in TRGL/Q175 mice at 2 months (not shown) or 7 months (e.g., Fig. 6B, which only exhibited a reduction in LY number) of age, indicating that even if these alterations signify an existence of impairments in the late phase of the ALP, they do not occur until a later age (e.g., 17 mo). Importantly, this pattern of late-onset ALP pathology progression, e.g., AL/pa-AL increases, is consistent with our findings from HD human brains (Berg, Veeranna et al. 2024) where enlarged CTSD-positive AL accumulate and cluster in affected neuronal populations at the late disease stages (HD3 and HD4). These findings suggest that the overall autophagy function in heterozygous Q175 is largely maintained until the relatively late disease stage and provide a rationale to stimulate autophagy as a therapeutic intervention when the autophagy machinery is still functional, which would generate beneficial effects. We obtained support for this concept in the subsequent study with mTORi INK when administered to TRGL/Q175 at 6 months of age for three weeks.

Autophagy as a primary mechanism for INK-enhanced clearance of small mHTT species

One of the key findings from this study is the reduction of mHTT-IR in neurons detected by immunolabeling after the 3-week oral treatment with mTORi INK (Fig. 7). This beneficial outcome from the drug treatment is interpreted as a primary result of the manipulation/stimulation of autophagy by the compound, leading to enhanced autophagy clearance of mHTT species including smaller, presumably soluble, species and larger aggregates such as the IBs. This interpretation is supported by the following considerations and evidence. First, in theory, the compound used for the treatment is a mTORi so is expected to enhance autophagy (Kim and Guan 2015, Bensalem, Fourrier et al. 2021). Second, target engagement is achieved as clearly reflected by the diminished levels of mTOR-mediated phosphorylation on mTOR itself (S2481), ULK1 (S757) and p-S6 (S240/S244), along with changes in proteins downstream of mTOR, such as p-ATG14 (S29) and LC3. Third, hue-angle based confocal image analysis of AL/LY subtypes reveals a reduced AL number and an increased LY number in INK-treated TRGL/Q175 compared

to Veh-treated TRGL/Q175. The findings from this analysis of AL/LY subtypes highlights the

advantages of crossing TRGL with Q175 mice to evaluate autophagy modulators on the ALP,

consistent with a notion of developing better tools capable of investigating autophagy flux *in vivo*

(Nixon 2013, Vodicka, Lim et al. 2014).

Our confocal microscopy analyses revealed mHTT signal in CTSD-positive vesicles, which

can be classified as AL based on their LC3 fluorescence, and IEM detected mHTT-silver-enhanced

gold particles in AP and AL including lipofuscin granules. Both findings indicate the existence of

mHTT species inside the ALP as autophagy substrates. The presence of low levels of mHTT in

CTSD-positive AL also implies a constitutive functional degradation event of mHTT in AL (Fig.

2B1, 10-mo-old mouse). This is in contrast to the case in young AD mouse models where early-

onset autophagy-stress exhibited severe substrate accumulation (e.g., reflected by strong LC3

signal) due to deficits of degradative functions within AL. (Lee, Yang et al. 2022). A reduced level

in the portion of AV-associated mHTT signals (Fig. 7B2) by mTORi INK further suggests that

mHTT degradation has been enhanced after manipulating the ALP. This AV-associated mHTT

pool is considered to be corresponding to the mHTT-silver-enhanced gold signal under IEM (Fig.

2B2), i.e., amorphous and existing within the lumen of AV. Such mHTT localization and

ultrastructural features are consistent with recent findings that mHTT proteins exogenously added

to Q175 brain sections for incubation are recruited to multivesicular bodies/amphisomes, AL and

residual bodies (lipofuscin granules), and ultrastructurally localized to non-fibrillar, electron-

lucent regions within the lumen of these organelles (Zhou, Peskett et al. 2021).

Possible autophagy-related mechanism for the clearance of large mHTT IBs

It is notable that the AV-non-associated form of mHTT (i.e., did not show colocalization with

the endogenous tfLC3 signal implying that they might not have a contact with AVs) decrease more

obviously after drug-treatment, along with the reductions in p62 and Ub signals (Fig. 7B3). These

mHTT species are mainly larger aggresomes/IBs, including those within the nuclei. Based on our

mHTT-IEM observation (Fig. 1B), the IBs in Q175 brain are fibrillar aggregates in the nucleus

and cytoplasm, exhibiting a homogeneous feature in shape (i.e., round/oval cotton-ball) and in

content (i.e., short fine fibrils), similar to the nuclear IBs we found in human HD brains (Berg,

Veeranna et al. 2024), whose cytoplasmic and neuritic IBs, however, also exhibit more

heterogeneous compositions such as a mixture of fine fibrils with AVs, fingerprint-like structures

or bundles of microtubules. In the current study, we did not observe that a whole IB is contained within an AP or AL in our EM examination, raising a question of how these larger aggregates could be degraded by the autophagy machinery after an autophagy activation by mTOR inhibition. This issue is informed by findings that the mHTT species are dynamic and the aggregates can form as different phases (liquid-like or solid-like), making it possible for mHTT species to move out or into IBs (Liu, Shyu et al. 2015, Peskett, Rau et al. 2018, Aktar, Burudpakdee et al. 2019, Riguet, Mahul-Mellier et al. 2021) and to shuttle between the nucleus and the cytoplasm (Xia, Lee et al. 2003, Atwal, Xia et al. 2007). Thus, it is speculated that under the autophagy activated condition induced by mTOR inhibition, the enhanced clearance of mHTT in the ALP pool would promote movements of mHTT species from the IB forms, leading to a reduction on the overall mHTT load.

UPS as an additional mechanism for the clearance of mHTT species

Even if INK is a mTORi mainly targeting autophagy, we cannot completely exclude additional contributions from the UPS for the observed reduction of mHTT species given the crosstalk between the two proteolytic systems. It is well accepted that autophagy plays a more important role in the clearance of larger protein aggregates such as aggresomes/IBs (Sarkar and Rubinsztein 2008). For example, autophagy is involved in the degradation of p62 bodies (Bjørkøy, Lamark et al. 2005) and autophagy markers are colocalized with cytoplasmic mHTT IBs (Iwata, Christianson et al. 2005). However, there is also evidence for degradation of polyQ aggregates in the nuclei (Iwata, Nagashima et al. 2009), which contain p62 condensates as a hub for UPS-mediated nuclear protein turnover (Fu, Cohen-Kaplan et al. 2021). We observed a high level of colocalization of mHTT with p62 and Ub in aggregates at both intranuclear and extranuclear locations (Fig. 2A), which is presumably more related to the UPS hub and autophagy, respectively. It is possible that stimulation of autophagy is reducing the mHTT in the cytoplasm and thereby partially relieves the burden of the proteasome both in the cytoplasm and in the nucleus so that the nuclear proteasome operates more effectively.

In summary, Q175 models, as detected in the current study, develop ALP alterations late in the disease progression. Although milder than what is seen in late-stage HD human brain (Berg, Veeranna et al. 2024), there are similarities in ALP pathobiology between human HD brains and Q175 brains. These include late-onset nature, mHTT as an ALP substrate, increased AL/pa-AL,

mouse IB ultrastructural feature similar to the common fine fibril type of human IBs. The late-onset mild ALP pathology in Q175 is different from the severe AV accumulation in mouse models of AD as mentioned above (Lee, Yang et al. 2022). Together, the mild and late-onset nature of ALP alterations in Q175 provides a basis for manipulating autophagy to be a promising therapeutic strategy if the manipulations (e.g., stimulating autophagy with mTORi as in this study) are applied when the autophagy machinery is largely undamaged. This has been validated to be successful in this study as demonstrated by the INK-induced reduction of mHTT species along with other beneficial effects, consistent with findings from similar studies and supporting the general notion of lowering mHTT as a therapeutic strategy for HD (Cortes and La Spada 2014, Barker, Fujimaki et al. 2020).

ACKNOWLEDGMENTS

This work was supported by the CHDI Foundation (R.A.N.) and the National Institute of Aging

(P01 AG017617 to R.A.N.). We are grateful to Dr. T Yoshimori (Osaka University, Japan) for the

mRFP-eEGFP-LC3 construct used in the TRGL mice.

CONFLICT OF INTEREST STATEMENT

All authors declare no conflict of interests.

 FIGURE LEGENDS

Fig. 1. Identification of inclusions and HTT molecular species in Q175 mice.

(A) IHC detects age-dependent increase in the number of mHTT inclusions. Brain sections from

6-mo- and 10-mo-old WT and Q175 mice were processed for IHC with antibody mEM48

(MAB5374) directed against mHTT (A). Images from sections without (A1-3) or with (A4) a

cresyl violet counterstain are shown where the dark-brown puncta represent mHTT-positive

inclusions. Arrowheads depict neuronal intranuclear inclusions (NIIs), determined with the

assistance from the nuclear labeling by cresyl violet, while arrows indicate extranuclear inclusions,

primarily the neuritic inclusion in the neuropil. Bars = 20 μm.

(B) mHTT inclusions are detected in nucleus, dendrites and axons of Q175 brains by IEM.

Sagittal vibratome brain sections of 17-mo-old Q175 were cut and went through EM processing.

Small blocks were obtained from the striatal areas for ultrathin sectioning. Tissue containing grids

were processed for immunogold labeling procedure with antibody mEM48, using 10 nm gold

followed by silver enhancement. Structures showing high level of silver-enhanced gold labeling

were considered as mHTT-positive.

(C) Various forms of HTT molecules are detected with different antibodies by immunoblotting.

Equal amounts of proteins from hemibrain homogenates of 17-mo-old WT, TRGL, Q175 and

TRGL/Q175 (labeled as "Cross") were subjected to SDS-PAGE and processed for WB with

different antibodies directed against HTT/mHTT, including MAB1574 (C1, C2), mAb PHP2 (C1)

and MAB5490 (C2). Images were collected by a digital gel imager (Syngene G:Box XX9). The

arrowhead and arrow (C2) depict a 120 kDa and a 48 kDa fragment, respectively. (C3)

Densitometry was performed with Image J for the blots shown in (C2) and the results were

normalized by the immunoblot(s) of given loading control protein(s) (e.g., GAPDH). Values are

the Mean +/- SEM for each group (n = 7 TRGL, 4 Q175 and 10 TRGL/Q175). Significant

differences among the groups were analyzed by One-Way ANOVA followed by Sidak's multiple

comparisons test. * P < 0.05, ** P < 0.01.

Fig. 2. Colocalization of mHTT with p62, Ub and CTSD.

(A) Triple labeling detects colocalization of mHTT with both autophagy adaptor proteins p62

and Ub. Brain sections from 6-mo- and 10-mo-old mice were immunostained with antibodies to

HTT (antibody MW8; red), p62 (green) or pan-Ub (blue), followed by an additional DAPI (cyan)

labeling, and confocal images from the striatum are shown. Boxed areas are enlarged and shown
in the last column. Arrows and arrowheads depict the white areas representing triply labeled
inclusions containing the signals of the three proteins, where arrows are for NIIs, determined with
the assistance from the nuclear DAPI labeling, while the arrowheads are for extranuclear
inclusions. Bar = 10 μm.

(B) mHTT is detected in vesicles of the ALP. (B1) Brain sections from 10-mo-old Q175 were
double-immunolabeled with antibodies to HTT (antibody MW8; green) and CTSD (red), followed
by an additional DAPI (blue) labeling, and a 3-color-merged confocal image from the striatum is
shown. Large and small green arrows depict HTT nuclear and extranuclear inclusions,
respectively, while yellow arrowheads depict yellow puncta showing HTT and CTSD signal
colocalization. Bar = 10 μm. (B2) IEM with anti-HTT antibody (MAB1574) specifically detects
HTT signal, represented by the silver-enhanced gold particles (red arrowheads), in AV/LY in cell
bodies, dendrites and axons. (B3) To demonstrate the labeling specificity of the HTT antibody in
this IEM study, the number of silver-enhanced gold particles in AV/LY existing in neuronal cell
bodies and neurites was counted from 69 EM images from two 10-mo-old Q175 mice against the
number of silver-enhanced gold particles in mitochondria on the same images, and the result is
shown in the bar graph. Statistical significances between the two groups were analyzed by
unpaired, two-tailed t-test. **** $P < 0.0001$.

Fig. 3. Mild late-onset alterations in the ALP in the striatum of 17-mo-old Q175.
(A) Quantitation of AV/LY subtypes of striatal neurons detect increases in AL, pa-AL and a
decrease in LY in 17-mo-old TRGL/Q175 vs. TRGL. (A1) Brain sections from TRGL and
TRGL/Q175 (4 sections/mouse, 10 mice/genotype) were immunostained with an anti-CTSD
antibody. Confocal images from the cranial-dorsal portion of the striatum (3 images at
120x/section) were collected and representative images for each eGFP-LC3 (green), mRFP-LC3
(red) and CTSD (blue) are shown. Arrowheads depict pa-AL. (A2) Hue angle-based analysis was
performed for AV/LY subtype determination using the methods described in Lee et al., 2019. Data
are presented as Vesicle #/Neuron (TRGL: n = 713 neurons; TRGL/Q175: n = 601 neurons).
Statistical significances between the two groups for each vesicle type were analyzed by unpaired
t-test. Two-tailed P value: *** $P<0.001$, **** $P<0.0001$.

(B) EM detects larger AL/lipofuscin granules in the Q175 striatum. Sagittal vibratome brain
sections from 17-mo-old mice were cut and went through EM processing. Small blocks were
obtained from the striatal area for ultrathin sectioning, followed by EM examinations of the grids.
The red circle depicts NII, and the arrow depicts larger sized (> 1µm) lipofuscin granules, which
were counted on randomly collected images of neurons from striatum of WT (420 neurons from n
= 6 mice) or Q175 (586 neurons from n = 9 mice).

Fig. 4. Decrease of DARPP-32-IR in striatal neurons of 17-mo-old TRGL/Q175 in the
absence of a NeuN signal reduction.
Brain sections from TRGL and TRGL/Q175 (n = 10 mice/genotype, 4 sections/mouse) were
double-immunostained with anti-DARPP-32 and -NeuN antibodies. Confocal images from the
cranial-dorsal portion of the striatum (3 images at 120x/section) were collected and representative
images are shown (A). Arrows depict examples of neurons showing strong NeuN signal while
minimal DARPP-32-IR. (B) Images were quantified by Image J for Integrated Density of DARPP-
32- (top) or NeuN-IR/Image field (middle), and for # of NeuN positive cells/Image field (bottom).
Statistical significances between the two groups were analyzed by unpaired, two-tailed t-test. * P
< 0.05.

Fig. 5. mTORi INK exhibits target engagement and induces downstream responses in the
ALP in 7-mo-old TRGL/Q175.
Equal amounts of proteins from brain homogenates of 7-mo-old TRGL/Q175 mice untreated
(labeled as "Veh") or mTORi INK treated [4 mg/kg (4-mpk), daily, 3 weeks; labeled as "INK"]
were subjected to SDS-PAGE and processed for WB with antibodies directed against several
marker proteins in the autophagy pathway, representing target engagement of INK or downstream
responses. Representative blots are shown on the left (A) while quantitative results of the blots are
shown on the right (B). The bottom LC3 blots represent a repeated immunoblotting experiment.
Values are the Mean +/- SEM for each group (n = 6-7 mice per condition). Significant differences
between the two groups were analyzed by unpaired, two-tailed t-test. * P<0.05, ** P<0.01, ***
P<0.001, **** P<0.0001. tfLC3 = mRFP-eGFP-LC3; Endo-LC3 = endogenous LC3.

Fig. 6. mTORi INK reverses the mild alteration of AV/LY subtypes in the striatum of 7-mo-old TRGL/Q175.

(A) Brain sections from untreated or INK (4-mpk, 3w)-treated TRGL/Q175 vs. TRGL (4 sections/mouse, 5-6 mice per condition) were immunostained with an anti-CTSD antibody. Confocal images from the cranial-dorsal portion of the striatum (3 images at 120x/section) were collected and representative images for each eGFP-LC3 (green), mRFP-LC3 (red) and CTSD (blue) are shown. (B) Hue angle-based analysis was performed for AV/LY subtype determination using the methods described in Lee et al., 2019. Data are presented as Vesicle #/Neuron (TRGL-Veh: n = 260 neurons; TRGL/Q175-Veh: n = 218 neurons; TRGL/Q175-INK: n = 287 neurons). Statistical significances among the groups were analyzed by One-Way ANOVA followed by Sidak's multiple comparisons test. ** $P<0.01$, **** $P<0.0001$.

Fig. 7. mTORi INK reduces HTT-, Ub- or p62-IR-covered areas parallelly in the striatum of 7-mo-old TRGL/Q175.

(A) Brain sections from INK (4-mpk, 3w)-treated or untreated TRGL/Q175 (4 sections/mouse, 6-7 mice/condition,) were immunostained with anti-HTT (MW8), -p62 or -pan-Ub antibodies. Confocal images from the cranial-dorsal portion of the striatum (3 images at 120x/section) were collected. Shown are single-channel images (i.e., without showing the eGFP and mRFP signals). (B1-B3) Areas covered by either the HTT-, p62- or Ub-IR on a per cell basis are quantified (for HTT-IR, TRGL/Q175-Veh: n = 196 neurons, TRGL/Q175-INK: n = 385 neurons; for p62-IR, TRGL/Q175-Veh: n = 311 neurons, TRGL/Q175-INK: n = 378 neurons; for Ub-IR, TRGL/Q175-Veh: n = 244 neurons, TRGL/Q175-INK: n = 347 neurons) and grouped as "Total Area" (B1), "AV-associated Form" (i.e., the IR which was associated with tfLC3 signals representing AVs)(B2) or "AV-unassociated Form" (i.e., the IR which was not associated with the tfLC3 signals)(B3). Statistical significances between the two groups were analyzed by unpaired t-test. Two-tailed P value: * $P < 0.05$, **** $P < 0.0001$.

Fig. 8. INK treatment does not reverse the reduction of DARPP-32-IR in the striatum of 7-mo-old TRGL/Q175.

(A) Sagittal brain sections containing the striatum area were immunostained with an anti-DARPP-32 antibody, and 10x images taken from the cranial-dorsal portion of the striatum are shown (1[st]

row). Boxed areas on the 1st row are enlarged and shown on the 2nd row for easy viewing of the immunostaining patterns for each condition. (B) For quantitation purpose, each whole 10x image (excluding the areas covered by the fiber bundles which usually exhibited minimal background staining – achieved by threshold setting) was quantified by Image J (1 image/section, 4 sections/mouse) and the results are expressed as the Integrated Density of DARPP-32-IR. n = 6-8 mice/condition. Statistical significances among the groups were analyzed by One-Way ANOVA followed by Sidak's multiple comparisons test. ** $P < 0.01$ compared to TRGL-Veh.

 SUPPLEMENTAL FIGURE LEGENDS

Fig. S1 (related to Fig. 2): Colocalization of HTT/Ub and Ub/p62 in IBs in the STR.

Brain sections from 10-mo-old mice were processed for double IF with antibodies against HTT

(antibody MW8; red) and pan-Ub (green), or pan-Ub (red) and p62 (green), and confocal images

from the striatum are shown. Arrowheads depict IBs showing colocalization signals.

Fig. S2 (related to Fig. 3): Molecules involved in various stages of the ALP are largely

unchanged in 17-mo-old TRGL/Q175.

Equal amounts of proteins from brain homogenates of 17-mo-old TRGL and TRGL/Q175 were

subjected to SDS-PAGE and processed for WB with antibodies directed against a number of

interested marker proteins in the ALP, grouped according to the functions of the proteins in the

autophagy process: autophagy induction signaling (A), membrane nucleation/AP formation (B),

autophagy adaptor proteins (C) and AL formation/substrate degradation (e.g., lysosomal

hydrolases or structural components)(D). The blot of the loading control protein GAPDH is boxed

and placed under each protein of interest. Images were collected by a digital gel imager (Syngene

G:Box XX9). Densitometry was performed with Image J and the result for each protein of interest

was normalized by its corresponding GAPDH blot and presented in the bar graph. Values are the

Mean +/- SEM for each group (n = 7 TRGL and 10 TRGL/Q175). Significant differences between

the 2 groups were analyzed by unpaired, two-tailed t-test. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$.

Fig. S3 (related to Fig. 5): Dosing tests demonstrate mTORi INK BBB penetration and target

engagement even at low dosages.

(A) Dosing tests for INK. INK or the vehicle was administered by oral gavage. Two dosing tests

were done in 6-mo-old WT mice, one with 10, 8 mg/kg (mpk) every other day or 4-mpk daily for

2 weeks and the other with 7.5, 5, 2.5-mpk daily for 2 weeks. Equal amounts of proteins from brain

homogenates of mice receiving vehicle (labeled as "Veh") or INK were subjected to SDS-PAGE

and processed for WB with antibodies directed against marker proteins representing target

engagement of INK, such as p-ULK1 (S757) and p-S6 – the drug effects are reflected by the

changes within the boxed areas compared to the signals of Veh outside the boxed areas.

(B) INK was detected in the brains of mice receiving a single dose via oral gavage. After 6, 12

or 24 hrs post INK administration to 6-mo-old WT mice, INK was detected in cerebellum of mice

treated with 1, 3 or 10-mpk INK. Total 60 mice: n = 6 mice/dose/time point, where the Vehicle

control group was just tested once at the 6-hr time point.

MATERIALS AND METHODS

The HD Q175 mouse model and the generation of a new Q175 model crossed with the autophagy reporter mouse TRGL (TRGL/Q175): Q175 mice (B6J.zQ175 Knock-In mice, CHDI-81003003)(Menalled, Sison et al. 2003, Menalled, Kudwa et al. 2012) were obtained from the Jackson Lab (Stock No. 027410) on a C57BL/6J background. The zQ175 KI allele has the mouse *Htt* exon 1 replaced by the human *HTT* exon 1 sequence containing a ~190 CAG repeat tract, and the average size for the mice used in this study was 200 CAG. Q175 genotyping and CAG sizing were conducted by Laragen, Inc. (Culver City, CA). The TRGL (**T**hy-1 m**R**FP-e**G**FP-**L**C3) mouse model, expressing tandem fluorescence-tagged LC3 (tfLC3 or mRFP-eGFP-LC3) in neurons, was generated, genotyped by PCR and maintained in a C57BL/6J background as described previously (Lee, Rao et al. 2019). The TRGL was crossed with Q175 to generate the new TRGL/Q175 model featuring the tfLC3 probe as an *in vivo* autophagy reporter.

The mice were maintained in the Nathan Kline Institute for Psychiatric Research (NKI) animal facility and housed in a 12 hrs light/dark cycle. All animal procedures were performed following the National Institutes of Health Guidelines for the Humane Treatment of Animals, with approval from the Institutional Animal Care and Use Committee at the NKI. Animals of both sexes were used in this study. Details for mouse ages are in the Figure Legend and/or Results. All efforts were made to minimize animal suffering and number of animals used.

mTORi INK administration to mice: INK-128 (ChemScene) was formulated in the sterile filtered vehicle (Veh) – 0.5% carboxymethyl cellulose (Sigma, Cat #5678) and 0.05% tween 80 in water – at different concentrations to achieve the various desired dosages. The mixture was homogenized with a tissue homogenizer, stored in a sterile container at room temperature for up to 2 weeks, and stirred prior to each dosing. It was administered to mice via oral gavage at a dose volume of 5 ml/kg, which was a relatively smaller volume of drug solution (generally 100-150 ul depending on the mouse body weight) aimed at avoiding the suppression of appetite. Mice receiving the Veh (also at a solution volume of 5 ml/kg) served as the control group. Oral gavage was conducted with 20G-38mm Cadence Science™ Malleable Stainless-Steel Animal Feeding Needles #9921 (Fisher Sci Cat #.14-825-275).

mTORi INK pharmacokinetics in mouse brains after oral administration: INK was formulated into 0.5% carboxymethyl cellulose and 0.05% tween-80 in water to be administered orally. 6 months old C57B6/J male mice were treated with compound at 1, 3 or 10 mg/kg (mpk) or vehicle. 6, 12 or 24 hours after administration, mice were terminally anesthetized with pentobarbital and cerebellum was dissected and fresh frozen on liquid nitrogen and stored at -80ºC until analysis. Tissue was homogenized and extracted with acetonitrile (containing 0.1% formic acid). Extract was analyzed using a LC-MS/MS (Waters Xevo TQ MS) method with a lower limit of quantitation of 13 nM.

Brain tissue preparation*:* To obtain tissues for experiments, the animals were anesthetized with a mixture of ketamine (100 mg/kg BW) and xylazine (10 mg/kg BW). Mice for light microscopic (LM) analyses were usually fixed by cardiac perfusion using 4% paraformaldehyde (PFA) in 0.1 M sodium cacodylate buffer [pH 7.4, Electron Microscopy Sciences (EMS), Hatfield, PA]. Following perfusion fixation, the brains were immersion-fixed in the same fixative overnight at 4°C. For transmission electron microscopic (EM) study, 4% PFA was supplemented with 2% glutaraldehyde (EMS). For biochemical analyses such as immunoblotting, the brains were flash frozen on dry ice and stored at –70°C. When both morphological and biochemical analyses were to be performed on the same brain, the brain was removed after brief perfusion with saline. One hemisphere was frozen at –70°C and the other half was immersion-fixed in 4% PFA for 3 days at 4°C.

Antibodies for immunohistochemistry (IHC) and western blotting (WB): The following primary antibodies were used in this study. (1) from Cell Signaling Technology: tHTT rabbit mAb (total HTTs, #5656), MTOR rabbit mAb (#2983); p-MTOR (S2481 autophosphorylation) pAb (#2974), p-MTOR (S2448) pAb (#2971), p70S6K rabbit mAb (#2708), p-p70S6K (S371) pAb (#9208), p-p70S6K (T389) rabbit mAb (#9235), S6 ribosomal protein mAb (#2317), p-S6 ribosomal protein (S240/244) rabbit mAb (#5364), ULK1 pAb (#4773), ULK1 rabbit mAb (#6439), p-ULK1 (S757) pAb (#6888), p-ULK1 (S317) pAb (#37762), ATG5 rabbit mAb (#12994), ATG14 pAb (#5504), p-ATG14 (S29) rabbit mAb (#92340), p-Beclin 1 (S30) rabbit mAb (#35955), VPS34 rabbit mAb (#4263), TRAF6 rabbit mAb (#8028). (2) from Millipore-Sigma: ntHTT mAb (mEM48, #MAB5374; N-Terminus-specific), HTT mAb (1C2, #MAB1574; epitope: N-terminal part of the

human TATA Box Binding Protein (TBP) containing a 38-glns stretch), HTT mAb (#MAB5490; epitope: human HTT aa115-129), ATG5 pAb (#ABC14), LC3 pAb (#ABC929), β–actin mAb (#A1978). (3) from other vendors: HTT mAb MW8 (epitope: HD exon 1 with 67Q; Develop Studies Hybridoma Bank, University of Iowa), HTT mAb PHP2 (CHDI-90001516-2, Coriell/CHDI), Beclin 1 mAb (BD Biosciences, #612113), LC3 pAb (Novus Biologics, #NB100-2220), p62/SQSTM1 mAb (BD Biosciences, #610832) or C-term-specific p62/SQSTM1 Guinea Pig pAb (Progen Biotechnik #C-1620); ubiquitin pAb (Dako Agilent, #Z0458), ubiquitin pAb (Abcam, #ab7780), LAMP1 rat mAb (Developmental Studies Hybridoma Bank, #H4A3), CTSB goat pAb (Neuromics, #GT15047). (4) In-house made: CTSD pAb (RU4). CTSD sheep pAb (D-2-3)(Cataldo, Thayer et al. 1990).

The following secondary antibodies and reagents for immunoperoxidase labeling were purchased from Vector Laboratories (Burlingame, CA): biotinylated goat anti-rabbit or -mouse IgG/IgM, Vectastain ABC kit (PK-4000), and DAB Peroxidase Substrate Kit (SK-4100). Mouse on Mouse (M.O.M) detection kit (BMK-2201), normal-goat (S-100) and normal-donkey (S-2000-20) serum blocking solution were also from Vector Lab. The following secondary antibodies for immunofluorescence were purchased from Thermo Fisher Scientific (Waltham, MA): Alexa Fluor 488-conjugated goat anti-rabbit IgG (A11034), Alexa Fluor 568- goat anti-rabbit IgG (A11036), Alexa Fluor 647- goat anti-rabbit (A21245), Alexa Fluor Plus 405- goat anti-rabbit (A48254), Alexa Fluor 568- goat anti-mouse IgG (A11031), Alexa Fluor 647- goat anti-mouse (A21235) and Alexa Fluor 647- goat anti-rat (A21247). HRP- linked secondary antibodies for immunoblotting were obtained from Jackson ImmunoResearch Laboratories (West Grove, PA): Rabbit IgG (711-035-152), Mouse IgG (711-035-150), Rat IgG (712-035-150), and Goat IgG (705-035-003).

Immunolabeling of brain sections: Immunoperoxidase and immunofluorescence IHC were performed according to the protocols previously described (Yang, Lee et al. 2009, Lee, Rao et al. 2019). Brain vibratome sections (40 μm) were blocked and incubated in primary antibody O/N (up to 3 days in some cases) at 4°C. ABC detection method was used for immunoperoxidase labeling visualized with DAB while Alexa-Fluor conjugated secondary antibodies were used for immunofluorescence. Autofluorescence was quenched with 1% Sudan black (Sigma-Aldrich; St. Louis, MO) in 70% ethanol for 20 minutes. DAB labeled sections was inspected on a Zeiss

AxioSkop II equipped with a HrM digital camera (Carl Zeiss, Germany). Immunofluoescently

labeled sections were examined on a Zeiss LSM880 confocal microscope.

Confocal image collection and hue-angle based quantitative analysis for AV/LY subtypes:

Confocal imaging was performed using a plan-Apochromat 20x or 40x/1.4 oil objective lens on a

LSM880 laser scanning confocal microscope with the following parameters: eGFP (ex: 488, em:

490-560 with MBS 488), mRFP (ex: 561, em: 582-640 with MBS 458/561), Alexafluor 647 (ex:

633, em: 640-710 with MBS 488/561/633) and DAPI (ex: 405, em: 410-483), with the 'best signal

scanning mode' which separates scanning tracks for each excitation and emission set to exclude

crosstalk between each fluorophore signal. The resolution of 40x images was 1024 x 1024 pixels

(corresponding to an area of 212.34 x 212.34 μm^2), and the resolution of 3x digitally zoomed

images was also 1024 x 1024 pixels (corresponding to an area of 70.78 x 70.78 μm^2). Detailed

settings for image collection were reported previously (Lee, Rao et al. 2019).

Hue angle-based vesicle analysis enables quantitative determination of AV/LY subtypes

including AP, AL, poorly acidified AL (pa-AL) and pure LY and the method was described in

detail previously (Lee, Rao et al. 2019). Briefly, high resolution confocal images containing the

LC3 (red and green) and CTSD (blue) punctate signals were analyzed with the Zen Blue Image

Analysis Module from Carl Zeiss Microscopy. Threshold for each of the three-color channels (red,

green, blue; RGB) was set by taking the average of intensity value from 20 neuronal perikarya.

The signal was segmented into discrete puncta by using the automatic watershed function to

separate clumped vesicles into individual puncta. Background signal was eliminated using the size

exclusion function of Zen Blue. The R, G and B intensity values of each vesicle were calculated

using the profile function of Zen Blue. The RGB ratio of each vesicle was converted into a hue

angle and saturation range – which we assigned to each AV subtype and should match the desired

AV color range as perceived visually (i.e., yellow for AP, blue for Ly, purple for AL and white for

de-acidified AL) – by entering the values of R, G, and B for a given punctum into the formula, as

follows: Hue° = IF(180/PI()*ATAN2(2*R-G-B,SQRT(3)*(G-B)) < 0,180/PI()*ATAN2(2*R-G-

B,SQRT(3) *(G-B)) + 360,180/PI()*ATAN2(2*R-G-B,SQRT(3)*(G-B))). Saturation percent of

the hue angle was calculated by entering the values of R, G, and B for a given punctum into the

following   formula   =   (MAX(RGB)-MIN(RGB))/SUM(MAX(RGB)+MIN(RGB))*100

(www.workwithcolor.com), provided lightness is less than 1, which is the usual case for our data.

Hue angle was converted to color using the Hue color wheel. The data was pooled and categorized
in Excel spreadsheets.

Ultrastructural analyses: Vibratome brain sections (50 μm) were treated with 1% osmium
tetroxide in 100 mM sodium cacodylate buffer pH 7.4 for 30 minutes, washed in distilled water
four times (10 min/wash), and then treated with 2% aqueous uranyl acetate overnight at 4°C in the
dark. Samples were then washed and sequentially dehydrated with increasing concentrations of
ethanol (20, 30, 50, 70, 90, and 100%) for 30 min each, followed by three additional treatments
with 100% ethanol for 20 min each. Samples were then infiltrated with increasing concentrations
of Spurr's resin (25% for 1 h, 50% for 1 h, 75% for 1 h, 100% for 1 h, 100% overnight at room
temperature), and then incubated overnight at 70°C in a resin mold. For TEM ultrastructural
analysis, 70 nm sections were cut using a Leica Reichert Ultracut S ultramicrotome and a Diatome
diamond knife, placed on to grids and then post stained with 2% uranyl acetate and lead citrate.
Images were taken using a Ceta Camera on a ThermoFisher Talos L120C transmission electron
microscope operating at 120kV.

For post-embedding IEM, ultrathin sections were placed on to carbon formvar 75 mesh nickel
grids and etched using 4% sodium metaperidotate for 10 minutes before being washed twice in
distilled water and then blocked for one hour. Grids were incubated in the primary antibodies at 4
°C overnight. Next day grids underwent seven washes in 1xPBS and were then incubated in anti-
mouse or anti rabbit 10 nm gold secondary (1/50 dilution) for 1 hour. After this the grid was
washed seven times in 1xPBS and twice in distilled water. Grids were then silver enhanced for 5
minutes (Nanoprobes). Grids were finally post stained with 1% uranyl acetate for 5 minutes
followed by two washes in water and then stained with lead citrate for 5 minutes followed by a
final two washes in distilled water. Samples were then imaged on a ThermoFisher Talos L120C
operating at 120kV.

Western blotting*:* Samples for WB were prepared by homogenizing brains in a tissue-
homogenizing buffer (250 mM sucrose, 20 mM Tris pH 7.4, 1 mM EDTA, 1 mM EGTA)
containing protease and phosphatase inhibitors as previously described. (Schmidt, Nixon et al.
2012) Following electrophoresis, proteins were transferred onto 0.2 μm-pore nitrocellulose
membranes (Whatman, Florham Park, NJ) at 100 mA for 8-12 hours depending on the target

protein. The blots were blocked for 1 hour in 5% non-fat milk in TBS, rinsed in TBST (TBS +

0.1% Tween-20), then incubated with a primary antibody in 1% BSA/TBST overnight at 4°C. The

membrane was washed and incubated in a HRP conjugated goat-anti rabbit or mouse secondary

antibody, diluted 1:5000 in 5% milk for 1 hr at room temperature. The membrane was again

washed and then incubated in a Novex ECL (Invitrogen) for 1 min. The detection of the signals

was achieved through either exposure to a film or scanning by a digital gel imager (Syngene G:Box

XX9) as specified in the figure legends. Densitometry was performed with Image J and the results

were normalized by the immunoblot(s) of given loading control protein(s) (usually GAPDH,

unless otherwise noted).

824   References

826   Abd-Elrahman, K. S., A. Hamilton, S. R. Hutchinson, F. Liu, R. C. Russell and S. S. G. Ferguson (2017).
827   "mGluR5 antagonism increases autophagy and prevents disease progression in the zQ175 mouse
828   model of Huntington's disease." <u>Sci Signal</u> **10**(510).
829   Aktar, F., C. Burudpakdee, M. Polanco, S. Pei, T. C. Swayne, P. N. Lipke and L. Emtage (2019). "The
830   huntingtin inclusion is a dynamic phase-separated compartment." <u>Life Sci Alliance</u> **2**(5).
831   Ashkenazi, A., C. F. Bento, T. Ricketts, M. Vicinanza, F. Siddiqi, M. Pavel, F. Squitieri, M. C. Hardenberg, S.
832   Imarisio, F. M. Menzies and D. C. Rubinsztein (2017). "Polyglutamine tracts regulate beclin 1-
833   dependent autophagy." <u>Nature</u> **545**(7652): 108-111.
834   Atwal, R. S., J. Xia, D. Pinchev, J. Taylor, R. M. Epand and R. Truant (2007). "Huntingtin has a membrane
835   association signal that can modulate huntingtin aggregation, nuclear entry and toxicity." <u>Hum Mol</u>
836   <u>Genet</u> **16**(21): 2600-2615.
837   Baiamonte, B. A., F. A. Lee, S. T. Brewer, D. Spano and G. J. LaHoste (2013). "Attenuation of Rhes activity
838   significantly delays the appearance of behavioral symptoms in a mouse model of Huntington's
839   disease." <u>PLoS One</u> **8**(1): e53606.
840   Barker, R. A., M. Fujimaki, P. Rogers and D. C. Rubinsztein (2020). "Huntingtin-lowering strategies for
841   Huntington's disease." <u>Expert Opin Investig Drugs</u> **29**(10): 1125-1132.
842   Bates, G. P., R. Dorsey, J. F. Gusella, M. R. Hayden, C. Kay, B. R. Leavitt, M. Nance, C. A. Ross, R. I. Scahill,
843   R. Wetzel, E. J. Wild and S. J. Tabrizi (2015). "Huntington disease." <u>Nat Rev Dis Primers</u> **1**: 15005.
844   Bensalem, J., C. Fourrier, L. K. Hein, S. Hassiotis, C. G. Proud and T. J. Sargeant (2021). "Inhibiting mTOR
845   activity using AZD2014 increases autophagy in the mouse cerebral cortex." <u>Neuropharmacology</u>
846   **190**: 108541.
847   Berg, M. J., Veeranna, C. M. Peterhoff, A. Kumar, P. S. Mohan, P. Stavrides, D. M. Marchionini, D. S. Yang
848   and R. A. Nixon (2024). "Pathobiology of the autophagy-lysosomal pathway in the Huntington's
849   disease brain." <u>bioRxiv</u>.
850   Bjørkøy, G., T. Lamark, A. Brech, H. Outzen, M. Perander, A. Øvervatn, H. Stenmark and T. Johansen (2005).
851   "p62/SQSTM1 forms protein aggregates degraded by autophagy and has a protective effect on
852   huntingtin-induced cell death." <u>J Cell Biol</u> **171**(4): 603-614.
853   Boland, B., W. H. Yu, O. Corti, B. Mollereau, A. Henriques, E. Bezard, G. M. Pastores, D. C. Rubinsztein, R.
854   A. Nixon, M. R. Duchen, G. R. Mallucci, G. Kroemer, B. Levine, E. L. Eskelinen, F. Mochel, M.
855   Spedding, C. Louis, O. R. Martin and M. J. Millan (2018). "Promoting the clearance of neurotoxic
856   proteins in neurodegenerative disorders of ageing." <u>Nat Rev Drug Discov</u> **17**(9): 660-688.
857   Carty, N., N. Berson, K. Tillack, C. Thiede, D. Scholz, K. Kottig, Y. Sedaghat, C. Gabrysiak, G. Yohrling, H. von
858   der Kammer, A. Ebneth, V. Mack, I. Munoz-Sanjuan and S. Kwak (2015). "Characterization of HTT
859   inclusion size, location, and timing in the zQ175 mouse model of Huntington's disease: an in vivo
860   high-content imaging study." <u>PLoS One</u> **10**(4): e0123527.
861   Cataldo, A. M., C. Y. Thayer, E. D. Bird, T. R. Wheelock and R. A. Nixon (1990). "Lysosomal proteinase
862   antigens are prominently localized within senile plaques of Alzheimer's disease: evidence for a
863   neuronal origin." <u>Brain Res</u> **513**(2): 181-192.
864   Cortes, C. J. and A. R. La Spada (2014). "The many faces of autophagy dysfunction in Huntington's disease:
865   from mechanism to therapy." <u>Drug Discov Today</u> **19**(7): 963-971.
866   Croce, K. R. and A. Yamamoto (2019). "A role for autophagy in Huntington's disease." <u>Neurobiol Dis</u> **122**:
867   16-22.
868   Deng, Y., H. Wang, M. Joni, R. Sekhri and A. Reiner (2021). "Progression of basal ganglia pathology in
869   heterozygous Q175 knock-in Huntington's disease mice." <u>J Comp Neurol</u> **529**(7): 1327-1371.

DiFiglia, M., E. Sapp, K. O. Chase, S. W. Davies, G. P. Bates, J. P. Vonsattel and N. Aronin (1997).
"Aggregation of huntingtin in neuronal intranuclear inclusions and dystrophic neurites in brain."
Science **277**(5334): 1990-1993.
Franklin, G. L., H. A. G. Teive, F. S. Tensini, C. H. F. Camargo, N. S. C. de Lima, D. C. de Dos Santos, A. T.
Meira and S. J. Tabrizi (2024). "The Huntington's Disease Gene Discovery." Mov Disord **39**(2): 227-
234.
Fu, A., V. Cohen-Kaplan, N. Avni, I. Livneh and A. Ciechanover (2021). "p62-containing, proteolytically
active nuclear condensates, increase the efficiency of the ubiquitin-proteasome system." Proc
Natl Acad Sci U S A **118**(33).
Gutekunst, C. A., S. H. Li, H. Yi, J. S. Mulroy, S. Kuemmerle, R. Jones, D. Rye, R. J. Ferrante, S. M. Hersch
and X. J. Li (1999). "Nuclear and neuropil aggregates in Huntington's disease: relationship to
neuropathology." J Neurosci **19**(7): 2522-2534.
Heikkinen, T., T. Bragge, J. Kuosmanen, T. Parkkari, S. Gustafsson, M. Kwan, J. Beltran, A. Ghavami, S.
Subramaniam, N. Shahani, U. N. Ramirez-Jarquin, L. Park, I. Munoz-Sanjuan and D. M. Marchionini
(2021). "Global Rhes knockout in the Q175 Huntington's disease mouse model." PLoS One **16**(10):
e0258486.
Hickey, M. A., A. Kosmalska, J. Enayati, R. Cohen, S. Zeitlin, M. S. Levine and M. F. Chesselet (2008).
"Extensive early motor and non-motor behavioral deficits are followed by striatal neuronal loss in
knock-in Huntington's disease mice." Neuroscience **157**(1): 280-295.
Iwata, A., J. C. Christianson, M. Bucci, L. M. Ellerby, N. Nukina, L. S. Forno and R. R. Kopito (2005).
"Increased susceptibility of cytoplasmic over nuclear polyglutamine aggregates to autophagic
degradation." Proc Natl Acad Sci U S A **102**(37): 13135-13140.
Iwata, A., Y. Nagashima, L. Matsumoto, T. Suzuki, T. Yamanaka, H. Date, K. Deoka, N. Nukina and S. Tsuji
(2009). "Intranuclear degradation of polyglutamine aggregates by the ubiquitin-proteasome
system." J Biol Chem **284**(15): 9796-9803.
Jamilloux, Y., B. Lagrange, A. Di Micco, E. Bourdonnay, A. Provost, R. Tallant, T. Henry and F. Martinon
(2018). "A proximity-dependent biotinylation (BioID) approach flags the p62/sequestosome-1
protein as a caspase-1 substrate." J Biol Chem **293**(32): 12563-12575.
Jiang, A., R. R. Handley, K. Lehnert and R. G. Snell (2023). "From Pathogenesis to Therapeutics: A Review
of 150 Years of Huntington's Disease Research." Int J Mol Sci **24**(16).
Jiang, Y., J. J. Alam, S. N. Gomperts, P. Maruff, A. W. Lemstra, U. A. Germann, P. H. Stavrides, S. Darji, S.
Malampati, J. Peddy, C. Bleiwas, M. Pawlik, A. Pensalfini, D. S. Yang, S. Subbanna, B. S.
Basavarajappa, J. F. Smiley, A. Gardner, K. Blackburn, H. M. Chu, N. D. Prins, C. E. Teunissen, J. E.
Harrison, P. Scheltens and R. A. Nixon (2022). "Preclinical and randomized clinical evaluation of
the p38alpha kinase inhibitor neflamapimod for basal forebrain cholinergic degeneration." Nat
Commun **13**(1): 5308.
Kim, A., K. Lalonde, A. Truesdell, P. Gomes Welter, P. S. Brocardo, T. R. Rosenstock and J. Gil-Mohapel
(2021). "New Avenues for the Treatment of Huntington's Disease." Int J Mol Sci **22**(16).
Kim, Y. C. and K. L. Guan (2015). "mTOR: a pharmacologic target for autophagy regulation." J Clin Invest
125(1): 25-32.
Kim, Y. J., Y. Yi, E. Sapp, Y. Wang, B. Cuiffo, K. B. Kegel, Z. H. Qin, N. Aronin and M. DiFiglia (2001). "Caspase
3-cleaved N-terminal fragments of wild-type and mutant huntingtin are present in normal and
Huntington's disease brains, associate with membranes, and undergo calpain-dependent
proteolysis." Proc Natl Acad Sci U S A **98**(22): 12784-12789.
Klionsky, D. J., G. Petroni, R. K. Amaravadi, E. H. Baehrecke, A. Ballabio, P. Boya, J. M. Bravo-San Pedro, K.
Cadwell, F. Cecconi, A. M. K. Choi, M. E. Choi, C. T. Chu, P. Codogno, M. I. Colombo, A. M. Cuervo,
V. Deretic, I. Dikic, Z. Elazar, E. L. Eskelinen, G. M. Fimia, D. A. Gewirtz, D. R. Green, M. Hansen, M.

Jaattela, T. Johansen, G. Juhasz, V. Karantza, C. Kraft, G. Kroemer, N. T. Ktistakis, S. Kumar, C. Lopez-Otin, K. F. Macleod, F. Madeo, J. Martinez, A. Melendez, N. Mizushima, C. Munz, J. M. Penninger, R. M. Perera, M. Piacentini, F. Reggiori, D. C. Rubinsztein, K. M. Ryan, J. Sadoshima, L. Santambrogio, L. Scorrano, H. U. Simon, A. K. Simon, A. Simonsen, A. Stolz, N. Tavernarakis, S. A. Tooze, T. Yoshimori, J. Yuan, Z. Yue, Q. Zhong, L. Galluzzi and F. Pietrocola (2021). "Autophagy in major human diseases." EMBO J **40**(19): e108863.

Ko, J., J. M. Isas, A. Sabbaugh, J. H. Yoo, N. K. Pandey, A. Chongtham, M. Ladinsky, W. L. Wu, H. Rohweder, A. Weiss, D. Macdonald, I. Munoz-Sanjuan, R. Langen, P. H. Patterson and A. Khoshnan (2018). "Identification of distinct conformations associated with monomers and fibril assemblies of mutant huntingtin." Hum Mol Genet **27**(13): 2330-2343.

Kolla, R., P. Gopinath, J. Ricci, A. Reif, I. Rostami and H. A. Lashuel (2021). "A New Chemoenzymatic Semisynthetic Approach Provides Insight into the Role of Phosphorylation beyond Exon1 of Huntingtin and Reveals N-Terminal Fragment Length-Dependent Distinct Mechanisms of Aggregation." J Am Chem Soc **143**(26): 9798-9812.

Lee, J. H., M. V. Rao, D. S. Yang, P. Stavrides, E. Im, A. Pensalfini, C. Huo, P. Sarkar, T. Yoshimori and R. A. Nixon (2019). "Transgenic expression of a ratiometric autophagy probe specifically in neurons enables the interrogation of brain autophagy in vivo." Autophagy **15**(3): 543-557.

Lee, J. H., L. Tecedor, Y. H. Chen, A. M. Monteys, M. J. Sowada, L. M. Thompson and B. L. Davidson (2015). "Reinstating aberrant mTORC1 activity in Huntington's disease mice improves disease phenotypes." Neuron **85**(2): 303-315.

Lee, J. H., D. S. Yang, C. N. Goulbourne, E. Im, P. Stavrides, A. Pensalfini, H. Chan, C. Bouchet-Marquis, C. Bleiwas, M. J. Berg, C. Huo, J. Peddy, M. Pawlik, E. Levy, M. Rao, M. Staufenbiel and R. A. Nixon (2022). "Faulty autolysosome acidification in Alzheimer's disease mouse models induces autophagic build-up of Abeta in neurons, yielding senile plaques." Nat Neurosci **25**(6): 688-701.

Li, H., S. H. Li, A. L. Cheng, L. Mangiarini, G. P. Bates and X. J. Li (1999). "Ultrastructural localization and progressive formation of neuropil aggregates in Huntington's disease transgenic mice." Hum Mol Genet **8**(7): 1227-1236.

Lie, P. P. Y., D. S. Yang, P. Stavrides, C. N. Goulbourne, P. Zheng, P. S. Mohan, A. M. Cataldo and R. A. Nixon (2021). "Post-Golgi carriers, not lysosomes, confer lysosomal properties to pre-degradative organelles in normal and dystrophic axons." Cell Rep **35**(4): 109034.

Lie, P. P. Y., L. Yoo, C. N. Goulbourne, M. J. Berg, P. Stavrides, C. Huo, J. H. Lee and R. A. Nixon (2022). "Axonal transport of late endosomes and amphisomes is selectively modulated by local Ca(2+) efflux and disrupted by PSEN1 loss of function." Sci Adv **8**(17): eabj5716.

Liu, K. Y., Y. C. Shyu, B. A. Barbaro, Y. T. Lin, Y. Chern, L. M. Thompson, C. K. James Shen and J. L. Marsh (2015). "Disruption of the nuclear membrane by perinuclear inclusions of mutant huntingtin causes cell-cycle re-entry and striatal cell death in mouse and cell models of Huntington's disease." Hum Mol Genet **24**(6): 1602-1616.

Marchionini, D. M., J. P. Liu, A. Ambesi-Impiombato, K. Kerker, K. Cirillo, M. Bansal, R. Mushlin, D. Brunner, S. Ramboz, M. Kwan, K. Kuhlbrodt, K. Tillack, F. Peters, L. Rauhala, J. Obenauer, J. R. Greene, C. Hartl, V. Khetarpal, B. Lager, J. Rosinski, J. Aaronson, M. Alam, E. Signer, I. Munoz-Sanjuan, D. Howland and S. O. Zeitlin (2022). "Benefits of global mutant huntingtin lowering diminish over time in a Huntington's disease mouse model." JCI Insight **7**(20).

Mario Isas, J., N. K. Pandey, H. Xu, K. Teranishi, A. K. Okada, E. K. Fultz, A. Rawat, A. Applebaum, F. Meier, J. Chen, R. Langen and A. B. Siemer (2021). "Huntingtin fibrils with different toxicity, structure, and seeding potential can be interconverted." Nat Commun **12**(1): 4272.

Martin, D. D., S. Ladha, D. E. Ehrnhoefer and M. R. Hayden (2015). "Autophagy in Huntington disease and huntingtin in autophagy." Trends Neurosci **38**(1): 26-35.

Martinez-Vicente, M., Z. Talloczy, E. Wong, G. Tang, H. Koga, S. Kaushik, R. de Vries, E. Arias, S. Harris, D. Sulzer and A. M. Cuervo (2010). "Cargo recognition failure is responsible for inefficient autophagy in Huntington's disease." Nat Neurosci **13**(5): 567-576.

Menalled, L. B., A. E. Kudwa, S. Miller, J. Fitzpatrick, J. Watson-Johnson, N. Keating, M. Ruiz, R. Mushlin, W. Alosio, K. McConnell, D. Connor, C. Murphy, S. Oakeshott, M. Kwan, J. Beltran, A. Ghavami, D. Brunner, L. C. Park, S. Ramboz and D. Howland (2012). "Comprehensive behavioral and molecular characterization of a new knock-in mouse model of Huntington's disease: zQ175." PLoS One **7**(12): e49838.

Menalled, L. B., J. D. Sison, I. Dragatsis, S. Zeitlin and M. F. Chesselet (2003). "Time course of early motor and neuropathological anomalies in a knock-in mouse model of Huntington's disease with 140 CAG repeats." J Comp Neurol **465**(1): 11-26.

Mende-Mueller, L. M., T. Toneff, S. R. Hwang, M. F. Chesselet and V. Y. Hook (2001). "Tissue-specific proteolysis of Huntingtin (htt) in human brain: evidence of enhanced levels of N- and C-terminal htt fragments in Huntington's disease striatum." J Neurosci **21**(6): 1830-1837.

Nixon, R. A. (2013). "The role of autophagy in neurodegenerative disease." Nat Med **19**(8): 983-997.

Norman, J. M., G. M. Cohen and E. T. Bampton (2010). "The in vitro cleavage of the hAtg proteins by cell death proteases." Autophagy **6**(8): 1042-1056.

Ochaba, J., T. Lukacsovich, G. Csikos, S. Zheng, J. Margulis, L. Salazar, K. Mao, A. L. Lau, S. Y. Yeung, S. Humbert, F. Saudou, D. J. Klionsky, S. Finkbeiner, S. O. Zeitlin, J. L. Marsh, D. E. Housman, L. M. Thompson and J. S. Steffan (2014). "Potential function for the Huntingtin protein as a scaffold for selective autophagy." Proc Natl Acad Sci U S A **111**(47): 16889-16894.

Ouimet, C. C., K. C. Langley-Gullion and P. Greengard (1998). "Quantitative immunocytochemistry of DARPP-32-expressing neurons in the rat caudatoputamen." Brain Res **808**(1): 8-12.

Park, J. M., C. H. Jung, M. Seo, N. M. Otto, D. Grunwald, K. H. Kim, B. Moriarity, Y. M. Kim, C. Starker, R. S. Nho, D. Voytas and D. H. Kim (2016). "The ULK1 complex mediates MTORC1 signaling to the autophagy initiation machinery via binding and phosphorylating ATG14." Autophagy **12**(3): 547-564.

Peng, Q., B. Wu, M. Jiang, J. Jin, Z. Hou, J. Zheng, J. Zhang and W. Duan (2016). "Characterization of Behavioral, Neuropathological, Brain Metabolic and Key Molecular Changes in zQ175 Knock-In Mouse Model of Huntington's Disease." PLoS One **11**(2): e0148839.

Peskett, T. R., F. Rau, J. O'Driscoll, R. Patani, A. R. Lowe and H. R. Saibil (2018). "A Liquid to Solid Phase Transition Underlying Pathological Huntingtin Exon1 Aggregation." Mol Cell **70**(4): 588-601.e586.

Pressl, C., K. Mätlik, L. Kus, P. Darnell, J. D. Luo, M. R. Paul, A. R. Weiss, W. Liguore, T. S. Carroll, D. A. Davis, J. McBride and N. Heintz (2024). "Selective vulnerability of layer 5a corticostriatal neurons in Huntington's disease." Neuron.

Pryor, W. M., M. Biagioli, N. Shahani, S. Swarnkar, W. C. Huang, D. T. Page, M. E. MacDonald and S. Subramaniam (2014). "Huntingtin promotes mTORC1 signaling in the pathogenesis of Huntington's disease." Sci Signal **7**(349): ra103.

Ravikumar, B., C. Vacher, Z. Berger, J. E. Davies, S. Luo, L. G. Oroz, F. Scaravilli, D. F. Easton, R. Duden, C. J. O'Kane and D. C. Rubinsztein (2004). "Inhibition of mTOR induces autophagy and reduces toxicity of polyglutamine expansions in fly and mouse models of Huntington disease." Nat Genet **36**(6): 585-595.

Riguet, N., A. L. Mahul-Mellier, N. Maharjan, J. Burtscher, M. Croisier, G. Knott, J. Hastings, A. Patin, V. Reiterer, H. Farhan, S. Nasarov and H. A. Lashuel (2021). "Nuclear and cytoplasmic huntingtin inclusions exhibit distinct biochemical composition, interactome and ultrastructural properties." Nat Commun **12**(1): 6579.

Rose, C., F. M. Menzies, M. Renna, A. Acevedo-Arozena, S. Corrochano, O. Sadiq, S. D. Brown and D. C. Rubinsztein (2010). "Rilmenidine attenuates toxicity of polyglutamine expansions in a mouse model of Huntington's disease." Hum Mol Genet **19**(11): 2144-2153.

Rothe, T., M. Deliano, A. M. Wojtowicz, A. Dvorzhak, D. Harnack, S. Paul, T. Vagner, I. Melnick, H. Stark and R. Grantyn (2015). "Pathological gamma oscillations, impaired dopamine release, synapse loss and reduced dynamic range of unitary glutamatergic synaptic transmission in the striatum of hypokinetic Q175 Huntington mice." Neuroscience **311**: 519-538.

Rui, Y. N., Z. Xu, B. Patel, Z. Chen, D. Chen, A. Tito, G. David, Y. Sun, E. F. Stimming, H. J. Bellen, A. M. Cuervo and S. Zhang (2015). "Huntingtin functions as a scaffold for selective macroautophagy." Nat Cell Biol **17**(3): 262-275.

Sanchez-Garrido, J., V. Sancho-Shimizu and A. R. Shenoy (2018). "Regulated proteolysis of p62/SQSTM1 enables differential control of autophagy and nutrient sensing." Sci Signal **11**(559).

Sarkar, S. and D. C. Rubinsztein (2008). "Huntington's disease: degradation of mutant huntingtin by autophagy." FEBS J **275**(17): 4263-4270.

Sarkar, S. and D. C. Rubinsztein (2008). "Small molecule enhancers of autophagy for neurodegenerative diseases." Mol Biosyst **4**(9): 895-901.

Schmidt, S. D., R. A. Nixon and P. M. Mathews (2012). "Tissue processing prior to analysis of Alzheimer's disease associated proteins and metabolites, including Abeta." Methods Mol Biol **849**: 493-506.

Sotrel, A., P. A. Paskevich, D. K. Kiely, E. D. Bird, R. S. Williams and R. H. Myers (1991). "Morphometric analysis of the prefrontal cortex in Huntington's disease." Neurology **41**(7): 1117-1123.

Tanaka, M., Y. Machida, S. Niu, T. Ikeda, N. R. Jana, H. Doi, M. Kurosawa, M. Nekooki and N. Nukina (2004). "Trehalose alleviates polyglutamine-mediated pathology in a mouse model of Huntington disease." Nat Med **10**(2): 148-154.

The-Huntington's-Disease-Collaborative-Research-Group (1993). "A novel gene containing a trinucleotide repeat that is expanded and unstable on Huntington's disease chromosomes. ." Cell **72**(6): 971-983.

Trottier, Y., Y. Lutz, G. Stevanin, G. Imbert, D. Devys, G. Cancel, F. Saudou, C. Weber, G. David, L. Tora and et al. (1995). "Polyglutamine expansion as a pathological epitope in Huntington's disease and four dominant cerebellar ataxias." Nature **378**(6555): 403-406.

Valionyte, E., Y. Yang, S. A. Griffiths, A. T. Bone, E. R. Barrow, V. Sharma, B. Lu and S. Luo (2022). "The caspase-6-p62 axis modulates p62 droplets based autophagy in a dominant-negative manner." Cell Death Differ **29**(6): 1211-1227.

Vodicka, P., J. Lim, D. T. Williams, K. B. Kegel, K. Chase, H. Park, D. Marchionini, S. Wilkinson, T. Mead, H. Birch, D. Yates, K. Lyons, C. Dominguez, M. Beconi, Z. Yue, N. Aronin and M. DiFiglia (2014). "Assessment of chloroquine treatment for modulating autophagy flux in brain of WT and HD mice." J Huntingtons Dis **3**(2): 159-174.

Vonsattel, J. P. (2008). "Huntington disease models and human neuropathology: similarities and differences." Acta Neuropathol **115**(1): 55-69.

Vonsattel, J. P., R. H. Myers, T. J. Stevens, R. J. Ferrante, E. D. Bird and E. P. Richardson, Jr. (1985). "Neuropathological classification of Huntington's disease." J Neuropathol Exp Neurol **44**(6): 559-577.

Wold, M. S., J. Lim, V. Lachance, Z. Deng and Z. Yue (2016). "ULK1-mediated phosphorylation of ATG14 promotes autophagy and is impaired in Huntington's disease models." Mol Neurodegener **11**(1): 76.

Xia, J., D. H. Lee, J. Taylor, M. Vandelft and R. Truant (2003). "Huntingtin contains a highly conserved nuclear export signal." Hum Mol Genet **12**(12): 1393-1403.

Yang, D. S., J. H. Lee and R. A. Nixon (2009). "Monitoring autophagy in Alzheimer's disease and related
neurodegenerative diseases." Methods Enzymol **453**: 111-144.
Yang, J. and C. Zhang (2023). "Targeting the autophagy-lysosomal pathway in Huntington disease: a
pharmacological perspective." Front Aging Neurosci **15**: 1175598.
Zhou, Y., T. R. Peskett, C. Landles, J. B. Warner, K. Sathasivam, E. J. Smith, S. Chen, R. Wetzel, H. A. Lashuel,
G. P. Bates and H. R. Saibil (2021). "Correlative light and electron microscopy suggests that mutant
huntingtin dysregulates the endolysosomal pathway in presymptomatic Huntington's disease."
Acta Neuropathol Commun **9**(1): 70.
1064

Fig. 1. Identification of inclusions and HTT molecular species in Q175 mice.

Fig. 2. HTT colocalizes with p62, Ub and CTSD

Fig. 3. Mild late-onset alterations in the ALP in the striatum of 17-mo-old Q175.

Fig. 4. DARPP-32-IR decreases in striatal neurons of 17-mo-old TRGL/Q175 in the absence of a NeuN signal reduction.

Fig. 5. mTORi INK exhibits target engagement and induces downstream responses in the ALP in 7-mo-old TRGL/Q175.

Fig. 6. mTORi INK reverses the mild alteration of AV/LY subtypes in the striatum of 7-mo-old TRGL/Q175.

Fig. 7. mTORi INK reduces HTT-, Ub- or p62-IR-covered areas parallelly in the striatum of 7-mo-old TRGL/Q175.

Fig. 8. INK treatment does not reverse the reduction of DARPP-32-IR in the striatum of 7-mo-old TRGL/Q175.

Fig. S1. Colocalization of HTT/Ub and Ub/p62 in IBs in the STR.

Fig. S2 (related to Fig. 3). Molecules involved in various stages of the ALP are largely unchanged in 17-mo-old TRGL/Q175.

Fig. S2 (related Fig. 3). Molecules involved in various stages of the ALP are largely unchanged in 17-mo-old TRGL/Q175. (Cont'd)

Fig. S2 (related to Fig. 3). Molecules involved in various stages of the ALP are largely unchanged in 17-mo-old TRGL/Q175. (Cont'd)

Fig. S3 (related to Fig. 5). Dosing tests demonstrate mTORi INK BBB penetration and target engagement even at low dosages.