Rapid Changes in Chromosome Counts in Fishes from the Spiral Egg Clade within the Gourami Family (Osphronemidae).

Brendan Mobley and Andrew P. Anderson

Biology Department, Reed College, Portland, Oregon 97202, USA

Abstract

Identifying clades with numerous and noticeable changes in chromosome counts is an important step in unraveling the evolutionary mechanisms that shape cytogenetic processes. Here, we describe low chromosome counts in a group of teleost fishes delimited by their unique spiral egg structure and with a species with a known low chromosome count within the labyrinthine clade (Osphronemidae). We sampled seven of nine known species within this spiral egg clade, reporting novel chromosome counts for five species and confirming two others. Overall, we find high variability in both chromosome count and arm number, which suggests a rapid loss of chromosomes during the emergence of the clade and numerous large-scale mutations occurring across evolutionary time. Lastly, we offer some possible explanations for these changes based on current and ongoing empirical and theoretical research. These data provide important information in cataloguing rapid chromosomal shifts in teleost fishes and highlights this group for further study in chromosomal and genomic evolution due to their karyotypic heterogeneity.

Keywords

Cytogenetics, evolution, genomics, karyotype, fish

Introduction

Variability of chromosome numbers across vertebrates and the evolutionary mechanisms that create it is an active point of inquiry for evolutionary biologists (Martinez et al. 2015). Finding clades with high variance or rapid changes presents valuable data points in resolving the various hypotheses on how karyotypes change over time. Some patterns can include changes due to long stretches of repeat content creating opportunities for mismatched recombination (Amores et al. 2014) such as found in mammals which range in chromosome counts from 2N=6 in the female muntjac deer (*Muntiacus muntjac*) (Wurster and Benirschke 1970; Graphodatsky et al. 2020) to 2N=102 in the plains viscacha rat (*Tympanoctomys barrerae*) (Gallardo et al. 2006; Stanyon and Graphodatsky 2012; Lebeda et al. 2020). Other patterns can involve microchromosomes as typically seen in birds which have a range from 2N=40 in *Ceratogymna bucinator* to 2N=136–142 in *Corythaixoides concolor* (Christidis 1990; Kretschmer et al. 2018). Conversely, non-avian reptiles are karyologically heterogeneous and exhibit distinct evolutionary trends between lineages (Deakin and Ezaz 2019) and have a narrower range of diploid chromosome counts than other groups (2N=24–70, Olmo 2005). Teleost fishes have been found to have notable karyotype evolution patterns, particularly with regard to extreme chromosome counts, rapid cytogenetic changes, or both (e.g. *Nothobranchius* 2N=16–50:Krysanov et al. 2023 and *Corydoras* 2N=40–134: Shimabukuro-Dias et al. 2004) and have the widest 2N range of all vertebrates, ranging from 2N=12 in the marine species *Gonostoma bathyphilum* to 2N=446 in the freshwater species *Ptychobarbus dipogon* (Lebeda et al. 2020). Paradoxically, teleost fishes have a strong trend of conserved karyotypes (Galetti et al. 2000; Mank and Avise 2006; Nakatani et al. 2007) with over half of all karyotyped fish species having

diploid chromosome counts of 48 or 50 (Mank and Avise 2006; Arai 2011), which has changed

little from the proposed karyotype for the ancestor of all teleost fish (2N=52, Nakatani et al.

2007). With such diverse patterns across vertebrates, finding clades with unusual changes in

chromosome counts is valuable to understanding chromosome evolution as whole.

An intriguing group of fishes with apparent chromosomal variance is within the family

Osphronemidae, commonly called gouramis. The chocolate gourami, *Sphaerichthys*

osphromenoides, has the lowest recorded chromosome count among freshwater fishes

(Lehmann et al. 2021) with 2n=16 (Calton and Denton 1974). A species in the neighboring

genus, the pikehead gourami, *Luciocephalus pulcher*, was reported to have 2n=20 (Arai 2011).

Chromosome counts this low are exceedingly rare in fishes, as there are only thirteen fish

species with a diploid chromosome count lower than 2N=22 (Lehmann et al. 2021).

Furthermore, these counts are highly derived from the other Osphronemidae species, which

generally have 2N values between 46 and 48 (Supplemental Table 1).

Both *S. osphromenoides* and *L. pulcher* are members of the "spiral egg" clade, a monophyletic

group within the family Osphronemidae that includes the genera *Sphaerichthys*, *Luciocephalus*,

Parasphaerichthys, and *Ctenops*. The monophyly was proposed based on the unique

morphology of their eggs, which are covered in projections arranged in a spiral pattern, and later

confirmed and refined with molecular evidence (Britz et al. 1995; Rüber et al. 2006). Another

differentiating feature of the spiral egg clade is an angular jaw shape, which is taken to the

extreme in the highly derived pike-like morphology of the piscivorous genus *Luciocephalus*. The

spiral egg clade is also notable for having the only species, *S. osphromenoides* and *S.*

selatanensis, in the family Osphronemidae with female broodcare via mouthbrooding compared

to the overwhelmingly male mouthbrooders or bubble nesters in the family (Rüber et al. 2006),

although recent evidence has called into question the sex of caring parent in *S.*

osphromenoides (Zworykin et al. 2024). Chromosomes of the spiral egg clade remain largely

uninvestigated; besides *S. osphromenoides* and *L. pulcher*, only one other species has been

studied cytogenetically (*Ctenops nobilis*, 2N=44: Rishi et al. 1997). Given the low chromosome

counts of *S. osphromenoides* and *L. pulcher* and the large 2N decrease relative to the wider

family, we aim to characterize the karyotypes of additional members within the spiral egg clade.

With this information we will describe the karyotypic diversity and evolutionary history for this

remarkable group of fishes, thereby adding an extraordinary example to the chromosome count

diversity in fishes specifically and animals in general.

82   Methods

Fishes were sourced from the aquarium trade (Wet Spot Tropical Fish, Portland, Oregon, USA;

Nationwide Aquatics, Tinley Park, Illinois, USA; Aqua Imports, Boulder, Colorado, USA), then

held in species-specific tanks (110 liters) on a shared flow-through system (pH 7.0, GH 30 ppm,

KH 40 ppm) with a 12/12 hour light/dark cycle with 30 minutes of dim light to simulate dawn and

dusk. Specimens were housed for a minimum of one week before sampling to ensure good

health for optimal cell proliferation.

Chromosome preparations were made following Kligerman and Bloom (1977) with the indicated

modifications. Specimens were incubated in 0.005% colchicine solution for 6-7 hours, then

euthanized and dissected to remove gill arches. Sex determination was conducted by gross

examination of gonads with pictures taken throughout. Dissected specimens were stored in -80

°C for future molecular analyses. Gill arches were incubated in 0.4% KCl solution for 20-30

minutes, then fixed in two changes of 3:1 ethanol:acetic acid fixative for at least 30 minutes

each, followed by an overnight fixation period at 4 °C. To prepare slides, tissue was

homogenized into suspension by mincing in 50% acetic acid, then dropped onto a slide warmed

to 30-40 °C and air dried. Slides were examined under phase contrast microscopy for quality

control, then aged for at least one day at room temperature before being stained for 10 minutes

in 10% Giemsa in pH 6.8 phosphate buffer (Gibco™ Gurr Buffer Tablets) and air-dried.

Chromosomes were examined under a Nikon Eclipse Ti-E microscope driven by Nikon NIS

Elements AR software, then photographed with an oil immersion objective at 100x magnification

and green color filtering using a Hamatsu ORCA-Flash4.0 camera. Digital images were

optimized, then homologous chromosomes were paired by size and morphology and arranged

in decreasing order of size using ImageJ v1.52v and Adobe Photoshop 24.3.0. At least 35

complete metaphase spreads were photographed from each specimen with completeness

defined as the highest consistently observed chromosome count. Chromosomes were classified

as metacentric (m), submetacentric (sm), subtelocentric (st), or acrocentric (a) according to their

arm ratios (Levan et al. 1964). Chromosome arm number (Fundamental Number, FN) was

calculated by considering metacentric and submetacentric chromosomes as biarmed and

subtelocentric and acrocentric chromosomes as uniarmed.

Results

We describe karyotypes for the first time in six species (Fig 1): *S. selatanensis*, *S. vaillanti*, *S.*

acrostoma, *L. aura*, and *P. ocellatus*. We also confirmed the karyotypes of an additional two

species (*S. osphromenoides*, *L. pulcher*) which matched those established in the literature

(Calton and Denton 1974; Arai 2011). All species in the genus *Sphaerichthys* had different

chromosome counts, with 2N ranging from 16–28 (Table 1). The number of chromosome arms

(fundamental number, FN) showed less variation, with a range of 30–38. Notably, the sister

species *S. osphromenoides* and *S. selatanensis* had a primarily biarmed and primarily uniarmed

karyotype respectively, resulting in a nearly identical FN despite a 2N difference of 12. The

other two sister species in the genus (*S. acrostoma* and *S. vaillanti*) had karyotypes that were a

mix of biarmed and uniarmed chromosomes and had different values for both 2N and FN. We

confirmed that *L. pulcher* had an entirely uniarmed karyotype of 20 acrocentric chromosomes.

Luciocephalus aura had a primarily biarmed karyotype with six fewer chromosomes and eight

more chromosome arms. *Parasphaerichthys ocellatus* had a primarily uniarmed karyotype with

higher 2N and FN that were higher than any *Sphaerichthys* or *Luciocephalus* species, but lower

than was reported for *C. nobilis* in the literature (Arai 2011).

The two *S. selatanensis* that we sampled had different karyotypes (Supplemental Table 2). One

had 28 uniarmed chromosomes (2N=28, karyotype 2st+26a), while the other had 26 uniarmed

chromosomes and an unpaired metacentric chromosome (2N=27, karyotype 1m+2st+24a). The

unpaired metacentric chromosome was approximately twice the size of the largest acrocentric

chromosomes and may have been caused by a fused acrocentric pair. We cannot say whether

this is a sex chromosome because we could not confidently determine the sex of either

individual.

Species	2N	FN	Karyotype
Sphaerichthys osphromenoides	16	30	10m + 4sm + 2a
Sphaerichthys selatanensis	28	28	2st + 26a
Sphaerichthys vaillanti	20	34	6m + 8sm + 2st + 4a
Sphaerichthys acrostoma	24	38	4m + 10sm + 10a
Luciocephalus pulcher	20	20	20a
Luciocephalus aura	14	24	8m + 2sm + 4st
Parasphaerichthys ocellatus	34	44	6m + 4sm + 22st + 2a
Ctenops nobilis	44	60	8m + 8sm + 28a
Trichopodus leerii	46	46	46a
Osphronemus goramy	48	50	2sm + 46a
Helostoma temminckii	48	48	48a

5 MYA

Figure 1. Selected anabantoid karyotypes. Phylogenetic relationships are from Ruber et al. (2006) and are shown to scale for the
spiral egg clade (red) but not the selected species in the family Osphronemidae (black) or the outgroup (Helostomatidae, grey).
Values for 2N, FN, and Karyotype for four species not generated in this study can be found in Arai (2011) and Grazyna et al. (2008).
Karyotype formula describes number of metacentric (m), submetacentric (sm), subtelocentric (st), and acrocentric (a) chromosomes.
Note that chromosome sizes vary widely between spreads and are thus not directly comparable except within the same spread. Live
specimens were photographed with iPhone 13.

Discussion

We found that the genera *Sphaerichthys*, *Luciocephalus*, and *Parasphaerichthys* have low

chromosome counts (2N≤34) with high intra-genus variation in both 2N and FN. These trends

are not observed in the karyotypes of the broader family Osphronemidae, which mostly have

karyotypes similar to the hypothesized ancestral state for all teleost fishes (2N=52, Nakatani et

al. 2007), thereby suggesting that rapid karyotype evolution occurred since the divergence of

the spiral egg clade about 25 million years ago (Rüber et al. 2006). Karyotype evolution of this

rate and magnitude has not been reported in teleost fishes. The drastic differences between

karyotypes within the *Sphaerichthys* and *Luciocephalus* genera suggests that karyotype

evolution may have played a role in speciation process by creating post-zygotic isolation (Canitz

et al. 2016; Jackson et al. 2016; Mezzasalma et al. 2017; Romanenko et al. 2018), but it is also

possible that the observed karyotype diversity happened alongside the speciation process

instead of driving it (Krysanov et al. 2023). The observed karyotype pattern could have been

created by the influence of genetic drift or other forms of neutral selection, extremely strong

meiotic drive, the evolution of a trait that stimulates chromosome evolution, or a combination of

these factors.

Genetic drift has been proposed to be the driving force behind the fixation of highly

differentiated karyotypes in some clades of freshwater fishes, including the annual killifishes in

the genera *Nothobranchius* (2N=16-50) (Krysanov et al. 2016, 2023; Krysanov and Demidova

2018) and *Aphyosemion* (2N=20-40) (Völker et al. 2005, 2007, 2008). These species tend to

live in small, biogeographically isolated populations and frequently experience genetic

bottlenecks and founder effects due to the ephemeral nature of their habitats (Völker et al. 2006;

Krysanov and Demidova 2018; Krysanov et al. 2023), which may have sped up the

accumulation of both intra-and inter-chromosomal mutations, with centric fusions being

responsible for most decreases in chromosome count (Völker et al. 2005, 2008; Krysanov et al.

2016). *Sphaerichthys* and *Luciocephalus* species have very limited geographical ranges and,

given that most of them are threatened or endangered according to the IUCN Red List, these

populations are likely small; however, it is difficult to precisely estimate the strength of genetic

drift on the evolution of the karyotypes in our study group because there is little information

about their distribution and population structure.

An alternative explanation to genetic drift is meiotic drive. Species evolving under the influence

of strong meiotic drive will tend to reach karyotypes of predominantly biarmed or uniarmed

chromosomes (de Villena and Sapienza 2001; Molina et al. 2014). This has been observed in

fishes and mammals, with the note that groups with high rates of mismatched karyotypes

tended to have higher rates of chromosome evolution (Blackmon et al. 2019). The decrease in

chromosome number in our study group relative to the rest of the family may have been caused

by a meiotic drive toward biarmed chromosomes that rapidly fixed fusion mutations.

Additionally, the near-complete inversions in biarmed proportion between sister species (*S.*

osphromenoides and *S. selatanensis*, *L. aura* and *L. pulcher*) are consistent with an inversion in

the directionality of meiotic drive after or during the divergence of these species from their

common ancestor, such that either fusions or fissions were preferentially fixed in one species

but not the other. The karyotypes of *S. vaillanti* and *acrostoma*, which have a mix of biarmed

and uniarmed chromosomes, may be partway through a shift to a completely biarmed or

uniarmed karyotype. Additionally, changes in the arm number may have been caused by

pericentric inversions, which may also be subject to the force of meiotic drive (Molina et al.

2014). The large differences in karyotype between recently diverged species indicates that the

karyotypes were fixed extremely quickly, suggesting that meiotic drive would have to have been

extremely strong if it was driving these changes. There are several counterbalancing forces that

we would expect to weaken the strength of meiotic drive, including the relatively stronger force

of genetic drift, as well as the general trend of changes in chromosome count tending to be

slightly deleterious (King 1995).

Chromosomal rearrangements can have phenotypic impacts, particularly inversions, which can

suppress recombination by capturing multiple locally adapted alleles (Kirkpatrick and Barton

2006; Berg et al. 2016; da Silva et al. 2021). Additionally, low chromosome number has been

found to have correlations with phenotypic effects related to genome size (Gold 1979), including

specialization (defined as being highly phenotypically derived from their close evolutionary

relatives), tightening linkage groups, and occupying a narrower ecological niche (Gold 1979;

Hardie and Hebert 2004). The observed rearrangements and reductions in the spiral egg clade

may have played a role in acquiring highly specialized adaptations such as the ability of

Sphaerichtys and Luciocephalus to live in peat swamp forests and the associated blackwater

habitats, which are oligotrophic, sparsely inhabited, and highly acidic (pH < 4) (Polgar and

Jaafar 2018). By contrast, *P. ocellatus* and *C. nobilis* are not adapted to such harsh conditions

and are typically found in small muddy streams and pools. Additionally, it is possible that the

rapid genomic rearrangements observed in this group may have contributed to the observed

phenotypic differences in this subfamily, such as the highly derived morphology in

Luciocephalus.

The spiral egg clade presents an excellent opportunity to understand how these exceptionally

differentiated karyotypes arose and could give insight into larger patterns of chromosomal

evolution. Advanced cytogenetic techniques could help clarify which types of chromosomal

rearrangements occurred (ex. Ag-NOR staining, c-banding, FISH, etc.) as has been done in

other species in the family Osphronemidae (Grazyna et al. 2008; Pazza et al. 2009; Chaiyasan

et al. 2021; Supiwong et al. 2021), and measuring genome size of our study species would

allow testing for non-conservative mechanisms of chromosome evolution. To test the influence

of meiotic drive, work could be done to examine kinetochore protein levels during meiosis

(Chmátal et al. 2014), as well as the amount of minor satellite DNA repeats on the centromere ,

which have been associated with the action of meiotic drive in the western house mouse (Iwata-

Otsubo et al. 2017; Dudka and Lampson 2022). Other factors could be investigated that are

known to stimulate chromosomal rearrangements such as repetitive DNA content (King 1995;

Martinez et al. 2017) which was also implicated in the high incidence of chromosomal mutations

in *Nothobranchius* (Krysanov et al. 2023). There are other monophyletic clades in the family

Osphronemidae that have unusually differentiated chromosomes (Srisamoot et al. 2021),

suggesting that the underlying mechanism driving karyotypic change in the spiral egg clade may

be a shared ancestral trait and allowing for comparative genomic studies between the spiral egg

clade and closely related groups. Finally, it is also worth noting that most Osphronemidae

species have not been examined cytogenetically, hence karyotyping more species in the family

Osphronemidae could reveal more clades with high karyotype differentiation. Further attention

should be pair to this cytogenetically diverse group, as they could help resolve outstanding

evolutionary questions of chromosomal rearrangements and diversity.

AUTHOR CONTRIBUTIONS

Mobley: Conceptualization; Investigation; Methodology, Data analysis and curation; Writing –

original draft. **Anderson**: Conceptualization; Methodology; Data analysis and curation; Writing –

review and editing; Supervision.

ACKNOWLEDGEMENTS

This work was supported by National Science Foundation Grant #2147567. AP Anderson is

supported by the NSF Postdoctoral Research Fellowship in Biology Grant #2010841. B Mobley

was supported by a REP supplement to NSF grant #2147567.

DATA AVAILABILITY

Raw images of chromosome spreads and finalized karyotype images can be found here:

https://github.com/AndersonDrew/Gourami_Chromosome Contact authors for any additional

data/information.

CONFLICTS OF INTEREST STATEMENT

The authors declare that they have no competing interests.

References

Amores, A., J. Catchen, I. Nanda, W. Warren, R. Walter, M. Schartl, and J. H. Postlethwait. 2014. A RAD-tag genetic map for the platyfish (Xiphophorus maculatus) reveals mechanisms of karyotype evolution among teleost fish. Genetics 197:625–641. Oxford University Press.

Arai, R. 2011. Fish karyotypes: a check list. Springer Science & Business Media.

Berg, P. R., B. Star, C. Pampoulie, M. Sodeland, J. M. Barth, H. Knutsen, K. S. Jakobsen, and S. Jentoft. 2016. Three chromosomal rearrangements promote genomic divergence between migratory and stationary ecotypes of Atlantic cod. Sci. Rep. 6:23246. Nature Publishing Group UK London.

Blackmon, H., J. Justison, I. Mayrose, and E. E. Goldberg. 2019. Meiotic drive shapes rates of karyotype evolution in mammals. Evolution 73:511–523. Blackwell Publishing Inc Malden, USA.

Britz, R., M. Kokoscha, and R. Riehl. 1995. The anabantoid genera Ctenops, Luciocephalus, Parasphaerichthys, and Sphaerichthys (Teleostei: Perciformes) as a monophyletic group: evidence from egg surface structure and reproductive behaviour. Jpn. J. Ichthyol. 42:71–79.

Calton, M. S., and T. E. Denton. 1974. Chromosomes of the Chocolate Gourami: A Cytogenetic Anomaly. Science 185:618–619.

Canitz, J., F. Kirschbaum, and R. Tiedemann. 2016. Karyotype description of the African weakly electric fish Campylomormyrus compressirostris in the context of chromosome evolution in Osteoglossiformes. J. Physiol.-Paris 110:273–280. Elsevier.

CHAIYASAN, P., B. MINGKWAN, S. JANTARAT, C. SUWANNAPOOM, M. CIOFFI, T. LIEHR, S. TALUMPHAI, A. TANOMTONG, and W. Supiwong. 2021. Classical and molecular

cytogenetics of Belontia hasselti (Perciformes: Osphronemidae): Insights into the ZZ/ZW

sex chromosome system. Biodiversitas J. Biol. Divers. 22.

Chmátal, L., S. I. Gabriel, G. P. Mitsainas, J. Martínez-Vargas, J. Ventura, J. B. Searle, R. M.

Schultz, and M. A. Lampson. 2014. Centromere strength provides the cell biological

basis for meiotic drive and karyotype evolution in mice. Curr. Biol. 24:2295–2300.

Elsevier.

Christidis, L. 1990. Chordata 3. B. Aves. Schweizerbart'sche Verlagsbuchhandlung.

da Silva, S. A. S., P. A. de Lima-Filho, C. C. da Motta-Neto, G. W. W. F. da Costa, M. de B.

Cioffi, L. A. C. Bertollo, and W. F. Molina. 2021. High chromosomal evolutionary

dynamics in sleeper gobies (Eleotridae) and notes on disruptive biological factors in

Gobiiformes karyotypes (Osteichthyes, Teleostei). Mar. Life Sci. Technol. 1–10.

Springer.

de Villena, F. P.-M., and C. Sapienza. 2001. Female meiosis drives karyotypic evolution in

mammals. Genetics 159:1179–1189. Oxford University Press.

Deakin, J. E., and T. Ezaz. 2019. Understanding the evolution of reptile chromosomes through

applications of combined cytogenetics and genomics approaches. Cytogenet. Genome

Res. 157:7–20. S. Karger AG.

Dudka, D., and M. A. Lampson. 2022. Centromere drive: model systems and experimental

progress. Chromosome Res. 30:187–203. Springer.

Galetti, P. M., C. T. Aguilar, and W. F. Molina. 2000. An overview of marine fish cytogenetics.

Mar. Genet. 55–62. Springer.

Gallardo, M. H., C. A. González, and I. Cebrián. 2006. Molecular cytogenetics and

allotetraploidy in the red vizcacha rat, Tympanoctomys barrerae (Rodentia,

Octodontidae). Genomics 88:214–221. Elsevier.

Gold, J. R. 1979. Fish cytogenetics. Fish Physiol. 7–405. Academic Press.

Graphodatsky, A. S., P. L. Perelman, and S. J. O'Brien. 2020. Atlas of mammalian
chromosomes. No Title.

Grazyna, F.-S., D. Fopp-Bayat, M. Jankun, S. Krejszeff, and A. Mamcarz. 2008. Note on the
karyotype and NOR location of Siamese fighting fish Betta splendens (Perciformes,
Osphronemidae). Caryologia 61:349–353. Taylor & Francis.

Hardie, D. C., and P. D. Hebert. 2004. Genome-size evolution in fishes. Can. J. Fish. Aquat.
Sci. 61:1636–1646. NRC Research Press Ottawa, Canada.

Iwata-Otsubo, A., J. M. Dawicki-McKenna, T. Akera, S. J. Falk, L. Chmátal, K. Yang, B. A.
Sullivan, R. M. Schultz, M. A. Lampson, and B. E. Black. 2017. Expanded satellite
repeats amplify a discrete CENP-A nucleosome assembly site on chromosomes that
drive in female meiosis. Curr. Biol. 27:2365-2373. e8. Elsevier.

Jackson, B., R. K. Butlin, A. Navarro, and R. Faria. 2016. Chromosomal rearrangements and
speciation. Encycl. Evol. Biol. Elsevier Sci. Technol. Amst. Neth. 149–158.

King, M. 1995. Species evolution: the role of chromosome change. Cambridge University Press.

Kirkpatrick, M., and N. Barton. 2006. Chromosome inversions, local adaptation and speciation.
Genetics 173:419–434. Oxford University Press.

Kligerman, A. D., and S. E. Bloom. 1977. Rapid chromosome preparations from solid tissues of
fishes. J. Fish. Board Can. 34:266–269. NRC Research Press Ottawa, Canada.

Kretschmer, R., M. A. Ferguson-Smith, and E. H. C. De Oliveira. 2018. Karyotype evolution in
birds: From conventional staining to chromosome painting. Genes 9:181. MDPI.

Krysanov, E., and T. Demidova. 2018. Extensive karyotype variability of African fish genus
Nothobranchius (Cyprinodontiformes). Comp. Cytogenet. 12:387. Pensoft Publishers.

Krysanov, E., T. Demidova, and B. Nagy. 2016. Divergent karyotypes of the annual killifish
genus Nothobranchius (Cyprinodontiformes, Nothobranchiidae). Comp. Cytogenet.
10:439. Pensoft Publishers.

Krysanov, E. Y., B. Nagy, B. R. Watters, A. Sember, and S. A. Simanovsky. 2023. Karyotype

differentiation in the Nothobranchiusugandensis species group (Teleostei,

Cyprinodontiformes), seasonal fishes from the east African inland plateau, in the context

of phylogeny and biogeography. Comp. Cytogenet. 17:13. Pensoft Publishers.

Lebeda, I., P. Ráb, Z. Majtánová, and M. Flajšhans. 2020. Artificial whole genome duplication in

paleopolyploid sturgeons yields highest documented chromosome number in

vertebrates. Sci. Rep. 10:19705. Nature Publishing Group UK London.

Lehmann, R., A. Kovařík, K. Ocalewicz, L. Kirtiklis, A. Zuccolo, J. N. Tegner, J. Wanzenböck, L.

Bernatchez, D. K. Lamatsch, and R. Symonová. 2021. DNA transposon expansion is

associated with genome size increase in mudminnows. Genome Biol. Evol. 13:evab228.

Oxford University Press.

Levan, A., K. Fredga, and A. A. Sandberg. 1964. Nomenclature for centromeric position on

chromosomes.

Mank, J. E., and J. C. Avise. 2006. Phylogenetic conservation of chromosome numbers in

Actinopterygiian fishes. Genetica 127:321–327. Springer.

Martinez, P. A., U. P. Jacobina, R. V. Fernandes, C. Brito, C. Penone, T. F. Amado, C. R.

Fonseca, and C. J. Bidau. 2017. A comparative study on karyotypic diversification rate in

mammals. Heredity 118:366–373. Nature Publishing Group.

Martinez, P. A., J. P. Zurano, T. F. Amado, C. Penone, R. Betancur-R, C. J. Bidau, and U. P.

Jacobina. 2015. Chromosomal diversity in tropical reef fishes is related to body size and

depth range. Mol. Phylogenet. Evol. 93:1–4. Elsevier.

Mezzasalma, M., F. Andreone, G. Aprea, F. Glaw, G. Odierna, and F. M. Guarino. 2017. When

can chromosomes drive speciation? The peculiar case of the Malagasy tomato frogs

(genus Dyscophus). Zool. Anz. 268:41–46. Elsevier.

Molina, W. F., P. A. Martinez, L. A. C. Bertollo, and C. J. Bidau. 2014. Evidence for meiotic drive

as an explanation for karyotype changes in fishes. Mar. Genomics 15:29–34. Elsevier.

Nakatani, Y., H. Takeda, Y. Kohara, and S. Morishita. 2007. Reconstruction of the vertebrate

ancestral genome reveals dynamic genome reorganization in early vertebrates. Genome

Res. 17:1254–1265. Cold Spring Harbor Lab.

Olmo, E. 2005. Rate of chromosome changes and speciation in reptiles. Genetica 125:185–203.

Springer.

Pazza, R., K. F. Kavalco, P. R. Penteado, S. A. F. Kavalco, and L. F. de Almeida-Toledo. 2009.

Gene mapping of 18S and 5S rDNA genes in the karyotype of the three-spot gourami

Trichogaster trichopterus (Perciformes, Osphronemidae). Zebrafish 6:219–222. Mary

Ann Liebert, Inc. 140 Huguenot Street, 3rd Floor New Rochelle, NY 10801 USA.

Polgar, G., and Z. Jaafar. 2018. Endangered Forested Wetlands of Sundaland. Springer.

Rishi, S., P. K. Gill, and K. K. Rishi. 1997. Karyotypic studies in eight species of fishes

belonging to family Belontidae (Pisces, Perciformes). Chromosome Sci. 1:55–59. The

Society of Chromosome Research.

Romanenko, S. A., N. A. Serdyukova, P. L. Perelman, V. A. Trifonov, F. N. Golenishchev, N. S.

Bulatova, R. Stanyon, and A. S. Graphodatsky. 2018. Multiple intrasyntenic

rearrangements and rapid speciation in voles. Sci. Rep. 8:14980. Nature Publishing

Group UK London.

Rüber, L., R. Britz, and R. Zardoya. 2006. Molecular phylogenetics and evolutionary

diversification of labyrinth fishes (Perciformes: Anabantoidei). Syst. Biol. 55:374–397.

Shimabukuro-Dias, C. K., C. Oliveira, and F. Foresti. 2004. Cytogenetic analysis of five species

of the subfamily Corydoradinae (Teleostei: Siluriformes: Callichthyidae). Genet. Mol.

Biol. 27:549–554. SciELO Brasil.

Srisamoot, T., W. Supiwong, N. Sriwattanarothai, B. Panijpan, N. Srisamoot, and A. Tanomtong.

2021. Karyotype of four mouth-brooding Betta fishes in Thailand. Food Agric. Sci.

Technol. FAST 7:22–31.

Stanyon, R., and A. Graphodatsky. 2012. Evolutionary dynamics of mammalian karyotypes.

Cytogenet Genome Res 137:79–286.

Supiwong, W., P. Wongchantra, W. Thongnetr, B. Mingkwan, P. Chaiyasan, S.

Pinmongkhonkul, K. Pinthong, and A. Tanomtong. 2021. Comparative cytogenetic

analysis of fishes in the genus Trichopodus (Osphronemidae) in Thailand. Biodiversitas

J. Biol. Divers. 22.

Völker, M., P. Ráb, and H. Kullmann. 2005. Karyotype differentiation in Chromaphyosemion

killifishes (Cyprinodontiformes, Nothobranchiidae). I: Chromosome banding patterns of

C. alpha, C. kouamense and C. lugens. Genetica 125:33–41. Springer.

Völker, M., P. Rab, and H. Kullmann. 2008. Karyotype differentiation in Chromaphyosemion

killifishes (Cyprinodontiformes, Nothobranchiidae): patterns, mechanisms, and

evolutionary implications. Biol. J. Linn. Soc. 94:143–153. Oxford University Press.

Völker, M., R. Sonnenberg, P. Ráb, and H. Kullmann. 2006. Karyotype differentiation in

Chromaphyosemion killifishes (Cyprinodontiformes, Nothobranchiidae). II: Cytogenetic

and mitochondrial DNA analyses demonstrate karyotype differentiation and its

evolutionary direction in C. riggenbachi. Cytogenet. Genome Res. 115:70–83.

Völker, M., R. Sonnenberg, P. Ráb, and H. Kullmann. 2007. Karyotype differentiation in

Chromaphyosemion killifishes (Cyprinodontiformes, Nothobranchiidae). III: Extensive

karyotypic variability associated with low mitochondrial haplotype differentiation in C.

bivittatum. Cytogenet. Genome Res. 116:116–126. S. Karger AG.

Wurster, D. H., and K. Benirschke. 1970. Indian Muntjac, Muntiacus muntjak: A Deer with a Low

Diploid Chromosome Number. Science 168:1364–1366. American Association for the

Advancement of Science.

Zworykin, D., J. Müller, H. Grundmeijer, and E. Pavlov. 2024. Paternal mouthbrooding in the

chocolate gourami Sphaerichthys osphromenoides (Osphronemidae). Environ. Biol.

Fishes 107:381–389.