Rapid Changes in Chromosome Counts in Fishes from the Spiral Egg Clade within the Gourami Family (Osphronemidae).

Brendan Mobley and Andrew P. Anderson

Biology Department, Reed College, Portland, Oregon 97202, USA

Abstract

Identifying clades with numerous and noticeable changes in chromosome counts is an important step in unraveling the evolutionary mechanisms that shape cytogenetics. Here, we describe chromosome counts in a group of teleost fishes delimited by their egg structure and with a species with a known low chromosome count within the labyrinthine clade (Osphronemidae). We sampled seven of nine known species within the spiral egg clade, reporting novel chromosome counts for five species while confirming two others. Overall, we find a rapid loss of chromosomes during the clade's emergence and high variability in both chromosome count and centricity suggesting multiple fusion/fissions across evolutionary time. Lastly, we offer some possible explanations for these changes based on current and ongoing empirical and theoretical research. These data provide important information in cataloguing rapid chromosomal shifts and we call attention to this group of fishes for further study in chromosomal and genomic evolution due to their karyotypic variability.

Keywords

Cytogenetics, evolution, genomics, karyotype, fish

Introduction

Variability of chromosome numbers across vertebrates and the evolutionary mechanisms that

create it is an active point of inquiry for evolutionary biologists (Martinez et al. 2015). Finding

clades with high variance or rapid changes presents more data points in resolving the various

hypotheses on how chromosome number changes over evolutionary time. Some patterns can

include changes due to long stretches of repeat content which can create opportunities for

mismatched recombination (Amores et al. 2014) such as found in mammals with a range of

chromosome counts from 2N=6 in the female muntjac deer (*Muntiacus muntjac*) (Wurster and

Benirschke 1970; Graphodatsky et al. 2020) to 2N=102 in the plains viscacha rat

(*Tympanoctomys barrerae*) (Gallardo et al. 2006; Stanyon and Graphodatsky 2012; Lebeda et

al. 2020). Other patterns can involve microchromosomes as typically seen in birds which have a

range from 2N=40 in *Ceratogymna bucinator* to 2N=136-142 in *Corythaixoides concolor*

(Christidis 1990; Kretschmer et al. 2018). Conversely, reptiles are karyologically heterogeneous

and distinct evolutionary trends are exhibited by different orders and suborders despite having

highly conserved karyotypes (Stanyon and Graphodatsky 2012) ranging from 24–70

chromosomes (Olmo 2005). Fishes have been found to have notable karyotype evolution

patterns, particularly with regard to extreme chromosome counts, rapid cytogenetic changes, or

both (e.g. *Nothobranchius* 2N=16–50:Krysanov et al. 2023 and *Corydoras* 2N=40–134:

Shimabukuro-Dias et al. 2004) and span the widest range of all vertebrates, ranging from

2N=12 in the marine species *Gonostoma bathyphilum* to 2N=446 in the freshwater species

Ptychobarbus dipogon (Lebeda et al. 2020). Paradoxically, there is a strong trend of conserved

karyotypes (Galetti et al. 2000; Mank and Avise 2006; Nakatani et al. 2007) with over half of all

karyotyped fish species having diploid chromosome counts of 48 or 50 (Mank and Avise 2006;

Arai 2011), which has changed little from the proposed karyotype for the ancestor of all teleost

fish (2N=52—Nakatani et al. 2007). With such diverse patterns across vertebrates, finding

clades with detectable changes in chromosome counts is valuable to understanding

chromosome evolution as whole.

An intriguing group of fishes with potential chromosomal variance is within the family

Osphronemidae, commonly called gouramis. The chocolate gourami, *Sphaerichthys*

osphromenoides, has the lowest recorded chromosome count among freshwater fishes

(Lehmann et al. 2021) with 2n=16 (Arai 2011). A species in the neighboring genus, the

pikehead gourami, *Luciocephalus pulcher*, was reported to have 2n=20 (Arai 2011).

Chromosome counts this low are exceedingly rare in fishes, as there are only thirteen fish

species with a diploid chromosome count lower than 2N=22 (Lehmann et al. 2021).

Furthermore, these counts are highly derived from the other Osphronemidae species, which

generally have 2N values between 46 and 48.

Both *S. osphromenoides* and *L. pulcher* are members of the "spiral egg" clade, a monophyletic

group within the family Osphronemidae that includes the genera *Sphaerichthys*, *Luciocephalus*,

Parasphaerichthys, and *Ctenops*. The monophyly was proposed based on the unique

morphology of their eggs, which are covered in projections arranged in a spiral pattern, and later

confirmed and refined with molecular evidence (Britz et al. 1995; Rüber et al. 2006). Another

differentiating feature of the spiral egg clade is an angular jaw shape, which is taken to the

extreme in the highly derived pike-like morphology of the piscivorous genus *Luciocephalus*. The

spiral egg clade is also notable for having the only species, *S. osphromenoides* and *S.*

selatanensis, in Osphronemidae with female broodcare via mouthbrooding compared to the

overwhelmingly male mouthbrooders or bubble nesters (Rüber et al. 2006) in the family,

although recent evidence has called into question the sex of caring parent in *S.*

osphromenoides (Zworykin et al. 2024). Chromosomes of the spiral egg clade remain largely

uninvestigated; besides *S. osphromenoides* and *L. pulcher*, only one other species has been

studied cytogenetically (*Ctenops nobilis*, 2N=44: Rishi et al. 1997). Given the low chromosome

counts of *S. osphromenoides* and *L. pulcher* and the large decrease relative to the wider family,

we aim to characterize the karyotypes of additional members within the spiral egg clade. With

this information we will describe the karyotypic diversity and evolutionary history for this

remarkable group of fishes, thereby adding an extraordinary example to the chromosome count

diversity in fishes specifically and animals in general.

Methods

Fishes were sourced from the aquarium trade (Wet Spot Tropical Fish, Portland, Oregon, USA;

Nationwide Aquatics, Tinley Park, Illinois, USA; Aqua Imports, Boulder, Colorado, USA), then

held in species-specific tanks (110 liters) on a shared flow-through system (pH 7.0, GH 30 ppm,

KH 40 ppm) with a 12/12 hour light/dark cycle with 30 minutes of dim light to simulate dawn and

dusk. Specimens were housed for a minimum of one week before sampling to ensure good

health for optimal cell proliferation.

Chromosome preparations were made following Kligerman and Bloom (1977) with the indicated

modifications. Specimens were incubated in 0.005% colchicine solution for 6-7 hours, then

euthanized and dissected to remove gill arches. Sex determination was conducted by gross

examination of gonads with pictures taken throughout. Dissected specimens were stored in -80

°C for future molecular analyses. Gill arches were incubated in 0.4% KCl solution for 20-30

minutes, then fixed in two changes of 3:1 ethanol:acetic acid fixative for at least 30 minutes

each, followed by an overnight fixation period at 4 °C. To prepare slides, tissue was dissociated

into suspension by mincing in 50% acetic acid, then dropped onto a slide warmed to 30-40 °C

and air dried. Slides were examined under phase contrast microscopy to ensure quality, then

aged for at least one day at room temperature before being stained for 10 minutes in 10%

Giemsa in pH 6.8 phosphate buffer (Gibco™ Gurr Buffer Tablets) and air-dried.

Chromosomes were examined under a Nikon Eclipse Ti-E microscope driven by Nikon NIS

Elements AR software, then photographed with an oil immersion objective at 100x magnification

and green color filtering using a Hamatsu ORCA-Flash4.0 camera. Digital images were

optimized, then homologous chromosomes were paired by size and morphology and arranged

in decreasing order of size using ImageJ v1.52v and Adobe Photoshop 24.3.0. At least 20

complete metaphase spreads were photographed from each specimen. Chromosomes were

classified as metacentric (m), submetacentric (sm), subtelocentric (st), or acrocentric (a)

according to their arm ratios (Levan et al. 1964). Chromosome arm number (Fundamental

Number, FN) was calculated by considering metacentric and submetacentric chromosomes as

biarmed and subtelocentric and telocentric chromosomes as uniarmed.

Results

We describe karyotypes for the first time in six species (Fig 1): *S. selatanensis*, *S. vaillanti*, *S.*

acrostoma, *L. aura*, and *P. ocellatus*. We also confirmed the karyotypes of an additional two

species (*S. osphromenoides*, *L. pulcher*) which matched those established in the literature

(Calton and Denton 1974; Arai 2011). All species in the genus *Sphaerichthys* had different

chromosome counts, with 2N ranging from 16–28 (Table 1). The number of chromosome arms

(fundamental number, FN) showed less variation, with a range of 30–38. Notably, the sister

species *S. osphromenoides* and *S. selatanensis* had a primarily biarmed and primarily uniarmed

karyotype respectively, resulting in a shared FN despite a 2N difference of 12. The other two

sister species in the genus (*S. acrostoma* and *S. vaillanti*) had karyotypes that were a mix of

biarmed and uniarmed chromosomes and had different values for both 2N and FN. We

confirmed that *L. pulcher* had an entirely uniarmed karyotype of 20 acrocentric chromosomes.

Luciocephalus aura had a primarily biarmed karyotype with six fewer chromosomes and eight

more chromosome arms. *Parasphaerichthys ocellatus* had a primarily biarmed karyotype with

higher 2N and FN than any *Sphaerichthys* or *Luciocephalus* species, though the 2N, but not the

FN, was lower than reported for *C. nobilis* in the literature (Arai 2011).

The two *S. selatanensis* that we sampled had different karyotypes (Table 1). One had 2N=28

with a karyotype formula 2st+26a, but the other had a single unpaired metacentric and one less

large acrocentric pair, giving 2N=27 and a karyotype formula of 1m+2st+24a. The unpaired

metacentric was approximately twice the size of the largest acrocentric chromosomes in the

karyotype. We cannot say whether this is a sex chromosome because we could not confidently

determine the sex of either individual.

Table 1: Summary of sampling of chromosome spreads. Data provided include the species of the individuals
sampled, the number of spreads collected and analyzed for that individual and descriptive statistics of that data set.

Genus	species	ID	# Spreads	Mode 2N	Median 2N	Mean 2N	Lowest 2N	Highest 2N
Sphaerichthys	*osphromenoides*	So1	47	16	16	15.34	12	17
Sphaerichthys	*selatanensis*	Ss1	39	27	27	25.54	16	28
Sphaerichthys	*selatanensis*	Ss2	37	28	27	26.57	18	28
Sphaerichthys	*vaillanti*	Sv1	81	20	20	18.91	12	20
Sphaerichthys	*vaillanti*	Sv2	50	20	20	19.68	14	23
Sphaerichthys	*vaillanti*	Sv8	35	20	20	19.14	12	20
Sphaerichthys	*acrostoma*	Sa3	44	24	24	24.00	22	26
Luciocephalus	*pulcher*	Lp4	40	20	20	19.43	16	20
Luciocephalus	*aura*	La1	88	14	14	13.73	12	16
Parasphaerichhtys	*ocellatus*	Po2	39	34	34	33.67	30	36
Parasphaerichhtys	*ocellatus*	Po4	86	34	34	33.26	26	36
Parasphaerichhtys	*ocellatus*	Po5	42	34	34	32.14	30	36

Species	2N	FN	Karyotype
Sphaerichthys osphromenoides	16	30	10m + 4sm + 2a
Sphaerichthys selatanensis	28	30	2st + 26a
Sphaerichthys vaillanti	20	36	4m + 4sm + 10st + 4a
Sphaerichthys acrostoma	24	38	4m + 10sm + 10a
Luciocephalus pulcher	20	20	20a
Luciocephalus aura	14	28	8m + 2sm + 4st
Parasphaerichthys ocellatus	34	64	6m + 4sm + 22st + 2a
Ctenops nobilis	44	60	8m + 8sm + 28a
Trichopodus leerii	46	46	46a
Osphronemus goramy	48	50	2sm + 46a
Helostoma temminckii	48	48	48a

5 MYA

Discussion

We found that the genera *Sphaerichthys*, *Luciocephalus*, and *Parasphaerichthys* have low chromosome counts (2N≤34) with high intra-genus variation in both 2N and FN. These trends are not observed in the karyotypes of the broader family Osphronemidae, which mostly has a karyotype similar (2N=52) to the hypothesized ancestral state for all teleost fishes (Nakatani et al. 2007), thereby suggesting that rapid karyotype evolution occurred since the divergence of the spiral egg clade about 25 million years ago (Rüber et al. 2006). Karyotype evolution of this rate and magnitude has not been reported in teleost fishes. The drastic differences between karyotypes within the *Sphaerichthys* and *Luciocephalus* genera suggests that karyotype evolution may have played a role in speciation process by creating post-zygotic isolation (Canitz et al. 2016; Jackson et al. 2016; Mezzasalma et al. 2017; Romanenko et al. 2018), but it is also possible that the observed karyotype diversity happened alongside the speciation process instead of driving it (Krysanov et al. 2023). Mechanisms of chromosomal rearrangements can include the influence of genetic drift or other forms of neutral selection, extremely strong meiotic drive, the evolution of a trait that stimulates chromosome evolution, or a combination of these factors.

Genetic drift has been proposed to be the driving force behind the fixation of highly differentiated karyotypes in other clades of freshwater fishes, including the annual killifishes in the genera *Nothobranchius* (2N=16-50) (Krysanov et al. 2016, 2023; Krysanov and Demidova 2018) and *Aphyosemion* (2N=20-40) (Völker et al. 2005, 2007, 2008). These species tend to live in small, biogeographically isolated populations and frequently experience genetic bottlenecks and founder effects due to the ephemeral nature of their habitats (Völker et al. 2006;

Krysanov and Demidova 2018; Krysanov et al. 2023), which may have sped up the

accumulation of both intra-and inter-chromosomal mutations, with centric fusions being

responsible for most decreases in chromosome count (Völker et al. 2005, 2008; Krysanov et al.

2016). *Sphaerichthys* and *Luciocephalus* species have very limited geographical ranges and,

given that most of them are threatened or endangered according to the IUCN Red List, these

populations are likely small; however, it is difficult to precisely estimate the strength of genetic

drift on the evolution of the karyotypes in our study group because there is little information

about their distribution and population structure.

An alternative and complementary explanation to genetic drift is meiotic drive. Species evolving

under the influence of strong meiotic drive will tend to reach karyotypes of predominantly

biarmed or uniarmed chromosomes (de Villena and Sapienza 2001; Molina et al. 2014). This

has been observed in fishes and mammals, with the note that groups with high rates of

mismatched karyotypes tended to have higher rates of chromosome evolution (Blackmon et al.

2019). The decrease in chromosome number in our study group relative to the rest of the family

may have been caused by meiotic drive with a polarity for biarmed chromosomes that rapidly

fixed fusion mutations. Additionally, the inversions in biarmed proportion between sister species

(*S. osphromenoides* and *S. selatanensis*, *L. aura* and *L. pulcher*) are consistent with an

inversion in the polarity of meiotic drive after or during the divergence of these species from

their common ancestor, such that either fusions or fissions were preferentially fixed in one

species but not the other. The karyotypes of *S. vaillanti* and *acrostoma*, which have a

combination of biarmed and uniarmed chromosomes, may be partway through a shift to a

completely biarmed or uniarmed karyotype. The changes in FN may have been caused by

pericentric inversions, which have been found to be the most prominent mechanism of

karyotype evolution in fish , and may also be subject to the force of meiotic drive (Molina et al.

2014). The large differences in karyotype between sister species indicates that the karyotypes

were fixed extremely quickly, indicating meiotic drive would have to have been extremely strong

if it was sole mechanism of these differences. There are several counterbalancing forces that

we would expect to weaken the strength of meiotic drive, including the relatively stronger force

of genetic drift, as well as the general trend of changes in chromosome count tending to be

slightly deleterious .

Chromosomal rearrangements can have phenotypic impacts, most notably chromosomal

inversions, which can suppress recombination by capturing multiple locally adapted alleles

(Kirkpatrick and Barton 2006; Berg et al. 2016; da Silva et al. 2021). Additionally, low

chromosome number has been found to have correlations with phenotypic effects related to

genome size (Gold 1979), including specialization (defined as being highly phenotypically

derived from their close evolutionary relatives), tightening linkage groups, and occupying a

narrower ecological niche (Gold 1979; Hardie and Hebert 2004). The observed rearrangements

and reductions in the spiral egg clade, may have played a role in acquiring highly specialized

adaptations such as the ability *Sphaerichtys and Luciocephalus* to live in peat swamp forests

and the associated blackwater habitats, which are oligotrophic and highly acidic (pH < 4)

(Polgar and Jaafar 2018). By contrast, *P. ocellatus* and *C. nobilis* are not adapted to such harsh

conditions and are typically found in small muddy streams and pools. Additionally, it is possible

that the rapid genomic rearrangements observed in this group may have contributed to the

observed phenotypic differences in this subfamily, such as the highly derived morphology in

Luciocephalus.

The spiral egg clade presents an excellent opportunity to understand how these exceptionally

differentiated karyotypes arose and how that may be applied to larger patterns of chromosomal

evolution. Advanced cytogenetic techniques could help clarify which types of chromosomal

rearrangements occurred (ex. Ag-NOR staining, c-banding, FISH, etc.) as has been done in

other species in the family Osphronemidae (Grazyna et al. 2008; Pazza et al. 2009; Chaiyasan

et al. 2021; Supiwong et al. 2021), and measuring genome size of our study species would

allow testing for non-conservative mechanisms of chromosome evolution. To test the influence

of meiotic drive, work could be done by examining kinetochore protein levels during meiosis

(Chmátal et al. 2014), as well as the number of satellite DNA repeats on the centromere , which

have been associated with the action of meiotic drive in the western house mouse. Other factors

could be investigated that are known to stimulate chromosomal rearrangements such as

repetitive DNA content (King 1995; Martinez et al. 2017) which was also implicated in the high

incidence of chromosomal mutations in *Nothobranchius* (Krysanov et al. 2023). There are other

monophyletic clades in the family Osphronemidae that have unusually differentiated

chromosomes (Srisamoot et al. 2021), suggesting that the underlying mechanism driving

karyotypic change in the spiral egg clade may be a shared ancestral trait and allowing for

comparative genomic studies between the spiral egg clade and closely related groups. Finally, it

is also worth noting that most Osphronemidae species have not been examined cytogenetically.

Karyotyping more species in the family Osphronemidae could reveal more clades with high

karyotype differentiation. Further attention should be pair to this cytogenetically diverse group,

as they could help resolve outstanding evolutionary questions of chromosomal rearrangements

and diversity.

AUTHOR CONTRIBUTIONS

Mobley: Conceptualization; Investigation; Methodology, Data analysis and curation; Writing –

original draft. **Anderson**: Conceptualization; Methodology; Supervision; Writing – review and

editing; Data analysis and curation.

ACKNOWLEDGEMENTS

This work was supported by National Science Foundation Grant #2147567. AP Anderson is

supported by the NSF Postdoctoral Research Fellowship in Biology Grant #2010841. B Mobley

was supported by a REP supplement to NSF grant #2147567.

DATA AVAILABILITY

Raw images of chromosome spreads and finalized karyotype images can be found here:

https://github.com/AndersonDrew/Gourami_Chromosome Contact authors for any additional

data/information.

CONFLICTS OF INTEREST STATEMENT

The authors declare that they have no competing interests.

References

Amores, A., J. Catchen, I. Nanda, W. Warren, R. Walter, M. Schartl, and J. H. Postlethwait. 2014. A RAD-tag genetic map for the platyfish (Xiphophorus maculatus) reveals mechanisms of karyotype evolution among teleost fish. Genetics 197:625–641. Oxford University Press.

Arai, R. 2011. Fish karyotypes: a check list. Springer Science & Business Media.

Berg, P. R., B. Star, C. Pampoulie, M. Sodeland, J. M. Barth, H. Knutsen, K. S. Jakobsen, and S. Jentoft. 2016. Three chromosomal rearrangements promote genomic divergence between migratory and stationary ecotypes of Atlantic cod. Sci. Rep. 6:23246. Nature Publishing Group UK London.

Blackmon, H., J. Justison, I. Mayrose, and E. E. Goldberg. 2019. Meiotic drive shapes rates of karyotype evolution in mammals. Evolution 73:511–523. Blackwell Publishing Inc Malden, USA.

Britz, R., M. Kokoscha, and R. Riehl. 1995. The anabantoid genera Ctenops, Luciocephalus, Parasphaerichthys, and Sphaerichthys (Teleostei: Perciformes) as a monophyletic group: evidence from egg surface structure and reproductive behaviour. Jpn. J. Ichthyol. 42:71–79.

Calton, M. S., and T. E. Denton. 1974. Chromosomes of the Chocolate Gourami: A Cytogenetic Anomaly. Science 185:618–619.

Canitz, J., F. Kirschbaum, and R. Tiedemann. 2016. Karyotype description of the African weakly electric fish Campylomormyrus compressirostris in the context of chromosome evolution in Osteoglossiformes. J. Physiol.-Paris 110:273–280. Elsevier.

CHAIYASAN, P., B. MINGKWAN, S. JANTARAT, C. SUWANNAPOOM, M. CIOFFI, T. LIEHR, S. TALUMPHAI, A. TANOMTONG, and W. Supiwong. 2021. Classical and molecular

280   cytogenetics of Belontia hasselti (Perciformes: Osphronemidae): Insights into the ZZ/ZW

281   sex chromosome system. Biodiversitas J. Biol. Divers. 22.

282 Chmátal, L., S. I. Gabriel, G. P. Mitsainas, J. Martínez-Vargas, J. Ventura, J. B. Searle, R. M.

283   Schultz, and M. A. Lampson. 2014. Centromere strength provides the cell biological

284   basis for meiotic drive and karyotype evolution in mice. Curr. Biol. 24:2295–2300.

285   Elsevier.

286 Christidis, L. 1990. Chordata 3. B. Aves. Schweizerbart'sche Verlagsbuchhandlung.

287 da Silva, S. A. S., P. A. de Lima-Filho, C. C. da Motta-Neto, G. W. W. F. da Costa, M. de B.

288   Cioffi, L. A. C. Bertollo, and W. F. Molina. 2021. High chromosomal evolutionary

289   dynamics in sleeper gobies (Eleotridae) and notes on disruptive biological factors in

290   Gobiiformes karyotypes (Osteichthyes, Teleostei). Mar. Life Sci. Technol. 1–10.

291   Springer.

292 de Villena, F. P.-M., and C. Sapienza. 2001. Female meiosis drives karyotypic evolution in

293   mammals. Genetics 159:1179–1189. Oxford University Press.

294 Galetti, P. M., C. T. Aguilar, and W. F. Molina. 2000. An overview of marine fish cytogenetics.

295   Mar. Genet. 55–62. Springer.

296 Gallardo, M. H., C. A. González, and I. Cebrián. 2006. Molecular cytogenetics and

297   allotetraploidy in the red vizcacha rat, Tympanoctomys barrerae (Rodentia,

298   Octodontidae). Genomics 88:214–221. Elsevier.

299 Gold, J. R. 1979. Fish cytogenetics. Fish Physiol. 7–405. Academic Press.

300 Graphodatsky, A. S., P. L. Perelman, and S. J. O'Brien. 2020. Atlas of mammalian

301   chromosomes. No Title.

302 Grazyna, F.-S., D. Fopp-Bayat, M. Jankun, S. Krejszeff, and A. Mamcarz. 2008. Note on the

303   karyotype and NOR location of Siamese fighting fish Betta splendens (Perciformes,

304   Osphronemidae). Caryologia 61:349–353. Taylor & Francis.

Hardie, D. C., and P. D. Hebert. 2004. Genome-size evolution in fishes. Can. J. Fish. Aquat.

Sci. 61:1636–1646. NRC Research Press Ottawa, Canada.

Jackson, B., R. K. Butlin, A. Navarro, and R. Faria. 2016. Chromosomal rearrangements and

speciation. Encycl. Evol. Biol. Elsevier Sci. Technol. Amst. Neth. 149–158.

King, M. 1995. Species evolution: the role of chromosome change. Cambridge University Press.

Kirkpatrick, M., and N. Barton. 2006. Chromosome inversions, local adaptation and speciation.

Genetics 173:419–434. Oxford University Press.

Kligerman, A. D., and S. E. Bloom. 1977. Rapid chromosome preparations from solid tissues of

fishes. J. Fish. Board Can. 34:266–269. NRC Research Press Ottawa, Canada.

Kretschmer, R., M. A. Ferguson-Smith, and E. H. C. De Oliveira. 2018. Karyotype evolution in

birds: From conventional staining to chromosome painting. Genes 9:181. MDPI.

Krysanov, E., and T. Demidova. 2018. Extensive karyotype variability of African fish genus

Nothobranchius (Cyprinodontiformes). Comp. Cytogenet. 12:387. Pensoft Publishers.

Krysanov, E., T. Demidova, and B. Nagy. 2016. Divergent karyotypes of the annual killifish

genus Nothobranchius (Cyprinodontiformes, Nothobranchiidae). Comp. Cytogenet.

10:439. Pensoft Publishers.

Krysanov, E. Y., B. Nagy, B. R. Watters, A. Sember, and S. A. Simanovsky. 2023. Karyotype

differentiation in the Nothobranchiusugandensis species group (Teleostei,

Cyprinodontiformes), seasonal fishes from the east African inland plateau, in the context

of phylogeny and biogeography. Comp. Cytogenet. 17:13. Pensoft Publishers.

Lebeda, I., P. Ráb, Z. Majtánová, and M. Flajšhans. 2020. Artificial whole genome duplication in

paleopolyploid sturgeons yields highest documented chromosome number in

vertebrates. Sci. Rep. 10:19705. Nature Publishing Group UK London.

Lehmann, R., A. Kovařík, K. Ocalewicz, L. Kirtiklis, A. Zuccolo, J. N. Tegner, J. Wanzenböck, L.

Bernatchez, D. K. Lamatsch, and R. Symonová. 2021. DNA transposon expansion is

associated with genome size increase in mudminnows. Genome Biol. Evol. 13:evab228.

Oxford University Press.

Levan, A., K. Fredga, and A. A. Sandberg. 1964. Nomenclature for centromeric position on

chromosomes.

Mank, J. E., and J. C. Avise. 2006. Phylogenetic conservation of chromosome numbers in

Actinopterygiian fishes. Genetica 127:321–327. Springer.

Martinez, P. A., U. P. Jacobina, R. V. Fernandes, C. Brito, C. Penone, T. F. Amado, C. R.

Fonseca, and C. J. Bidau. 2017. A comparative study on karyotypic diversification rate in

mammals. Heredity 118:366–373. Nature Publishing Group.

Martinez, P. A., J. P. Zurano, T. F. Amado, C. Penone, R. Betancur-R, C. J. Bidau, and U. P.

Jacobina. 2015. Chromosomal diversity in tropical reef fishes is related to body size and

depth range. Mol. Phylogenet. Evol. 93:1–4. Elsevier.

Mezzasalma, M., F. Andreone, G. Aprea, F. Glaw, G. Odierna, and F. M. Guarino. 2017. When

can chromosomes drive speciation? The peculiar case of the Malagasy tomato frogs

(genus Dyscophus). Zool. Anz. 268:41–46. Elsevier.

Molina, W. F., P. A. Martinez, L. A. C. Bertollo, and C. J. Bidau. 2014. Evidence for meiotic drive

as an explanation for karyotype changes in fishes. Mar. Genomics 15:29–34. Elsevier.

Nakatani, Y., H. Takeda, Y. Kohara, and S. Morishita. 2007. Reconstruction of the vertebrate

ancestral genome reveals dynamic genome reorganization in early vertebrates. Genome

Res. 17:1254–1265. Cold Spring Harbor Lab.

Olmo, E. 2005. Rate of chromosome changes and speciation in reptiles. Genetica 125:185–203.

Springer.

Pazza, R., K. F. Kavalco, P. R. Penteado, S. A. F. Kavalco, and L. F. de Almeida-Toledo. 2009.

Gene mapping of 18S and 5S rDNA genes in the karyotype of the three-spot gourami

Trichogaster trichopterus (Perciformes, Osphronemidae). Zebrafish 6:219–222. Mary

Ann Liebert, Inc. 140 Huguenot Street, 3rd Floor New Rochelle, NY 10801 USA.

Polgar, G., and Z. Jaafar. 2018. Endangered Forested Wetlands of Sundaland. Springer.

Rishi, S., P. K. Gill, and K. K. Rishi. 1997. Karyotypic studies in eight species of fishes

belonging to family Belontidae (Pisces, Perciformes). Chromosome Sci. 1:55–59. The

Society of Chromosome Research.

Romanenko, S. A., N. A. Serdyukova, P. L. Perelman, V. A. Trifonov, F. N. Golenishchev, N. S.

Bulatova, R. Stanyon, and A. S. Graphodatsky. 2018. Multiple intrasyntenic

rearrangements and rapid speciation in voles. Sci. Rep. 8:14980. Nature Publishing

Group UK London.

Rüber, L., R. Britz, and R. Zardoya. 2006. Molecular phylogenetics and evolutionary

diversification of labyrinth fishes (Perciformes: Anabantoidei). Syst. Biol. 55:374–397.

Shimabukuro-Dias, C. K., C. Oliveira, and F. Foresti. 2004. Cytogenetic analysis of five species

of the subfamily Corydoradinae (Teleostei: Siluriformes: Callichthyidae). Genet. Mol.

Biol. 27:549–554. SciELO Brasil.

Srisamoot, T., W. Supiwong, N. Sriwattanarothai, B. Panijpan, N. Srisamoot, and A. Tanomtong.

2021. Karyotype of four mouth-brooding Betta fishes in Thailand. Food Agric. Sci.

Technol. FAST 7:22–31.

Stanyon, R., and A. Graphodatsky. 2012. Evolutionary dynamics of mammalian karyotypes.

Cytogenet Genome Res 137:79–286.

Supiwong, W., P. Wongchantra, W. Thongnetr, B. Mingkwan, P. Chaiyasan, S.

Pinmongkhonkul, K. Pinthong, and A. Tanomtong. 2021. Comparative cytogenetic

analysis of fishes in the genus Trichopodus (Osphronemidae) in Thailand. Biodiversitas

J. Biol. Divers. 22.

Völker, M., P. Ráb, and H. Kullmann. 2005. Karyotype differentiation in Chromaphyosemion

killifishes (Cyprinodontiformes, Nothobranchiidae). I: Chromosome banding patterns of

C. alpha, C. kouamense and C. lugens. Genetica 125:33–41. Springer.

Völker, M., P. Rab, and H. Kullmann. 2008. Karyotype differentiation in Chromaphyosemion

killifishes (Cyprinodontiformes, Nothobranchiidae): patterns, mechanisms, and

evolutionary implications. Biol. J. Linn. Soc. 94:143–153. Oxford University Press.

Völker, M., R. Sonnenberg, P. Ráb, and H. Kullmann. 2006. Karyotype differentiation in

Chromaphyosemion killifishes (Cyprinodontiformes, Nothobranchiidae). II: Cytogenetic

and mitochondrial DNA analyses demonstrate karyotype differentiation and its

evolutionary direction in C. riggenbachi. Cytogenet. Genome Res. 115:70–83.

Völker, M., R. Sonnenberg, P. Ráb, and H. Kullmann. 2007. Karyotype differentiation in

Chromaphyosemion killifishes (Cyprinodontiformes, Nothobranchiidae). III: Extensive

karyotypic variability associated with low mitochondrial haplotype differentiation in C.

bivittatum. Cytogenet. Genome Res. 116:116–126. S. Karger AG.

Wurster, D. H., and K. Benirschke. 1970. Indian Muntjac, Muntiacus muntjak: A Deer with a Low

Diploid Chromosome Number. Science 168:1364–1366. American Association for the

Advancement of Science.

Zworykin, D., J. Müller, H. Grundmeijer, and E. Pavlov. 2024. Paternal mouthbrooding in the

chocolate gourami Sphaerichthys osphromenoides (Osphronemidae). Environ. Biol.

Fishes 107:381–389.

398