ORIGINAL ARTICLE

The subcortical correlates of self-reported sleep quality

Martin M. Monti [1,2]*

[1]Department of Psychology, University of California Los Angeles, 502 Portola Plaza, Los Angeles, 90095, CA, USA and [2]Brain Injury Research Center (BIRC), Department of Neurosurgery, University of California Los Angeles, 300 Stein Plaza Driveway, Los Angeles, 90095, CA, USA

*Corresponding author: monti@ucla.edu

Abstract

Study objectives: To assess the association between self-reported measures of sleep quality and cortical and subcortical local morphometry.

Methods: Sleep quality, operationalized with the Pittsburgh Sleep Quality Index (PSQI), and neuroanatomical data from the full release of the young adult Human Connectome Project dataset were analyzed (N=1,112; 46% female; mean age: 28.8 years old). Local cortical and subcortical morphometry was measured with subject-specific segmentations resulting in voxelwise gray matter difference (i.e., voxel based morephometry) measurements for cortex and local shape measurements for subcortical regions. Associations between the total score of PSQI, two statistical groupings of its subcomponents (obtained with a principal component analysis), and their interaction with demographic (i.e., sex, age, handedness, years of education) and biometric (i.e., BMI) variables were assessed using a general linear model and a nonparametric permutation approach.

Results: Sleep quality-related variance was significantly associated with subcortical morphometry, particularly in the bilateral caudate, putamen, and left pallidum, where smaller shape measures correlated with worse sleep quality. Notably, these associations were independent of demographic and biometric factors. In contrast, cortical morphometry, along with additional subcortical sites, showed no direct associations with sleep quality but demonstrated interactions with demographic and biometric variables.

Conclusions: This study reveals a specific link between self-reported sleep quality and subcortical morphometry, particularly within the striatum and pallidum, reinforcing the role of these regions in sleep regulation. These findings underscore the importance of considering subcortical morphology in sleep research and highlight potential neuromodulatory targets for sleep-related interventions.

Key words: Sleep and Brain, Neuroanatomy, Neuroimaging, Caudate, Putamen, Pallidum

Statement of Significance

This study provides evidence for the role of subcortical structures in sleep regulation, demonstrating that poorer self-reported sleep quality is associated with reduced shape measures in the bilateral caudate, putamen, and left pallidum. These associations were independent of demographic and biometric factors, distinguishing subcortical morphometry as a unique correlate of sleep quality. In contrast, cortical morphology exhibited no direct relationship with sleep quality but showed interactions with factors such as sex, BMI, and age, highlighting the complexity of brain-sleep relationships. These findings contribute to a growing body of literature emphasizing the basal ganglia's role in non-motor functions, including sleep regulation, and suggest potential neuromodulatory targets for sleep-related interventions, such as transcranial focused ultrasound or deep brain stimulation.

Introduction

Highly conserved across the animal kingdom [1], sleep is important for cognitive, social, and emotional functioning [2–4] across the lifespan [5, 6]. Indeed, sleep deprivation has been associated with cognitive and emotional dysfunction [7–12], sleep disturbances are associated with psychiatric illnesses [13–16] and are part of the known non-motor symptomatology of multiple neurodegenerative disorders [17–22].

Despite the importance of sleep, our understanding of its relationship with brain morphology remains in the early phases [23, 24], and has often returned conflicting findings. Poor sleep quality, in the context of insomnia disorder, for example, has been associated with morphometric differences (e.g., thinner cortex and/or smaller volumes) in orbitofrontal, mesial, and superior frontal cortices, precuneus, hippocampus, thalamus [25–29], although with substantial variance and inconsistencies across studies [30–35], potentially due to methodological differences and relatively small samples. Self-reported sleep quality in healthy individuals, across the lifespan, has been associated cross-sectionally with a number of regions, including frontal and parietal cortices, hippocampus, amygdala, and

thalamus [36–38], and longitudinally with changes spanning multiple cortices (i.e., frontal, temporal, and parietal) [36, 39] including hippocampus [40] and the posterior cingulate cortex [41] specifically (see [42] for a recent overview of findings). Further conflating the issue, recent work uncovered moderating effects of several variables including sex [34], age (at least during adolescence) [43], chronotype [44], and developmental socioeconomic context [45] on the neural correlates of sleep quality. Even as large sample studies emerge [34, 39, 40, 46–50], results are yet to coalesce and remain variable, perhaps also due also to different methodological choices [51], and are often associated with relatively small effect sizes. Nonetheless, understanding the relationship between brain and sleep is key to the development of novel interventions aimed at mitigating sleep-related disorders [52].

In what follows, the full release of the Human Connectome Project (HCP) [53] is leveraged to assess the relationship between self-reported sleep quality measures [54] and brain morphometry using local (i.e., voxelwise) cortical [55, 56] and subcortical [57] measures, as opposed to the global and regional features (i.e., average regional volumes, areas, and thickness) used in some prior work [40, 48, 50]. In particular, while prior studies have leveraged a subset of the same data for assessing functional and structural associations with sleep quality [34, 58], the approach used here adopts a more sensitive (i.e., voxelwise) approach at for both cortex, assessing differences in local concentration of gray matter [55], and subcortex, assessing differences in local shape of deep brain nuclei [57]. Importantly, these voxelwise methods have previously been shown to be more sensitive and better able to capture differences between groups than more global (e.g., regional) measures [59, 60].

As described below, this sensitive approach uncovers systematic associations between self-reported sleep quality and subcortical shape. In addition, variance in cortical and subcortical neuroanatomy is also associated with the interaction of sleep quality related measures and demographic and biometric variables, highlighting the multifaceted nature of brain-sleep relationships.

Methods

Sample description

This project is based on the Q3 release of the Human Connectome Project (HCP [53]). Out of the 1206 total participants, 93 could not be included in the analysis due to failure to acquire at least one T1-weighted MRI, resulting in an included sample of 1,113 observations. (As described below, one additional observation was excluded due segmentation failure, resulting in a final analyzed sample of 1,112 participants). The final analyzed sample included data from 605 female and 507 male participants, with a median age of 29 years. As described in the statistical analysis section, further demographic (i.e., handedness, years of education) and biometric (i.e., body mass index; BMI) factors, were also collected and used as covariates in assessing the relationshp between self-reported sleep quality and neuroanatomy (See Tab. 1 and Tab. SOM1). Because BMI was not recorded for one participant, and years of education was not recorded for two (different) participants, data was imputed with a median imputation strategy, which has been previously been shown to be equally effective as more complex approaches [61], particularly in datasets with low degrees of data missingness [62]. While this approach could decrease covariate variability, the standard deviation for each variable is

only decreased by 0.05% (i.e., from 1.799 to 1.798) and 0.04% (i.e., from 5.199 to 5.197) for years of education and BMI, respectively. This is consistent with the fact that less than 0.2% of the data for each variable is being imputed (in fact, 0.18% and 0.09% for years of education and BMI, respectively).

Fig. 1. PSQI data. (a) Distribution of PSQI total scores in the analyzed sample. The vertical black line indicates the conventional criterion discriminating "poor sleepers" (i.e., PSQI total score > 5) from "good sleepers" (i.e., PSQI total score ≤ 5; [54]). (b-d) Distribution of PSQI total score, PSQI PCA PC1, and PC2 by sex and age group. (No significant differences were observed in the distribution of PSQI total score, or either PCA component, across sex or age group.)

PSQI data

For each participant, the total score and the individual component scores of the Pittsburgh sleep Quality Index (PSQI [54]) were collected from the most recent HCP data distribution. Due to extensive correlations across the seven components of the PSQI (see SOM2), prior to analyzing their relationship to brain structures, a data reduction was performed using a Principal Component Analysis (PCA; [63]) approach. The PCA was performed in JASP 2024 (v. 0.19.3; [64]), using an orthogonal varimax rotation and retaining all components with an eigenvalue greater than 1. This data reduction resulted in two orthogonal components explaining 50% of the variance. As shown in Tab. 1, the first principal component was loaded upon mainly by subjective sleep quality, sleep duration, and habitual sleep efficiency (with loadings equal to 0.62, 0.83, and 0.74, respectively). The second principal component was loaded upon by sleep onset latency, sleep disturbances, and use of sleep medications (with loadings equal to 0.67, 0.61, and 0.69, respectively). The daytime dysfunction PSQI component loaded weakly (< 0.36) on both both PCs. As is often the case with this measure, the PSQI total score, as well as the two PCs, are positively skewed (skewness coefficients: 1.029, 1.237, and 1.252, respectively) and not normally distributed, as assessed with a Shapiro-Wilk test of normality (W=0.931, p<0.001; W=0.907, p<0.001; and W=0.927, p<0.001, for the total score, PC1 and PC2, respectively). The ensuing exploration of associations between PSQI, the two principal components, demographic and biometric variables (i.e., sex, age, years of

education, handedness, and BMI), and their interactions, are thus performed with a non-parametric approach (as described in the MRI data analysis section).

Subcortical shape

Thal
Caud
Putm
GP
Hipp
BrStem

T1-weighted data segmentations

Cortical voxel based morphometry

Fig. 2. Methods. Left: Illustration of cortical and subcortical segmentations of the T1-weighted data. Right: Illustration of 3D subcortical (top) and cortical (bottom) reconstructed meshes. (Image partially adapted from [65])

MRI data

A total of 1,113 T1-weighted data were obtained from the HCP Young Adult repository. For each participant, the "T1w_MPR1" dataset acquired as part of the "structural session" was employed. As described in the HCP Q3 release appendix (available at http://www.humanconnectome.org) these data were acquired with a Magnetization-Prepared Rapid Acquisition Gradient Echo (MPRAGE) sequence (TR = 2,400 ms, TE = 2.14 ms, TI = 1,000 ms, GRAPPA acceleration factor = 2). The acquisition resulted in an image size of $256 \times 320 \times 243$ voxels (in the x, y, and z directions, respectively) and an isovoxel resolution of 0.7 mm.

MRI data analysis

T1-weighted data preprocessing and segmentation

All MRI data analyses were performed with the FMRIB Software Library (FSL; [66]). T1-weighted data were first preprocessed using the `fsl_anat` script, which provides a general pipeline for processing anatomical images. For each T1-weighted image, the script reorients images to the standard (MNI) orientation (`fslreorient2std`), crops the image (`robustfov`), performs bias-field correction (RF/B1-inhomogeneity-correction; `FAST`; [67]), registration to standard space (linear and non-linear; `FLIRT` and `FNIRT`, [68, 69]), brain-extraction (`FNIRT`-based), tissue-type segmentation

Table 1. Sample demographic, biometric, and PSQI information. (Abbreviations: Med., median; sd., standard deviation; PC1, PSQI Principal Component 1; PC2, PSQI Principal Component 2; Subj. sleep qual., subjective sleep quality; sleep dur., sleep duration; sleep eff., sleep efficiency; sleep ons. lat., sleep onset latency; sleep disturb., sleep disturbances; sleep med., use of sleep medications; daytime dysf., daytime dysfunction;. See Tab. SOM1 for the same information broken down by sex.)

Variables	Med	Mean	sd	Range
PSQI Total Score	4	4.804	2.759	[0, 19]
Subj. sleep qual. (PC1)	1	0.894	0.635	[0,3]
Sleep dur. (PC1)	0	0.569	0.815	[0,3]
Sleep eff. (PC1)	0	0.456	0.795	[0,3]
Sleep ons. lat. (PC2)	1	0.967	0.817	[0,3]
Sleep disturb. (PC2)	1	1.093	0.485	[0,3]
Sleep med. (PC2)	0	0.237	0.677	[0,3]
Daytime dysf. ($-$)	1	0.588	0.635	[0,3]
Age	29	28.799	3.696	[22,37]
Handedness	80	65.845	44.208	[-100,100]
Education	16	14.927	1.799	[11,17]
BMI	25.53	26.534	5.199	[16.5,47.8]

(`FAST`; [67]), and subcortical structure segmentation (`FIRST`; [57]). Once subcortical segmentations were completed, and before any analysis, the resulting subcortical meshes were individually inspected by the author to ensure correct structure segmentation (cf., Fig. 2). During this control, it emerged that segmentation for one participant failed, not producing any mesh output. That datapoint was thus excluded from any further analysis (leading to a final analyzed sample of 1,112 observations). Cortical segmentations were obtained by feeding the bias-corrected T1-weighted data to FSL Voxel Based Morphometry (VBM; [70]) scripts, an optimized VBM protocol [56] enclosed in the FSL distribution. Again, segmentations were visually inspected for accuracy prior to any analysis. All 1,112 T1-weighted images return satisfactory segmentations of the cortical gray matter (cf., Fig. 2).

T1-weighted data analysis

The single-smoothed subject cortical data (Gaussian smoothing with $\sigma = 2$, resulting in an approximate 4.71 mm full width half max [FWHM] Gaussian kernel) and subcortical meshes were entered into a group analysis. While both cortical and subcortical tissues are segmented in the subject's native space, subcortical meshes are then co-registered to the standard MNI space for the purposes of group analysis. Conversely, cortical segmentations are co-registered to an *ad-hoc* sample-specific cortical template created, automatically, as an average of all the $1,112$ input data, for the purposes of performing the VBM group analysis. Each set of segmentations (i.e., cortical, subcortical) was entered into two group analyses. In the first analysis (henceforth, PSQI total score analysis), segmentations were regressed on the PSQI total score. In the second analysis (henceforth, PSQI PC analysis), segmentations were regressed on the two PC components described above. Each of these two statistical models also included a number of demographic (i.e., sex, age, handedness, years of education) and biometric (body mass index, BMI) sample characteristic variables. In addition, each analysis also included the interaction of each sample characteristic covarariate with the PSQI total score (in the PSQI total score analysis) and with PC1 and PC2 (in the PSQI PC analysis). Finally, each regression also included as an independent variable the subject's total normalized brain

Fig. 3. Correlates of sleep quality measures. Depiction of all voxels significantly associated with PSQI total score regressor (left), in the PSQI total score analysis, and with the PSQI PC1 regressor (right), in the PSQI PC analysis. As shown, only negative associations were detected between subcortical shape and the two sleep quality variables. No significant associations were detected in cortex for any of the sleep quality measures, neither positive nor negative, and no significant associations were observed for the PC2 component, in either cortex or subcortical regions, either positive or negative. (Note: colored areas imply a significant association, corrected for multiplicity; gray areas imply no significant associations.) See Tabs. SOM3 and SOM4 for detailed localization of each area of significant association.

volume (NBV; as calculated with `sienax`; [71]), in order to account for the effect of head size variability across individuals and to ensure that results reflect local shape change [65, 72].

Group analyses were performed with a non-parametric permutation testing approach [73] using `randomise` [74]. Significance was assessed against a criterion of $\alpha \leq 0.05$ corrected for multiplicity with a family-wise cluster correction approach with Threshold Free Cluster Enhancement (`TFCE`; [75, 76]).

Results

PSQI scores

As shown in Fig. 1 and in Tab. 1, the PSQI total score in the analyzed sample had a mean of 4.8 and a median of 4. Using the conventional PSQI > 5 criterion [54], 31.8% of participants in the analyzed sample were classified as having poor sleep quality. The PSQI total score was not significantly different across sex (Mann-Whitney test $[U_{MW}]$=147,234, p=0.246, effect size [rank-biserial correlation, r_{rb}]=-0.04) and was not significantly correlated with age (Kendall's Tau B $[\tau_B]$=-0.019, p=0.379, effect size [Fischer's z]=-0.019). However, PSQI was significantly associated with handedness (τ_B=-0.064, p=0.003, Fisher's z=-0.064), with greater right-handed lateralization showing smaller PSQI total scores (i.e., better sleep quality), years of education (τ_B=-0.108, p<0.001, Fisher's z=-0.108), with more years of education being associated with better sleep quality, and BMI (τ_B=0.072, p<0.001, Fisher's z=0.072), with greater biometric values being associated with worse sleep quality (see Fig. SOM1 for a depiction of the corrrelations between PSQI total score, PSQI PC1, PC2, and all sample characteristic covariates).

With respect to the two PSQI PCs, in terms of their interpretation, both scores were positively associated with higher PSQI scores, and therefore worse sleep quality (τ_B=0.584, p<0.001, Fisher's z=0.669 and τ_B=0.478, p<0.001, Fisher's z=0.521, for PC1 and PC2, respectively). Similarly to the PSQI total score, neither PC was significantly different across sex (U_{MW}=152,589, p=0.884, r_{rb}=-0.005 and U_{MW}=146,676, p=0.209, r_{rb}=-0.044, for PC1 and PC2, respectively) and neither was significantly correlated with age (τ_B=-0.001, p=0.953, Fischer's z=-0.001 and τ_B=-0.008, p=0.689, Fischer's z=-0.008, for PC1 and PC2, respectively). Interestingly, PC1's associations with the remaining covariate measures paralleled those reported for the PSQI total score (i.e., handedness: τ_B=-0.058, p=0.006, Fischer's z=-0.058; BMI: τ_B=0.103, p<0.001, Fischer's z=0.104; education: τ_B=0.103, p<0.001, Fischer's z=0.103). In contrast, no significant associations were observed between PC2 and any of the remaining covariates (i.e., handedness: τ_B=-0.012, p=0.566, Fischer's z=-0.012; BMI: τ_B=0.003, p=0.884, Fischer's z=0.003; education: τ_B=-0.023, p=0.303, Fischer's z=-0.023). (See Fig. SOM1 for a graphical depiction of the PCs correlations with PSQI total score and covariates.)

MRI results

As shown in Fig. 3, the PSQI total score was associated, inversely, with subcortical shape in portions of the bilateral caudate and putamen, implying an association between poorer sleep and smaller shapes in these regions (cf., Tab. SOM3 for precise neuroanatomical localization of significant clusters). No positive associations were detected between the PSQI total score and subcortical shape. No significant associations were

Fig. 4. Subcortical correlates of interactions with sample characteristic variables. Left: Depiction of all voxels significantly associated with the interaction of PSQI total score with each demographic and biometric covariate (i.e., sample characteristics variables). Right: Depiction of all voxels significantly associated with the interaction of PC1 and PC2, respectively, with each demographic and biometric covariate. See Tab. SOM3 and SOM4, respectively, for anatomical localization and statistics of each cluster. (Areas shown in color indicate significant voxels after correction for multiple comparisons. Areas shown in gray imply no significant result. '+' indicates to a positive association; '-' indicates a negative association.)

detected, either positive or negative, between voxelwise cortical morphometry and the PSQI total score.

Mirroring the correlation observed for the PSQI total score, as shown in Fig. 3, the PSQI PC1 was also negatively associated with subcortical shape in similar portions of the bilateral caudate and putamen, as well as additional portions of the left pallidum (see Tab. SOM4 for precise neuroanatomical localization of significant clusters), again implying an association between smaller subcortical shapes and poorer sleep quality, particularly with respect to subjective sleep quality, sleep duration, and habitual sleep efficiency. No positive associations were detected between PC1 and subcortical shape. As for the PSQI total score, no significant associations were detected, either positive or negative, with cortical morphometry. Finally, PC2, the PSQI principal component which was loaded upon maximally by sleep onset

latency, sleep disturbance, and use of sleep medications, did not exhibit any significant association, either positive or negative, with either subcortical shape or cortical morphometry.

In addition to the associations reported above for the regressors of interest (i.e., PSQI total score and the PSQI PCs), significant cortical and subcortical associations were also detected when assessing the interaction of each regressors of interest with demographic and biometric sample characteristics. As shown in Fig. 4, significant associations were uncovered between subcortical shape and the interaction of the PSQI total score with BMI, education, and handedness (see also Tab. SOM3). Specifically, a negative relationship was observed between the interaction of PSQI and BMI and shape in portions of the bilateral pallidum and hippocampus, as well as between the interaction of PSQI and handedness and portions of the bilateral caudate, hippocampus, and right amygdala. Conversely, a positive association was observed between the interaction of PSQI and years of education, and portions of the bilateral caudate, right thalamus and right pallidum.

Significant associations with interaction terms were also uncovered in the PSQI PC analysis (cf., Fig. 4 and Tab. SOM4). Specifically, the interaction of PC1 and BMI was positively correlated with subcortical shape in a small portion of the left amygdala. PC2, despite showing no direct association with any of the sleep quality regressors (i.e., PSQI total score, PSQI PCs), did exhibit multiple significant associations with the interaction of the sleep quality regressors with age, BMI, handedness, and sex. Specifically, the interaction of PC2 and age associated negatively with subcortical shape in portions of the left hippocampus, right caudate and amygdala. The interaction of PC2 and BMI revealed extensive negative associations with subcortical shape in large portions of the bilateral thalamus, caudate, putamen, pallidum, hippocampus, as well as the left amygdala and the brainstem. The interaction of PC2 and handedness also revealed a negative association with shape across bilateral caudate, hippocampus, right thalamus, and left amygdala. Similarly, the interaction of PC2 and sex was also associated with subcortical shape in broad portions of the bilateral pallidum, left caudate and putamen, as well as the brainstem. Finally, a small positive association was detected between subcortical shape in the left putamen and the interaction of PC2 and education.

As shown in Fig. 5, despite the absence of any significant association between cortical voxel based morephometry and the main sleep quality regressors (i.e., PSQI total score and PSQI PCs), a number of significant associations were detected between cortical morphometry and PC interaction terms. Specifically, the interaction of PC1 and age revealed positive associations in the bilateral insula, the left parahippocampal gyrus and temporal pole, as well as medial occipital cortex, in the occipito-parietal sulcus and lingual gyrus. In addition, two small negative associations were found, between PC1 and age, in left posterior parietal cortex. The interaction of PC2 and education was negatively associated with with morphometry measures in the right anterior insular cortex, while the interaction of PC2 and sex was associated with morphometry measures in two foci in the medial wall of frontal cortex, in the regions of the supplementary motor area and the paracingulate gyrus. (See Tab. SOM5 for precise neuroanatomical localization of significant clusters.)

Fig. 5. Cortical correlates of interactions with sample characteristic variables. Depiction of all voxels significantly associated with the interaction of PSQI PC1 and PC2 with each demographic and biometric covariate (i.e., sample characteristics variables). (Note that no significant associations were found between cortical morphometry and the interaction of PSQI total score and sample characteristic variables.) See Tab. SOM5 for anatomical localization and statistics of each significant cluster. (Areas shown in color indicate significant voxels after correction for multiple comparisons. Areas shown in gray imply no significant result. '+' indicates to a positive association; '-' indicates a negative association.)

Discussion

The present work shows that lower self-reported sleep quality and, in particular, subjective sleep quality, sleep duration, and habitual sleep efficiency, is associated with smaller subcortical shapes in the bilateral striatum and the left pallidum, after parceling out any variance explained by sample characteristics (i.e., demographic and biometric covariates) and their interaction with sleep quality measures.

Long known to serve a key role in the control of motor behavior [77], the contributions of the basal ganglia to non-motor function have only come to the forefront relatively recently [52, 78, 79], together with a renewed understanding of

the complex circuitry tying these regions together and within broader cortical-subcortical circuits [80, 81]. Indeed, clinical data show that patients with striatal dysfunction syndromes, such as Parkinson's and Huntington Disease, exhibit high rates of sleep disturbances [17–20, 82], something that is likely to be mediated, at least in part, by dopamine mechanisms [83]. Patients suffering from insomnia disorders show an association between shape change in many of these regions and sleep quality measures [60]. Interestingly, the availability of striatal dopamine in D2/D3 receptors has been associated with sleep quality in healthy adults [84]. Recent work in a large cohort of healthy volunteers also highlighted the association between neural tissue density in these regions (as assessed with diffusion tensor imaging) and sleep duration [85]. Consistent with these results, and the present findings, lesion studies in the rodent model have documented alteration of sleep-wake behavior following striatal damage [86, 87]. In particular, striatal lesions have been shown to lead to greater sleep-wake behavior fragmentation, including greater state transitions, loss of ultra-long wake bouts, and diminished diurnal sleep-wake variation [88]. It is not surprising, then, that despite significant heterogeneity in findings and measures (e.g., PSQI versus non-normative sleep duration), neuroanatomical variability in these regions has been previously associated to measures of sleep quality in healthy individuals [48–50]. Also conventionally conceived as a relay in the control of motor behavior [89–91], a growing amount of evidence links the pallidum – and its external portion in particular – to the regulation of sleep-wake behavior and electrophysiological arousal [78, 79], likely *via* dopaminergic neuromodulation [92]. Indeed, as found with the striatum, excitotoxic lesions of the external portion of the pallidum, in rodent models, lead to increased sleep fragmentation, with increased sleep-wake transitions and decreased sleep bouts [88, 93]. Conversely, optogenetic and electical neuromodulation of this region results in a broad increase of time spent sleeping [93, 94]. Consistent with these findings, in humans, the pallidum has been associated with behavioral and/or electrocortical arousal in healthy volunteers undergoing general anesthesia [95], and in severe brain injury patients [65, 72]. Intriguingly, and mirroring the rodent work, deep brain stimulation of the external portion of the pallidum in a single parkinsoninan patient dramatically reduced their sleep-related symptomatology [96].

The interaction of sleep quality, demographic, and biometric factors highlighted several additional associations with brain neuroanatomy in both subcortical (i.e., hippocampus, thalamus, amygdala, and brainstem) and cortical regions (i.e., sublobar, insular, and mesial cortices). Many of these areas have been previously associated with aspects of sleep quality. Hippocampal morphometry, for example, has been previously shown to have a negative relation with sleep quality indices, both cross-sectionally [49, 50] and longitudinally [40]. Similarly, prior work using diffusion tensor imaging also highlighted an association between (right) hippocampus and greater tissue density (as measured with mean diffusivity) [85]. Thalamus, and its interactions with cortex, have long been implicated in sleep-wake transitions, as a key target of ascending arousal system [97, 98]. Large sample population studies have revealed associations between non-normative sleep duration and macrostructural and microstructural brain outcomes, including thalamus (as well as hippocampus, putamen, and accumbens) [48, 49]. Sleep disturbances have also been associated with thalamic abnormalities and/or dysfunction in healthy volunteers [99] and across many

mental health and neuropsychiatric disorders [49, 100–102]. Furthermore, thalamic stroke and infraction are known to lead to sleep impairments [103, 104] and long-term sleep deprivation has been associated with decreased thalamic volume [105]. Importantly, in patients with insomnia, clinical features of the pathology are also associated with the connectivity between thalamus and many of the regions uncovered in the present work including cingualte cortex, hippocampus, and striatum [100]. Atrophy and connectivity dysfunction in the amygdala has also been associated with sleep disturbances across a number of conditions [50, 106–109], dissociably from its co-morbid association with anxiety [110]. Finally, structural and functional aspects of insular cortex, including its interactions with cingulate cortex [111], have been previously shown to be disrupted by sleep deprivation [112] as well as sleep disorders [113–116].

Despite this superficial consistency with prior work, it cannot be ignored that in the present work these associations are specific to the interaction of sleep quality measures and sample characteristics (i.e., demographic and biometric variables). This raises an important methodological issue in addition to those previously discussed [51]: while almost all studies make use of a number – often large – of covariates in assessing their effects of interest, many do not assess any interaction [e.g., 38, 49, 85, 99, 117] and, those that do, only include one or two interactions of interest (e.g., sex, age) [e.g., 34, 40, 48, 118]. In the context of a regression-type analysis, this can mean that variance might be misallocated to some extent, potentially resulting in spurious results. One interesting case in point is the previously reported association between the PSQI×Sex interaction and local neuroanatomical features in frontal cortex, along with sex-specific differences in the parahippocampus and parietal cortices [34]. In the present analysis no such effect was observed. Yet, as shown in Fig. 5, a subset of these regions were observed to associate – albeit with important lateralization differences – with the interaction of PSQI and age. Of course, whether the discrepancy is due to the use of different statistical models (e.g., inclusion of different interaction terms) or to other factors remains to be understood. Nonetheless, it is noteworthy that, while the PSQI total score by sex interaction did not exhibit any significant association with either cortical or subcortical morphometry in the analysis presented above (cf., Figs. 4,5 and Tabs. SOM3-SOM5), when the analysis was re-run omitting all interaction terms with the exception of one interaction of interest (i.e., PSQI×Sex), significant associations were detected in multiple subcortical regions (see Fig. SOM2 for a depiction of the result). This finding further highlights the multifaceted nature of brain-sleep quality relations as well as the need for considering more sophisticated approaches, in testing such associations, that can better model complex structures (e.g., structural equation modeling [119, 120], causal discovery analysis [121], Bayesian network analysis [122]).

Although not an aim of the present work, the results reported above also contribute to the discussion concerning whether the factor structure of the PSQI includes more than one dimension and, if it does, how individual items/components of the instrument distribute across dimensions [123–125]. Here, an exploratory PCA with varimax rotation was employed as a statistical device for the sole purposes of obtaining (rotated) components which, if more than one, would be uncorrelated with each other, thus avoiding the issue of excessive multicollinearity in the general linear model analysis [126]. Nonetheless, it is interesting to observe that

the two components derived from this approach exhibited a different correlational structure with demographic and biometric variables, brain measures, and their interactions. PC1, which was loaded maximally upon by subjective sleep quality, sleep duration, and habitual sleep efficiency, exhibited a similar correlational structure with demographic and biometric covariates to that of the PSQI total score. Both were similarly and significantly associated with handedness, BMI, and years of education. Conversely, PC2, which was loaded upon maximally by sleep onset latency, sleep disturbance, and use of sleep medications, exhibited no such associations (cf., Tab. 1 and Fig. SOM1). The two PCs also exhibited a different correlational structure with brain measures (cf., Fig. 4,5 and Tab. SOM3-SOM5). Indeed, PC1 exhibited a similar association with subcortical areas as that observed for the PSQI total score. Conversely, PC2 did not show any significant associations with cortical or subcortical morphometry. It is also relevant that while the PSQI total score failed to show any significant association with cortical morphometry, both in the context of the main PSQI total score regressor as well as its interaction with sample characteristics, PC1 and PC2 did return significant cortical morphometry associations with select interactions, indicating that they are capturing different aspects of the relationship between neuroanatomical variability and self-reported sleep quality measures. It should be highlighted, however, that all the above considerations result from data-driven observations, not systematic assessments of the dimensionality of the PSQI, and should thus be interpreted as such.

Finally, in assessing the present findings, one must take stock of a number of limitations tied to the approach employed. First, the segmentation technique employed here [57] cannot distinguish a number of regions and subregions (e.g., parabrachial nucleus [127, 128], internal versus external segments of the globus pallidus [129–131], reticular [132–134], anterior and mediodorsal [135] nuclei), cell populations (e.g., cell versus matrix thalamic neurons [136]), as well as neurotransmetter-specific (e.g., glutamatergic, GABAergic [137]) neuromodulatory systems, that are believed to play an important role in the regulation of sleep-wake behavior. Second, the morphometric approach employed centers on local cortical and subcortical structures, but is blind to other factors (e.g., white matter integrity, glymphatic system, perivascular space), which have been proposed to play a role in sleep [117, 138–141]. Third, the present work is focused entirely on neuroanatomical variability, yet there exist known associations between changes in brain function and sleep quality [58, 142–147]. It will be interesting, in future work, to delineate the degree to which (if any) the known associations between brain network function and perceived sleep quality [e.g., 58, 148] is related to the neuroanatomical variability observed here. Finally, while longitudinal studies have reported atrophy and smaller volumes over time to be associated with decreasing sleep quality [e.g., 36, 40, 118], the present findings are cross-sectional in nature and do not permit claims of causality in the uncovered brain-sleep quality associations.

Conclusion

This study demonstrates a robust association between conventional self-reported measures of sleep quality [54] and subcortical morphometry, particularly in the bilateral caudate, putamen, and left pallidum. These findings provide evidence for the importance of the basal ganglia in sleep regulation in healthy, young, individuals, reinforcing prior work linking striatal function to sleep disturbances in both healthy and clinical populations [52, 78, 79]. While cortical morphometry did not directly correlate with sleep quality measures, interactions with demographic and biometric factors were observed, emphasizing the multifaceted nature of brain-sleep relationships. These results underscore the importance of considering both direct and interaction effects when studying neuroanatomical correlates of sleep. Finally, in light of the rising availability of non-invasive deep brain neuromodulatory techniques (e.g., low-intensity transcranial focused ultrasound [149–151]), the potential identification of subcortical structures as sleep-related hubs could be investigated causally and, should such a link be established, could present an opportunity for targeted interventions aimed at improving sleep quality in both healthy individuals and clinical populations [52, 96].

Data availability statement

The data underlying this article are publicly available at https://www.humanconnectome.org/study/hcp-young-adult. This manuscript is made available on BiorXiv as preprint https://doi.org/10.1101/2024.05.29.596530.

Funding

This work is supported in part by funds from the NIH NIGMS (R01GM135420), DOD CDMRP (HT9425-24-1-1081), and the Brain Injury Research Center at UCLA. Data were provided by the Human Connectome Project, WU-Minn Consortium (Principal Investigators: David Van Essen and Kamil Ugurbil; 1U54MH091657) funded by the 16 NIH Institutes and Centers that support the NIH Blueprint for Neuroscience Research; and by the McDonnell Center for Systems Neuroscience at Washington University.

Acknowledgments

The author would like to acknowledge the UCLA Institute for Digital Research and Education (IDRE) for making temporarily available to investigators large computational space and resources for allowing processing and analysis of large datasets.

Disclosures

No competing interest is declared.

Author contributions statement

M.M.M. conceived the experiment, analysed the data, and wrote the manuscript.

References

1. Keene Alex C., Duboue Erik R.. The origins and evolution of sleep *Journal of Experimental Biology.* 2018;221:jeb159533.

2. Walker Matthew P. The role of sleep in cognition and emotion *Annals of the New York Academy of Sciences.* 2009;1156:168–197.

3. Goldstein Andrea N, Walker Matthew P. The role of sleep in emotional brain function *Annual review of clinical psychology.* 2014;10:679–708.

4. Djonlagic Ina, Mariani Sara, Fitzpatrick Annette L, et al. Macro and micro sleep architecture and cognitive performance in older adults *Nature human behaviour.* 2021;5:123–145.

5. Yaffe Kristine, Falvey Cherie M, Hoang Tina. Connections between sleep and cognition in older adults *The Lancet Neurology.* 2014;13:1017–1028.

6. Galván Adriana. The need for sleep in the adolescent brain *Trends in cognitive sciences.* 2020;24:79–89.

7. Lim Julian, Dinges David F. A meta-analysis of the impact of short-term sleep deprivation on cognitive variables. *Psychological bulletin.* 2010;136:375.

8. Krause Adam J, Simon Eti Ben, Mander Bryce A, et al. The sleep-deprived human brain *Nature Reviews Neuroscience.* 2017;18:404–418.

9. Simon Eti Ben, Vallat Raphael, Barnes Christopher M, Walker Matthew P. Sleep loss and the socio-emotional brain *Trends in Cognitive Sciences.* 2020;24:435–450.

10. García Aída, Del Angel Jacqueline, Borrani Jorge, Ramirez Candelaria, Valdez Pablo. Sleep deprivation effects on basic cognitive processes: which components of attention, working memory, and executive functions are more susceptible to the lack of sleep? *Sleep Science.* 2021;14:107.

11. Tomaso Cara C, Johnson Anna B, Nelson Timothy D. The effect of sleep deprivation and restriction on mood, emotion, and emotion regulation: three meta-analyses in one *Sleep.* 2021;44:zsaa289.

12. Li Jiahui, Cao Yixuan, Ou Simei, Jiang Tianxiang, Wang Ling, Ma Ning. The effect of total sleep deprivation on working memory: evidence from diffusion model *Sleep.* 2024:zsae006.

13. Koyanagi Ai, Stickley Andrew. The Association between Sleep Problems and Psychotic Symptoms in the General Population: A Global Perspective *Sleep.* 2015;38:1875-1885.

14. Freeman Daniel, Sheaves Bryony, Waite Felicity, Harvey Allison G, Harrison Paul J. Sleep disturbance and psychiatric disorders *The Lancet Psychiatry.* 2020;7:628–637.

15. Cheng Wei, Rolls Edmund, Gong Weikang, et al. Sleep duration, brain structure, and psychiatric and cognitive problems in children *Molecular psychiatry.* 2021;26:3992–4003.

16. Sun Xiaohui, Liu Bin, Liu Sitong, et al. Sleep disturbance and psychiatric disorders: a bidirectional Mendelian randomisation study *Epidemiology and Psychiatric Sciences.* 2022;31:e26.

17. Emser W, Brenner M, Stober T, Schimrigk K. Changes in nocturnal sleep in Huntington's and Parkinson's disease *Journal of neurology.* 1988;235:177–179.

18. Morton A Jennifer, Wood Nigel I, Hastings Michael H, Hurelbrink Carrie, Barker Roger A, Maywood Elizabeth S. Disintegration of the sleep-wake cycle and circadian timing in Huntington's disease *Journal of Neuroscience.* 2005;25:157–163.

19. Zhang Ye, Ren Rong, Yang Linghui, et al. Sleep in Huntington's disease: a systematic review and meta-analysis of polysomongraphic findings *Sleep.* 2019;42:zsz154.

20. Stefani Ambra, Högl Birgit. Sleep in Parkinson's disease *Neuropsychopharmacology.* 2020;45:121–128.

21. Romanella SM, Roe D, Tatti E, et al. The sleep side of aging and Alzheimer's disease *Sleep medicine.* 2021;77:209–225.

22. Oh Jun Y., Walsh Christine M., Ranasinghe Kamalini, et al. Subcortical Neuronal Correlates of Sleep in Neurodegenerative Diseases *JAMA Neurology.* 2022;79:498-508.

23. Lima Santos João Paulo, Hayes Rebecca, Franzen Peter L, et al. The association between cortical gyrification and sleep in adolescents and young adults *Sleep.* 2024;47:zsad282.

24. Wei Ruoqi, Ganglberger Wolfgang, Sun Haoqi, et al. Linking brain structure, cognition, and sleep: insights from clinical data *Sleep.* 2024;47:zsad294.

25. Altena Ellemarije, Vrenken Hugo, Van Der Werf Ysbrand D, Heuvel Odile A, Van Someren Eus JW. Reduced orbitofrontal and parietal gray matter in chronic insomnia: a voxel-based morphometric study *Biological psychiatry.* 2010;67:182–185.

26. Riemann Dieter, Voderholzer Ulrich, Spiegelhalder Kai, et al. Chronic insomnia and MRI-measured hippocampal volumes: a pilot study *Sleep.* 2007;30:955–958.

27. Neylan Thomas C, Mueller Susanne G, Wang Zhen, et al. Insomnia severity is associated with a decreased volume of the CA3/dentate gyrus hippocampal subfield *Biological psychiatry.* 2010;68:494–496.

28. Joo Eun Yeon, Kim Hosung, Suh Sooyeon, Hong Seung Bong. Hippocampal substructural vulnerability to sleep disturbance and cognitive impairment in patients with chronic primary insomnia: magnetic resonance imaging morphometry *Sleep.* 2014;37:1189–1198.

29. Li Min, Wang Ruonan, Zhao Meng, et al. Abnormalities of thalamus volume and resting state functional connectivity in primary insomnia patients *Brain Imaging and Behavior.* 2019;13:1193–1201.

30. Winkelman John W, Plante David T, Schoerning Laura, et al. Increased rostral anterior cingulate cortex volume in chronic primary insomnia *Sleep.* 2013;36:991–998.

31. Spiegelhalder Kai, Regen Wolfram, Baglioni Chiara, et al. Insomnia does not appear to be associated with substantial structural brain changes *Sleep.* 2013;36:731–737.

32. Li Meng, Yan Jianhao, Li Shumei, et al. Altered gray matter volume in primary insomnia patients: a DARTEL-VBM study *Brain Imaging and Behavior.* 2018;12:1759–1767.

33. Yu Siyi, Feng Fen, Zhang Qi, et al. Gray matter hypertrophy in primary insomnia: a surface-based morphometric study *Brain imaging and behavior.* 2020;14:1309–1317.

34. Neumann Nicola, Lotze Martin, Domin Martin. Sex-specific association of poor sleep quality with gray matter volume *Sleep.* 2020;43:zsaa035.

35. Aquino Giulia, Benz Fee, Dressle Raphael, et al. Towards the neurobiology of insomnia: A systematic review of neuroimaging studies *Sleep Medicine Reviews.* 2023:101878.

36. Sexton Claire E, Storsve Andreas B, Walhovd Kristine B, Johansen-Berg Heidi, Fjell Anders M. Poor sleep quality is associated with increased cortical atrophy in community-dwelling adults *Neurology.* 2014;83:967–973.

37. Liu Yun-Rui, Fan Dong-Qiong, Gui Wen-Jun, Long Zhi-Liang, Lei Xu, Yu Jing. Sleep-related brain atrophy

and disrupted functional connectivity in older adults *Behavioural brain research.* 2018;347:292–299.

38. Alperin Noam, Wiltshire John, Lee Sang H, et al. Effect of sleep quality on amnestic mild cognitive impairment vulnerable brain regions in cognitively normal elderly individuals *Sleep.* 2019;42:zsy254.

39. Fjell Anders M, Sørensen Øystein, Amlien Inge K, et al. Poor self-reported sleep is related to regional cortical thinning in aging but not memory decline—results from the Lifebrain Consortium *Cerebral Cortex.* 2021;31:1953–1969.

40. Fjell Anders M, Sørensen Øystein, Amlien Inge K, et al. Self-reported sleep relates to hippocampal atrophy across the adult lifespan: results from the Lifebrain consortium *Sleep.* 2020;43:zsz280.

41. Liu Che, Lee Sang H, Loewenstein David A, Galvin James E, Camargo Christian J, Alperin Noam. Poor sleep accelerates hippocampal and posterior cingulate volume loss in cognitively normal healthy older adults *Journal of sleep research.* 2022;31:e13538.

42. André Claire, Laniepce Alice, Chételat Gaël, Rauchs Géraldine. Brain changes associated with sleep disruption in cognitively unimpaired older adults: a short review of neuroimaging studies *Ageing research reviews.* 2021;66:101252.

43. Jalbrzikowski Maria, Hayes Rebecca A, Scully Kathleen E, et al. Associations between brain structure and sleep patterns across adolescent development *Sleep.* 2021;44:zsab120.

44. Rosenberg Jessica, Jacobs Heidi IL, Maximov Ivan I, Reske Martina, Shah NJ. Chronotype differences in cortical thickness: grey matter reflects when you go to bed *Brain Structure and Function.* 2018;223:3411–3421.

45. Hansen Melissa, Simon Katrina R, Strack Jordan, He Xiaofu, Noble Kimberly G, Merz Emily C. Socioeconomic disparities in sleep duration are associated with cortical thickness in children *Brain and Behavior.* 2023;13:e2859.

46. Westwood Andrew J, Beiser Alexa, Jain Nikita, et al. Prolonged sleep duration as a marker of early neurodegeneration predicting incident dementia *Neurology.* 2017;88:1172–1179.

47. Toschi Nicola, Passamonti Luca, Bellesi Michele. Sleep quality relates to emotional reactivity via intracortical myelination *Sleep.* 2021;44:zsaa146.

48. Schiel Julian E, Tamm Sandra, Holub Florian, et al. Associations between sleep health and grey matter volume in the UK Biobank cohort (n= 33 356) *Brain communications.* 2023;5:fcad200.

49. Wang Jian, Huang Cigui, Luo Guowei, et al. Reduced sleep quality defines a subtype of obsessive-compulsive disorder with lower Glx levels in the resting thalamus and worse response inhibition *Journal of Psychiatric Research.* 2024;173:14–24.

50. Stolicyn Aleks, Lyall Laura M, Lyall Donald M, et al. Comprehensive assessment of sleep duration, insomnia, and brain structure within the UK Biobank cohort *Sleep.* 2024;47:zsad274.

51. Spiegelhalder Kai, Riemann Dieter, Schiel Julian. Are sleep and brain structure associated in the UK Biobank? A letter to the editor regarding "Comprehensive assessment of sleep duration, insomnia and brain structure within the UK Biobank cohort." by Stolicyn et al. *Sleep.* 2024;47:zsad317.

52. Hasegawa Harutomo, Selway Richard, Gnoni Valentina, et al. The subcortical belly of sleep: New possibilities in neuromodulation of basal ganglia? *Sleep medicine reviews.* 2020;52:101317.

53. Van Essen David C, Smith Stephen M, Barch Deanna M, et al. The WU-Minn human connectome project: an overview *Neuroimage.* 2013;80:62–79.

54. Buysse Daniel J, Reynolds III Charles F, Monk Timothy H, Berman Susan R, Kupfer David J. The Pittsburgh Sleep Quality Index: a new instrument for psychiatric practice and research *Psychiatry research.* 1989;28:193–213.

55. Ashburner John, Friston Karl J. Voxel-based morphometry—the methods *Neuroimage.* 2000;11:805–821.

56. Good Catriona D, Johnsrude Ingrid S, Ashburner John, Henson Richard NA, Friston Karl J, Frackowiak Richard SJ. A voxel-based morphometric study of ageing in 465 normal adult human brains *Neuroimage.* 2001;14:21–36.

57. Patenaude Brian, Smith Stephen M, Kennedy David N, Jenkinson Mark. A Bayesian model of shape and appearance for subcortical brain segmentation *Neuroimage.* 2011;56:907–922.

58. Zhou Zhongxing, Cai Biao, Zhang Gemeng, Zhang Aiying, Calhoun Vince D, Wang Yu-Ping. Prediction and classification of sleep quality based on phase synchronization related whole-brain dynamic connectivity using resting state fMRI *NeuroImage.* 2020;221:117190.

59. Muñoz-Ruiz Miguel Ángel, Hartikainen Päivi, Koikkalainen Juha, et al. Structural MRI in frontotemporal dementia: comparisons between hippocampal volumetry, tensor-based morphometry and voxel-based morphometry *PloS one.* 2012;7:e52531.

60. Emamian Farnoosh, Mahdipour Mostafa, Noori Khadijeh, et al. Alterations of subcortical brain structures in paradoxical and psychophysiological insomnia disorder *Frontiers in psychiatry.* 2021;12:661286.

61. Berkelmans Gijs FN, Read Stephanie H, Gudbjörnsdottir Soffia, et al. Population median imputation was noninferior to complex approaches for imputing missing values in cardiovascular prediction models in clinical practice *Journal of Clinical Epidemiology.* 2022;145:70–80.

62. Salgado Cátia M, Azevedo Carlos, Proença Hugo, Vieira Susana M. Missing data *Secondary analysis of electronic health records.* 2016:143–162.

63. Pearson Karl. LIII. On lines and planes of closest fit to systems of points in space *The London, Edinburgh, and Dublin philosophical magazine and journal of science.* 1901;2:559–572.

64. JASP Team . JASP (Version 0.18.3)[Computer software] 2024.

65. Lutkenhoff Evan S, Wright Matthew J, Shrestha Vikesh, et al. The subcortical basis of outcome and cognitive impairment in TBI: A longitudinal cohort study *Neurology.* 2020;95:e2398–e2408.

66. Smith Stephen M, Jenkinson Mark, Woolrich Mark W, et al. Advances in functional and structural MR image analysis and implementation as FSL *Neuroimage.* 2004;23:S208–S219.

67. Zhang Yongyue, Brady Michael, Smith Stephen. Segmentation of brain MR images through a hidden

Markov random field model and the expectation-maximization algorithm *IEEE transactions on medical imaging.* 2001;20:45–57.

68. Jenkinson Mark, Smith Stephen. A global optimisation method for robust affine registration of brain images *Medical image analysis.* 2001;5:143–156.

69. Jenkinson Mark, Bannister Peter, Brady Michael, Smith Stephen. Improved optimization for the robust and accurate linear registration and motion correction of brain images *Neuroimage.* 2002;17:825–841.

70. Douaud Gwenaëlle, Smith Stephen, Jenkinson Mark, et al. Anatomically related grey and white matter abnormalities in adolescent-onset schizophrenia *Brain.* 2007;130:2375–2386.

71. Smith Stephen M, Zhang Yongyue, Jenkinson Mark, et al. Accurate, robust, and automated longitudinal and cross-sectional brain change analysis *Neuroimage.* 2002;17:479–489.

72. Lutkenhoff Evan S, Chiang Jeffrey, Tshibanda Luaba, et al. Thalamic and extrathalamic mechanisms of consciousness after severe brain injury *Annals of neurology.* 2015;78:68–76.

73. Nichols Thomas E, Holmes Andrew P. Nonparametric permutation tests for functional neuroimaging: a primer with examples *Human brain mapping.* 2002;15:1–25.

74. Winkler Anderson M, Ridgway Gerard R, Webster Matthew A, Smith Stephen M, Nichols Thomas E. Permutation inference for the general linear model *Neuroimage.* 2014;92:381–397.

75. Smith Stephen M, Nichols Thomas E. Threshold-free cluster enhancement: addressing problems of smoothing, threshold dependence and localisation in cluster inference *Neuroimage.* 2009;44:83–98.

76. Salimi-Khorshidi Gholamreza, Smith Stephen M, Nichols Thomas E. Adjusting the effect of nonstationarity in cluster-based and TFCE inference *Neuroimage.* 2011;54:2006–2019.

77. Wilson SA Kinnier. Modern problems in neurology *British Medical Journal.* 1928;2:914.

78. Lazarus Michael, Huang Zhi-Li, Lu Jun, Urade Yoshihiro, Chen Jiang-Fan. How do the basal ganglia regulate sleep–wake behavior? *Trends in neurosciences.* 2012;35:723–732.

79. Lazarus Michael, Chen Jiang-Fan, Urade Yoshihiro, Huang Zhi-Li. Role of the basal ganglia in the control of sleep and wakefulness *Current opinion in neurobiology.* 2013;23:780–785.

80. Gerfen Charles R, Bolam J Paul. The neuroanatomical organization of the basal ganglia in *Handbook of behavioral neuroscience*;24:3–32Elsevier 2016.

81. Zheng Zhong S, Monti Martin M. Cortical and thalamic connections of the human globus pallidus: Implications for disorders of consciousness *Frontiers in Neuroanatomy.* 2022;16:960439.

82. Menza Matthew, Dobkin Roseanne DeFronzo, Marin Humberto, Bienfait Karina. Sleep disturbances in Parkinson's disease *Movement Disorders.* 2010;25:S117–S122.

83. Lima Marcelo MS. Sleep disturbances in Parkinson's disease: the contribution of dopamine in REM sleep regulation *Sleep medicine reviews.* 2013;17:367–375.

84. Lee Bi-Fang, Chiu Nan-Tsing, Yang Yen Kuang, Chu Ching Lin. The relation between striatal dopamine D2/D3 receptor availability and sleep quality in healthy adults *Nuclear Medicine Communications.* 2007;28:401–406.

85. Takeuchi Hikaru, Taki Yasuyuki, Nouchi Rui, et al. Shorter sleep duration and better sleep quality are associated with greater tissue density in the brain *Scientific reports.* 2018;8:5833.

86. Corsi-Cabrera M, Grinberg-Zylberbaum J, Arditti LS. Caudate nucleus lesion selectively increases paradoxical sleep episodes in the rat *Physiology & behavior.* 1975;14:7–11.

87. Mena-Segovia Juan, Cintra León, Prospéro-García Oscar, Giordano Magda. Changes in sleep–waking cycle after striatal excitotoxic lesions *Behavioural brain research.* 2002;136:475–481.

88. Qiu Mei-Hong, Vetrivelan Ramalingam, Fuller Patrick M, Lu Jun. Basal ganglia control of sleep–wake behavior and cortical activation *European Journal of Neuroscience.* 2010;31:499–507.

89. Albin Roger L, Young Anne B, Penney John B. The functional anatomy of basal ganglia disorders *Trends in neurosciences.* 1989;12:366–375.

90. Alexander Garrett E, Crutcher Michael D. Functional architecture of basal ganglia circuits: neural substrates of parallel processing *Trends in neurosciences.* 1990;13:266–271.

91. Nambu Atsushi, Tokuno Hironobu, Takada Masahiko. Functional significance of the cortico–subthalamo–pallidal 'hyperdirect'pathway *Neuroscience research.* 2002;43:111–117.

92. Qiu Mei-Hong, Zhong Zhi-Gang, Chen Michael C, Lu Jun. Nigrostriatal and mesolimbic control of sleep–wake behavior in rat *Brain Structure and Function.* 2019;224:2525–2535.

93. Qiu Mei-Hong, Yao Qiao-Ling, Vetrivelan Ramalingam, Chen Michael C, Lu Jun. Nigrostriatal dopamine acting on globus pallidus regulates sleep *Cerebral cortex.* 2016;26:1430–1439.

94. Qiu MH, Chen Michael C, Wu Jianping, Nelson Dwight, Lu Jun. Deep brain stimulation in the globus pallidus externa promotes sleep *Neuroscience.* 2016;322:115–120.

95. Crone Julia Sophia, Lutkenhoff Evan Scott, Bio Branden Joseph, Laureys Steven, Monti Martin Max. Testing proposed neuronal models of effective connectivity within the cortico-basal ganglia-thalamo-cortical loop during loss of consciousness *Cerebral Cortex.* 2017;27:2727–2738.

96. Castillo Pablo R, Middlebrooks Erik H, Grewal Sanjeet S, et al. Globus pallidus externus deep brain stimulation treats insomnia in a patient with Parkinson disease in *Mayo Clinic Proceedings*;95:419–422Elsevier 2020.

97. Saper Clifford B, Fuller Patrick M, Pedersen Nigel P, Lu Jun, Scammell Thomas E. Sleep state switching *Neuron.* 2010;68:1023–1042.

98. Whyte Christopher J., Redinbaugh Michelle J., Shine James M., Saalmann Yuri B.. Thalamic contributions to the state and contents of consciousness *Neuron.* 2024;112:1611-1625.

99. Liu Chun-Hong, Liu Cun-Zhi, Zhang Jihui, et al. Reduced spontaneous neuronal activity in the insular cortex and thalamus in healthy adults with insomnia symptoms *Brain research.* 2016;1648:317–324.

100. Li Min, Wang Ruonan, Zhao Meng, et al. Abnormalities of thalamus volume and resting state functional connectivity in primary insomnia patients *Brain Imaging and Behavior.* 2019;13:1193–1201.

101. Zou Guangyuan, Li Yuezhen, Liu Jiayi, et al. Altered thalamic connectivity in insomnia disorder during wakefulness and sleep *Human brain mapping.* 2021;42:259–270.

102. Ma Xueying, Zhang Yan, Ma Shaohui, et al. Association between abnormal thalamic metabolites and sleep disturbance in patients with end-stage renal disease *Metabolic brain disease.* 2018;33:1641–1648.

103. Wu Wei, Cui Linyang, Fu Ying, et al. Sleep and cognitive abnormalities in acute minor thalamic infarction *Neuroscience bulletin.* 2016;32:341–348.

104. Li Q, Zhao X, Gong DH, Geng YM, Zhang HL, Bi PX. Sleep disorders of acute thalamic stroke and its influence on plasma IL-17. *Journal of Biological Regulators and Homeostatic Agents.* 2017;31:745–751.

105. Liu Chunlei, Kong Xiang-zhen, Liu Xueyong, Zhou Renlai, Wu Bin. Long-term total sleep deprivation reduces thalamic gray matter volume in healthy men *Neuroreport.* 2014;25:320–323.

106. Poulin Stéphane P, Dautoff Rebecca, Morris John C, et al. Amygdala atrophy is prominent in early Alzheimer's disease and relates to symptom severity *Psychiatry Research: Neuroimaging.* 2011;194:7–13.

107. Zhang Rui, Tomasi Dardo, Manza Peter, et al. Sleep disturbances are associated with cortical and subcortical atrophy in alcohol use disorder *Translational psychiatry.* 2021;11:428.

108. Yu Honghui, Chen Liting, Li Haijun, et al. Abnormal resting-state functional connectivity of amygdala subregions in patients with obstructive sleep apnea *Neuropsychiatric disease and treatment.* 2019;977–987.

109. Burdová K, Filip P, Šonka K, et al. Hippocampal and amygdala atrophy rates correlate with the severity of sleep apnoea in Parkinson's disease *Parkinsonism & Related Disorders.* 2023;113.

110. Gong Liang, Liao Tianzhi, Liu Duan, et al. Amygdala changes in chronic insomnia and their association with sleep and anxiety symptoms: insight from shape analysis *Neural plasticity.* 2019;2019.

111. Zhang Liwei, Shao Yongcong, Jin Xueguang, Cai Xiaoping, Du Feng. Decreased effective connectivity between insula and anterior cingulate cortex during a working memory task after prolonged sleep deprivation *Behavioural Brain Research.* 2021;409:113263.

112. Chen Lingling, Qi Xueliang, Zheng Jiyong. Altered regional cortical brain activity in healthy subjects after sleep deprivation: a functional magnetic resonance imaging study *Frontiers in Neurology.* 2018;9:588.

113. Li Chao, Dong Mengshi, Yin Yi, Hua Kelei, Fu Shishun, Jiang Guihua. Aberrant effective connectivity of the right anterior insula in primary insomnia *Frontiers in Neurology.* 2018;9:317.

114. Liu Chun-Hong, Guo Jing, Lu Shun-Li, et al. Increased salience network activity in patients with insomnia complaints in major depressive disorder *Frontiers in Psychiatry.* 2018;9:93.

115. Yin Huazhan, Zhang Li, Li Dan, Xiao Lu, Cheng Mei. The gray matter volume of the right insula mediates the relationship between symptoms of depression/anxiety and sleep quality among college students *Journal of health psychology.* 2021;26:1073–1084.

116. Wu Huawang, Zheng Yingjun, Zhan Qianqian, et al. Covariation between spontaneous neural activity in the insula and affective temperaments is related to sleep disturbance in individuals with major depressive disorder *Psychological Medicine.* 2021;51:731–740.

117. Hidese Shinsuke, Ota Miho, Matsuo Junko, et al. Association between the Pittsburgh sleep quality index and white matter integrity in healthy adults: a whole-brain magnetic resonance imaging study *Sleep and Biological Rhythms.* 2023;21:249–256.

118. Namsrai Tergel, Ambikairajah Ananthan, Cherbuin Nicolas. Poorer sleep impairs brain health at midlife *Scientific Reports.* 2023;13:1874.

119. Chung Min-Huey, Liu Wen-I, Lee Hui-Ling, Hsu Nanly. Selected neurophysiological, psychological, and behavioral influences on subjective sleep quality in nurses: a structure equation model *PloS one.* 2013;8:e79529.

120. Moussa-Chamari Imen, Farooq Abdulaziz, Romdhani Mohamed, et al. The relationship between quality of life, sleep quality, mental health, and physical activity in an international sample of college students: A structural equation modeling approach *Frontiers in Public Health.* 2024;12:1397924.

121. Andersen Thea Otte, Sejling Christoffer, Jensen Andreas Kryger, et al. Nighttime smartphone use, sleep quality, and mental health: investigating a complex relationship *Sleep.* 2023;46:zsad256.

122. Xu Selene, Thompson Wesley, Ancoli-Israel Sonia, Liu Lianqi, Palmer Barton, Natarajan Loki. Cognition, quality-of-life, and symptom clusters in breast cancer: using Bayesian networks to elucidate complex relationships *Psycho-oncology.* 2018;27:802–809.

123. Manzar Md Dilshad, Zannat Wassilatul, Moiz Jamal Ali, et al. Factor scoring models of the Pittsburgh sleep quality index: a comparative confirmatory factor analysis *Biological Rhythm Research.* 2016;47:851–864.

124. Manzar Md Dilshad, BaHammam Ahmed S, Hameed Unaise Abdul, et al. Dimensionality of the Pittsburgh Sleep Quality Index: a systematic review *Health and quality of life outcomes.* 2018;16:1–22.

125. Liu Dan, Kahathuduwa Chanaka, Vazsonyi Alexander T. The Pittsburgh Sleep Quality Index (PSQI): Psychometric and clinical risk score applications among college students. *Psychological Assessment.* 2021;33:816.

126. Monti Martin M. Statistical analysis of fMRI time-series: a critical review of the GLM approach *Frontiers in human neuroscience.* 2011;5:28.

127. Fuller Patrick, Sherman David, Pedersen Nigel P, Saper Clifford B, Lu Jun. Reassessment of the structural basis of the ascending arousal system *Journal of Comparative Neurology.* 2011;519:933–956.

128. Benarroch Eduardo E. Parabrachial nuclear complex: multiple functions and potential clinical implications *Neurology.* 2016;86:676–683.

129. Castillo Pablo R, Benarroch Eduardo E. How could the basal ganglia control sleep? *Neurology.* 2020;95:302–304.

130. Zheng Zhong Sheng, Reggente Nicco, Monti Martin M. Arousal Regulation by the External Globus Pallidus: A New Node for the Mesocircuit Hypothesis *Brain Sciences.* 2023;13:146.

131. Yin Zixiao, Jiang Yin, Merk Timon, et al. Pallidal activities during sleep and sleep decoding in dystonia, Huntington's, and Parkinson's disease *Neurobiology of Disease.* 2023;182:106143.

132. Steriade M. Thalamic origin of sleep spindles: Morison and Bassett (1945) *Journal of neurophysiology.* 1995;73:921–922.

133. Steriade Mircea. Sleep, epilepsy and thalamic reticular inhibitory neurons *Trends in neurosciences.* 2005;28:317–324.

134. Crunelli Vincenzo, Hughes Stuart W. The slow (¡ 1 Hz) rhythm of non-REM sleep: a dialogue between three cardinal oscillators *Nature neuroscience.* 2010;13:9–17.

135. Schreiner Thomas, Kaufmann Elisabeth, Noachtar Soheyl, Mehrkens Jan-Hinnerk, Staudigl Tobias. The human thalamus orchestrates neocortical oscillations during NREM sleep *Nature communications.* 2022;13:5231.

136. Bonjean Maxime, Baker Tanya, Bazhenov Maxim, Cash Sydney, Halgren Eric, Sejnowski Terrence. Interactions between core and matrix thalamocortical projections in human sleep spindle synchronization *Journal of Neuroscience.* 2012;32:5250–5263.

137. Saper Clifford B, Fuller Patrick M. Wake–sleep circuitry: an overview *Current opinion in neurobiology.* 2017;44:186–192.

138. Reddy Oliver Cameron, Werf Ysbrand D. The sleeping brain: harnessing the power of the glymphatic system through lifestyle choices *Brain sciences.* 2020;10:868.

139. Jamieson Daniel, Broadhouse Kathryn M, McLoughlin Larisa T, et al. Investigating the association between sleep quality and diffusion-derived structural integrity of white matter in early adolescence *Journal of Adolescence.* 2020;83:12–21.

140. Siow Tiing Yee, Toh Cheng Hong, Hsu Jung-Lung, et al. Association of sleep, neuropsychological performance, and gray matter volume with glymphatic function in community-dwelling older adults *Neurology.* 2022;98:e829–e838.

141. Shih Nien-Chu, Barisano Giuseppe, Lincoln Karen D, Mack Wendy J, Sepehrband Farshid, Choupan Jeiran. Effects of sleep on brain perivascular space in a cognitively healthy population *Sleep Medicine.* 2023;111:170–179.

142. Klumpp Heide, Hosseini Bobak, Phan K Luan. Self-reported sleep quality modulates amygdala resting-state functional connectivity in anxiety and depression *Frontiers in Psychiatry.* 2018;9:220.

143. Amorim Liliana, Magalhães Ricardo, Coelho Ana, et al. Poor sleep quality associates with decreased functional and structural brain connectivity in normative aging: a MRI multimodal approach *Frontiers in aging neuroscience.* 2018;10:375.

144. Dai Xi-Jian, Peng De-Chang, Gong Hong-Han, et al. Altered intrinsic regional brain spontaneous activity and subjective sleep quality in patients with chronic primary insomnia: a resting-state fMRI study *Neuropsychiatric disease and treatment.* 2014:2163–2175.

145. Komarov Oleksii, Ko Li-Wei, Jung Tzyy-Ping. Associations among emotional state, sleep quality, and resting-state EEG spectra: a longitudinal study in graduate students *IEEE Transactions on Neural Systems and Rehabilitation Engineering.* 2020;28:795–804.

146. Venkataraman Arun, Zhuang Yuchuan, Marsella Jennifer, et al. Functional MRI correlates of sleep quality in HIV *Nature and science of sleep.* 2021:291–301.

147. He Dongmei, Ren Dongmei, Guo Zhiwei, Jiang Binghu. Insomnia disorder diagnosed by resting-state fMRI-based SVM classifier *Sleep Medicine.* 2022;95:126–129.

148. Picchioni Dante, Pixa Morgan L, Fukunaga Masaki, et al. Decreased connectivity between the thalamus and the neocortex during human nonrapid eye movement sleep *Sleep.* 2014;37:387–397.

149. Cain Joshua A, Visagan Shakthi, Johnson Micah A, et al. Real time and delayed effects of subcortical low intensity focused ultrasound *Scientific reports.* 2021;11:6100.

150. Cain Joshua A, Spivak Norman M, Coetzee John P, et al. Ultrasonic thalamic stimulation in chronic disorders of consciousness *Brain Stimulation: Basic, Translational, and Clinical Research in Neuromodulation.* 2021;14:301–303.

151. Dell'Italia John, Sanguinetti Joseph L, Monti Martin M, Bystritsky Alexander, Reggente Nicco. Current state of potential mechanisms supporting low intensity focused ultrasound for neuromodulation *Frontiers in human neuroscience.* 2022;16:872639.

Supplementary Online Materials for
The subcortical correlates of self-reported sleep quality

Contents

Table SOM1. Sample demographic and PSQI information broken down by sex. (Abbreviations: Med., median; sd., standard deviation; PC1, PSQI Principal Component 1; PC2, PSQI Principal Component 2; subj. sleep qual., subjective sleep quality; sleep dur., sleep duration; sleep eff., sleep efficiency; sleep ons. lat., sleep onset latency; sleep disturb., sleep disturbances; sleep med., use of sleep medications; daytime dysf., daytime dysfunction; U_{MW}, Mann-Whitney U test for independent samples; r_{rb}, rank-biserial correlation. Significance level of the Mann-Whitney test between female and male: *** indicates p<0.001; ** indicates p<0.01; * indicates p<0.05. See Tab. 1 for sample-wide statistics.)

Variables	Female (N=605)			Male (N=507)			U_{MW}	r_{rb}
	Med	Mean	sd	Med	Mean	sd		
PSQI Total Score	4	4.942	2.933	4	4.639	2.528	147,234	-0.04
Subj. sleep qual. (PC1)	1	0.899	0.66	1	0.888	0.604	153,910	0.004
Sleep dur. (PC1)	0	0.559	0.817	0	0.582	0.814	156,488	0.02
Sleep eff. (PC1)	0	0.534	0.846	0	0.363	0.719	137209***	-0.105
Sleep ons. lat. (PC2)	1	0.975	0.833	1	0.957	0.799	152,612	-0.005
Sleep disturb. (PC2)	1	1.129	0.492	1	1.049	0.474	143149**	-0.067
Sleep med. (PC2)	0	0.261	0.717	0	0.209	0.625	149,911	-0.023
Daytime dysf. (−)	1	0.585	0.642	1	0.592	0.626	154,947	0.01
Age	30	29.55	3.6	28	27.901	3.611	114060***	-0.256
Handedness	80	70.388	43.761	75	60.424	44.17	116345***	-0.241
Education	16	15.015	1.815	16	14.822	1.776	140547*	-0.08
BMI	24.92	26.279	5.785	26.11	26.836	4.388	173375***	0.132

Table SOM2. Correlations between the PSQI components. (Abbreviations: τ_B, Kendall's Tau B; Fisher's z (SE), standard error of the Fisher's z effect size. All other abbreviations are as in SOM1.)

		PSQI_Comp1 Subj. sleep qual.	PSQI_Comp2 Sleep ons. lat.	PSQI_Comp3 Sleep dur.	PSQI_Comp4 Sleep eff.	PSQI_Comp5 Sleep disturb.	PSQI_Comp6 Sleep med.	PSQI_Comp7 Daytime dysf.
PSQI_Comp1 Subj. sleep qual.	τ_B	—						
	p-value	—						
	Fisher's z	—						
	Fisher's z (SE)	—						
PSQI_Comp2 Sleep ons. lat.	τ_B	0.2818 ***	—					
	p-value	2.489×10^{-25}	—					
	Fisher's z	0.2896	—					
	Fisher's z (SE)	0.01825	—					
PSQI_Comp3 Sleep dur	τ_B	0.325 ***	0.0803 **	—				
	p-value	3.494×10^{-32}	0.002883	—				
	Fisher's z	0.3372	0.08047	—				
	Fisher's z (SE)	0.01766	0.01988	—				
PSQI_Comp4 Sleep eff.	τ_B	0.2576 ***	0.1881 ***	0.3573 ***	—			
	p-value	1.498×10^{-20}	4.225×10^{-12}	2.092×10^{-38}	—			
	Fisher's z	0.2636	0.1903	0.3738	—			
	Fisher's z (SE)	0.01854	0.01923	0.01718	—			
PSQI_Comp5 Sleep disturb.	τ_B	0.2978 ***	0.2465 ***	0.09726 ***	0.1821 ***	—		
	p-value	2.968×10^{-26}	3.230×10^{-19}	4.998×10^{-4}	9.840×10^{-11}	—		
	Fisher's z	0.3071	0.2517	0.09757	0.1841	—		
	Fisher's z (SE)	0.01804	0.01866	0.01981	0.01928	—		
PSQI_Comp6 Sleep med.	τ_B	0.1293 ***	0.1856 ***	0.01306	0.06918 *	0.08825 **	—	
	p-value	4.621×10^{-6}	1.838×10^{-11}	0.6416	0.0144	0.002068	—	
	Fisher's z	0.13	0.1878	0.01306	0.06929	0.08848	—	
	Fisher's z (SE)	0.01965	0.01925	0.02002	0.01992	0.01985	—	
PSQI_Comp7 Daytime dysf.	τ_B	0.226 ***	0.08373 **	0.1358 ***	0.08785 **	0.2039 ***	0.03508	—
	p-value	7.805×10^{-16}	0.002294	1.114×10^{-6}	0.001768	8.119×10^{-13}	0.22	—
	Fisher's z	0.23	0.08393	0.1367	0.08807	0.2068	0.03509	—
	Fisher's z (SE)	0.01888	0.01987	0.01961	0.01985	0.01909	0.02	—

Table SOM3. Subcortical Results. Significant associations between variables included in the PSQI total score analysis and subcortical shape. See Fig. 4. For each cluster, volume, peak Z-score and MNI coordinates, hemisphere, and anatomical label are shown. (Abbreviations and symbols: Vol, volume; Coord, coordinates; Hem, hemisphere; L, left; R, right; '-' represents a negative association; '+' represents a positive association.)

Vol	Cluster Peak			
(mm^3)	Z-score	MNI Coord	Hem	Region
PSQI (-)				
1018	3.2	$-22, 2, -8$	L	Putamen
705	2.9	$10, 19, 10$	R	Caudate
646	3.5	$21, 4, -8$	R	Putamen
216	3.5	$-11, 19, -3$	L	Caudate
169	2.7	$17, 19, -3$	R	Caudate
71	2.3	$-28, -13, -4$	L	Putamen
PSQI × BMI (-)				
527	2.8	$33, -21, -12$	R	Hippocampus
488	2.7	$-18, 3, 4$	L	Pallidum
197	2.4	$17, 6, 2$	R	Pallidum
111	3.1	$-28, -24, -20$	L	Hippocampus
51	3.1	$-23, -10, -19$	L	Hippocampus
41	2.3	$15, 1, -5$	R	Pallidum
PSQI × Education (+)				
1118	3.5	$10, 17, 11$	R	Caudate
718	2.9	$13, -3, 12$	R	Thalamus
630	3.1	$10, -33, 3$	R	Thalamus
338	3.1	$-13, 12, 17$	L	Caudate
262	2.8	$20, 5, 0$	R	Pallidum
128	2.6	$24, -15, -3$	R	Pallidum
74	3.2	$20, -21, 20$	R	Caudate
PSQI × Handedness (-)				
1040	3.5	$-10, 3, 17$	L	Caudate
536	3.0	$8, 12, 12$	R	Caudate
460	3.3	$-23, -30, -14$	L	Hippocampus
328	3.2	$19, 18, 5$	R	Caudate
266	3.2	$-28, -13, -17$	L	Hippocampus
162	3.3	$12, 16, -10$	R	Amygdala
65	3.1	$29, -14, -14$	R	Hippocampus
77	3.0	$22, -23, -16$	R	Hippocampus
47	2.7	$7, 7, -9$	R	Amygdala

Table SOM4. Subcortical Results. Significant associations between variables included in the PSQI PC analysis and subcortical shape. See Fig. 4. (Abbreviations are the same as in Tab. SOM3.)

Vol. (mm^3)	Z-score	Cluster Peak MNI Coord	Hem	Region
PC1 (-)				
1187	3.5	$-24, 3, -8$	L	Putamen
421	3.7	$21, 5, -8$	R	Caudate
277	2.5	$11, 16, 14$	R	Caudate
180	2.4	$19, 19, 4$	R	Caudate
180	2.2	$-10, 19, 8$	L	Caudate
160	2.8	$-23, -10, 5$	L	Pallidum
63	2.4	$-15, 17, -4$	L	Caudate
35	2.5	$-28, -13, -4$	L	Putamen
PC1 × BMI (+)				
158	2.5	$-30, -1, -23$	L	Amygdala
PC2 × Age (-)				
977	3.4	$26, 17, -3$	R	Caudate
466	2.8	$-28, -39, -5$	L	Hippocampus
124	2.4	$18, -1, -23$	R	Amygdala
117	2.4	$-19, -8, -26$	L	Hippocampus
105	3.0	$22, -12, -13$	R	Amygdala
PC2 × BMI (-)				
1417	4.1	$19, -37, -4$	R	Hippocampus
1158	4.0	$-4, -14, -33$	–	Brainstem
760	3.3	$22, 20, -4$	R	Putamen
757	4.0	$-24, -14, -4$	L	Pallidum
717	3.6	$19, -9, -5$	R	Pallidum
602	4.5	$-7, -46, -48$	–	Brainstem
510	3.6	$-13, 0, 12$	L	Caudate
483	3.8	$17, 4, 23$	R	Caudate
424	3.1	$15, -5, 13$	R	Thalamus
358	3.0	$7, -24, -16$	–	Brainstem
349	3.2	$-23, 11, 7$	L	Putamen
333	3.0	$-16, 6, 21$	L	Caudate
302	2.7	$-17, -35, -1$	L	Thalamus
298	3.9	$23, -8, -20$	R	Hippocampus
276	3.2	$-27, -13, 10$	L	Putamen
257	4.2	$-30, -22, -22$	L	Hippocampus
248	3.3	$11, 22, -4$	R	Caudate
196	3.8	$-20, -37, -6$	L	Hippocampus
175	3.4	$-10, 17, -2$	L	Caudate
128	3.4	$-24, -13, -17$	L	Hippocampus
109	2.9	$-17, -3, -23$	L	Amygdala
92	2.7	$-25, -6, -14$	L	Amygdala
35	2.6	$-1, -48, -32$	–	Brainstem
PC2 × Education (+)				
346	2.8	$-30, 2, 1$	L	Putamen
PC2 × Handedness (-)				
610	3.6	$-10, 4, 18$	L	Caudate
516	3.3	$10, 15, 14$	R	Caudate
338	3.9	$-28, -12, -19$	L	Hippocampus
466	3.7	$-23, -34, -10$	L	Hippocampus
270	2.8	$19, 18, 8$	R	Caudate
266	2.5	$19, -33, 6$	R	Thalamus
231	2.4	$10, -1, 8$	R	Thalamus
147	3.7	$28, -12, -17$	R	Hippocampus
65	4.0	$21, -28, -11$	R	Hippocampus
39	2.7	$-18, -3, -16$	L	Amygdala
35	2.4	$-18, 16, 5$	L	Caudate
PC2 × Sex (-)				
2507	3.0	$10, -35, -28$	R	Brainstem
1580	3.1	$21, -10, -5$	R	Pallidum

558	3.0	$-15, 10, 19$	L	Caudate
371	2.7	$-23, 10, 6$	L	Putamen
260	2.9	$-15, 1, 2$	L	Pallidum
151	3.1	$-20, -9, -6$	L	Pallidum
117	3.0	$-16, -7, 17$	L	Caudate
107	3.5	$-18, 18, -3$	L	Caudate

Table SOM5. Cortical results. Significant associations between variables included in the PSQI PC analysis and cortical morphometry (note that no significant associations were found between cortical morphometry and any of the variables in the PSQI total score analysis). See Fig. 5. (Abbreviations are the same as in Tab. SOM3.)

Vol. (mm³)	Z-score	Cluster Peak MNI Coord	Hem	Region
PC1 × Age (+)				
14371	3.9	$-19, -0, -21$	L	Parahipp. Gyrus
1822	3.1	$35, 3, 4$	R	Rolandic Operc.
672	3.2	$-34, 20, -1$	L	Ant. Insula
220	2.7	$19, -52, -9$	R	Lingual Gyrus
116	2.6	$-48, 3, -12$	L	Sup. Temp. Pole
65	2.3	$33, -3, -11$	R	Ant. Insula
33	2.2	$-30, 8, 10$	L	Ant. Insula
32	1.7	$-4, -65, 14$	L	Occipitopar. Sulc.
PC1 × Age (−)				
370	4.0	$-17, -71, 48$	L	Sup. Par. Gyrus
349	4.3	$-33, -68, 44$	L	Intrapar. Sulc.
PC2 × Education (−)				
861	3.2	$33, 12, 3$	R	Ant. Insula
189	3.2	$3, 6, -13$	R	Ant. Insula
PC2 × Sex (+)				
3412	5.1	$-2, -7, 60$	L	Supp. Motor Area
852	4.1	$5, 18, 41$	R	Paracing. Gyrus

Fig. SOM1. PSQI and covariates correlation structure. Pairwise correlation (Kendall's tau (τ) between PSQI, PSQI PC1, PSQI PC2 and covariate measures (i.e., age, handedness, BMI, years of education). ('*' p ¡ 0.05; '**' p ¡ 0.01; '***' p ¡ 0.001).

PSQI Total Score Analysis

PSQI x Sex (-)

Fig. SOM2. Subcortical result for the PSQI × Sex contrast when including only one interaction in the analysis.