Gastrin Releasing Peptide Signaling in the Nucleus Accumbens Medial Shell Regulates Neuronal Excitability and Motivation

Erin E. Aisenberg[1], Hongli Wang[2], Atehsa A. Sahagun[1], Emilie M. Tu[2], and Helen S. Bateup[1,2,*]

[1]Helen Wills Neuroscience Institute, University of California, Berkeley, Berkeley, CA, USA

[2]Department of Molecular and Cell Biology, University of California, Berkeley, Berkeley, CA USA

*Correspondence to: bateup@berkeley.edu

Summary

Neuropeptides are the largest class of neuromodulators, which can be co-released together with classical neurotransmitters. It has been shown that subpopulations of dopamine neurons express mRNA for the neuropeptide Gastrin-releasing peptide (GRP); however, its functional relevance in dopaminergic circuits is unknown. Here, we find that the GRP receptor (GRPR) is present in the nucleus accumbens medial shell (NAc MSh), which is targeted by GRP-expressing midbrain dopamine neurons as well as glutamatergic inputs from the hippocampus and amygdala. We show that the NAc MSh GRPR-positive cells are a subpopulation of D2 receptor-expressing neurons, which have high intrinsic excitability and can be activated by GRP *in vivo*. Selective deletion of *Grpr* from the NAc MSh increases motivation in a progressive ratio test, indicating a role for GRPR in motivated behaviors. These experiments establish GRP/GRPR signaling as a novel regulator of mesolimbic circuits and advance our understanding of neuropeptides in the striatum.

Keywords

Gastrin-releasing peptide, GRP, GRPR, dopamine neurons, striatum, nucleus accumbens medial shell, intrinsic excitability, striatal projection neurons, neuropeptides, motivation, progressive ratio test

Acknowledgments

This work was supported by a Chan Zuckerberg Biohub Investigator grant to H.S.B. We thank Dr. Daniel Kramer for his contribution to the early phases of the project. We thank Dr. Hendrik Wildner and Dr. Hanns Ulrich Zeilhofer for sending us tissue samples for mouse validation. We thank Kamran Ahmed for help with schematics for figures. We thank Dr. Dirk Hockemeyer for help designing a genotyping strategy for the *Grpr* KO mice. Select confocal imaging experiments were conducted at the CRL Molecular Imaging Center, RRiD:SCR_017852, supported by the Gordon and Betty Moore Foundation. We thank Holly Aaron and Feather Ives for their microscopy advice and support. We thank Mahmoud Farhan for help with mouse colony maintenance and genotyping. We thank members of the Bateup lab for feedback on this project.

Author contributions

Conceptualization – E.E.A. and H.S.B.; Formal analysis – E.E.A. and H.W.; Funding acquisition – H.S.B.; Investigation – E.E.A., H.W., A.A.S., and E.M.T.; Methodology – E.E.A., H.W., and E.M.T.; Project administration – E.E.A. and H.S.B., Software – H.W.; Supervision – H.S.B.; Visualization – E.E.A.; Writing – original draft – E.E.A.; Writing – review & editing – E.E.A., H.W., A.A.S., E.M.T., and H.S.B.

Declaration of interests

The authors declare no competing interests.

Supplemental information

Supplementary Figures S1-S8.

Introduction

Neuromodulators play a critical role in nervous system activity via their regulation of neuronal excitability, synaptic transmission, and plasticity (Guillaumin and Burdakov, 2021; Nusbaum, 2017). The largest family of known neuromodulators are neuropeptides—small short-chain polypeptides that act via metabotropic G-protein-coupled receptors (GPCRs) (Castro and Bruchas, 2019). Compared to classical neurotransmitters, neuropeptides lead to diverse effects on their target neurons and regulate numerous physiological functions such as eating, sleeping, sexual behaviors, stress response, and pain perception (Hermes et al., 2013; Hokfelt et al., 2003). Additionally, neuropeptides have distinct machinery for their synthesis and transport and cannot be synthesized or recycled in axons (Russo, 2017). It has recently become appreciated that many neurons co-express genes for both neurotransmitters and one or more neuropeptides (Kramer et al., 2018; Smith et al., 2019). In particular, a recent single-cell RNA-sequencing (scRNA-seq) study found that 98% of mouse cortical neurons express at least one neuropeptide or neuropeptide receptor gene (Smith et al., 2019). As such, neuropeptides are poised to make a major contribution to brain function and behavior; however, an understanding of their breadth, scope and mechanisms of action is currently limited.

One such neuropeptide is Gastrin-releasing peptide (GRP), a bombesin-like neuropeptide that signals through a G_q-coupled GPCR (gastrin releasing peptide receptor, GRPR, also called BB2) (Moody et al., 1981; Moody and Merali, 2004). GRP was originally named for its role in stimulating gastrin release from the G cells in the stomach (Merali et al., 1999). In the central nervous system, GRP has been studied in the spinal cord in the context of itch and pain processing (Pagani et al., 2019; Sun and Chen, 2007; Yu et al., 2017). In the brain, several studies have demonstrated a role for GRP/GRPR signaling in emotionally-motivated and fear memories as well as responses to stress (Roesler et al., 2014; Roesler and Schwartsmann, 2012). For example, in the lateral nucleus of the amygdala, *Grp* provides negative feedback regulating fear (Shumyatsky et al., 2002). In this region, GRP excites the GABAergic interneurons that express GRPR, increasing their inhibition of the principle neurons expressing *Grp* (Shumyatsky et al., 2002). In the auditory cortex, *Grpr* is expressed in vasoactive intestinal polypeptide (VIP) neurons and GRP signaling recruits disinhibitory cortical microcircuits and enhances auditory fear memories (Melzer et al., 2021). At the cellular level, GRP wash-on in brain slices leads to depolarization, and in some cases rhythmic or tonic firing, in a variety of neuron types (Hermes et al., 2013; Lee et al., 1999; Piggins and Rusak, 1993; Pinnock and Woodruff, 1991).

A recent scRNA-seq study of mouse midbrain dopamine (DA) neurons found that 30% of midbrain DA neurons in the ventral tegmental area (VTA) and the substantia nigra pars compacta (SNc) express the mRNA for *Grp* (Kramer et al., 2018), consistent with prior studies demonstrating GRP immunoreactivity in the SNc (Larsen et al., 1989). Additionally, *Grp* has been identified as a differentially expressed gene between VTA and SNc DA neurons (Chung et al., 2005; Viereckel et al., 2016). *GRP*-expressing DA neurons have also been observed in post-mortem human brains and are suggested to be relatively resistant to degeneration in Parkinson's disease (Viereckel et al., 2016). When examining the projection targets of the *Grp*+ neurons, Kramer et al. (2018) found two populations of *Grp*+ DA neurons, one that resides in the ventral SNc and projects to the dorsal medial striatum (DMS) and the other that is in the medial VTA and targets the nucleus accumbens medial shell (NAc MSh) (Kramer et al., 2018). These findings suggest

that GRP may be co-released by DA neurons and may be a functionally relevant signaling molecule in the striatum.

The striatum is the main input structure of the basal ganglia, which integrates glutamatergic inputs from the cortex and thalamus, DA from the midbrain, GABAergic inhibition from multiple intra-striatal sources, and acetylcholine from cholinergic interneurons (Gerfen and Surmeier, 2011). GABAergic striatal projection neurons (SPNs) are the major output neuron of the striatum, comprising 90-95% of the neuronal population (Kemp and Powell, 1971). SPNs can be divided into direct-pathway and indirect-pathway neurons based on their primary projections to either the substantia nigra pars reticulata and globus pallidus internal segment (SNr/GPi) or globus pallidus external segment (GPe), respectively (Gerfen, 2000). Direct pathway SPNs (dSPNs) typically express type 1 DA receptors (D1) while indirect pathway SPNs (iSPNs) generally express type 2 DA receptors (D2) (Gerfen, 2000). The dorsal striatum (DS) is important for motor functions including action selection and motor learning, while the ventral striatum controls motivated behaviors and reward learning (Voorn et al., 2004). In the DS, activation of dSPNs with optogenetics promotes locomotor activity while bulk activation of iSPNs inhibits movement (Bateup et al., 2010; Kravitz et al., 2013). During normal movement, however, both pathways are active and must work in a coordinated manner (Cox and Witten, 2019; Gerfen and Bolam, 2016). The ventral striatum (nucleus accumbens, NAc) also contains direct- and indirect-pathway SPNs. In the NAc, the dSPNs have canonically been thought to directly project to and inhibit the VTA, promoting motivated behavior through disinhibition of VTA GABA neurons that synapse onto VTA DA neurons and project back to the NAc (Kupchik and Kalivas, 2017). NAc iSPNs first project to and inhibit the ventral pallidum (VP). VP neurons then project to the VTA and ultimately the thalamus, resulting in inhibition of motivated behavior (Kupchik and Kalivas, 2017).

With increasing use of cell type-specific approaches, it has become clear that our canonical divisions of the striatum: D1 vs D2 SPNs, dorsal vs ventral striatum, indirect vs direct pathway are oversimplified and there is likely further complexity to discover within these circuits (Kupchik et al., 2015; Puighermanal et al., 2020). While neuropeptide signaling has been known to occur in the striatum through opioid receptors, these receptors are all $G_{i/o}$-coupled, and the opioids themselves are synthesized within the striatum (Castro and Bruchas, 2019). In contrast, the *Grp+* DA neurons described above would bring in a neuropeptide from outside the striatum and are hypothesized to increase the excitability of GRPR-expressing neurons.

The goal of this study was to assess a possible role of GRP-GRPR signaling in the NAc. We found that *Grpr* is expressed in the NAc medial shell (MSh), mainly in D2R-expressing SPNs. While the genetic profile of *Grpr+* neurons in the NAc MSh was consistent with an iSPN identity, we found that they have unique electrophysiological properties. Specifically, a majority of *Grpr+* cells are spontaneously active in slice recordings, and they have high intrinsic excitability. We also found that in addition to the *Grp+* DA neurons, the NAc MSh receives glutamatergic *Grp+* inputs from the hippocampus and amygdala. We show that GRP activates and increases the excitability of GRPR neurons in the NAc MSh both in slice and *in vivo*. Finally, we show that specific deletion of *Grpr* in the NAc MSh increases motivation in a progressive ratio test. These findings identify a previously uncharacterized cell type in the NAc MSh and map out a functional GRP-GRPR circuit that mediates striatal excitability and motivated behaviors.

Results

Grpr is expressed in striatal neurons

To investigate whether GRP may be a functionally relevant neuromodulator in nigrostriatal and mesolimbic dopamine circuits, we used fluorescent *in situ* hybridization (FISH) to probe for expression of mRNA encoding *Grpr* in the DS and NAc MSh (Fig. 1A). Indeed, we found *Grpr* expression in subpopulations of cells in the DS and NAc MSh (Fig. 1B). These cells were concentrated in the anterior portion of the striatum, with the highest concentration of *Grpr+* cells found around A/P +1.18 mm from bregma (Fig. 1B). Our findings are consistent with prior autoradiographic studies showing a high density of binding sites for bomesin-like peptides in the NAc and striatum (Wolf et al., 1983; Zarbin et al., 1985).

To determine the identity of *Grpr*-expressing cells, we used multi-plex FISH with markers for known striatal cell types (Fig. 1C-L). In the NAc MSh, we found that the vast majority of *Grpr+* cells were positive for *Gad2* mRNA (96.29%) (Fig. 1D), and no *Grpr+* cells were *Chat+* (Fig. 1H). This suggests that *Grpr+* cells are GABA-ergic and do not express the primary marker for cholinergic interneurons, the most prevalent interneuron type in the NAc MSh (Castro and Bruchas, 2019). In the DS, we found similar expression patterns (Fig. S1). Here, 92.66% of *Grpr+* cells were *Gad2+* (Fig. S1A,B) compared to only 1.90% co-expressing *Chat* mRNA (Fig. S1C,F). To determine whether the *Grpr+* cells were SPNs or a type of GABAergic interneuron, we probed for *Ppp1r1b*, which encodes the SPN-marker DARPP-32 (Fig. 1E,I). Nearly all *Grpr*-expressing cells co-expressed *Ppp1r1b* in the NAc MSh (93.76%) (Fig. 1I) and DS (75.52%) (Fig. S1C,G), suggesting they were likely SPNs.

To determine the specific SPN identity of the *Grpr+* cells, we probed for mRNA encoding D1 (*Drd1*) and D2 (*Drd2*) DA receptors (Fig. 1F), which are typically expressed in dSPNs and iSPNs, respectively. The majority of *Grpr+* cells in the NAc MSh expressed *Drd2* mRNA (77.26%) (Fig. 1J), suggesting that they could be iSPNs. However, a small subpopulation of *Grpr+* cells in the NAc MSh expressed *Drd1* (26.71%) (Fig. 1K). Multiple studies have reported that a proportion (5-30%) of SPNs in the NAc MSh express both *Drd1* and *Drd2* (Al-Hasani et al., 2015; Bertran-Gonzalez et al., 2008; Kupchik et al., 2015); therefore, we examined whether the *Grpr+/Drd1+* population in the NAc MSh were dSPNs or SPNs co-expressing both DA receptors. We found that 61.51% of *Grpr+* cells in the NAc MSh expressed *Drd2*, but not *Drd1*, 14.10% expressed *Drd1* only, and 12.61% expressed both *Drd1* and *Drd2* (Fig. 1M). Given the preferential expression of *Drd2* in *Grpr+* cells, we probed for the iSPN marker *Penk* (Fig. 1G) and found that it was expressed by the majority (67.38%) of *Grpr+* cells in the NAc MSh (Fig. 1L). We also examined the co-localization between *Drd2* and *Penk* mRNA and found that 22.17% of the *Grpr+* cells in the NAc MSh expressed *Drd2*, but not *Penk*, 4.26% expressed *Penk*, but not *Drd2*, and *63.12%* of *Grpr+* cells in the NAc MSh expressed both markers (Fig. 1N). Together this analysis indicates that *Grpr+* cells in the NAc MSh are likely to be GABA-ergic D2R-expressing SPNs. We note that only a small percentage of all *Prpp1r1b* (1.54%), *Drd2* (3.94%), or *Penk*-expressing (2.79%) cells in the MSh were *Grpr* positive, indicating that *Grpr+* cells are a select subpopulation of SPNs in the NAc MSh (Fig. 1O-Q).

We next examined the identity of the *Grpr+* cells in the DS that we identified in Fig. 1B (Fig. S1). While a majority of *Grpr+* cells in the DS expressed SPN markers (Fig. S1C-J), the SPN sub-type

classification was less clear. *Drd2* mRNA was expressed in 32.74% of DS *Grpr+* cells, while *Drd1* mRNA was expressed in 43.28% of *Grpr+* cells, (Fig. S1H,I). We also found that *Penk* mRNA was found in 38.03% of DS *Grpr+* cells, consistent with the proportion expressing *Drd2* mRNA (Fig. S1J). Together this suggests that *Grpr* is expressed in both types of SPNs in the DS.

Given the more consistent D2R SPN identity of *Grpr+* cells in the NAc MSh and the fact that the NAc was previously identified as a region with high GRPR expression (Wolf et al., 1983; Zarbin et al., 1985), we focused further analysis on the *Grpr*-expressing cell population in the NAc MSh, which receives dopaminergic inputs from the VTA.

GRPR-expressing neurons in the NAc MSh are morphologically similar to neighboring SPNs
To study the properties of GRPR-expressing cells in the NAc MSh, we obtained a transgenic mouse line that fluorescently labels GRPR+ cells with eGFP (Fig. 2A-C). *Grpr*::eGFP mice (Gensat [MMRRC #036178-UCD](Gong et al., 2003)) have previously been shown to express *eGFP* exclusively in *Grpr+* cells (Pagani et al., 2019).

To characterize the morphology of GRPR+ cells in the NAc MSh, we filled cells with neurobiotin through a patch pipette and performed cellular reconstructions of eGFP+ and eGFP- cells (Fig. 2D,E). We found that GRPR+ cells had similar dendritic complexity as neighboring cells as measured by Sholl analysis (Fig. 2F). Additionally, we observed no differences in the total dendritic arbor length or the number of primary dendrites in eGFP+ vs eGFP- cells (Fig. 2G,H). We also examined dendritic spines in cells transduced with an mCherry-expressing virus and observed that GRPR+ cells were spiny and had a similar spine density as neighboring neurons (Fig. 2I-K). This analysis further indicates that GRPR+ cells in the NAc MSh are likely a subpopulation of SPNs, as striatal interneurons have few or no spines (Tepper et al., 2010).

GRPR+ neurons in the NAc MSh have high intrinsic excitability
We next assessed the functional and physiological properties of GRPR-expressing neurons in the NAc MSh using *Grpr*::eGFP;D1-tdTomato mice (Ade et al., 2011) (Fig. 3A-D). We found that consistent with our FISH results, only ~5% of eGFP+ cells in the NAc MSh were tdTomato+, indicating that the eGFP+ cells in the NAc MSh are likely D2R SPNs (Fig. 3C,D). By contrast, 81.21% of eGFP+ cells in the DS were D1-tdTomato positive (Fig. 3B,D).

We used whole-cell electrophysiology to record from *Grpr*-eGFP+ cells and compared them to neighboring *Grpr*-eGFP- neurons in the NAc MSh (Fig. 3E,F). We also recorded from genetically identified dSPNs (D1-tdTomato+, Fig. 3G) and presumed iSPNs that were GRPR negative (D1-tdTomato-;*Grpr*-eGFP-, Fig. 3H). Interestingly, we found that the baseline physiological properties of *Grpr*-eGFP+ cells differed significantly from other SPNs in the NAc MSh. Approximately 75% of the *Grpr*-eGFP+ cells were spontaneously active at baseline (Fig. 3E), which was not characteristic of eGFP- cells or other SPN subtypes (Fig. 3F-H). We also found that compared to neighboring cell populations, *Grpr*-eGFP+ neurons were significantly more depolarized (Fig. 3I), had higher membrane resistance (Fig. 3J), and had significantly wider action potentials (Fig. 3K). *Grpr*-eGFP+ cells also differed in their response to current injection compared to neighboring

SPNs, exhibiting increased action potential firing in response to small currents and a transition to depolarization block with higher currents (Fig. 3L).

Given that the *Grpr*-eGFP+ cells were spontaneously active and had properties that were distinct from other SPNs in the area, we assessed how they compared to cholinergic interneurons, which are known to be tonically active (Zhou et al., 2002) (Fig. S3A,B). We recorded from tdTomato-labeled neurons in *Chat*-Cre;Ai9 mice (Madisen et al., 2010; Rossi et al., 2011) and found that while the *Chat*-Cre+ cells tended to be spontaneously active at baseline (53%, Fig. S2B), their membrane potential, membrane resistance, and action-potential half-width were all significantly different from *Grpr*-eGFP+ cells (Fig. S2C-E). Additionally, with increasing positive current injection, the *Chat*+ cells were able to continue to fire action potentials, while the *Grpr*-eGFP+ cells went into depolarization block with +60 pA current injection (Fig. S2F). This, together with the fact that we did not find any co-localization between *Chat* and *Grpr* mRNA (see Fig. 1H), rules out the possibility that GRPR+ cells are cholinergic interneurons.

Multiple inputs to the NAc MSh express Grp mRNA
While it was previously established that *Grp*+ DA neurons project to the NAc MSh from the VTA (Kramer et al., 2018), the NAc MSh receives inputs from a number of other brain regions including the hippocampus, amygdala, thalamus, entorhinal cortex, and others (Li et al., 2018). Many of these brain areas have been reported to express the mRNA for *Grp* (Shumyatsky et al., 2002). Therefore, we asked whether other inputs into the NAc MSh might also have the potential to co-release GRP. To investigate this, we injected retrobeads into the NAc MSh and performed FISH for *Grp* in the ten primary brain regions that send projections to the NAc MSh (Fig. 4A,B). Notably, we found a high percentage of retrobead+ cells that were also *Grp*+ in the VTA, hippocampus (subiculum and CA1), and basolateral amygdala (BLA) (Fig. 4C,D). We found little to no expression of *Grp* in other regions projecting to the NAc MSh, including the lateral entorhinal cortex, anterior piriform cortex, and thalamus, amongst others (Fig. 4C,D). This suggests that a select group of afferents to the NAc MSh could have the capacity to co-release GRP.

We summed together all the *Grp*+ inputs quantified and found that the main sources of *Grp* into the NAc MSh were from the subiculum (51%), CA1 (34%), VTA (6%), and BLA (5%) (Fig. 4D-H). We further probed these four regions for *Vglut* (excitatory neurons) and *Gad2* (inhibitory neurons) and found that in CA1, the subiculum, and the BLA, nearly all the *Grp* inputs were glutamatergic (Fig. 4I-K). This suggests that in addition to the dopaminergic inputs, several different glutamatergic inputs to the NAc MSh may have the ability to use GRP as a neuromodulator.

GRP depolarizes and increases the excitability of GRPR-expressing neurons
After establishing that GRPR+ cells in the NAc MSh have distinct electrophysiological properties and mapping the *Grp*+ inputs, we sought to determine how GRP affects the functional properties of these cells. To do this, we washed on GRP to NAc MSh slices and measured membrane potential and resistance in *Grpr*-eGFP+ cells with current clamp recordings. We found that 300 nM GRP caused an approximately 6 mV depolarization in *Grpr*-eGFP+ cells and an average 140% increase in membrane resistance (Fig. 5A,C,D), consistent with the effects of GRP in other cell types (Hermes et al., 2013; Melzer et al., 2021; Pagani et al., 2019). Importantly, these

changes were blocked in the presence of the GRPR-specific antagonist, DPDMB (Fig. 5B,E,F). As a control, we performed the same experiment on D2-GFP+ cells in D2-GFP mice (Gong et al., 2003), the majority of which are not expected to express GRPR (see Fig. 1P). We did not detect any change in membrane potential or resistance with GRP wash-on (Fig. 5G,I,J). While co-application of DPDMB did not affect membrane potential in D2-GFP+ cells (Fig. 5H,K), we did observe a slight decrease in membrane resistance in the presence of the antagonist in this experiment (Fig. 5L).

We observed that the *Grpr*-eGFP+ cells had high baseline excitability (see Fig. 3) and activation of GRPR increased excitability; therefore, we asked whether inhibition of GRPR would affect the baseline properties of *Grpr*-eGFP+ neurons. We compared the membrane potential and resistance of *Grpr*-eGFP+ and D2-GFP+ cells with and without DPDMB, prior to the addition of GRP. We found a small decrease in the membrane resistance of *Grpr*-eGFP+ cells treated with DPDMB compared to untreated cells, which did not occur in D2-GFP+ cells (Fig. S3A-D). This suggests that GRPR either has basal activity that can be blocked with DPDMB or that there may be tonic GRP-GRPR signaling in the slice preparation that increases the excitability of *Grpr*-eGFP+ cells.

GRPR+ neurons in the NAc MSh are more likely to be Fos+ in vivo
Given the high excitability of *Grpr*-eGFP+ cells in our slice recordings, we examined whether they may be more active *in vivo*. To assess this, we performed FISH in naïve wild-type (WT) mice and probed for *Drd2, Grpr*, and *Fos* mRNA. We assessed the percentage of *Drd2*+ cells in the NAc MSh that were *Fos*+ under baseline conditions, using *Fos* as a proxy for neuronal activity. We found that *Drd2*+/*Grpr*+ cells were over twice as likely to be *Fos*+ compared to neighboring *Drd2*+/*Grpr*- cells (18.73% compared to 8.60% *Fos*+) (Fig. 6A,B). This suggests that expression of *Grpr* boosts the excitability of SPNs such that they are more likely to be activated *in vivo*.

To probe whether the increased excitability seen both in slice and *in vivo* was due to GRPR activation or some other mechanism unique to *Grpr*-eGFP+ cells, we generated *Grpr* knockout (KO) mice. To do this, we obtained floxed *Grpr* (*Grpr*^fl^) mice (Yu et al., 2017) and bred them to CMV-Cre mice (Schwenk et al., 1995) to induce constitutive deletion of *Grpr* throughout the body. Figure S4A-D shows PCR genotyping results identifying mice with exon 2 deletion and WT controls. We verified that the *Grpr* FISH probe binding site was not located within exon 2, and thus could be used to identify cells that normally expressed *Grpr*, even if the gene was no longer functional (we term all cells that typically express *Grpr* as "*Grpr*-lineage cells"). We first examined whether there was any difference in the percentage of *Grpr*-lineage cells that were *Fos*+ between WT and *Grpr* KO mice (Fig. S4E-G). We found that disrupting *Grpr* expression significantly reduced the percentage of *Grpr*-lineage cells that were *Fos*+ in naïve mice (Fig. S4E). Furthermore, in *Grpr* KO mice, there was no longer a significant difference in the percentage of *Fos*+ cells between *Drd2*+*;Grpr*-lineage cells and *Drd2*+*;Grpr*- cells (Fig. 6C-D), suggesting that expression of *Grpr* is necessary for this effect. Together, these results indicate that *Grpr*+ cells in the NAc MSh are more active *in vivo*, at least in part due to GRPR signaling.

GRP injection in vivo increases Fos expression
We next asked whether GRP application would further increase the activity of GRPR+ cells and/or have broader network effects. We injected 3 μM GRP unilaterally into the NAc MSh of anesthetized WT mice and harvested the brains 45 minutes later (Fig. 6E) (Melzer et al., 2021). We performed FISH and probed for *Drd2, Grpr,* and *Fos* mRNA and compared the NAc MSh of the injected hemisphere to the uninjected hemisphere (Fig. 6F,G). We found an increase in the total number of *Fos*+ cells in the GRP-injected hemisphere compared to the control, indicating an increase in activated cells upon GRP injection (Fig. 6H). The total number of *Grpr*+ cells was unchanged, as expected (Fig. 6I). Importantly, *Fos* was induced in *Grpr*+ cells as we found an increase in the total number of *Grpr*+ cells that were *Fos*+ as well as an increase in the proportion of *Grpr*+ cells that were *Fos*+ following GRP injection (Fig. 6J,K). Given that *Fos* is a proxy for neuronal activity, these data imply that GRP increases the activity of *Grpr*+ cells in the NAc MSh. We also observed that while the percentage of *Fos*+ cells that were *Drd2*+/*Grpr*+ increased (Fig. 6L), the percentage *Fos*+ cells that were *Drd2*+/*Grpr*- did not change (Fig. 6M), showing that the GRP-induced activation was selective to GRPR-expressing cells and did not induce larger network-level activation.

As a control to verify that this effect was induced by GRP and not due to the injection itself, we repeated the above experiment, but injected sterile saline instead of GRP (Fig. S5). While we saw a small increase in the total number of *Fos*+ cells in the saline-injected hemisphere, there was no increase in the total number of *Grpr*+ cells or the *Grpr*+ cells that were *Fos*+ (Fig. S5A-E). Additionally, we saw no increase in the percentage of *Grpr*+ cells that were *Fos*+ (Fig. S5F). Together, these results indicate that while saline injection can acutely increase neuronal activity in the NAc MSh, it does not preferentially activate *Grpr*+ cells.

Selective deletion of Grpr in the NAc MSh increases motivation in the progressive ratio test
Given that GRP-GRPR signaling had a strong effect on modulating the excitability of a subset of NAc MSh neurons, we tested whether NAc MSh-specific deletion of *Grpr* affected striatal-dependent behaviors. To generate conditional *Grpr* knock-out (cKO) mice, we injected *Grpr*[fl/fl] or [fl/y] mice (Yu et al., 2017) bilaterally with an AAV expressing CRE recombinase, to specifically delete *Grpr* in the NAc MSh (Fig. 7A,B). A GFP-expressing virus was used as a control. *Grpr* cKO mice and GFP-injected control littermates were then run through the open field test, accelerating rotarod, progressive ratio (PR) lever-pressing task, and sucrose preference test (Fig. S6A) to assess motor and reward-related behaviors.

The open field assay was used to measure general locomotor behavior and was analyzed using DeepLabCut (Fig. S6B) (Mathis et al., 2018). We found that *Grpr* cKO mice travelled a greater total distance during the one-hour session (Fig. S6C), suggesting a mild increase in locomotor activity. We assessed the time spent in the center, as a proxy for avoidance behavior, and number of grooming bouts, as a measure of repetitive behavior, and found no differences (Fig. S6D,E). We used Keypoint-MoSeq (Weinreb et al., 2023; Wiltschko et al., 2020) to perform unbiased identification of spontaneous behavior syllables used by the mice in the open field. The behavior syllables detected by this approach included movements like forward run, right or left turn, and grooming (Fig. S6F,G). We found no significant differences between groups in their usage of any

of the top 28 most frequent syllables (Fig. S6F,G). Together, this suggests that disruption of GRPR signaling specifically in the NAc MSh does not have a major impact on general mouse behavior patterns.

Both the DS and ventral striatum have been implicated in motor learning. To examine whether *Grpr* deletion impacts motor coordination or motor routine learning, we tested the mice on the accelerating rotarod with three trials per day across four consecutive testing days. This task is known to require plasticity in striatal circuits, involving both the dorsal and ventral striatum (Cording and Bateup, 2023; Costa et al., 2004; Durieux et al., 2012; Platt et al., 2017; Rothwell et al., 2014; Yin et al., 2009). We saw no differences in motor coordination or learning between *Grpr* cKO mice and GFP-injected controls (Fig. S7A-D).

The NAc is a core component of the mesolimbic reward processing pathway. Alterations in NAc activity can therefore affect the hedonic value of natural rewards (Berridge and Kringelbach, 2015; Klawonn and Malenka, 2018). To test whether *Grpr* signaling is involved in this, we compared *Grpr* cKO and control mice in the sucrose preference test (Fig. S7E). This test is a measure of hedonia, or the ability to derive pleasure from rewarding stimuli (Bessa et al., 2013; Hurley and Carelli, 2020; Martínez-Hernández et al., 2012). We first verified there was no difference in starting weight between groups (Fig. S7F). During the test sessions, we found that *Grpr* cKO and control mice showed a similar preference for the sucrose solution over plain water (Fig. S7G). These results suggest that GRPR signaling in the NAc MSh is not required for hedonic behavior and reward value processing.

In addition to processing reward, the NAc is implicated in the control of motivated behavior. In particular, modulation of D2 SPNs in the NAc MSh has previously been shown to regulate motivation (Gallo et al., 2018; Olivetti et al., 2020). Specifically, chemogenetic inhibition of the indirect pathway NAc outputs to the VP is sufficient to enhance motivation (Gallo et al., 2018), suggesting an inhibitory role for this pathway in motivated behavior. To examine how *Grpr* deletion impacts motivation, we tested the mice on a PR lever pressing task (Fig. 7C). Prior to training, mice were food restricted to 90-95% of their body weight. The mice were trained to press the lever for a food pellet reward in daily sessions with a fixed-ratio of 1 (FR1) paradigm until they successfully received 20 or more rewards in a one-hour session on two consecutive days. No differences were seen in the number of days it took mice to learn the task (Fig. 7D). We also observed no differences in lever-pressing behavior across FR1 training days (Fig. S8A,B). Additionally, there was no difference in the starting weight of mice or the percentage of their initial body weight on the PR day between groups (Fig. 7E,F). During the two-hour PR session, mice had to press the lever an increasing number of times to earn food pellet rewards. The PR session was considered over after three minutes had elapsed without a lever press, after the mouse had engaged with the lever at least once. Survival function analysis revealed a significant difference in the number of mice that continued to engage in the task between groups, with more *Grpr* cKO mice demonstrating prolonged pressing (Fig. 7G). Additionally, *Grpr* deletion in the NAc MSh led to a significant increase in the total number of lever presses during the PR session (Fig. 7H), as well as the breakpoint, which is the maximum number of lever presses a mouse is willing to press for a single pellet reward (Fig. 7I). *Grpr* cKO mice also showed a significant increase in the total PR session time (Fig. 7J,K). These results indicate that disruption of GRP-GRPR signaling in the NAc MSh enhances motivation.

Discussion

Neuropeptides are powerful neuromodulations that regulate neuronal activity and communication in a variety of ways (Guillaumin and Burdakov, 2021; Nusbaum, 2017). It has recently become appreciated that many neurons co-express genes for both classical neurotransmitters and one or more neuropeptides (Kramer et al., 2018; Smith et al., 2019). As such, neuropeptides are poised to be key regulators of neural circuit function. Here we have identified a novel role for the neuropeptide GRP and its receptor GRPR in mediating the excitability and activity of neurons in the NAc MSh. Specifically, we find that *Grpr+* neurons in the NAc MSh are a subpopulation of D2R SPNs with distinct electrophysiological properties. We mapped the *Grp*-expressing inputs into the NAc MSh and showed that in addition to the previously identified DA neurons from the VTA, the NAc MSh receives significant *Grp+* inputs from glutamatergic neurons in the hippocampus and amygdala. Furthermore, we show both in slice and *in vivo* that the *Grpr+* neurons in the NAc MSh have high basal excitability and can be further excited by GRP. Finally, we eliminate NAc GRP-GRPR signaling by generating *Grpr* cKO mice and show that this not only reduces the baseline activation of the *Grpr+* neuronal subpopulation, but also leads to increased motivation in a PR task. Together, these experiments identify and characterize a new SPN subtype and circuit in the NAc MSh with unique physiological properties and an important role in motivated behavior.

Grpr-expressing neurons are a novel subtype of SPN in the NAc MSh
The NAc MSh is composed of two main types of SPNs that make up 90-95% of the cells in the region (Kemp and Powell, 1971). These SPNs can be divided into indirect- and direct-pathway based on the type of DA receptor they express and their output pathway to other basal ganglia structures (Castro and Bruchas, 2019; Gerfen et al., 1990; Gerfen and Surmeier, 2011; Kravitz et al., 2010; Tai et al., 2012). In the NAc MSh, dSPNs express dopamine D1 receptors, dynorphin, and substance P and have canonically been thought to promote motivated behaviors. In contrast, iSPNs express dopamine D2 receptors, adenosine 2A receptors, and enkephalin and generally inhibit it (Anderson and Reiner, 1990; Besson et al., 1990; Kupchik and Kalivas, 2017). The other 5% of cells in the NAc MSh are local interneuron populations, with the two main interneuron types in the NAc MSh being tonically active cholinergic interneurons and low-threshold spiking interneurons (Castro and Bruchas, 2019; Tepper et al., 2010). In comparison to the DS, SPNs in the NAc MSh are smaller, slightly less spiny and more active than their DS counterparts (Meredith et al., 1992; Roitman et al., 2005). Additionally, between 5-30% of SPNs in the NAc MSh have been reported to express both types of dopamine receptors (Al-Hasani et al., 2015; Bertran-Gonzalez et al., 2008; Kupchik et al., 2015).

Here we report that a subpopulation of cells in the NAc MSh express *Grpr*. When characterizing the *Grpr+* cells, we found that they co-expressed many canonical iSPN markers including *Drd2*, *Penk*, *Gad2*, and *Ppp1r1b*. We also found they had spines, and their dendritic morphology was similar to neighboring cells in the NAc MSh, the vast majority of which are expected to be SPNs. Yet, when we performed whole-cell current clamp recordings, the *Grpr*-eGFP+ cells were quite different from all other examined cell types in the NAc MSh. While iSPNs in the NAc MSh are considered the most depolarized and excitable cells at baseline (Castro and Bruchas, 2019; Ma et al., 2012; Roitman et al., 2005; Taha and Fields, 2005; Tejeda et al., 2017), *Grpr*-eGFP+ cells were both more depolarized and had an approximately 160% greater membrane resistance than

Grpr-negative *Drd2+* SPNs. We found parallel results *in vivo* when we probed for *Fos* mRNA in naïve mice, and found that *Grpr+* cells were over twice as likely to be *Fos+* than *Grpr-* iSPNs in the NAc MSh. Interestingly, when we disrupted *Grpr* expression in these cells, *Fos* was no longer preferentially expressed in *Grpr+* compared to *Grpr-* SPNs, indicating that GRP-GRPR signaling contributes to the baseline excitability of this novel cell type. Parallel results were observed in slice recordings, where incubation with DPDMB, a GRPR-specific antagonist, led to reduced baseline membrane resistance of *Grpr+* cells.

Since we observed that approximately 75% of *Grpr*-eGFP+ cells were spontaneously active at baseline in slice, this suggested that they could be a type of tonically active cholinergic interneuron (Castro and Bruchas, 2019; Zhou et al., 2002). However, *Grpr+* cells did not express *Chat* mRNA, which encodes the synthetic enzyme for acetylcholine and is a marker of these cells (Bolam et al., 1984). In addition, *Grpr*-eGFP+ cells had distinct electrophysiological properties across all measurements compared to *Chat+* cells. While the *Chat+* cell recordings were from a different mouse line, given the *in situ* and dendritic spine data we can convincingly rule out the possibility that GRPR+ cells are cholinergic interneurons. These experiments therefore identify a novel, highly excitable, and spontaneously active SPN subtype in the NAc MSh that expresses *Grpr*.

Both glutamatergic and dopaminergic inputs into the NAc MSh have Grp
This study was initially based on prior work showing that a subpopulation of midbrain DA neurons express *Grp* (Greene et al., 2005; Poulin et al., 2014; Viereckel et al., 2016) and project to the NAc MSh (Kramer et al., 2018). However, many other brain regions are also known to project to the NAc MSh (Li et al., 2018). We found that additional *Grp*-expressing inputs come into the NAc MSh from the hippocampus and amygdala and are glutamatergic. This raises the possibility of co-release of GRP with either DA or glutamate from multiple distinct inputs. Additionally, while neuropeptide signaling has been known to occur in the NAc MSh through opioid receptors, these receptors are all $G_{i/o}$-coupled, and the opioids themselves are synthesized within the NAc (Castro and Bruchas, 2019). In contrast, the *Grp+* inputs we identified are projections that come into the NAc MSh from other brain regions.

GRPR is a G_q-coupled receptor and therefore activation leads to an increase in intracellular calcium (Kroog et al., 1995). Previous studies that have examined GRP-GRPR signaling in other brain regions found that GRP excites GRPR-expressing neurons in the auditory cortex, paraventricular thalamus, lateral amygdala, hippocampus and spinal cord (Hermes et al., 2013; Lee et al., 1999; Melzer et al., 2021; Pagani et al., 2019; Piggins and Rusak, 1993; Pinnock and Woodruff, 1991; Shumyatsky et al., 2002). In the spinal cord, it was shown that GRP excites neurons via blockade of $K_{ir}2$ potassium channels (Pagani et al., 2019). We found that while the *Grpr+* neurons in the NAc MSh were already spontaneously active at baseline, application of GRP in slice caused further depolarization and increased the membrane resistance by 140%. These effects could be blocked by a GRPR-specific antagonist and the presence of the antagonist alone led to a decrease in baseline membrane resistance, raising the possibility that there is basal GRP-GRPR activity in slice. Together these experiments map out a novel mesolimbic GRP-GRPR circuit whereby GRP comes into the NAc MSh from glutamatergic and dopaminergic neurons from the hippocampus, amygdala, and VTA and acutely excites GRPR+ cells, which are GABAergic.

GRP-GRPR signaling affects striatal associated behaviors
GRP-GRPR signaling in other brains regions is important for a variety of different behaviors (Roesler et al., 2014; Roesler and Schwartsmann, 2012). For example, in the amygdala, *Grpr* KO mice showed enhanced long-term fear memories (Shumyatsky et al., 2002). In the spinal cord, GRP-GRPR signaling is the basis of itch (Majumdar and Weber, 2011; Pagani et al., 2019; Sun et al., 2017; Sun and Chen, 2007; Sun et al., 2009; Yu et al., 2017). More recently, *Grpr* KO in the auditory cortex decreased auditory fear memories (Melzer et al., 2021). Given that GRP-GRPR signaling in other brain regions is sufficient to modulate behavior, we generated *Grpr* cKO mice to test whether NAc-associated behaviors were impacted.

We first tested *Grpr* cKO mice in the open field test to assess general locomotion and used Keypoint-MoSeq (Weinreb et al., 2023; Wiltschko et al., 2020) to identify behavioral syllables in an unbiased manner. We observed a slight increase in the total distance travelled in *Grpr* cKO mice, but found no differences in center entries, grooming bouts, or in the frequency of any of the top syllables identified using MoSeq. This indicates that GRP-GRPR signaling in the NAc MSh does not grossly affect behavior.

In addition to the open field test, we tested mice on the accelerating rotarod, an operant conditioning protocol, and the sucrose preference test, which have all been associated with the ventral striatum (Cording and Bateup, 2023; Costa et al., 2004; Durieux et al., 2012; Gallo et al., 2018; Liu et al., 2018; Platt et al., 2017; Rothwell et al., 2014; Serchov et al., 2016; Trifilieff et al., 2013; Wulff et al., 2023; Yin et al., 2009). While we saw no differences in the accelerating rotarod or sucrose preference test, we found that *Grpr* cKO mice showed increased motivation in the PR task. We do not believe this is due to increased lever pressing in general, as there were no significant differences in lever-press behavior during FR1 training. The NAc MSh is an important regulator of motivation and, more specifically, iSPNs have been shown to inhibit motivation (Gallo et al., 2018; Kupchik et al., 2015). In a prior study, the authors used chemogenetics to inhibit iSPN outputs to the VP in the NAc MSh and found this increased motivation. This is consistent with studies showing that iSPNs typically inhibit motivated behaviors (Castro and Bruchas, 2019; Kupchik and Kalivas, 2017; Voorn et al., 2004). Here, by eliminating GRP-GRPR signaling specifically in the NAc MSh, we expected to reduce the excitability of a small subpopulation of iSPNs. Consistent with this, we found that the percentage of *Fos*-positive *Grpr*-lineage iSPNs was reduced in *Grpr* KO mice. The inability to activate *Grpr*-expressing iSPNs led to increased motivation, consistent with studies broadly manipulating iSPNs in this region. This suggests that GRP-GRPR signaling in the NAc MSh normally constrains motivation and suggests that the relatively small population of *Grpr*+ NAc neurons can have an impact on motivated behaviors.

Limitations and future directions
This study identified and characterized a previously undescribed subpopulation of SPNs in the NAc MSh that express *Grpr* and *Drd2*. We provide a genetic, morphologic, and electrophysiological characterization of these cells and probe a possible behavioral role of GRP-GRPR through *Grpr* deletion. These experiments lay the groundwork for further characterization of GRP-GRPR signaling in the striatum and in the brain, in general. We note that although it was beyond the scope of the present study, *Grpr*+ cells also exist in the DS. Given the unique electrophysiological properties of the GRPR+ cells in the NAc MSh, the GRPR+ in the DS warrant their own characterization. Additionally, while we identified the *Grp*+ inputs into the NAc MSh and

noted that the *Grpr+* cells themselves were GABAergic, we were unable to address whether the *Grpr+* cells project to and inhibit the ventral pallidum. This was largely due to limitations in the available transgenic mouse lines for manipulating *Grpr*-expressing cells. Recent evidence has indicated that unlike the DS, the canonical projection pathways from the NAc are oversimplified and both D1- and D2-SPNs can participate in indirect and direct pathways depending on their downstream target from the ventral pallidum (Kupchik et al., 2015). Therefore, the outputs of *Grpr+* cells should be examined in future studies. Additionally, while we perform cellular-level characterization of the *Grpr*-expressing cells and test a subset of behaviors that have been associated with the NAc MSh, future studies could assess a possible role of GRP-GRPR signaling in additional behaviors. One such direction could be utilizing tasks that assess context-induced reinstatement of drug seeking for drugs of abuse. These tests have been associated specifically with ventral subiculum to NAc MSh projections, where we find a majority of the *Grp* inputs to the NAc MSh (Bossert et al., 2016; Martin and Ono, 2000). Finally, given the G_q-coupling, a possible role of GRP in modulating synaptic transmission or plasticity could be examined.

Overall, our study identifies a new cell type in the NAc MSh and describes a mesolimbic circuit defined by the expression and functionality of GRP to GRPR signaling.

Methods

Mice

Animal experiments were performed in accordance with protocols approved by the University of California, Berkeley, Institutional Animal Care and Use Committee (protocol #: AUP-2016-04-8684-2). Both male and female mice were used across all experiments except for the knock-out FISH experiments where male littermate WT and *Grpr* KO mice were used. Mice were postnatal day P30-P90 unless otherwise stated. The following mouse lines were used in this study: C57BL/6J, JAX strain #000664, *Grpr*::eGFP (GENSAT [MMRRC #036178-UCD]), *Drd2*-eGFP (GENSAT [MMRRC #000230-UNC]), *Drd1a*-tdTomato (JAX strain #016204) (Ade et al., 2011), CMV-Cre (JAX strain #006054) (Schwenk et al., 1995), *Grpr^fl* (JAX strain # 033148) (Yu et al., 2017), ChAT-IRES-Cre (JAX strain # 06410) (Rossi et al., 2011), and Ai9 (JAX strain # 07909) (Madisen et al., 2010). *Grpr* KO mice were generated by breeding CMV-Cre mice to *Grpr^fl* mice to knock out *Grpr* in all cells.

Fluorescent in situ hybridization

Fluorescent *in situ* hybridization was performed to identify and characterize *Grpr*-expressing cells in the NAc MSh and DS as well as their *Grp*-expressing inputs. Brains were harvested, flash-frozen in OCT mounting medium (Fisher Scientific #23-730-571) on dry ice and stored at -80 °C for up to 6 months. 16 μm-thick coronal sections were collected using a cryostat, mounted directly onto 75x25mm Superfrost® Plus glass slides (VWR #48311-703) and stored at -80°C for up to 1 month. Fluorescent *in situ* hybridization was performed according to protocols provided with the RNAscope® Multiplex Fluorescent Reagent Kit (ACD #320850) and RNAscope® Multiplex Fluorescent Reagent Kit V2 (ACD #323100). *Grpr* mRNA was visualized with a probe in channel 2 (ACD #317871-C2), *Gad2* mRNA in channel 1 (ACD #439371), *Drd2* mRNA in channel 1 (ACD #406501), *Drd1* mRNA in channel 3 (ACD #461901-C3), *Chat* mRNA in channel 1 (ACD #408731), *Ppp1r1b* mRNA in channel 3 (ACD #405901-C3), *Grp* mRNA in channel 2 (ACD #317861-C2), *Slc1717* mRNA in channel 1 (ACD #481851), and *Fos* mRNA in channel 3 (ACD #316921-C3).

Sections were imaged using an Olympus FluoView 3000 confocal microscope equipped with 405, 488, 561, and 640 nm lasers and a motorized stage for tile imaging. Z stack images captured the entire thickness of the section at 1-2.5 μm steps for images taken with 20x (Olympus #UCPLFLN20X) or 10x air objectives (UPLXAPO10X). Additional images were acquired on a Zeiss LSM 710 AxioObserver with Zeiss 10x, 20x, and 63x objectives housed in the Molecular Imaging Center at UC Berkeley. Images were analyzed using FIJI. Cells were considered positive for *Grpr* if they contained two or more fluorescent puncta within the cellular boundary created by DAPI. All images for a single dataset were acquired on the same microscope.

Immunohistochemistry

Male and female mice (P30-P90) were deeply anesthetized by isoflurane and transcardially perfused with ice cold 1X PBS (~5–7 ml) followed by 4% paraformaldehyde (PFA) solution (Electron Microscopy Sciences: 15713) in 1X PBS (~5–10 ml) using a peristaltic pump (Instech). The brains were removed and post-fixed by immersion in 4% PFA in 1X PBS solution overnight at 4°C. Brains were suspended in 30% sucrose in 0.1 M PBS for cryoprotection. After brains descended to the bottom of the vial (typically 24–48 hr), 40 μm coronal sections of the NAc MSh

were cut on a freezing microtome (American Optical AO 860), collected into serial wells, and stored at 4°C in 1X PBS containing 0.02% (w/v) sodium azide (NaN3; Sigma Aldrich).

Free-floating sections were washed with gentle shaking, 3 × 5 min in 1X PBS followed by 1 hr incubation at RT with BlockAid blocking solution (Life Tech: B10710). Primary antibodies were applied at 4°C in 1X PBS containing 0.25% (v/v) Trinton-X-100 (PBS-Tx) overnight to ensure penetration of the antibody throughout the slice. Sections were then washed with cold 1X PBS-Tx 3 × 10 min and incubated for 2 hr at RT with secondary antibodies in PBS-Tx. Sections were washed in cold 1X PBS 3 × 10 min, mounted on SuperFrost slides (VWR: 48311–703), and coverslipped with Prolong Gold antifade mounting media (Life Tech: P36935).

We used a primary antibody against GFP (1:5000, Chicken, Abcam: 13970) and an Alexa Fluor 488 goat anti-chicken secondary antibody (1:500, Invitrogen: A-11039).

Images of 40 µm sections processed for immunohistochemistry were acquired using an Olympus FluoView 3000 confocal microscope (described above). Z-stack images captured the entire thickness of the section at 1–2 µm steps for images taken with a 20X air (Olympus #UCPLFLN20X) or a 10X air Olympus #UPLXAPO10X) objective.

Neurobiotin-filled neuron reconstruction for Sholl analysis
Male and female mice were deeply anesthetized by isoflurane, transcardially perfused with ice cold ACSF using a peristaltic pump (Instech) and decapitated. 275-µm-thick coronal striatal slices were prepared on a vibratome (Leica VT1000 S) in oxygenated ice-cold choline-based external solution (pH 7.8) containing 110 mM choline chloride, 25 mM NaHCO3, 1.25 mM NaHPO4, 2.5 mM KCl, 7 mM MgCl2, 0.5 mM CaCl2, 25 mM glucose, 11.6 mM sodium ascorbate, and 3.1 mM sodium pyruvate. Slices were recovered in ACSF at 34°C for 15 min and then kept at room temperature (RT) for the duration of the recordings. All solutions were continuously bubbled with 95% O2 and 5% CO2. For whole cell recordings, 3–5 mΩ borosilicate glass pipettes (Sutter Instrument: BF150-86-7.5) were filled with a potassium-based internal solution (pH 7.4) containing 135 mM KMeSO4, 5 mM KCl, 5 mM HEPES, 4 mM Mg-ATP, 0.3 mM Na-GTP, 10 mM phosphocreatine, 1 mM EGTA, and 4 mg/mL neurobiotin (Vector laboratories: SP-1120).

275 µm slices containing *Grpr*-eGFP+/- NAc MSh neurons were filled with neurobiotin-containing internal solution (4 mg/mL) during patch-clamp experiments and then fixed in 4% paraformaldehyde solution (Electron Microscopy Sciences: 15713) in 1X PBS for 24–48 hr at 4°C. With continuous gentle shaking, slices were washed in 1X PBS 3 × 5 min and incubated with BlockAid blocking solution (Life Tech: B10710) for 1 hr at RT. Streptavidin Alexa Flour 633 conjugate (1:750, Invitrogen: S21375) was applied overnight at 4°C in 1X PBS containing 0.50% (v/v) Triton-X-100 (PBS-Tx). The following day, slices were washed in 1X PBS 3 × 10 min, mounted on SuperFrost slides (VWR: 48311–703) with the cell-containing side of the slice facing up, and coverslipped with Prolong Gold antifade mounting media (Life Tech: P36935).

Filled cells in the mounted sections were imaged on an Olympus FluoView 3000 confocal microscope at 20x magnification with 2X digital zoom for a final magnification of 40X. 1 µm steps were used to acquire a z-stack spanning the entirety of the neurobiotin-filled cell body and dendritic arbors. 3D reconstructions of the cells were done using IMARIS 9.2.1 software (Bitplane) with automated filament tracing and manual editing. The mask generated by the automated

filament tracing algorithm was continuously cross-referenced with the original z-stack image to ensure accuracy. Spurious segments created by the automated filament tracer due to background noise were removed, while processes with incomplete reconstruction were manually edited to incorporate missing segments.

Sparse viral injections for spine density quantification
Neonatal (P0) or juvenile (P28-30) *Grpr*::eGFP mice were injected bilaterally with 250 nL (neonates) or 100 nL (juveniles) AAV5-hSyn-mCherry (Addgene: 114422-AAV5) diluted 1:200 to label neurons in the NAc MSh. Neonatal mice were cryoanesthetized and injected bilaterally at 200 nL/min. Injections were targeted to the NAc MSh, with coordinates approximately 0.75 mm lateral to midline, 1.0 mm posterior to bregma, and 3.0 mm ventral to the head surface. For juvenile injections, mice were briefly anesthetized with 3% isoflurane (Piramal Healthcare: PIR001710) and oxygen. Mice were mounted on a stereotaxic frame (Kopf instruments Model 940) with stabilizing ear cups and a nose cone delivering constant 1.5% isoflurane in medical oxygen. Viruses were injected using a pulled glass capillary (Drummond Scientific Company: 5-000-2005) at a rate of 100 nL/minute. Following injection, the capillary remained in place for 1 minute per every minute spent injecting to allow the tissue to recover and prevent virus backflow up the injection tract upon retraction. Coordinates from bregma for juvenile injections were as follows: M/L +/- 1.00 mm, A/P +1.90 mm, D/V -4.50 mm.

All mice were perfused at P56. Mice were perfused and brains were post-fixed with 4% paraformaldehyde overnight, then sectioned at 80 µm. Sections were blocked for 1 hr at RT in BlockAid (ThermoFisher: B10710) and incubated for 16 hr at 4°C with antibodies against RFP (1:500, rabbit, Rockland (VWR): RL600-401-379) and GFP (1:500, chicken, Abcam: 13970). Sections were washed 3 × 10 min in PBS-Tx and incubated for 2 hr at RT with Alexa Fluor 488 and Alexa Fluor 546 secondary antibodies (1:500, Invitrogen: A-11039 and A-11001). Sections were washed 3 × 10 min in 1X PBS and mounted onto slides using Prolong Gold antifade mounting media (Life Tech: P36935).

Dendrites were imaged on an Olympus FluoView 3000 scanning confocal microscope at 60x magnification with 2.5x digital zoom for a final magnification of 150x. Images were deconvoluted in the CellSens4 software using a built-in advanced maximum likelihood algorithm. Dendritic spine reconstructions were generated using the filament tracer feature in Imaris 9.3.1. 1-9 dendrites were reconstructed per cell, with the detection parameters for thinnest spine diameter set to 1.5 µm and the maximum spine length set to 3.5 µm. After automatic detection of spines, the digital reconstruction was compared to the original z-stack image to ensure accuracy. Detected spines were manually reviewed by the experimenter and added or removed as needed. Spine density of each dendrite was calculated by dividing the number of spines on that dendrite by the total dendrite length. The values of all dendrites quantified for a given neuron were averaged and the average spine density per dendrite is reported.

Electrophysiology
Mice (P30-90) were perfused transcardially with ice-cold ACSF (pH=7.4) containing (in mM): 127 NaCl, 25 NaHCO3, 1.25 NaH2PO4, 2.5 KCl, 1 MgCl2, 2 CaCl2, and 25 glucose, bubbled continuously with carbogen (95% O2 and 5% CO2). Brains were rapidly removed, and coronal slices (275 µm) were cut on a VT1000S vibratome (Leica) in oxygenated ice-cold choline-based

external solution (pH=7.8) containing (in mM): 110 choline chloride, 25 NaHCO3, 1.25 NaHPO4, 2.5 KCl, 7 MgCl2, 0.5 CaCl2, 25 glucose, 11.6 sodium ascorbate, and 3.1 sodium pyruvate. Slices were recovered in ACSF at 34°C for 15 min and then kept at RT before recording.

Recordings were made with a MultiClamp 700B amplifier (Molecular Devices) at RT using 3-5 MOhm glass patch electrodes (Sutter: BF150-86-7.5). Data were acquired using ScanImage software, written and maintained by Dr. Bernardo Sabatini (https://github.com/bernardosabatini/%20SabalabAcq). Traces were analyzed in Igor Pro (Wavemetrics). Recordings with a series resistance > 30 MOhms were rejected.

Current-clamp recordings
Current clamp recordings were made using a potassium-based internal solution (pH=7.4) containing (in mM): 135 KMeSO4, 5 KCl, 5 HEPES, 4 Mg-ATP, 0.3 Na-GTP, 10 phosphocreatine, and 1 ETGA. No synaptic blockers were included, and no holding current was applied to the membrane. For intrinsic excitability experiments, positive current steps (1s, +10 to +100 pA) were applied in 10 pA intervals to generate an input-output curve. Membrane potential was averaged over a 300 ms period between 50-350 ms of the 2 s sweep prior to current injection. Membrane resistance was calculated from voltage-clamp RC checks prior to switching to current-clamp. A cell was considered spontaneously active at baseline if it fired at least one action potential during the first two minutes after breaking in without current injection. Action potential half width was calculated as the width at half maximum amplitude of spikes induced by current injection.

GRP wash-on
Membrane potential changes upon bath application of 300 nM GRP (Anaspec: AS-24214) dissolved in ACSF were monitored in current-clamp mode. 5 s current-clamp recordings were made every 2 s. Prior to GRP wash-on, membrane potential was recorded for 10 minutes at baseline, and membrane potential was averaged over a 300 ms window between 50-350 ms of the 2 s sweep. 300 nM GRP was then bath applied for 10 minutes. At the conclusion of GRP wash-on, we briefly switched back to voltage-clamp to perform 5 RC checks to calculate membrane resistance and confirm that series resistance was stable and <30 MOhms. GRP was then washed-out. The baseline membrane potential was calculated as the average over the two minutes prior to GRP wash-on. The membrane potential following GRP application was calculated as the average over the last two minutes of wash-on. Baseline membrane resistance was calculated from the initial voltage-clamp RC check. Wash-on experiments in the presence of the GRPR antagonist, 1 µM DPDMB (Bachem: H-3042) were performed in the same way, except the slices were pre-incubated in DPDMB for at least 15 minutes prior to recording and DPDMB was present in the ACSF for the duration of the recordings.

Stereotaxic injections
Retrobead injections
P50-90 WT male and female mice were used for retrograde labeling experiments. Red retrobeads IX (Lumafluor: 78R180) were diluted 1:3 in sterile saline. 350 nL diluted beads were unilaterally injected into the NAc MSh using a pulled glass pipette. The following coordinates from bregma were used: M/L + 0.65 mm, A/P +1.90 mm, D/V -4.50 mm. To allow for sufficient labeling, mice were sacrificed 14 days post injection. Brains were harvested, flash-frozen in OCT mounting medium (Fisher Scientific: 23-730-571) on dry ice and stored at -80°C for up to 6 months. 16 µm-

thick coronal sections were collected using a cryostat, mounted directly onto 75x25mm Superfrost® Plus glass slides (VWR: 48311-703) and stored at -80°C for up to 1 month. Prior to sectioning the posterior part of the brain, the injection site was validated to ensure that retrobeads were contained in the NAc MSh. After confirming, the rest of the brain was sectioned and fluorescent *in situ* hybridization was performed on all sections containing brain regions with retrobead+ cells.

<u>GRP injections</u>
Male and female mice (P30-P90) underwent stereotaxic surgery to inject GRP into the NAc MSh. 800 nL of 3 µM GRP (dissolved in sterile saline) or sterile saline were injected unilaterally into the left NAc MSh. The stereotaxic coordinates used were: + 0.65 mm, A/P +1.90 mm, D/V -4.50 mm. 45 minutes after the end of the injection, mice were sacrificed, and their brains harvested and flash frozen for fluorescent *in situ* hybridization.

<u>CRE and GFP injections</u>
Male and female mice (P28-30) underwent stereotaxic surgery to inject pENN.AAV.hSyn.HI.eGFP-Cre.WPRE.SV40 (AAV5) or pAAV-hSyn-EGFP (AAV5) into the NAc MSh. 600 nL of 1:3 diluted CRE or GFP virus in sterile saline were injected bilaterally into the NAc MSh. The stereotaxic coordinates used were: +/- 0.85 mm, A/P +1.70 mm, D/V -4.50 mm. Mice were allowed to recover and viruses were given four weeks to express before mice underwent behavioral testing. At the end of behavioral testing, mice were sacrificed and the tissue was sliced to confirm proper viral targeting.

<u>Genotyping to validate global KO mice</u>
Multiple PCRs were used as a genotyping strategy to confirm the genotype of *Grpr* KO and WT control mice. The strategy involved primer pairs for the presence of the floxed or WT allele, the LoxP scar, and an internal region of *Grpr* exon 2. The following primer sets were used: *Grpr* floxed (WT- forward: 5' GGAGAGAGGTATAGAGGGGC 3', WT-R: 5' ACATGTAGGTTGCAGGGGAT 3', MUT-F 5' GGGTTATTGTCTCATGAGCGG 3'), LoxP scar (Forward 5' AGACAGCTCTATCACGGTCC 3', Reverse 5' GAGTTCAGTTCCCAGCAACC 3'), *Grpr* Exon 2 (Forward 5' TTTCTGACCTCCACCCCTTC 3', Reverse 5' CACGGGAAGATTGTAGGCAC 3'). With this strategy, we expected the following bands for *Grpr* KO mice: *Grpr* floxed (no band), CMV-Cre (521 bp internal control), LoxP scar (298 bp), *Grpr* Exon 2 (no band) and WT mice: *Grpr* floxed (573 bp WT band), CMV-Cre (521 bp internal control and 180 bp band), LoxP scar (no band), *Grpr* Exon 2 (206 bp).

Behavioral experiments
Behavior studies were carried out in the dark phase of the light cycle under red lights (open field) or white lights (rotarod and operant conditioning). Sucrose preference was carried out in the room where the mice are normally housed. Mice were habituated to the behavior testing room for at least 30 minutes prior to testing, and at least 24 hours elapsed between sessions. All behavioral equipment was cleaned between each trial with 70% ethanol. Equipment was rinsed with diluted soap and water at the end of each day. Behavioral experiments included both male and female mice and began four weeks after the stereotaxic injections at P54-P60. Male mice were trained

or tested before female mice each day. Experimenters were blinded to the genotype during behavioral testing.

Open field

Exploratory behavior in a novel environment and general locomotor activity were assessed by a 60 min session in an open field chamber (40 cm L x 40 cm W x 34 cm H) made of transparent plexiglass. The mouse was placed in the bottom right hand corner of the arena and behavior was recorded using an overhead monochrome camera (FLIR Grasshopper 3: GS3-U3-41C6NIR-C) with a 16 mm wide angle lens (Kowa: LM16HC) placed on top of the arena from a height of 50 cm. Data was analyzed using DeepLabCut (DLC) (Mathis et al., 2018; Nath et al., 2019) and Keypoint-MoSeq (Weinreb et al., 2023; Wiltschko et al., 2020).

Processing and analysis of open field data

To extract the body part (keypoint) coordinates from the video recordings, we used DLC 2.3.4 (Mathis et al., 2018; Nath et al., 2019). 14 body parts including nose, head, left ear, right ear, left forelimb, right forelimb, spine 1, spine 2, spine 3, left hindlimb, right hindlimb, tail 1, tail 2, and tail 3 were manually labelled on a small subset of the video frames. A DLC model was then trained using the annotated frames to label these 14 body parts for all videos recorded. The distance travelled and center entries were calculated with the coordinate of body part tail 1.

Discrete behavior syllables were extracted with Keypoint-MoSeq 0.4.4 (Weinreb et al., 2023). Syllable usage and transition data were obtained using built-in functions in the Keypoint-MoSeq package. Decoding analysis was done with customized Python 3.9 script. Code available upon request in Github.

Rotarod

The accelerating rotarod test was used to examine motor learning. Mice were trained on a rotarod apparatus (Ugo Basile: 47650) for four consecutive days. Three trials were completed per day with a 5 min break between trials. The rotarod was accelerated from 5-40 revolutions per minute (rpm) over 300 s for trials 1-6 (days 1 and 2), and from 10-80 rpm over 300 s for trials 7-12 (days 3 and 4). On the first testing day, mice were acclimated to the apparatus by being placed on the rotarod rotating at a constant 5 rpm for 60 s and returned to their home cage for 5 minutes prior to starting trial 1. Latency to fall, or to rotate off the top of the rotarod barrel, was measured by the rotarod stop-trigger timer.

Operant conditioning

Motivated behavior was assessed in an operant conditioning lever pressing task using a food pellet reinforcer. Animals were food restricted to ~90-95% of their body weight before commencing training in an operant conditioning chamber (Med Associates: ENV-307A). The chambers contained a single retractable lever on the left side of the food receptacle and a house light on the opposite end of the chamber. Each session began with illumination of the house light and the presentation of the left lever. The session ended with retraction of the lever and with the house light turning off. The lever remained present throughout the session. Mice were weighed before each session and habituated to the experiment room for at least 30 minutes prior to testing.

During the first training session, following two days of food restriction, a food pellet (14 mg regular "chow" pellet, Bio-Serve: F05684) was delivered on a random time schedule (RT), with a reinforcer delivered on average every 60 seconds for a total of 15 minutes with no levers present. On subsequent days, mice underwent 60-minute fixed ratio (FR1) sessions where mice were presented with only the left lever. Each lever press was rewarded with a single pellet reward. During initial FR1 sessions, crushed pellets were placed on the lever at the beginning of the task and each time 10 minutes elapsed without a lever press to prime the mice to press the lever. Mice underwent daily FR1 sessions until they had two consecutive sessions with 20 or more lever presses. After earning 20 rewards the first time, the levers were no longer primed with crushed pellets. The mice then moved on to the progressive ratio (PR) session the following day. The PR session ended when a mouse failed to press the lever for a period of at least three minutes, after initiating the lever at least once, or after two hours had elapsed, whichever came first. The number of lever presses required for a reward increased with each reward. The number of lever presses to earn a reward were as follows: 1, 2, 3, 4, 5, 6, 7, 8, 10, 12, 14, 16, 18, 20, 22, 24, 28, 32, 36, 40, 44, 48, 52, 56, 64, 72, 80, 88, 96, 104, 112, 120, 128, 136.

<u>Sucrose preference test</u>
Grpr cKO and GFP-injected control mice were assessed for normal hedonic behavior by using the sucrose preference test. Three days prior to the sucrose preference test, the water source in the home cage was switched from a Hydro Pac (plain water) to a water bottle containing the contents of the Hydro Pak. After three days of acclimating to the water bottles, a second bottle was added to the home cage containing 5% sucrose in Hydro Pac water. Mice had access to both bottles for 24 hours. After 24 hours, mice were individually housed, again with two bottles, one with 5% sucrose and the other with standard water. Mice were kept singly housed for another 24 hours. The volume of water and sucrose consumed during this 24-hour period was measured. Sucrose preference was calculated by taking the total volume of sucrose solution consumed and dividing it by the total liquid consumed in the 24-hour period. To avoid any confounds due to side-specific preferences, the bottles' location was randomly assigned (front or back) for each mouse. All experiments were conducted on weekends, ensuring a quiet environment for the tested animals.

Quantification and statistical analysis
GraphPad Prism 10 was used to perform statistical analyses. All datasets were first analyzed using D'Agostino and Pearson normality test, and then parametric or non-parametric two-tailed statistical tests were employed accordingly to determine significance. If the variances between two groups were significantly different, a Welch's correction was applied. Significance was set as *$p < 0.05$, **$p < 0.01$, ***$p < 0.001$, and ****$p < 0.0001$. All p values were corrected for multiple comparisons. Statistical details including sample sizes for each experiment are reported in the figure legends.

Figures and Legends

Figure 1

Figure 1: *Grpr* is expressed in striatal projection neurons in the NAc MSh.

(A) Representative image of a coronal brain section (A/P +1.18) from a wild-type (WT) mouse. Fluorescent *in situ* hybridization (FISH) for *Grpr* in magenta. Nuclei are labeled in blue with DAPI. Insets show higher magnification images for dorsal striatum (DS) and nucleus accumbens medial shell (NAc MSh). **(B)** Summary schematic of *Grpr* expression across the A/P axis of the DS and NAc MSh. Each dot represents the location of a *Grpr*+ cell. Dots are summed across 3 WT mice. **(C)** Representative image of the NAc MSh. FISH for *Gad2* in green and *Grpr* in magenta. Nuclei labeled with DAPI in blue. Dashed circles outline *Grpr*+ cells. **(D)** Mean +/- SEM of percentage of *Grpr*+ cells in the NAc MSh and that are *Gad2*+ in WT mice. (n= 4 mice). **(E-G)** Representative images of NAc MSh. Nuclei labeled with DAPI in blue. **(E)** FISH for *Chat* in green, *Grpr* in magenta, and *Ppp1r1b* in cyan. Dashed circles outline *Grpr*+ cells. **(F)** FISH for *Drd2* in green, *Grpr* in magenta, and *Drd1* in cyan. Dashed circles outline *Grpr*+ cells. **(G)** FISH for *Drd2* in green, *Grpr* in magenta, and *Penk* in cyan. Dashed circles outline *Grpr*+ cells. **(H-L)** Mean +/- SEM of percentage of *Grpr*+ cells in the NAc MSh of WT mice that are **(H)** *Chat*+ (n=7 mice), **(I)** *Ppp1r1b*+ (n=7 mice), **(J)** *Drd2*+ (n=12 mice), **(K)** *Drd1*+ (n=8 mice), or **(L)** *Penk*+ (n=8 mice). **(M)** Mean +/- SEM of percentage of *Grpr*+ cells in the NAc MSh that are *Drd2*+/*Drd1*- (green), *Drd2*-/*Drd1*+ (blue), or *Drd2*+/*Drd1*+ (black) in WT mice (n= 8 mice). **(N)** Mean +/- SEM of percentage of *Grpr*+ cells in the NAc MSh that are *Drd2*+/*Penk*- (green), *Drd2*-/*Penk*+ (blue), or *Drd2*+/*Penk*+ (black) in WT mice (n= 8 mice). **(O-Q)** Mean +/- SEM of percentage of **(O)** *Ppp1r1b*+ (n=7 mice), **(P)** *Drd2*+ (n=12 mice), or **(Q)** *Penk*+ cells (n=8 mice) in the NAc MSh that are *Grpr*+.

Figure 2

Figure 2: GRPR-expressing neurons in the NAc MSh are morphologically similar to neighboring striatal neurons.

(A) Representative image of a coronal brain section (A/P +1.18) from a *Grpr*::eGFP mouse that was used to identify GRPR+ cells for whole-cell recordings. eGFP in green, DAPI labeled nuclei in blue. **(B)** Zoom-in of the nucleus accumbens medial shell (NAc MSh) from panel **A**. **(C)** Zoom-in of panel **B** showing eGFP+ cells in grayscale. **(D-E)** Reconstructions of the dendrites and cell body of NAc MSh *Grpr*-eGFP+ **(D)** and *Grpr*-eGFP- **(E)** neurons. **(F)** Sholl analysis of NAc MSh neuron dendritic arbors. Dark colored lines are the mean, lighter color shading represents the SEM. *Grpr*-eGFP+ in green: n=18 neurons from 8 mice, *Grpr*-eGFP- in black: n=23 neurons from 14 mice. Two-way ANOVA p-values are shown. **(G)** Mean +/- SEM total dendritic arbor length per cell. *Grpr*-eGFP+ in green: n=18 neurons from 8 mice, *Grpr*-eGFP- in black: n=23 neurons from 14 mice; p=0.4452, Welch's two-tailed t-test. **(H)** Box plots of the number of primary dendrites per neuron for each genotype (boxes represent the interquartile range (25-75%), lines denote the median, whiskers represent mix to max values). *Grpr*-eGFP+ in green: n=18 neurons from 8 mice, *Grpr*-eGFP- in black: n=23 neurons from 14 mice; p=0.1151, Welch's two-tailed t-test. **(I-J)** Representative images of dendrites from a *Grpr*-eGFP+ **(I)** and *Grpr*-eGFP- **(J)** neuron in the NAc MSh. **(K)** Mean +/- SEM spine density in *Grpr*-eGFP+ neurons vs *Grpr*-eGFP- neurons. *Grpr*-eGFP+ in green: n=21 neurons from 11 mice, *Grpr*-eGFP- in black: n=18 neurons from 9 mice; p=0.1357, Welch's two-tailed t-test. For panels **G**, **H**, and **K**, dots represent values from individual neurons.

Figure 3

Figure 3: GRPR-expressing neurons in the NAc MSh have high intrinsic excitability.

(A) Representative coronal brain section (A/P +1.70) from a *Grpr*::eGFP;Drd1-tdTomato mouse that was used to identify GRPR+ cells for whole-cell recordings. **(B)** Zoom-in of the dorsal striatum (DS) from panel **A**. **(C)** Zoom-in of the nucleus accumbens medial shell (NAc MSh) from panel **A**. **(D)** Mean +/- SEM percent of eGFP+ cells that are tdTomato+ in *Grpr*::eGFP;Drd1-tdTomato mice in the DS (magenta) and NAc MSh (green) (n=6 mice). **(E-H)** Representative traces from **(E)** *Grpr*-eGFP+, **(F)** *Grpr*-eGFP-, **(G)** D1-tdTomato+, and **(H)** D1-tdTomato-/*Grpr*-eGFP- (presumed D2+) cells in the NAc MSh recorded at baseline (left) and following +40 pA current injection (right). **(I)** Mean +/- SEM membrane potential of NAc MSh neurons. *Grpr*-eGFP+ in green: n=30 cells from 15 mice, *Grpr*-eGFP- in black: n=34 cells from 11 mice, D1-tdTomato+ in magenta: n=31 cells from 5 mice, and D1-tdTomato-/*Grpr*-eGFP- in dark blue: n=27 cells from 3 mice. (****$p<0.0001$, one-way ANOVA; Holm-Sidak's multiple comparison tests, ***$p=0.0005$ *Grpr*-eGFP+ vs *Grpr*-eGFP-, ****$p<0.0001$ *Grpr*-eGFP+ vs D1-tdTomato+, ****$p<0.0001$ *Grpr*-eGFP+ vs D1-tdTomato-/*Grpr*-eGFP-). **(J)** Mean +/- SEM membrane resistance of NAc MSh neurons. *Grpr*-eGFP+ in green: n=30 cells from 15 mice, *Grpr*-eGFP- in black: n=34 cells from 11 mice, D1-tdTomato+ in magenta: n=31 cells from 5 mice, and D1-tdTomato-/*Grpr*-eGFP- in dark blue: n=27 cells from 3 mice. (****$p<0.0001$, one-way ANOVA; Holm-Sidak's multiple comparison tests, ***$p=0.0004$ *Grpr*-eGFP+ vs *Grpr*-eGFP-, ****$p<0.0001$ *Grpr*-eGFP+ vs D1-tdTomato+, ***$p=0.0003$ *Grpr*-eGFP+ vs D1-tdTomato-/*Grpr*-eGFP-). **(K)** Mean +/- SEM action potential (AP) half-width of NAc MSh neurons. *Grpr*-eGFP+ in green: n=23 cells from 10 mice, *Grpr*-eGFP- in black: n=20 cells from 7 mice, D1-tdTomato+ in magenta: n=28 cells from 5 mice, and D1-tdTomato-/*Grpr*-eGFP- in dark blue: n=26 cells from 3 mice. (****$p<0.0001$, one-way ANOVA; Holm-Sidak's multiple comparison tests, ****$p<0.0001$ *Grpr*-eGFP+ vs *Grpr*-eGFP-, ****$p<0.0001$ *Grpr*-eGFP+ vs D1-tdTomato+, ****$p<0.0001$ *Grpr*-eGFP+ vs D1-tdTomato-/*Grpr*-eGFP-). **(L)** Input-output curves showing the mean +/- SEM firing frequency of NAc MSh neurons in response to positive current steps of increasing amplitude. *Grpr*-eGFP+ in green: n=23 cells from 10 mice, *Grpr*-eGFP- in black: n=21 cells from 7 mice, D1-tdTomato+ in magenta: n=31 cells from 5 mice, and D1-tdTomato-/*Grpr*-eGFP- in dark blue: n=27 cells from 3 mice. For panels **I**, **J**, and **K**, + indicates the presence of a fluorophore labeling the given cell type, - indicates that the cell was negative for the fluorophore, and o indicates that the tissue did not have a fluorophore that would label the given cell type.

Figure 4

Figure 4: Glutamatergic cells from the hippocampus and amygdala that project to the NAc MSh express *Grp.*

(A) Schematic of retrograde tracing experiment. Red retrobeads were injected into the nucleus accumbens medial shell (NAc MSh) and the listed projection targets were examined for *Grp* mRNA and retrobead co-localization using fluorescent *in situ* hybridization (FISH). Projection targets: ventral tegmental area (VTA), lateral entorhinal cortex (LE), anterior piriform (APir), hippocampus subiculum (Sblm), hippocampus CA1 (CA1), lateral hypothalamus (LH), basolateral amygdala (BLA), paraventricular thalamus (PVA), ventral pallidum (VP), and dorsal lateral septum (LSD). **(B)** Pie chart displaying the percentage of total retrobead+ cells coming from each region. Retrobead+ cells were summed across all mice (n=4 injections from 3 mice) and then divided by the number of positive cells in each region. **(C)** Percentage of total retrobead+ cells that were *Grp*+ in each of the examined regions. Bars represent mean +/- SEM, dots represent summed values from all slices in that region per 4 unique injection sites from 3 mice. **(D)** Pie chart showing the percentage of total retrobead+/*Grp*+ cells coming from each region, quantified across all mice (n=4 injections from 3 mice). **(E-H)** Representative FISH images from the indicated brain regions showing *Grp*+ cells in green and retrobead+ cells in magenta; **(E)** VTA, **(F)** subiculum, **(G)** CA1, and **(H)** BLA. **(I)** Percentage of retrobead+/*Grp*+ cells that were *Gad2*+ or *Vglut*+ for the four main *Grp*+ inputs into the NAc MSh (mean +/- SEM, dots represent summed value from all slices in that region per 3 unique injection sites from 2 mice). **(J-K)** Representative FISH images of the subiculum showing *Grp*+ cells in green, retrobead+ cells in magenta, and **(J)** *Gad2*+ or **(K)** *Vglut*+ cells in blue.

Figure 5

Figure 5: GRP depolarizes and increases the excitability of GRPR-expressing neurons in the NAc MSh.

(A) Representative trace showing the membrane potential over time of a nucleus accumbens medial shell (NAc MSh) *Grpr*-eGFP+ cell. The blue bar represents a ten-minute wash-on of 300 nM GRP. **(B)** Representative trace from a NAc MSh *Grpr*-eGFP+ cell in which 300 nM GRP was washed-on for 10 minutes in the presence of 1 µM DPDMB (red bar, entirety of recording). **(C)** Membrane potential of *Grpr*-eGFP+ NAc MSh neurons at baseline (black) and after 300 nM GRP wash-on (blue). Baseline value is the average of the two-minute period prior to GRP wash-on and 300 nM GRP value is the average of the final two minutes of the ten-minute 300 nM GRP wash-on; (n=10 cells from 7 mice; **p=0.0059, Wilcoxon test). **(D)** Membrane resistance of *Grpr*-eGFP+ NAc MSh neurons at baseline and after GRP wash-on. Baseline value is the average of five sweeps of VC RC check at the beginning of the recording and just following 300 nM GRP wash-on. (n=10 cells from 7 mice; **p=0.0098, Wilcoxon test). **(E)** Membrane potential of *Grpr*-eGFP+ NAc MSh neurons at baseline and after GRP wash-on in the presence of 1 µM DPDMB; (n=10 cells from 8 mice; p=0.3750, Wilcoxon test). **(F)** Membrane resistance of *Grpr*-eGFP+ NAc MSh neurons at baseline and after GRP wash-on in the presence of DPDMB; (n=10 cells from 8 mice; p=0.6953, Wilcoxon test). **(G)** Representative trace showing the membrane potential over time of a NAc MSh D2-GFP+ cell. Blue bar represents ten-minute wash-on of 300 nM GRP. **(H)** Representative trace showing the membrane potential over time of a NAc MSh D2-GFP+ cell. 300 nM GRP (blue bar) was washed-on in the presence of 1 µM DPDMB (red bar, entirety of recording). **(I)** Membrane potential of D2-GFP+ NAc MSh neurons at baseline (black) and after 300 nM GRP wash-on (blue); (n=20 cells from 20 mice; p=0.1054, Wilcoxon test). **(J)** Membrane resistance of D2-GFP+ NAc MSh neurons at baseline and after GRP wash-on; (n=20 cells from 20 mice; p=0.1429, Wilcoxon test). **(K)** Membrane potential of D2-GFP+ NAc MSh neurons at baseline and after GRP wash-on in the presence of DPDMB; (n=22 cells from 17 mice; p=0.0501, Wilcoxon test). **(L)** Membrane resistance of D2-GFP+ NAc MSh neurons at baseline and after GRP wash-on in the presence of DPDMB; (n=22 cells from 17 mice; **p=0.0025, Wilcoxon test).

Figure 6

Figure 6: *Grpr+* cells are more likely to be *Fos+ in vivo* and GRP injection increases *Fos* expression in *Grpr+* cells in the NAc MSh.

(A) Mean +/- SEM percent of *Drd2+*/*Grpr+* (magenta) vs. *Drd2+*/*Grpr-* (green) cells in the nucleus accumbens medial shell (NAc MSh) of wild-type (WT) mice that are *Fos+*. (n=8 mice; **p=0.0078, Wilcoxon test). **(B)** Representative image of the NAc MSh. Fluorescent *in situ* hybridization (FISH) for *Drd2* in green, *Grpr* in magenta, and *Fos* in cyan. Dashed circles outline *Grpr+* cells, white=*Grpr+*/*Drd2+*/*Fos-*, cyan=*Grpr+*/*Drd2+*/*Fos+*. **(C-D)** Same as A-B, but in *Grpr* KO mice (n=7 mice, p=0.0781, Wilcoxon test). **(E)** Schematic of the experiment. 3 μM GRP was injected unilaterally into the NAc MSh of WT mice. 45 minutes post injection, the mice were sacrificed and brains were harvested for FISH. **(F-G)** Representative images of NAc MSh from the injected hemisphere **(F)** and un-injected hemisphere **(G)**. FISH for *Drd2* in green, *Grpr* in magenta, and *Fos* in cyan. Nuclei are labeled with DAPI in blue. Dashed circles outline *Grpr+* cells, white=*Grpr+*/*Fos-*, cyan=*Grpr+*/*Fos+*. **(H)** Total number of cells in the NAc MSh that were *Fos+* in the un-injected (black) vs GRP-injected hemisphere (blue) summed from two coronal sections per mouse; (n=10 mice; **p=0.0098, Wilcoxon test). **(I)** Total number of cells in the NAc MSh that were *Grpr+* in the un-injected vs GRP-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; p=0.9043, Wilcoxon test). **(J)** Total number of *Grpr+* cells in the NAc MSh that were *Fos+* in the un-injected vs GRP-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; **p=0.0020, Wilcoxon test). **(K)** Percentage of *Grp*r+ cells in the NAc MSh that were *Fos+* in the un-injected vs GRP-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; **p=0.0039, Wilcoxon test). **(L)** Percentage of *Fos+* cells in the NAc MSh that were *Drd2+* and *Grpr+* in the un-injected vs GRP-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; **p=0.0020, Wilcoxon test). **(M)** Percentage of *Fos+* cells in the NAc MSh that were *Drd2+* and *Grpr-* in the un-injected vs GRP-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; p=0.7695, Wilcoxon test).

Figure 7

Figure 7: *Grpr* deletion in the NAc MSh enhances motivation in a progressive ratio task.

(A) Schematic of the experiment. *Grpr*$^{fl/fl \text{ or } fl/y}$ mice were injected bilaterally at P28-30 with 600 nL of AAV5-hSyn-GFP or AAV5-hSyn-CRE-GFP in the nucleus accumbens medial shell (NAc MSh). **(B)** Sample image of injection site in the NAc MSh showing AAV-CRE-GFP in green with nuclei labeled in blue with DAPI. **(C)** Schematic of progressive ratio (PR) lever pressing task and timeline for operant conditioning. RT = random timing, FR1 = fixed ratio 1. **(D)** Box plots of the number of days it took control (GFP injected, green) and *Grpr* cKO (CRE injected, black) mice to reach criterion on the lever pressing task (boxes represent the interquartile range (25-75%), lines denote the median, whiskers represent mix to max values). Mice reached criterion after two consecutive days of 20 or more lever presses in the fixed-ratio (FR1) sessions. Mann-Whitney test, p=0.4522 (n=22 GFP and n=14 CRE mice for all panels). **(E)** Mean +/- SEM weight of GFP and CRE mice just prior to starting food deprivation. Mann-Whitney test, p=0.8917. **(F)** Mean +/- SEM percentage of starting weight per mouse on the day the mouse completed the PR task. Mann-Whitney test, p=0.4192. **(G)** Survival function of total session time in operant boxes on PR testing day. The session ended when a mouse went three minutes without pressing the lever after pressing the lever at least one time. Mantel-Cox test, **p=0.0054. **(H)** Mean +/- SEM total lever presses in the PR session before timeout. Mann-Whitney test, **p=0.0077. **(I)** Mean +/- SEM breakpoint in the PR session before timeout. Breakpoint was defined as the number of lever presses for a single reward at which the mouse stopped pressing the lever for more than three minutes. Mann-Whitney test, **p=0.0060. **(J)** Mean +/- SEM of total session time during the PR task before timeout. Mann-Whitney test, *p=0.0381. **(K)** Raster plot of lever presses per individual mouse during the PR task. Each line represents a single lever press. Each row represents the lever-pressing data of an individual mouse. Green=GFP-injected control mice (n=22), Black=CRE injected *Grpr* cKO mice (n=14).

Supplementary Figure 1

Supplementary Figure 1: *Grpr* is expressed in striatal projection neurons in the dorsal striatum.

(A) Representative image of the dorsal striatum (DS). Fluorescent *in situ* hybridization (FISH) for *Gad2* in green and *Grpr* in magenta. Nuclei are labeled with DAPI in blue. Right panels show zoomed-in images of the boxed region. Dashed circles outline *Grpr*+ cells. **(B)** Mean +/- SEM percentage of *Grpr*+ cells in the DS that are *Gad2*+ in wild-type (WT) mice. (n=4 mice). **(C-E)** Representative images of DS. Nuclei labeled with DAPI in blue. Dashed circles outline *Grpr*+ cells. **(C)** FISH for *Chat* in green, *Grpr* in magenta, and *Ppp1r1b* in cyan. **(D)** FISH for *Drd2* in green, *Grpr* in magenta, and *Drd1* in cyan. **(E)** FISH for *Drd2* in green, *Grpr* in magenta, and *Penk* in cyan. **(F-J)** Mean +/- SEM percentage of *Grpr*+ cells in the DS of WT mice that are **(F)** *Chat*+ (n=7 mice), **(G)** *Ppp1r1b*+ (n=7 mice), **(H)** *Drd2*+ (n=12 mice), **(I)** *Drd1*+ (n=8 mice), or **(J)** *Penk*+ (n=8 mice).

Supplementary Figure 2

Supplementary Figure 2: GRPR-expressing neurons in the NAc MSh have different electrophysiological properties from cholinergic interneurons.

(A-B) Representative traces from a **(A)** *Grpr*-eGFP+ and **(B)** *Chat*-Cre+ neuron in the nucleus accumbens medial shell (NAc MSh) recorded at baseline (left) and following +40 pA current injection (right). **(C)** Mean +/- SEM membrane potential of NAc MSh neurons. *Grpr*-eGFP+ in green: n=30 cells from 15 mice and *Chat*-Cre+ in light blue: n=34 cells from 7 mice. (Welch's two-tailed t-test, ***p=0.0009). **(D)** Mean +/- SEM membrane resistance of NAc MSh neurons. *Grpr*-eGFP+ in green: n=33 cells from 15 mice and *Chat*-Cre+ in light blue: n=34 cells from 7 mice. (Welch's two-tailed t-test, ****p<0.0001). **(E)** Mean +/- SEM action potential (AP) half-width of NAc MSh neurons. *Grpr*-eGFP+ in green: n=23 cells from 10 mice and *Chat*-Cre+ in light blue: n=34 cells from 7 mice. (Welch's two-tailed t-test, ****p<0.0009). **(F)** Input-output curves showing the mean +/- SEM firing frequency of NAc MSh neurons in response to positive current steps of increasing amplitude. *Grpr*-eGFP+ in green: n=23 cells from 10 mice and *Chat*-Cre+ in light blue: n=34 cells from 7 mice. For panels **C-D** dots represent values for individual neurons. Data and traces for *Grpr*-eGFP+ cells are duplicated from Figure 3 for comparison.

Supplementary Figure 3

Supplementary Figure 3: DPDMB decreases the baseline excitability of GRPR-expressing neurons in the NAc MSh.

(A) Mean +/- SEM membrane potential of *Grpr*-eGFP+ nucleus accumbens medial shell (NAc MSh) neurons at baseline without (black) and in the presence of 1 µM DPDMB (red). Baseline value is the average of the two-minute period prior to GRP wash-on; (n=10 cells per condition from 7 mice with no DPDMB and 8 mice with DPDMB; p=0.1968, Mann-Whitney test). **(B)** Mean +/- SEM membrane resistance of *Grpr*-eGFP+ NAc MSh neurons at baseline without and in the presence of 1 DPDMB. Baseline value is the average of five sweeps of VC RC check at the beginning of the recording; (n=10 cells per condition from 7 mice with no DPDMB and 8 mice with DPDMB; *p=0.0232, Mann-Whitney test). **(C)** Mean +/- SEM membrane potential of D2-GFP+ NAc MSh neurons at baseline without and in the presence of DPDMB (no DPDMB: n=20 cells from 20 mice, with DPDMB: n=22 cells from 17 mice; p=0.8129, Mann-Whitney test). **(D)** Mean +/- SEM membrane resistance of D2-GFP+ NAc MSh neurons at baseline without and in the presence of DPDMB. (no DPDMB: n=20 cells from 20 mice, with DPDMB: n=22 cells from 17 mice; p=0.6092, Mann-Whitney test). For all panels, dots represent values for individual neurons.

Supplementary Figure 4

Supplementary Figure 4: Mice lacking *Grpr* show reduced *Fos* expression in the NAc MSh.

Grpr floxed mice were bred with CMV-Cre mice to delete *Grpr*. *Grpr* is on the X chromosome and only males were used for this analysis (*Grpr*$^{fl/y}$). **(A)** Genotyping strategy for detection of the *Grpr* knock-out (KO), WT, and floxed alleles. Three primers (red) were designed to detect the presence of the loxP site in the floxed allele (1-WT forward, 2-floxed reverse, and 3-WT reverse). Two primers (blue) were designed to detect the loxP scar in the KO allele (1-loxP forward and 2-loxP reverse). Two primers (purple) were designed to detect exon 2 (1-Exon 2 forward and 2-Exon 2 reverse). **(B)** PCR detecting the presence of the floxed or WT allele for *Grpr*. WT mice were expected to have a band at 573 bp and KO mice to have no band. **(C)** PCR detecting the loxP scar, indicating successful recombination. WT mice were not expected to have a band as the sequence is too large to be amplified, while KO mice should have a band at 298 bp. **(D)** PCR for the presence of *Grpr* exon 2. WT mice were expected to have a band at 206 bp, while KO mice should not have a band. **(E)** Mean +/- SEM percent *Grpr*+ cells in the nucleus accumbens medial shell (NAc MSh) that are *Fos*+ in *Grpr* WT (black) and *Grpr* KO (red) mice. (n=4 WT and n=7 KO mice, **p=0.0061, Mann-Whitney test). **(F-G)** Representative images of NAc MSh from **(F)** *Grpr* WT and **(G)** *Grpr* KO mice. Fluorescent *in situ* hybridization for *Grpr* in magenta and *Fos* in cyan. Dashed circles outline *Grpr*+ cells, magenta=*Grpr*+/*Fos*- and cyan=*Grpr*+/*Fos*+.

Supplementary Figure 5

Supplementary Figure 5: Saline injection into the NAc MSh does not induce *Fos* in *Grpr*-expressing cells.

(A-B) Representative images of nucleus accumbens medial shell (NAc MSh) from the **(A)** saline injected hemisphere and **(B)** un-injected hemisphere. Fluorescent *in situ* hybridization (FISH) for *Drd2* in green, *Grpr* in magenta, and *Fos* in cyan. Nuclei are labeled with DAPI in blue. **(C)** Total number of cells in the NAc MSh that were *Fos+* in the un-injected (black) vs saline-injected hemisphere (light blue) summed from two coronal sections per mouse; (n=10 mice; **p=0.0098, Wilcoxon test). **(D)** Total number of cells in the NAc MSh that were *Grpr+* in the un-injected vs saline-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; p=0.9414, Wilcoxon test). **(E)** Total number of *Grpr+* cells in the NAc MSh that were *Fos+* in the un-injected vs saline-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; p=0.9570, Wilcoxon test). **(F)** Percentage of *Grp*r+ cells in the NAc MSh that were *Fos+* in the un-injected vs saline-injected hemisphere summed from two coronal sections per mouse; (n=10 mice; p=0.7695, Wilcoxon test).

Supplementary Figure 6

Supplementary Figure 6: Mice lacking *Grpr* in the NAc MSh have increased locomotor activity but no differences in syllable usage the open field.

(A) Timeline of behavioral assays. **(B)** Schematic of the open field. **(C)** Total distance travelled in the one-hour session for control (GFP injected, green) and *Grpr* cKO (CRE injected, black). Mann-Whitney test, *p=0.0234 (n=23 GFP and 12 CRE mice for all panels). **(D)** Mean +/- SEM of total time spent in the center of the open field during the one-hour session. Center was defined as a 20 cm x 20 cm square in the middle of the arena. Mann-Whiney test, p=0.8506. **(E)** Mean +/- SEM of the total number of grooming bouts during the one-hour open field session. Mann-Whitney test, p=0.1723. **(F)** Frequency of syllable usage for control (GFP injected, green) and *Grpr* cKO (CRE injected, black) mice. Syllable descriptions: 0-left turn, 1-static, 2-forward, 3-left turn 2, 4-grooming, 5-right turn, 6-forward 2, 7-left turn 3, 8-forward left, 9-right turn 2, 10-forward 3, 11-grooming 2, 12-grooming 3, 13-grooming 4, 14-backward right, 15-backward left, 16-forward 4, 17-left turn 4, 18-grooming 5, 19-forward 5, 20-forward 6, 21-forward 7, 22-backward, 23-forward 8, 24-forward right, 25-right turn 3, 26-grooming 6, 27-left turn 5. Kruskal-Wallis with Dunn's correction for multiple comparisons, 0: p=0.1724,1: p=0.1540, 2: p=0.5629, 3: p=0.1498, 4: p=0.2520, 5: p=0.8470, 6: p=0.6170, 7: p=0.2589, 8: p=0.7552, 9: p=0.4081, 10: p=0.5152, 11: p=0.2021, 12: p=0.1498, 13: p=0.8276, 14: p=0.1540, 15: p=0.2021, 16: p=0.7552, 17: p=0.6745, 18: p=0.1782, 19: p=0.1540, 20: p=0.5186, 21: p=0.2021, 22: p=0.2180, 23: p=0.5214, 24: p=0.1540, 25: p=0.5629, 26: p=0.4716, 27: p=0.2180. **(G)** Average pose trajectories for the top six Keypoint-MoSeq syllables. Each trajectory includes ten poses, starting 165 ms before and ending 500 ms after syllable onset.

Supplementary Figure 7

Supplementary Figure 7: Mice lacking *Grpr* in the NAc MSh show no differences in the rotarod or sucrose preference tests.

(A) Schematic of the accelerating rotarod. **(B)** Mean +/- SEM of performance on trial 1 of the rotarod in control (GFP-injected, green) vs *Grpr* cKO (CRE-injected, black) mice. Mann-Whitney test, p=0.6700 (n=24 GFP and 14 CRE mice). **(C)** Mean +/- SEM of learning rate, measured as the slope of the linear regression of the performance per mouse across all trials. Mann-Whitney test, p=0.5201. (n is the same as in panel **B**) **(D)** Rotarod performance for GFP-injected and CRE-injected mice across all trials, measured as terminal velocity (mean +/- SEM). Two-way repeated measures ANOVA p-values are shown in the graph. (n is the same as for panel **B**). **(E)** Schematic of sucrose preference test. **(F)** Mean +/- SEM weight of GFP-injected and CRE-injected mice just prior to the sucrose preference test. Mann-Whitney test, p=0.9300 (n=21 GFP and 11 CRE mice). **(G)** Mean +/- SEM sucrose preference measured as the percentage of total liquid consumed that was 5% sucrose. Welch's two-tailed t-test, p=0.6785; (n is the same as for panel **F**).

Supplementary Figure 8

Supplementary Figure 8: Mice lacking *Grpr* in the NAc MSh show no differences in operant conditioning.

(A) Number of lever presses on each day of the fixed-ratio 1 (FR1) training for individual mice. Control (GFP-injected) mice are shown in green and *Grpr* cKO (CRE-injected) mice are shown in black. (n=22 GFP and 14 CRE mice for all panels) **(B)** Mean +/- SEM of total lever presses on the final two days of FR1 training in control (GFP-injected, green) vs *Grpr* cKO (CRE-injected, black) mice. Two-way repeated measures ANOVA: day x genotype p=0.0671, day ****p<0.0001, genotype p=0.3845. Holm-Sidak's multiple comparisons test, day 1: p=0.8596, day 2: p=0.1278.

References

Ade, K.K., Wan, Y., Chen, M., Gloss, B., and Calakos, N. (2011). An Improved BAC Transgenic Fluorescent Reporter Line for Sensitive and Specific Identification of Striatonigral Medium Spiny Neurons. Front Syst Neurosci *5*, 32.

Al-Hasani, R., McCall, J.G., Shin, G., Gomez, A.M., Schmitz, G.P., Bernardi, J.M., Pyo, C.-O., Park, S.I., Marcinkiewcz, C.M., and Crowley, N.A. (2015). Distinct subpopulations of nucleus accumbens dynorphin neurons drive aversion and reward. Neuron *87*, 1063-1077.

Anderson, K.D., and Reiner, A. (1990). Extensive co-occurrence of substance P and dynorphin in striatal projection neurons: an evolutionarily conserved feature of basal ganglia organization. Journal of Comparative Neurology *295*, 339-369.

Bateup, H.S., Santini, E., Shen, W., Birnbaum, S., Valjent, E., Surmeier, D.J., Fisone, G., Nestler, E.J., and Greengard, P. (2010). Distinct subclasses of medium spiny neurons differentially regulate striatal motor behaviors. Proceedings of the National Academy of Sciences *107*, 14845-14850.

Berridge, K.C., and Kringelbach, M.L. (2015). Pleasure systems in the brain. Neuron *86*, 646-664.

Bertran-Gonzalez, J., Bosch, C., Maroteaux, M., Matamales, M., Hervé, D., Valjent, E., and Girault, J.A. (2008). Opposing patterns of signaling activation in dopamine D1 and D2 receptor-expressing striatal neurons in response to cocaine and haloperidol. J Neurosci *28*, 5671-5685.

Bessa, J., Morais, M., Marques, F., Pinto, L., Palha, J.A., Almeida, O., and Sousa, N. (2013). Stress-induced anhedonia is associated with hypertrophy of medium spiny neurons of the nucleus accumbens. Translational psychiatry *3*, e266-e266.

Besson, M., Graybiel, A., and Quinn, B. (1990). Co-expression of neuropeptides in the cat's striatum: an immunohistochemical study of substance P, dynorphin B and enkephalin. Neuroscience *39*, 33-58.

Bolam, J., Wainer, B., and Smith, A. (1984). Characterization of cholinergic neurons in the rat neostriatum. A combination of choline acetyltransferase immunocytochemistry, Golgi-impregnation and electron microscopy. Neuroscience *12*, 711-718.

Bossert, J.M., Adhikary, S., St. Laurent, R., Marchant, N.J., Wang, H.-L., Morales, M., and Shaham, Y. (2016). Role of projections from ventral subiculum to nucleus accumbens shell in context-induced reinstatement of heroin seeking in rats. Psychopharmacology *233*, 1991-2004.

Castro, D.C., and Bruchas, M.R. (2019). A motivational and neuropeptidergic hub: Anatomical and functional diversity within the nucleus accumbens shell. Neuron *102*, 529-552.

Chung, C.Y., Seo, H., Sonntag, K.C., Brooks, A., Lin, L., and Isacson, O. (2005). Cell type-specific gene expression of midbrain dopaminergic neurons reveals molecules involved in their vulnerability and protection. Hum Mol Genet *14*, 1709-1725.

Cording, K.R., and Bateup, H.S. (2023). Altered motor learning and coordination in mouse models of autism spectrum disorder. Frontiers in cellular neuroscience *17*.

Costa, R.M., Cohen, D., and Nicolelis, M.A. (2004). Differential corticostriatal plasticity during fast and slow motor skill learning in mice. Current Biology *14*, 1124-1134.

Cox, J., and Witten, I.B. (2019). Striatal circuits for reward learning and decision-making. Nature Reviews Neuroscience *20*, 482-494.

Durieux, P.F., Schiffmann, S.N., and de Kerchove d'Exaerde, A. (2012). Differential regulation of motor control and response to dopaminergic drugs by D1R and D2R neurons in distinct dorsal striatum subregions. The EMBO journal *31*, 640-653.

Gallo, E.F., Meszaros, J., Sherman, J.D., Chohan, M.O., Teboul, E., Choi, C.S., Moore, H., Javitch, J.A., and Kellendonk, C. (2018). Accumbens dopamine D2 receptors increase motivation by decreasing inhibitory transmission to the ventral pallidum. Nature communications *9*, 1-13.

Gerfen, C.R. (2000). Molecular effects of dopamine on striatal-projection pathways. Trends in neurosciences *23*, S64-S70.

Gerfen, C.R., and Bolam, J.P. (2016). The neuroanatomical organization of the basal ganglia. In Handbook of behavioral neuroscience (Elsevier), pp. 3-32.

Gerfen, C.R., Engber, T.M., Mahan, L.C., Susel, Z., Chase, T.N., Monsma Jr, F.J., and Sibley, D.R. (1990). D1 and D2 dopamine receptor-regulated gene expression of striatonigral and striatopallidal neurons. Science *250*, 1429-1432.

Gerfen, C.R., and Surmeier, D.J. (2011). Modulation of striatal projection systems by dopamine. Annu Rev Neurosci *34*, 441-466.

Gong, S., Zheng, C., Doughty, M.L., Losos, K., Didkovsky, N., Schambra, U.B., Nowak, N.J., Joyner, A., Leblanc, G., and Hatten, M.E. (2003). A gene expression atlas of the central nervous system based on bacterial artificial chromosomes. Nature *425*, 917-925.

Greene, J.G., Dingledine, R., and Greenamyre, J.T. (2005). Gene expression profiling of rat midbrain dopamine neurons: implications for selective vulnerability in parkinsonism. Neurobiol Dis *18*, 19-31.

Guillaumin, M.C.C., and Burdakov, D. (2021). Neuropeptides as Primary Mediators of Brain Circuit Connectivity. Front Neurosci *15*, 644313.

Hermes, M.L., Kolaj, M., Coderre, E.M., and Renaud, L.P. (2013). Gastrin-releasing peptide acts via postsynaptic BB2 receptors to modulate inward rectifier K+ and TRPV1-like conductances in rat paraventricular thalamic neurons. J Physiol *591*, 1823-1839.

Hokfelt, T., Bartfai, T., and Bloom, F. (2003). Neuropeptides: opportunities for drug discovery. Lancet Neurol *2*, 463-472.

Hurley, S.W., and Carelli, R.M. (2020). Activation of infralimbic to nucleus accumbens shell pathway suppresses conditioned aversion in male but not female rats. Journal of Neuroscience *40*, 6888-6895.

Kemp, J.M., and Powell, T.P.S. (1971). The structure of the caudate nucleus of the cat: light and electron microscopy. Philosophical Transactions of the Royal Society of London B, Biological Sciences *262*, 383-401.

Klawonn, A.M., and Malenka, R.C. (2018). Nucleus Accumbens Modulation in Reward and Aversion. Cold Spring Harb Symp Quant Biol *83*, 119-129.

Kramer, D.J., Risso, D., Kosillo, P., Ngai, J., and Bateup, H.S. (2018). Combinatorial expression of Grp and Neurod6 defines dopamine neuron populations with distinct projection patterns and disease vulnerability. Eneuro *5*.

Kravitz, A.V., Freeze, B.S., Parker, P.R., Kay, K., Thwin, M.T., Deisseroth, K., and Kreitzer, A.C. (2010). Regulation of parkinsonian motor behaviours by optogenetic control of basal ganglia circuitry. Nature *466*, 622-626.

Kravitz, A.V., Owen, S.F., and Kreitzer, A.C. (2013). Optogenetic identification of striatal projection neuron subtypes during in vivo recordings. Brain research *1511*, 21-32.

Kroog, G.S., Sainz, E., Worland, P.J., Akeson, M.A., Benya, R.V., Jensen, R.T., and Battey, J.F. (1995). The Gastrin-releasing Peptide Receptor Is Rapidly Phosphorylated by a Kinase Other Than Protein Kinase C After Exposure to Agonist (∗). Journal of Biological Chemistry *270*, 8217-8224.

Kupchik, Y.M., Brown, R.M., Heinsbroek, J.A., Lobo, M.K., Schwartz, D.J., and Kalivas, P.W. (2015). Coding the direct/indirect pathways by D1 and D2 receptors is not valid for accumbens projections. Nature neuroscience *18*, 1230-1232.

Kupchik, Y.M., and Kalivas, P.W. (2017). The direct and indirect pathways of the nucleus accumbens are not what you think. Neuropsychopharmacology *42*, 369.

Larsen, P., Saermark, T., and Mikkelsen, J. (1989). An immunohistochemical demonstration of gastrin-releasing peptide (GRP) in the rat substantia nigra. Journal of Chemical Neuroanatomy *2*, 83-93.

Lee, K., Dixon, A., Gonzalez, I., Stevens, E., McNulty, S., Oles, R., Richardson, P., Pinnock, R., and Singh, L. (1999). Bombesin-like peptides depolarize rat hippocampal interneurones through interaction with subtype 2 bombesin receptors. The Journal of Physiology *518*, 791.

Li, Z., Chen, Z., Fan, G., Li, A., Yuan, J., and Xu, T. (2018). Cell-type-specific afferent innervation of the nucleus accumbens core and shell. Frontiers in neuroanatomy *12*, 84.

Liu, M.-Y., Yin, C.-Y., Zhu, L.-J., Zhu, X.-H., Xu, C., Luo, C.-X., Chen, H., Zhu, D.-Y., and Zhou, Q.-G. (2018). Sucrose preference test for measurement of stress-induced anhedonia in mice. Nature protocols *13*, 1686-1698.

Ma, Y.Y., Cepeda, C., Chatta, P., Franklin, L., Evans, C.J., and Levine, M.S. (2012). Regional and cell-type-specific effects of DAMGO on striatal D1 and D2 dopamine receptor-expressing medium-sized spiny neurons. ASN Neuro *4*.

Madisen, L., Zwingman, T.A., Sunkin, S.M., Oh, S.W., Zariwala, H.A., Gu, H., Ng, L.L., Palmiter, R.D., Hawrylycz, M.J., and Jones, A.R. (2010). A robust and high-throughput Cre reporting and characterization system for the whole mouse brain. Nature neuroscience *13*, 133-140.

Majumdar, I.D., and Weber, H.C. (2011). Biology of mammalian bombesin-like peptides and their receptors. Current Opinion in Endocrinology, Diabetes and Obesity *18*, 68-74.

Martin, P.D., and Ono, T. (2000). Effects of reward anticipation, reward presentation, and spatial parameters on the firing of single neurons recorded in the subiculum and nucleus accumbens of freely moving rats. Behavioural brain research *116*, 23-38.

Martínez-Hernández, J., Lanuza, E., and Martínez-García, F. (2012). Lesions of the dopaminergic innervation of the nucleus accumbens medial shell delay the generation of preference for sucrose, but not of sexual pheromones. Behavioural brain research *226*, 538-547.

Mathis, A., Mamidanna, P., Cury, K.M., Abe, T., Murthy, V.N., Mathis, M.W., and Bethge, M. (2018). DeepLabCut: markerless pose estimation of user-defined body parts with deep learning. Nature neuroscience *21*, 1281-1289.

Melzer, S., Newmark, E.R., Mizuno, G.O., Hyun, M., Philson, A.C., Quiroli, E., Righetti, B., Gregory, M.R., Huang, K.W., and Levasseur, J. (2021). Bombesin-like peptide recruits disinhibitory cortical circuits and enhances fear memories. Cell *184*, 5622-5634. e5625.

Merali, Z., McIntosh, J., and Anisman, H. (1999). Role of bombesin-related peptides in the control of food intake. Neuropeptides *33*, 376-386.

Meredith, G.E., Agolia, R., Arts, M.P., Groenewegen, H.J., and Zahm, D.S. (1992). Morphological differences between projection neurons of the core and shell in the nucleus accumbens of the rat. Neuroscience *50*, 149-162.

Moody, T., O'donohue, T., and Jacobowitz, D. (1981). Biochemical localization and characterization of bombesin-like peptides in discrete regions of rat brain. Peptides *2*, 75-79.

Moody, T.W., and Merali, Z. (2004). Bombesin-like peptides and associated receptors within the brain: distribution and behavioral implications. Peptides *25*, 511-520.

Nath, T., Mathis, A., Chen, A.C., Patel, A., Bethge, M., and Mathis, M.W. (2019). Using DeepLabCut for 3D markerless pose estimation across species and behaviors. Nat Protoc *14*, 2152-2176.

Nusbaum, M.P. (2017). Neurotransmission: peptide transmitters turn 36. J Exp Biol *220*, 2492-2494.

Olivetti, P.R., Balsam, P.D., Simpson, E.H., and Kellendonk, C. (2020). Emerging roles of striatal dopamine D2 receptors in motivated behaviour: implications for psychiatric disorders. Basic & clinical pharmacology & toxicology *126*, 47-55.

Pagani, M., Albisetti, G.W., Sivakumar, N., Wildner, H., Santello, M., Johannssen, H.C., and Zeilhofer, H.U. (2019). How gastrin-releasing peptide opens the spinal gate for itch. Neuron *103*, 102-117. e105.

Piggins, H.D., and Rusak, B. (1993). Electrophysiological effects of pressure-ejected bombesin-like peptides on hamster suprachiasmatic nucleus neurons in vitro. Journal of neuroendocrinology *5*, 575-581.

Pinnock, R., and Woodruff, G. (1991). Bombesin excites a subpopulation of 5-hydroxytryptamine-sensitive neurones in the rat dorsal raphe nucleus in vitro. The Journal of Physiology *440*, 55-65.

Platt, R.J., Zhou, Y., Slaymaker, I.M., Shetty, A.S., Weisbach, N.R., Kim, J.-A., Sharma, J., Desai, M., Sood, S., and Kempton, H.R. (2017). Chd8 mutation leads to autistic-like behaviors and impaired striatal circuits. Cell reports *19*, 335-350.

Poulin, J.F., Zou, J., Drouin-Ouellet, J., Kim, K.Y., Cicchetti, F., and Awatramani, R.B. (2014). Defining midbrain dopaminergic neuron diversity by single-cell gene expression profiling. Cell Rep *9*, 930-943.

Puighermanal, E., Castell, L., Esteve-Codina, A., Melser, S., Kaganovsky, K., Zussy, C., Boubaker-Vitre, J., Gut, M., Rialle, S., and Kellendonk, C. (2020). Functional and molecular heterogeneity of D2R neurons along dorsal ventral axis in the striatum. Nature communications *11*, 1957.

Roesler, R., Kent, P., Luft, T., Schwartsmann, G., and Merali, Z. (2014). Gastrin-releasing peptide receptor signaling in the integration of stress and memory. Neurobiology of learning and memory *112*, 44-52.

Roesler, R., and Schwartsmann, G. (2012). Gastrin-releasing peptide receptors in the central nervous system: role in brain function and as a drug target. Frontiers in endocrinology *3*, 159.

Roitman, M.F., Wheeler, R.A., and Carelli, R.M. (2005). Nucleus accumbens neurons are innately tuned for rewarding and aversive taste stimuli, encode their predictors, and are linked to motor output. Neuron *45*, 587-597.

Rossi, J., Balthasar, N., Olson, D., Scott, M., Berglund, E., Lee, C.E., Choi, M.J., Lauzon, D., Lowell, B.B., and Elmquist, J.K. (2011). Melanocortin-4 receptors expressed by cholinergic neurons regulate energy balance and glucose homeostasis. Cell Metab *13*, 195-204.

Rothwell, P.E., Fuccillo, M.V., Maxeiner, S., Hayton, S.J., Gokce, O., Lim, B.K., Fowler, S.C., Malenka, R.C., and Südhof, T.C. (2014). Autism-associated neuroligin-3 mutations commonly impair striatal circuits to boost repetitive behaviors. Cell *158*, 198-212.

Russo, A.F. (2017). Overview of Neuropeptides: Awakening the Senses? Headache *57 Suppl 2*, 37-46.

Schwenk, F., Baron, U., and Rajewsky, K. (1995). A cre-transgenic mouse strain for the ubiquitous deletion of loxP-flanked gene segments including deletion in germ cells. Nucleic Acids Res *23*, 5080-5081.

Serchov, T., van Calker, D., and Biber, K. (2016). Sucrose preference test to measure anhedonic behaviour in mice. Bio-protocol *6*, e1958-e1958.

Shumyatsky, G.P., Tsvetkov, E., Malleret, G., Vronskaya, S., Hatton, M., Hampton, L., Battey, J.F., Dulac, C., Kandel, E.R., and Bolshakov, V.Y. (2002). Identification of a signaling network in lateral nucleus of amygdala important for inhibiting memory specifically related to learned fear. Cell *111*, 905-918.

Smith, S.J., Sumbul, U., Graybuck, L.T., Collman, F., Seshamani, S., Gala, R., Gliko, O., Elabbady, L., Miller, J.A., Bakken, T.E.*, et al.* (2019). Single-cell transcriptomic evidence for dense intracortical neuropeptide networks. Elife *8*.

Sun, S., Xu, Q., Guo, C., Guan, Y., Liu, Q., and Dong, X. (2017). Leaky gate model: intensity-dependent coding of pain and itch in the spinal cord. Neuron *93*, 840-853. e845.

Sun, Y.-G., and Chen, Z.-F. (2007). A gastrin-releasing peptide receptor mediates the itch sensation in the spinal cord. Nature *448*, 700-703.

Sun, Y.-G., Zhao, Z.-Q., Meng, X.-L., Yin, J., Liu, X.-Y., and Chen, Z.-F. (2009). Cellular basis of itch sensation. Science *325*, 1531-1534.

Taha, S.A., and Fields, H.L. (2005). Encoding of palatability and appetitive behaviors by distinct neuronal populations in the nucleus accumbens. J Neurosci *25*, 1193-1202.

Tai, L.H., Lee, A.M., Benavidez, N., Bonci, A., and Wilbrecht, L. (2012). Transient stimulation of distinct subpopulations of striatal neurons mimics changes in action value. Nat Neurosci *15*, 1281-1289.

Tejeda, H.A., Wu, J., Kornspun, A.R., Pignatelli, M., Kashtelyan, V., Krashes, M.J., Lowell, B.B., Carlezon, W.A., and Bonci, A. (2017). Pathway-and cell-specific kappa-opioid receptor modulation of excitation-inhibition balance differentially gates D1 and D2 accumbens neuron activity. Neuron *93*, 147-163.

Tepper, J.M., Tecuapetla, F., Koós, T., and Ibáñez-Sandoval, O. (2010). Heterogeneity and diversity of striatal GABAergic interneurons. Frontiers in neuroanatomy *4*, 150.

Trifilieff, P., Feng, B., Urizar, E., Winiger, V., Ward, R.D., Taylor, K.M., Martinez, D., Moore, H., Balsam, P.D., and Simpson, E.H. (2013). Increasing dopamine D2 receptor expression in the adult nucleus accumbens enhances motivation. Molecular psychiatry *18*, 1025-1033.

Viereckel, T., Dumas, S., Smith-Anttila, C.J., Vlcek, B., Bimpisidis, Z., Lagerström, M.C., Konradsson-Geuken, Å., and Wallén-Mackenzie, Å. (2016). Midbrain gene screening identifies a new mesoaccumbal glutamatergic pathway and a marker for dopamine cells neuroprotected in Parkinson's disease. Scientific Reports *6*, 35203.

Voorn, P., Vanderschuren, L.J., Groenewegen, H.J., Robbins, T.W., and Pennartz, C.M. (2004). Putting a spin on the dorsal–ventral divide of the striatum. Trends in neurosciences *27*, 468-474.

Weinreb, C., Pearl, J., Lin, S., Osman, M.A.M., Zhang, L., Annapragada, S., Conlin, E., Hoffman, R., Makowska, S., Gillis, W.F.*, et al.* (2023). Keypoint-MoSeq: parsing behavior by linking point tracking to pose dynamics. bioRxiv.

Wiltschko, A.B., Tsukahara, T., Zeine, A., Anyoha, R., Gillis, W.F., Markowitz, J.E., Peterson, R.E., Katon, J., Johnson, M.J., and Datta, S.R. (2020). Revealing the structure of pharmacobehavioral space through motion sequencing. Nature neuroscience *23*, 1433-1443.

Wolf, S.S., Moody, T.W., O'Donohue, T.L., Zarbin, M.A., and Kuhar, M.J. (1983). Autoradiographic visualization of rat brain binding sites for bombesin-like peptides. European Journal of Pharmacology *87*, 163-164.

Wulff, A.B., Cooper, P., Kodjo, E., Abel, E., and Thompson, S.M. (2023). How sucrose preference is gained and lost: An in-depth analysis of drinking behavior during the sucrose preference test in mice. Eneuro *10*.

Yin, H.H., Mulcare, S.P., Hilário, M.R., Clouse, E., Holloway, T., Davis, M.I., Hansson, A.C., Lovinger, D.M., and Costa, R.M. (2009). Dynamic reorganization of striatal circuits during the acquisition and consolidation of a skill. Nature neuroscience *12*, 333-341.

Yu, Y.-Q., Barry, D.M., Hao, Y., Liu, X.-T., and Chen, Z.-F. (2017). Molecular and neural basis of contagious itch behavior in mice. Science *355*, 1072-1076.

Zarbin, M.A., Kuhar, M.J., O'Donohue, T.L., Wolf, S.S., and Moody, T.W. (1985). Autoradiographic localization of (125I-Tyr4) bombesin-binding sites in rat brain. Journal of Neuroscience *5*, 429-437.

Zhou, F.M., Wilson, C.J., and Dani, J.A. (2002). Cholinergic interneuron characteristics and nicotinic properties in the striatum. Journal of neurobiology *53*, 590-605.