The genomic signature and transcriptional response of metal tolerance

in brown trout inhabiting metal-polluted rivers

Running title: Metal tolerance in brown trout

Authors:

Josephine R Paris[1], R Andrew King[1], Joan Ferrer Obiol[2], Sophie Shaw[,3,4], Anke Lange[1],

Vincent Bourret[5], Patrick B Hamilton[6], Darren Rowe[1], Lauren V Laing[1], Audrey Farbos[1],

Karen A Moore[1], Mauricio A Urbina[7,8], Ronny van Aerle[9,10] Julian M Catchen[11], Rod W

Wilson[1,10], Nicolas R Bury[12], Eduarda M Santos[1,10*] and Jamie R Stevens[1*]

Affiliations:

1. Department of Biosciences, University of Exeter, United Kingdom

2. Dipartimento di Scienze e Politiche Ambientali, Università degli Studi di Milano, Italy

3. Centre for Genome Enabled Biology and Medicine, University of Aberdeen, United
Kingdom

4. All Wales Medical Genomics Service, Institute of Medical Genomics and Vale University
Health Board, Cardiff, United Kingdom

5. Université du Québec à Trois-Rivières, Trois-Rivières, QC, Canada

6. Department of Clinical and Biomedical Sciences, University of Exeter, United Kingdom

7. Departamento de Zoología, Facultad de Ciencias Naturales y Oceanográficas,
Universidad de Concepción, Concepción, Chile

8. Instituto Milenio de Oceanografía (IMO), Universidad de Concepción, PO Box 1313,
Concepción, Chile

9. Centre for Environment, Fisheries and Aquaculture Science (Cefas), Weymouth, United
Kingdom

10. Sustainable Aquaculture Futures, University of Exeter, United Kingdom

11. Department of Evolution, Ecology, and Behavior, University of Illinois at Urbana-
Champaign, USA

12. Institute for Life Sciences, University of Southampton, United Kingdom

Corresponding Authors: Josephine R Paris (parisjosephine@gmail.com), Eduarda M

Santos * (e.santos@exeter.ac.uk) and Jamie R Stevens * (j.r.stevens@exeter.ac.uk);

*shared senior authors

Abstract

Industrial pollution is a major driver of ecosystem degradation, but it can also act as a driver of contemporary evolution. As a result of intense mining activity during the Industrial Revolution, several rivers across the southwest of England are polluted with high concentrations of metals. Despite the documented negative impacts of ongoing metal pollution, brown trout (*Salmo trutta* L.) survive and thrive in many of these metal-impacted rivers. We used population genomics, transcriptomics, and metal burdens to investigate the genomic and transcriptomic signatures of potential metal tolerance. RADseq analysis of six populations (originating from three metal-impacted and three control rivers) revealed strong genetic substructuring between impacted and control populations. We identified selection signatures at 122 loci, including genes related to metal homeostasis and oxidative stress. Trout sampled from metal-impacted rivers exhibited significantly higher tissue concentrations of cadmium, copper, nickel, and zinc, which remained elevated after 11 days in metal-free water. After depuration, we used RNAseq to quantify gene expression differences between metal-impacted and control trout, identifying 2,042 differentially expressed genes (DEGs) in the gill, and 311 DEGs in the liver. Transcriptomic signatures in the gill were enriched for genes involved in ion transport processes, metal homeostasis, oxidative stress, hypoxia and response to xenobiotics. Our findings reveal shared genomic and transcriptomic pathways involved in detoxification, oxidative stress responses, and ion regulation. Overall, our results demonstrate the diverse effects of metal pollution in shaping both neutral and adaptive genetic variation, whilst also highlighting the potential role of constitutive gene expression in promoting metal tolerance.

Keywords: adaptation, freshwater, pollution, RADseq, RNAseq, toxic metals

56   Introduction

Pollution causes long-term environmental impacts with severe implications for biodiversity
and human health (Schwarzenbach et al., 2010; Travis & Hester, 1991), yet growing
evidence of tolerance to toxic chemicals has resulted in a greater awareness of pollution in
mediating contemporary evolution (Hendry et al., 2017; Smith & Bernatchez, 2008;
Whitehead, 2014). Whilst organisms can respond to short-term stress via phenotypic
plasticity, understanding the mechanisms underpinning tolerance to chronic exposure
requires consideration of long-term adaptive responses (Badyaev, 2005; Ghalambor et al.,
2007). However, because pollutants are often not acutely toxic by design, have wide-ranging
physiological effects, and occur in complex mixtures in the environment (involving different
metals in varying concentrations and forms, which can interact with each other), dissecting
the role of pollutants in mediating adaptive responses can be difficult, especially in natural
populations (Whitehead et al., 2017).

Metals are naturally occurring elements, but are found at significantly higher concentrations
in ecosystems associated with certain human activities such as mining, agriculture and
industrial processes (Vareda et al., 2019). These activities have led to elevated levels in
aquatic environments, in some cases exceeding the mortality thresholds for many species
(see Kumar et al., 2019). Essential metals, such as copper, iron, manganese and zinc, are
crucial for life, acting as key components for various biological processes including
development, energy metabolism and immunity, but can become toxic when threshold
concentrations are exceeded (Chandrapalan & Kwong, 2021). On the other hand, non-
essential metals, such as arsenic and cadmium, lack known biological function, and thus
toxicity manifests when such metals are present in the environment in particular chemical
states at lower concentrations (Wood et al., 2012a). Metals exert their toxic effects through
different mechanisms, including the production of reactive oxygen species (ROS), DNA
damage, and the disruption or inhibition of enzymes, proteins, and subcellular organelles
(Hartwig et al., 2002; Valko et al., 2005; Vangronsveld & Clijsters, 1994). Therefore,
adaptation to metal polluted environments would require physiological mechanisms that
reduce toxicity while maintaining levels of essential metals.

Over the last four decades, a wealth of information has been published on the impact of
metals on fish including: discerning the role of different tissues in metal uptake, storage and
detoxification (*e.g.* Bradley & Morris, 1986; Eastwood & Couture, 2002); understanding the
physiological processes regulating homeostasis and toxicity (Shahjahan et al., 2022; Wood
et al., 2012a, 2012b); and characterising the molecular mechanisms underlying metal
homeostasis (Hamilton & Mehrle, 1986; Srikanth et al., 2013). At the genomic level, studies

have evaluated the impact of metal pollution on genetic structure and diversity (Bourret et

al., 2008; Maes et al., 2005), as well as genetic adaptation (Hamilton et al., 2017; Laporte et

al., 2016). Further, a number of studies have demonstrated that metal pollution also affects

epigenetic processes, with long-term impacts on the health of individuals and populations

(Abdelnour et al., 2024; Pierron et al., 2023). Meanwhile, at the transcriptomic level, several

studies have quantified differences in gene transcription between metal-impacted and

control fish (Bélanger-Deschênes et al., 2013; Coffin et al., 2022; Pierron et al., 2009; Uren

Webster et al., 2013), highlighting genes and pathways potentially involved in the mitigation

of metal toxicity. However, measuring gene expression from individuals sampled directly

from their native sites does not allow for a distinction between short-term responses to acute

stress, versus responses to chronic stress (López-Maury et al., 2008). In this regard,

constitutive gene expression, *i.e.*, the higher baseline expression of certain genes or

pathways, can confer protection to stressors by providing a pre-emptive adaptive response

(Hendry, 2016; Rivera et al., 2021). Investigations aimed at understanding the impacts of

metal pollution, and the evolution of potentially adaptive responses, would benefit from a

multifaceted approach combining physiology, genomics and transcriptomics.

Brown trout (*Salmo trutta* L.) in the southwest of England provide one of the best purported

cases of metal tolerance in aquatic environments. Mining in the area dates back to the

Bronze Age (ca. 2500-800 BCE), and later, during the Industrial Revolution (1850-1900),

the region dominated the global output of tin, copper and arsenic (Burt et al., 2014;

Timberlake, 2017). As a testament to its role in shaping Britain's industrial development and

its broader influence on the global mining industry, the area has been designated as a

UNESCO World Heritage Site (The Cornwall and West Devon Mining Landscape;

https://www.cornishmining.org.uk/). The longevity and intensity of mining has resulted in

elevated contemporary metal concentrations in several rivers in the region (Rainbow, 2020),

and consequently, negative ecological and biological effects including a reduction or total

extermination of aquatic life (Brown, 1977; Bryan et al., 1987; Hart et al., 2020). Yet brown

trout are abundant, and are able to reproduce and survive in many of these rivers, despite

the fact that they experience both chronic and seasonally-varying acute exposure to metals

at concentrations known to be lethal to naïve salmonids (see Table S1). Based on neutral

genetic markers, trout from metal-impacted rivers were shown to be genetically differentiated

from trout sampled from nearby control rivers, and displayed significantly reduced genetic

diversity (Osmond et al., 2024; Paris et al., 2015). Another study showed that trout from

metal-impacted rivers had significantly elevated tissue-metal burdens (Uren Webster et al.,

2013). This opens the question of how these brown trout cope with metal toxicity.

Transcriptomics analysis comparing metal-impacted trout and control fish sampled directly

from the river highlighted differentially expressed genes involved in metal and ion

homeostasis, oxidative stress, and immunity (Uren Webster et al., 2013). However, the

extent to which these signatures represent a short-term response to metal toxicity is

unknown. To investigate the signatures of potentially adaptive responses, genetic variation

should be surveyed using genome-wide markers, and transcriptional responses should be

quantified in fish outside of the metal-impacted environment.

Here, we investigate the signatures of potential long-term adaptation to metal pollution in

wild brown trout using a combination of genomics, transcriptomics, and tissue-metal

burdens. First, we use RADseq to genotype six brown trout populations (originating from

three metal-contaminated and three control rivers) to assess how metal pollution has shaped

the neutral and non-neutral genetic make-up of metal-impacted populations. We

hypothesised that if metals have caused population-level impacts, we would observe

distinctive patterns of genetic variation in metal-impacted populations. We also reasoned

that if metal pollution has driven an adaptive response, selection signatures at outlier loci

related to metal handling will be evident. Second, we aimed to evidence and assess, if and

how metals are absorbed into the tissues of metal-impacted brown trout. We address this by

quantifying differences in metal burdens across five tissues (gill, liver, gut, kidney, and

muscle) between metal-impacted trout and those from a matched control site. To refine our

understanding of metal handling, we quantified tissue-burdens from individuals sampled

directly from the river, and after an 11-day depuration in control (metal-free) water. Finally,

we quantified gene expression differences between metal-impacted and control trout

following the 11-day period of depuration in two key tissues known for their roles in metal

uptake and detoxification (gill and liver) to assess the molecular processes involved in

potential long-term responses to metal pollution. Integrating across these approaches

provides a broad understanding of adaptation to metals, and of the mechanisms

underpinning evolution to environmental stress in wild fish.

Materials and Methods

2.1 Study Sites and Sampling

The Cornwall and West Devon (United Kingdom) mining region contains many small rivers

affected by metal pollution. While many rivers are affected by high concentrations of metals,

individual rivers are polluted with different mixtures of metal pollutants (Figure 1A). For

clarity, we term trout sampled from these rivers as "metal-impacted". To reduce confounding

effects of biogeographic history, we sampled what we term "control" trout from rivers with

lower concentrations of metals sampled within the same region. Due to the nature of the

bedrock and extensive history of mining in the region, we note that control rivers may also

have been historically impacted by metals, but in recent years show comparatively lower

metal concentrations (Table S2).

For the population genomics analyses, a total of $n = 120$ trout were sampled across 14

sampling sites from a total of six rivers ($n = 20$ per river; Figure 1A, B; Table S2; Table S3).

The largest adult fish (> 150 mm fork-length) were sampled to ensure resident origin (*i.e.,*

not anadromous individuals straying from nearby rivers) and prolonged survival in the metal-

polluted rivers. Fish were anaesthetised using a stock solution of benzocaine (100 g/L in

ethanol) diluted to 50 mg/L in river water prior to adipose fin-clip removal. Tissue samples

were stored in 95% ethanol at 4° C prior to DNA extraction.

For the quantification of tissue burdens and transcriptomics analysis, trout (0+ parr) were

sampled from the River Hayle (metal-impacted, $n = 26$) and the River Fowey (control, $n =$

26). Fish from the River Hayle were chosen as representative metal-impacted trout as they

showed the most distinct genetic signatures in our analyses, suggesting minimal gene flow

with populations from neighbouring rivers over multiple generations. This population has also

been characterised and studied previously (Minghetti et al., 2014; Osmond et al., 2024; Paris

et al., 2015; Uren Webster et al., 2013). Trout from the River Fowey were selected as control

fish since the river exhibits relatively low metal contamination and the most comparable

water chemistry (which is a key determinant of metal bioavailability) to that of the River

Hayle (Table S4).

All fish were caught by backpack electrofishing, following Home Office (UK) guidelines,

Environment Agency (UK) authorisation, and relevant landowner permissions.

Figure 1. Schematic of collection sites and experimental design. (**A**) Location within the UK of the
Cornwall and West Devon mining region and location of sampling sites. Metal-impacted sites are
coloured in red and control sites are coloured in blue. Filled coloured shapes represent sites sampled
for tissue-metal burden analysis before and after a period of depuration. The heatmap shows the
scaled metal concentrations of each river (averaged over a 15-year period; full data available in Table
S2) (**B**) Timepoint 1: River. For population genomics, $n = 20$ brown trout were sampled from three
metal contaminated rivers: Hayle (red triangle); Red River (red circle); Gannel (red square) and three

control rivers: Camel (blue square); Fowey (blue circle) and Carrick Roads (blue triangle). (**C**)

Timepoint 1: River. For tissue burden analysis, metal-impacted trout were sampled from the Hayle

(red triangle; $n = 10$) and control trout were sampled from the Fowey (blue circle; $n = 10$). (**D**)

Timepoint 2: Depuration. Trout were sampled after 11 days of depuration in control water for tissue

burden analysis ($n = 10$ per group) and transcriptomic analysis of the gill and liver using RNAseq ($n =$

6 per group).

Trout were transported to the Aquatic Resources Centre, University of Exeter, in water from

their river of origin. Immediately upon arrival (on the same day of sampling), we randomly

selected 10 metal-impacted and 10 control fish (from the n = 26 fish sampled per site) to

represent the status of individuals in the river (Timepoint 1: River; Figure 1C). After 11 days

of depuration and acclimation to laboratory conditions (in control water), we randomly

sampled 10 metal-impacted and 10 control fish (Timepoint 2: Depuration; Figure 1D). The

11-day depuration period was chosen based on a balance between metal elimination

kinetics (*i.e.*, rate constants; see Veltman et al., 2008) and the challenges of maintaining wild

brown trout in aquaria.

Fish were humanely killed via an overdose of benzocaine (0.5 g/L), followed by destruction

of the brain. We recorded the wet body mass and fork length, and the condition factor was

calculated (K) = [mass (g) x 100] / [fork length (cm)^3] before dissecting and weighing tissues

for analysis. As a growing number of studies have demonstrated sex-related differences in

ecotoxicological responses (e.g. Burger et al., 2007), we attempted to account for any

potential sex-related biases in our metal-burden and RNAseq analyses. Due to the young

age of the fish, gonadal sex could not be reliably assessed, so additional muscle tissue was

collected and stored in 95 % ethanol at -20 °C for genetic sex determination (King &

Stevens, 2020). All animal work was carried out in accordance with the UK Animals

Scientific Procedures Act (ASPA) 1986 and was ethically approved by the Animal Welfare

and Ethical Review Body at the University of Exeter.

2.2 Laboratory depuration and acclimation: water chemistry and fish

maintenance

In the laboratory, 7-9 fish were housed in 39 L glass tanks. For the depuration period, fish

were held in synthetic control (metal-free) water to mimic the water chemistry of the River

Hayle (Minghetti et al., 2014) using salts of the following concentrations: 391 µM NaCl, 128

µM $KHCO_3$, 500 µM $MgSO_4$, 350 µM $Ca(SO_4)$, 75 µM $Ca(NO_3)_2$. Salts were made up to a

50x stock concentration, dosed into constant-flowing reverse osmosis purified water using a

conductivity pump set to dose between 265 and 270 µS/cm. Water chemistry of the synthetic

freshwater was assessed via daily pH and conductivity measurements. Freshwater was

delivered via a flow-through system at a flow rate of 100 ml/min. The laboratory was

temperature controlled at 15°C and a natural day-night 12h:12h light cycle was used.

Individuals were fed daily to satiation.

2.3 Population genomics using RADseq

2.3.1 DNA extraction and RADseq library preparation

DNA was extracted from adipose fin tissue using the QIAGEN DNeasy Blood & Tissue Kit

(QIAGEN). DNA concentrations were determined using the QuantiFluor ONE dsDNA assay

system (Promega). Single-digest *SbfI* RADseq libraries were prepared following the protocol

described by Etter et al. (2011). Libraries were amplified using 12 PCR cycles and were

sequenced on an Illumina HiSeq 2500 (100bp single-end) at a target coverage of 20x per

individual.

2.3.2 RADseq data pre-processing

Data were processed using Stacks v2.6 (Rochette et al., 2019). Raw reads were

demultiplexed and cleaned using *process_radtags*. Optimised parameters for construction of

de novo RAD loci were determined using the *r80* method (Paris et al., 2017). Seven of the

120 individuals were removed due to low coverage. We assembled loci *de novo* (minimum

stack depth (m) = 3, distance between stacks (M) = 2, distance between catalog loci (n) = 2)

and then aligned loci to the brown trout genome (Ensembl: version fSalTru1.1; Hansen et al.,

2021) using *stacks-integrate-alignments* (Paris et al., 2017) to provide positional information.

We used stringent parameters for filtering the loci, in order to detect top candidates for

potential metal adaptation. Loci were filtered with the following parameters in *populations*: *-r*

80, -p 6, -min-maf 0.1, -max-obs-het 0.8. For analyses sensitive to linkage, we used the –

write-single-snp option and pruned loci in linkage (LD dataset) using Plink v1.9 (Purcell et

al., 2007).

2.3.3 Population structure, genetic diversity, and effective population size estimation

Population structure was assessed by Principal Components Analysis (PCA) and the

neighbour-joining (NJ) tree method. PCA was conducted in Plink v1.9 (Purcell et al., 2007).

For constructing the NJ tree, the VCF file was converted to Phylip format using *vcf2phylip*

(Ortiz, 2019), requiring a minimum of 100 samples per SNP. Distances were calculated

using the *dist.dna* function in ape v5.6 (Paradis & Schliep, 2019), using the K80 substitution

model and removing pairwise missing data.

Expected heterozygosity (H_E), observed heterozygosity (H_O) and the inbreeding coefficient

(F_{IS}) were calculated for each population using Hierfstat (Goudet, 2005). To assess variance

in individual genetic diversity, we calculated the proportion of polymorphic sites across

positions that were variable in at least one population using the --het option in VCFtools

v0.1.16 (Danecek et al., 2011). Pairwise Weir and Cockerham's F_{ST} were computed in

Hierfstat (Goudet, 2005). To perform an assessment of fixed alleles within each population,

we calculated allele frequencies using VCFtools v0.1.16 (Danecek et al., 2011) and plotted

the number of fixed alleles per population in R v4.0 (R Core Team, 2021).

Biases in estimating the effective population size (N_e) using linkage-disequilibrium methods

are well-known (see Waples, 2024), especially when using RADseq and low sample sizes

(Marandel et al., 2020). To alleviate these biases, we used the recently described methods

implemented in currentNE, which performs particularly well in the small sample size scenario

(Santiago et al., 2024). currentNE was run by estimating the average number of full siblings

(k parameter) from the input data (a Hardy-Weinberg equilibrium and missing data filtered

VCF file per population), using 40 chromosomes as estimated from the brown trout

reference genome. 90% confidence intervals were provided by the neural network (ANN).

2.3.4 Identification of loci associated with metal pollution and water chemistry

To identify candidate genes potentially involved in metal adaptation, we used a combination

of statistical methods to detect loci showing high allele frequency differentiation: pcadapt

v4.3.3 (Luu et al., 2017) and OutFLANK v0.2 (Whitlock & Lotterhos, 2015), and genotype-

environment association (GEA) tests: Redundancy Analysis (RDA; Capblancq et al., 2018)

and Latent Factor Mixed Models (LFMM v1.5; Frichot et al., 2013). pcadapt was run using *K*

= 4, as identified using the *screeplot* function. p-values were corrected for multiple testing

using an alpha value of 0.05. For OutFLANK, left and right trim fractions were set to 0.35

and 0.06, respectively. We removed low heterozygosity variants (< 0.1) and used a q-value

threshold of 0.05.

For GEA tests, we used 41 water chemistry and metal contamination variables for each of

the six rivers (14 sample sites; Table S3). We used the average value for each variable (data

covers a 15-year period; Table S2). Due to microgeographic differences in metal

contamination within each river, we extracted allele frequency data for each of the 14

sampling sites to match the environmental data (4664 loci).

RDA was run adapting the approaches described by Bourret et al., (2014) and Forester et

al., (2018). To avoid collinearity amongst the environmental predictors, the environmental

data were reduced to 10 Rotated Components (RC) by running a PCA (10 factors, *varimax*

rotation). RCs were reduced to a number of factors identified by a parallel analysis criterion

(Horn, 1965) using 1000 bootstraps. We used a 0.01 significance threshold (0.68 loading

cutoff) to associate an environmental parameter to a factor. Following the RDA, a global

analysis of variance (*anova.cca*; 999 permutations) was performed to assess the global

significance of the RDA and an analysis by axis was run to determine if retained RCs were
significantly correlated with population-level allele frequencies. Candidates were identified as
variants in the tails of the RDA constrained axis distribution. A cutoff of 2.5 standard
deviations (two-tailed p-value of 0.012) was used.

LFMM was run using the RCs generated above. The number of latent factors was estimated
using the *snmf* function ($K = 4$). We ran 10,000 iterations with a burn-in of 5000. The Gibbs
sampler was run five independent times on the dataset in order to estimate the appropriate
average LFMM parameters (z-score values). p-values were adjusted using estimation of the
genomic inflation factor (λ) for each RC variable. Significance was tested using an FDR <
0.05.

Candidate loci (*i.e.*, all loci identified from the analyses described above) were assessed for
functional gene-level effects using the Variant Effect Predictor tool in Ensembl (McLaren et
al., 2016). We also classified the potential gene-level effects of the fixed alleles in each of
the metal-impacted populations. Variants were classified as intergenic, intronic,
synonymous, and non-synonymous, including the potential variant consequences.

2.4 Tissue-metal burden

2.4.1 Laboratory analysis

Five tissues (gill, liver, gut, posterior kidney, and muscle) were dissected, freeze-dried and
weighed, and stored at -20 °C. Tissues were digested in 500µl of nitric acid (70 %, purified
by redistillation, ≥ 99.999 % trace metals basis, Sigma Aldrich) for 48 hours at room
temperature (~20 °C), followed by digestion with 0.1 % hydrogen peroxide (Fisher; Hydrogen
Peroxide 30% (w/v) Extra Pure SLR) for a further 24 hours. Digested tissue was diluted 1:10
in ultrapure deionised H_2O and acidified with 0.1 % nitric acid. A standard of 10 µg/L yttrium
(Y) was added to each sample as an internal standard to perform within-sample corrections.
Metal concentrations (quantified as ng/mg) were measured by Inductively Coupled Plasma
Mass Spectrometry (ICP-MS) (E:AN 6100DRC, Perkin-Elmer, Cambridge, UK).

2.4.1 Statistical analysis

Data were checked for normality (Shapiro test) and homogeneity of variance (Levene test)
using the stats package (v3.6.1) in R v4.0 prior to conducting statistical tests and were log
transformed when necessary. Assessment of burdens for each tissue were statistically
assessed using a two-way ANOVA, with 'metal-impacted' and 'control' fish specified as
Group, and Timepoint ('River', or 'Depuration') specified as Factor. For analyses where
statistical differences were identified, post-hoc Tukey tests were performed to conduct
pairwise multiple comparisons, accepting a significance threshold of FDR ≤ 0.05.

2.5 Transcriptomics analysis

2.4.1 RNA extraction, library construction, and sequencing

Gill and liver tissues were dissected from euthanised fish ($n = 6$ metal-impacted; $n = 6$

control) and were snap-frozen in liquid nitrogen and stored at -80°C prior to analysis. To

avoid sex-biassed gene expression differences, we selected three (genetically-determined)

females and three males from each group. RNA was extracted using a QIAGEN RNeasy

extraction kit (QIAGEN), including an on-column DNAse digestion step. RNA quantification

and quality assessments (accepting a RIN > 8.0) were performed using a Bioanalyser

(Agilent RNA 6000 Nano). Libraries were prepared using the Illumina TruSeq Stranded

mRNA library preparation protocol. Concentrations of the cDNA libraries were quantified

using a Tapestation (D1000 ScreenTape) and final library pool concentrations were

determined by qPCR. Libraries were sequenced (125 bp paired-end) on an Illumina HiSeq

2500.

2.4.2 RNAseq data pre-processing

Raw reads were cleaned using Trimmomatic v0.27 (Bolger et al., 2014) and aligned to the

brown trout reference genome (Ensembl version fSalTru1.1; Hansen et al., 2021) using

STAR v2.5.4 (Dobin et al., 2012). Transcript abundances were calculated at gene regions

using featureCounts v2.0.1 (Liao et al., 2014). Genes with low read counts were removed by

only including genes with more than 10 counts in ≥ 6 samples. This reduced the initial

dataset of 43,946 genes to 28,261 and 21,820 expressed genes in the gill and liver,

respectively.

2.4.2 Differential gene expression analysis

Analysis of differentially expressed genes (DEGs) was performed using DESeq2 v1.40.2

(Love et al., 2014). For assessing clustering of the biological replicates, we performed PCA

on a variance stabilising transformed (VST) normalised count matrix. Size factors and

dispersion were estimated using the default DESeq2 parameters. A negative binomial GLM

was used to test for DEGs, retaining only those genes with a significance threshold FDR ≤

0.05 and log2foldchange values of ≤ -1 or ≥ 1.

All genes (including non-significant DEGs, but with dispersion outliers removed) were used

to test for a statistical enrichment in Gene Ontology (GO) terms via Gene Set Enrichment

Analysis (GSEA) using clusterProfiler v3.16.1 (Yu et al., 2012). Lists containing GO

categorisation of brown trout terms (using the Brown Trout Genes fSalTru1.1 database;

AH85410) were downloaded from Ensembl using BioMart v2.44 (Smedley et al., 2009).

Genes were ranked by a single ranking metric by multiplying the log fold-change by the

negative logarithm (base 10) of the adjusted p-value. This was done to ensure that genes

with both large fold-changes and low p-values receive higher combined ranks. Minimum and

maximum gene set sizes of 20 and 200 were used. GO terms with a Benjamini–Hochberg

adjusted p-value ≤ 0.25 were classified as being significantly enriched or underrepresented.

367   Results

368   3.1 Population genomics

369   3.1.1 RADseq libraries and loci construction

370   We generated an average of 2,279,882 reads (SEM ± 119,860) per individual at an average

371   of 20X coverage, assembling 10,115 SNPs in 8,718 loci genotyped across 80% of the

372   individuals from each of the six rivers. Details of the RADseq libraries, including number of

373   reads and coverage obtained for each individual are shown in Table S5.

374   3.1.2 Population genetics characterisation of metal-impacted and control trout

Overall, we observed strong genetic sub-structuring with the highest pairwise F_{ST} values

detected in comparisons to metal-impacted trout populations, in particular those sampled

from the River Hayle ($F_{ST} > 0.1$ in all comparisons) and the lowest pairwise F_{ST} values

(range: River Camel-River Fowey $F_{ST} = 0.015$ to Carrick Roads-River Fowey; $F_{ST} = 0.03$)

were observed between control trout populations (Figure 2A).

PCA separated the metal-impacted trout populations from one another, and also from the

control trout (Figure 2B). PC1 (18.8%) defined trout from the River Hayle, PC2 (8.97%)

separated the Red River and River Gannel trout. PC3 (8.09%) further separated the River

Gannel trout and also revealed patterns of IBD in the control trout populations. In the NJ

tree, metal-impacted trout showed three distinctive branches, with the River Hayle and Red

River individuals originating from a shared branch in the NJ tree, whereas the River Gannel

formed its own group (Figure 2C). The River Hayle and Red River showed particularly long

branches, indicative of higher substitution rates. We also observed genetic substructuring

within the River Hayle (upstream and downstream of a highly metal-contaminated middle

region; Durrant et al., 2011; Paris et al., 2015) and the Red River (by tributary) in the NJ tree.

We observed differences in the genetic diversity and effective population size (N_e) estimates

between metal-impacted and control trout (Figure 2D; Figure 2E). Compared to control sites,

genetic diversity was significantly lower (Mann-Whitney U $p < 0.005$) in metal-impacted trout

sampled from the Red River ($H_O = 0.28$) and the River Hayle ($H_O = 0.23$), but not in the River

Gannel ($H_O = 0.31$). The proportion of polymorphic sites also showed a more even

distribution in the control trout, whereas long tails were observed in the River Hayle and Red

River, indicative of high variance among the individuals sampled from these sites (Figure

2D). Further investigation of the heterozygosity per sampling site showed that this high

variance was not caused by microgeographic differences in the upstream and downstream

reaches of the River Hayle, or by the two sampled tributaries in the Red River. Differences in

observed heterozygosity were reflected in the inbreeding coefficient (F_{IS}), which was overall

higher in metal-impacted trout compared to control trout, being highest in trout sampled from

the River Hayle (F_{IS} = 0.25), and lowest in the River Fowey (F_{IS} = 0.04).

Figure 2. Population genomic data describing the structure and diversity of metal-impacted and

control trout. (**A**) Heatmap of pairwise F_{ST} calculated between all sampling sites. (**B**) Principal

Component Analysis (PCA) showing PC1 (~19% of the variance) against PC2 (~9% of the variance)

and PC1 against PC3 (~8% of the variance). Individuals are coded by sampling site as in panel A. (**C**)

Neighbour-joining tree from aligned SNPs, computed using the K80 substitution model. (**D**) Proportion

of polymorphic sites for each individual sampled from each of the six sampling sites. (**E**) Estimates of

the effective population size (N_e) for each of the six sampling sites. (**F**) Quantification of fixed alleles

calculated at each of the six sampling sites. (**G**) Redundancy Analysis (RDA; p = 0.007, R^2 = 0.2145)

showing RDA1 (p = 0.037) and RDA2 (p = 0.1). Red vectors represent RC1 (water chemistry features;

Table S7) and RC2 (Cd, Cu, Ni, Zn). In all panels, sampling sites are coloured by metal-impacted

(red) or control (blue). For panels A, B, C, and F the sampling sites are coloured and shaped as:

Hayle (red triangle); Red River (red circle); Gannel (red square); Camel (blue square); Fowey (blue

circle) and Carrick Roads (blue triangle).

N_e estimates (Figure 2E) for the River Hayle (N_e = 68; 90% CI: 43-106) and Red River (N_e =

64; 90% CI: 40-100) were low, but the River Gannel (N_e = 366; 90% CI: 157-863) had a

higher N_e than the River Fowey (N_e = 186; 90% CI: 87-398) and Carrick Roads (N_e = 137;

90% CI: 77-243), and was comparable to that of the River Camel (N_e = 372; 90% CI: 157-

881). Overall, metal-impacted trout did not show a higher number of fixed alleles, except for

the River Hayle, which showed a much higher number compared to trout from all other sites

(Figure 2F).

3.1.3 Identification of adaptive loci

Using four methods, we identified a total of 122 candidate loci potentially involved in metal

adaptation (Table S6). Of the pcadapt candidates (n = 40), the majority were associated with

PC1 (n = 18, which separates the River Hayle). Remaining outliers were associated with

PC2 (n = 14; separates the Red River), and PC3 (n = 8; separates the River Gannel).

OutFLANK identified eight candidates, which were primarily driven by divergent F_{ST} values

observed in comparisons to the River Hayle.

The RDA was globally significant (p = 0.007; R^2 = 0.2145) and both axes were also

significant (RDA axis 1: p = 0.037; RDA axis 2: p = 0.1, accepting a significance threshold ≤

0.1; Figure 2G). We identified a total of 30 loci which were significantly associated with these

axes (two-tailed p = 0.012). RDA axis 1 (five loci) separated the River Hayle, and to a lesser

extent, the Red River and River Gannel from the control group. RDA axis 2 (25 loci)

separated the Red River and River Gannel populations from the control group, suggesting

this axis represents loci which are diverging in these populations, but not in the River Hayle.

Correlations between the loadings on the RCs and the RDA axes showed that RDA axis 1

was primarily associated with the environmental features of RC2: Cd, Cu, Ni and Zn (Table

S7), which are the primary metal pollutants of the River Hayle. RDA axis 2 was primarily

associated with the environmental features of RC1 (water chemistry features; Table S7).

Using LFMM, we uncovered significant associations of 44 loci (RC1: 22 loci; RC2: 14 loci;

RC3: 8 loci. We found no overlap amongst the loci identified using the four outlier

approaches.

Of the 122 candidates, 60 loci showed good-quality alignments (MQ=60) to genes in the

brown trout genome (Table S6). Twenty-eight of the variants were intronic, 21 variants were

located upstream or downstream of genes, three were synonymous variants, and nine

variants had functional gene-level predictor effects. Gene functions were varied, but we

identified candidates potentially involved in metal homeostasis, and response to oxidative

stress and xenobiotics, as well as genes involved in calcium homeostasis, tumour regulation,

transcriptional regulation, and immunity (see Discussion). Of the 1348 fixed variants

identified in each of the metal-impacted populations, 42 aligned to genes in the brown trout

genome. All variants were found to be intronic, except for a missense variant in the River

Hayle, located in an open reading frame (Table S8). Amongst the intronic variants, we

identified a gene involved in protein ubiquitination (*usp2b*), a potassium channel regulator

(*kcnab1b*), an iron ion binding gene (*iscub*), and an aryl hydrocarbon receptor protein

(BMAL2).

3.2 Laboratory maintenance and fish morphometrics

In the laboratory, all fish were observed feeding; there were no mortalities and no changes in

behaviour were observed. Stable water chemistry of the synthetic freshwater was

maintained throughout the experiment (Table S9). Metal-impacted trout were significantly

larger than control fish; however, there was no significant difference in condition factor

(Table S10). The metal-impacted and control groups showed uneven numbers of males and

females and this was taken into consideration during the statistical comparisons (Timepoint

1: metal-tolerant M:F 2.67, control M:F 1.5; Timepoint 2: metal-tolerant M:F 1.4, control M:F

1.17).

3.3 Tissue-metal burden analysis

No differences were observed between Timepoint 1 and Timepoint 2 for any metal in any

tissue in the control trout. Sex did not affect the statistical significance of the metal-burden

data. Full statistical results for the metal-burden data can be found in Table S11.

At Timepoint 1 (river), gill metal concentrations were 7-, 6-, 1- and 2- fold higher in metal-

impacted trout compared to control trout for Cd ($p < 0.0001$), Cu ($p < 0.0001$), Ni ($p = 0.01$)

and Zn ($p < 0.0001$), respectively (Figure 3A). In metal-tolerant trout, the concentrations of

Cd, Cu and Ni decreased during the depuration period (Timepoint 2) (Cd, $p = 0.0003$; Cu, $p

< 0.0001$; Ni, $p = 0.01$), but Zn did not ($p = 1$). Although the concentration of Cd decreased in

Timepoint 2, it was still higher than the concentrations found in control trout at the same

timepoint ($p < 0.0001$). Zn burdens also remained higher than control trout at Timepoint 2 ($p

< 0.0001$).

In the gut, metal burdens at Timepoint 1 were 14-, 11- and 2-fold higher in metal-impacted

trout for Cd ($p < 0.0001$), Cu ($p < 0.0001$) and Zn ($p < 0.0001$), respectively (Figure 3B). Ni

burden was not higher ($p = 0.99$) and did not decrease at Timepoint 2 ($p = 0.98$). The

concentrations of both Cu and Zn decreased at Timepoint 2 (Cu, $p < 0.0001$, Zn, $p < 0.0001$

to concentrations comparable to those observed in control trout (Cu, $p = 0.83$; Zn, $p = 0.68$).

Cd concentrations did not decrease (p = 0.56), with 18-fold higher concentrations being

measured in metal impacted fish compared to control trout after Timepoint 2 (p < 0.0001).

In the kidney, metal burdens measured in fish at Timepoint 1 were 27- and 23-fold higher in

metal-impacted trout for Cd (p < 0.0001) and Cu (p < 0.0001) (Figure 3C). Although both

metals showed a significant depuration (Cd, p < 0.0001; Cu, p = 0.0002), they remained 17-

and 11-fold higher than control trout at Timepoint 2 (Cd, p < 0.0001; Cu, p = 0.006). Ni was

higher in metal-impacted trout, but this was marginally non-significant in comparison to

control trout (p = 0.09). No statistical differences were observed for Zn.

In the liver, metal burdens were 21-, 30-, 2-, and 2-fold higher in metal-impacted trout at

Timepoint 1 for Cd (p < 0.0001), Cu (p < 0.0001), Ni (p = 0.001) and Zn (p < 0.0001),

respectively (Figure 3D). Metal-impacted trout showed a depuration of all metals at

Timepoint 2 (Cd, p < 0.0001, Cu, p < 0.0001, Ni, p = 0.002; Zn, p < 0.0001). However,

concentrations of Cd and Cu remained 7- and 17-fold higher than control trout at Timepoint 2

(Cd, p = 0.003; Cu, p < 0.0001).

In the muscle, metal-impacted trout showed 6-fold higher Cd burden concentrations

compared to control trout (Cd, p < 0.0001) when measured at Timepoint 1, and Cd did not

depurate (Cd, p = 0.05) (Figure 3E). Although slightly higher burdens of Cu were observed in

metal-impacted trout, this was not significant (p = 0.41); Cu did however depurate (p = 0.03).

Ni and Zn did not show any significant differences.

Figure 3. Tissue-metal burden data of cadmium (Cd), copper (Cu), nickel (Ni), and zinc (Zn)
measured in metal-impacted (in red) and control (in blue) trout measured at Timepoint 1 (River) and
Timepoint 2 (Depuration). Metal burden is measured as ng/mg of dry weight tissue for the (**A**) Gill; (**B**)
Gut; (**C**) Kidney; (**D**) Liver; and (**E**) Muscle. Significance was measured using a Two-way ANOVA (full
details in Table S11). Only significant results from pairwise Tukey tests are included where ****
indicates p≤0.0001, *** indicates p≤0.001, ** indicates p≤0.01, and * indicates p≤0.05.

510 *3.3 Comparison of the transcriptional landscapes between metal-*
511 *impacted and control trout following a period of depuration*

512 *3.3.1 RNAseq libraries and gene expression profiles*

Across the six samples per group (metal-impacted and control), we generated an average of
16,549,620 (SEM ± 1,341,005) reads for the gill and 11,588,849 (SEM ± 1,329,081) reads
for the liver (Table S12). PCA revealed a clear separation between metal-impacted and
control trout in their transcription profiles after depuration. In the gill, replicates from each
group were separated along PCA axis 1 (49% of the variation; Figure 4A), whereas in the
liver they were separated along PCA axis 2 (21% of the variation; Figure 4B).

519 *3.3.2 Identification of differentially expressed genes*

Analysis of the differentially expressed genes (DEGs) revealed 2,042 DEGs (FDR ≤ 0.05) in
the gill, of which 1,180 were over-expressed (log2FC ≥ 1) and 862 were under-expressed
(log2FC ≤ -1) in metal-impacted trout compared to control trout (Figure 4C; see Table S13
for the full list of significant DEGs). We identified several metal-binding and antioxidant
genes that remained over-expressed after depuration in the gill tissue of metal-impacted
trout, including metallothionein 2 (*mt2*), metal response element binding transcription factor 2
(*mtf2*), metalloreductase (*steap4*), methionine sulfoxide reductase B2 (*msrb2*), thioredoxin-
related transmembrane protein 2 (*tmx2a*), three glutathione-dependent enzymes (*tstd1*,
mgst3, *gstz1*), and lysyl oxidase (*loxl2a*, *loxl2b*). We also identified genes with well-known
roles in response to oxidative stress and response to xenobiotics, including an epoxide
hydrolase, two cytochrome p450 genes, five cytochrome c oxidase subunits, a cytochrome
b-245 chaperone (*cybc1*), both aryl hydrocarbon receptors (*ahr1a*, *ahr1b*), and fourteen
heat-shock proteins. Related specifically to the known activity of the gill in response to
metals, we also identified ATP-binding cassette (ABC) transporters (*abca4a*, *abcg2a*,
abcc5), and two transcripts encoding carbonic anhydrase 4 (*ca4*). Amongst the under-
expressed genes, we observed a high number of immune-system related genes, including
interferon-induced proteins, interleukins, and E3 ubiquitin protein ligases.

In the liver, 311 DEGs (FDR ≤ 0.05) were identified, of which 200 were over-expressed
(log2FC ≥ 1), and 111 were under-expressed (log2FC ≤ -1) in metal-impacted trout
compared to control trout after the depuration period (Figure 4D; see Table S14 for the full
list of DEGs). We did not observe the same level of transcriptional changes in the liver, with
very few over-expressed metal-binding, oxidative stress response, or xenobiotic handling
genes. However, the metal-binding protein *heavy metal-binding protein HIP-like*, the
oxidative stress gene superoxide dismutase 3 (*sod3*), and the well-known xenobiotic

metabolism gene, cytochrome P450 1A (*cyp1a*) were over-expressed in the metal-impacted
group.

Intersecting the gill and liver DEGs, we identified 25 shared over-expressed genes, and eight
shared under-expressed genes (Table S15). Of note, *DENN domain-containing protein 5B-*
like was over-expressed in both tissues and was the most strongly over-expressed gene in
both tissues.

Figure 4. Differential gene expression between metal-impacted and control trout. (**A**) Principal
Components Analysis (PCA) of the variance stabilising transformation (VST) normalised counts in the
gill. (**B**) Principal Components Analysis (PCA) of the VST normalised counts in the liver. Coloured
dots represent the transcriptomic profile of individual fish from the metal impacted (red) and control
(blue) rivers. (**C**) Volcano plot of the 28,261 genes measured in the gill. (**D**) Volcano plot of the 21,820
genes measured in the liver. Coloured dots in the volcano plots correspond to differentially expressed

genes (DGEs) with FDR ≤ 0.05 and a log2foldchange of 1 (red) or -1 (blue), meaning genes are over-
expressed, or under-expressed in metal-impacted trout, respectively.

3.3.3 Gene Set Enrichment Analysis (GSEA) on Gene Ontology (GO) terms

To further investigate the transcriptomic landscapes of metal-impacted trout following
depuration, we explored enriched and underrepresented GO terms via GSEA. In the gill, we
identified several enriched biological processes indicative of changes in ionoregulation
including chloride, calcium, sodium, and potassium ion transporters, and response to
oxidative stress, including regulation of small GTPase mediated signal transduction.
Underrepresented biological processes included microtubule movement, DNA repair and
replication, immune responses, protein folding, catabolism, and ubiquitination. Amongst the
enriched molecular function pathways, we identified ion channel activity, including voltage-
gated potassium and calcium channel activity, metallocarboxypeptidase activity,
cytochrome-c oxidase activity, and copper ion binding and iron sulphur cluster binding. The
identified GO terms associated with both enriched and underrepresented processes in the
gill dataset are detailed in Table S16.

In the liver, we found a weak concerted biological signal related to metal tolerance. We
found no significantly enriched or underrepresented biological pathways, but cytochrome-c
oxidase activity was a significantly enriched molecular function. The identified GO terms
associated with both enriched and underrepresented processes in the liver dataset are
detailed in Table S17.

Discussion

We applied an integrative approach, using genomics, transcriptomics, and tissue-metal
burdens, to investigate the impacts of, and potential long-term adaptation to metal pollutants
in wild brown trout. Population genomics analysis confirmed that metal-impacted trout are
highly genetically differentiated from control trout, and that two of the three populations show
significantly lower genetic diversity and low N_e, highlighting the effects of metal pollutants on
the genetic make-up of these populations. Using a series of outlier tests, we identified
selection signatures for loci associated with metal homeostasis, and response to oxidative
stress and xenobiotics. Measured immediately after removal from the river, metal-impacted
trout showed considerably higher concentrations of metals across multiple tissues,
suggesting the requirement of mechanisms to cope with increased metal burden. After 11
days in control water, patterns of metal depuration were tissue- and metal-specific, pointing
towards a complex network of biological responses involved in metal handling. Analysis of
gill transcriptional profiles revealed over-expressed genes and enriched biological processes

in metal-tolerant trout involved in ionoregulation, metal homeostasis and oxidative stress
providing a mechanistic understanding of metal tolerance.

4.1 Evidence of high tissue concentrations of metals in metal-impacted
trout

Tissue-metal burden analysis in individuals sampled at their river of origin showed
significantly higher concentrations of metals in the tissues of metal-impacted trout compared
to control fish, similarly to that reported in Uren Webster et al., (2013) on fish collected from
comparable locations. The varying levels of metal concentrations across the different tissues
points to the diverse roles of these organs in organismal level metal uptake, storage, and
detoxification. The gills, being in intimate contact with the water, and the gut, via ingestion of
metal-contaminated prey, are well-characterised as the main routes of metal uptake in fish
(Wood, 2012). In the gills, we found elevated concentrations of all metals measured in this
study (Cd, Cu, Ni and Zn), pointing to this tissue as the main entry site. Meanwhile, elevated
concentrations of Cd, Cu and Zn in the gut suggests brown trout are also acquiring metals
from ingested food in the river. However, Ni was not elevated in the gut; previous research
has shown that pre-exposure to Ni reduces gastrointestinal uptake, suggesting a
homeostatic interaction might exist for this metal (Chowdhury et al., 2008).

The liver and kidney provide key functions in the detoxification and excretion of metals in fish
(Wood, 2012). All metals measured (Cd, Cu, Ni and Zn) were significantly elevated in the
liver, supporting its role as the main tissue of metal storage and detoxification in vertebrates.
In contrast, only Cd and Cu were elevated in the kidney. As reported here, the kidney has
been previously found to accumulate the highest concentrations of Cd (Andres et al., 2000;
De Smet et al., 2001). Muscle generally shows a weak accumulating potential for metals (Jia
et al., 2017), and we found that only Cd was significantly elevated within this tissue.

Overall, the observation of elevated tissue burdens of metals in trout living in metal-impacted
environments supports the hypothesis that physiological mechanisms, likely influenced by
genomic and/or transcriptomic mechanisms must be involved in the dynamics of internal
metal storage and detoxification, and in preventing the harmful effects associated with such
high metal loads.

4.2 Metal pollutants impact the genetic structure and reduce genetic
diversity of metal-impacted trout

We observed strong genetic sub-structuring between metal-impacted and control trout
across all analyses. Such patterns have been observed previously using both neutral
microsatellites (Paris et al., 2015) and SNPs (Osmond et al., 2024). Despite their

geographic proximity, metal-impacted trout populations showed strong genetic separation,

both from one another, and also from control trout. In contrast, differentiation was low

among control trout occupying clean rivers, which are geographically much more distant

from one another. For example, the mouths of the River Hayle and Red River (metal-

impacted) are separated by just ~9 km, whereas the River Camel and the Carrick Roads

(control) are separated by ~270 km of coastline. Population genetic studies on salmonids

often report a high incidence of population structure (Bekkevold et al., 2020; King et al.,

2020; King et al., 2024), frequently attributed to processes of local adaptation (Fraser et al.,

2011; Primmer, 2011; Taylor, 1991). However, identifying what proportion of the genetic

structure is explained solely by local adaptation is confounded by neutral demographic

processes. Yet it can also be argued that if populations are geographically close, but

experience substantially different environmental pressures, the expected differentiation due

to non-selective factors is small (Wadgymar & DeMarche, 2022; Whitlock & Lotterhos,

2015). The low genetic differentiation observed among control populations also supports

the local adaptation hypothesis. Moreover, there are no known physical barriers separating

the mouths of the metal-impacted rivers, and thus the strong population structure could also

indicate that non-adapted individuals which stray into metal-impacted rivers suffer adverse

toxic effects (Sergeant et al., 2022). Over time, a reduction in gene flow would be expected

to further increase genetic drift and strengthen genetic differentiation.

The effect (neutral and/or non-neutral) of metal pollutants was particularly marked in the

River Hayle, where we observed micrographic structuring based on metal contamination

levels, with trout sampled upstream and downstream of an extremely toxic middle section of

the river (~1.5km), showing strong genetic differentiation (Paris et al., 2015). As well as

indicating that metal toxicity may act as a chemical barrier to migration within the river

(Hecht et al., 2007; Williams & Gallagher, 2013), these microgeographic differences could

also indicate finer-scale adaptive responses. Indeed, primary gill cell cultures showed

higher metallothionein gene expression patterns when exposed to River Hayle water

downstream from the toxic section (Minghetti et al., 2014).

Metal-impacted populations do not appear to be experiencing the same genetic impact in

response to metal pollutants. The River Hayle, the Red River, and the River Gannel, are all

impacted by different combinations of metal pollutants and also differ in their water

chemistry features (which affect metal bioavailability). This result could stem from a number

of factors: metal toxicity at different concentration thresholds within each river; disparate

adaptive responses in each of the metal-impacted populations; differences in the population

size and demographic history of these populations. Previous analysis using Approximate

Bayesian Computation (ABC) showed that bottlenecks occurred in all metal-impacted

populations (Paris et al., 2015). The River Hayle and Red River trout appeared to be most

affected by metal pollution, with these populations showing the strongest population

structure and particularly long branches in the neighbour-joining tree. The latter observation

indicates higher substitution rates in these populations, which can be the result of

multifarious biological processes, including higher levels of inbreeding, drift, episodes of

heightened adaptive evolution, but also higher mutation rates, or differences in life-history

traits such as generation time (Mendes & Hahn, 2016). Increased mutation rates have been

evidenced in metal-impacted sunfish (*Lepomis auratus*; Theodorakis et al., 2006), and

metal-impacted yellow perch (*Perca flavescens*) had lower longevity than control

conspecifics (Couture & Pyle, 2008). Genetic diversity and N_e were also lower in the River

Hayle and Red River populations, but particularly so in the former, which also had a higher

inbreeding coefficient. Such observations support the hypothesis that environmental

pollution can cause significant genetic erosion (van Straalen & Timmermans, 2002).

Nevertheless, the fact that trout remain in these rivers, and show evidence of extremely

high concentrations of metals in their tissues, suggests an adaptive component to metal

tolerance.

4.3 Assessment of candidate loci involved in local adaptation to metal

pollution

To detect candidate loci underpinning potential metal tolerance we applied several

statistical tests, each with different assumptions, using a particularly rigorously set of filtered

loci and stringent cut-off criteria for significance. This was to reduce spurious correlations

between loci, environmental variables, and the strong population structure we observed

(Ahrens et al., 2018; Hoban et al., 2016). Amongst the 122 outlier loci, we identified 12

variants with potential gene-level effects, several of which have functional annotations with

potential relevance to metal adaptation. We identified a splice-gene variant in *map35k*,

which is known to be activated by various stressors including ROS and calcium overload

(Ichijo et al., 1997) and is regulated in response to copper (O'Doherty et al., 2020). In

addition, we identified a synonymous variant in *scara3* which is involved in scavenging

oxidative molecules (Han et al., 1998; Sone et al., 2010) and a missense variant in *ferritin,*

heavy subunit-like which is involved in the storage of iron in a nontoxic state (Munro &

Linder, 1978), both potentially contributing to a reduction of oxidative stress caused by

elevated intracellular metal ion concentrations. We also identified a stop-loss mutation in

cyp7a1, one of the cytochrome P450 enzymes involved in mono-oxygenation reactions in a

wide range of endogenous and exogenous compounds (Uno et al., 2012); and which has

been shown to be regulated in response to toxic metals (Chi et al., 2019). Genes involved

in transcriptional regulation were also identified including a 3'-UTR variant in ZNF239 and a

missense variant in PCGF3. Further, we identified a splice donor variant in *hmmr*, which

regulates cell growth (He et al., 2020) and is modified in exposure to arsenic (Dodmane et

al., 2013), and missense variants in *mfhas1* and *inavaa* which both function in innate

immunity: *mfhas1* is a regulator of *tlr2* and *tlr4* (Shi et al., 2017; Zhong et al., 2015), and

inavaa is stimulated by pattern recognition receptors including *tlr2* and *tlr4* (Yan et al.,

2017). Whilst these genes represent promising candidates underpinning local adaptation to

metals, we recognise that we have not surveyed the full variation potentially responsible for

metal tolerance due to, for example, limitations in the RADseq approach (Lowry et al.,

2017, but see Catchen et al., 2017), and neglecting the role of structural variants (SVs;

Mérot et al., 2020), especially duplications arising from the salmonid whole-genome

duplication event (Moriyama & Koshiba-Takeuchi, 2018). Further research using whole-

genome sequencing methods are required to increase our understanding of potential

candidates involved in adaptation to metals in these trout populations.

4.4 Metal- and tissue-specific depuration patterns over time

We observed both metal- and tissue- specific patterns of depuration, pointing to differential

mechanisms for each of the measured metals for deposition and excretion. The patterns of

depuration in the gill, liver and kidney are of particular interest. In the gill, Cd, Cu and Ni

decreased in concentration over time, but Zn did not, and although Cd depurated from the

gill, its concentration remained higher than that observed in control fish. Toxic

concentrations of all metals can have deleterious effects on the gill, including oxidative

stress (Sevcikova et al., 2011), impaired ionoregulation (Wood et al., 2012a, 2012b),

damage and mucus accumulation (Playle, 1998), necessitating their rapid detoxification and

removal (Saibu et al., 2018). Rapid clearance (within 16h) of Cd and Cu from the gill has

been reported previously in rainbow trout (*Oncorhynchus mykiss*), with accompanying

increases in the liver (Handy, 1992). Indeed, metal-impacted trout continued to show high

burdens of all metals in the liver, but especially Cd and Cu, both of which showed evidence

of depuration, but not to comparable concentrations observed in the control fish. We

observed similar patterns for Cd and Cu in the kidney, suggesting overall that the gill is

essential in detoxifying these metals, and that they are subsequently transported to the liver

and kidney for excretion (Garai et al., 2021). We also observed a high retention of Cd

across all tissues, which has been observed previously, and over much longer time periods

than our experiment allowed for (Defo et al., 2015; Kraemer et al., 2005).

As others have shown, these results indicate that tissue metal concentrations can be high,

but the metals within the tissue are likely present in a non-toxic form (*e.g.*, Campbell et al.,

2008; Leonard et al., 2014; Shekh et al., 2021; Wang & Rainbow, 2006). The physiological

processes governing this are attributed to subcellular partitioning, whereby metals are

compartmentalised into either a metal-sensitive fraction (MSF), which is considered the

target of attack by metals in cells, or the biologically detoxified metal (BDM) portion, which

is considered to alleviate toxicity (see review by Wang, 2013). Notably, there is evidence

that over time, metals are moved from the MSF to the BDM, and that fish surviving long-

term in chronically metal-impacted environments have partitioned the majority of their tissue

metal burdens into the BDM (Wood et al., 2012b). Overall, the observation of tissue-specific

shifts in metal burdens, as well as the retention of high metal burdens among particular

tissues, emphasises the importance of detoxification and excretion in metal-impacted brown

trout, necessitating molecular mechanisms underpinning these processes.

Our study design used only two timepoints to protect the welfare of experimental fish (Díaz

et al., 2020), particularly as we were removing wild fish from relatively small natural

populations. This strategy limited our ability to quantify depuration over both: i) additional

timepoints, which would have allowed a more detailed insight into the journey of metal

depuration; and ii) a longer depuration period, permitting a more confident determination of

"true" depuration (including our insights into constitutive gene expression). However, we

highlight the challenges of maintaining wild fish under laboratory conditions (Huntingford et

al., 2006), including acclimation to captivity (Baillon et al., 2016), and perhaps, more

importantly, uncertainty regarding whether metal-impacted trout would ever completely

depurate to reach the background tissue-metal burdens observed in the control population.

4.5 Constitutive gene expression as a potential mechanism of adaptation

to metal pollution

We observed distinct transcriptional profiles between metal-impacted and control trout after

11 days of depuration, suggesting a role for constitutive gene regulation in mitigating metal

toxicity. In particular, the gill had a higher number of DEGs (2,042), 58% of which were

over-expressed in comparison to the control trout, including those involved in metal

detoxification, oxidative stress and response to xenobiotics. These included the well-known

metal-binding gene metallothionein (*mt2*), a key protein involved in the homeostasis and

detoxification of Cd, Cu and Zn (Hamilton & Mehrle, 1986). In a previous study quantifying

transcriptional differences measured from trout sampled directly from the river, *mt2* was

over-expressed in both the gill (log2FC: 8.2) and liver (log2FC: 5.6) (Uren Webster et al.,

2013), whereas after depuration, we found that this gene was over-expressed (albeit to a

lesser extent) only in the gill (log2FC: 1.6). We also identified genes with well-known roles

in stress and xenobiotic response, including heat-shock proteins (Gupta et al., 2010). We

further identified the overexpression of both isoforms of the aryl hydrocarbon receptor

(*ahr1a, ahr1b*), which mediate the biochemical response to toxic compounds including

metals (reviewed in Kou et al., 2024) and have been implicated in toxicant adaptation in

killifish (*Fundulus heteroclitus*; Aluru et al., 2015; Whitehead et al., 2017) and Atlantic

tomcod (*Microgadus tomcod*; Wirgin et al., 2011). In our RADseq data, we also identified an

intronic SNP in an aryl hydrocarbon receptor protein. Related specifically to the activity of

the gill in response to metals, we identified the overexpression of several ABC transporters,

which are known to be involved in the uptake, transfer and elimination of xenobiotics (Kropf

et al., 2020). We observed little overlap in the DEGs identified in the gill and liver, pointing

to the distinctive roles of these tissues in metal homeostasis. However, the most highly

over-expressed gene in both tissues was *DENN domain-containing protein 5B-like*. The

DENN domain is an evolutionarily conserved module found in all eukaryotes (Zhang et al.,

2012). DENN domain-containing proteins directly interact with Rab GTPases (Marat et al.,

2011), acting as molecular switches in the regulation of multiple intracellular trafficking

processes (Homma et al., 2021), that respond to stress (Ganotra et al., 2023) and interact

with metallothionein (Knipp et al., 2005; Wolff et al., 2008).

Overall, despite the high number of identified DEGs, we identified a relatively low number of

significantly enriched or underrepresented GO terms. This could indicate low concordance

in the function of the genes which govern the recovery process from metal exposure.

Although we sampled individuals of the same age, and of equal sex ratios (three males and

three females per group), this result could also indicate the stochasticity in using wild fish in

our experiment, and moreover, that individual wild trout might not experience metal toxicity

in the same way, nor bear the same metal tolerance phenotype. Finally, variability in the

process of acclimation to laboratory conditions could also confound the identification of

metal detoxification pathways. Related to this, although we identified some GO terms

related specifically to metal homeostasis in the gill, we identified fewer than anticipated, and

the liver showed very little signal related to metal homeostasis. A recent study in

mosquitofish (*Gambusia affinis*) inhabiting metal-polluted environments, also found that

many DEGs were not associated with known metal response pathways, highlighting the

multifariousness of metal response pathways and indicating that undocumented pathways

could be playing a role in metal tolerance (Coffin et al., 2022). Generally, these results

highlight the complexity of understanding metal tolerance, and calls attention to our lack of

knowledge surrounding adaptation and acclimation to metals in wild vertebrates.

In the gills, we identified a number of pathways involved in ion transport. An important
mechanism of metal toxicity is the disruption of ion transport pathways in the gill, which
eventually causes disruptions of ionic balance in the body (Evans, 1987; Wood, 1992). In
particular, Cd and Zn are known to share uptake routes in the gill (Ca transport pathways),
where acute toxicity occurs as a result of the disruption of Ca homeostasis (Hogstrand et
al., 1996; Niyogi & Wood, 2004). On the other hand, Cu uptake in the gill is known to occur
mainly via Na transport pathways, and acute toxicity manifests via the disruption of Na
homeostasis (Grosell & Wood, 2002). We identified Ca ion transmembrane transport,
voltage-gated calcium channel activity, and Na ion transport as significantly enriched
processes in the gill. Additionally, metals can cause the inhibition of the Na+/K+-ATPase
pump (Glover et al., 2016; Grosell & Wood, 2002; Haverinen & Vornanen, 2023), and Cu
can increase Cl loss (Mazon et al., 2002). Both potassium and chloride ion transporters
were enriched, as well as voltage-gated potassium channels. We also identified an intronic
variant in a potassium channel regulator (*kcnab1b*) in the RADseq data. Together, this
suggests that compensatory increases in ion transport activity may underpin responses to
limit the toxic effects of metals in the gill of metal-impacted trout.

Our results suggest that constitutive gene expression is likely an important adaptive
mechanism permitting metal tolerance. Plasticity theory predicts that when populations
experience consistent inducing conditions over multiple generations, a plastic response can
eventually become constitutively expressed, even in the absence of the original cues
inducing the change (*i.e.* genetic assimilation; Pigliucci et al., 2006; Waddington, 1961). It
has also been suggested that constitutive expression can provide higher fitness than short-
term responsive expression (Evans, 2015; Geisel, 2011; Snell-Rood et al., 2010). We note
that we are limited in fully deciphering what proportion of the DEGs are constitutively
expressed solely due to metal handling, as the DEG signatures could also be indicative of
response to other environmental factors. However, the clear separation of the gene
expression profiles, the identification of relevant genes and pathways in the gill,
accompanied by the observed differences in tissue-metal concentrations, and their
depuration, support an important role of constitutive gene expression in promoting metal
tolerance.

829   5 Considerations and conclusions

We provide new insights into the evolution of metal tolerance in brown trout originating from
historically metal-polluted rivers. The observation of elevated tissue-metal burdens confirms
that metal pollution affects the trout inhabiting these rivers and supports the hypothesis that

adaptive strategies to cope with the insult of elevated metals within internal tissues have
evolved. Across our genomic and transcriptomic analyses, we identified signatures of
genetic differentiation in metal-impacted rivers and a number of expressed or positively
selected genes, which are functionally linked to known metal response molecular pathways.
Specifically, several RADseq outlier loci were identified as genes with known roles in the
response to oxidative stress and we also identified transcriptomic pathways associated with
this process, including the regulation of small GTPases (Ferro et al., 2012), heat shock
proteins (Kalmar & Greensmith, 2009), AhR pathway activation (Kou et al., 2024), and
cytochrome-c oxidase activity (Srinivasan & Avadhani, 2012).

The high number of differentially expressed genes in the gill, accompanied by the strong
patterns of metal accumulation and depuration, highlights this organ as being of key
functional importance for metal tolerance in this population with other organs playing also a
role in the dynamics of metal homeostasis. Overall, our data contributes to an in-depth
understanding of the molecular pathways allowing brown trout to live in highly contaminated
environments and the evolutionary processes driving these adaptations.

Data Availability Statement

De-multiplexed RADseq data for all samples are available in the European Nucleotide
Archive (ENA) under the project accession: PRJEB78481. Note that data for the following
samples (cam01, cam02, fal01, gan01, gan02, hay06 - hay13) were published in Paris et al.
2017 (doi.org/10.1111/2041-210X.12775) and are also available at the NCBI under
BioProject PRJNA379215. RNAseq data is available from ArrayExpress accession E-MTAB-
10209.

Author contributions

JRP, RWW, NRB, EMS and JRS conceived the study, designed, and performed research.
JRP, RAK, SS, AL, VB, JFO, PBH, DR, LVL, MAU, RvA and JMC analysed data. KAM and
AF contributed analytical tools. JRP, EMS and JRS wrote the paper with contributions from
co-authors. All authors read the manuscript and provided comments.

Acknowledgements and Funding

Environment Agency provided assistance in the collection of water chemistry features and
selection of appropriate sampling sites. Simon Toms (UK Environment Agency) granted
permission to collect wild brown trout for the physiology experiment. Aquatic Research
Centre (ARC) staff at the University of Exeter provided assistance with fish husbandry and
experimental design, in particular Gregory Paull and Charles Tyler who oversaw the ethical
aspects of the laboratory experiment and who provided guidelines on Home Office
Regulations. Jennifer Fitzgerald, Rebecca Millard and Hannah Littler assisted with
experimental sampling. RAD-seq and RNA-seq sequencing was performed at the University
of Exeter Sequencing Service, which utilised equipment funded by the Wellcome Trust
Institutional Strategic Support Fund (WT097835MF), Wellcome Trust Multi-user Equipment
Award (WT101650MA) and BBSRC LOLA award (BB/K003240/1). Research was co-funded
by the Environment Agency, the Westcountry Rivers Trust, and the University of Exeter.

References

Abdelnour, S. A., Naiel, M. A. E., Said, M. B., Alnajeebi, A. M., Nasr, F. A., Al-Doaiss, A.
A., Mahasneh, Z. M. H., & Noreldin, A. E. (2024). Environmental epigenetics: Exploring
phenotypic plasticity and transgenerational adaptation in fish. *Environmental Research*,
252(Pt 1), 118799.
Ahrens, C. W., Rymer, P. D., Stow, A., Bragg, J., Dillon, S., Umbers, K. D. L., &
Dudaniec, R. Y. (2018). The search for loci under selection: trends, biases and
progress. *Molecular Ecology*, *27*(6), 1342–1356.
Aluru, N., Karchner, S. I., Franks, D. G., Nacci, D., Champlin, D., & Hahn, M. E. (2015).
Targeted mutagenesis of aryl hydrocarbon receptor 2a and 2b genes in Atlantic killifish
(*Fundulus heteroclitus*). *Aquatic Toxicology*, *158*, 192–201.
Andres, S., Ribeyre, F., Tourencq, J.-N., & Boudou, A. (2000). Interspecific comparison of
cadmium and zinc contamination in the organs of four fish species along a polymetallic
pollution gradient (Lot River, France). *The Science of the Total Environment*, *248*(1),
11–25.
Badyaev, A. V. (2005). Stress-induced variation in evolution: from behavioural plasticity to
genetic assimilation. *Proceedings. Biological Sciences / The Royal Society*, *272*(1566),
877–886.
Baillon, L., Pierron, F., Pannetier, P., Normandeau, E., Couture, P., Labadie, P., Budzinski,
H., Lambert, P., Bernatchez, L., & Baudrimont, M. (2016). Gene transcription profiling in
wild and laboratory-exposed eels: Effect of captivity and in situ chronic exposure to
pollution. *Science of the Total Environment*, *571*, 92–102.
Bekkevold, D., Höjesjö, J., Nielsen, E. E., Aldvén, D., Als, T. D., Sodeland, M., Kent, M. P.,

Lien, S., & Hansen, M. M. (2020). Northern European *Salmo trutta* (L.) populations are
genetically divergent across geographical regions and environmental gradients.
Evolutionary Applications, *13*(2), 400–416.

Bélanger-Deschênes, S., Couture, P., Campbell, P. G. C., & Bernatchez, L. (2013).
Evolutionary change driven by metal exposure as revealed by coding SNP genome
scan in wild yellow perch (*Perca flavescens*). *Ecotoxicology*, *22*(5), 938–957.

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics*, *30*(15), 2114–2120.

Bourret, V., Couture, P., Campbell, P. G. C., & Bernatchez, L. (2008). Evolutionary
ecotoxicology of wild yellow perch (*Perca flavescens*) populations chronically exposed
to a polymetallic gradient. *Aquatic Toxicology*, *86*(1), 76–90.

Bourret, V., Dionne, M., & Bernatchez, L. (2014). Detecting genotypic changes associated
with selective mortality at sea in Atlantic salmon: polygenic multilocus analysis
surpasses genome scan. *Molecular Ecology*, *23*(18), 4444–4457.

Bradley, R. W., & Morris, J. R. (1986). Heavy metals in fish from a series of metal-
contaminated lakes near Sudbury, Ontario. *Water, Air, and Soil Pollution*, *27*(3), 341–
354.

Brown, B. E. (1977). Effects of mine drainage on the River Hayle, Cornwall a) factors
affecting concentrations of copper, zinc and iron in water, sediments and dominant
invertebrate fauna. *Hydrobiologia*, *52*(2), 221–233.

Bryan, G. W., Gibbs, P. E., Hummerstone, L. G., & Burt, G. R. (1987). Copper, zinc, and
organotin as long-term factors governing the distribution of organisms in the Fal estuary
in Southwest England. *Estuaries*, *10*(3), 208–219.

Burger, J., Fossi, C., McClellan-Green, P., & Orlando, E. F. (2007). Methodologies,
bioindicators, and biomarkers for assessing gender-related differences in wildlife
exposed to environmental chemicals. *Environmental Research*, *104*(1), 135–152.

Burt, R., Burnley, R., Gill, M., & Neill, A. (2014). *Mining in Cornwall and Devon: Mines and
Men*. Exeter: University of Exeter Press.

Campbell, P. G. C., Kraemer, L. D., Giguère, A., Hare, L., & Hontela, A. (2008). Subcellular
distribution of cadmium and nickel in chronically exposed wild fish: inferences regarding
metal detoxification strategies and implications for setting water quality guidelines for
dissolved metals. *Human and Ecological Risk Assessment: An International Journal*,
14(2), 290–316.

Capblancq, T., Luu, K., Blum, M. G. B., & Bazin, E. (2018). Evaluation of redundancy
analysis to identify signatures of local adaptation. *Molecular Ecology Resources*, *18*(6),
1223–1233.

Catchen, J. M., Hohenlohe, P. A., Bernatchez, L., Funk, W. C., Andrews, K. R., & Allendorf,

F. W. (2017). Unbroken: RADseq remains a powerful tool for understanding the
genetics of adaptation in natural populations. *Molecular Ecology Resources*, *17*(3), 362–
365.
Chandrapalan, T., & Kwong, R. W. M. (2021). Functional significance and physiological
regulation of essential trace metals in fish. *The Journal of Experimental Biology*,
224(24), jeb238790.
Chi, L., Lai, Y., Tu, P., Liu, C.-W., Xue, J., Ru, H., & Lu, K. (2019). Lipid and cholesterol
homeostasis after arsenic exposure and antibiotic treatment in mice: Potential role of
the Microbiota. *Environmental Health Perspectives*, *127*(9), 97002.
Chowdhury, M. J., Bucking, C., & Wood, C. M. (2008). Pre-exposure to waterborne nickel
downregulates gastrointestinal nickel uptake in rainbow trout: indirect evidence for
nickel essentiality. *Environmental Science & Technology*, *42*(4), 1359–1364.
Coffin, J. L., Kelley, J. L., Jeyasingh, P. D., & Tobler, M. (2022). Impacts of heavy metal
pollution on the ionomes and transcriptomes of Western mosquitofish (*Gambusia
affinis*). *Molecular Ecology*, *31*(5), 1527–1542.
Couture, P., & Pyle, G. (2008). Live fast and die young: Metal effects on condition and
physiology of wild yellow perch from along two metal contamination gradients. *Human
and Ecological Risk Assessment: An International Journal*, *14*(1), 73–96.
Danecek, P., Auton, A., Abecasis, G., Albers, C. A., Banks, E., DePristo, M. A., Handsaker,
R. E., Lunter, G., Marth, G. T., Sherry, S. T., McVean, G., Durbin, R., & 1000 Genomes
Project Analysis Group. (2011). The variant call format and VCFtools. *Bioinformatics*,
27(15), 2156–2158.
Defo, M. A., Bernatchez, L., Campbell, P. G. C., & Couture, P. (2015). Transcriptional and
biochemical markers in transplanted *Perca flavescens* to characterize cadmium- and
copper-induced oxidative stress in the field. *Aquatic Toxicology*, *162*, 39–53.
De Smet, H., De Wachter, B., Lobinski, R., & Blust, R. (2001). Dynamics of (Cd,Zn)-
metallothioneins in gills, liver and kidney of common carp *Cyprinus carpio* during
cadmium exposure. *Aquatic Toxicology, 52*(3-4), 269–281.
Díaz, L., Zambrano, E., Flores, M. E., Contreras, M., Crispín, J. C., Alemán, G., Bravo, C.,
Armenta, A., Valdés, V. J., Tovar, A., Gamba, G., Barrios-Payán, J., & Bobadilla, N. A.
(2020). Ethical considerations in animal research: The principle of 3R's. *Revista de
Investigacion Clinica; Organo Del Hospital de Enfermedades de La Nutricion*, *73*(4),
199–209.
Dobin, A., Davis, C. A., Schlesinger, F., Drenkow, J., Zaleski, C., Jha, S., Batut, P.,
Chaisson, M., & Gingeras, T. R. (2012). STAR: ultrafast universal RNA-seq aligner.
Bioinformatics, *29*(1), 15–21.
Dodmane, P. R., Arnold, L. L., Kakiuchi-Kiyota, S., Qiu, F., Liu, X., Rennard, S. I., & Cohen,

S. M. (2013). Cytotoxicity and gene expression changes induced by inorganic and
organic trivalent arsenicals in human cells. *Toxicology*, *312*, 18–29.

Durrant, C. J., Stevens, J. R., Hogstrand, C., & Bury, N. R. (2011). The effect of metal
pollution on the population genetic structure of brown trout (*Salmo trutta* L.) residing in
the River Hayle, Cornwall, UK. *Environmental Pollution*, *159*(12), 3595–3603.

Eastwood, S., & Couture, P. (2002). Seasonal variations in condition and liver metal
concentrations of yellow perch (*Perca flavescens*) from a metal-contaminated
environment. *Aquatic Toxicology, 58*(1-2), 43–56.

Etter, P. D., Bassham, S., Hohenlohe, P. A., Johnson, E. A., & Cresko, W. A. (2011). SNP
discovery and genotyping for evolutionary genetics using RAD sequencing. *Methods in
Molecular Biology, 772*, 157–178.

Evans, D. H. (1987). The fish gill: site of action and model for toxic effects of environmental
pollutants. *Environmental Health Perspectives*, *71*, 47–58.

Evans, T. G. (2015). Considerations for the use of transcriptomics in identifying the "genes
that matter" for environmental adaptation. *The Journal of Experimental Biology*, *218*(Pt
12), 1925–1935.

Ferro, E., Goitre, L., Retta, S. F., & Trabalzini, L. (2012). The interplay between ROS and
Ras GTPases: physiological and pathological Implications. *Journal of Signal
Transduction*, *2012*, 365769.

Forester, B. R., Lasky, J. R., Wagner, H. H., & Urban, D. L. (2018). Comparing methods for
detecting multilocus adaptation with multivariate genotype-environment associations.
Molecular Ecology, *27*(9), 2215–2233.

Fraser, D. J., Weir, L. K., Bernatchez, L., Hansen, M. M., & Taylor, E. B. (2011). Extent and
scale of local adaptation in salmonid fishes: review and meta-analysis. *Heredity*, *106*(3),
404–420.

Frichot, E., Schoville, S. D., Bouchard, G., & François, O. (2013). Testing for associations
between loci and environmental gradients using latent factor mixed models. *Molecular
Biology and Evolution*, *30*(7), 1687–1699.

Ganotra, J., Sharma, B., Biswal, B., Bhardwaj, D., & Tuteja, N. (2023). Emerging role of
small GTPases and their interactome in plants to combat abiotic and biotic stress.
Protoplasma, *260*(4), 1007–1029.

Garai, P., Banerjee, P., Mondal, P., Ch, N., & Saha, R. (2021). Effect of heavy metals on
fishes: Toxicity and bioaccumulation. *Journal of Clinical Toxicology*, *11*(S18), 1–10.

Geisel, N. (2011). Constitutive versus responsive gene expression strategies for growth in
changing environments. *PloS One*, *6*(11), e27033.

Ghalambor, C. K., McKAY, J. K., Carroll, S. P., & Reznick, D. N. (2007). Adaptive versus
non-adaptive phenotypic plasticity and the potential for contemporary adaptation in new

environments. *Functional Ecology*, *21*(3), 394–407.

Glover, C. N., Urbina, M. A., Harley, R. A., & Lee, J. A. (2016). Salinity-dependent

mechanisms of copper toxicity in the galaxiid fish, *Galaxias maculatus*. *Aquatic*

Toxicology, *174*, 199–207.

Goudet, J. (2005). Hierfstat, a package for R to compute and test hierarchical F-statistics.

Molecular Ecology Notes, *5*(1), 184–186.

Grosell, M., & Wood, C. M. (2002). Copper uptake across rainbow trout gills: mechanisms of

apical entry. *The Journal of Experimental Biology*, *205*(Pt 8), 1179–1188.

Gupta, S. C., Sharma, A., Mishra, M., Mishra, R. K., & Chowdhuri, D. K. (2010). Heat shock

proteins in toxicology: How close and how far? *Life Sciences*, *86*(11-12), 377–384.

Hamilton, P. B., Rolshausen, G., Uren Webster, T. M., & Tyler, C. R. (2017). Adaptive

capabilities and fitness consequences associated with pollution exposure in fish.

Philosophical Transactions of the Royal Society of London. Series B, Biological

Sciences, *372*(1712), 20160042.

Hamilton, S. J., & Mehrle, P. M. (1986). Metallothionein in fish: Review of its importance in

assessing stress from metal contaminants. *Transactions of the American Fisheries*

Society, *115*(4), 596–609.

Handy, R. D. (1992). The assessment of episodic metal pollution. I. Uses and limitations of

tissue contaminant analysis in rainbow trout (*Oncorhynchus mykiss*) after short

waterborne exposure to cadmium or copper. *Archives of Environmental Contamination*

and Toxicology, *22*(1), 74–81.

Han, H. J., Tokino, T., & Nakamura, Y. (1998). CSR, a scavenger receptor-like protein with a

protective role against cellular damage caused by UV irradiation and oxidative stress.

Human Molecular Genetics, *7*(6), 1039–1046.

Hansen, T., Fjelldal, P. G., Lien, S., Smith, M., Corton, C., Oliver, K., Skelton, J., Betteridge,

E., Doulcan, J., Fedrigo, O., Mountcastle, J., Jarvis, E., McCarthy, S. A., Chow, W.,

Howe, K., Torrance, J., Wood, J., Sims, Y., Haggerty, L., … Blaxter, M. (2021). The

genome sequence of the brown trout, *Salmo trutta* Linnaeus 1758. *Wellcome Open*

Research, *6*, 108.

Hart, M. B., Molina, G. S., & Smart, C. W. (2020). Estuarine foraminifera from South West

England: impact of metal pollution in a mining heritage area. *Journal of Sedimentary*

Environments, *5*(1), 1–16.

Hartwig, A., Asmuss, M., Ehleben, I., Herzer, U., Kostelac, D., Pelzer, A., Schwerdtle, T., &

Bürkle, A. (2002). Interference by toxic metal ions with DNA repair processes and cell

cycle control: molecular mechanisms. *Environmental Health Perspectives*, *110*(Suppl 5),

797–799.

Haverinen, J., & Vornanen, M. (2023). Dual effect of metals on branchial and renal Na,K-

ATPase activity in thermally acclimated crucian carp (*Carassius carassius*) and rainbow
trout (*Oncorhynchus mykiss*). *Aquatic Toxicology*, *254*, 106374.

Hecht, S. A., Baldwin, D. H., Mebane, C. A., Hawkes, T., Gross, S. J., & Scholz, N. L.
(2007). An overview of sensory effects on juvenile salmonids exposed to dissolved
copper: Applying a benchmark concentration approach to evaluate sublethal
neurobehavioral toxicity. NOAA Technical Memorandum NMFS-NWFSC-83,
https://repository.library.noaa.gov/view/noaa/3524

Hendry, A. P. (2016). Key questions on the role of phenotypic plasticity in eco-evolutionary
dynamics. *The Journal of Heredity*, *107*(1), 25–41.

Hendry, A. P., Gotanda, K. M., & Svensson, E. I. (2017). Human influences on evolution,
and the ecological and societal consequences. *Philosophical Transactions of the Royal
Society of London. Series B, Biological Sciences*, *372*(1712), 20160028.

He, Z., Mei, L., Connell, M., & Maxwell, C. A. (2020). Hyaluronan mediated motility receptor
(HMMR) encodes an evolutionarily conserved homeostasis, mitosis, and meiosis
regulator rather than a hyaluronan receptor. *Cells*, *9*(4), 819.

Hoban, S., Kelley, J. L., Lotterhos, K. E., Antolin, M. F., Bradburd, G., Lowry, D. B., Poss, M.
L., Reed, L. K., Storfer, A., & Whitlock, M. C. (2016). Finding the genomic basis of local
adaptation: Pitfalls, practical solutions, and future directions. *The American Naturalist*,
188(4), 379–397.

Hogstrand, C., Verbost, P. M., Bonga, S. E., & Wood, C. M. (1996). Mechanisms of zinc
uptake in gills of freshwater rainbow trout: interplay with calcium transport. *The
American Journal of Physiology*, *270*(5 Pt 2), R1141–R1147.

Homma, Y., Hiragi, S., & Fukuda, M. (2021). Rab family of small GTPases: an updated view
on their regulation and functions. *The FEBS Journal*, *288*(1), 36–55.

Horn, J. L. (1965). A rationale and test for the number of factors in factor analysis.
Psychometrika, *30*, 179–185.

Huntingford, F. A., Adams, C., Braithwaite, V. A., Kadri, S., Pottinger, T. G., Sandøe, P., &
Turnbull, J. F. (2006). Current issues in fish welfare. *Journal of Fish Biology*, *68*(2), 332–
372.

Ichijo, H., Nishida, E., Irie, K., ten Dijke, P., Saitoh, M., Moriguchi, T., Takagi, M.,
Matsumoto, K., Miyazono, K., & Gotoh, Y. (1997). Induction of apoptosis by ASK1, a
mammalian MAPKKK that activates SAPK/JNK and p38 signaling pathways. *Science*,
275(5296), 90–94.

Jia, Y., Wang, L., Qu, Z., Wang, C., & Yang, Z. (2017). Effects on heavy metal accumulation
in freshwater fishes: species, tissues, and sizes. *Environmental Science and Pollution
Research International*, *24*(10), 9379–9386.

Kalmar, B., & Greensmith, L. (2009). Induction of heat shock proteins for protection against

oxidative stress. *Advanced Drug Delivery Reviews*, *61*(4), 310–318.

King, R. A., Ellis, C. D., Bekkevold, D., Ensing, D., Lecointre, T., Osmond, D. R., Piper,

A., Roberts, D. E., Launey, S., & Stevens, J. R. (2024). Leveraging the genetic diversity

of trout in the rivers of the British Isles and northern France to understand the

movements of sea trout (*Salmo trutta* L.) around the English Channel. *Evolutionary*

Applications, *17*(7), e13759.

King, R. A., & Stevens, J. R. (2020). An improved genetic sex test for Atlantic salmon (*Salmo*

salar L.). *Conservation Genetics Resources*, *12*(2), 191–193.

King, R. A., Stockley, B., & Stevens, J. R. (2020). Small coastal streams - critical reservoirs

of genetic diversity for trout (*Salmo trutta* L.) in the face of increasing anthropogenic

stressors. *Ecology and Evolution*, *10*(12), 5651–5669.

Knipp, M., Meloni, G., Roschitzki, B., & Vasák, M. (2005). Zn_7metallothionein-3 and the

synaptic vesicle cycle: interaction of metallothionein-3 with the small GTPase Rab3A.

Biochemistry, *44*(9), 3159–3165.

Kou, Z., Tran, F., & Dai, W. (2024). Heavy metals, oxidative stress, and the role of AhR

signaling. *Toxicology and Applied Pharmacology*, *482*, 116769.

Kraemer, L. D., Campbell, P. G. C., & Hare, L. (2005). A field study examining metal

elimination kinetics in juvenile yellow perch (*Perca flavescens*). *Aquatic Toxicology*,

75(2), 108–126.

Kropf, C., Fent, K., Fischer, S., Casanova, A., & Segner, H. (2020). ABC transporters in gills

of rainbow trout (*Oncorhynchus mykiss*). *The Journal of Experimental Biology*, *223*(15),

jeb221069.

Kumar, V., Parihar, R. D., Sharma, A., Bakshi, P., Singh Sidhu, G. P., Bali, A. S., Karaouzas,

I., Bhardwaj, R., Thukral, A. K., Gyasi-Agyei, Y., & Rodrigo-Comino, J. (2019). Global

evaluation of heavy metal content in surface water bodies: A meta-analysis using heavy

metal pollution indices and multivariate statistical analyses. *Chemosphere*, *236*, 124364.

Laporte, M., Pavey, S. A., Rougeux, C., Pierron, F., Lauzent, M., Budzinski, H., Labadie, P.,

Geneste, E., Couture, P., Baudrimont, M., & Bernatchez, L. (2016). RAD sequencing

reveals within-generation polygenic selection in response to anthropogenic organic and

metal contamination in North Atlantic Eels. *Molecular Ecology*, *25*(1), 219–237.

Leonard, E. M., Banerjee, U., D'Silva, J. J., & Wood, C. M. (2014). Chronic nickel

bioaccumulation and sub-cellular fractionation in two freshwater teleosts, the round

goby and the rainbow trout, exposed simultaneously to waterborne and dietborne nickel.

Aquatic Toxicology, *154*, 141–153.

Liao, Y., Smyth, G. K., & Shi, W. (2014). featureCounts: an efficient general purpose

program for assigning sequence reads to genomic features. *Bioinformatics*, *30*(7), 923–

930.

López-Maury, L., Marguerat, S., & Bähler, J. (2008). Tuning gene expression to changing
environments: from rapid responses to evolutionary adaptation. *Nature Reviews.*
Genetics, *9*(8), 583–593.
Love, M. I., Huber, W., & Anders, S. (2014). Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. *Genome Biology*, *15*(12), 550.
Lowry, D. B., Hoban, S., Kelley, J. L., Lotterhos, K. E., Reed, L. K., Antolin, M. F., & Storfer,
A. (2017). Breaking RAD: an evaluation of the utility of restriction site-associated DNA
sequencing for genome scans of adaptation. *Molecular Ecology Resources*, *17*(2), 142–
152.
Luu, K., Bazin, E., & Blum, M. G. B. (2017). pcadapt: an R package to perform genome
scans for selection based on principal component analysis. *Molecular Ecology*
Resources, *17*(1), 67–77.
Maes, G. E., Raeymaekers, J. A. M., Pampoulie, C., Seynaeve, A., Goemans, G., Belpaire,
C., & Volckaert, F. A. M. (2005). The catadromous European eel *Anguilla anguilla* (L.)
as a model for freshwater evolutionary ecotoxicology: Relationship between heavy
metal bioaccumulation, condition and genetic variability. *Aquatic Toxicology*, *73*(1), 99–
114.
Marandel, F., Charrier, G., Lamy, J.-B., Le Cam, S., Lorance, P., & Trenkel, V. M. (2020).
Estimating effective population size using RADseq: Effects of SNP selection and sample
size. *Ecology and Evolution*, *10*(4), 1929–1937.
Marat, A. L., Dokainish, H., & McPherson, P. S. (2011). DENN domain proteins: regulators of
Rab GTPases. *The Journal of Biological Chemistry*, *286*(16), 13791–13800.
Mazon, A. F., Cerqueira, C. C. C., & Fernandes, M. N. (2002). Gill cellular changes induced
by copper exposure in the South American tropical freshwater fish *Prochilodus scrofa*.
Environmental Research, *88*(1), 52–63.
McLaren, W., Gil, L., Hunt, S. E., Riat, H. S., Ritchie, G. R. S., Thormann, A., Flicek, P., &
Cunningham, F. (2016). The Ensembl Variant Effect Predictor. *Genome Biology*, *17*(1),
122.
Mendes, F. K., & Hahn, M. W. (2016). Gene tree discordance causes apparent substitution
rate variation. *Systematic Biology*, *65*(4), 711–721.
Mérot, C., Oomen, R. A., Tigano, A., & Wellenreuther, M. (2020). A roadmap for
understanding the evolutionary significance of structural genomic variation. *Trends in*
Ecology & Evolution, *35*(7), 561–572.
Minghetti, M., Schnell, S., Chadwick, M. A., Hogstrand, C., & Bury, N. R. (2014). A primary
FIsh Gill Cell System (FIGCS) for environmental monitoring of river waters. *Aquatic*
Toxicology, *154*, 184–192.
Moriyama, Y., & Koshiba-Takeuchi, K. (2018). Significance of whole-genome duplications on

the emergence of evolutionary novelties. *Briefings in Functional Genomics*, *17*(5), 329–
338.

Munro, H. N., & Linder, M. C. (1978). Ferritin: structure, biosynthesis, and role in iron
metabolism. *Physiological Reviews*, *58*(2), 317–396.

Niyogi, S., & Wood, C. M. (2004). Kinetic analyses of waterborne Ca and Cd transport and
their interactions in the gills of rainbow trout (*Oncorhynchus mykiss*) and yellow perch
(*Perca flavescens*), two species differing greatly in acute waterborne Cd sensitivity.
Journal of Comparative Physiology. B, Biochemical, Systemic, and Environmental
Physiology, *174*(3), 243–253.

O'Doherty, C., Keenan, J., O'Neill, F., Clynes, M., Sinkunaite, I., Horgan, K., Murphy, R., &
O'Sullivan, F. (2020). Gene expression profiling of copper-resistant Caco-2 clones.
Metallomics: Integrated Biometal Science, *12*(10), 1521–1529.

Ortiz, E. M. (2019). vcf2phylip v2. 0: convert a VCF matrix into several matrix formats for
phylogenetic analysis. *URL https://doi org/105281/zenodo*, *2540861*.

Osmond, D. R., King, R. A., Russo, I. M., Bruford, M., & Stevens, J. R. (2024). Living in
a post□industrial landscape: repeated patterns of genetic divergence in brown trout
(*Salmo trutta* L.) across the British Isles. *Diversity & Distributions*, *30*(7), e13854.

Paradis, E., & Schliep, K. (2019). ape 5.0: an environment for modern phylogenetics and
evolutionary analyses in R. *Bioinformatics*, *35*(3), 526–528.

Paris, J. R., King, R. A., & Stevens, J. R. (2015). Human mining activity across the ages
determines the genetic structure of modern brown trout (*Salmo trutta* L.) populations.
Evolutionary Applications, *8*(6), 573–585.

Paris, J. R., Stevens, J. R., & Catchen, J. M. (2017). Lost in parameter space: a road map
for stacks. *Methods in Ecology and Evolution / British Ecological Society*, *8*(10), 1360–
1373.

Pierron, F., Daffe, G., Daramy, F., Heroin, D., Barré, A., Bouchez, O., Clérendeau, C.,
Romero-Ramirez, A., & Nikolski, M. (2023). Transgenerational endocrine disruptor
effects of cadmium in zebrafish and contribution of standing epigenetic variation to
adaptation. *Journal of Hazardous Materials*, *455*(131579), 131579.

Pierron, F., Bourret, V., St-Cyr, J., Campbell, P. G. C., Bernatchez, L., & Couture, P. (2009).
Transcriptional responses to environmental metal exposure in wild yellow perch (*Perca*
flavescens) collected in lakes with differing environmental metal concentrations (Cd, Cu,
Ni). *Ecotoxicology*, *18*(5), 620–631.

Pigliucci, M., Murren, C. J., & Schlichting, C. D. (2006). Phenotypic plasticity and evolution
by genetic assimilation. *The Journal of Experimental Biology*, *209*(Pt 12), 2362–2367.

Playle, R. C. (1998). Modelling metal interactions at fish gills. *Science of the Total*
Environment, *219*(2), 147–163.

Primmer, C. R. (2011). Genetics of local adaptation in salmonid fishes. *Heredity*, *106*(3),
401–403.
Purcell, S., Neale, B., Todd-Brown, K., Thomas, L., Ferreira, M. A. R., Bender, D., Maller, J.,
Sklar, P., de Bakker, P. I. W., Daly, M. J., & Sham, P. C. (2007). PLINK: a tool set for
whole-genome association and population-based linkage analyses. *American Journal of*
Human Genetics, *81*(3), 559–575.
Rainbow, P. S. (2020). Mining-contaminated estuaries of Cornwall – field research
laboratories for trace metal ecotoxicology. *Journal of the Marine Biological Association*
of the United Kingdom. Marine Biological Association of the United Kingdom, *100*(2),
195–210.
R Core Team. (2021). *R: A Language and Environment for Statistical Computing*. R
Foundation for Statistical Computing. https://www.R-project.org/
Rivera, H. E., Aichelman, H. E., Fifer, J. E., Kriefall, N. G., Wuitchik, D. M., Wuitchik, S. J. S.,
& Davies, S. W. (2021). A framework for understanding gene expression plasticity and
its influence on stress tolerance. *Molecular Ecology*, *30*(6), 1381–1397.
Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2: Analytical methods
for paired-end sequencing improve RADseq-based population genomics. *Molecular*
Ecology, *28*(21), 4737–4754.
Saibu, Y., Kumar, S., Jamwal, A., Peak, D., & Niyogi, S. (2018). A FTIRM study of the
interactive effects of metals (zinc, copper and cadmium) in binary mixtures on the
biochemical constituents of the gills in rainbow trout (*Oncorhynchus mykiss*).
Comparative Biochemistry and Physiology. Toxicology & Pharmacology: CBP, *211*, 48–
56.
Santiago, E., Caballero, A., Köpke, C., & Novo, I. (2024). Estimation of the contemporary
effective population size from SNP data while accounting for mating structure. *Molecular*
Ecology Resources, *24*(1), e13890.
Schwarzenbach, R. P., Egli, T., Hofstetter, T. B., von Gunten, U., & Wehrli, B. (2010). Global
water pollution and human health. *Annual Review of Environment and Resources*,
35(1), 109–136.
Sergeant, C. J., Sexton, E. K., Moore, J. W., Westwood, A. R., Nagorski, S. A., Ebersole, J.
L., Chambers, D. M., O'Neal, S. L., Malison, R. L., Hauer, F. R., Whited, D. C., Weitz, J.,
Caldwell, J., Capito, M., Connor, M., Frissell, C. A., Knox, G., Lowery, E. D., Macnair,
R., … Skuce, N. (2022). Risks of mining to salmonid-bearing watersheds. *Science*
Advances, *8*(26), eabn0929.
Sevcikova, M., Modra, H., Slaninova, A., Svobodova, Z., & Others. (2011). Metals as a
cause of oxidative stress in fish: a review. *Veterinary Medicine*, *56*(11), 537–546.
Shahjahan, M., Taslima, K., Rahman, M. S., Al-Emran, M., Alam, S. I., & Faggio, C. (2022).

Effects of heavy metals on fish physiology - a review. *Chemosphere*, *300*(134519),

134519.

Shekh, K., Saeed, H., Kodzhahinchev, V., Brinkmann, M., Hecker, M., & Niyogi, S. (2021).

Differences in the subcellular distribution of cadmium and copper in the gills and liver of

white sturgeon (*Acipenser transmontanus*) and rainbow trout (*Oncorhynchus mykiss*).

Chemosphere, *265*, 129142.

Shi, Q., Xiong, B., Zhong, J., Wang, H., Ma, D., & Miao, C. (2017). MFHAS1 suppresses

TLR4 signaling pathway via induction of PP2A C subunit cytoplasm translocation and

inhibition of c-Jun dephosphorylation at Thr239. *Molecular Immunology*, *88*, 79–88.

Smedley, D., Haider, S., Ballester, B., Holland, R., London, D., Thorisson, G., & Kasprzyk, A.

(2009). BioMart - biological queries made easy. *BMC Genomics*, *10*(1), 1–12.

Smith, T. B., & Bernatchez, L. (2008). Evolutionary change in human-altered environments.

Molecular Ecology, *17*(1), 1–8.

Snell-Rood, E. C., Van Dyken, J. D., Cruickshank, T., Wade, M. J., & Moczek, A. P. (2010).

Toward a population genetic framework of developmental evolution: the costs, limits,

and consequences of phenotypic plasticity. *BioEssays: News and Reviews in Molecular,*

Cellular and Developmental Biology, *32*(1), 71–81.

Sone, H., Akanuma, H., & Fukuda, T. (2010). Oxygenomics in environmental stress. *Redox*

Report: Communications in Free Radical Research, *15*(3), 98–114.

Srikanth, K., Pereira, E., Duarte, A. C., & Ahmad, I. (2013). Glutathione and its dependent

enzymes' modulatory responses to toxic metals and metalloids in fish - a review.

Environmental Science and Pollution Research International, *20*(4), 2133–2149.

Srinivasan, S., & Avadhani, N. G. (2012). Cytochrome c oxidase dysfunction in oxidative

stress. *Free Radical Biology & Medicine*, *53*(6), 1252–1263.

Taylor, E. B. (1991). A review of local adaptation in Salmonidac, with particular reference to

Pacific and Atlantic salmon. *Aquaculture*, *98*(1), 185–207.

Theodorakis, C. W., Lee, K.-L., Adams, S. M., & Law, C. B. (2006). Evidence of altered gene

flow, mutation rate, and genetic diversity in redbreast sunfish from a pulp-mill-

contaminated river. *Environmental Science & Technology*, *40*(1), 377–386.

Timberlake, S. (2017). New ideas on the exploitation of copper, tin, gold, and lead ores in

Bronze Age Britain: The mining, smelting, and movement of metal. *Materials and*

Manufacturing Processes, *32*(7-8), 709–727.

Travis, C. C., & Hester, S. T. (1991). Global chemical pollution. *Environmental Science &*

Technology, *25*(5), 814–819.

Uno, T., Ishizuka, M., & Itakura, T. (2012). Cytochrome P450 (CYP) in fish. *Environmental*

Toxicology and Pharmacology, *34*(1), 1–13.

Uren Webster, T. M., Bury, N., van Aerle, R., & Santos, E. M. (2013). Global transcriptome

profiling reveals molecular mechanisms of metal tolerance in a chronically exposed wild

population of brown trout. *Environmental Science & Technology*, *47*(15), 8869–8877.

Valko, M., Morris, H., & Cronin, M. T. D. (2005). Metals, toxicity and oxidative stress. *Current*

Medicinal Chemistry, *12*(10), 1161–1208.

Vangronsveld, J., & Clijsters, H. (1994). Toxic effects of metals. *Plants and the Chemical*

Elements: Biochemistry, Uptake, Tolerance and Toxicity, 149–177.

van Straalen, N. M., & Timmermans, M. J. T. N. (2002). Genetic variation in toxicant-

stressed populations: an evaluation of the "Genetic Erosion" hypothesis. *Human and*

Ecological Risk Assessment: An International Journal, *8*(5), 983–1002.

Vareda, J. P., Valente, A. J. M., & Durães, L. (2019). Assessment of heavy metal pollution

from anthropogenic activities and remediation strategies: A review. *Journal of*

Environmental Management, *246*, 101–118.

Veltman, K., Huijbregts, M. A. J., Van Kolck, M., Wang, W.-X., & Hendriks, A. J. (2008).

Metal bioaccumulation in aquatic species: quantification of uptake and elimination rate

constants using physicochemical properties of metals and physiological characteristics

of species. *Environmental Science & Technology*, *42*(3), 852–858.

Waddington, C. H. (1961). Genetic assimilation. *Advances in Genetics*, *10*, 257–293.

Wadgymar, S. M., & DeMarche, M. L. (2022). Local adaptation: causal agents of selection

and adaptive trait divergence. *Annual Review of Ecology, Evolution & Systematics, 53,*

87-111.

Wang, W.-X. (2013). Prediction of metal toxicity in aquatic organisms. *Chinese Science*

Bulletin, 58, 194–202.

Wang, W.-X., & Rainbow, P. S. (2006). Subcellular partitioning and the prediction of

cadmium toxicity to aquatic organisms. *Environmental Chemistry*, *3*(6), 395–399.

Waples, R. S. (2024). Practical application of the linkage disequilibrium method for

estimating contemporary effective population size: A review. *Molecular Ecology*

Resources, *24*(1), e13879.

Whitehead, A. (2014). Evolutionary genomics of environmental pollution. *Advances in*

Experimental Medicine and Biology, *781*, 321–337.

Whitehead, A., Clark, B. W., Reid, N. M., Hahn, M. E., & Nacci, D. (2017). When evolution is

the solution to pollution: Key principles, and lessons from rapid repeated adaptation of

killifish (*Fundulus heteroclitus*) populations. *Evolutionary Applications*, *10*(8), 762–783.

Whitlock, M. C., & Lotterhos, K. E. (2015). Reliable detection of loci responsible for local

adaptation: inference of a null model through trimming the distribution of F(ST). *The*

American Naturalist, *186 Suppl 1*, S24–S36.

Williams, C. R., & Gallagher, E. P. (2013). Effects of cadmium on olfactory mediated

behaviors and molecular biomarkers in coho salmon (*Oncorhynchus kisutch*). *Aquatic*

Toxicology, *140-141*, 295–302.

Wirgin, I., Roy, N. K., Loftus, M., Chambers, R. C., Franks, D. G., & Hahn, M. E. (2011).

Mechanistic basis of resistance to PCBs in Atlantic tomcod from the Hudson River.

Science, *331*(6022), 1322–1325.

Wolff, N. A., Lee, W.-K., Abouhamed, M., & Thévenod, F. (2008). Role of ARF6 in

internalization of metal-binding proteins, metallothionein and transferrin, and cadmium-

metallothionein toxicity in kidney proximal tubule cells. *Toxicology and Applied*

Pharmacology, *230*(1), 78–85.

Wood, C. M. (1992). Flux measurements as indices of H+ and metal effects on freshwater

fish. *Aquatic Toxicology*, *22*(4), 239–263.

Wood, C. M. (2012). An Introduction to Metals in Fish Physiology and Toxicology: Basic

Principles. In Wood, C.M., Farrell, A.P., Brauner, C.J. (Ed.), *Homeostasis and*

Toxicology of Non-Essential Metals (Vol. 31B, pp. 2–40). Cambridge, Massachusetts:

Academic Press.

Wood, C. M., Farrell, A. P., & Brauner, C. J. (2012a). *Homeostasis and Toxicology of Non-*

Essential Metals (Vol. 31B). Cambridge, Massachusetts: Academic Press.

Wood, C. M., Farrell, A. P., & Brauner, C. J. (2012b). *Homeostasis and Toxicology of*

Essential Metals (Vol. 31A). Cambridge, Massachusetts: Academic Press.

Yan, J., Hedl, M., & Abraham, C. (2017). An inflammatory bowel disease-risk variant in

INAVA decreases pattern recognition receptor-induced outcomes. *The Journal of*

Clinical Investigation, *127*(6), 2192–2205.

Yu, G., Wang, L.-G., Han, Y., & He, Q.-Y. (2012). clusterProfiler: an R Package for

comparing biological themes among gene clusters. *Omics: A Journal of Integrative*

Biology, *16*(5), 284–287.

Zhang, D., Iyer, L. M., He, F., & Aravind, L. (2012). Discovery of novel DENN proteins:

Implications for the evolution of eukaryotic intracellular membrane structures and human

disease. *Frontiers in Genetics*, *3*, 283.

Zhong, J., Shi, Q.-Q., Zhu, M.-M., Shen, J., Wang, H.-H., Ma, D., & Miao, C.-H. (2015).

MFHAS1 is associated with sepsis and stimulates TLR2/NF-κB signaling pathway

following negative regulation. *PloS One*, *10*(11), e0143662.

1339