Single-nucleus transcriptomics reveal the cytological mechanism of conjugated

linoleic acids in regulating intramuscular fat deposition

Liyi Wang [1, 2, 3], Shiqi Liu [1, 2, 3], Shu Zhang [1, 2, 3], Yizhen Wang [1, 2, 3], Yanbing Zhou [1, 2,

3, *], Tizhong Shan [1, 2, 3, *]

[1] College of Animal Sciences, Zhejiang University, Hangzhou, China

[2] Key Laboratory of Molecular Animal Nutrition (Zhejiang University), Ministry of

Education, China

[3] Key Laboratory of Animal Feed and Nutrition of Zhejiang Province, Hangzhou,

China

* Correspondence:

Tizhong Shan, **ORCID**: 0000-0002-4738-414x

tzshan@zju.edu.cn, Zhejiang University,

Yuhangtang Road 866, Hangzhou, CN 310058

Yanbing Zhou, **ORCID**: 0000-0002-4196-5947

zhouyanbing@zju.edu.cn, Zhejiang University,

Yuhangtang Road 866, Hangzhou, CN 310058

Abstract

Conjugated linoleic acids (CLAs) can serve as a nutritional intervention to regulate quality, function and fat infiltration in skeletal muscles but the specific cytological mechanisms are still unknown. Here, we applied single-nucleus RNA-sequencing (snRNA-seq) to characterize the cytological mechanism of CLAs regulates fat infiltration in skeletal muscles based on pig models. We investigated the regulatory effects of CLAs on cell populations and molecular characteristics in pig muscles and found CLAs could promote the transformation of fast glycolytic myofibers into slow oxidative myofibers. We also observed three subpopulations including $SCD^+/DGAT2^+$, $FABP5^+/SIAH1^+$, and $PDE4D^+/PDE7B^+$ subclusters in adipocytes and CLAs could increase the percentage of $SCD^+/DGAT2^+$ adipocytes. RNA velocity analysis showed $FABP5^+/SIAH1^+$ and $PDE4D^+/PDE7B^+$ adipocytes could differentiate into $SCD^+/DGAT2^+$ adipocytes. We further verified the differentiated trajectory of mature adipocytes and identified $PDE4D^+/PDE7B^+$ adipocytes could differentiate into $SCD^+/DGAT2^+$ and $FABP5^+/SIAH1^+$ adipocytes by using high IMF content Laiwu pig models. The cell-cell communication analysis identified the interaction network between adipocytes and other subclusters such as fibro/adipogenic progenitors (FAPs). Pseudotemporal trajectory analysis and RNA velocity analysis also showed FAPs could differentiate into $PDE4D^+/PDE7B^+$ preadipocytes and we discovered the differentiated trajectory of preadipocytes into mature adipocytes. Besides, we found CLAs could promote FAPs differentiate into

$SCD^+/DGAT2^+$ adipocytes via inhibiting c-Jun N-terminal kinase (JNK) signalling

pathway *in vitro*. This study provides a foundation for regulating fat infiltration in

skeletal muscles by using nutritional strategies and provides potential opportunities to

serve pig as an animal model to study human fat infiltrated diseases.

Keywords: Single-nucleus RNA-sequencing; Conjugated linoleic acid; Skeletal

muscle; Fibro/adipogenic progenitors; Fat infiltration; Pig

Introduction

Meat is one of the most important sources of animal protein for humans, and its quality is associated with human health. Recently, due to the development of economic levels and the improvement of living standards, people are seeking for 'less but better' meat (Sahlin & Trewern, 2022). Intramuscular fat (IMF) deposition is a key factor positively related to meat quality traits and lipo-nutritional values of meat, such as flavor, tenderness, and juiciness(Hausman, Basu, Du, Fernyhough-Culver, & Dodson, 2014; W. Yi, Huang, Wang, & Shan, 2023). Besides, fat infiltration in skeletal muscle (also known as myosteatosis) is the pathologic fat accumulation in skeletal muscle with poor metabolic and musculoskeletal health, it always accompanied by the decline of muscle quality and function(Biltz et al., 2020; Jiang, Marriott, & Maly, 2019). Myosteatosis is now considered as a common feature of ageing and is also related to some diseases(Wang, Valencak, & Shan, 2024). However, the occurrence mechanism and cell sources of fat accumulation in skeletal muscle is very complicated. Recently, with the rapid development of multi-omics including single-cell RNA sequencing (scRNA-seq)(Tabula Muris et al., 2018), single-nucleus RNA-seq (snRNA-seq)(Petrany et al., 2020), and spatial transcriptomics (ST)(Jin et al., 2021), more and more cell types have been found to contribute to lipid deposition in skeletal muscle including myogenic cells (e.g., satellite cells (SCs)(Asakura, Komaki, & Rudnicki, 2001) and myogenic factor 5 (Myf5)[+] mesenchymal stem cells (MSCs)(Yin et al., 2013)) and non-myogenic cells (e.g., fibro/adipogenic progenitors

(FAPs)(Uezumi, Fukada, Yamamoto, Takeda, & Tsuchida, 2010), fibroblasts,

myeloid-derived cells(Z. Xu et al., 2020), pericytes(Farrington-Rock et al., 2004),

endothelial cells (ECs)(Lang et al., 2008), $PW1^+/Pax7^-$ interstitial cells

(PICs)(Mitchell et al., 2010), and side population cells (SPs)(Tamaki et al., 2002))

based on animal models. Hence, more and more researches have been focusing on

exploring the regulatory mechanism of myosteatosis at the cytological levels. Besides,

fat infiltration in skeletal muscle is regulated by many influential triggers, including

ageing, metabolic and nonmetabolic diseases, disuse and inactivity, and muscle

injury(Wang et al., 2024). Many genes and signaling pathways participate in the

formation and regulation of fat infiltration in skeletal muscle (Biferali et al., 2021;

Wosczyna et al., 2021). However, the specific mechanism of nutrients regulates fat

infiltration in muscle is still unknown.

Nutritional regulation strategy is one of the most vital strategies to regulate lipid

accumulation in skeletal muscle based on some animal models and clinic trials,

including vitamins(Gilsanz, Kremer, Mo, Wren, & Kremer, 2010; L. Zhao et al.,

2020), conjugated linoleic acid (CLAs)(van Vliet, Fappi, Reeds, & Mittendorfer,

2020), linseed(Wei et al., 2016), plant extract(You et al., 2023), and so on. Hence,

exploring the potential nutritional strategies to regulate fat accumulation in skeletal

muscle is valuable for animal production and human health. Pigs are not only an

important source of animal protein in the human diet but also serve as a valuable

model for human medical biology due to their similar physiological structure in terms

of size, metabolic characteristics, and cardiovascular system (Groenen et al., 2012;

Lunney et al., 2021). Chinese local pig species could be used as excellent animal

models to study the mechanism of lipid deposition due to they have better meat

quality and high IMF content. A high IMF content always results in better juiciness,

tenderness, and flavor of pork. There are different myogenesis potential in neonatal

skeletal muscle between Laiwu pigs and Duroc pigs(D. D. Xu et al., 2023). Our

previous study has revealed the cell heterogeneity and transcriptional dynamics of

lipid deposition in skeletal muscle of Laiwu pigs and we found high IMF content

Laiwu pigs had lower muscle fiber diameter(Wang, Zhao, et al., 2023). Heigai pig is a

model of Chinese indigenous pig breeds, which has advantages including high

farrowing rate, good pork quality, and strong disease resistance(Liyi Wang et al.,

2022). Specially, we have found CLAs can improve meat quality especially increase

IMF content both in lean type pig breeds and fat type pig breeds(Wang, Huang, Wang,

& Shan, 2021; L. Wang et al., 2022). However, although many studies have discussed

the effects of CLA on IMF deposition, there are still gaps in the cytological

mechanism of CLAs in regulating lipid deposition in skeletal muscle.

Here, we present a snRNA-seq dataset collected from *longissimus dorsi muscle*

(LDM) of Heigai pigs after feeding CLAs supplement to allow for analysing

heterogeneity of transcriptional states in muscles. We investigated the regulatory

effects of CLAs on the muscle fiber type transformation and IMF deposition. We also

identified the differentiation trajectories of three subclusters in adipocytes based on

Heigai pig models and high IMF content Laiwu pig models. Based on the

pseudotemporal trajectories analysis, we found CLAs could promote FAPs

differentiate into SCD$^+$/DGAT2$^+$ adipocytes via regulating mitogen-activated protein

kinase (MAPK) signalling pathway. This study paves a way to regulate lipid

accumulation in muscles by using nutritional strategies and provides theoretical basis

on using pig as an animal model to study human muscle-related diseases.

Results

CLAs changed cell populations and transcriptional dynamics in LDM

Our previous study discovered CLAs improved IMF content in LDM of Heigai

pigs(L. Wang et al., 2022), here, we also found CLAs significantly increased TG

content but significantly decreased TC content in LDM of pigs (Figure. 1A).

Meanwhile, immunofluorescence staining results showed more lipid droplets in the

LDM of the CLAs group (Figure. 1B). To investigate the changes of cell

heterogeneity in pig muscles after CLAs treatment at the cellular level, we performed

snRNA-seq from LDM tissue of Heigai pigs using the 10× Genomics Chromium

platform (Figure. 1C). First, after Cell Ranger analyses, the estimated number of cells,

fraction of reads in cells, mean reads per cell, median genes per cell and median UMI

counts per cell in LDM were showed in Supplementary Fig. S1a. We obtained 25507

cells from 2 individual libraries, comprising 10835 cells from CON group and 11705

cells from CLA group for the downstream analysis after the quality control of

snRNA-seq data (Figure. S1B). Based on the Seurat package, we used Uniform

Manifold Approximation and Projection (UMAP) plots to show the different

subclusters (Figure. 1D). We identified 8 different clusters in two groups of pig

muscles, including myofibers (*CAPN3*), FAPs/fibroblasts (*PDGFRA*), ECs (*CD34*),

adipocytes (*PPARG*), immune cells (*PTPRC*), muscle satellite cells (MuSCs) (*PAX7*),

myeloid derived cells (*MRC1*) and pericytes (*RGS5*) using the expression of marker

genes (Figure. 1E). Next, the percentage of these cell types showed differences in

different group (Figure. 1F). Compared with the CON group, the FAPs/fibroblasts

(3.39% *vs.* 8.31%), ECs (1.26% *vs.* 2.94%), adipocytes (1.74% *vs.* 2.37%), myeloid

derived cells (0.93% *vs.* 2.17%) and pericytes (0.45% *vs.* 0.7%) had a higher

proportion in the CLA group. However, CLA decreased the proportion of myofibers

(90.46% vs. 81.1%). The top 10 most differentially expressed genes (DEGs) were

showed in the heatmap and the bar plot showed the top 3 KEGG enrichment of DEGs

between the 8 different cell types in LDM (Figure. 1G). Besides, violin plots

displayed CLAs upregulated the expression of mature adipocyte master genes

including *ADIPOQ*, *FABP4*, *PLIN1*, and *LIPE*, adipogenic marker genes including

PPARG, *PPARA*, *CEBPA*, and *CEBPB*, and lipid metabolism-related genes including

LPL, *ELOVL4*, *ACAA2* and *HACD2* in pig muscles (Figures. S1C-E). These results

indicated the cell types in LDM of Heigai pigs had significant differences after

feeding CLAs which might induce the alterations in lipid deposition.

Characterization of myofibers after CLAs treatment through clustering analysis

To explore the changes of myofibers after CLA treatment, we next carried out a

subcluster analysis and investigated the cell heterogeneity of myofibers. UMAP plots

displayed the distribution in different subsets of myofibers (Figure. 2A). Based on our

previous study(Wang, Zhao, et al., 2023) and different gene expression in myofibers,

we also characterized 6 different cell types in myofibers, including I myofibers

(*MYH7*), IIA myofibers (*MYH2*), IIX myofibers (*MYH1*), IIB myofibers (*MYH4*),

myotendinous junctions (MTJ, *ANKRD1*), and neuromuscular junction (NMJ,

ABLIM2) (Figure. 2B). The bar plot showed the proportion of type I myofibers (17.54%

vs. 22.01%) and IIA myofibers (3.64% *vs.* 7.52%) had an increased tendency while

the proportion of IIX myofibers (10.09% *vs.* 3.87%), IIB myofibers (63.88% *vs.*

59.35%), MTJ (6.41% *vs.* 0.28%) and NMJ (5.32% *vs.* 0.09%) had a reduced

tendency in CLA group (Figure. 2C). Besides, violin plot displayed CLAs increased

the expression of myofiber type marker genes (*MYH7*, *MYH2*, *MYH1*, and *MYH4*),

myofiber type transformation-related genes (*PPARGC1A*, *STK11*, and *HDAC1*) and

oxidation-related genes (*COX5A*, *COX5B*, and *COX8A*) but decreased

glycolysis-related genes (*PFKM*, *HK2*, and *LDHC*) (Figure. 2D). Additionally, the

expression of *MYHCI* and *COX5B* in LDM of Heigai pigs was also significantly

increased after CLA treatment (Figure. 2E). The top 10 DEGs between the 6 cell

types in myofibers were showed in heatmap (Figure. 2F). Functional enrichment

analyses revealed the enrichment of metabolic pathways, oxidative phosphorylation,

and thermogenesis in I myofibers and calcium, cGMP-PKG, and MAPK signalling

pathway in IIB myofibers by using KEGG pathways (Figure. 2G). These data

discovered the significantly heterogeneity in myofibers between two groups and

CLAs could promote slow oxidative myofibers switch into fast glycolic myofibers in

pig muscles.

Clustering and RNA velocity analysis revealed subpopulations and cellular

dynamics of adipocytes after CLAs treatment

To investigate the regulatory mechanism of CLAs in IMF deposition, we next

performed a subset analysis on adipocytes. Our previous study had discovered there

are three subclusters in adipocytes nuclei of Laiwu pigs(Wang, Zhao, et al., 2023). In

this study, we also characterized 3 subpopulations including SCD$^+$/DGAT2$^+$

adipocytes, FABP5$^+$/SIAH1$^+$ adipocytes, and PDE4D$^+$/PDE7B$^+$ adipocytes according

to the most DEGs (Figures. 3A and S2A). Interestingly, we discovered CLAs

increased the amounts of SCD$^+$/DGAT2$^+$ adipocytes (Figure. 3B) and the proportion

of SCD$^+$/DGAT2$^+$ adipocytes (79.37% *vs.* 82.31%) but decreased the proportion of

PDE4D$^+$/PDE7B$^+$ adipocytes (14.81% *vs.* 11.19%) (Figure. S2B). We also found

CLAs increased the expression of SCD$^+$/DGAT2$^+$ adipocytes marker genes (*SCD* and

ARHGAP31) and FABP5$^+$/SIAH1$^+$ adipocytes marker genes (*SIAH1* and *COX1*) but

decreased the expression of PDE4D$^+$/PDE7B$^+$ adipocytes marker genes (*PDE4D*) in

adipocytes (Figure. 3C). Similarly, we also discovered *SCD* and *DGAT2* expression

was remarkably increased in LDM after feeding with CLAs (Figure. 3D). Meanwhile,

immunofluorescence results showed more SCD1$^+$ adipocytes in the LDM of the

CLAs group (Figure. 3E). The top 10 most DEGs were displayed in the heatmap and

the bar plot showed the top 3 KEGG enrichment of DEGs between the 3 subclusters

in LDM (Figure. 3F). To explore the effects of CLAs on differentiated trajectory of

adipocytes, we carried out the RNA velocity analysis of adipocytes (Figure. S2C).

RNA velocity results showed the differentiated trajectory of mature adipocytes in

muscles of Heigai pigs (Figure. 3G). Transcriptional dynamics of *PDE4D* and *CAPN3*

were showed in Figure. S2D based on RNA velocity analysis. These results indicated

in mature adipocytes, PDE4D$^+$/PDE7B$^+$ adipocytes and FABP5$^+$/SIAH1$^+$ adipocytes

could differentiate into SCD$^+$/DGAT2$^+$ adipocytes (Figure. 3H).

The verification of differentiated trajectories of adipocytes by using Laiwu pig

models

To further verify the differentiated trajectory of adipocytes, we next carried out a

pseudotemporal trajectory analysis and RNA velocity analysis of adipocytes by using

our previous high IMF content pig models based on Monocle 2 and scVelo (Wang,

Zhao, et al., 2023) (Figure. 4A). In adipocytes of Laiwu pigs, high IMF content Laiwu

pigs (HLW) group had the higher proportion of SCD$^+$ /DGAT2$^+$ adipocytes but the

lower proportion of PDE4D$^+$/PDE7B$^+$ adipocytes (Figure. 4B). Meanwhile,

immunofluorescence staining results also showed the more SCD1$^+$ adipocytes in

HLW group (Figure. 4C). The pseudotemporal trajectory and RNA velocity analysis

verified that PDE4D$^+$/PDE7B$^+$ subclusters could differentiate in two different

directions, SCD$^+$/DGAT2$^+$ and FABP5$^+$/SIAH1$^+$ subclusters with two bifurcations

(Figures. 4D-F). The pseudotemporal heatmap showed gene expression dynamics

including *COX1*, *ANO4*, *PPARG*, *ADIPOQ*, *ACACA*, and *ELOVL6* at Point 2 (Figure.

S3A). UMAP plots also showed transcriptional dynamics of marker genes such as

FABP4, *ADIPOQ*, *MYBPC1*, and *EYV4* (Figures. S3B-D). Also, we discovered the

expression of preadipocytes related genes including *PDGFRA*, *CD34*, *CD38*, and

WT1 was enriched in PDE4D$^+$/PDE7B$^+$ adipocytes and mature adipocytes related

genes including *FABP4*, *ADIPOQ*, *LIPE*, *PLIN1*, *PPARG*, and *AGT* were enriched in

SCD$^+$/DGAT2$^+$ and FABP5$^+$/SIAH1$^+$ adipocytes (Figure. 4G). Hence, the

differentiated trajectory of mature adipocytes in muscles was displayed in Figure. 4H

based on above results, which showed that PDE4D$^+$/PDE7B$^+$ adipocytes could

differentiate into SCD$^+$/DGAT2$^+$ and FABP5$^+$/SIAH1$^+$ adipocytes, and

FABP5$^+$/SIAH1$^+$ adipocytes also can differentiate into SCD$^+$/DGAT2$^+$ adipocytes.

Additionally, the expression of *SCD*, *DGAT2*, *FABP5*, and *SIAH1* was upregulated

but *PDE4D* and *PDE7B* expression were downregulated in HLW group (Figure. 4I).

These data indicated the differentiated trajectory of mature adipocytes in muscle of

pigs and the percentage of SCD$^+$/DGAT2$^+$ and FABP5$^+$/SIAH1$^+$ subclusters was

higher in HLW group.

Transcriptional dynamics of glycerophospholipid metabolism in high IMF

deposition pigs

Our previous studies have discovered the changes of glycerophospholipid

metabolism in muscles after CLA treatment(L. Wang et al., 2022) and in high IMF

content Laiwu pigs(Wang, Zhao, et al., 2023). To further investigate the

transcriptional dynamics of glycerophospholipid metabolism in high IMF deposition

pigs, we then compare the gene program in Heigai pigs and Laiwu pigs. After CLA

treatment, the snRNA-seq dataset the expression of genes involved in the

glycerophospholipid metabolism in different groups and subclusters was displayed

slight differences in (Figures. S4A-C). Also, in Laiwu pigs, there are differences in

gene program involved in the glycerophospholipid metabolism between two groups

(Figures. S4D-F). Interestingly, we found *LCLAT1* was enriched in PDE4D$^+$/PDE7B$^+$

subcluster and *AGPAT3* and *AGPAT5* was enriched in SCD$^+$/DGAT2$^+$ subcluster

(Figures. S5B and S5E). We also discovered the increase of diglycerides and

phosphatidylinositols   and   the   decrease   of   phosphatidic   acids   and

phosphatidylethanolamines in high IMF deposition pigs might due to the changes of

AGPAT3, *AGPAT4*, *AGPAT5*, *CEPT1*, and *CDIPT1* (Figures. S5C and S5F). These

data revealed the significant differences in lipid composition and distribution in LDM

of high IMF deposition pigs might be due to the different expression levels of

glycerophospholipid metabolism-related genes.

Cell-cell communication analysis showed the interaction between adipocytes and

FAPs/fibroblasts

To further explore the association between adipocytes nuclei and other cell

clusters, we applied cell-cell communication analysis by using CellPhoneDB on 8 cell

types in muscles of Heigai pigs and found adipocytes mainly interacted with ECs,

FAPs/fibroblasts, MuSCs, and pericytes (Figure. 5A). In addition, dot plot represented

stronger communication from adipocytes to other subclusters through LRP6, FGFR1,

and COL4A2 pathways in LDM of Heigai pigs (Figure. 5C). Next, we also applied

cell-cell communication analysis by using CellPhoneDB on 9 cell types in muscles of

Laiwu pigs and found adipocytes mainly interacted with SPs, FAPs/fibroblasts, ECs,

and pericytes (Figure. 5B). Also, dot plot showed stronger communication from

adipocytes to other subclusters through COL6A3, LAMC1, and THBS1 pathways in

LDM of Laiwu pigs (Figure. 5D). These results suggested adipocytes has tight

association with FAPs/fibroblasts.

Characterization of FAPs after CLAs treatment through clustering and

pseudotime analysis

Previous studies have demonstrated that FAPs could differentiate into mature

adipocytes and are the main cell sources of IMF cells(Joe et al., 2010; Uezumi et al.,

2010). To further investigate the effects of CLAs on occurrence mechanism of IMF

deposition, we then performed subcluster and pseudotemporal trajectory analysis on

FAPs/fibroblasts. First, UMAP plots displayed the cell distribution in different

subpopulations of FAPs/fibroblasts (Figure. 6A) and we identified 3 subpopulations

in FAPs/fibroblasts based on the expression of marker genes, including FAPs

(*PDGFRA*), Fibroblasts (*COL1A1*), and PDE4D$^+$/PDE7B$^+$ subclusters (*PDE4D* and

PDE7B) (Figure. 6B). In CLA group, we found the proportion of FAPs (70.40% *vs*.

37.06%) and PDE4D$^+$/PDE7B$^+$ (4.11% *vs*. 2.18%) was higher than that in CON group

but the proportion of in Fibroblasts (60. 76% *vs*. 25.49%) was lower (Figure. 6C). The

top 10 most DEGs between the 3 subpopulations were showed in the heatmap and the

bar plot displayed the significant enrichment of the signalling pathways in muscles by

using KEGG enrichment analyses and we also found calcium and cGMP-PKG

signalling pathway were enriched in PDE4D$^+$/PDE7B$^+$ subclusters (Figure. 6D).

Besides, dot plot showed CLAs upregulated the expression of preadipocytes-related

genes including *CD38* and *CD34*, adipogenic master genes including *ADIPOQ*,

FABP4, *PLIN1*, and *LIPE*, mature adipocyte marker genes including *CEBPA*, and

lipid metabolism-related genes including *LPL*, *ELOVL4*, *ACAA2* and *HACD2* in

FAPs/Fibroblasts (Figure. S5A). To further explore FAPs' differentiated trajectory,

we applied a pseudotemporal trajectory analysis and RNA velocity analysis of

FAPs/Fibroblasts. According to the results, we found FAPs could differentiate into

PDE4D$^+$/PDE7B$^+$ and Fibroblasts subpopulations (Figures. 6E and S5C). The

pseudotemporal heatmap also displayed transcriptional dynamics including *TIMP3*

and *THBS1* at Point 1 (Figure. 6F). UMAP plots also showed transcriptional

dynamics of marker genes in three subpopulations such as *ITGA5*, *HSPH1*, and *ETF1*

(Figures. S5B-E). To further identify the differentiated trajectory of IMF cells, we

isolated primary FAPs from pigs and found the expression of *FABP4*, *ADIPOQ*, *SCD*,

and *DGAT2* was significantly increased but *PDE4D* expression was significantly

downregulated during adipogenic differentiation *in vitro* (Figure. 6G). Besides, the

expression of *FABP5* and *SIAH1* was first significantly increased then significantly

decreased during adipogenic differentiation (Figure. 6G). Hence, FAPs may first

differentiate into PDE4D$^+$/PDE7B$^+$ preadipocytes and then differentiate into

PDE4D$^+$/PDE7B$^+$ adipocytes (Figure. 6H). These results displayed the differentiated

trajectory of preadipocytes into mature adipocytes in pig muscles and CLAs might

influence this process then affect IMF deposition.

CLAs promoted FAPs directed differentiation into SCD$^+$/DGAT2$^+$ subclusters

To further explore the cytological mechanism of CLAs regulating IMF

deposition, we next investigated the regulatory effects of CLAs on the differentiation

trajectory of FAPs differentiate into adipocytes. First, to explore the role of CLAs in

the adipogenic differentiation of FAPs, we isolated primary FAPs from piglets and

induced these adipogenic differentiation. Nile Red staining and OD490 results

revealed CLAs can promote the adipogenic differentiation of FAPs after adipogenic

differentiation for 8 days *in vitro* (Figures. 7A-B). Besides, the mRNA expression of

SCD, *SIAH1* and adipogenic genes, including *FABP4*, *PPARG*, and *FASN* were

significantly upregulated but *PDE4D* was significantly downregulated (Figure. 7C).

The protein levels of FABP4 and SCD1 were significantly upregulated and PDE4D

were significantly downregulated (Figure. 7D). In addition, the marker genes'

expression at key points of adipogenic differentiation such as *ADIPOQ*, *ELOVL6*,

ACACA, *ARBB1*, *NEB*, and *MYBPC1* were significantly upregulated and *THBS1* and

TIMP3 were significantly downregulated (Figure. S6A). Interestingly, we next found

MAPK signaling pathway were enriched in adipocytes nuclei especially

$SCD^+/DGAT2^+$ subcluster (Figure. 7E). The expression of MAPK signaling pathway

including ERK, c-Jun N-terminal kinase (JNK), and p38 signaling pathway related

genes were changed in muscle nuclei (Figures. 7F and S6B-C). Furthermore, we

observed that the protein levels of JNK phosphorylation were significantly decreased

after CLA treatment during FAPs adipogenic differentiation (Figure. 7G). Hence, we

next used JNK activator Anisomysin to explore the regulatory mechanism (Figure.

7H). Oil Red O staining result showed that after 24h 20 nM Anisomysin treatment, the

increased adipogenic differentiation of FAPs by CLA were significantly inhibited

(Figure. 7I). Also, CLA + Anisomysin group had the lower OD 490 value compared

with CLA group (Figure. 7J). Nile Red staining result also showed that Anisomysin

significantly inhibited the enhanced lipid droplets in FAPs after CLA treatment

(Figure. 7K). MAP2K4 expression significantly upregulated and the expression of

FABP4 and *SCD* significantly decrease after Anisomysin treatment (Figure. 7L).

These data indicated that CLAs may promote the directed differentiation of FAPs into

$SCD^+/DGAT2^+$ subclusters via inhibiting JNK signaling pathway.

Discussion

CLAs can serve as a nutritional intervention to regulate lipid deposition in skeletal muscle of human according to clinic trials(van Vliet et al., 2020). In animal production, CLAs could regulate meat quality in pigs and cattle, especially improve IMF content (L. Wang et al., 2022; Zhang et al., 2016). These studies pointed out that the CLAs plays a vital role in regulating fat infiltration in skeletal muscle. However, to date, the cellular mechanism of CLAs regulates lipid deposition has not been studied. Here, we utilized the 10x Genomics platform to identify the cell heterogeneity and transcriptional changes in muscles after CLAs treatment based on pig models. This study revealed the effects of CLAs on cell populations and molecular characteristics of muscles and highlighted the cytological mechanism of CLAs regulates pork quality in skeletal muscle.

CLAs is always found in ruminant animals and dairy products, it is a class of positional and geometric isomers of linoleic acids with a conjugated double bond. CLAs not only has anti-cancer, anti-hypertension, anti-adipogenic, and anti-diabetic effects, but also can improve muscle function and decrease body fat percentage. For example, adding 3.2 g/day CLAs significantly increased muscle mass in higher body fat percentage Chinese adults (Chang et al., 2020). After 0.9 g/day CLAs supplementation, body weight variation and muscle mass significantly increased and body fat percentage variation decreased in student athletes (Terasawa, Okamoto,

Nakada, & Masuda, 2017). LC-MS metabolomics results discovered CLAs changed

57 metabolites which enriched in lipids/lipid-like molecules in plasma of humans (He

et al., 2022). However, another study found that for sedentary older adults, CLAs had

no significant influence on muscle anabolic effects (van Vliet et al., 2020). Therefore,

the specific influences of CLAs on skeletal muscle is still disputed and the function on

lipid deposition in human skeletal muscle needs further investigation. In animal

models, many studies have demonstrated the important effects of CLAs on regulating

fat accumulation in skeletal muscles. In porcine models, our foregoing studies have

discovered that adding CLAs into the pig diet could significantly increase IMF

contents in LDM of lean pig breeds and Heigai pigs(Wang et al., 2021; L. Wang et al.,

2022). Zhang *et al* (Zhang et al., 2016)found 2% dietary CLAs significantly increased

IMF deposition and reduced subcutaneous fat deposition in cattle. However, in mouse

model, 0.5% mixed isomer CLAs did not lead to lipid accumulation in muscle of

mice(M, S, & M, 2013). Hence, CLAs supplementation positively affect IMF

deposition in muscles of pigs and ruminants but the effects of CLAs may have

species-specific. In our study, we identified 8 cell types in skeletal muscles of Heigai

pigs, including myofibers, FAPs/fibroblasts, ECs, adipocytes, immune cells, MuSCs,

myeloid derived cells and pericytes. Recently, a variety of single cell studies have

been performed on mouse/human skeletal muscles and they also identified some cell

populations such as myofibers, FAPs/fibroblasts, ECs, adipocytes, immune cells,

MuSCs, myeloid derived cells, tenocytes, SPs, and pericytes(Petrany et al., 2020; D.

D. Xu et al., 2023; Z. Xu et al., 2020). We also found CLAs improved TG content and

increased the percentage of adipocytes in LDM. Previous study demonstrated there

are three subclusters in adipocytes and the formation and deposition of IMF mainly

relied on $DGAT2^+/SCD^+$ adipocytes and $FABP5^+/SIAH1^+$ adipocytes(Wang, Zhao, et

al., 2023). Specially, we found CLAs enhanced the percentage of $SCD^+/DGAT2^+$

subclusters. These indicated CLAs improve IMF deposition might through increasing

$SCD^+/DGAT2^+$subpopulations of adipocytes. Our findings could provide a foundation

for using nutritional strategies to increase pork quality especially IMF deposition.

However, the specific function of CLAs on fat infiltration and deposition of skeletal

muscle in people and rodents needs further study.

Skeletal muscle contains slow and fast muscle fibers and there are four major

muscle fiber types in mice, including slow muscle fibers with type I muscle fibers

(*Myh7*), fast muscle fibers with type IIA muscle fibers (*Myh2*), type IIX muscle fibers

(*Myh1*), and type IIB muscle fibers (*Myh4*)(Dos Santos et al., 2020; Petrany et al.,

2020). In pigs, illustrated there are three myofiber composition including type I, type

IIA, and type IIB in skeletal muscles by ST technology(Jin et al., 2021). In this study,

we identified six different subpopulations in myofibers including I myofibers, IIA

myofibers, IIX myofibers, IIB myofibers, MTJ, and NMJ and IIB myofibers had the

highest percentage in pig muscles. MTJ are known to exhibit structural specialization;

NMJ are responsible for formation and maintenance of the synaptic apparatus and

previous studies have identified these cell populations in murine skeletal muscles by

using snRNA-seq(Dos Santos et al., 2020; Petrany et al., 2020). We also identified

MTJ and NMJ cell populations in LDM of Laiwu pigs(Wang, Zhao, et al., 2023).

These results indicate it also exist MTJ and NMJ cell populations in porcine skeletal

muscles. In our previous study, we also found the percentage of type IIa myofibers

had an increased tendency and type IIb myofibers had a decreased tendency in high

IMF content Laiwu pigs(Wang, Zhao, et al., 2023). These results suggest that IMF

content is closely related to muscle fiber type. Besides, slow muscle fibers always

called slow-twitch oxidative muscle fibers like I myofibers have higher activities in

mitochondrial oxidative metabolic enzymes and myoglobin while fast muscle fibers

always called fast-twitch glycolytic muscle fibers like II myofibers have higher levels

of glycolytic enzymes and glycogen (Schiaffino & Reggiani, 2011). Similarly, we also

found I myofibers enriched in metabolic pathways, oxidative phosphorylation, and

thermogenesis. In recent years, studies have focused on exploring the influences of

CLAs on regulating muscle fiber type. In commercial pigs, the MyHC I mRNA

abundance were improved in LDM of the CLAs group (Men, Deng, Xu, Tao, & Qi,

2013). In mice, t10, c12-CLAs, but not c9, t11-CLAs can increase oxidative skeletal

muscle fiber type in gastrocnemius muscle and C2C12 myoblasts (Duan et al., 2021).

CLAs have been found to prevent sarcopenia by maintaining redox balance during

aging, actively regulating mitochondrial adaptation, improve muscle metabolism, and

inducing hypertrophy of type IIX myofibers after endurance exercise(Barone et al.,

2017; Chen, Yang, & Park, 2018). Also, we discovered CLAs enhanced the

percentage of I and IIA myofibers but reduced the percentage of IIB myofibers.

Previous study has found PPARγ coactivator-1α (PGC1α) serves a valuable role in

skeletal muscle metabolism and is a master regulator of oxidative phosphorylation

genes and could regulate muscle fiber type transformation(Handschin & Spiegelman,

2011). In our study, the PGC1α expression was also increased after CLA treatment in

myofiber. These results suggested CLAs can promote glycolytic skeletal muscle fiber

types switching into oxidative skeletal muscle fiber types through upregulating

PGC1α expression.

Numerous studies have discovered that the cell sources of IMF cells and found

several cell subsets lead to the ectopic IMF formation and deposition including SCs,

Myf5[+] MSCs, FAPs, ECs, pericytes, Fibroblasts, myeloid-derived cells, SPs, and PICs

(Sciorati, Clementi, Manfredi, & Rovere-Querini, 2015; Z. Xu et al., 2020). In this

study, we found CLAs increased the percentage of preadipocytes such as FAPs, ECs,

myeloid-derived cells, and pericytes. Importantly, FAPs are the major source of IMF

cells(Joe et al., 2010; Uezumi et al., 2010) and our previous study also verified the

adipogenic capacity of FAPs in 2D and 3D culture models(Wang, Zhao, et al., 2023).

Xu *et al.* found that FAPs serve as a cellular interaction hub in skeletal muscle of

pigs(D. D. Xu et al., 2023). We used pseudotemporal trajectory and RNA velocity

analysis combined with *in vitro* study to investigate that FAPs could first differentiate

into PDE4D[+]/PDE7B[+] adipocytes and then differentiate into DGAT2[+]/ SCD[+] and

FABP5[+]/SIAH1[+] adipocytes. However, the regulatory mechanism of FAPs directional

differentiation still needs to be further explored. *In vitro* studies demonstrated trans-10,

cis-12 CLAs inhibited skeletal muscle differentiation in C2C12 cells and inhibited

3T3-L1 adipocyte adipogenesis (Hommelberg et al., 2010; Yeganeh, Taylor, Poole,

Tworek, & Zahradka, 2016). However, the influences of CLAs on regulating the

adipogenic differentiation of FAPs are still unclear. In this study, we found CLAs

facilitated FAPs differentiating into $SCD^+/DGAT2^+$ adipocytes. Previous studies have

found mice orally treated with CLAs mixture upregulated *Scd1* expression in

muscle(Parra, Serra, & Palou, 2012). Besides, SCD1 expression is modulated by

mTOR signaling pathway in cancer cells(J. M. Yi, Zhu, Wu, Thompson, & Jiang,

2020; S. H. Zhao et al., 2021). Moreover, MAPK signaling pathway were enriched in

adipogenic differentiation of FAPs after CLAs treatment. Previous studies have

discovered JNK had negative effects on regulating the adipogenic differentiation of

human mesenchymal stem cells(Jang et al., 2015) and FAPs could prevent skeletal

muscle regeneration after muscle injury by ST2/JNK signaling pathways(Yamakawa

et al., 2023). In this study, based on JNK signing pathway activator treatment and *in*

vitro experiment, we found CLAs may promote FAPs directed differentiation into

$SCD^+/DGAT2^+$ adipocytes via inhibiting JNK signaling pathway. These results may

provide new targets for treating human fat infiltrated diseases by nutritional strategies.

However, we did not further explore the mechanistic action and the downstream

transcriptional regulators need to be discussed.

In a word, we provide detailed insights into the cytological mechanism of CLAs

regulates fat infiltration in skeletal muscles based on pig models via using snRNA-seq.

We analysed the effects of CLAs on the cell heterogeneity and transcriptional

dynamics in pig muscles and discovered CLAs could promote glycolytic muscle fiber

types switching into oxidative muscle fiber types through regulating PGC1α. We also

identified the differentiation trajectories of adipocytes and FAPs. Our data also

demonstrated CLAs could promote FAPs differentiate into DGAT2$^+$/SCD$^+$ adipocytes

via inhibiting JNK signalling pathway. This study provides a new way of developing

nutritional strategies to combat myosteatosis and other muscle-related diseases and

also offers potential opportunities to promote the utilization of pigs as animal models

to study human diseases.

Materials and methods

Animals and samples

The Zhejiang University Animal Care and Use Committee approved all

procedures and housing (ZJU20170466). 56 Heigai pigs (average body weight: 85.58

± 10.39 kg) were divided randomly into CON group (added 1% soyabean oil) and

CLA group (added 1% CLAs) for 40 days (5 days pre-feeding period and 35 days

formal test period) and the nutritional levels and the feeding process as we previously

reported(L. Wang et al., 2022; Wang, Zhang, Huang, Zhou, & Shan, 2023). At the end

of experiment, we collected LDM from the right side of the carcass for subsequent

immunofluorescence staining, biochemical assay, and snRNA-seq analyses. For

Laiwu pigs, based on the determination of IMF content in Laiwu pigs, we divided

them into two groups: HLW group and low IMF content Laiwu pigs (LLW) group.

The two most representative samples from each group were selected for later

snRNA-seq and datasets generated from muscle of high IMF content pig samples

were downloaded from the Genome Sequence Archive (Genomics, Proteomics &

Bioinformatics 2021) in National Genomics Data Center (Nucleic Acids Res 2022),

China National Center for Bioinformation/Beijing Institute of Genomics, Chinese

Academy of Sciences (GSA: CRA011059; https://ngdc.cncb.ac.cn/gsa) as we

previously discussed (Wang, Zhao, et al., 2023).

Triglycerides (TG) and total cholesterol (TC) determination

The contents of TG and TC in LDM were determined by commercial kits (TG,

E1025-105; TC, E1015-50) bought from Beijing APPLYGEN Gene Technology Co.,

LTD.

Immunofluorescence staining

The paraffin section was dewaxed and immersed in pre-heated sodium citrate,

then placed in a microwave oven and heated for 15 min to perform antigen retrieval.

Fixed the sections that cooled to room temperature in 4% paraformaldehyde for 10

min, followed by 10 min permeated with 0.5% Triton-X100and 1 h blocked with

blocking buffer (5% goat serum and 2% BSA). Sections were then incubated

overnight at 4 °C with Perilipin (Abcam, ab16667, 1:500), MF20 (Developmental

Studies Hybridoma Bank, 1:50), and SCD1 (HuaBio, ER1916-26, 1:500) primary

antibodies. Then the primary antibody was discarded and the sections were washed

three times with PBS for 5 min each time. Incubated the sections with secondary

antibodies for 1h and DAPI for 5 min, then washed with PBS. Sealed cell with

glycerol and used fluorescent microscope to capture images.

LDM nuclei isolation and 10X Genomics Chromium library and sequencing

LDM nuclei isolation and 10X Genomics Chromium library and sequencing

were performed by LC-Bio Technology Co., Ltd. (Hangzhou, China) as previous

published paper(Wang, Zhao, et al., 2023). Briefly, nuclei of LDM samples were

isolated, then homogenized and incubated for 5 min on ice. The homogenate was then

filtered, centrifuged and collected. The pellet was then resuspended, washed by the

buffer, incubated and centrifuged. After centrifugation, the pellet was resuspended,

filtered, and counted. Single-cell suspensions were loaded onto 10X Genomics

Chromium for capturing 5000 single cells, followed by cDNA amplification and

library construction steps were performed. Libraries were sequenced using the

Illumina NovaSeq 6000 sequencing system.

Bioinformatics analysis

SnRNA-seq results were demultiplexed and converted to FASTQ format by

using Illumina bcl2fastq software and followingly processed by the Cell Ranger. Then,

the Seurat packages was used to analysis the cell Ranger output. After the quality

control, 22540 cells were obtained. DoubletFinder package was used to remove

doublets and Harmony package was used to performed batch correction of data

integration between samples. We further used Seurat, UMAP, the FindAllMarkers

function to visualize the data, find clusters, and select marker genes. Monocle 2

package was used to perform trajectory analysis and model differentiation trajectories.

RNA velocity analysis was independently performed in FAPs by SAMTools and the

Velocyto (Bergen, Lange, Peidli, Wolf, & Theis, 2020). CellPhoneDB package was

used to cell communication analysis and make further speculations about potential

cellular interaction mechanisms.

Primary FAPs isolation, magnetic cell sorting and cell culture

Primary FAPs isolation were performed as previously described (Wang, Zhao, et

al., 2023). Briefly, a piece of muscle from a 3-day-old piglet was minced, and added 5

times the volume of 0.2% collagenase type I, then digested at 37 °C for 1 hour., After

filtering and centrifuging, adding red blood cell lysate to split for 5 min at 4 °C

followed by incubated with a Dead Cell Removal Kit at room temperature for 15 min.

Then adding CD140a antibody, incubating and centrifugating. Next, added 20 μL

antibiotin microbeads, incubation and centrifugation. After passing through the

magnetic column, the cells on the adsorption column were PDGFRα⁺ cells. For FAPs

adipogenic differentiation, when the cells confluence reached 90%, the10% FBS

growth medium was replaced with induction medium after 4 days, the medium was

changed to differentiation mediumand cultured for another 4 days until the adipocytes

were mature.

Nile Red staining

Rinse cultured FAPs with 1 x PBS 3 times, discard PBS, fix FAPs with 4%

formaldehyde for 15 min, repeat the rinsing step, and then add Nile Red solution

(1:500 for lipid droplet staining) and DAPI (1:500 for nuclei staining) for 5 min.

Sealed cell with glycerol and used fluorescent microscope to capture images.

Total RNA extraction and quantitative real-time PCR (qPCR)

Total RNA extraction and qPCR were conducted as described before (Shan et al.,

2016). Briefly, total RNA of FAPs were extracted by using TRIzol and the

Spectrophotometer and a ReverAid First Strand cDNA Synthesis Kit were used to

measure the purity and concentration of total RNA and reversed RNA samples. qPCR

was performed by using Applied Biosystems StepOnePlus Real-Time PCR System

with Hieff qPCR SYBR® Green Master Mix and gene-specific primers

(Supplementary Table 1). Relative changes in gene expression were analysed using

the $2^{-\Delta\Delta CT}$ method and normalized using 18S ribosomal RNA as an internal control.

Protein extraction and western blotting

Protein extraction and western blotting were carried out as mentioned previously (Shan et al., 2016). In brief, total proteins were isolated from cells or tissues with RIPA buffer. After measuring the concentrations, proteins were separated using SDS-PAGE and subsequently transferred to a polyvinylidene fluoride membrane (PVDF, Millipore Corporation). Then blocked PVDF membrane with blocking buffer (5% fat-free milk) for 1 h and incubated with primary antibodies overnight at 4 °C. The peroxisome proliferator-activated receptorγ (PPARγ) (C26H12, 1:1000) were purchased from Cell Signaling Technology (CST). The β-actin (M1210-2, 1:10000), FABP4 (E71703-98, 1:2000), SCD1 (ER1916-26, 1:1000), PDE4D (ER1916-26, 1:500), p-JNK1/2/3(T183+T183+T221) (ET1609-42, 1:2000), and JNK1/2/3 (ET1601-28, 1:2000) antibody were from HuaBio. Dilution of the secondary antibody was 500-fold. The ChemiScope3500 Mini-System was used for protein detected.

Statistical analysis

GraphPad (Prism 8.3.0) was used for data analyses and R software (version 4.3.2) was used for data visualization. Data comparisons were made by unpaired two-tailed Student's t tests and one-way ANOVA analysis. Differences were considered significant at $P < 0.05$.

Author contribution

Liyi Wang: conceptualization, data curation, formal analysis, investigation,

methodology, software, validation, visualization, writing-original draft. **Shiqi Liu:**

investigation, methodology, writing-review & editing. **Shu Zhang:** investigation.

Yizhen Wang: resources, supervision. **Yanbing Zhou:** conceptualization,

investigation, methodology, supervision. **Tizhong Shan:** conceptualization, funding

acquisition, project administration, resources, supervision, writing- review & editing.

Acknowledgement

We thank members of the Shan Laboratory for comments and this work was

partially supported by the National Natural Science Foundation of China (32272887),

the Natural Science Foundation of Zhejiang Province (LZ22C170003), and the

"Hundred Talents Program" funding from Zhejiang University to TZS.

Data availability

Data required to reanalyse the data reported in this paper is available from the

lead contact (tzshan@zju.edu.cn) upon request.

Conflict of interest

The authors declare no conflict of interest with the contents of this article.

References

Asakura, A., Komaki, M., & Rudnicki, M. A. (2001) **Muscle satellite cells are multipotential stem cells that exhibit myogenic, osteogenic, and adipogenic differentiation** *Differentiation* **68**: 245-253. https://doi:10.1046/j.1432-0436.2001.680412.x

Barone, R., Sangiorgi, C., Marino Gammazza, A., D'Amico, D., Salerno, M., Cappello, F., . . . Macaluso, F. (2017) **Effects of Conjugated Linoleic Acid Associated With Endurance Exercise on Muscle Fibres and Peroxisome Proliferator-Activated Receptor gamma Coactivator 1 alpha Isoforms** *J Cell Physiol* **232**: 1086-1094. https://doi:10.1002/jcp.25511

Bergen, V., Lange, M., Peidli, S., Wolf, F. A., & Theis, F. J. (2020) **Generalizing RNA velocity to transient cell states through dynamical modeling** *Nat Biotechnol* **38**: 1408-1414. https://doi:10.1038/s41587-020-0591-3

Biferali, B., Bianconi, V., Perez, D. F., Kronawitter, S. P., Marullo, F., Maggio, R., . . . Mozzetta, C. (2021) **Prdm16-mediated H3K9 methylation controls fibro-adipogenic progenitors identity during skeletal muscle repair** *Sci Adv* **7**: eabd9371. https://doi:10.1126/sciadv.abd9371

Biltz, N. K., Collins, K. H., Shen, K. C., Schwartz, K., Harris, C. A., & Meyer, G. A. (2020) **Infiltration of intramuscular adipose tissue impairs skeletal muscle contraction** *J Physiol* **598**: 2669-2683. https://doi:10.1113/JP279595

Chang, H., Gan, W., Liao, X., Wei, J. X., Lu, M. N., Chen, H. T., . . . Liu, X. (2020) **Conjugated linoleic acid supplements preserve muscle in high-body-fat**

adults: A double-blind, randomized, placebo trial *Nutrition Metabolism and Cardiovascular Diseases* **30**: 1777-1784. https://doi:10.1016/j.numecd.2020.05.029

Chen, P. B., Yang, J. S., & Park, Y. (2018) **Adaptations of Skeletal Muscle Mitochondria to Obesity, Exercise, and Polyunsaturated Fatty Acids** *Lipids* **53**: 271-278. https://doi:10.1002/lipd.12037

Dos Santos, M., Backer, S., Saintpierre, B., Izac, B., Andrieu, M., Letourneur, F., . . . Maire, P. (2020) **Single-nucleus RNA-seq and FISH identify coordinated transcriptional activity in mammalian myofibers** *Nat Commun* **11**: 5102. https://doi:10.1038/s41467-020-18789-8

Duan, C., Yin, C., Ma, Z., Li, F., Zhang, F., Yang, Q., . . . Wang, S. (2021) **trans 10, cis 12, but Not cis 9, trans 11 Conjugated Linoleic Acid Isomer Enhances Exercise Endurance by Increasing Oxidative Skeletal Muscle Fiber Type via Toll-like Receptor 4 Signaling in Mice** *J Agric Food Chem* **69**: 15636-15648. https://doi:10.1021/acs.jafc.1c06280

Farrington-Rock, C., Crofts, N. J., Doherty, M. J., Ashton, B. A., Griffin-Jones, C., & Canfield, A. E. (2004) **Chondrogenic and adipogenic potential of microvascular pericytes** *Circulation* **110**: 2226-2232. https://doi:10.1161/01.CIR.0000144457.55518.E5

Gilsanz, V., Kremer, A., Mo, A. O., Wren, T. A., & Kremer, R. (2010) **Vitamin D**

status and its relation to muscle mass and muscle fat in young women *J*

Clin Endocrinol Metab **95**: 1595-1601. https://doi:10.1210/jc.2009-2309

Groenen, M. A. M., Archibald, A. L., Uenishi, H., Tuggle, C. K., Takeuchi, Y.,

Rothschild, M. F., . . . Schook, L. B. (2012) **Analyses of pig genomes provide**

insight into porcine demography and evolution *Nature* **491**: 393-398.

https://doi:10.1038/nature11622

Handschin, C., & Spiegelman, B. M. (2011) **PGC-1 coactivators and the regulation**

of skeletal muscle fiber-type determination *Cell Metab* **13**: 351.

https://doi:10.1016/j.cmet.2011.03.008

Hausman, G. J., Basu, U., Du, M., Fernyhough-Culver, M., & Dodson, M. V. (2014)

Intermuscular and intramuscular adipose tissues: Bad vs. good adipose

tissues *Adipocyte* **3**: 242-255. https://doi:10.4161/adip.28546

He, Y. F., Xu, K., Li, Y. F., Chang, H., Liao, X., Yu, H., . . . Shi, L. (2022)

Metabolomic Changes Upon Conjugated Linoleic Acid Supplementation

and Predictions of Body Composition Responsiveness *J Clin Endocrinol*

Metab **107**: 2606-2615. https://doi:10.1210/clinem/dgac367

Hommelberg, P. P. H., Plat, J., Remels, A. H. V., van Essen, A. L. M., Kelders, M. C. J.

M., Mensink, R. P., . . . Langen, R. C. J. (2010) **Trans-10, cis-12 conjugated**

linoleic acid inhibits skeletal muscle differentiation and GLUT4

651  **expression independently from NF-kappa B activation** *Mol Nutr Food Res*

652  **54**: 1763-1772. https://doi:10.1002/mnfr.201000103

653 Jang, Y. J., Koo, H. J., Sohn, E. H., Kang, S. C., Rhee, D. K., & Pyo, S. (2015)

654  **Theobromine inhibits differentiation of 3T3-L1 cells during the early**

655  **stage of adipogenesis via AMPK and MAPK signaling pathways** *Food*

656  *Funct* **6**: 2365-2374. https://doi:10.1039/c5fo00397k

657 Jiang, Z., Marriott, K., & Maly, M. R. (2019) **Impact of Inter- and Intramuscular**

658  **Fat on Muscle Architecture and Capacity** *Crit Rev Biomed Eng* **47**: 515-533.

659  https://doi:10.1615/CritRevBiomedEng.2020031124

660 Jin, L., Tang, Q., Hu, S., Chen, Z., Zhou, X., Zeng, B., . . . Li, M. (2021) **A pig**

661  **BodyMap transcriptome reveals diverse tissue physiologies and**

662  **evolutionary dynamics of transcription** *Nat Commun* **12**: 3715.

663  https://doi:10.1038/s41467-021-23560-8

664 Joe, A. W., Yi, L., Natarajan, A., Le Grand, F., So, L., Wang, J., . . . Rossi, F. M. (2010)

665  **Muscle injury activates resident fibro/adipogenic progenitors that**

666  **facilitate myogenesis** *Nat Cell Biol* **12**: 153-163. https://doi:10.1038/ncb2015

667 Lang, I., Schweizer, A., Hiden, U., Ghaffari-Tabrizi, N., Hagendorfer, G., Bilban,

668  M., . . . Desoye, G. (2008) **Human fetal placental endothelial cells have a**

669  **mature arterial and a juvenile venous phenotype with adipogenic and**

670  **osteogenic differentiation potential** *Differentiation* **76**: 1031-1043.

671  https://doi:10.1111/j.1432-0436.2008.00302.x

Lunney, J. K., Van Goor, A., Walker, K. E., Hailstock, T., Franklin, J., & Dai, C. H.

(2021) **Importance of the pig as a human biomedical model** *Sci Transl Med*

13: eabd5758. https://doi:10.1126/scitranslmed.abd5758

M, K. K., S, I., & M, B. K. (2013) **Mice do not accumulate muscle lipid in response**

to dietary conjugated linoleic acid. *J Anim Sci* **91**: 4705-4712.

Men, X. M., Deng, B., Xu, Z. W., Tao, X., & Qi, K. K. (2013) **Age-related changes**

and nutritional regulation of myosin heavy-chain composition in

longissimus dorsi of commercial pigs *Animal* **7**: 1486-1492.

https://doi:10.1017/S1751731113000992

Mitchell, K. J., Pannerec, A., Cadot, B., Parlakian, A., Besson, V., Gomes, E. R., . . .

Sassoon, D. A. (2010) **Identification and characterization of a non-satellite**

cell muscle resident progenitor during postnatal development *Nat Cell*

Biol **12**: 257-U256. https://doi:10.1038/ncb2025

Parra, P., Serra, F., & Palou, A. (2012) **Transcriptional analysis reveals a high**

impact of conjugated linoleic acid on stearoyl-Coenzyme A desaturase 1

mRNA expression in mice gastrocnemius muscle *Genes Nutr* **7**: 537-548.

https://doi:10.1007/s12263-011-0279-x

Petrany, M. J., Swoboda, C. O., Sun, C., Chetal, K., Chen, X., Weirauch, M. T., . . .

Millay, D. P. (2020) **Single-nucleus RNA-seq identifies transcriptional**

heterogeneity in multinucleated skeletal myofibers *Nat Commun* **11**: 6374.

https://doi:10.1038/s41467-020-20063-w

Sahlin, K. R., & Trewern, J. (2022) **A systematic review of the definitions and**
interpretations in scientific literature of 'less but better' meat in
high-income settings *Nature Food* **3**: 454-460.
https://doi:10.1038/s43016-022-00536-5

Schiaffino, S., & Reggiani, C. (2011) **Fiber types in mammalian skeletal muscles**
Physiol Rev **91**: 1447-1531. https://doi:10.1152/physrev.00031.2010

Sciorati, C., Clementi, E., Manfredi, A. A., & Rovere-Querini, P. (2015) **Fat**
deposition and accumulation in the damaged and inflamed skeletal
muscle: cellular and molecular players *Cell Mol Life Sci* **72**: 2135-2156.
https://doi:10.1007/s00018-015-1857-7

Shan, T., Xiong, Y., Zhang, P., Li, Z., Jiang, Q., Bi, P., . . . Kuang, S. (2016) **Lkb1**
controls brown adipose tissue growth and thermogenesis by regulating the
intracellular localization of CRTC3 *Nat Commun* **7**: 12205.
https://doi:10.1038/ncomms12205

Tabula Muris, C., Overall, c., Logistical, c., Organ, c., processing, Library, p., . . .
Principal, i. (2018) **Single-cell transcriptomics of 20 mouse organs creates a**
Tabula Muris *Nature* **562**: 367-372. https://doi:10.1038/s41586-018-0590-4

Tamaki, T., Akatsuka, A., Ando, K., Nakamura, Y., Matsuzawa, H., Hotta, T., . . .
Edgerton, V. R. (2002) **Identification of myogenic-endothelial progenitor**
cells in the interstitial spaces of skeletal muscle *J Cell Biol* **157**: 571-577.
https://doi:10.1083/jcb.200112106

Terasawa, N., Okamoto, K., Nakada, K., & Masuda, K. (2017) **Effect of Conjugated**

Linoleic Acid Intake on Endurance Exercise Performance and

Anti-fatigue in Student Athletes *J Oleo Sci* **66**: 723-733.

https://doi:10.5650/jos.ess17053

Uezumi, A., Fukada, S., Yamamoto, N., Takeda, S., & Tsuchida, K. (2010)

Mesenchymal progenitors distinct from satellite cells contribute to ectopic

fat cell formation in skeletal muscle *Nat Cell Biol* **12**: 143-152.

https://doi:10.1038/ncb2014

van Vliet, S., Fappi, A., Reeds, D. N., & Mittendorfer, B. (2020) **No independent or**

combined effects of vitamin D and conjugated linoleic acids on muscle

protein synthesis in older adults: a randomized, double-blind,

placebo-controlled clinical trial *Am J Clin Nutr* **112**: 1382-1389.

https://doi:10.1093/ajcn/nqaa240

Wang, L., Huang, Y., Wang, Y., & Shan, T. (2021) **Effects of Polyunsaturated Fatty**

Acids Supplementation on the Meat Quality of Pigs: A Meta-Analysis

Front Nutr **8**: 746765. https://doi:10.3389/fnut.2021.746765

Wang, L., Nong, Q., Zhou, Y., Sun, Y., Chen, W., Xie, J., . . . Shan, T. (2022) **Changes**

in Serum Fatty Acid Composition and Metabolome-Microbiome

Responses of Heigai Pigs Induced by Dietary N-6/n-3 Polyunsaturated

Fatty Acid Ratio *Front Microbiol* **13**: 917558.

https://doi:10.3389/fmicb.2022.917558

Wang, L., Valencak, T. G., & Shan, T. (2024) **Fat infiltration in skeletal muscle:**

Influential triggers and regulatory mechanism *iScience* **27**: 109221.

https://doi:10.1016/j.isci.2024.109221

Wang, L., Zhang, S., Huang, Y., You, W., Zhou, Y., Chen, W., . . . Shan, T. (2022)

CLA improves the lipo-nutritional quality of pork and regulates the gut

microbiota in Heigai pigs *Food Funct*12093-12104.

https://doi:10.1039/d2fo02549c

Wang, L., Zhang, S., Huang, Y., Zhou, Y., & Shan, T. (2023) **Conjugated linoleic**

acids inhibit lipid deposition in subcutaneous adipose tissue and alter lipid

profiles in serum of pigs *J Anim Sci* **101**: skad294.

https://doi:10.1093/jas/skad294

Wang, L., Zhao, X., Liu, S., You, W., Huang, Y., Zhou, Y., . . . Shan, T. (2023)

Single-nucleus and bulk RNA sequencing reveal cellular and

transcriptional mechanisms underlying lipid dynamics in high marbled

pork *NPJ Sci Food* **7**: 23. https://doi:10.1038/s41538-023-00203-4

Wei, H., Zhou, Y., Jiang, S., Huang, F., Peng, J., & Jiang, S. (2016) **Transcriptional**

response of porcine skeletal muscle to feeding a linseed-enriched diet to

growing pigs *J Anim Sci Biotechnol* **7**: 6.

https://doi:10.1186/s40104-016-0064-1

Wosczyna, M. N., Perez Carbajal, E. E., Wagner, M. W., Paredes, S., Konishi, C. T.,

Liu, L., . . . Rando, T. A. (2021) **Targeting microRNA-mediated gene**

repression limits adipogenic conversion of skeletal muscle mesenchymal

stromal cells *Cell Stem Cell* **28**: 1323-1334 e1328.

https://doi:10.1016/j.stem.2021.04.008

Xu, D. D., Wan, B. Y., Qiu, K., Wang, Y. B., Zhang, X., Jiao, N., . . . Yin, J. D. (2023)

Single-Cell RNA-Sequencing Provides Insight into Skeletal Muscle

Evolution during the Selection of Muscle Characteristics *Advanced Science*.

https://doi:10.1002/advs.202305080

Xu, Z., You, W., Chen, W., Zhou, Y., Nong, Q., Valencak, T. G., . . . Shan, T. (2020)

Single□cell RNA sequencing and lipidomics reveal cell and lipid dynamics

of fat infiltration in skeletal muscle *Journal of Cachexia, Sarcopenia and*

Muscle **12**: 109-129. https://doi:10.1002/jcsm.12643

Yamakawa, D., Tsuboi, J., Kasahara, K., Matsuda, C., Nishimura, Y., Kodama, T., . . .

Inagaki, M. (2023) **Cilia-Mediated Insulin/Akt and ST2/JNK Signaling**

Pathways Regulate the Recovery of Muscle Injury *Adv Sci* **10**: e2202632.

https://doi:10.1002/advs.202202632

Yeganeh, A., Taylor, C. G., Poole, J., Tworek, L., & Zahradka, P. (2016) **Trans10,**

cis12 conjugated linoleic acid inhibits 3T3-L1 adipocyte adipogenesis by

elevating beta-catenin levels *Biochimica Et Biophysica Acta-Molecular and*

Cell Biology of Lipids **1861**: 363-370. https://doi:10.1016/j.bbalip.2016.01.004

Yi, J. M., Zhu, J. J., Wu, J., Thompson, C. B., & Jiang, X. J. (2020) **Oncogenic**

activation of PI3K-AKT-mTOR signaling suppresses ferroptosis via

SREBP-mediated lipogenesis *Proc Natl Acad Sci U S A* **117**: 31189-31197. https://doi:10.1073/pnas.2017152117

Yi, W., Huang, Q., Wang, Y., & Shan, T. (2023) **Lipo-nutritional quality of pork: The lipid composition, regulation, and molecular mechanisms of fatty acid deposition** *Anim Nutr* **13**: 373-385. https://doi:10.1016/j.aninu.2023.03.001

Yin, H., Pasut, A., Soleimani, V. D., Bentzinger, C. F., Antoun, G., Thorn, S., . . . Rudnicki, M. A. (2013) **MicroRNA-133 controls brown adipose determination in skeletal muscle satellite cells by targeting Prdm16** *Cell Metab* **17**: 210-224. https://doi:10.1016/j.cmet.2013.01.004

You, W. J., Liu, S. Q., Ji, J. F., Ling, D. F., Tu, Y., Zhou, Y. B., . . . Shan, T. Z. (2023) **Growth arrest and DNA damage-inducible alpha regulates muscle repair and fat infiltration through ATP synthase F1 subunit alpha** *Journal of Cachexia Sarcopenia and Muscle* **14**: 326-341. https://doi:10.1002/jcsm.13134

Zhang, H., Dong, X., Wang, Z., Zhou, A., Peng, Q., Zou, H., . . . Wang, L. (2016) **Dietary conjugated linoleic acids increase intramuscular fat deposition and decrease subcutaneous fat deposition in Yellow Breed x Simmental cattle** *Anim Sci J* **87**: 517-524. https://doi:10.1111/asj.12447

Zhao, L., Son, J. S., Wang, B., Tian, Q., Chen, Y., Liu, X., . . . Du, M. (2020) **Retinoic acid signalling in fibro/adipogenic progenitors robustly enhances muscle regeneration** *EBioMedicine* **60**: 103020. https://doi:10.1016/j.ebiom.2020.103020

798   Zhao, S. H., Cheng, L., Shi, Y., Li, J., Yun, Q. H., & Yang, H. (2021) **MIEF2**

799   **reprograms lipid metabolism to drive progression of ovarian cancer**

800   **through ROS/AKT/mTOR signaling pathway** *Cell Death Dis* **12**: 18.

801   https://doi:10.1038/s41419-020-03336-6

 Figure Legends

Figure. 1. SnRNA-seq identifies distinct cell populations after CLAs treatment in

pig muscles. **(A)** TG and TC content of LDM tissues in control and CLAs groups

(n=5). **(B)** LDM tissues stained with the adipogenic marker perilipin (red), muscle

fiber marker MyHC (green) and DAPI (blue) in different groups. Scale bars, 200 and

100 μm, respectively. **(C)** Scheme of the experimental design for snRNA-seq on

different muscles. **(D)** UMAP visualization of all of the isolated single nuclei from

Heigai pig muscles colored by cluster identity. **(E)** UMAP and violin plot displaying

the expression of selected marker genes for each cluster in pigs. **(F)** Nuclear

proportion in each cluster in pig muscles of control and CLAs groups. Each cluster is

color-coded. **(G)** Left, heatmap showing the top 10 most differentially expressed

genes between cell types identified. Right, KEGG enrichment for marker genes of

each cell type in muscles. Each lane represents a subcluster. Error bars represent SEM.

* $P < 0.05$, ** $P < 0.01$, two-tailed Student's t-test.

Figure. 2. Cell and transcriptional heterogeneity in myofibers. **(A)** UMAP plot

showing six subclusters of the isolated single nuclei from the control and CLAs

muscles. **(B)** UMAP and violin plot displaying the expression of selected marker

genes for each subcluster. **(C)** Cell proportion in each subcluster in different groups.

Each cluster is colour-coded. **(D)** Violin plot showing the expression of myofiber type

marker genes (*MYH7*, *MYH2*, *MYH1* and *MYH4*), myofiber type

transformation-related genes (*PPARGC1A*, and *STK11*), oxidation-related genes

(*COX5A*, *COX5B*, and *COX8A*), and glycolysis-related genes (*PFKM*, *HK2*, and

LDHC) after CLAs treatment. **(E)** The mRNA expression of myofiber type related

genes in LDM muscles after different treatment (n=6). **(F)** Heatmap representing the

top 10 most differently expressed genes between cell subclusters identified. **(G)**

KEGG enrichment for marker genes of each cell type in myofibers. I, type I

myonuclei; IIa, type IIa myonuclei; IIx, type IIx myonuclei; IIb, type IIb myonuclei;

MTJ, myotendinous junction nuclei; NMJ, neuromuscular junction nuclei. Error bars

represent SEM. *$P < 0.05$, two-tailed Student's t-test.

Figure. 3. Clustering and transcriptional dynamics of adipocytes. (A) UMAP plot

displaying the isolated single nuclei in three subclusters of adipocytes. **(B)** Bar plot

displaying the cell amounts in each subcluster in different groups. **(C)** Dot plot

showing the expression of three subcluster marker genes in muscle nuclei of Heigai

pigs. **(D)** The mRNA expression of three subcluster marker genes in LDM muscles

after different treatment (n=6). **(E)** LDM tissues stained with the adipogenic marker

perilipin (red), muscle fiber marker MyHC (green), SCD1 (pink), and DAPI (blue) in

different groups. Scale bars, 100 μm. **(F)** Left, heatmap showing the top 10 most

differentially expressed genes between cell types identified. Right, KEGG enrichment

for marker genes of each cell type in muscles. **(G)** Unsupervised pseudotime

trajectory of the three subtypes of adipocytes by RNA velocity analysis. Trajectory is

colored by cell subtypes. The arrow indicates the direction of cell pseudo-temporal

differentiation. **(H)** Scheme of the differentiation trajectories in mature adipocytes.

Error bars represent SEM. *$P < 0.05$, ** $P < 0.01$, two-tailed Student's t-test.

Figure. 4. Pseudotemporal and differentiated trajectories of adipocytes in high

IMF content Laiwu pig muscles. (A) Scheme of the experimental design for

snRNA-seq on adipocytes of high IMF content Laiwu pig muscles. **(B)** Cell

proportion of adipocytes subclusters in HLW and LLW groups. Each cluster is

color-coded. **(C)** LDM tissues stained with the adipogenic marker perilipin (red),

muscle fiber marker MyHC (green), SCD1 (pink), and DAPI (blue) in HLW and

LLW groups. Scale bars, 100 μm.   **(D-E)** Pseudotime ordering of all of adipocytes of

subcluster DGAT2$^+$/SCD$^+$, FABP5$^+$/SIAH1$^+$, and PDE4D$^+$/PDE7B$^+$. Each dot

represents one nucleus (color-coded by its identity), and each branch represents one

cell state. Pseudotime is shown colored in a gradient from dark to light blue, and the

start of pseudotime is indicated. Activation of the PDE4D$^+$/PDE7B$^+$ cluster can lead

to DGAT2$^+$/SCD$^+$ and FABP5$^+$/SIAH1$^+$ fate. **(F)** Unsupervised pseudotime trajectory

of the three subtypes of adipocytes by RNA velocity analysis. Trajectory is colored by

cell subtypes. The arrow indicates the direction of cell pseudo-temporal

differentiation. **(G)** Dot plot showing the expression of preadipocytes and mature

adipocytes- related genes in different subclusters. **(H)** Scheme of the differentiation

trajectories in mature adipocytes of Laiwu pigs. **(I)** Violin plot showing the

expression of three subcluster marker genes in different groups.

Figure. 5. Cell-cell communication analysis of adipocytes in pig muscles. (A)

Cell-cell communication analysis showed the network between adipocytes and other

clusters in muscles of Heigai pigs. **(B)** Cell-cell communication analysis showed the

network between adipocytes and other clusters in muscles of Laiwu pigs. **(C)** Dotplot

representing the gene expression and significance of the receptor-ligand relationship

in different cell population in muscles of Heigai pigs. **(D)** Dotplot representing the

gene expression and significance of the receptor-ligand relationship in different cell

population in muscles of Laiwu pigs. The larger the circle, the smaller the *P* value of

the relationship in the corresponding cell population, the more significant it is.

Figure. 6. Clustering and pseudotemporal trajectories of FAPs. (A) UMAP plot

showing three subclusters of the isolated single nuclei from control and CLAs muscle.

(B) Violin plot displaying the expression of selected marker genes for each subcluster.

(C) Cell proportion in each subcluster in different group. Each cluster is color-coded.

(D) Left, heatmap showing the top 10 most differentially expressed genes between

cell types identified. Right, KEGG enrichment for marker genes of each cell type in

muscles. **(E)** Pseudotime ordering of all of the FAP/fibroblast of subcluster FAPs,

Fibroblasts, and PDE4D$^+$/PDE7B$^+$. Each dot represents one nucleus (color-coded by

its identity), and each branch represents one cell state. Pseudotime is shown colored in

a gradient from dark to light blue, and the start of pseudotime is indicated. Activation

of the FAP cluster can lead to fibroblast fate or PDE4D$^+$/PDE7B$^+$ fate. **(F)**

Pseudotemporal heatmap showing gene expression dynamics for significant marker

genes. Genes (rows) were clustered into three modules, and cells (columns) were

ordered according to pseudotime in different groups. **(G)** The expression of

adipogenesis and three subcluster marker genes in differentiated FAPs in different

differentiation stage (n=6). **(H)** Scheme of the differentiation trajectories of

preadipocytes into mature adipocytes. Error bars represent SEM. *$P < 0.05$, ** $P <$

0.01, *** $P < 0.001$, two-tailed Student's t-test.

Figure. 7. The cytological mechanism of CLAs regulates FAPs differentiation. (A)

Dfferentiated FAPs stained with Nile Red (red) and DAPI (blue) in different groups.

Scale bars, 200 and 100 μm, respectively. **(B)** OD490 levels of total lipids in

differentiated FAPs after different treatment (n=4). **(C)** The mRNA expression of

three subcluster marker genes and adipogenic marker genes in differentiated FAPs

after different treatment (n=5). **(D)** Protein levels of FABP4, SCD1, and PDE4D were

detected by western blot. **(E)** MAPK signalling pathway enrichment in different cells.

(F) Dot plot showing the expression of MAPK signalling pathway related genes after

CLA treatment in FAPs/Fibroblasts. **(G)** Protein levels of P-JNK and JNK were

detected by western blot. **(H)** Scheme of CLAs regulating the differentiation

trajectories of FAPs into mature adipocytes. **(I)** Dfferentiated FAPs stained with Oil

Red O in different groups after treating with 20 nM Anisomysin. **(J)** OD490 levels of

total lipids in differentiated FAPs after different treatment (n=4). **(K)** Dfferentiated

FAPs stained with Nile Red (red) and DAPI (blue) after 20 nM Anisomysin treatment.

Scale bars, 200 μm. **(L)** The mRNA expression of adipogenic related genes in

differentiated FAPs after different treatment (n=4). Error bars represent SEM. *$P <$

0.05, ** $P < 0.01$, *** $P < 0.001$, two-tailed Student's t-test and one-way ANOVA

analysis. Letters represent statistical significance. Lowercase letters represent $P < 0.05$

and uppercase letters represent $P < 0.01$.

Figure 7