Immunopeptidomics-guided identification of functional neoantigens in non-small cell lung cancer

*Ben Nicholas[1,2], *Alistair Bailey[1,2], Katy J McCann[3], Oliver Wood[3], Eve Currall[4], Peter Johnson[5], Tim Elliott[2,6], Christian Ottensmeier[3,4] and Paul Skipp[1]

[1]Centre for Proteomic Research, Biological Sciences and Institute for Life Sciences, Building 85, University of Southampton, Southampton, United Kingdom

[2]Centre for Cancer Immunology and Institute for Life Sciences, Faculty of Medicine, University of Southampton, Southampton, United Kingdom

[3]School of Cancer Sciences, Faculty of Medicine, University of Southampton, Southampton, United Kingdom

[4]Institute of Systems, Molecular and Integrative Biology, University of Liverpool, Liverpool, United Kingdom

[5]Cancer Research UK Clinical Centre, University of Southampton, Southampton, United Kingdom

[6]Oxford Cancer Centre for Immuno-Oncology and CAMS-Oxford Institute, Nuffield Department of Medicine, University of Oxford, Oxford, United Kingdom

*These authors contributed equally

ORCIDs:

Ben Nicholas: 0000-0003-1467-9643

Alistair Bailey: 0000-0003-0023-8679

Katy J McCann: 0000-0003-4388-7265

Eve Currall: 0009-0008-0400-2376

Peter Johnson: 0000-0003-2306-4974

Tim Elliott: 0000-0003-1097-0222

Christian Ottensmeier: 0000-0003-3619-1657

Paul Skipp: 0000-0002-2995-2959

Corresponding author: Dr Ben Nicholas email: bln1@soton.ac.uk

Running title:

Proteogenomics identification of neoantigens in lung cancer

Keywords:

HLA, immunopeptidome, non-small cell lung cancer, antigen presentation

Abstract

Non-small cell lung cancer (NSCLC) has poor survival even with modern checkpoint inhibitor

therapies. Personalised vaccines based on short peptide neoantigens containing tumour

mutations are an attractive precision medicine strategy, but identifying therapeutically relevant

neoantigens remains challenging, with existing methods yielding positive responses in only 6%

of candidates tested.

We developed an immunopeptidomics approach to improve neoantigen identification in 24

NSCLC patients (15 adenocarcinoma, 9 squamous cell carcinoma). We directly identified one

neoantigen and using whole exome sequencing, transcriptomics and mass spectrometry-based

immunopeptidomics, we filtered predicted neoantigens based on observed cohort HLA peptide

presentation. This approach achieved positive functional responses in 5 of 6 patients tested

(83% success rate) with 13% of putative neoantigens (9 out of 70) eliciting strong responses.

Bayesian modelling of our initial rules-based neoantigen selection further revealed patient

specific peptide presentation patterns and propensities.

Our findings demonstrate that incorporating donor specific HLA peptide presentation data

substantially improves neoantigen identification success rates and immune response specificity,

advancing personalised cancer vaccine development.

Introduction

Lung cancer is the second most common cancer in the UK and despite recent advances in

checkpoint inhibitor therapies fewer than 20% of patients survive 5 years, with most surviving

less than one year post-diagnosis[1,2].The majority are diagnosed at advanced stages (III-IV) with

non-small cell lung cancer (NSCLC) accounting for 85-90% of cases. NSCLC comprises three

histological subtypes: adenocarcinoma (LUAD, most common, peripheral), squamous cell

carcinoma (LUSC, central), and large cell undifferentiated carcinoma (least common, can form

throughout the lung).

Four immunotherapies targeting PD-1/PD-L1 are licensed for NSCLC but show reduced efficacy

in patients with EGFR or ALK mutations[3,4]. Consequently, immunotherapy typically follows

chemotherapy and/or targeted therapy. While LUAD generally shows more favourable survival

than LUSC, treatment resistance remains a major challenge.

The TRACERx study revealed evolutionary processes underlying this resistance. Whole

genome doubling, driven by tobacco smoke and cytidine deaminase activity, protects tumours

against high mutation burdens and chromosomal instability[5]. Smoking-induced truncal mutations

and deaminase-driven branch mutations create extensive intratumour heterogeneity that

predicts recurrence and death, yet confounds biomarker utility for immunotherapy response

prediction[6].

An attractive strategy for NSCLC treatment is vaccination targeting on HLA presented

neoantigens. This approach assumes neoantigens can be identified that expand tumour killing

T-cell populations and/or modulate the tumour microenvironment to make T-cell infiltration or

checkpoint inhibitors more effective. The personalised nature of neoantigens minimise the risk

of off-target effects and autoimmunity. However, direct identification of neoantigens is rare[7,8]

and most approaches to neoantigen discovery rely on predicting that a given mutation leads to

protein synthesis, antigen processing and HLA presentation. Direct observation is rare in part

due to limits in the sensitivity of the mass spectrometry proteomic detection of HLA ligands,

known as immunopeptidomics. Moreover, it is estimated that only a small fraction of mutations

are actually presented, possibly as low as 0.5% of non-silent mutations[9]. For example, a

NSCLC tumour with 600 missense variants might yield only 3 presented neoantigens amongst a

lung tissue immunopeptidome of around 60,000 unique class I and II HLA peptides[10]. A typical

experiment may identify 3,000 of these peptides. Assuming a hypergeometric distribution, the

probability of observing one class I or II HLA neoantigen is about 14%. Or to put it another way,

there is around an 86% chance of not seeing any neoantigens in any single mass spectrometry

proteomics experiment.

Given these odds, much effort has been put into the in silico prediction of mutations that will

give rise to neoantigens that would make effective vaccines. There are well established

algorithms that can predict the likelihood of a peptide of given amino acid sequence binding to

an HLA molecule, and immunopeptidomic evidence of the peptide length preference of peptide

for different HLA allotypes[11], and preferential regions of proteins favourable for presentation[12–14].

However, even with this knowledge prediction is stymied by the number of potential neoantigen

candidates each mutation might yield, creating large lists of candidate peptides. Moreover, the

key biochemical and structural parameters of immunogenic neoantigens remain unknown. The

best neoantigen prediction models have a success rate such that around 6% of their putative

neoantigens are T-cell reactive[15,16], although recent machine learning models report increased

predictive power[17–20].

Here we adopted a patient-specific approach where, rather than relying solely on various

general HLA peptide characteristics, we leveraged each patient's HLA-I and -II

immunopeptidomes alongside the cohort level immunopeptidomes to understand their individual

presentation propensity. Thus, immunopeptidomics was used as circumstantial evidence of the

biological availability of a mutated protein for presentation by HLA. First we mapped the

mutational and immunopeptidome landscapes of a cohort of LUAD and LUSC patients. We then

predicted HLA-restricted neoantigens using existing algorithms and used immunopeptidomic

data from their individual tumours to filter those predictions on the basis of evidence that they

could be presented. Neoantigen selection success was increased to 13%, with strong functional

responses confirmed in 5 of 6 patients (83.3%) for our neoantigen panels selected for each

donor. Finally, we used Bayesian modelling of the cohort's immunopeptidomes to quantify how

donor-specific presentation propensity and protein-level evidence combine to determine peptide

presentation, providing a statistical framework for understanding our rules-based selection

success.

These proof-of-concept results demonstrate how the information contained within the

immunopeptidome has the potential to enhance proteogenomics strategies for identifying

neoantigens for every patient according to their specific presentation propensity, enabling

precision neoantigen selection that accounts for each patient's unique presentation biology.

Results

A proteogenomics workflow for neoantigen identification

Our NSCLC cohort consisted of 24 patients, 15 LUAD (8 female, 7 male) and 9 LUSC (5 female, 4 male). Median age at diagnosis was 69 (See Table 1 and Supplementary Data S1). Tumour tissue and PBMCs were used for HLA typing, whole exome sequencing, RNA sequencing and mass spectrometry proteomics of the HLA immunopeptidome (Figure 1). To identify candidate neoantigens for each patient we developed a workflow that surveyed both the genomic and immunopeptidomic landscapes. Somatic missense variants called from the whole exome sequencing (WES, Supplementary Data S2 and S3) were used to generate a mutanome for each individual against which the HLA immunopeptidome could be searched for direct observation of neoantigens. Variants, gene expression and the patient HLA allotypes were also used for the prediction of putative neoantigens using existing tools[21].

Table 1: Clinical summary of patients in this study with non-small cell lung cancer

Donor	Smoking status	Cancer subtype
A113	Current smoker	Squamous
A114	Ex smoker	Adenocarcinoma
A115	Ex smoker	Squamous
A116	Never smoker	Squamous
A117	Current smoker	Adenocarcinoma
A118	Ex smoker	Adenocarcinoma
A119	Ex smoker	Adenocarcinoma
A120	Ex smoker	Adenocarcinoma
A133	Ex smoker	Squamous
A134	Current smoker	Squamous
A136	Never smoker	Adenocarcinoma
A137	Ex smoker	Adenocarcinoma

A139	Ex smoker	Adenocarcinoma
A140	Ex smoker	Squamous
A141	Ex smoker	Adenocarcinoma
A142	Current smoker	Adenocarcinoma
A143	Ex smoker	Adenocarcinoma
A144	Ex smoker	Squamous
A145	Ex smoker	Squamous
A146	Current smoker	Adenocarcinoma
A147	Current smoker	Adenocarcinoma
A148	Ex smoker	Adenocarcinoma
A152	Ex smoker	Squamous
A153	Ex smoker	Adenocarcinoma

A

1.

Patient cohort:
Adenocarcinoma, n = 15
Squamous, n = 9
Sample collection:
Tumour/Normal, PBMCs

2. HLA typing

Whole exome sequencing

RNAseq

Mass spectrometry proteomics

3. Somatic variants

| A | T | G | C | A | C | T |
| A | T | G | G | A | C | T |

Gene expression

HLA I & II
immunopeptidomes

4. Neoantigen selection

Binding
algorithms *pVACseq*

Proteogenomics
guided selection

5. Neoantigen evaluation

B

Patient Samples

Tumor Tissue and Normal Adjacent Tissue

Whole Exome Sequencing

RNA Sequencing

Alignment to 1000 Genomes GRCh38/hg38 reference

Alignment and transcript quantification: HISAT2, 1000 Genomes GRCh38/hg38 reference & StringTie

Somatic Variant Calling: Mutect2, Strelka & Varscan Annotation with Variant Effect Predictor

PBMCs

HLA Typing

Variant Call File

pVACseq Phased proximal variants included 4 binding algorithms for HLA-I predictions: MHCflurry, MHCnuggetsI, NetMHC, PickPocket & 4 for HLA-II predictions: MHCnuggetsII, NetMHCIIpan, NNalign, SMMalign

Generate Patient Mutanome

Mass Spectrometry Proteomics

$Score = binding\ score + fold\ change + (variant\ expression \times fold\ change) + (tumour\ VAF/\ 2)$

HLA Immunopeptidome Analysis

HLA-I: 8mer peptides

HLA-I: 9mer peptides

HLA-I: 10mer peptides

HLA-I: 11mer peptides

HLA-II: 15mer peptides

Search HLA Immunopeptidome for Neoantigens

Binding affinity < 500 nM?

Yes   Yes   Yes   Yes   Yes

HLA-I: 8mer peptides filtered results

HLA-I: 9mer peptides filtered results

HLA-I: 10mer peptides filtered results

HLA-I: 11mer peptides filtered results

HLA-II: 15mer peptides filtered results

Source protein in immunopeptidome?

No   Yes

Correct HLA Pathway?

No   Yes

Preferred HLA Length?

No   Yes

Reject

Re-Ranking

Check Biological Relevance: Manual curation

COSMIC Database

Human Protein Atlas

Final selection

Yes

Functional T-cell assay: IFN-γ ELISpot

Figure 1: Integrated proteogenomics workflow. (A) HLA typing, whole exome sequencing, RNASeq and mass spectrometry-based proteomic of the HLA immunopeptidome were collected for 24 lung cancer patients providing mutational, gene expression and immunopeptidomic data from which to identify candidate neoantigens using binding algorithms and manual inspection of the combined proteogenomic data. (B) Flowchart of neoantigen selection workflow

The mutational landscape of NSCLC in the studied cohort

To assess the likelihood of identifying HLA presented neoantigens we first examined the mutational landscape of the NSCLC cohort and found it to be consistent with previous reports[22,23]. Somatic variants were identified by WES of tumour and matched normal adjacent tissues. Tumour mutational burden (TMB) quantifies the number of mutations per million bases (Mb). From WES it is calculated as the number of variants divided by the size of the exome targets; here the target size was 35.7 Mb and the number of variants were either the total number of all variants, or only the protein coding missense variants: $N_{vars}/(35.7) = N_{vars}/Mb$.

This revealed that both cancer types have relatively high mutational burdens calculated from all variants, ranging from 27 to 280 mutations per Mb (Mt/Mb) with similar median mutational burdens of 109 Mt/Mb for LUAD and 104 Mt/Mb LUSC, but a broader range for LUAD (Figure 2 A, Supplementary Data S1). In terms of missense variants alone, this scales as ranging from 5 to 43 Mt/Mb and medians of 20 Mt/Mb for LUAD and 16 Mt/Mb LUSC

Approximately one third of all nucleotide transitions and transversions were C>A transversions in both LUSC and LUAD (Figure 2 B), a known mutational signature of smoking[22]. Of the 51,810 LUAD and 32,344 LUSC single nucleotide variants, approximately 20% were missense variants (10,565 LUAD, 6,772). These missense SNVs along with approximately 15% insertion/deletion variants (9,697 LUAD, 6,780 LUSC) predict amino acid changes at the protein level and are therefore potential sources of HLA neoantigens (Figure 2 C).

For each cancer subtype, patterns of single base substitutions created by the somatic mutations were extracted to identify mutational signatures that were fitted to those identified in COSMIC[24-26] (Figure 2 D-F). LUAD signature A fit SBS36 indicating base excision repair deficiency characterised by C>A transversions. LUAD signature C fit SBS2, which is common in lung cancer and thought to indicate APOBEC cytidine deaminase activity as characterised by C>T

transitions. LUSC signature C fit SBS29, another signature characterised by C>A transversions

and linked to tobacco chewing.

In addition to examining the potential for neoantigen generation at the exon level, we sought to

examine the potential for neoantigen recognition within the tumours using bulk gene expression

data from RNAseq to assess the fractions of immune cells present in the tumours[27] (Figure 2

G). This estimation also provides an indication of the tumour sample purity. All expressed genes

not used as markers for immune cells are labelled as 'otherCells' and we would expect this

category to comprise the largest proportion of cells in a tumour sample. Therefore if a sample

has lower proportion of 'otherCells' it is indicative of a less pure tumour sample. For LUAD and

LUSC, the median proportions of 'otherCells' are one third. In cases with very low proportions of

'otherCells' such as A134 and A145, the corresponding histology reports indicate these were

fibrotic samples consistent with the very high proportions of cancer associated fibroblasts

identified by RNAseq. However at the cohort level, proportions of T-cells estimated capable of

responding to neoantigens presented by HLA were estimated with similar medians for CD4+ T-

cells of 18% and 15% for LUAD and LUSC respectively, and medians for CD8+ T-cells of 2%

and less than 1% for LUAD and LUSC respectively.

In summary, the mutational landscape of the NSLC cohort is characterised by a high tumour

mutational burden in both cancer subtypes, the largest proportion of variants with the potential

for generation of neoantigens arising from C>A transversions. Furthermore, gene expression

data estimates the presence of limited populations of T-cells with the potential to recognise HLA

presented neoantigens.

Figure 2: The mutational landscape of lung cancer in the studied cohort. (A) The mutational burden of each cancer type: squamous (n=9) and adenocarcinoma (n=15). (B) Mutation frequency of six transition and transversion categories for each cancer type. (C) Mutation frequencies each cancer type. (D-F) Mutational signatures identified in each cancer subtype. (G) The proportions of immune cells estimated from bulk tumor RNASeq in each tumour sample.

The peptidome landscape of NSCLC in the studied cohort

Mass spectrometry proteomics of the HLA immunopeptidomes identified large distributions of peptides with their characteristic modes of 9 amino acids (AA) and 15 AA for class I and II HLA peptides respectively (Figure 3 A). Median class I immunopeptidome sizes were 5422 and 2998 for Adenocarcinoma and Squamous NSCLC respectively. Median class II immunopeptidome sizes were 2849 and 1125 for Adenocarcinoma and Squamous NSCLC respectively.

The lung cancer peptidome resembles the healthy lung tissue peptidome

We compared the distinct source protein populations yielding the class I and II HLA peptidomes between our LUAD, LUSC samples and healthy lung tissues from the Human HLA Ligand Atlas[10] to examine their similarities and differences (Figure 3 B-C, Supplementary Data S5) , considering only proteins present in at least two-thirds of our samples peptidomes. Our analysis suggests that healthy lung and tumour tissues immunopeptidomes sample largely the same protein populations. 92% of HLA-I proteins and 52% of HLA-II proteins were common to all three tissue types. The remaining proteins most likely represent experimental variation.

187   **Direct identification of a neoantigen in a LUAD patient immunopeptidome**

Across the cohort of 24 patients we identified a single missense variant product by direct mass

spectrometric observation in the class I HLA immunopeptidome of one LUAD patient (A147)

(Figure S1-S3). This derived from a C>A variant in the ALYREF gene yielding an Asp10Tyr

mutation in its protein product THO complex subunit 4 (Uniprot: Q86V81). This mutation yielded

seven nested 15-18mer peptides with the mutation Y before the start of core sequence of

MSLDDIIKL. No wild type peptides were observed for this protein in either the HLA I or II

immunopeptidomes, suggesting this mutation altered either the binding affinity of these peptides

or the source protein processing in the antigen processing pathway. The rarity of this

observation is in keeping with estimates of frequencies in the order of 0.5% of missense

variants encoding presented neoantigens[7,9]. The length of the ALYREF peptides suggested

these may be class II HLA peptides that we had captured by chance in this assay. The motif

most closely matched the patients HLA-DRB1*03:01 allotype with peptide AYKMDMSLDDIIKLN

predicted as a weakly binding peptide[28]. We identified 1135 missense mutations for patient

A147 (Supplementary Data S1). An estimate of 0.5% missense yielding neoantigens would

represent 6 neoantigens. Here we observed one.

Figure 3: The peptidome landscape of lung cancer. (A) Length distributions of

immunopeptides from tumour tissues. (B-C) Upset plots of proteins presented by HLA molecules (class I left, class II right) comparing proteins between cancer subtypes and healthy lung tissues from the HLA ligand atlas. Proteins were included if observed in at least two-thirds of the peptidomes for each cancer subtype.

A proteogenomics view of NSCLC in the studied cohort

Consistent with the view that non-silent mutations rarely encode HLA presented neoantigens[9], we observed that LUAD and LUSC driver genes[29] are not mutated and presented by HLA molecules with the same frequencies. Some drivers are frequently mutated, but rarely presented e.g. APC, whereas others are rarely mutated, but frequently present in the HLA immunopeptidomes e.g. KEAP1 (Figure 4 A). TP53 is both frequently mutated and presented in the class I HLA peptidomes of both NSCLC subtypes (Figure 4 A).

We found that loss of class I HLA heterozygosity in the genome[30] is reflected in the peptidome. In heterozygous patients, immunopeptides identified as presented by HLA molecules from the retained allele were observed at higher proportions in the peptidome than from the lost allele for HLA-A and B allotypes (Figure 4 B, Supplementary Data S4).

We also found that mutations are distributed across the cellular compartments at the same frequencies as the genes are expressed (Figure 4 C left), but the HLA pathways sample the compartments preferentially. Class I HLA immunopeptides are derived preferentially from nuclear and cytosolic proteins, whilst class II HLA immunopeptides are derived preferentially from membrane and extracellular proteins (Figure 4 C right).

Collectively, these observations imply that the likelihood of a putative neoantigen being presented by either HLA class is influenced by the cellular compartment origin of the source protein and secondly, putative neoantigens with motifs[31] for the retained class I HLA allotypes are more likely to be presented than those from the lost allotype.

Figure 4: Integrating the mutational and immunopeptidome landscape reveals previously unclear relationships between mutations and peptide presentation. (A) Parity plots of the frequency of mutated initiating driver genes for each cancer type plotted against the

frequency of observed presentation from the corresponding protein in either the HLA-I (left) or HLA-II peptidome (right). Parity of frequency is indicated by the dotted line. The colour indicates which cancer type the gene was identified as an initiating driver in[5]. (B) The relative proportions of immunopeptides assigned to each of the two HLA-A, B and C allotypes for heterozygous patients.[7,31,32]. The colour represents whether an allele is predicted to have loss of heterozygosity in the genome[30]. (C) Comparison of the frequency between the cellular compartments in which gene expression and somatic mutations occur, and those from which HLA peptides are observed in each cancer type.

Proteogenomics guided NSCLC neoantigen selection and testing

In selecting neoantigens we initially used the pVACseq tool to create a list of putative neoantigens for each patient and HLA allotype and different peptide lengths[33]. Briefly, we used pVACseq with the whole exome and transcriptome outputs and patient HLA allotypes to predict 8-11mer peptides for class I HLA and 15-mer peptides for class II HLA-DRB allotypes across eight binding algorithms. This combined genomic and binding score creates an overall score for each peptide (Details in Section 1.8.0.10). For our 24 patients this comprises 524 HLA class I tables and 74 HLA class II tables of ranked predictions. Discarding any prediction for a peptide with >500 nM binding affinity, pVACseq yielded 27,466 class I HLA and 127,015 class II HLA predicted neoantigen peptides (Supplementary Data S6-S8)

We were able to test predictions for six patients, but this still required selecting from thousands of possible candidate peptides. We consequently filtered the candidate peptides according to whether peptides arising from the gene product with a missense variant were already present in the patients' respective class I or class II HLA peptidome (Supplementary Data S3) and

according to HLA peptide length preferences[34]. This reduced the number of candidates to a few

hundred peptides for each patient. We finally manually curated the ranked peptide candidates

for biological relevance using auxiliary information from the literature, the Human Protein Atlas

and COSMIC. (Figure 1, Section 1.8.0.10).

Our exploratory filtering process for candidate neoantigens can be summarised as: Does a

missense mutation exist? Is there evidence that the mutated gene product enters the antigen

processing pathway for presentation, and if so in which HLA pathway? Is the candidate

neoantigen of the preferred HLA allotype length? Is the candidate neoantigen predicted to bind

to the HLA allotype according to pVACseq? To preferentially select one neoantigen candidate

over another in the final ranking we looked for evidence of biological relevance for the candidate

in COSMIC[26] and the HPA[35].

As HLA peptidome observation took precedence over pVACseq rank, some candidates such as

peptide 08-FAT1 were low ranking (70th percentile) but still with a predicted binding affinity

lower than 500 nM (Table 2).

For six patients, 3 LUAD and 3 LUSC, we selected 9 to 14 putative neoantigens per patient (70

in total) and synthesised the specific putative HLA-I or HLA-II peptides in the mutant neoantigen

and wildtype forms (Supplementary Data S9). Using their donor matched PBMCs, we identified

nine strong neoantigen specific responses to putative neoantigens in five out of six patients,

including for the directly observed ALYREF peptide (Figure 5 A-F, Table 2). This represents an

83% success in achieving a donor functional response and a 13% response rate in antigenic

peptide selection, twice the genomics-based peptide prediction rate of 6% reported in the

literature[15]. We observed responses to both class I and class II HLA candidate neoantigens in

LUAD (Figure 5 A-C), but only class II HLA candidate neoantigens in LUSC (Figure 5 E-F).

LUSC patient A116 yielded no responses (Figure 5 D).

Table 2: Peptides yielding IFN-γ ELISPOT responses

Tissue	ID / HLA / Peptide Length[a]	Peptide[b]	Rank %[c]	Peptidome support[d]	Auxiliary support[e]	ELISpot response
LUAD	A119 / DRB1*04:04 / 15	01-CANT1	11	II	HPA	Strong
LUAD	A119 / HLA-A*31:01 / 10	12-PTPRT	10	I	COSMIC	Strong
LUAD	A147 / Observed / 15	01-ALYREF	-	I	-	Strong
LUAD	A147 / DRB1*04:01 / 15	08-FAT1	70	I+II	COSMIC	Strong
LUAD	A147 / HLA-A*01:01 / 9	14-TP53	7	I	COSMIC	Weak
LUAD	A148 / HLA-A*26:01 / 9	01-KMT2C	20	I	COSMIC	Strong
LUAD	A148 / DRB1*01:01 / 15	05-NT5E	6	I+II	HPA	Strong
LUSC	A134 / DRB1*01:03 / 15	06-KRT8	10	I+II	-	Strong
LUSC	A144 / DRB1*04:01 / 15	04-FAT1	24	I+II	COSMIC	Strong
LUSC	A144 / DRB1*04:01 / 15	08-NF1	3	I+II	COSMIC	Strong

[a]The patient ID, predicted HLA allotype for the peptide and peptide length. ALYREF was an observed peptide.

[b]The peptide identifier and gene name corresponding with those in Figure 5.

[c]Rank % is the rank of the peptide in the table for that Donor and HLA allotype, a lower rank corresponds with better pVACseq score as detailed in the materials and methods.

[d]Peptidome support indicates from which class HLA peptidome source protein peptides were observed.

[e]Auxiliary support indicates support for biological relevance either from the lung cancer associated proteins in the Human Protein Atlas (HPA) or the Top 20 mutated genes in COSMIC.

Figure 5: Proteogenomics guided NSCLC neoantigen selection identifies nine strong candidates (A-F) IFN-γ ELISPOT of putative neoantigens three LUAD and three LUSC patients donor matched PBMCs. Wildtype peptides are represented by blue bars, putative HLA-I neoantigens by red bars, putative HLA-II neoantigens by orange bars, the observed ALYREF neoantigen in purple and the CEFT control peptide mix in grey.

Model of peptide presentation

To retrospectively examine the mechanisms underlying our rules based selection approach and quantify the relative importance of factors affecting peptide presentation, we created Bayesian multilevel regression models of the NSCLC immunopeptidomes[36]. The goal was to understand the utility of immunopeptidomes in the context of neoantigen selection by quantifying what determines observed peptide presentation.

Several features have been identified as causal: protein abundance, protein hotspots, peptide

length preferences of HLA molecules, peptide binding motifs and other features of antigen

processing proteins such as tapasin, TAPBR and TAP on HLA-I peptide selection[37–41].

Our findings indicated that driver genes do not correlate with peptide presentation (Figure 4 A)

and that the presented proteome of LUAD, LUSC and healthy lung tissues are highly similar

(Figure 3 B-C). Additionally, as others have also found, direct observations of neoantigens are

rare[7,9]. Although not exclusively, we found that proteins were preferentially presented via HLA

pathways corresponding with the cellular compartment they occupy (Figure 4 C).

To construct the models we began with the source proteins, which we did not observe directly.

Proteins from the underlying proteome were latent variables from which peptides derived, bound

to and were presented by HLA molecules. Proteins were represented in the form of peptide

counts for each protein observed, for each HLA allotype and donor. We assumed that there

were relationships between donor or NSCLC subtype cohort and protein properties

e.g. abundance, with other peptides on binding to HLA and presentation, represented as a

causal model by a directed acyclic graph (DAG) (Figure 6 A). Arrows indicated the direction of

the causal effects the model assumed. The DAG indicated several sources of confounding,

specifically the need for the model to adjust during fitting for protein, donor, cohort, peptide and

HLA specific effects in estimating the probability of each peptide being presented. Within the

limits of detection, a peptide was either presented or not, hence we assumed this outcome

variable (yes or no) as being a Bernoulli distribution arising from a linear combination of our

predictor variables: protein, donor, cohort, peptide and HLA properties. Importantly, having

assumed the outcome distribution, Bayes theorem was used to pool observations across each

subtype cohort and share information to adjust the estimation for each peptide as indicated by

the DAG. Pooled adjustment helped to account for differing immunopeptidome sizes between

donors and also enabled estimation of the probability of presentation of a peptide even if it was

missing in any given immunopeptidome. These models quantified the relative importance of

donor specific versus protein specific factors in determining peptide presentation.

Model diagnostics confirmed robust estimation with high signal-to-noise ratios (SNR > 10),

adequate effective sample sizes, and good MCMC convergence across all models

(Supplementary Tables S1-S2, Figures S4-S11).

To examine the relative importance of donor differences, protein properties and donor-specific

protein preferences, we calculated the contribution to the total variance of each model

component (Figure 6 B). We found moderately greater contributions of donor effects to protein

effects for LUAD HLA-I and HLA-II presentation, ratios of 1.3 and 3 times greater respectively

(Figure 6 C). For the LUSC cohort, donor effects were much larger contributors to presentation

variance than protein effects for both HLA-I and II, ratios of 8 and 22 times respectively

(Figure 6 C). Donor-specific protein contributions varied and were greatest for HLA-II

immunopeptidomes, and was the largest variance component for LUAD HLA-II.

To further quantify these effects, we calculated the average difference in peptide presentation

probability for each donor relative to median donor presentation (Figure 6 D). For LUAD, most

donors showed HLA-I presentation close to the median, but differences in the order of 50% for

some donors with respect to median presentation for HLA-II, albeit with much greater

uncertainty. The model estimated donors such as A148 and A147 were more likely to present

peptides, despite A148 and A147 having only the 6th and 10th highest number of HLA-I and 8th

and 9th number of HLA-II peptide observations respectively (Supplementary Data S1). The

estimates for LUSC donors showed greater variation and uncertainty, with most donors below

the median. Donor A116 had the greatest number of LUSC HLA-I peptide observations, and 3rd

greatest LUSC HLA-II peptide observations (Supplementary Data S1), but the model estimated

A116 below the median for presentation for both HLA classes in LUSC, and we were unable to

select a reactive neoantigen peptide for this donor.

In our rules-based selection of neoantigens we ranked a prediction more highly if a donor

presented peptides from the same protein as the putative neoantigen (Figure 1 B). To quantify

this effect and examine our rules-based assumption, we calculated the average effect on

peptide presentation of observing multiple peptides from the same protein across each NSCLC

subtype cohort versus the effect of observing more than one peptide from the same protein

presented by a donor.

The model identified competing effects such that when a protein presented multiple peptides

across the cohort, this decreased the probability of observing an additional peptide by between

18-23% depending on NSCLC subtype and HLA class (Figure 6 E). In contrast, a donor

presenting one peptide from a protein increased the probability of observing another peptide

from the same protein by between 23-41% depending on NSCLC subtype and HLA class

(Figure 6 E), supporting our rules-based assumption. The model quantified that donor-specific

protein preferences substantially influenced presentation probability, but also revealed that

cohort-level presentation patterns created competing effects.

In summary, by pooling information across immunopeptidomes in the NSCLC cohorts, the

models identified NSCLC subtype variation and donor-specific effects on peptide presentation:

each donor's general propensity to present peptides and their protein-specific preferences.

Figure 6: Bayesian multilevel regression model of HLA peptide presentation (A) Directed

Acyclic Graph of HLA presentation for NSCLC cohorts. Variables adjusted for

confounding in dark blue, unadjusted variables in light blue. (B) Contributions of donor

differences, protein properties and donor specific proteins to the total variance in

peptide presentation in Bayesian multilevel regression models of NSCLC immunopeptidomes. (C) Barplot of the ratio between donor and protein variance components in the peptide presentation models. The red dashed line indicates where these components are of equal value. (D) Comparison of the average propensity of donor to present peptides with respect to median presentation for each NSCLC subtype and HLA class. (E) Comparison of the average effect on the probability of presentation of an additional peptide from the same protein if a donor has already presented a peptide versus multiple peptides from that protein being presented across the cohort for each NSCLC subtype and HLA class.

Discussion

The strategy of using HLA presented peptides as a basis for immunotherapy is long

standing[42,43]. Researchers have sought to identify either peptides common to a cancer type, so-

called tumour associated antigens, or peptides unique to a patient's tumour, so called

neoantigens. Here we sought to improve our ability to identify HLA presented neoantigens in

two NSCLC sub-types using a proteogenomics approach that combines exome sequencing,

transcriptomics, mass spectrometry immunopeptidomics and mechanistic modelling. In tumour

samples from cohort of 24 NSCLC patients we found relatively high mutational burdens with

exonic mutations characterised by a predominance of C>A transversions and containing small

populations of T-cells. Consistent with previous reports[8,9] using mass spectrometry

immunopeptidomics, we only directly identified one neoantigen amongst tens of thousands of

peptide identifications. However, following a rules based approach, we utilised the remaining

observations to inform our selection of neoantigens from ranked lists generated by in silico

prediction algorithms[33] to the extent that we were able to identify positive functional assay

neoantigens for 5 out of the 6 patients (83%) we were able to test. This included a positive

response for the directly observed neoantigen. These findings also represent a two-fold

improvement over previous reports for neoantigen prediction and identification[15,16]. We then

applied Bayeisan inference to retrospectively examine our rules based neoantigen selection and

quantify patient specific differences in peptide presentation propensity.

Although identifying patient specific neoantigens and was the primary goal of our study, the data

included some interesting related observations: With the exception of TP53 and FAT1 in the

HLA-I and HLA-II immunopeptidomes respectively, there was no correlation between driver

gene mutation frequency and their peptide presentation frequency. This provides some

circumstantial support for these two genes as sources for neoantigens[44,45]. TP53 mutations can

be either truncal or late-stage[46], but a third of TP53 mutations occur in so-called hotspot

regions[47], making them of interest both for early detection and as targets for immunotherapy[48,49].

Overall we found that the tumour peptidomes contained peptides derived from the same source

proteins as healthy tissue. Furthermore, although somatic mutations are not random, as seen in

the mutational signatures and driver genes, their distribution amongst cell compartments

corresponds with gene expression frequency. There is no enrichment for mutations in genes

expressing proteins in specific cell compartments. This implies a connection between the cell

compartment from where the source protein derives and the subsequent HLA antigen

processing pathway it primarily feeds, consistent with previous reports[50]. TP53 is a

predominantly a nuclear protein, whilst FAT1 is predominantly extracellular, hence their higher

frequencies in HLA I and II immunopeptidomes respectively. Whilst this might seem tautological,

it does indicate that it would be unwise to preferentially select class I neoantigen predictions for

FAT1 and vice versa for TP53, and yet this is not explicitly considered in existing neoantigen

prediction algorithms. Hence, we selected source protein cell compartment as a relevant

neoantigen parameter in our rules based approach. This choice was supported by the

mechanistic model prediction of an increased probability of observing another peptide from the

same protein if a patient has presented one peptide already by between 23-41%. However, our

mechanistic model also identified competing effects between proteins that were widely

presented in NSCLC subtype and those proteins specific to each donor, highlighting the need

for patient specific predictions that aren't solely reliant on general HLA peptide characteristics.

Loss of class I HLA heterozygosity in the genome was reflected in the proportions of peptides

observed for each HLA-I allotype, and although we did not use this information to select

neoantigens, this might be another useful parameter when ranking candidates.

There are various limitations in our study which might be addressed by future studies. Perhaps

most significantly from a methodological perspective, mass spectrometry as a technique does

not have the amplification step found in many genomic sequencing methodologies. Therefore

the strength of the input signal arises almost entirely from sample quality and preparation and

sensitivity is determined by the mass spectrometer itself. The complexity of the input mixture

and the differential ability of peptides to ionise, along with their relative abundances all affect

what fraction of the immunopeptidome is observed. Various single molecule technologies are

being developed that may address this problem, of which pore-based technologies, possibly in

combination with fluorescence fingerprinting, seem well suited to identification of short

peptides[51–53]. Sequencing peptides using pore technologies offers the tantalising prospect of

providing much greater coverage of the immunopeptidome, and therefore direct observation of

neoantigens. There are many challenges to this approach, not least post-translational

modifications and the non-polar nature of protein peptides, but much progress has already been

made[54–56].

In our study we only considered canonical neoantigens arising from missense variants. This

was a limitation largely arising from choosing whole exome sequencing, but there is increasing

evidence for non-canonical antigens arising from non-coding regions of the genome[57–59]. There

are several sources of non-canonical peptides such as circRNA translation, unannotated open

reading frames (nuORFs), lncRNA and alternative reading frames. As yet, clinical efficacy is

unknown and it is also unclear how abundant non-canonical peptides are in the

immunopeptidome due their precursor instability and/or relative abundance[60]. Likewise,

detection of non-canonical peptides is further hampered by the increase in search space they

represent and subsequent impact on detection sensitivity, though this is being addressed[61].

Immunopeptidomic identification of circRNA and cancer specific nuORFs peptides have been

reported[63] and immune responses in mice to pancreatic cancer nuORF peptides[64]. Therefore

we'd expect inclusion of non-canonical targets to be part of future immunopeptidomic studies.

Here we have identified potential candidates for personalised vaccines that elicit strong positive

responses in functional T-cell assays, however further evaluation is required to determine their

clinical efficacy, and their effectiveness as vaccines cannot be guaranteed. The stage of the

cancer at which the patient receives the vaccine may be crucial for efficacy. Chronic neoantigen

exposure driving T-cells to dysfunctional states, late-differentiated T-cells dominating the tumour

microenvironment, and loss of HLA heterozygosity are all reasons NSCLC may become harder

to treat with neoantigen vaccines at later stages[6,65–67]. Heterogeneity in NSCLC tumours is likely

to influence the efficacy of neoantigen based vaccines[29].

Personalised neoantigen vaccines are already being trialled for the treatment of melanoma,

glioblastoma and pancreatic cancer[68–70]. These trials rely on the delivery of mRNA containing a

number of long sequences predicted to be processed into the final HLA presented neoantigens.

The vaccine response rate is in currently the order of 50% of patients when multivalent epitope

vaccines in combination with an checkpoint inhibitor adjuvant are used[71]. So whilst these results

are extremely promising, there is clearly room for improvement, including in the neoantigen

selection process. Immunogenic peptides are predicted by algorithms that incorporate machine

learnt parameters such as peptide binding affinity[72] or proteasomal cleavage[73], or more recently

using machine learning to identify features such as protein hotspots from large mass

spectrometry immunopeptidomics datasets[17,18].

The principal difference in our approach is one of tactics rather than strategy. Our tactical

difference being to look at which proteins yield peptides presented by HLA molecules and then

manually identifying supporting evidence for each neoantigen candidate protein in the literature.

To understand the mechanistic basis for this approach, we retrospectively applied Bayesian

inference to quantify how donor-specific and protein-specific factors contribute to peptide

presentation. This analysis validated our empirical observation that donor-presented source

proteins are informative for candidate ranking, while revealing competing cohort-level effects

that could inform parameter weighting in future automated selection algorithms. This tactic has

some similarity to the 'Tübingen approach' for identification of tumour associated neoantigens

which uses mass spectrometry proteomics identifications of HLA peptides to rank candidates[74],

as used in the glioblastoma vaccine[69].

Related immunopeptidomic-guided approaches have been reported. Tokita et al. used pooled

wildtype immunopeptidomes as "surrogate" immunopeptidomes to filter neoantigen candidates

against exonic mutations, successfully identifying two neoantigens in colorectal cancer[75]. A

large-scale immunopeptide-guided selection study used machine learning to analyse

approximately 0.3 million peptides[19]. In concordance with estimates that only 0.5% of mutations

yield presented neoantigen peptides[9], they predicted 0.7% of peptides as immunogenic (2,523

peptides total: 1,476 canonical and 1,047 noncanonical), with three canonical peptides

validated. Our approach differs in using patient-specific tumour immunopeptidomes rather than

surrogate immunopeptidomes, systematically testing multiple candidates per patient (70

peptides across 6 patients) and using mechanistic modelling to identify relevant parameters for

machine learning models.

Although still far from perfect with 87% of our predictions failing, our approach outperformed

current machine learning models and, in the tested patients, identified peptides that elicited a

strong functional response in ~83% of cases tested. Perhaps most importantly, our findings

indicate that patient specific knowledge beyond their mutational differences, incorporating

knowledge of each individual's HLA-presented peptides are critical parameters in improving

neoantigen selection workflows and precision therapies.

Materials and Methods

Ethics statement

This study was performed in accordance with the Declaration of Helsinki. Ethical approval was obtained from the local research ethics committee (LREC reference 14-SC-0186 150975) and written informed consent was provided by the patients.

Tissue preparation

Tumours were excised from resected lung tissue post-operatively by pathologists and processed either for histological evaluation of tumour type and stage, or snap frozen at −80°C. Whole blood samples were obtained, and PBMCs were isolated by density gradient centrifugation over Lymphoprep prior to storage at −80°C.

HLA typing

HLA typing was performed by Next Generation Sequencing by the NHS Blood and Transplant Histocompatibility and Immunogenetics Laboratory, Colindale, UK.

DNA and RNA extraction

DNA and RNA were extracted from tumor tissue that had been obtained fresh and immediately snap frozen in liquid nitrogen. Ten to twenty 10 µm cryosections were used for nucleic acid extraction using the automated Maxwell® RSC instrument (Promega) with the appropriate sample kit and according to the manufacturer's instructions: Maxwell RSC Tissue DNA tissue kit and Maxwell RSC simplyRNA tissue kit, respectively. Similarly, DNA was extracted from snap frozen normal adjacent tissue as described above. DNA and RNA were quantified using Qubit fluorometric quantitation assay (ThermoFisher Scientific) according to the manufacturer's

477   instructions. RNA quality was assessed using the Agilent 2100 Bioanalyzer generating an RNA

478   integrity number (RIN; Agilent Technologies UK Ltd.).

Whole exome sequencing

The tumor and normal adjacent samples were prepared using SureSelect Human All Exon V7

library (Agilent, Santa Clara USA). 100 bp paired end reads sequencing was performed using

the Illumina NovaSeq 6000 system by Edinburgh Genomics (Edinburgh, UK) providing ~100X

depth. Reads were aligned to the 1000 genomes project version of the human genome

reference sequence (GRCh38/hg38) using the Burrows-Wheeler Aligner (BWA; version 0.7.17)

using the default parameters with the addition of using soft clipping for supplementary

alignments. Following GATK Best Practices, aligned reads were merged[76], queryname sorted,

de-duplicated and position sorted[77] prior to base quality score recalibration[78].

Tumour purity and ploidy

489   Tumour purity and ploidy were estimated from WES using the ASCAT R package v3.3.0

490   following the recommended steps on the Github repository: https://github.com/VanLoo-

491   lab/ascat[79].

Somatic variant calling

Somatic variant calling was performed using three variant callers: Mutect2 (version 4.1.2.0)[80],

Varscan (version 2.4.3)[81], and Strelka (version 2.9.2)[82]. For Mutect2, a panel of normals was

created using 40 samples (20 male and 20 female) from the GBR dataset. Variants were

combined using gatk GenomeAnalysisTK (version 3.8-1) with a priority order of Mutect2,

Varscan, Strelka. Variants were then left aligned and trimmed, and multi-allelic variants split[83].

Hard filtering of variants was performed such that only variants that had a variant allele fraction

> 5%, a total coverage > 20 and variant allele coverage > 5 were kept. Filtered variants were

annotated using VEP (version 97)[84] and with their read counts (https://github.com/genome/bam-

readcount) to generate the final filtered and annotated variant call files (VCF).

RNA sequencing

Samples were prepared as TruSeq stranded mRNA libraries (Illumina, San Diego, USA) and

100bp paired end sequencing was performed using the Illumina NovaSeq 6000 system by

Edinburgh Genomics (Edinburgh, UK). Raw reads were pre-processed to using fastp (version

0.20.0)[85]. Filtered reads were aligned to the 1000 genomes project version of the human

genome reference sequence (GRCh38/hg38) using hisat2 (version 2.1.0)[86], merged and then

transcripts assembled and gene expression estimated with stringtie2 (version 1.3.5)[87] using

reference guided assembly.

Mutanome generation

511   The annotated and filtered VCFs were processed using Variant Effect Predictor (version 97)[84]

512   plugin ProteinSeqs to derive the amino acid sequences arising from missense mutations for

513   each sample for use in immunopeptide analyses.

Neoantigen prediction

Variant call files were prepared for the pvacseq neoantigen prediction pipeline (version

1.5.10)[21,33] by adding tumor and normal DNA coverage, and tumor transcript and gene

expression estimates using vatools (version 4.1.0) (http://www.vatools.org/). Variant call files of

phased proximal variants were also created for use with the pipeline[88]. Prediction of

neoantigens arising from somatic variants was then performed using pvacseq with the patient

HLA allotypes to predict 8-11mer peptides for class I HLA and 15-mer peptides for class II HLA-

DRB allotypes. Four binding algorithms were used for class I predictions (MHCflurry,

MHCnuggetsI, NetMHC, PickPocket) and four for class II predictions (MHCnuggetsII,

NetMHCIIpan, NNalign, SMMalign). Unfiltered outputs were post-processed in R[89] and split into

individual tables for each peptide length and HLA allotype for each patient, and each table was

then ranked according to the pvacseq score, where:

$$score = binding\ score\ +\ fold\ change\ +\ (variant\ expression\ \times\ fold\ change)$$

$$+\ (tumour\ VAF\,/\,2)$$

Here $binding\ score$ is 1/median neoantigen binding affinity, $fold\ change$ is the difference in

median binding affinity between neoantigen and wildtype peptide (agretopicity).

Each table was then filtered according to whether wildtype peptide(s) from the same protein as

predicted neoantigen was present in the individual's peptidome, and further filtered manually

according to biological relevance e.g. the ontology of the protein and its likely presence in the

relevant HLA pathway, for example a cytoplasmic resident protein would be considered more

likely to yield a HLA-I neoantigen than a HLA-II one. The Human Protein Atlas list of 354 genes

identified for unfavourable prognosis in lung cancer, the COSMIC top 20 mutated genes and

literature searches were also used as a screen for genes/proteins/peptides of biological

relevance.

Immunopeptidomics

Snap frozen tissue samples were briefly thawed and weighed prior to 30s of mechanical

homogenization (Fisher, using disposable probes) in 4 mL lysis buffer (0.02M Tris, 0.5% (w/v)

IGEPAL, 0.25% (w/v) sodium deoxycholate, 0.15mM NaCl, 1mM EDTA, 0.2mM iodoacetamide

supplemented with EDTA-free protease inhibitor mix). Homogenates were clarified for 10 min at

2,000g, 4°C and then for a further 60 min at 13,500g, 4°C. 2 mg of anti-MHC-I mouse

monoclonal antibodies (W6/32) covalently conjugated to Protein A sepharose (Repligen) using

DMP as previously described[90,91] were added to the clarified supernatants and incubated with

constant agitation for 2 h at 4°C. The captured MHC-I/β_2m/immunopeptide complex on the

beads was washed sequentially with 10 column volumes of low (isotonic, 0.15M NaCl) and high

(hypertonic, 0.4M NaCl) TBS washes prior to elution in 10% acetic acid and dried under

vacuum. The MHC-I-depleted lysate was then incubated with anti-MHC-II mouse monoclonal

antibodies (IVA12) and MHC-II bound peptides were captured and eluted in the same

conditions.

Immunopeptides were separated from MHC-I/β_2m or MHC-II heavy chain using offline HPLC on

a C18 reverse phase column, as previously described[90]. Briefly, dried immunoprecipitates were

reconstituted in buffer (1% acetonitrile,0.1% TFA) and applied to a 10cm RP-18e 100-4.6

chromolith column (Merck) using an Ultimate 3000 HPLC equipped with UV monitor.

Immunopeptides were then eluted using a 15 min 0-40% linear acetonitrile gradient at a flow

rate of 1 mL/min. Peptide fractions were eluted and pooled at between 0 and 30% acetonitrile,

and the β_2m and MHC heavy chains eluted at >40% acetonitrile.

HLA peptides were separated by an Ultimate 3000 RSLC nano system (Thermo Scientific)

using a PepMap C18 EASY-Spray LC column, 2 μm particle size, 75 μm x 75 cm column

(Thermo Scientific) in buffer A (0.1% Formic acid) and coupled on-line to an Orbitrap Fusion

Tribrid Mass Spectrometer (Thermo Fisher Scientific,UK) with a nano-electrospray ion source.

Peptides were eluted with a linear gradient of 3%-30% buffer B (Acetonitrile and 0.1% Formic

acid) at a flow rate of 300 nL/min over 110 minutes. Full scans were acquired in the Orbitrap

analyser using the Top Speed data dependent mode, performing a MS scan every 3 second

cycle, followed by higher energy collision-induced dissociation (HCD) MS/MS scans. MS

spectra were acquired at resolution of 120,000 at 300 m/z, RF lens 60% and an automatic gain

control (AGC) ion target value of 4.0e5 for a maximum of 100 ms. MS/MS resolution was 30,000

at 100 m/z. Higher energy collisional dissociation (HCD) fragmentation was induced at an

energy setting of 28 for peptides with a charge state of 2–4, while singly charged peptides were

fragmented at an energy setting of 32 at lower priority. Fragments were analysed in the Orbitrap

at 30,000 resolution. Fragmented m/z values were dynamically excluded for 30 seconds. Data

desposited in PRIDE[92], see Data Availability for details.

Proteomic data analysis

Raw spectrum files were analyzed using Peaks Studio 10.0 build 20190129[93,94] and the data

processed to generate reduced charge state and deisotoped precursor and associated product

ion peak lists which were searched against the UniProt database (20,350 entries, 2020-04-07)

plus the corresponding mutanome for each sample (~1,000-5,000 sequences) and

contaminants list in unspecific digest mode. Parent mass error tolerance was set a 5ppm and

fragment mass error tolerance at 0.03 Da. Variable modifications were set for N-term acetylation

(42.01 Da), methionine oxidation (15.99 Da), carboxyamidomethylation (57.02 Da) of cysteine.

As previously described, carbamidomethylated cysteines were treated as variable modifications

due to the low concentration of 0.2 mM of iodoacetamide used in the lysis buffer to inhibit

cysteine proteases[95]. A maximum of three variable modifications per peptide was set. The false

discovery rate (FDR) was estimated with decoy-fusion database searches[93] and were filtered to

1% FDR. Downstream analysis and data visualizations of the Peaks Studio identifications was

performed in R using associated packages[89,96].

Immunopeptide HLA assignment

589 Identified immunopeptides were assigned to their HLA allotype for each patient using motif

590 deconvolution tools and manual inspection. For class I HLA peptides initial assignment used

591 MixMHCp (version 2.1)[7,31] and for class II HLA peptides initial assignment used MoDec (version

592 1.1)[32]. Downstream analysis and data visualizations was performed in R using associated

593 packages[89,96,97].

594   Synthetic peptides

595   Peptides for functional T-cell assays and spectra validation were synthesised using standard

596   solid phase Fmoc chemistry (Peptide Protein Research Ltd, Fareham, UK).

597   Functional T-cell assay

PBMC (2×10^6 per well) were stimulated in 24-well plates with peptide (individual/pool) plus

recombinant IL-2 (R&D Systems Europe Ltd.) at a final concentration of 5µg/mL and 20IU/mL,

respectively, and incubated at 37°C with 5% CO2; final volume was 2mL. Media containing

additional IL-2 (20IU/mL) was refreshed on days 4, 6, 8 and 11 and on day 13 cells were

harvested. Expanded cells (1×10^5 cell/well) were incubated in triplicate with peptide (individual)

at 5µg/mL final concentration for 22 hours at 37°C in 5% CO2; phytohemagglutinin (PHA;

Sigma-Aldrich Company Ltd.) and CEFT peptide mix (JPT Peptide Technologies GmbH, Berlin,

Germany), a pool of 27 peptides selected from defined HLA Class I- and II-restricted T-cell

epitopes, were used as positive controls. Spot forming cells (SFC) were counted using the AID

ELISpot plate reader system ELR04 and software (AID Autoimmun Diagnostika GmbH) and

positivity calling for ELISpot data used the runDFR(x2) online tool

(http://www.scharp.org/zoe/runDFR/). Downstream analysis and data visualizations was

performed in R using associated packages[89,96].

611   Peptide presentation model

612   Peptide presentation was modelled in R using Bayesian Multilevel Regression Models[36].

613   For all simulations the Monte Carlo Markov Chain settings were:

```
614    # MCMC settings

615   chains = 4,

616   cores = 4,
```

```
617  iter = 4000,

618  warmup = 1000,

619  control = list(

620      adapt_delta = 0.95,

621      max_treedepth = 15)
```

Model quality control was assessed using trace plots to assess convergence and effective

parameter sampling (Figure S4-S7), and caterpillar plots to assess the uncertainty of model

parameters (Figure S8-S11).

HLA Class I model

A peptide is either presented or not, so peptide presentation is 1 or 0. The model is therefore

defined using a Binomial distribution with a logistic link function:

`peptide_presentation ~ Binomial(1, p) logit(p) = η`

The linear predictor η is defined as:

`η = β₀ + β₁·binding_pre_threshold + β₂·binding_post_threshold +`

`β₃·length_diff_from_9 + Σⱼ β₁ⱼ·(binding_pre_threshold × I(hla = j)) + Σⱼ`

`β₂ⱼ·(binding_post_threshold × I(hla = j)) + Σⱼ β₃ⱼ·(length_diff_from_9 ×`

`I(hla = j)) + β₇·z_sqrt_peptides_donor + β₈·z_sqrt_peptides_cohort + u_donor`

`+ u_donor:uniprot_id + u_uniprot_id`

Where:

• `j` is the index for each HLA allotypes (`j = 1, 2, ..., J`)

• `I(hla = j)` equals 1 when the observed peptide has HLA allotype j, 0 otherwise

• `β₁ⱼ, β₂ⱼ, β₃ⱼ` are HLA-specific interaction coefficients

This allows binding and length preferences to vary by HLA allotype.

There are three levels of random effects:

1. Donor-level effects: `u_donor ~ Normal(0, σ²_donor)`

2. Donor-protein interaction effects: `u_donor:protein ~ Normal(0,`

`σ²_donor:protein)`

3. Protein-level effects: `u_protein ~ Normal(0, σ²_protein)`

Partial pooling via these random effects helps to account for donor heterogeneity and any

protein-specific effects.

The variables are:

HLA-I binding scores were estimated using MixMHCpred3.0 and the resulting

`score_best_allele` scores for each peptide were centred and split around 0 to capture strong

and weak binding effects.

• `binding_pre_threshold`: Weak binding peptide score

• `binding_post_threshold`: Strong binding peptide score

• `length_diff_from_9`: Difference in peptide length from 9 amino acids

• `hla`: HLA allotype

• `z_sqrt_peptides_donor`: Standardised square root of number of peptides per protein

observed per donor

• `z_sqrt_peptides_cohort`: Standardised square root of number peptides peptides per

protein observed per cohort

• `donor`: Donor identifier

• `uniprot_id`: UNIPROT protein identifier

Prior Distributions:

• β_0 ~ `Normal(-1, 1)` Intercept

• β_1 ~ `Normal(1, 0.5)` binding_pre_threshold (main effect)

• β_2 ~ `Normal(1, 0.5)` binding_post_threshold (main effect)

• β_3 ~ `Normal(0, 0.5)` length_diff_from_9 (main effect)

• β_{1j} ~ `Normal(0, 0.5)` binding_pre_threshold:hla interactions

• β_{2j} ~ `Normal(0, 0.5)` binding_post_threshold:hla interactions

• β_{3j} ~ `Normal(0, 0.5)` length_diff_from_9:hla interactions

• β_7 ~ `Normal(0.5, 0.8)` z_sqrt_peptides_donor

• β_8 ~ `Normal(-0.6, 0.8)` z_sqrt_peptides_cohort

• `σ_donor` ~ `Exponential(2)` Donor random effect SD

• `σ_donor:uniprot_id` ~ `Exponential(2)` Donor-protein random effect SD

• `σ_uniprot_id` ~ `Exponential(2)` Protein random effect SD

HLA Class II model

As for HLA-I, peptide is either presented or not, so peptide presentation is 1 or 0. The model is

therefore defined using a Binomial distribution with a logistic link function:

`peptide_presentation ~ Binomial(1, p) logit(p) = η`

The linear predictor η is defined as:

`η = β₀ + Σⱼ sⱼ(binding_score_centered) × I(hla = j)       +`

`β₁·length_diff_from_15 + Σⱼ β₁ⱼ·(length_diff_from_15 × I(hla = j))   +`

`β₂·z_sqrt_peptides_donor + β₃·z_sqrt_peptides_cohort  + u_donor +`

`u_donor:uniprot_id + u_uniprot_id`

For HLA-II peptides, MixMHC2pred was used and produces scores as a binding rank and these

were modelled as smooth splines:

Σ_j `s`$_j$`(binding_score_centered)` × `I(hla = j)`

Where:

•   `s`$_j$`(·)` are HLA allotype specific thin plate regression splines

•   `j` indexes the HLA Class II allotypes (`j = 1, 2, ..., J`)

•   `I(hla = j)` equals 1 when the observed peptide has HLA allotype `j`, 0 otherwise

There are three levels of random effects:

1. Donor-level effects: `u_donor ~ Normal(0, σ²_donor)`

2. Donor-protein interaction effects: `u_donor:protein ~ Normal(0,`

`σ²_donor:protein)`

3. Protein-level effects: `u_protein ~ Normal(0, σ²_protein)`

Partial pooling via these random effects helps to account for donor heterogeneity and any

protein-specific effects.

The variables are:

HLA-II binding scores were estimated using MixMHC2pred and the resulting

`percent_rank_best` scores for each peptide were centred:

•   `binding_score_centered`: Centred binding peptide rank score

•   `length_diff_from_15`: Difference in peptide length from 15 amino acids

•   `hla`: HLA allotype

•   `z_sqrt_peptides_donor`: Standardised square root of number of peptides per protein

observed per donor

•   `z_sqrt_peptides_cohort`: Standardised square root of number peptides peptides per

protein observed per cohort

•   `donor`: Donor identifier

•   `uniprot_id`: UNIPROT protein identifier

Prior Distributions:

• β_0 ~ `Normal(-2, 1)` Intercept

• β_1 ~ `Normal(-0.25, 0.5)` length_diff_from_15

• β_{1j} ~ `Normal(0, 0.5)` length_diff_from_15:hla interactions

• β_2 ~ `Normal(0.5, 0.7)` z_sqrt_peptides_donor

• β_3 ~ `Normal(-0.5, 0.7)` z_sqrt_peptides_cohort

• s_j`(binding_score_centered)` ~ `Normal(0, 1)` Smooth spline prior (sds)

•   `σ_donor` ~ `Exponential(2)` Donor random effect SD

•   `σ_donor:uniprot_id` ~ `Exponential(2)` Donor-protein random effect SD

•   `σ_uniprot_id` ~ `Exponential(2)` Protein random effect SD

Author Contributions

Conceptualisation by PS, TE, CO, PJ, BN and AB. Project administration by PS, TE, CO, PJ and KM. Supervision by PS, TE and CO. Methodology, investigation, visualisation and formal analysis performed by BN, AB, KM, OW and EC. AB, BN, PJ, TE and CO wrote the manuscript. All authors reviewed the manuscript.

Funding

This study was funded by a Cancer Research UK Centres Network Accelerator Award Grant (A21998). Instrumentation in the Centre for Proteomic Research is funded by the Biotechnology and Biological Sciences Research Council, Grant/Award Number: BM/M012387/1. The funders played no role in study design, data collection, analysis and interpretation of data, or the writing of this manuscript.

Competing Interests

All authors declare no financial or non-financial competing interests.

Data availability

Whole Exome Sequencing and RNAseq data have been deposited at the European Genome-phenome Archive (EGA) under EGA Study ID: EGAS00001005499

The mass spectrometry proteomics data have been deposited to the ProteomeXchange Consortium via the PRIDE partner repository with the dataset identifier PXD028990 and 10.6019/PXD028990.

739 Supplementary Data are available on Github at https://github.com/ab604/lung-neoantigen-

740 supplement and Zenodo https://zenodo.org/doi/10.5281/zenodo.12820423

References

1. Cancer survival in england - office for national statistics.

2. Cancer Survival in England, cancers diagnosed 2016 to 2020, followed up to 2021.

3. Gainor, J. F. *et al.* EGFR Mutations and ALK Rearrangements Are Associated with Low

Response Rates to PD-1 Pathway Blockade in NonSmall Cell Lung Cancer: A Retrospective

Analysis. *Clinical Cancer Research* **22**, 4585–4593 (2016).

4. Mazieres, J. *et al.* Efficacy of immune-checkpoint inhibitors (ICI) in non-small cell lung

cancer (NSCLC) patients harboring activating molecular alterations (ImmunoTarget). *Journal of*

Clinical Oncology **36**, 9010–9010 (2018).

5. Jamal-Hanjani, M. *et al.* Tracking the Evolution of Non–Small-Cell Lung Cancer. *New*

England Journal of Medicine **376**, 2109–2121 (2017).

6. Neoantigen-directed immune escape in lung cancer evolution | nature.

7. Bassani-Sternberg, M. *et al.* Direct identification of clinically relevant neoepitopes

presented on native human melanoma tissue by mass spectrometry. *Nature Communications* **7**,

13404 (2016).

8. Nicholas, B. *et al.* Identification of neoantigens in oesophageal adenocarcinoma.

Immunology **168**, 420–431 (2023).

9. Newey, A. *et al.* Immunopeptidomics of colorectal cancer organoids reveals a sparse

HLA class I neoantigen landscape and no increase in neoantigens with interferon or MEK-

inhibitor treatment. *Journal for ImmunoTherapy of Cancer* **7**, 309 (2019).

10.   Marcu, A. *et al.* The HLA Ligand Atlas - A resource of natural HLA ligands presented on

benign tissues. *bioRxiv* 778944 (2020) doi:10.1101/778944.

11.   Andreatta, M. & Nielsen, M. Gapped sequence alignment using artificial neural networks:

Application to the MHC class i system. *Bioinformatics* **32**, 511–7 (2016).

12.   Juncker, A. S. *et al.* Systematic Characterisation of Cellular Localisation and Expression

Profiles of Proteins Containing MHC Ligands. *PLoS ONE* **4**, e7448 (2009).

13.   Müller, M., Gfeller, D., Coukos, G. & Bassani-Sternberg, M. 'Hotspots' of antigen

presentation revealed by human leukocyte antigen ligandomics for neoantigen prioritization.

Frontiers in Immunology **8**, (2017).

14.   Gfeller, D., Liu, Y. & Racle, J. Contemplating immunopeptidomes to better predict them.

Seminars in Immunology **66**, 101708 (2023).

15.   Wells, D. K. *et al.* Key Parameters of Tumor Epitope Immunogenicity Revealed Through

a Consortium Approach Improve Neoantigen Prediction. *Cell* **183**, 818–834.e13 (2020).

16.   Buckley, P. R. *et al.* Evaluating performance of existing computational models in

predicting CD8+ t cell pathogenic epitopes and cancer neoantigens. *Briefings in Bioinformatics*

23, bbac141 (2022).

17.   Gartner, J. J. *et al.* A machine learning model for ranking candidate HLA class I

neoantigens based on known neoepitopes from multiple human tumor types. *Nature Cancer* **2**,

563–574 (2021).

18.   Müller, M. *et al.* Machine learning methods and harmonized datasets improve

immunogenic neoantigen prediction. *Immunity* **56**, 2650–2663.e6 (2023).

19.   Cai, Y. *et al.* Immunopeptidomics-guided discovery and characterization of neoantigens

for personalized cancer immunotherapy. *Science Advances* **11**, (2025).

20.   Huber, F. *et al.* A comprehensive proteogenomic pipeline for neoantigen discovery to

advance personalized cancer immunotherapy. *Nature Biotechnology* **43**, 1360–1372 (2024).

21.   Hundal, J. *et al.* pVACtools: A Computational Toolkit to Identify and Visualize Cancer

Neoantigens. *Cancer Immunology Research* **8**, 409–420 (2020).

22.   Alexandrov, L. B. *et al.* Signatures of mutational processes in human cancer. *Nature*

500, 415–421 (2013).

23.   Kandoth, C. *et al.* Mutational landscape and significance across 12 major cancer types.

Nature **502**, 333–339 (2013).

24.   Gori, K. & Baez-Ortega, A. *Sigfit: Flexible Bayesian Inference of Mutational Signatures*.

372896 https://www.biorxiv.org/content/10.1101/372896v2 (2020) doi:10.1101/372896.

25.   Alexandrov, L. B. *et al.* The repertoire of mutational signatures in human cancer. *Nature*

578, 94–101 (2020).

26.   Tate, J. G. *et al.* COSMIC: The catalogue of somatic mutations in cancer. *Nucleic Acids*

Research **47**, D941–D947 (2019).

27.   Racle, J., Jonge, K. de, Baumgaertner, P., Speiser, D. E. & Gfeller, D. Simultaneous

enumeration of cancer and immune cell types from bulk tumor gene expression data. *eLife* **6**,

e26476 (2017).

28.   Nilsson, J. B. *et al.* Accurate prediction of HLA class II antigen presentation across all

loci using tailored data acquisition and refined machine learning. *Science Advances* **9**, (2023).

29.   Jamal-Hanjani, M. *et al.* Tracking the evolution of nonsmall-cell lung cancer. *New*

England Journal of Medicine **376**, 2109–2121 (2017).

30.   McGranahan, N. *et al.* Allele-Specific HLA Loss and Immune Escape in Lung Cancer

Evolution. *Cell* **171**, 1259–1271.e11 (2017).

31.   Gfeller, D. *et al.* The Length Distribution and Multiple Specificity of Naturally Presented

HLA-I Ligands. *The Journal of Immunology* (2018) doi:10.4049/jimmunol.1800914.

32.   Racle, J. *et al.* Robust prediction of HLA class II epitopes by deep motif deconvolution of

immunopeptidomes. *Nature Biotechnology* **37**, 1283–1286 (2019).

33.   Hundal, J. *et al.* pVAC-seq: A genome-guided in silico approach to identifying tumor

neoantigens. *Genome Medicine* **8**, 11 (2016).

34.   Gfeller, D. *et al.* The Length Distribution and Multiple Specificity of Naturally Presented

HLA-I Ligands. *The Journal of Immunology* (2018) doi:10.4049/jimmunol.1800914.

35.   Uhlén, M. *et al.* Tissue-based map of the human proteome. *Science* **347**, (2015).

36.   Bürkner, P.-C. **brms**: An *R* Package for Bayesian Multilevel Models Using *Stan*. *Journal*

of Statistical Software **80**, (2017).

37.   Howarth, M., Williams, A., Tolstrup, A. B. & Elliott, T. Tapasin enhances MHC class I

peptide presentation according to peptide half-life. *Proceedings of the National Academy of*

Sciences **101**, 11737–11742 (2004).

38.   Boyle, L. H. *et al.* Tapasin-related protein TAPBPR is an additional component of the

MHC class I presentation pathway. *Proceedings of the National Academy of Sciences* **110**,

3465–3470 (2013).

39.   Müller, M., Gfeller, D., Coukos, G. & Bassani-Sternberg, M. 'Hotspots' of antigen

presentation revealed by human leukocyte antigen ligandomics for neoantigen prioritization.

Frontiers in Immunology **8**, (2017).

40.   Bassani-Sternberg, M., Pletscher-Frankild, S., Jensen, L. J. & Mann, M. Mass

Spectrometry of Human Leukocyte Antigen Class I Peptidomes Reveals Strong Effects of

Protein Abundance and Turnover on Antigen Presentation. *Molecular & Cellular Proteomics* **14**,

658–673 (2015).

41.   Gfeller, D. *et al.* The Length Distribution and Multiple Specificity of Naturally Presented

HLA-I Ligands. *The Journal of Immunology* (2018) doi:10.4049/jimmunol.1800914.

42.   Rosenberg, S. A. *et al.* Immunologic and therapeutic evaluation of a synthetic peptide

vaccine for the treatment of patients with metastatic melanoma. *Nature Medicine* **4**, 321–327

(1998).

43.   Weinschenk, T. *et al.* Integrated functional genomics approach for the design of patient-

individual antitumor Vaccines1. *Cancer Research* **62**, 5818–5827 (2002).

44.   Hsiue, E. H.-C. *et al.* Targeting a neoantigen derived from a common *TP53* mutation.

Science **371**, (2021).

45.   Castle, J. C. *et al.* Exploiting the mutanome for tumor vaccination. *Cancer Research* **72**,

1081–1091 (2012).

46.   Levine, A. J., Jenkins, N. A. & Copeland, N. G. The Roles of Initiating Truncal Mutations

in Human Cancers: The Order of Mutations and Tumor Cell Type Matters. *Cancer Cell* **35**, 10–

15 (2019).

47.   Hollstein, M., Sidransky, D., Vogelstein, B. & Harris, C. C. p53 Mutations in Human

Cancers. *Science* **253**, 49–53 (1991).

48.   Lin, M. J. *et al.* Cancer vaccines: the next immunotherapy frontier. *Nature Cancer* **3**,

911–926 (2022).

49.   Vadakekolathu, J. *et al.* Multi-Omic Analysis of Two Common P53 Mutations: Proteins

Regulated by Mutated P53 as Potential Targets for Immunotherapy. *Cancers* **14**, 3975 (2022).

50.   Pearson, H. *et al.* MHC class Iassociated peptides derive from selective regions of the

human genome. *Journal of Clinical Investigation* **126**, 4690–4701 (2016).

51.   Restrepo-Pérez, L., Joo, C. & Dekker, C. Paving the way to single-molecule protein

sequencing. *Nature Nanotechnology* **13**, 786–796 (2018).

52.   Alfaro, J. A. *et al.* The emerging landscape of single-molecule protein sequencing

technologies. *Nature Methods* **18**, 604–617 (2021).

53.   Lucas, F. L. R., Versloot, R. C. A., Yakovlieva, L., Walvoort, M. T. C. & Maglia, G.

Protein identification by nanopore peptide profiling. *Nature Communications* **12**, 5795 (2021).

54.   Motone, K. & Nivala, J. Not if but when nanopore protein sequencing meets single-cell

proteomics. *Nature Methods* **20**, 336–338 (2023).

55.   Martin-Baniandres, P. *et al.* Enzyme-less nanopore detection of post-translational

modifications within long polypeptides. *Nature Nanotechnology* **18**, 1335–1340 (2023).

56.   Motone, K. *et al.* Multi-pass, single-molecule nanopore reading of long protein strands

with single-amino acid sensitivity. (2023).

57.   Laumont, C. M. *et al.* Noncoding regions are the main source of targetable tumor-

specific antigens. *Science Translational Medicine* **10**, (2018).

58.   Chong, C. *et al.* Integrated proteogenomic deep sequencing and analytics accurately

identify non-canonical peptides in tumor immunopeptidomes. *Nature Communications* **11**,

(2020).

59.   Ruiz Cuevas, M. V. *et al.* Most non-canonical proteins uniquely populate the proteome or

immunopeptidome. *Cell Reports* **34**, 108815 (2021).

60.   Jeck, W. R. *et al.* Circular RNAs are abundant, conserved, and associated with ALU

repeats. *RNA* **19**, 141–157 (2012).

61.   Horokhovskyi, Y. *et al.* An Automated Workflow to Address Proteome Complexity and

the Large Search Space Problem in Proteomics and HLA-I Immunopeptidomics. *Molecular &*

Cellular Proteomics **24**, 101039 (2025).

62.   Ferreira, H. J. *et al.* Immunopeptidomics-based identification of naturally presented non-

canonical circRNA-derived peptides. *Nature Communications* **15**, (2024).

63.   Ouspenskaia, T. *et al.* Unannotated proteins expand the MHC-I-restricted

immunopeptidome in cancer. *Nature Biotechnology* **40**, 209–217 (2021).

64.   Ely, Z. A. *et al.* Pancreatic cancerrestricted cryptic antigens are targets for T cell

recognition. *Science* **388**, (2025).

65.   Rooney, M. S., Shukla, S. A., Wu, C. J., Getz, G. & Hacohen, N. Molecular and genetic

properties of tumors associated with local immune cytolytic activity. *Cell* **160**, 48–61 (2015).

66.   Joshi, K. *et al.* Spatial heterogeneity of the T cell receptor repertoire reflects the

mutational landscape in lung cancer. *Nature Medicine* **25**, 1549–1559 (2019).

67.   Ghorani, E. *et al.* The T cell differentiation landscape is shaped by tumour mutations in

lung cancer. *Nature Cancer* **1**, 546–561 (2020).

68.   Sahin, U. *et al.* Personalized RNA mutanome vaccines mobilize poly-specific therapeutic

immunity against cancer. *Nature* **547**, 222–226 (2017).

69.   Hilf, N. *et al.* Actively personalized vaccination trial for newly diagnosed glioblastoma.

Nature **565**, 240–245 (2019).

70.   Rojas, L. A. *et al.* Personalized RNA neoantigen vaccines stimulate T cells in pancreatic

cancer. *Nature* **618**, 144–150 (2023).

71.   Sethna, Z. *et al.* RNA neoantigen vaccines prime long-lived CD8+ T cells in pancreatic

cancer. *Nature* **639**, 1042–1051 (2025).

72.   Andreatta, M. & Nielsen, M. Gapped sequence alignment using artificial neural networks:

Application to the MHC class i system. *Bioinformatics* **32**, 511–7 (2016).

73.   Nielsen, M., Lundegaard, C., Lund, O. & Keşmir, C. The role of the proteasome in

generating cytotoxic T-cell epitopes: insights obtained from improved predictions of proteasomal

cleavage. *Immunogenetics* **57**, 33–41 (2005).

74.   Singh-Jasuja, H., Emmerich, N. P. N. & Rammensee, H.-G. The Tübingen approach:

identification, selection, and validation of tumor-associated HLA peptides for cancer therapy.

Cancer Immunology, Immunotherapy **53**, 187–195 (2004).

75.   Tokita, S. *et al.* Identification of immunogenic HLA class I and II neoantigens using

surrogate immunopeptidomes. *Science Advances* **10**, (2024).

76.   Danecek, P. *et al.* Twelve years of SAMtools and BCFtools. *GigaScience* **10**, (2021).

77.   *Picard Toolkit*. (Broad Institute, 2019).

78.   Genomics in the Cloud [Book].

79.   Ross, E. M., Haase, K., Van Loo, P. & Markowetz, F. Allele-specific multi-sample copy

number segmentation in ASCAT. *Bioinformatics* **37**, 1909–1911 (2021).

80.   Benjamin, D. *et al.* Calling Somatic SNVs and Indels with Mutect2. *bioRxiv* 861054

(2019) doi:10.1101/861054.

81.   Koboldt, D. C. *et al.* VarScan 2: Somatic mutation and copy number alteration discovery

in cancer by exome sequencing. *Genome Research* **22**, 568–576 (2012).

82.   Kim, S. *et al.* Strelka2: fast and accurate calling of germline and somatic variants. *Nature*

Methods **15**, 591–594 (2018).

83.   Bonfield, J. K. *et al.* HTSlib: C library for reading/writing high-throughput sequencing

data. *GigaScience* **10**, (2021).

84.   McLaren, W. *et al.* The ensembl variant effect predictor. *Genome Biology* **17**, 122

(2016).

85.   Chen, S., Zhou, Y., Chen, Y. & Gu, J. Fastp: An ultra-fast all-in-one FASTQ

preprocessor. *Bioinformatics* **34**, i884–i890 (2018).

86.   Kim, D., Paggi, J. M., Park, C., Bennett, C. & Salzberg, S. L. Graph-based genome

alignment and genotyping with HISAT2 and HISAT-genotype. *Nature Biotechnology* **37**, 907–

915 (2019).

87.   Kovaka, S. *et al.* Transcriptome assembly from long-read RNA-seq alignments with

StringTie2. *Genome Biology* **20**, 278 (2019).

88.   Hundal, J. *et al.* Accounting for proximal variants improves neoantigen prediction. *Nature*

Genetics **51**, 175–179 (2019).

89.   R Core Team. *R: A Language and Environment for Statistical Computing*. (R Foundation

for Statistical Computing, Vienna, Austria, 2018).

90.   Purcell, A. W., Ramarathinam, S. H. & Ternette, N. Mass spectrometrybased

identification of MHC-bound peptides for immunopeptidomics. *Nature Protocols* **14**, 1687–1707

(2019).

91.   Bailey, A. *et al.* Characterization of the class i MHC peptidome resulting from DNCB

exposure of HaCaT cells. *Toxicological Sciences* (2020) doi:10.1093/toxsci/kfaa184.

92.   Perez-Riverol, Y. *et al.* The PRIDE database resources in 2022: a hub for mass

spectrometry-based proteomics evidences. *Nucleic Acids Research* **50**, D543–D552 (2021).

93.   Zhang, J. *et al.* PEAKS DB: De novo sequencing assisted database search for sensitive

and accurate peptide identification. *Molecular & Cellular Proteomics* **11**, M111010587 (2012).

94.   Tran, N. H., Zhang, X., Xin, L., Shan, B. & Li, M. De novo peptide sequencing by deep

learning. *Proceedings of the National Academy of Sciences of the United States of America*

(2017) doi:10.1073/pnas.1705691114.

95.   Chong, C. *et al.* High-throughput and Sensitive Immunopeptidomics Platform Reveals

Profound Interferony-Mediated Remodeling of the Human Leukocyte Antigen (HLA)

Ligandome*. *Molecular & Cellular Proteomics* **17**, 533–548 (2018).

96.   Wickham, H. *et al.* Welcome to the tidyverse. *Journal of Open Source Software* **4**, 1686

(2019).

97.   Jessen, L. E. *PepTools - an r-Package for Making Immunoinformatics Accessible*.

(2018).