Ancestral SARS-CoV-2 immune imprinting persists on RBD but not NTD after

sequential Omicron infections

Zuowei Wang[1,2], Ling Li[1,2], Ruiping Du[1], Xixian Chen[1], Yi Sun[1], Rongrong Qin[1], Yunjian

Li[1], Hualong Feng[1], Lin Hu[1], Xuanyi Chen[1], Maosheng Lu[1], Liwei Jiang[1,*], Teng Zuo[1,*]

[1]Laboratory of Immunoengineering, Institute of Health and Medical Technology, Hefei

Institutes of Physical Science, Chinese Academy of Sciences, Hefei 230031, China

[2]These authors contributed equally to this work

*Correspondence: jlw0531@cmpt.ac.cn (L.J.); zuot@cmpt.ac.cn (T.Z.)

Summary

Whether Omicron exposures could overcome ancestral SARS-CoV-2 immune imprinting remains controversial. Here we analyzed B cell responses evoked by sequential Omicron infections in vaccinated and unvaccinated individuals. Plasma neutralizing antibody titers against ancestral SARS-CoV-2 and variants indicate that immune imprinting is not consistently induced by inactivated or recombinant protein vaccines. However, once induced, immune imprinting is not countered by successive Omicron challenges. We compared binding specificities, neutralizing capacities, developing origins and targeting epitopes of monoclonal antibodies from individuals with or without immune imprinting. Although receptor-binding domain (RBD) and N-terminal domain (NTD) of spike are both primary targets for neutralizing antibodies, immune imprinting only shapes antibody responses to RBD by impeding the production of Omicron-specific neutralizing antibodies while facilitating the development of broadly neutralizing antibodies. We propose that immune imprinting can be either neglected by NTD-based vaccines to induce variant-specific antibodies or leveraged by RBD-containing vaccines to induce broadly neutralizing antibodies.

Highlights

- Immune imprinting is not consistently induced by vaccination

- Repeated Omicron infections could not counter immune imprinting

- Immune imprinting shapes antibody responses to RBD but not NTD

- Immune imprinting leads to broadly neutralizing antibodies

Introduction

Immunological memory is a hallmark of adaptive immunity, which consists of humoral and cellular immunity[1]. Upon primary exposure to an antigen, humoral immune responses typically start from priming of naive B cells through cognate interaction between B cell receptor (BCR) and antigen in follicles of secondary lymphoid organs[2]. Primed B cells migrate to the border between B cell follicle and T cell zone, where they encounter activated CD4 T cells. Cognate T-B interaction leads to clonal expansion of B cells and formation of germinal centers (GCs). Within GCs, B cells undergo somatic hypermutation (SHM) on variable region of BCR to generate a pool of mutated B cells, which are then selected based on their affinities to the antigen. After repeated rounds of mutation and selection, affinity-matured GC B cells differentiate into plasmablasts (PBs), plasma cells (PCs) or memory B cells. PB/PCs constantly secret antibodies and maintain high levels of serum antibodies for months to years. Meanwhile, memory B cells circulate throughout the body until they encounter cognate antigens. Activated memory B cells rapidly differentiate into PB/PCs to secret antibodies, or to a lesser extent, into GC B cells to undergo another round of affinity maturation[3].

As the basic principle of long-term immune protection against pathogens after infection or vaccination, immunological memory ensures much faster and stronger immune responses upon antigen re-exposure[1]. However, the effects of immunological memory could be undesirable when antigens in prior and subsequent exposures are heterologous but antigenically related. One example of such scenario is that individuals' antibody responses against influenza viruses are largely biased towards

the predominant strains in their early lives[4]. This phenomenon is referred to as original

antigenic sin or immune imprinting, which describes the effects of immunity elicited by

prior antigen exposures on subsequent immune responses to antigenically related

antigens.

Since the outbreak of COVID-19 pandemic, the ancestral SARS-CoV-2 has kept

evolving and generated hundreds of thousands of variants. Among them, Omicron

emerged in November 2021 and accumulated more than 30 mutations on spike, the

target of neutralizing antibodies[5]. Over the past two and a half years, the predominant

strains have been rapidly iterated by Omicron-derived variants, including BA.1, BA.2,

BA.4, BA.5, BQ.1, XBB.1.5, XBB.1.16, XBB.1.9, EG.5.1 and BA.2.86. These variants

exhibit unprecedented neutralization evasion from antibodies elicited by ancestral

SARS-CoV-2 infection or vaccines based on ancestral SARS-CoV-2 [6-11]. Consequently,

several waves of infection by Omicron variants have occurred and updated vaccines

have been developed to match those variants [12-14].

Analysis of B cell responses in individuals who experienced breakthrough infection of

Omicron variants or received Omicron booster vaccines demonstrates the induction of

immune imprinting by ancestral SARS-CoV-2[15-20]. Moreover, the imprinted antibody

responses display substantially reduced neutralizing activities against later Omicron

variants compared with ancestral SARS-CoV-2 and earlier Omicron variants,

highlighting that immune imprinting may impede the generation of neutralizing

antibodies against future SARS-CoV-2 variants or other related viruses[21]. Notably, a

recent study reported that sequential Omicron exposures could overcome immune

imprinting and induce a large portion of *de novo* Omicron-specific antibodies[22]. In

contrast, other studies concluded that a single exposure to antigenically distant

XBB.1.5 or double exposures to Omicron variants rarely elicited Omicron-specific

antibody responses in the presence of immune imprinting[23-25]. Therefore, it is

controversial whether Omicron exposures could override ancestral SARS-CoV-2

immune imprinting.

In this study, we compared human B cell responses elicited by sequential Omicron

infections in the presence or absence of ancestral SARS-CoV-2 immune imprinting.

By measuring plasma neutralizing antibody titers against ancestral SARS-CoV-2 and

variants, we find that immune imprinting is not consistently induced by inactivated or

recombinant protein vaccines. However, once induced, immune imprinting is not

countered by repeated Omicron exposures. Comprehensive analysis of monoclonal

antibodies from individuals with or without immune imprinting demonstrates that RBD

and NTD are both primary targets for neutralizing antibodies, whereas only RBD-

specific antibody responses are imprinted. Consequently, Omicron-specific

neutralizing antibodies to RBD are hindered while broadly neutralizing antibodies to

RBD are elicited. These findings provide new insights into the development of next-

generation SARS-CoV-2 vaccines.

Results

Sequential Omicron infections could not overcome ancestral SARS-CoV-2

immune imprinting induced by inactivated or recombinant RBD vaccines.

To examine whether Omicron exposures could overcome ancestral SARS-CoV-2

immune imprinting, we collected peripheral blood from 12 donors of sequential

Omicron infections (Figure 1A). They were first infected in December 2022 or January

2023, when BA.5 and BF.7 were predominant strains[26]. Around half a year later, they

were infected again when XBB lineages (XBB*) were predominant strains[27]. Before the

infections, Donor A and B were not administrated with any vaccines based on ancestral

SARS-CoV-2, while the other donors received two or three doses of inactivated

vaccine or recombinant RBD vaccine or in combination.

We measured plasma neutralizing antibody titers against a panel of 18 pseudoviruses

bearing spikes from ancestral SARS-CoV-2 (WT), Alpha, Beta, Gamma, Delta, BA.1,

BA.4/5, BA.2.75, XBB, XBB.1.5, EG.5, EG.5.1, FL.1.5, FL.1.5.1, HK.3, BA.2.86, JN.1

and SARS-CoV-1 (Figure S1A). Donor A-D show poor or lower neutralizing titers

against earlier SARS-CoV-2 strains from WT to BA.1 while higher titers against strains

from BA.4/5 to HK.3 (Figure 1B). As Donor A and B were unvaccinated, their patterns

represent antibody responses induced by sequential Omicron exposures in the

absence of immune imprinting. Despite the fact that Donor C and D were vaccinated,

they display similar patterns with Donor A and B, suggesting that immune imprinting is

not induced by vaccination in Donor C and D. Donor E-K maintain higher titers against

earlier strains from WT to BA.2.75 while lower titers against XBB and later strains,

indicating the presence of immune imprinting in these individuals (Figure 1C). Different

from above donors, Donor L exhibits a quite chaotic pattern of neutralizing titers and

thus is excluded for further analysis (Figure S1B).

We compared neutralizing titers of Donor A-D (referred to as unimprinted group, G1)

and Donor E-K (referred to as imprinted group, G2) (Figure 1D). For early strains from

WT to BA.1, imprinted group has significantly higher titers than unimprinted group. On

the contrary, for more recent Omicron variants, including XBB, XBB.1.5, EG.5, EG.5.1,

FL.1.5 and HK.3, unimprinted group surpasses the imprinted group. Taken together,

these data suggest that immune imprinting is not consistently induced by vaccination.

However, once induced, immune imprinting is not countered by sequential Omicron

infections.

Immune imprinting leads to more frequent cross-reactive antibodies to RBD but

not NTD

To decipher the effects of immune imprinting on human B cell responses elicited by

sequential Omicron infections, we moved on to isolate and characterize monoclonal

antibodies (mAbs) from those donors. As they were sequentially exposed to BA.5/BF.7

and XBB*, we sorted single memory B cells or plasmablasts with XBB.1.5 spike as bait

(Figure S2). After cloning antibody variable regions from single cells into corresponding

antibody-expressing vectors, we transfected HEK293T cells with paired plasmids and

collected supernatant for ELISA and neutralization assay. The supernatant was tested

against spike (S), S1, RBD, NTD and S2 of XBB.1.5 and WT by ELISA, and tested

against XBB.1.5 and WT pseudoviruses by neutralization assay (Figure 2A). In total,

we identified 364 clones positive for XBB.1.5 spike from Donor A, B, C, D, E, F, G, I, J

and K. The proportions of antibodies against each domain vary in a wide range and no

significant differences are observed between unimprinted group (G1) and imprinted

group (G2) (Figure 2B). With 50% as cutoff for neutralization of supernatant, 84 clones

were identified as neutralizing antibodies against XBB.1.5 or WT or both (Figure 2A,

B). Notably, cross-neutralizing antibodies are all from imprinted group (Figure 2A). In

addition, the vast majority of neutralizing antibodies target either RBD or NTD and

there are no significant differences between the two groups (Figure 2B).

We further analyzed the frequency of cross-reactive antibodies (XBB.1.5$^+$ WT$^+$) (Figure

2C, D). Most S2-specific antibodies are cross-reactive, which should be due to the

conservation of this domain. For NTD-specific antibodies, the frequencies range from

0 to 50% and there is no difference between G1 and G2. In contrast, for RBD-specific

antibodies, G2 has remarkably higher frequency than G1, which represents recall of

cross-reactive memory B cells induced by vaccination. Overall, these results indicate

that immune imprinting mainly influences antibody responses to RBD after sequential

Omicron infections.

Immune imprinting facilitates the development of broadly neutralizing

antibodies against RBD

To determine the effects of immune imprinting on neutralizing antibodies, we purified

RBD- or NTD-specific mAbs with supernatant neutralization above or close to 50%. A

total number of 87 mAbs, including 16 RBD-specific and 32 NTD-specific mAbs from

G1 and 33 RBD-specific and 6 NTD-specific mAbs from G2, were tested against a

panel of SARS-CoV-2 pseudoviruses (Figure 3A). Similar to the neutralization patterns

of plasma from G1, RBD-specific mAbs from G1 (G1-RBD) poorly neutralize early

strains from WT to BA.1 while potently neutralize later Omicron variants. RBD-specific

mAbs from G2 can be divided into two subgroups according to their neutralization

patterns. One subgroup resembles mAbs from G1-RBD in neutralization pattern and

is named as non-WT-neutralizing (Abbreviated as non-WT-neut). The other subgroup

replicates the neutralization patterns of plasma from G2 and is termed as WT-

neutralizing (Abbreviated as WT-neut). It is worth mentioning that there are several

broadly neutralizing antibodies in this group, including KXD-E1, KXD-F3, KXD-F1,

KXD-G10 and KXD-G5. Most NTD-specific mAbs are able to neutralize BA.4/5, BQ.1,

XBB, XBB.1.5, EG.5.1 and HK.3, and the patterns are similar between G1 and G2.

Regarding to the neutralizing potency (IC50s) against different strains, mAbs from G1-

RBD are more potent to BA.4/5 than later variants whereas mAbs from G2-RBD-non-

WT-neut exhibit similar neutralizing activity against BA.4/5 and later strains (Figure

S3A). Antibodies from G2-RBD-WT-neut have equivalent potency to early strains from

WT to BA.4/5 while decreased potency to later variants (Figure S3A). Different from

RBD-specific mAbs, NTD-specific mAbs from both G1 and G2 are most potent to XBB

and XBB.1.5 (Figure S3B).

We compared the neutralizing potency of mAbs from different groups against the same

strain (Figure 3B,3C). Overall, mAbs from G1-RBD have similar neutralizing potency

with mAbs from G2-RBD-non-WT-neut. However, mAbs from G1-RBD are more potent

than mAbs from G2-RBD-WT-neut against variants from BA.4/5 to JN.1 except

BA.2.75 and XBB. On the other hand, NTD-specific mAbs from both groups have

analogous potency against all tested strains except HK.3. These data further confirm

that antibody responses to RBD while not NTD are affected by immune imprinting.

Moreover, the imprinted RBD-specific antibodies tend to have broadly neutralizing

activity although they generally show suboptimal neutralizing potency against Omicron

variants.

Repeated Omicron infections induce de novo Omicron-specific antibody

responses in the presence of immune imprinting

To examine whether repeated Omicron infections could induce de novo antibody

responses from naïve B cells in the presence of immune imprinting, we move on to

investigate the origins of neutralizing mAbs from each group. In published literatures,

it is generally assumed that cross-reactive antibodies derive from memory B cells

induced by WT whereas omicron-specific antibodies originate from naïve B cells

activated by Omicron[22-25]. However, these assumptions remain to be confirmed by

experiments to exclude possibilities that memory B cells induced by WT may lose

cross-reactivity and Omicron-specific naïve B cells may develop cross-reactivity during

affinity maturation. Here, we constructed germline versions of representative mAbs

from each group and tested their binding to a panel of spikes expressed on HEK293T

cells by FACS (Figure 4A). Germline antibodies from G1-RBD, G2-RBD-non-WT-neut,

G1-NTD and G2-NTD bind Omicron spikes but not WT spike. Oppositely, germline

antibodies from G2-RBD-WT-neut mostly recognize WT spike but not Omicron spikes.

These data confirm that mAbs from G1-RBD, G2-RBD-non-WT-neut, G1-NTD and G2-

NTD are de novo antibodies induced by Omicron, whereas mAbs from G2-RBD-WT-

neut are recalled antibodies initially induced by WT.

Correlated with their origins, mAbs from G2-RBD-WT-neut show higher mutation levels

than other mAbs, which reflects extensive affinity maturation sequentially driven by WT

and Omicron (Figure 4B). On the other hand, we also analyzed the repertoire of each

group (Figure 4C and Table 1). In G2-RBD-WT-neut group, IGHV1-18 and IGLV1-44

are most frequent as a consequence of B cell clonal expansion in Donor J. Moreover,

IGHV3-53, IGHV3-66 and IGHV1-69, which have been identified as frequent VH genes

for RBD-specific antibodies from individuals exposed to WT spike[28,29], are frequently

used in G2-RBD-WT-neut but not in G1-RBD and G2-RBD-non-WT-neut. For NTD-

specific antibodies, IGHV4-39 is most frequently used and is identified from multiple

donors in both groups. Therefore, these IGHV4-39 antibodies can be taken as a class

of public antibody against NTD. In sum, these data suggest that repeated Omicron

exposures could induce de novo antibody responses to RBD and NTD in the presence

of immune imprinting.

Omicron-specific and WT-cross-reactive antibodies recognize similar epitopes

To determine whether de novo antibody responses induced by repeated Omicron

infections recognize distinct epitopes compared with WT-cross-reactive antibodies, we

performed competition ELISA (Figure 5A, 5B). Three broadly neutralizing antibodies

targeting RBD, including BD55-1205, BD56-1854 and BD55-5483 were taken as

reference[30]. A major cluster is formed by the majority of RBD-specific antibodies. Most

antibodies in this cluster also compete with BD55-1205, BD56-1854 and ACE2.

Outside the major cluster, a minor cluster is formed by 6 antibodies. They compete

with BD55-5483 but not with BD55-1205, BD56-1854 and ACE2. The rest 3 RBD-

specific antibodies do not exhibit apparent competition with other antibodies or ACE2.

On the other hand, NTD-specific antibodies show competition with each other and form

a single cluster, suggesting that NTD-specific neutralizing antibodies target a supersite.

Taken together, these data provide explanation for the suppressed Omicron-specific

antibody responses to RBD in the presence of immune imprinting.

Discussion

In this study, we compared B cell responses evoked by sequential Omicron infections

in individuals vaccinated or unvaccinated with inactivated or recombinant RBD

vaccines. Unexpectedly, two vaccinees show no hints of immune imprinting, according

to their polyclonal and monoclonal antibody responses against ancestral SARS-CoV-

2 and variants. Therefore, they are classified into unimprinted group along with two

unvaccinated individuals throughout this study. Considering the factor of individual

variations, we believe that this classification is more reasonable than simply grouping

the cohorts based on antigen exposure history, especially for cohorts with limited size.

We speculate that no immune imprinting in these two individuals is probably due to

weak immune responses stimulated by vaccination. Moreover, whether this

phenomenon is common remains to be determined with larger cohorts.

In contrast to individuals without immune imprinting, individuals with immune imprinting

have lower plasma neutralizing antibody titers against multiple Omicron variants after

XBB, along with more frequent WT-cross-reactive antibodies and less frequent

Omicron-specific antibodies to RBD. Therefore, we conclude that immune imprinting

induced by inactivated or recombinant RBD vaccines is not countered by sequential

Omicron infections, which is inconsistent to a study performed by Yisimayi et al[22]. We

suppose this contradiction is due to following factors. First, the infections were

sequentially caused by BA.1/BA.2 and BA.5/BF.7 in their study whereas by BA.5/BF.7

and XBB* in our study. We notice that individuals with or without immune imprinting

have equivalent plasma neutralizing antibody titers to BA.4/5 and BA.2.75. Moreover,

individuals with immune imprinting show unaffected neutralizing titers against early

Omicron variants including BA.1, BA.2.75 and BA.4/5 compared with WT. These data

suggest that immune imprinting does not compromise antibody responses to early

Omicron variants after repeated Omicron exposures. Second, all vaccinees were

considered as imprinted in their study while two vaccinees were taken as unimprinted

in our study. As vaccinees without immune imprinting are able to generate abundant

Omicron-specific antibodies, excluding them from imprinted cohort may lead to more

clear results.

A major finding of our study is that antibody responses to NTD are not affected by

immune imprinting. As immunological memory and cross-reactivity are the basis of

immune imprinting[31], one possible explanation for this finding is that NTD is less

immunogenic than RBD because of its extensive glycan shielding[32,33], and thus few

NTD-specific memory B cells are induced. Another possible explanation is that NTD-

specific memory B cells induced by WT are not cross-reactive to Omicron variants.

However, consistent with previous studies[34-39], our results argue that NTD is highly

immunogenic and able to elicit antibodies cross-reactive to XBB.1.5 and WT, which

meet the preconditions for immune imprinting. Therefore, the mechanisms for the

unimprinted antibody responses to NTD remain to be explored.

Similar to previous studies[15-18,20,22-24], we prove that immune imprinting persistently

shapes antibody responses to RBD. As a result, RBD-specific antibodies can be

divided into two groups: WT-cross-reactive antibodies and Omicron-specific antibodies.

Compared with Omicron-specific antibodies, WT-cross-reactive antibodies tend to

have broader neutralization spectrum. For example, KXD-F1 neutralize all tested

pseudoviruses from WT to JN.1 with IC50s ranging from 0.006 to 0.084 µg/ml. As

germlines of those WT-cross-reactive antibodies generally have limited breadth, there

is no doubt that the breadth of WT-cross-reactive antibodies is attributed to affinity

maturation. However, the underlying mechanisms are not fully understood. According

to the model of affinity maturation in GCs, it is possible that WT-cross-reactive memory

B cells become Omicron-specific after affinity maturation. By comparing the binding

specificities of paired germline and mature antibodies, we find that all WT-reactive

germline antibodies consistently recognize WT spike after affinity maturation, which

negate the possibility like our previous study[40]. Oppositely, it is also possible that

Omicron-specific naïve B cells may become cross-reactive after affinity maturation.

However, we have not found evidence to support the possibility, despite that fact that

infection or vaccination with WT could elicit broadly neutralizing antibodies against

Omicron variants[30,41-43]. More studies are required to solve these puzzles on affinity

maturation.

Currently, ancestral SARS-CoV-2 immune imprinting has been acknowledged as a

great barrier for vaccine update. According to our findings, we propose that NTD-based

vaccines could be designed to bypass immune imprinting, since NTD is also a major

target of neutralizing antibodies. On the other hand, vaccines containing RBD could

be designed to leverage the potential of immune imprinting in eliciting broadly

neutralizing antibodies.

Limitations of the study

Although we included unvaccinated and vaccinated individuals in this study, both

cohorts are relatively limited in size. Moreover, the numbers of antibodies cloned from

different individuals are not equivalent, with a range from 6 to 89. In addition, only

XBB.1.5 spike was used to stain the B cells for single cell sorting, and thus the

frequencies of cross-reactive and Omicron-specific B cells were not determined by

FACS.

Materials and methods

Ethics statement

This study was approved by the Ethics Committee of Hefei Institutes of Physical Science, Chinese Academy of Sciences (Approval Number: YXLL-2023-47). All donors provided written informed consent for collection of information, analysis of plasma and PBMCs, and publication of data generated from their samples.

Human samples

Peripheral blood samples were collected from 12 donors (Figure1A). Plasma and peripheral blood mononuclear cells (PBMCs) were separated from blood by Ficoll density gradient centrifugation.

Cell lines

HEK293T cells expressing human ACE2 (HEK293T-hACE2) were kindly provided by Prof. Ji Wang at Sun Yat-Sen University. HEK293T cells were from ATCC. HEK293T and HEK293T-hACE2 cells were cultured in DMEM with 10% fetal bovine serum (FBS, VivaCell, Shanghai, China, C04001) and 1% penicillin/streptomycin (pen/strep). FreeStyle 293F cells (Thermo Fisher Scientific) were cultured in SMM 293-TII Expression Medium (Sino Biological Inc., M293TII). All cells were maintained in a 37˚C incubator at 5% CO_2

Protein expression and purification

The genes encoding the ectodomain of SARS-CoV-2 spike including WT and XBB.1.5 were constructed with foldon trimerization motif, His tag, tandem strep-tag II and FLAG tag at C-terminal. The genes encoding the S1, NTD and RBD of WT and XBB.1.5 were

constructed with a strep-tag II at C-terminal. The genes encoding the S2 of WT and

XBB.1.5 were constructed with a His tag at C-terminal. Spike trimer, S1, S2, NTD and

RBD were expressed in the FreeStyle 293F cells and purified by Streptactin Agarose

Resin 4FF (Yeasen, 20495ES60) or Ni-NTA Agarose (Qiagen, 30210).

Spike-specific Single B cell sorting and PCR

PBMCs were incubated with 200nM XBB.1.5 spike for 30 min at 4°C. After wash, they

were stained with cell-surface antibodies: CD3-BV510 (BioLegend, 317331), CD19-

PE/Cy7 (BioLegend, 302215), CD27-APC (BioLegend, 356409), CD38-APC/Cy7

(BioLegend, 356615), human IgM-AF700 (BioLegend, 314537), human IgD-

perCP/Cy5.5 (BioLegend, 348207), anti-His-AF488 (Proteintech, CL488-66005), anti-

FLAG-PE (BioLegend, 637309) and DAPI. The stained cells were washed with FACS

buffer (PBS containing 2% FBS) and resuspended in 500μl FACS buffer. Spike-specific

single B cells were gated as DAPI$^-$CD3$^-$CD19$^+$CD27$^+$IgD$^-$His$^+$FLAG$^+$ and sorted into

96-well PCR plates containing 4μl lysis buffer (0.5×PBS, 0.1M DTT and RNase

inhibitor) per well. After reverse transcription reaction, variable regions of heavy and

light chains were amplified by nested PCR and cloned into human IgG1 expression

vectors. Plasmids for paired heavy and light chains were co-transfected into HEK293T

cells or FreeStyle 293F cells. Antibodies were purified with Protein A magnetic beads

(GenScript, L00273).

ELISA

Spike, S1, S2, NTD or RBD were coated onto 96-well ELISA plates (100 ng/well) and

incubated at 4°C overnight. After blocking with PBS containing 10% FBS, HEK293T

supernatant was added to the wells and incubated at 37°C for 1hr. HRP-conjugated

goat anti-human IgG antibodies (Zen-bio, 550004; 1:2000 dilution) were added to the

wells and incubated at 37°C for 1hr. TMB substrate (Sangon Biotech, E661007-0100)

was added to the wells and incubated at room temperature for 5 mins. The reaction

was stopped by TMB Stop Solution (Sangon Biotech, E661006-0500) and absorbance

at 450 nm was measured.

Pseudovirus neutralization assay

To generate pseudoviruses, HEK293T cells were transfected with psPAX2, pLenti-

luciferase and spike-encoding plasmids using polyetherimide (PEI). Supernatants with

pseudoviruses were collected 48hrs after transfection. 3-fold serially diluted plasma

(starting at 1:20), HEK293T supernatant (starting at 1:4.5), or mAbs (starting at

10μg/ml) were mixed with pseudoviruses at 37°C for 1hr. HEK293T-hACE2 cells

($1.5×10^4$ per well) were added into the mixture and incubated at 37°C for 48hrs. Cells

were lysed to measure luciferase activity by Bright-Lite Luciferase Assay System

(Vazyme Biotech, DD1204-02). The percentages of neutralization were determined by

comparing with the virus control. The plasmids encoding spike of WT, Alpha, Beta,

Gamma, Delta, BA.1, BA.4/5, XBB, XBB.1.5 and SARS-CoV-1 were kindly provided

by Prof. Linqi Zhang at Tsinghua University. The plasmids encoding spike of BA.2.75

was kindly provided by Prof. Zezhong Liu at Fudan University. The sequences of EG.5,

EG.5.1, FL.1.5, FL.1.5.1, HK.3, BA.2.86 and JN.1 spike were synthesized and cloned

into expression vectors.

Competition ELISA

NTD-specific and RBD-specific neutralizing mAbs were labeled with HRP (ProteinTech,

PK20001). XBB.1.5 spike was coated onto 96-well ELISA plates (50 ng/well) and

incubated at 4°C overnight. After blocking with PBS containing 10% FBS, blocking

mAbs were added to the wells and incubated at 37°C for 1hr. HRP-labeled detecting

mAbs were added to the wells in the presence of competing mAbs. The ratio of

blocking antibody and detecting antibody is 100:1. After 1hr incubation at 37°C, TMB

substrate (Sangon Biotech, E661007-0100) was added to the wells and incubated at

room temperature for 30 mins. The reaction was stopped by TMB Stop Solution

(Sangon Biotech, E661006-0500) and absorbance at 450 nm were measured. The

percentages of signal decrease caused by blocking mAbs were calculated.

ACE2 competition FACS

5nM biotinylated XBB.1.5 spike was incubated with 10µg/ml competing mAbs for

30min at 4°C, then the mixtures were added to HEK293T-hACE2 cells. After 30min at

4°C, the cells were washed with FACS buffer. Streptavidin-APC (BioLegend, 405207)

and DAPI were added to the wells at 4°C for 30min. The stained cells were washed

with FACS buffer and resuspended in 200µl FACS buffer. Positive cells were gated as

DAPI$^-$XBB.1.5 spike$^+$. The percentages of signal decrease caused by competing mAbs

were calculated.

Germline antibodies binding FACS

The plasmids encoding spikes of SARS-CoV-2 WT, Alpha, Beta, Gamma, Delta, BA.1,

BA.4/5, BF.7, BA.2.75, BQ.1, XBB.1.5, EG.5.1, BA.2.86 were transfected into

HEK293T cells. After 48hr, cells were harvested and incubated with 50µg/ml germline

antibodies for 30min at 4°C. After washed with FACS buffer, cells were stained with

goat anti-human IgG FITC (Proteintech, SA00003-12) and DAPI. The stained cells

were washed with FACS buffer and resuspended in 200µl FACS buffer. Positive cells

were gated as DAPI$^-$IgG$^+$.

Acknowledgements

The authors thank all donors for providing peripheral blood. This work was supported

by the National Key Plan for Scientific Research and Development of China (Grant No.

2022YFC2305800) and the National Natural Science Foundation of China (Grant No.

32200765, Grant No. 32370941).

Author contributions

Conceptualization, L.J. and T.Z.; Methodology, L.J. and T.Z.; Validation, Z.W., L.L., R.D.,

X.C., Y.S.; Formal Analysis, Z.W.; Investigation, Z.W., L.L., R.D., X.C., Y.S., R.Q., Y.L.,

H.F., L.H., X.C., M.L., L.J. and T.Z.; Writing – Original Draft, L.J. and T.Z.; Writing –

Review & Editing, L.J., T.Z., Z.W., L.L.; Visualization, T.Z., Z.W.; Supervision, L.J. and

T.Z.; Funding Acquisition, L.J. and T.Z.;

Competing interests

L.J., T.Z., L.L. and Z.W. are inventors on a patent application for antibodies including

KXD-E1, KXD-F1, KXD-F3, KXD-G5 and KXD-G10. The other authors declare that

they have no competing interests.

Supplemental information

Figures S1–S3 and Table S1

Data S1. Excel spreadsheet containing numerical data for figures.

Figure 1. Sequential Omicron infections could not overcome ancestral SARS-CoV-2 immune imprinting induced by inactivated or recombinant RBD vaccines.

(A) Vaccination and infection information of 12 donors. (B, C) Summary of plasma neutralizing antibody titers (NT50). For plasma with no neutralizing activity at 20-fold dilution, a number between 10–20 is given as titer to separate the curves. The numbers below the line are geometric mean titers against the pseudoviruses. The titers against WT are compared with other variants and the fold changes are indicated above the line. For panel B, statistical analysis was performed by two-tailed paired T test. For panel C, statistical analysis was performed by two-tailed Wilcoxon matched-pairs signed rank test. $*P < 0.05$, $**P < 0.01$, $**** P < 0.0001$. (D) Comparison of neutralizing antibody titers of G1 (Donor A-D, unimprinted group) and G2 (Donor E-K, imprinted group). The numbers on top are fold changes. Statistical analysis was performed by two-tailed unpaired T test. $*P < 0.05$, $**P < 0.01$. The neutralizing titers are calculated with results from two independent experiments, in which duplicates are performed.

Figure 2. Immune imprinting leads to more frequent cross-reactive antibodies to RBD but not NTD. (A) ELISA and neutralization results of 364 mAbs. Supernatant of HEK293T cells was tested against Spike, S1, S2, NTD and RBD of WT and XBB.1.5 by ELISA, and tested against WT and XBB.1.5. pseudoviruses by neutralization assay. For ELISA, supernatant was not diluted. For neutralization, supernatant was diluted by 4.5-fold. The results are mean values of two independent experiments. For neutralization, duplicates are performed in each experiment. (B) Frequency of binding

mAbs and neutralizing mAbs against different domains in each individual. (C)

Frequencies of WT-cross-reactive antibodies (XBB.1.5$^+$ WT$^+$) and Omicron-specific

antibodies (XBB.1.5$^+$ WT$^-$) in each individual. (D) Comparison of WT-cross-reactive

antibodies between G1 and G2. Data are represented as mean ± SD. Statistical

analysis was performed by two-tailed unpaired T test. NS (Not significant) p > 0.05, **P

< 0.01.

Figure 3. Immune imprinting facilitates the development of broadly neutralizing

antibodies against RBD. (A) IC50s of mAbs against WT, Alpha, Beta, Gamma, Delta,

BA.1, BA.4/5, BA.2.75, BQ.1, XBB, XBB.1.5, EG.5.1, HK.3, BA.2.86 and JN.1. The

highest antibody concentration used to determine IC50 is 10µg/ml. (B) Comparing

IC50s of RBD-specific mAbs from three groups. (C) Comparing IC50s of NTD-specific

mAbs from two groups. Data are represented as the mean ± SD. Statistical analysis

was performed by two-tailed unpaired T test. NS (Not significant) p > 0.05, *P < 0.05,

**P < 0.01. IC50s are calculated with results from two independent experiments, in

which duplicates are performed.

Figure 4. Repeated Omicron infections induce de novo Omicron-specific

antibody responses in the presence of immune imprinting. (A) Binding

specificities of germline antibodies measured by FACS. HEK293T cells expressing

SARS-CoV-2 spikes were incubated with 50µg/ml germline antibodies and analyzed

by FACS. The percentages of positive cells are calculated. The results are mean

values of two independent experiments. (B) Somatic hypermutations of VH and VL.

DNA sequences were analyzed. Data are represented as mean ± SD. Statistical

analysis was performed by two-tailed unpaired T test. NS (Not significant) p > 0.05, **P

< 0.01, ***P < 0.001, ****P < 0.0001. (C) Frequencies of V genes in each group.

Frequently used V genes are highlighted by colored background.

Figure 5. Omicron-specific and WT-cross-reactive antibodies recognize similar

epitopes. (A) Competition ELISA of RBD-specific mAbs. Competition with ACE2 was

performed by FACS. (B) Competition ELISA of NTD-specific mAbs. The results are

mean values of two independent experiments.

Figure S1. Plasma neutralizing antibody titers. (A) Measurement of neutralizing

antibody titers against a panel of pseudoviruses. Data are represented as mean ± SD.

(B) Neutralizing antibody titers of donor L.

Figure S2. FACS plots representing gating strategies for XBB.1.5 spike-specific

single B cell sorting.

Figure S3. Neutralizing potency of mAbs against different variants. (A) IC50s of

RBD-specific mAbs. The numbers below the line are geometric mean IC50 of mAbs

against the pseudoviruses. IC50s against BA.4/5 are compared with other variants. (B)

IC50s of NTD-specific mAbs. The numbers below the line are geometric mean IC50 of

mAbs against the pseudoviruses. IC50s against XBB are compared other variants.

Statistical analysis was performed by two-tailed Wilcoxon matched-pairs signed rank

test. NS (Not significant) p > 0.05, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

References

1. Gray, D. (1993). Immunological memory. Annu Rev Immunol *11*, 49-77. 10.1146/annurev.iy.11.040193.000405.

2. Victora, G.D., and Nussenzweig, M.C. (2022). Germinal Centers. Annu Rev Immunol *40*, 413-442. 10.1146/annurev-immunol-120419-022408.

3. Mesin, L., Schiepers, A., Ersching, J., Barbulescu, A., Cavazzoni, C.B., Angelini, A., Okada, T., Kurosaki, T., and Victora, G.D. (2020). Restricted Clonality and Limited Germinal Center Reentry Characterize Memory B Cell Reactivation by Boosting. Cell *180*, 92-106 e111. 10.1016/j.cell.2019.11.032.

4. Francis, T. (1960). On the Doctrine of Original Antigenic Sin. Proceedings of the American Philosophical Society *104*, 572-578.

5. Viana, R., Moyo, S., Amoako, D.G., Tegally, H., Scheepers, C., Althaus, C.L., Anyaneji, U.J., Bester, P.A., Boni, M.F., Chand, M., et al. (2022). Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. Nature *603*, 679-686. 10.1038/s41586-022-04411-y.

6. He, Q., Wu, L., Xu, Z., Wang, X., Xie, Y., Chai, Y., Zheng, A., Zhou, J., Qiao, S., Huang, M., et al. (2023). An updated atlas of antibody evasion by SARS-CoV-2 Omicron sub-variants including BQ.1.1 and XBB. Cell Rep Med *4*, 100991. 10.1016/j.xcrm.2023.100991.

7. Kurhade, C., Zou, J., Xia, H., Liu, M., Chang, H.C., Ren, P., Xie, X., and Shi, P.Y. (2023). Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1 and XBB.1 by parental mRNA vaccine or a BA.5 bivalent booster. Nat Med *29*, 344-347. 10.1038/s41591-022-02162-x.

8. Wang, Q., Iketani, S., Li, Z., Liu, L., Guo, Y., Huang, Y., Bowen, A.D., Liu, M., Wang, M., Yu, J., et al. (2023). Alarming antibody evasion properties of rising SARS-CoV-2 BQ and XBB subvariants. Cell *186*, 279-286 e278. 10.1016/j.cell.2022.12.018.

9. Miller, J., Hachmann, N.P., Collier, A.Y., Lasrado, N., Mazurek, C.R., Patio, R.C., Powers, O., Surve, N., Theiler, J., Korber, B., and Barouch, D.H. (2023). Substantial Neutralization Escape by SARS-CoV-2 Omicron Variants BQ.1.1 and XBB.1. N Engl J Med *388*, 662-664. 10.1056/NEJMc2214314.

10. Wang, Q., Guo, Y., Zhang, R.M., Ho, J., Mohri, H., Valdez, R., Manthei, D.M., Gordon, A., Liu, L., and Ho, D.D. (2023). Antibody neutralisation of emerging SARS-CoV-2 subvariants: EG.5.1 and XBC.1.6. Lancet Infect Dis *23*, e397-e398. 10.1016/S1473-3099(23)00555-8.

11. Yang, S., Yu, Y., Xu, Y., Jian, F., Song, W., Yisimayi, A., Wang, P., Wang, J., Liu, J., Yu, L., et al. (2024). Fast evolution of SARS-CoV-2 BA.2.86 to JN.1 under heavy immune pressure. Lancet Infect Dis *24*, e70-e72. 10.1016/S1473-3099(23)00744-2.

12. Chalkias, S., Eder, F., Essink, B., Khetan, S., Nestorova, B., Feng, J., Chen, X., Chang, Y., Zhou, H., Montefiori, D., et al. (2022). Safety, immunogenicity and antibody persistence of a bivalent Beta-containing booster vaccine against COVID-19: a phase 2/3 trial. Nat Med *28*, 2388-2397. 10.1038/s41591-022-02031-7.

13. Chalkias, S., Harper, C., Vrbicky, K., Walsh, S.R., Essink, B., Brosz, A., McGhee, N., Tomassini, J.E., Chen, X., Chang, Y., et al. (2022). A Bivalent Omicron-Containing Booster Vaccine against Covid-19. N Engl J Med *387*, 1279-1291. 10.1056/NEJMoa2208343.

14. Collier, A.Y., Miller, J., Hachmann, N.P., McMahan, K., Liu, J., Bondzie, E.A., Gallup, L., Rowe, M., Schonberg, E., Thai, S., et al. (2023). Immunogenicity of BA.5 Bivalent mRNA Vaccine

Boosters. N Engl J Med *388*, 565-567. 10.1056/NEJMc2213948.

15.   Park, Y.J., Pinto, D., Walls, A.C., Liu, Z., De Marco, A., Benigni, F., Zatta, F., Silacci-Fregni, C.,
Bassi, J., Sprouse, K.R., et al. (2022). Imprinted antibody responses against SARS-CoV-2
Omicron sublineages. Science *378*, 619-627. 10.1126/science.adc9127.

16.   Wang, Z., Zhou, P., Muecksch, F., Cho, A., Ben Tanfous, T., Canis, M., Witte, L., Johnson, B.,
Raspe, R., Schmidt, F., et al. (2022). Memory B cell responses to Omicron subvariants after
SARS-CoV-2 mRNA breakthrough infection in humans. J Exp Med *219*.
10.1084/jem.20221006.

17.   Kaku, C.I., Bergeron, A.J., Ahlm, C., Normark, J., Sakharkar, M., Forsell, M.N.E., and Walker,
L.M. (2022). Recall of preexisting cross-reactive B cell memory after Omicron BA.1
breakthrough infection. Sci Immunol *7*, eabq3511. 10.1126/sciimmunol.abq3511.

18.   Quandt, J., Muik, A., Salisch, N., Lui, B.G., Lutz, S., Kruger, K., Wallisch, A.K., Adams-Quack,
P., Bacher, M., Finlayson, A., et al. (2022). Omicron BA.1 breakthrough infection drives
cross-variant neutralization and memory B cell formation against conserved epitopes. Sci
Immunol *7*, eabq2427. 10.1126/sciimmunol.abq2427.

19.   Weber, T., Dahling, S., Rose, S., Affeldt, P., Vanshylla, K., Ullrich, L., Gieselmann, L., Teipel,
F., Gruell, H., Di Cristanziano, V., et al. (2023). Enhanced SARS-CoV-2 humoral immunity
following breakthrough infection builds upon the preexisting memory B cell pool. Sci
Immunol *8*, eadk5845. 10.1126/sciimmunol.adk5845.

20.   Addetia, A., Piccoli, L., Case, J.B., Park, Y.J., Beltramello, M., Guarino, B., Dang, H., de Melo,
G.D., Pinto, D., Sprouse, K., et al. (2023). Neutralization, effector function and immune
imprinting of Omicron variants. Nature *621*, 592-601. 10.1038/s41586-023-06487-6.

21.   Wang, Q., Guo, Y., Tam, A.R., Valdez, R., Gordon, A., Liu, L., and Ho, D.D. (2023). Deep
immunological imprinting due to the ancestral spike in the current bivalent COVID-19
vaccine. Cell Rep Med *4*, 101258. 10.1016/j.xcrm.2023.101258.

22.   Yisimayi, A., Song, W., Wang, J., Jian, F., Yu, Y., Chen, X., Xu, Y., Yang, S., Niu, X., Xiao, T.,
et al. (2024). Repeated Omicron exposures override ancestral SARS-CoV-2 immune
imprinting. Nature *625*, 148-156. 10.1038/s41586-023-06753-7.

23.   Johnston, T.S., Li, S.H., Painter, M.M., Atkinson, R.K., Douek, N.R., Reeg, D.B., Douek, D.C.,
Wherry, E.J., and Hensley, S.E. (2024). Immunological imprinting shapes the specificity of
human antibody responses against SARS-CoV-2 variants. Immunity *57*, 912-925 e914.
10.1016/j.immuni.2024.02.017.

24.   Tortorici, M.A., Addetia, A., Seo, A.J., Brown, J., Sprouse, K., Logue, J., Clark, E., Franko, N.,
Chu, H., and Veesler, D. (2024). Persistent immune imprinting occurs after vaccination with
the COVID-19 XBB.1.5 mRNA booster in humans. Immunity *57*, 904-911 e904.
10.1016/j.immuni.2024.02.016.

25.   Liang, C.Y., Raju, S., Liu, Z., Li, Y., Asthagiri Arunkumar, G., Case, J.B., Scheaffer, S.M., Zost,
S.J., Acreman, C.M., Gagne, M., et al. (2024). Imprinting of serum neutralizing antibodies
by Wuhan-1 mRNA vaccines. Nature. 10.1038/s41586-024-07539-1.

26.   Pan, Y., Wang, L., Feng, Z., Xu, H., Li, F., Shen, Y., Zhang, D., Liu, W.J., Gao, G.F., and Wang,
Q. (2023). Characterisation of SARS-CoV-2 variants in Beijing during 2022: an
epidemiological and phylogenetic analysis. Lancet *401*, 664-672. 10.1016/S0140-
6736(23)00129-0.

27.   Feng, Y., Su, Q., Li, L., He, X., Niu, P., Guo, X., Zhao, X., Tang, J., Jia, Z., and Wang, J. (2024).

Genomic Surveillance for SARS-CoV-2 Variants: Dominance of XBB Replacement—China,
January–June 2023. China CDC Weekly *6*, 324-331.

28.   Wang, Y., Yuan, M., Lv, H., Peng, J., Wilson, I.A., and Wu, N.C. (2022). A large-scale
systematic survey reveals recurring molecular features of public antibody responses to
SARS-CoV-2. Immunity *55*, 1105-1117 e1104. 10.1016/j.immuni.2022.03.019.

29.   Yan, Q., Hou, R., Huang, X., Zhang, Y., He, P., Zhang, Y., Liu, B., Wang, Q., Rao, H., Chen,
X., et al. (2022). Shared IGHV1-69-encoded neutralizing antibodies contribute to the
emergence of L452R substitution in SARS-CoV-2 variants. Emerg Microbes Infect *11*,
2749-2761. 10.1080/22221751.2022.2140611.

30.   Cao, Y., Jian, F., Wang, J., Yu, Y., Song, W., Yisimayi, A., Wang, J., An, R., Chen, X., Zhang,
N., et al. (2023). Imprinted SARS-CoV-2 humoral immunity induces convergent Omicron
RBD evolution. Nature *614*, 521-529. 10.1038/s41586-022-05644-7.

31.   Koutsakos, M., and Ellebedy, A.H. (2023). Immunological imprinting: Understanding
COVID-19. Immunity *56*, 909-913. 10.1016/j.immuni.2023.04.012.

32.   Walls, A.C., Park, Y.J., Tortorici, M.A., Wall, A., McGuire, A.T., and Veesler, D. (2020).
Structure, Function, and Antigenicity of the SARS-CoV-2 Spike Glycoprotein. Cell *181*,
281-292 e286. 10.1016/j.cell.2020.02.058.

33.   Wrapp, D., Wang, N., Corbett, K.S., Goldsmith, J.A., Hsieh, C.L., Abiona, O., Graham, B.S.,
and McLellan, J.S. (2020). Cryo-EM structure of the 2019-nCoV spike in the prefusion
conformation. Science *367*, 1260-1263. 10.1126/science.abb2507.

34.   Chi, X., Yan, R., Zhang, J., Zhang, G., Zhang, Y., Hao, M., Zhang, Z., Fan, P., Dong, Y., Yang,
Y., et al. (2020). A neutralizing human antibody binds to the N-terminal domain of the
Spike protein of SARS-CoV-2. Science *369*, 650-655. 10.1126/science.abc6952.

35.   Liu, L., Wang, P., Nair, M.S., Yu, J., Rapp, M., Wang, Q., Luo, Y., Chan, J.F., Sahi, V., Figueroa,
A., et al. (2020). Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2
spike. Nature *584*, 450-456. 10.1038/s41586-020-2571-7.

36.   McCallum, M., De Marco, A., Lempp, F.A., Tortorici, M.A., Pinto, D., Walls, A.C., Beltramello,
M., Chen, A., Liu, Z., Zatta, F., et al. (2021). N-terminal domain antigenic mapping reveals
a site of vulnerability for SARS-CoV-2. Cell *184*, 2332-2347 e2316.
10.1016/j.cell.2021.03.028.

37.   Suryadevara, N., Shrihari, S., Gilchuk, P., VanBlargan, L.A., Binshtein, E., Zost, S.J., Nargi,
R.S., Sutton, R.E., Winkler, E.S., Chen, E.C., et al. (2021). Neutralizing and protective human
monoclonal antibodies recognizing the N-terminal domain of the SARS-CoV-2 spike
protein. Cell *184*, 2316-2331 e2315. 10.1016/j.cell.2021.03.029.

38.   Wang, Z., Muecksch, F., Cho, A., Gaebler, C., Hoffmann, H.H., Ramos, V., Zong, S., Cipolla,
M., Johnson, B., Schmidt, F., et al. (2022). Analysis of memory B cells identifies conserved
neutralizing epitopes on the N-terminal domain of variant SARS-Cov-2 spike proteins.
Immunity *55*, 998-1012 e1018. 10.1016/j.immuni.2022.04.003.

39.   Cerutti, G., Guo, Y., Wang, P., Nair, M.S., Wang, M., Huang, Y., Yu, J., Liu, L., Katsamba, P.S.,
Bahna, F., et al. (2021). Neutralizing antibody 5-7 defines a distinct site of vulnerability in
SARS-CoV-2 spike N-terminal domain. Cell Rep *37*, 109928.
10.1016/j.celrep.2021.109928.

40.   Li, L., Chen, X., Wang, Z., Li, Y., Wang, C., Jiang, L., and Zuo, T. (2023). Breakthrough
infection elicits hypermutated IGHV3-53/3-66 public antibodies with broad and potent

neutralizing activity against SARS-CoV-2 variants including the emerging EG.5 lineages.
PLoS Pathog *19*, e1011856. 10.1371/journal.ppat.1011856.

41.　Ju, B., Zhang, Q., Wang, Z., Aw, Z.Q., Chen, P., Zhou, B., Wang, R., Ge, X., Lv, Q., Cheng, L.,
et al. (2023). Infection with wild-type SARS-CoV-2 elicits broadly neutralizing and
protective antibodies against omicron subvariants. Nat Immunol *24*, 690-699.
10.1038/s41590-023-01449-6.

42.　Liu, Y., Wang, Z., Zhuang, X., Zhang, S., Chen, Z., Zou, Y., Sheng, J., Li, T., Tai, W., Yu, J., et
al. (2023). Inactivated vaccine-elicited potent antibodies can broadly neutralize SARS-
CoV-2 circulating variants. Nat Commun *14*, 2179. 10.1038/s41467-023-37926-7.

43.　Ju, B., Fan, Q., Liu, C., Shen, S., Wang, M., Guo, H., Zhou, B., Ge, X., and Zhang, Z. (2023).
Omicron BQ.1.1 and XBB.1 unprecedentedly escape broadly neutralizing antibodies
elicited by prototype vaccination. Cell Rep *42*, 112532. 10.1016/j.celrep.2023.112532.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure S1

Figure S2

Figure S3