Ancestral SARS-CoV-2 immune imprinting persists on RBD but not NTD after

sequential Omicron infections

Zuowei Wang[1,2], Ling Li[1,2], Ruiping Du[1], Xixian Chen[1], Yi Sun[1], Rongrong Qin[1], Yunjian

Li[1], Hualong Feng[1], Lin Hu[1], Xuanyi Chen[1], Maosheng Lu[1], Liwei Jiang[1,*], Teng Zuo[1,*]

[1]Laboratory of Immunoengineering, Institute of Health and Medical Technology, Hefei

Institutes of Physical Science, Chinese Academy of Sciences, Hefei 230031, China

[2]These authors contributed equally to this work

*Correspondence: jlw0531@cmpt.ac.cn (L.J.); zuot@cmpt.ac.cn (T.Z.)

Abstract

Whether Omicron exposures could overcome ancestral SARS-CoV-2 immune imprinting remains controversial. Here we analyzed B cell responses evoked by sequential Omicron infections in vaccinated and unvaccinated individuals. Plasma neutralizing antibody titers against ancestral SARS-CoV-2 and variants indicate that immune imprinting is not consistently induced by inactivated or recombinant protein vaccines. However, once induced, immune imprinting is not countered by successive Omicron challenges. We compared binding specificities, neutralizing capacities, developing origins and targeting epitopes of monoclonal antibodies from individuals with or without immune imprinting. Although receptor-binding domain (RBD) and N-terminal domain (NTD) of spike are both primary targets for neutralizing antibodies, immune imprinting only shapes antibody responses to RBD by impeding the production of Omicron-specific neutralizing antibodies while facilitating the development of broadly neutralizing antibodies. We propose that immune imprinting can be either neglected by NTD-based vaccines to induce variant-specific antibodies or leveraged by RBD-based vaccines to induce broadly neutralizing antibodies.

Introduction

Immunological memory is a hallmark of adaptive immunity, which consists of humoral and cellular immunity(1). Upon primary exposure to an antigen, humoral immune responses typically start from priming of naive B cells through cognate interaction between B cell receptor (BCR) and antigen in follicles of secondary lymphoid organs(2). Primed B cells migrate to the border between B cell follicle and T cell zone, where they encounter activated CD4 T cells. Cognate T-B interaction leads to clonal expansion of B cells and formation of germinal centers (GCs). Within GCs, B cells undergo somatic hypermutation (SHM) on variable region of BCR to generate a pool of mutated B cells, which are then selected based on their affinities to the antigen. After repeated rounds of mutation and selection, affinity-matured GC B cells differentiate into plasmablasts (PBs), plasma cells (PCs) or memory B cells. PB/PCs constantly secrete antibodies and maintain high levels of serum antibodies for months to years. Meanwhile, memory B cells circulate throughout the body until they encounter cognate antigens. Activated memory B cells rapidly differentiate into PB/PCs to secrete antibodies, or to a lesser extent, into GC B cells to undergo another round of affinity maturation(3).

As the basic principle of long-term immune protection against pathogens after infection or vaccination, immunological memory ensures much faster and stronger immune responses upon antigen re-exposure(1). However, the effects of immunological memory could be undesirable when antigens in prior and subsequent exposures are heterologous but antigenically related. One example of such scenario is that individuals' antibody responses against influenza viruses are largely biased towards

the predominant strains in their early lives(4). This phenomenon is referred to as

original antigenic sin or immune imprinting, which describes the effects of immunity

elicited by prior antigen exposures on subsequent immune responses to antigenically

related antigens.

Since the outbreak of COVID-19 pandemic, the ancestral SARS-CoV-2 has kept

evolving and generated hundreds of thousands of variants. Among them, Omicron

emerged in November 2021 and accumulated more than 30 mutations on spike, the

target of neutralizing antibodies(5). Over the past two and a half years, the

predominant strains have been rapidly iterated by Omicron-derived variants, including

BA.1, BA.2, BA.4, BA.5, BQ.1, XBB.1.5, XBB.1.16, XBB.1.9, EG.5.1 and BA.2.86.

These variants exhibit unprecedented neutralization evasion from antibodies elicited

by ancestral SARS-CoV-2 infection or vaccines based on ancestral SARS-CoV-2 (6-

11). Consequently, several waves of infection by Omicron variants have occurred and

updated vaccines have been developed to match those variants (12-14).

Analysis of B cell responses in individuals who experienced breakthrough infection of

Omicron variants or received Omicron booster vaccines demonstrates the induction of

immune imprinting by ancestral SARS-CoV-2(15-20). Moreover, the imprinted antibody

responses display substantially reduced neutralizing activities against later Omicron

variants compared with ancestral SARS-CoV-2 and earlier Omicron variants,

highlighting that immune imprinting may impede the generation of neutralizing

antibodies against future SARS-CoV-2 variants or other related viruses(21). Notably,

a recent study reported that sequential Omicron exposures could overcome immune

imprinting and induce a large portion of *de novo* Omicron-specific antibodies(22). In contrast, other studies concluded that a single exposure to antigenically distant XBB.1.5 or double exposures to Omicron variants rarely elicited Omicron-specific antibody responses in the presence of immune imprinting(23-25). Therefore, it is controversial whether Omicron exposures could override ancestral SARS-CoV-2 immune imprinting.

In this study, we compared human B cell responses elicited by sequential Omicron infections in the presence or absence of ancestral SARS-CoV-2 immune imprinting. By measuring plasma neutralizing antibody titers against ancestral SARS-CoV-2 and variants, we find that immune imprinting is not consistently induced by inactivated or recombinant protein vaccines. However, once induced, immune imprinting is not countered by repeated Omicron exposures. Comprehensive analysis of monoclonal antibodies from individuals with or without immune imprinting demonstrates that RBD and NTD are both primary targets for neutralizing antibodies, whereas only RBD-specific antibody responses are imprinted. Consequently, Omicron-specific neutralizing antibodies to RBD are hindered while broadly neutralizing antibodies to RBD are elicited. These findings provide new insights into the development of next-generation SARS-CoV-2 vaccines.

Results

Sequential Omicron infections could not overcome ancestral SARS-CoV-2 immune imprinting induced by inactivated or recombinant RBD vaccines.

To examine whether Omicron exposures could overcome ancestral SARS-CoV-2 immune imprinting, we collected peripheral blood from 12 donors of sequential Omicron infections (Figure 1A). They were first infected in December 2022 or January 2023, when BA.5 and BF.7 were predominant strains(26). Around half a year later, they were infected again when XBB lineages (XBB*) were predominant strains(27). Before the infections, Donor A and B were not administrated with any vaccines based on ancestral SARS-CoV-2, while the other donors received two or three doses of inactivated vaccine or recombinant RBD vaccine or in combination.

We measured plasma neutralizing antibody titers against a panel of 18 pseudoviruses bearing spikes from ancestral SARS-CoV-2 (WT), Alpha, Beta, Gamma, Delta, BA.1, BA.4/5, BA.2.75, XBB, XBB.1.5, EG.5, EG.5.1, FL.1.5, FL.1.5.1, HK.3, BA.2.86, JN.1 and SARS-CoV-1 (Figure S1A). Donor A-D show poor or lower neutralizing titers against earlier SARS-CoV-2 strains from WT to BA.1 while higher titers against strains from BA.4/5 to HK.3 (Figure 1B). As Donor A and B were unvaccinated, their patterns represent antibody responses induced by sequential Omicron exposures in the absence of immune imprinting. Despite the fact that Donor C and D were vaccinated, they display similar patterns with Donor A and B, suggesting that immune imprinting is not induced by vaccination in Donor C and D. Donor E-K maintain higher titers against earlier strains from WT to BA.2.75 while lower titers against XBB and later strains, indicating the presence of immune imprinting in these individuals (Figure 1C). Different from above donors, Donor L exhibits a quite chaotic pattern of neutralizing titers and thus is excluded for further analysis (Figure S1B).

We compared neutralizing titers of Donor A-D (referred to as unimprinted group, G1)

and Donor E-K (referred to as imprinted group, G2) (Figure 1D). For early strains from

WT to BA.1, imprinted group has significantly higher titers than unimprinted group. On

the contrary, for more recent Omicron variants, including XBB, XBB.1.5, EG.5, EG.5.1,

FL.1.5 and HK.3, unimprinted group surpasses the imprinted group. Taken together,

these data suggest that immune imprinting is not consistently induced by vaccination.

However, once induced, immune imprinting is not countered by sequential Omicron

infections.

Immune imprinting leads to more frequent cross-reactive antibodies to RBD but

not NTD

To decipher the effects of immune imprinting on human B cell responses elicited by

sequential Omicron infections, we moved on to isolate and characterize monoclonal

antibodies (mAbs) from those donors. As they were sequentially exposed to BA.5/BF.7

and XBB*, we sorted single memory B cells or plasmablasts with XBB.1.5 spike as bait

(Figure S2). After cloning antibody variable regions from single cells into corresponding

antibody-expressing vectors, we transfected HEK293T cells with paired plasmids and

collected supernatant for ELISA and neutralization assay. The supernatant was tested

against spike (S), S1, RBD, NTD and S2 of XBB.1.5 and WT by ELISA, and tested

against XBB.1.5 and WT pseudoviruses by neutralization assay (Figure 2A). In total,

we identified 364 clones positive for XBB.1.5 spike from Donor A, B, C, D, E, F, G, I, J

and K. The proportions of antibodies against each domain vary in a wide range and no

significant differences are observed between unimprinted group (G1) and imprinted

group (G2) (Figure 2B). With 50% as cutoff for neutralization of supernatant, 84 clones were identified as neutralizing antibodies against XBB.1.5 or WT or both (Figure 2A, B). Notably, cross-neutralizing antibodies are all from imprinted group (Figure 2A). In addition, the vast majority of neutralizing antibodies target either RBD or NTD and there are no significant differences between the two groups (Figure 2B).

We further analyzed the frequency of cross-reactive antibodies (XBB.1.5$^+$ WT$^+$) (Figure 2C, D). Most S2-specific antibodies are cross-reactive, which should be due to the conservation of this domain. For NTD-specific antibodies, the frequencies range from 0 to 50% and there is no difference between G1 and G2. In contrast, for RBD-specific antibodies, G2 has remarkably higher frequency than G1, which represents recall of cross-reactive memory B cells induced by vaccination. Overall, these results indicate that immune imprinting mainly influences antibody responses to RBD after sequential Omicron infections.

Immune imprinting facilitates the development of broadly neutralizing antibodies against RBD

To determine the effects of immune imprinting on neutralizing antibodies, we purified RBD- or NTD-specific mAbs with supernatant neutralization above or close to 50%. A total number of 87 mAbs, including 16 RBD-specific and 32 NTD-specific mAbs from G1 and 33 RBD-specific and 6 NTD-specific mAbs from G2, were tested against a panel of SARS-CoV-2 pseudoviruses (Figure 3A). Similar to the neutralization patterns of plasma from G1, RBD-specific mAbs from G1 (G1-RBD) poorly neutralize early strains from WT to BA.1 while potently neutralize later Omicron variants. RBD-specific

mAbs from G2 can be divided into two subgroups according to their neutralization

patterns. One subgroup resembles mAbs from G1-RBD in neutralization pattern and

is named as non-WT-neutralizing (Abbreviated as non-WT-neut). The other subgroup

replicates the neutralization patterns of plasma from G2 and is termed as WT-

neutralizing (Abbreviated as WT-neut). It is worth mentioning that there are several

broadly neutralizing antibodies in this group, including KXD-E1, KXD-F3, KXD-F1,

KXD-G10 and KXD-G5. Most NTD-specific mAbs are able to neutralize BA.4/5, BQ.1,

XBB, XBB.1.5, EG.5.1 and HK.3, and the patterns are similar between G1 and G2.

Regarding to the neutralizing potency (IC50s) against different strains, mAbs from G1-

RBD are more potent to BA.4/5 than later variants whereas mAbs from G2-RBD-non-

WT-neut exhibit similar neutralizing activity against BA.4/5 and later strains (Figure

S3A). Antibodies from G2-RBD-WT-neut have equivalent potency to early strains from

WT to BA.4/5 while decreased potency to later variants (Figure S3A). Different from

RBD-specific mAbs, NTD-specific mAbs from both G1 and G2 are most potent to XBB

and XBB.1.5 (Figure S3B).

We compared the neutralizing potency of mAbs from different groups against the same

strain (Figure 3B,3C). Overall, mAbs from G1-RBD have similar neutralizing potency

with mAbs from G2-RBD-non-WT-neut. However, mAbs from G1-RBD are more potent

than mAbs from G2-RBD-WT-neut against variants from BA.4/5 to JN.1 except

BA.2.75 and XBB. On the other hand, NTD-specific mAbs from both groups have

analogous potency against all tested strains except HK.3. These data further confirm

that antibody responses to RBD while not NTD are affected by immune imprinting.

Moreover, the imprinted RBD-specific antibodies tend to have broadly neutralizing

activity although they generally show suboptimal neutralizing potency against Omicron

variants.

Repeated Omicron infections induce de novo Omicron-specific antibody

responses in the presence of immune imprinting

To examine whether repeated Omicron infections could induce de novo antibody

responses from naïve B cells in the presence of immune imprinting, we move on to

investigate the origins of neutralizing mAbs from each group. In published literatures,

it is generally assumed that cross-reactive antibodies derive from memory B cells

induced by WT whereas omicron-specific antibodies originate from naïve B cells

activated by Omicron(22-25). However, these assumptions remain to be confirmed by

experiments to exclude possibilities that memory B cells induced by WT may lose

cross-reactivity and Omicron-specific naïve B cells may develop cross-reactivity during

affinity maturation. Here, we constructed germline versions of representative mAbs

from each group and tested their binding to a panel of spikes expressed on HEK293T

cells by FACS (Figure 4A). Germline antibodies from G1-RBD, G2-RBD-non-WT-neut,

G1-NTD and G2-NTD bind Omicron spikes but not WT spike. Oppositely, germline

antibodies from G2-RBD-WT-neut mostly recognize WT spike but not Omicron spikes.

These data confirm that mAbs from G1-RBD, G2-RBD-non-WT-neut, G1-NTD and G2-

NTD are de novo antibodies induced by Omicron, whereas mAbs from G2-RBD-WT-

neut are recalled antibodies initially induced by WT.

Correlated with their origins, mAbs from G2-RBD-WT-neut show higher mutation levels

than other mAbs, which reflects extensive affinity maturation sequentially driven by WT

and Omicron (Figure 4B). On the other hand, we also analyzed the repertoire of each

group (Figure 4C and Table 1). In G2-RBD-WT-neut group, IGHV1-18 and IGLV1-44

are most frequent as a consequence of B cell clonal expansion in Donor J. Moreover,

IGHV3-53, IGHV3-66 and IGHV1-69, which have been identified as frequent VH genes

for RBD-specific antibodies from individuals exposed to WT spike(28, 29), are

frequently used in G2-RBD-WT-neut but not in G1-RBD and G2-RBD-non-WT-neut.

For NTD-specific antibodies, IGHV4-39 is most frequently used and is identified from

multiple donors in both groups. Therefore, these IGHV4-39 antibodies can be taken as

a class of public antibody against NTD. In sum, these data suggest that repeated

Omicron exposures could induce de novo antibody responses to RBD and NTD in the

presence of immune imprinting.

Omicron-specific and WT-cross-reactive antibodies recognize similar epitopes

To determine whether de novo antibody responses induced by repeated Omicron

infections recognize distinct epitopes compared with WT-cross-reactive antibodies, we

performed competition ELISA (Figure 5A, 5B). Three broadly neutralizing antibodies

targeting RBD, including BD55-1205, BD56-1854 and BD55-5483 were taken as

reference(30). A major cluster is formed by the majority of RBD-specific antibodies.

Most antibodies in this cluster also compete with BD55-1205, BD56-1854 and ACE2.

Outside the major cluster, a minor cluster is formed by 6 antibodies. They compete

with BD55-5483 but not with BD55-1205, BD56-1854 and ACE2. The rest 3 RBD-

specific antibodies do not exhibit apparent competition with other antibodies or ACE2.

On the other hand, NTD-specific antibodies show competition with each other and form

a single cluster, suggesting that NTD-specific neutralizing antibodies target a supersite.

Taken together, these data provide explanation for the suppressed Omicron-specific

antibody responses to RBD in the presence of immune imprinting.

Discussion

In this study, we compared B cell responses evoked by sequential Omicron infections

in individuals vaccinated or unvaccinated with inactivated or recombinant RBD

vaccines. Unexpectedly, two vaccinees show no hints of immune imprinting, according

to their polyclonal and monoclonal antibody responses against ancestral SARS-CoV-

2 and variants. Therefore, they are classified into unimprinted group along with two

unvaccinated individuals throughout this study. Considering the factor of individual

variations, we believe that this classification is more reasonable than simply grouping

the cohorts based on antigen exposure history, especially for cohorts with limited size.

We speculate that no immune imprinting in these two individuals is probably due to

weak immune responses stimulated by vaccination. Moreover, whether this

phenomenon is common remains to be determined with larger cohorts.

In contrast to individuals without immune imprinting, individuals with immune imprinting

have lower plasma neutralizing antibody titers against multiple Omicron variants after

XBB, along with more frequent WT-cross-reactive antibodies and less frequent

Omicron-specific antibodies to RBD. Therefore, we conclude that immune imprinting

induced by inactivated or recombinant RBD vaccines is not countered by sequential

Omicron infections, which is inconsistent to a study performed by Yisimayi et al(22).

We suppose this contradiction is due to following factors. First, the infections were

sequentially caused by BA.1/BA.2 and BA.5/BF.7 in their study whereas by BA.5/BF.7

and XBB* in our study. We notice that individuals with or without immune imprinting

have equivalent plasma neutralizing antibody titers to BA.4/5 and BA.2.75. Moreover,

individuals with immune imprinting show unaffected neutralizing titers against early

Omicron variants including BA.1, BA.2.75 and BA.4/5 compared with WT. These data

suggest that immune imprinting does not compromise antibody responses to early

Omicron variants after repeated Omicron exposures. Second, all vaccinees were

considered as imprinted in their study while two vaccinees were taken as unimprinted

in our study. As vaccinees without immune imprinting are able to generate abundant

Omicron-specific antibodies, excluding them from imprinted cohort may lead to more

clear results.

A major finding of our study is that antibody responses to NTD are not affected by

immune imprinting. As immunological memory and cross-reactivity are the basis of

immune imprinting(31), one possible explanation for this finding is that NTD is less

immunogenic than RBD because of its extensive glycan shielding(32, 33), and thus

few NTD-specific memory B cells are induced. Another possible explanation is that

NTD-specific memory B cells induced by WT are not cross-reactive to Omicron

variants. However, consistent with previous studies(34-39), our results argue that NTD

is highly immunogenic and able to elicit antibodies cross-reactive to XBB.1.5 and WT,

which meet the preconditions for immune imprinting. Therefore, the mechanisms for

the unimprinted antibody responses to NTD remain to be explored.

Similar to previous studies(15-18, 20, 22-24), we prove that immune imprinting

persistently shapes antibody responses to RBD. As a result, RBD-specific antibodies

can be divided into two groups: WT-cross-reactive antibodies and Omicron-specific

antibodies. Compared with Omicron-specific antibodies, WT-cross-reactive antibodies

tend to have broader neutralization spectrum. For example, KXD-F1 neutralize all

tested pseudoviruses from WT to JN.1 with IC50s ranging from 0.006 to 0.084 μg/ml.

As germlines of those WT-cross-reactive antibodies generally have limited breadth,

there is no doubt that the breadth of WT-cross-reactive antibodies is attributed to

affinity maturation. However, the underlying mechanisms are not fully understood.

According to the model of affinity maturation in GCs, it is possible that WT-cross-

reactive memory B cells become Omicron-specific after affinity maturation. By

comparing the binding specificities of paired germline and mature antibodies, we find

that all WT-reactive germline antibodies consistently recognize WT spike after affinity

maturation, which negate the possibility like our previous study(40). Oppositely, it is

also possible that Omicron-specific naïve B cells may become cross-reactive after

affinity maturation. However, we have not found evidence to support the possibility,

despite that fact that infection or vaccination with WT could elicit broadly neutralizing

antibodies against Omicron variants(30, 41-43). More studies are required to solve

these puzzles on affinity maturation.

Currently, ancestral SARS-CoV-2 immune imprinting has been acknowledged as a

great barrier for vaccine update. According to our findings, we propose that NTD-based

vaccines could be designed to bypass immune imprinting, since NTD is also a major

target of neutralizing antibodies. On the other hand, vaccines containing RBD could

be designed to leverage the potential of immune imprinting in eliciting broadly

neutralizing antibodies.

Materials and methods

Ethics statement

This study was approved by the Ethics Committee of Hefei Institutes of Physical

Science, Chinese Academy of Sciences (Approval Number: YXLL-2023-47). All donors

provided written informed consent for collection of information, analysis of plasma and

PBMCs, and publication of data generated from their samples.

Human samples

Peripheral blood samples were collected from 12 donors (Figure1A). Plasma and

peripheral blood mononuclear cells (PBMCs) were separated from blood by Ficoll

density gradient centrifugation.

Cell lines

HEK293T cells expressing human ACE2 (HEK293T-hACE2) were kindly provided by

Prof. Ji Wang at Sun Yat-Sen University. HEK293T cells were from ATCC. HEK293T

and HEK293T-hACE2 cells were cultured in DMEM with 10% fetal bovine serum (FBS,

VivaCell, Shanghai, China, C04001) and 1% penicillin/streptomycin (pen/strep).

FreeStyle 293F cells (Thermo Fisher Scientific) were cultured in SMM 293-TII

Expression Medium (Sino Biological Inc., M293TII). All cells were maintained in a 37°C

incubator at 5% CO_2

Protein expression and purification

The genes encoding the ectodomain of SARS-CoV-2 spike including WT and XBB.1.5 were constructed with foldon trimerization motif, His tag, tandem strep-tag II and FLAG tag at C-terminal. The genes encoding the S1, NTD and RBD of WT and XBB.1.5 were constructed with a strep-tag II at C-terminal. The genes encoding the S2 of WT and XBB.1.5 were constructed with a His tag at C-terminal. Spike trimer, S1, S2, NTD and RBD were expressed in the FreeStyle 293F cells and purified by Streptactin Agarose Resin 4FF (Yeasen, 20495ES60) or Ni-NTA Agarose (Qiagen, 30210).

Spike-specific Single B cell sorting and PCR

PBMCs were incubated with 200nM XBB.1.5 spike for 30 min at 4°C. After wash, they were stained with cell-surface antibodies: CD3-BV510 (BioLegend, 317331), CD19-PE/Cy7 (BioLegend, 302215), CD27-APC (BioLegend, 356409), CD38-APC/Cy7 (BioLegend, 356615), human IgM-AF700 (BioLegend, 314537), human IgD-perCP/Cy5.5 (BioLegend, 348207), anti-His-AF488 (Proteintech, CL488-66005), anti-FLAG-PE (BioLegend, 637309) and DAPI. The stained cells were washed with FACS buffer (PBS containing 2% FBS) and resuspended in 500µl FACS buffer. Spike-specific single B cells were gated as DAPI$^-$CD3$^-$CD19$^+$CD27$^+$IgD$^-$His$^+$FLAG$^+$ and sorted into 96-well PCR plates containing 4µl lysis buffer (0.5×PBS, 0.1M DTT and RNase inhibitor) per well. After reverse transcription reaction, variable regions of heavy and light chains were amplified by nested PCR and cloned into human IgG1 expression vectors. Plasmids for paired heavy and light chains were co-transfected into HEK293T cells or FreeStyle 293F cells. Antibodies were purified with Protein A magnetic beads

(GenScript, L00273).

ELISA

Spike, S1, S2, NTD or RBD were coated onto 96-well ELISA plates (100 ng/well) and

incubated at 4°C overnight. After blocking with PBS containing 10% FBS, HEK293T

supernatant was added to the wells and incubated at 37°C for 1hr. HRP-conjugated

goat anti-human IgG antibodies (Zen-bio, 550004; 1:2000 dilution) were added to the

wells and incubated at 37°C for 1hr. TMB substrate (Sangon Biotech, E661007-0100)

was added to the wells and incubated at room temperature for 5 mins. The reaction

was stopped by TMB Stop Solution (Sangon Biotech, E661006-0500) and absorbance

at 450 nm was measured.

Pseudovirus neutralization assay

To generate pseudoviruses, HEK293T cells were transfected with psPAX2, pLenti-

luciferase and spike-encoding plasmids using polyetherimide (PEI). Supernatants with

pseudoviruses were collected 48hrs after transfection. 3-fold serially diluted plasma

(starting at 1:20), HEK293T supernatant (starting at 1:4.5), or mAbs (starting at

10μg/ml) were mixed with pseudoviruses at 37°C for 1hr. HEK293T-hACE2 cells

($1.5×10^4$ per well) were added into the mixture and incubated at 37°C for 48hrs. Cells

were lysed to measure luciferase activity by Bright-Lite Luciferase Assay System

(Vazyme Biotech, DD1204-02). The percentages of neutralization were determined by

comparing with the virus control. The plasmids encoding spike of WT, Alpha, Beta,

Gamma, Delta, BA.1, BA.4/5, XBB, XBB.1.5 and SARS-CoV-1 were kindly provided

by Prof. Linqi Zhang at Tsinghua University. The plasmids encoding spike of BA.2.75

was kindly provided by Prof. Zezhong Liu at Fudan University. The sequences of EG.5,

EG.5.1, FL.1.5, FL.1.5.1, HK.3, BA.2.86 and JN.1 spike were synthesized and cloned

into expression vectors.

Competition ELISA

NTD-specific and RBD-specific neutralizing mAbs were labeled with HRP (ProteinTech,

PK20001). XBB.1.5 spike was coated onto 96-well ELISA plates (50 ng/well) and

incubated at 4°C overnight. After blocking with PBS containing 10% FBS, blocking

mAbs were added to the wells and incubated at 37°C for 1hr. HRP-labeled detecting

mAbs were added to the wells in the presence of competing mAbs. The ratio of

blocking antibody and detecting antibody is 100:1. After 1hr incubation at 37°C, TMB

substrate (Sangon Biotech, E661007-0100) was added to the wells and incubated at

room temperature for 30 mins. The reaction was stopped by TMB Stop Solution

(Sangon Biotech, E661006-0500) and absorbance at 450 nm were measured. The

percentages of signal decrease caused by blocking mAbs were calculated.

ACE2 competition FACS

5nM biotinylated XBB.1.5 spike was incubated with 10μg/ml competing mAbs for

30min at 4°C, then the mixtures were added to HEK293T-hACE2 cells. After 30min at

4°C, the cells were washed with FACS buffer. Streptavidin-APC (BioLegend, 405207)

and DAPI were added to the wells at 4°C for 30min. The stained cells were washed

with FACS buffer and resuspended in 200μl FACS buffer. Positive cells were gated as

DAPI$^-$XBB.1.5 spike$^+$. The percentages of signal decrease caused by competing mAbs

were calculated.

Germline antibodies binding FACS

The plasmids encoding spikes of SARS-CoV-2 WT, Alpha, Beta, Gamma, Delta, BA.1, BA.4/5, BF.7, BA.2.75, BQ.1, XBB.1.5, EG.5.1, BA.2.86 were transfected into HEK293T cells. After 48hr, cells were harvested and incubated with 50µg/ml germline antibodies for 30min at 4°C. After washed with FACS buffer, cells were stained with goat anti-human IgG FITC (Proteintech, SA00003-12) and DAPI. The stained cells were washed with FACS buffer and resuspended in 200µl FACS buffer. Positive cells were gated as DAPI$^-$IgG$^+$.

Acknowledgements

The authors thank all donors for providing peripheral blood. This work was supported by the National Key Plan for Scientific Research and Development of China (Grant No. 2022YFC2305800) and the National Natural Science Foundation of China (Grant No. 32200765, Grant No. 32370941).

References

1. D. Gray, Immunological memory. *Annu Rev Immunol* **11**, 49-77 (1993).
2. G. D. Victora, M. C. Nussenzweig, Germinal Centers. *Annu Rev Immunol* **40**, 413-442 (2022).
3. L. Mesin *et al.*, Restricted Clonality and Limited Germinal Center Reentry Characterize Memory B Cell Reactivation by Boosting. *Cell* **180**, 92-106 e111 (2020).
4. T. Francis, On the Doctrine of Original Antigenic Sin. *Proceedings of the American Philosophical Society* **104**, 572-578 (1960).
5. R. Viana *et al.*, Rapid epidemic expansion of the SARS-CoV-2 Omicron variant in southern Africa. *Nature* **603**, 679-686 (2022).
6. Q. He *et al.*, An updated atlas of antibody evasion by SARS-CoV-2 Omicron sub-variants including BQ.1.1 and XBB. *Cell Rep Med* **4**, 100991 (2023).
7. C. Kurhade *et al.*, Low neutralization of SARS-CoV-2 Omicron BA.2.75.2, BQ.1.1 and XBB.1 by parental mRNA vaccine or a BA.5 bivalent booster. *Nat Med* **29**, 344-347 (2023).
8. Q. Wang *et al.*, Alarming antibody evasion properties of rising SARS-CoV-2 BQ and XBB subvariants. *Cell* **186**, 279-286 e278 (2023).
9. J. Miller *et al.*, Substantial Neutralization Escape by SARS-CoV-2 Omicron Variants BQ.1.1

and XBB.1. *N Engl J Med* **388**, 662-664 (2023).

10. Q. Wang *et al.*, Antibody neutralisation of emerging SARS-CoV-2 subvariants: EG.5.1 and
XBC.1.6. *Lancet Infect Dis* **23**, e397-e398 (2023).

11. S. Yang *et al.*, Fast evolution of SARS-CoV-2 BA.2.86 to JN.1 under heavy immune
pressure. *Lancet Infect Dis* **24**, e70-e72 (2024).

12. S. Chalkias *et al.*, Safety, immunogenicity and antibody persistence of a bivalent Beta-
containing booster vaccine against COVID-19: a phase 2/3 trial. *Nat Med* **28**, 2388-2397
(2022).

13. S. Chalkias *et al.*, A Bivalent Omicron-Containing Booster Vaccine against Covid-19. *N
Engl J Med* **387**, 1279-1291 (2022).

14. A. Y. Collier *et al.*, Immunogenicity of BA.5 Bivalent mRNA Vaccine Boosters. *N Engl J Med*
388, 565-567 (2023).

15. Y. J. Park *et al.*, Imprinted antibody responses against SARS-CoV-2 Omicron sublineages.
Science **378**, 619-627 (2022).

16. Z. Wang *et al.*, Memory B cell responses to Omicron subvariants after SARS-CoV-2 mRNA
breakthrough infection in humans. *J Exp Med* **219** (2022).

17. C. I. Kaku *et al.*, Recall of preexisting cross-reactive B cell memory after Omicron BA.1
breakthrough infection. *Sci Immunol* **7**, eabq3511 (2022).

18. J. Quandt *et al.*, Omicron BA.1 breakthrough infection drives cross-variant neutralization
and memory B cell formation against conserved epitopes. *Sci Immunol* **7**, eabq2427
(2022).

19. T. Weber *et al.*, Enhanced SARS-CoV-2 humoral immunity following breakthrough
infection builds upon the preexisting memory B cell pool. *Sci Immunol* **8**, eadk5845 (2023).

20. A. Addetia *et al.*, Neutralization, effector function and immune imprinting of Omicron
variants. *Nature* **621**, 592-601 (2023).

21. Q. Wang *et al.*, Deep immunological imprinting due to the ancestral spike in the current
bivalent COVID-19 vaccine. *Cell Rep Med* **4**, 101258 (2023).

22. A. Yisimayi *et al.*, Repeated Omicron exposures override ancestral SARS-CoV-2 immune
imprinting. *Nature* **625**, 148-156 (2024).

23. T. S. Johnston *et al.*, Immunological imprinting shapes the specificity of human antibody
responses against SARS-CoV-2 variants. *Immunity* **57**, 912-925 e914 (2024).

24. M. A. Tortorici *et al.*, Persistent immune imprinting occurs after vaccination with the
COVID-19 XBB.1.5 mRNA booster in humans. *Immunity* **57**, 904-911 e904 (2024).

25. C. Y. Liang *et al.*, Imprinting of serum neutralizing antibodies by Wuhan-1 mRNA vaccines.
Nature 10.1038/s41586-024-07539-1 (2024).

26. Y. Pan *et al.*, Characterisation of SARS-CoV-2 variants in Beijing during 2022: an
epidemiological and phylogenetic analysis. *Lancet* **401**, 664-672 (2023).

27. Y. Feng *et al.*, Genomic Surveillance for SARS-CoV-2 Variants: Dominance of XBB
Replacement—China, January–June 2023. *China CDC Weekly* **6**, 324-331 (2024).

28. Y. Wang *et al.*, A large-scale systematic survey reveals recurring molecular features of
public antibody responses to SARS-CoV-2. *Immunity* **55**, 1105-1117 e1104 (2022).

29. Q. Yan *et al.*, Shared IGHV1-69-encoded neutralizing antibodies contribute to the
emergence of L452R substitution in SARS-CoV-2 variants. *Emerg Microbes Infect* **11**,
2749-2761 (2022).

30.   Y. Cao *et al.*, Imprinted SARS-CoV-2 humoral immunity induces convergent Omicron RBD
evolution. *Nature* **614**, 521-529 (2023).

31.   M. Koutsakos, A. H. Ellebedy, Immunological imprinting: Understanding COVID-19.
Immunity **56**, 909-913 (2023).

32.   A. C. Walls *et al.*, Structure, Function, and Antigenicity of the SARS-CoV-2 Spike
Glycoprotein. *Cell* **181**, 281-292 e286 (2020).

33.   D. Wrapp *et al.*, Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation.
Science **367**, 1260-1263 (2020).

34.   X. Chi *et al.*, A neutralizing human antibody binds to the N-terminal domain of the Spike
protein of SARS-CoV-2. *Science* **369**, 650-655 (2020).

35.   L. Liu *et al.*, Potent neutralizing antibodies against multiple epitopes on SARS-CoV-2 spike.
Nature **584**, 450-456 (2020).

36.   M. McCallum *et al.*, N-terminal domain antigenic mapping reveals a site of vulnerability
for SARS-CoV-2. *Cell* **184**, 2332-2347 e2316 (2021).

37.   N. Suryadevara *et al.*, Neutralizing and protective human monoclonal antibodies
recognizing the N-terminal domain of the SARS-CoV-2 spike protein. *Cell* **184**, 2316-
2331 e2315 (2021).

38.   Z. Wang *et al.*, Analysis of memory B cells identifies conserved neutralizing epitopes on
the N-terminal domain of variant SARS-Cov-2 spike proteins. *Immunity* **55**, 998-1012
e1018 (2022).

39.   G. Cerutti *et al.*, Neutralizing antibody 5-7 defines a distinct site of vulnerability in SARS-
CoV-2 spike N-terminal domain. *Cell Rep* **37**, 109928 (2021).

40.   L. Li *et al.*, Breakthrough infection elicits hypermutated IGHV3-53/3-66 public antibodies
with broad and potent neutralizing activity against SARS-CoV-2 variants including the
emerging EG.5 lineages. *PLoS Pathog* **19**, e1011856 (2023).

41.   B. Ju *et al.*, Infection with wild-type SARS-CoV-2 elicits broadly neutralizing and protective
antibodies against omicron subvariants. *Nat Immunol* **24**, 690-699 (2023).

42.   Y. Liu *et al.*, Inactivated vaccine-elicited potent antibodies can broadly neutralize SARS-
CoV-2 circulating variants. *Nat Commun* **14**, 2179 (2023).

43.   B. Ju *et al.*, Omicron BQ.1.1 and XBB.1 unprecedentedly escape broadly neutralizing
antibodies elicited by prototype vaccination. *Cell Rep* **42**, 112532 (2023).

Figures and tables

Figure 1. Sequential Omicron infections could not overcome ancestral SARS-

CoV-2 immune imprinting induced by inactivated or recombinant RBD vaccines.

(A) Vaccination and infection information of 12 donors. (B, C) Summary of plasma

neutralizing antibody titers (NT50). For plasma with no neutralizing activity at 20-fold

dilution, a number between 10–20 is given as titer to separate the curves. The numbers

below the line are geometric mean titers against the pseudoviruses. The titers against

WT are compared with other variants and the fold changes are indicated above the

line. For panel B, statistical analysis was performed by two-tailed paired T test. For

panel C, statistical analysis was performed by two-tailed Wilcoxon matched-pairs

signed rank test. *P < 0.05, **P < 0.01, **** P < 0.0001. (D) Comparison of neutralizing

antibody titers of G1 (Donor A-D, unimprinted group) and G2 (Donor E-K, imprinted

group). The numbers on top are fold changes. Statistical analysis was performed by

two-tailed unpaired T test. *P < 0.05, **P < 0.01. The neutralizing titers are calculated

with results from two independent experiments, in which duplicates are performed.

Figure 2. Immune imprinting leads to more frequent cross-reactive antibodies to RBD but not NTD. (A) ELISA and neutralization results of 364 mAbs. Supernatant of HEK293T cells was tested against Spike, S1, S2, NTD and RBD of WT and XBB.1.5 by ELISA, and tested against WT and XBB.1.5. pseudoviruses by neutralization assay. For ELISA, supernatant was not diluted. For neutralization, supernatant was diluted by 4.5-fold. The results are mean values of two independent experiments. For neutralization, duplicates are performed in each experiment. (B) Frequency of binding mAbs and neutralizing mAbs against different domains in each individual. (C) Frequencies of WT-cross-reactive antibodies (XBB.1.5+ WT+) and Omicron-specific antibodies (XBB.1.5+ WT-) in each individual. (D) Comparison of WT-cross-reactive antibodies between G1 and G2. Data are represented as mean ± SD. Statistical analysis was performed by two-tailed unpaired T test. NS (Not significant) p > 0.05, **P < 0.01.

Figure 3. Immune imprinting facilitates the development of broadly neutralizing

antibodies against RBD. (A) IC50s of mAbs against WT, Alpha, Beta, Gamma, Delta,

BA.1, BA.4/5, BA.2.75, BQ.1, XBB, XBB.1.5, EG.5.1, HK.3, BA.2.86 and JN.1. The

highest antibody concentration used to determine IC50 is 10μg/ml. (B) Comparing

IC50s of RBD-specific mAbs from three groups. (C) Comparing IC50s of NTD-specific

mAbs from two groups. Data are represented as the mean ± SD. Statistical analysis

was performed by two-tailed unpaired T test. NS (Not significant) p > 0.05, *P < 0.05,

**P < 0.01. IC50s are calculated with results from two independent experiments, in

which duplicates are performed.

Figure 4. Repeated Omicron infections induce de novo Omicron-specific antibody responses in the presence of immune imprinting. (A) Binding specificities of germline antibodies measured by FACS. HEK293T cells expressing SARS-CoV-2 spikes were incubated with 50µg/ml germline antibodies and analyzed by FACS. The percentages of positive cells are calculated. The results are mean values of two independent experiments. (B) Somatic hypermutations of VH and VL. DNA sequences were analyzed. Data are represented as mean ± SD. Statistical analysis was performed by two-tailed unpaired T test. NS (Not significant) p > 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001. (C) Frequencies of V genes in each group. Frequently used V genes are highlighted by colored background.

Figure 5. Omicron-specific and WT-cross-reactive antibodies recognize similar

epitopes. (A) Competition ELISA of RBD-specific mAbs. Competition with ACE2 was

performed by FACS. (B) Competition ELISA of NTD-specific mAbs. The results are

mean values of two independent experiments.

Figure S1. Plasma neutralizing antibody titers. (A) Measurement of neutralizing

antibody titers against a panel of pseudoviruses. Data are represented as mean ± SD.

(B) Neutralizing antibody titers of donor L.

Figure S2. FACS plots representing gating strategies for XBB.1.5 spike-specific

single B cell sorting.

Figure S3. Neutralizing potency of mAbs against different variants. (A) IC50s of

RBD-specific mAbs. The numbers below the line are geometric mean IC50 of mAbs

against the pseudoviruses. IC50s against BA.4/5 are compared with other variants. (B)

IC50s of NTD-specific mAbs. The numbers below the line are geometric mean IC50 of

mAbs against the pseudoviruses. IC50s against XBB are compared other variants.

Statistical analysis was performed by two-tailed Wilcoxon matched-pairs signed rank

563   test. NS (Not significant) p > 0.05, *P < 0.05, **P < 0.01, ***P < 0.001, ****P < 0.0001.

Table S1. Origins, targets and sequence information of 87 neutralizing mAbs.

mAbs	Donor	Target	VH V	VH D	VH J	VH CDR3	VH CDR3 (aa)	VH SHM(%)	VL V	VL J	VL CDR3	VL CDR3 (aa)	VL SHM(%)
KXD-A10	A	NTD	IGHV4-39	IGHD5-12	IGHJ4	ARLRDWDIVAGGGVFDY	17	6.7	IGLV3-1	IGLJ3	QSWDSSTAV	9	2.9
KXD-A21	A	RBD	IGHV1-24	IGHD6-13	IGHJ6	TTDRLGQQLAKGYYYYGMDV	20	2.7	IGLV3-1	IGLJ2	QAWDRTTPV	9	2.8
KXD-A22	A	NTD	IGHV4-59	IGHD2-15	IGHJ6	ARGGYDYYGMDV	12	5.9	IGLV1-40	IGLJ2	QSFDNSLSVHVL	12	3.4
KXD-A23	A	RBD	IGHV3-15	IGHD3-22	IGHJ4	KTEFGTYYYDTNGYDY	16	3.0	IGLV3-1	IGLJ2	LAWDRSTGV	9	3.6
KXD-A24	A	NTD	IGHV4-39	IGHD6-13	IGHJ4	ARQRVGSSWYYSDS	14	6.4	IGLV1-44	IGLJ1	AAWDDSLNGSYV	12	2.0
KXD-A25	A	RBD	IGHV3-43	IGHD2-15	IGHJ4	ATSEVDYFEY	10	3.8	IGKV4-1	IGKJ2	QQYYSTPPYT	10	3.0
KXD-A31	A	NTD	IGHV4-39	IGHD2-15	IGHJ4	ARDSPLGFCSASSCYFVDS	19	3.3	IGLV1-36	IGLJ1	AAWDDSLNGFV	11	3.4
KXD-A33	A	NTD	IGHV4-38	IGHD2-15	IGHJ5	ARQDAGSTQFDL	12	6.1	IGLV1-40	IGLJ3	QSYDGSLSGWV	11	2.3
KXD-A34	A	NTD	IGHV3-49	IGHD3-9	IGHJ6	TREDPRYNFDWLPSGMDV	18	4.0	IGKV1-39	IGKJ2	QQSYSTLYT	9	2.9
KXD-A43	A	NTD	IGHV4-39	IGHD5-12	IGHJ4	ARLREWDIVAGGGVFDY	17	6.1	IGLV3-1	IGLJ3	QSWDSSTAV	9	2.9
KXD-A48	A	NTD	IGHV4-39	IGHD2-15	IGHJ4	TRDSPLGHCGGSSCYYVGY	19	6.4	IGLV1-36	IGLJ1	AAWDDSLNGYV	11	3.1
KXD-A49	A	RBD	IGHV1-18	IGHD3-3	IGHJ4	ARDRGFWSGFDS	12	3.1	IGLV6-57	IGLJ2	QSYDSGIVV	9	2.8
KXD-A5	A	NTD	IGHV1-18	IGHD1-26	IGHJ4	ARVGESSGYSRSWRNEFDY	19	5.5	IGKV2-40	IGKJ1	MQRLDSPWT	9	4.6
KXD-A51	A	NTD	IGHV2-5	IGHD3-22	IGHJ4	ARSSPEIYYYDSSDYYQYYFDY	22	5.6	IGLV1-47	IGLJ2	AAWDDSLSGPYV	12	1.0
KXD-A52	A	NTD	IGHV3-49	IGHD3-9	IGHJ6	TRGDPRYNFDWLPDGMDV	18	2.3	IGKV1-39	IGKJ2	QQSYSTLYT	9	2.5
KXD-A56	A	NTD	IGHV3-49	IGHD3-9	IGHJ6	SREDPRYNFDWVPDGMDV	18	6.3	IGKV1-39	IGKJ2	QQSYSTLYT	9	7.5
KXD-A57	A	NTD	IGHV4-38	IGHD2-15	IGHJ5	ARQDAGSTQFDP	12	7.4	IGLV1-40	IGLJ3	QSYDGRLSGWV	11	3.1
KXD-A60	A	NTD	IGHV4-39	IGHD5-12	IGHJ4	ARLREWDIVAGGGVFDY	17	6.0	IGLV3-1	IGLJ3	QSWDSSTAV	9	1.8
KXD-A63	A	NTD	IGHV4-39	IGHD6-13	IGHJ4	ARERVGSSWYYSDY	14	1.3	IGLV1-44	IGLJ1	AAWDDGLNGSNV	12	0.7
KXD-A67	A	NTD	IGHV4-39	IGHD2-15	IGHJ4	TRDSPLGYCSGSSCYFVDY	19	5.4	IGLV1-36	IGLJ1	ATWDDSLNGYV	11	2.4
KXD-A69	A	RBD	IGHV1-24	IGHD5-24	IGHJ6	TTNGLQQKAVGVYYYYYGLDV	21	4.7	IGLV3-1	IGLJ2	QAWDSSTAV	9	2.2
KXD-A70	A	NTD	IGHV4-39	IGHD3-10	IGHJ5	ARMVQPENWFDP	12	7.4	IGLV1-36	IGLJ2	AAWDHSLNGVI	11	2.7
KXD-A77	A	NTD	IGHV1-18	IGHD6-13	IGHJ4	ARVGESSGYSRSWRNEIDY	19	5.8	IGKV2-40	IGKJ1	MQRIEFPWT	9	2.6
KXD-A78	A	NTD	IGHV4-39	IGHD2-15	IGHJ4	TRDSPLGFCRGGSCYFVDY	19	3.4	IGLV1-36	IGLJ1	AAWDDSLNAYV	11	1.7
KXD-A8	A	RBD	IGHV3-74	IGHD3-22	IGHJ6	ARFSGLSGGYYYGMDV	16	2.7	IGLV6-57	IGLJ2	QSYDNSDLV	9	4.2
KXD-A82	A	NTD	IGHV4-39	IGHD2-15	IGHJ4	ARDSPLGSCSDSSCYFVDS	19	8.2	IGLV1-36	IGLJ1	AAWDDNLNGFV	11	2.4
KXD-A84	A	NTD	IGHV4-59	IGHD2-15	IGHJ6	ARGGYDYYGMDV	12	3.8	IGLV1-40	IGLJ2	QSYDDSLSVHVV	12	3.1
KXD-B6	B	NTD	IGHV3-30	IGHD2-2	IGHJ6	AGEGEDGFCSSLSCYSSPGGV	21	8.1	IGKV2-30	IGKJ4	MQGTHWPPLT	10	3.6
KXD-C10	C	NTD	IGHV4-39	IGHD3-9	IGHJ4	ARVVYYDILTGQKTFDY	17	5.1	IGLV1-51	IGLJ3	GTWDSSLSAGV	11	3.0
KXD-C12	C	NTD	IGHV4-39	N/A	IGHJ5	ARQLPTENWFDP	12	7.0	IGLV3-21	IGLJ2	QVWDNRSHHLVS	12	3.9
KXD-C14	C	NTD	IGHV3-33	IGHD3-3	IGHJ4	ARDSSYYDLWSGLDY	15	5.4	IGKV1-16	IGKJ2	QQYYSFPYT	9	5.3
KXD-C15	C	RBD	IGHV5-51	IGHD5-18	IGHJ4	ARPTSTDTASDY	12	5.1	IGKV1-39	IGKJ1	QQSYSPLWT	9	2.6
KXD-C18	C	NTD	IGHV3-33	IGHD1-26	IGHJ6	ARESEDSGGPGI	12	6.1	IGKV2-28	IGKJ2	MQALQTPRCT	10	1.3
KXD-C19	C	NTD	IGHV3-7	IGHD2-15	IGHJ6	ARERFCSAGSCYSFLGRRVYYYDGMDV	27	2.7	IGKV1-39	IGKJ1	QQSYSIPPT	9	4.7
KXD-C21	C	RBD	IGHV3-21	N/A	IGHJ3	ARGHAEEGDAFDT	13	5.8	IGKV1-5	IGKJ1	QQYNGFPWT	9	3.6
KXD-C23	C	RBD	IGHV3-48	IGHD6-13	IGHJ5	ARDPVCGYSSSCPLFGS	17	4.7	IGLV1-40	IGLJ3	QSYDSSLSGNWV	12	4.7
KXD-C24	C	RBD	IGHV5-51	IGHD3-9	IGHJ4	ARLPSVGHFDWGYFDY	16	6.0	IGKV3-20	IGKJ1	QQYGGSPPWT	10	5.2
KXD-C25	C	NTD	IGHV4-30	IGHD3-3	IGHJ5	ARTIFGDPNDWFDP	14	5.4	IGLV1-40	IGLJ2	QSYDTSLNGWDVV	13	2.0
KXD-C26	C	NTD	IGHV4-39	IGHD6-13	IGHJ4	ARHDFDIATADTSFHY	16	3.7	IGLV3-1	IGLJ3	QTWDNSAVV	9	6.9
KXD-C27	C	NTD	IGHV4-39	N/A	IGHJ5	ARQTPPNCFDP	11	2.0	IGLV3-21	IGLJ3	QVWDNGNDHPGV	12	3.2
KXD-C28	C	RBD	IGHV3-66	IGHD2-21	IGHJ5	ARGPFRKFDP	10	6.9	IGKV1-33	IGKJ2	QQYDNIPYT	9	3.9
KXD-C29	C	NTD	IGHV4-39	IGHD2-2	IGHJ4	ARHDFDIVTANTIFHY	16	5.1	IGLV3-1	IGLJ2	QTWDSSAVV	9	4.6
KXD-C31	C	NTD	IGHV4-39	IGHD7-27	IGHJ6	ARSGDSLFYYYYYYMDV	17	5.7	IGKV4-1	IGKJ4	QQYYSVPLT	9	4.4
KXD-C36	C	NTD	IGHV4-39	IGHD3-9	IGHJ5	ARHVDYDIVTDSGRFDP	17	10.0	IGLV3-1	IGLJ3	QTWDSGSGV	9	3.6
KXD-C4	C	RBD	IGHV3-30	IGHD5-18	IGHJ4	AREHESETAVDK	12	5.1	IGKV1-12	IGKJ4	QQANNFHLT	9	5.3
KXD-C40	C	NTD	IGHV4-39	IGHD2-2	IGHJ4	ARHDFDIVTANTIFHY	16	6.4	IGLV3-1	IGLJ2	QTWDSSAVV	9	4.9
KXD-C44	C	RBD	IGHV3-11	IGHD6-13	IGHJ1	ASLGYSTIWDGSAEYFQH	18	2.7	IGKV1-39	IGKJ4	QQSYSTPRT	9	0.7
KXD-D3	D	RBD	IGHV4-4	IGHD1-14	IGHJ6	ARDGPHTTSPQDDYYYMDV	19	6.4	IGKV2-29	IGKJ5	MQGRLSPRT	9	5.3
KXD-E1	E	RBD	IGHV3-53	IGHD5-18	IGHJ4	ARLGGYGRKDYDH	13	7.2	IGKV1-33	IGKJ2	HQYDNVPPT	9	5.6
KXD-E2	E	RBD	IGHV1-69	IGHD2-2	IGHJ3	AHTPLPSPCHADALDI	16	13.1	IGKV1-39	IGKJ3	QQSHGSPFT	9	9.2
KXD-E3	E	RBD	IGHV1-69	IGHD4-17	IGHJ3	ARERVDYGDRLLNGSAFDI	19	9.8	IGKV3-20	IGKJ3	QLCGGSAPT	9	6.2
KXD-E4	E	RBD	IGHV3-53	IGHD1-26	IGHJ4	VKMEWDLL	8	6.9	IGKV1-5	IGKJ1	QQYIVYPWT	9	6.1
KXD-F1	F	RBD	IGHV4-34	IGHD3-10	IGHJ4	ARGYRDYGRGMFPLGN	16	11.3	IGKV3-11	IGKJ1	QQRDNWPPT	9	5.0
KXD-F10	F	RBD	IGHV4-30	IGHD3-9	IGHJ6	ARANLLTFHVDV	12	8.4	IGLV6-57	IGLJ2	QSYDGSKVV	9	3.2
KXD-F12	F	RBD	IGHV4-30	IGHD5-12	IGHJ5	ARLRLLVNWFDP	12	4.0	IGLV6-57	IGLJ3	QSYDGSNVV	9	2.1
KXD-F3	F	RBD	IGHV3-9	IGHD1-1	IGHJ4	AKDFHNNWYPGRFDF	15	2.0	IGKV1-33	IGKJ5	QQYDNVPL	8	3.8
KXD-F6	F	RBD	IGHV3-33	IGHD2-2	IGHJ6	AKDMAYYLLVPEARPPYGMDV	21	2.7	IGKV3-20	IGKJ3	HQYGGSPFT	9	4.3
KXD-G10	G	RBD	IGHV4-30	IGHD7-27	IGHJ4	ARALGIIDY	9	8.1	IGKV1-NL1	IGKJ5	QQYYSTPT	8	4.9
KXD-G23	G	RBD	IGHV4-4	IGHD3-10	IGHJ4	ARARGIIDY	9	10.5	IGKV1-NL1	IGKJ4	QQYYSTPS	8	5.1
KXD-G26	G	RBD	IGHV1-69	IGHD2-2	IGHJ4	ATNAPYCTSTACFNF	15	10.3	IGLV1-40	IGLJ2	QSYDSGLSASV	11	7.7
KXD-G32	G	NTD	IGHV4-39	IGHD1-26	IGHJ5	ARLAWDFALSFHWFDP	16	12.5	IGLV2-11	IGLJ2	CSYGGADSWV	10	3.3
KXD-G33	G	RBD	IGHV1-24	IGHD3-16	IGHJ4	AAGGLHMSLGIMPFDY	16	8.5	IGLV1-47	IGLJ3	ASWDDSLSGRGQGV	14	7.8
KXD-G44	G	NTD	IGHV4-39	IGHD3-3	IGHJ4	ARHPAHDFWSGYFDN	15	7.1	IGKV3-15	IGKJ5	QQYNEWPRFS	10	2.9
KXD-G5	G	RBD	IGHV4-39	IGHD7-27	IGHJ4	ARTLGMFDH	9	8.1	IGKV1-NL1	IGKJ3	QQYYDTPFT	9	5.8
KXD-G6	G	RBD	IGHV4-4	IGHD6-13	IGHJ4	TSVLGTFDY	9	12.2	IGKV1-NL1	IGKJ5	QQYYTTRT	8	6.0
KXD-I29	I	RBD	IGHV3-9	IGHD3-10	IGHJ5	TKGQVGELPFPLDS	14	8.5	IGLV1-44	IGLJ2	AAWDDTLNSLV	11	4.7
KXD-I39	I	NTD	IGHV3-15	IGHD3-9	IGHJ4	NTERDNLNFGDY	12	5.6	IGKV1-33	IGKJ2	QQYANLPRT	9	5.6
KXD-I42	I	RBD	IGHV3-9	IGHD3-10	IGHJ4	AKGQLGEFPFPVDF	14	5.8	IGLV1-44	IGLJ3	AVWDDRVNGLV	11	7.1
KXD-J16	J	RBD	IGHV1-46	IGHD4-11	IGHJ5	ARGVYSNYEDWFDP	14	3.4	IGKV4-1	IGKJ5	QQYYTSPLT	9	1.3
KXD-J21	J	RBD	IGHV1-46	IGHD4-11	IGHJ4	ARGVYSNYEDWFDP	14	3.1	IGKV4-1	IGKJ4	QQYYTTPLT	9	3.0
KXD-J23	J	RBD	IGHV3-33	IGHD3-3	IGHJ2	AREGYDFWGDSPENWYFDF	19	4.7	IGLV2-11	IGLJ1	CSYTGTYTYV	10	3.8
KXD-J28	J	RBD	IGHV1-18	IGHD2-2	IGHJ6	GRVESHIVLVPSAGPYYYYMDV	22	11.6	IGLV1-44	IGLJ1	AAWDDSLNSYV	11	8.1
KXD-J33	J	NTD	IGHV3-33	IGHD4-17	IGHJ4	ARDFYWSQADDYGDPHWPGG	20	4.1	IGLV3-10	IGLJ2	YSADGTAYHRGV	12	4.8
KXD-J37	J	RBD	IGHV3-53	IGHD3-16	IGHJ4	TRVLPGGQYFDY	12	9.9	IGKV3-20	IGKJ2	QQYESPAMYI	10	4.0
KXD-J39	J	RBD	IGHV3-33	IGHD3-3	IGHJ2	AKEGYDFWGADPENWYFDL	19	0.7	IGLV2-11	IGLJ1	CSYAGPYTYV	10	4.7
KXD-J42	J	RBD	IGHV1-18	IGHD2-2	IGHJ6	ARVDSHIVLVTAAIPDYYYMDV	22	9.5	IGLV1-44	IGLJ1	AAWDVSLNGYV	11	8.8
KXD-J44	J	RBD	IGHV1-18	IGHD2-21	IGHJ6	ARVNSHIVMVASATPYYYYMDV	22	11.3	IGLV1-44	IGLJ1	AAWDASLNAYV	11	7.5
KXD-J47	J	RBD	IGHV1-18	IGHD2-21	IGHJ6	ARVDSHIVAVPAAIPDYFYMDV	22	11.5	IGLV1-44	IGLJ1	AGWVHTLNGYV	11	11.2
KXD-J5	J	RBD	IGHV1-18	IGHD2-21	IGHJ6	ARVDSHIVMVASAIPYYYHLDA	22	9.9	IGLV1-44	IGLJ1	AAWDASLNAYV	11	7.9
KXD-J6	J	RBD	IGHV1-18	IGHD2-2	IGHJ6	ARVDSHIVSVPSAIPFYYYMDV	22	9.6	IGLV1-44	IGLJ1	AAWDASLNAYV	11	3.4
KXD-J8	J	RBD	IGHV3-33	IGHD3-3	IGHJ2	AKEGYDFWGADPEDWYFDL	19	2.4	IGLV2-11	IGLJ1	CSYAGPYTYV	10	4.2
KXD-K19	K	RBD	IGHV3-66	IGHD3-16	IGHJ4	ARDLLEAGGTDY	12	6.5	IGKV3-20	IGKJ3	QFYSTIAVT	9	5.9
KXD-K20	K	RBD	IGHV3-9	IGHD3-22	IGHJ4	AKAGDASGYYSDFVY	15	5.1	IGLV2-11	IGLJ2	CSFAGGYTWV	10	2.7
KXD-K29	K	RBD	IGHV3-9	IGHD2-21	IGHJ4	ARLSVGTASTYFDY	14	6.1	IGLV1-44	IGLJ3	AAWDDSLNGHWV	12	2.0
KXD-K3	K	RBD	IGHV3-66	IGHD3-22	IGHJ4	TRDLGGYYDQ	10	9.9	IGKV1-5	IGKJ1	QQYNTDSRT	9	3.9
KXD-K40	K	RBD	IGHV3-66	IGHD3-22	IGHJ4	ARDLGGYFDH	10	8.9	IGKV1-5	IGKJ1	QQYSTDSRT	9	5.6
KXD-K48	K	NTD	IGHV4-39	IGHD5-12	IGHJ5	ARHVWDIALRKDWFDP	16	7.1	IGKV3-20	IGKJ1	QQYGSSRT	8	2.1