Energy Deficit is a Key Driver of Sleep Homeostasis

Scarlet J. Park[1-3], Keith R. Murphy[1,4,5], William W. Ja[1,2,*]

[1] Department of Neuroscience, The Herbert Wertheim UF Scripps Institute for Biomedical Innovation & Technology, Jupiter, FL 33458, USA

[2] Skaggs Graduate School of Chemical and Biological Sciences, Jupiter, FL 33458, USA

[3] Current affiliation: Nova Southeastern University, Palm Beach Gardens, FL 33410, USA

[4] Integrative Biology and Neuroscience Program, Florida Atlantic University, Jupiter FL 33458, USA

[5] Current affiliation: Department of Psychiatry and Behavioral Sciences, Stanford University, Stanford, CA 94305, USA

*Correspondence: w.ja@ufl.edu (W.W.J.)

Abstract

Sleep and feeding are vital homeostatic behaviors, and disruptions in either can result in substantial metabolic consequences. Distinct neuronal manipulations in *Drosophila* can dissociate sleep loss from subsequent homeostatic rebound, offering an optimal platform to examine the precise interplay between these fundamental behaviors. Here, we investigate concomitant changes in sleep and food intake in individual animals, as well as respiratory metabolic expenditure, that accompany behavioral and genetic manipulations that induce sleep loss in *Drosophila melanogaster*. We find that sleep disruptions resulting in energy deficit through increased metabolic expenditure and manifested as increased food intake were consistently followed by rebound sleep. In contrast, "soft" sleep loss, which does not induce rebound sleep, is not accompanied by increased metabolism and food intake. Our results demonstrate that homeostatic sleep rebound is linked to energy deficit accrued during sleep loss. Collectively, these findings support the notion that sleep functions to conserve energy and highlight the need to examine the effects of metabolic therapeutics on sleep.

Introduction

Mindless grazing following a sleepless night is a prevalent phenomenon in modern society[1,2]. Sleep and eating—typically mutually exclusive behaviors that are amongst those most vital for survival and prosperity—are intricately linked. Food deprivation increases activity levels in many species, including flies[3], whereas food ingestion temporarily elevates immediate sleepiness—a phenomenon colloquially known as "food coma"—in flies[4], laboratory rodents[5,6], and humans[7,8]. Conversely, acute and chronic sleep deprivation is associated with altered taste perception, food cravings, increased appetite and food intake, rapid weight gain, and detrimental metabolic changes in laboratory rodents[9] and humans[10].

Sleep homeostasis has been hypothesized to serve as an adaptive response to energy expenditure, purposed towards reducing and recovering from energy use[11]. This is consistent with findings that the intensity of neural activity during wakefulness may contribute to sleep drive[12]. This may also explain the greater sleep need of nematodes during energetically expensive molting periods[13]. In at least some mammals, brown fat metabolism is necessary for sleep rebound, further supporting the idea that energy expenditure generates sleep-promoting signals[14]. If this model is correct, then "hard" sleep loss—i.e. homeostatic sleep loss that accompanies a subsequent, compensatory increase in sleep (as opposed to "soft" sleep loss, which is not succeeded by sleep rebound)—should be associated with increased metabolic activity and/or hunger.

Sleep curtailment has been consistently linked to increased food intake[1,15,16] and reduced energy expenditure[17]. However, understanding the precise mechanisms that drive changes in feeding and metabolism, and that regulate the sequence of events

following sleep disruption, remains challenging due to the complex relationship between these factors. Experimental isolation of specific variables is particularly difficult in observational human sleep studies. Development of an effective model system for controlled manipulations and measurements of sleep and feeding would enable research on the regulatory pathways linking these processes.

Studies on the characteristics of sleep and its genetic and neuronal control have yielded diverse methods to manipulate sleep in *Drosophila melanogaster*[15,18,19]. These tools, combined with the scalability and genetic tractability of the model, provide unique opportunities for studying homeostatic sleep regulation and the underlying circuit logic[19,20]. Despite recent advances in understanding the interaction between sleep regulation and other homeostatic processes in flies, including metabolism and food intake[10], studies on the relationship between sleep and feeding behavior have typically relied on measurements of each behavior in separate groups of animals, mainly due to the lack of techniques that enable paired measurements. We have previously shown that flies exhibit postprandial sleep in the Activity Recording CAFE (ARC), an assay that allows concurrent, longitudinal measurements of sleep and food intake in freely-moving animals[4]. The ARC can thus account for individual variability, making correlational studies more powerful and offering more in-depth insights into the relationship between short-, intermediate-, and long-term food intake and sleep.

To investigate the link between sleep deprivation, energy balance, and sleep rebound, we measured changes in sleep, feeding, and metabolic expenditure with different methods of sleep restriction, including starvation-induced sleep suppression and thermogenetic stimulation of various wake-promoting neurons. Manipulations that elicit

sleep rebound are accompanied by increased food intake during or following sleep deprivation, as well as increased respiratory metabolism—two dominant variables in the energy balance equation. These results suggest that homeostatic regulation of sleep is a result or a readout of energy balance.

Results

We first tested the effect of mechanical sleep deprivation (SD) on food intake using a custom-made fly vial agitator (Supplemental Movie 1), which effectively prevented sleep as indicated by subsequent sleep rebound (Fig. 1a). Additionally, sleep-deprived flies consumed more food the following day despite the homeostatic increase in sleep, which correspondingly limits dining hours (Fig. 1a). Importantly, increased consumption was not indirectly due to compromised feeding during the deprivation treatment, as food intake was not significantly different during SD (Fig. 1b).

Manipulating food intake reciprocally affected sleep. Depriving flies of food overnight increased not only food intake but also total sleep the following day (Fig. 1c), consistent with previous studies on the antihypnotic effects of starvation[3]. In short, loss of either sleep or food intake led to an increase in both.

What is the relationship between sleep loss, rebound sleep, and food intake? Are homeostatic sleep rebound and hyperphagia controlled by the same mechanisms and, if not, can they be experimentally decoupled? Does sleep loss have direct orexigenic effects, or is the increased feeding a response to altered energy states following sleep deprivation? To explore these questions, we thermogenetically activated several broad

groups of neurons that cause soft or hard sleep loss[20], allowing us to dissect the components of sleep homeostasis and their potential effects on feeding.

TrpA1-mediated stimulation of cholinergic neurons suppressed sleep and led to sleep rebound (Fig. 2a-b), as previously reported[20]. Additionally, the compensatory sleep rebound was accompanied by post-SD hyperphagia (Fig. 2b). Sleep rebound and hyperphagia were observed in both sexes (Fig. S1a). In contrast, thermogenetic activation of octopaminergic (Fig. 2c) or dopaminergic (Fig. 2d) neurons induced sleep suppression without statistically significant changes in rebound sleep or feeding.

Cholinergic neuron stimulation, which induces rebound sleep, resulted in greater sleep loss than the octopaminergic and dopaminergic manipulations (Fig. 2e). To test whether the degree of sleep loss drives rebound sleep, a longer, 12-hour stimulation was tested. Extended activation of octopaminergic neurons further suppressed sleep but had no effect on sleep rebound or food consumption afterwards (Fig. S1b). Dopaminergic neurons, when stimulated for longer, suppressed sleep and elicited sleep rebound, but had no effect on post-SD food intake (Fig. S1b). Instead, hyperphagia was observed during the treatment period itself. These results suggest that hard sleep loss is not only due to the degree of sleep loss, but that both a threshold level and "class" of SD is necessary to trigger sleep rebound.

To better understand the relationship between sleep homeostasis and feeding behavior, we examined the effects of activating more restricted subsets of cholinergic neuronal populations implicated in various aspects of sleep regulation: sleep homeostasis; arousal (suppresses sleep but does not elicit sleep rebound); or sleep pressure

(increases sleep following neuron activation with no sleep loss during treatment)[19,20]. Consistent with previous findings[20], activation of neurons labeled by 24C10-GAL4 or the intersection of 24C10-GAL4 and ppk-GAL4 (24C10∩ppk) suppressed sleep and elicited sleep rebound (Fig. 3a, Fig. S2a). There was no post-SD hyperphagia, but food intake was elevated during neuron stimulation, similar to the feeding phenotype that accompanied the longer activation of dopaminergic neurons (Fig. S1b). Activation of neurons labeled by 44F01- or 60D04-GAL4 suppressed sleep during the heat phase but did not evoke homeostatic sleep rebound (Fig. 3b), consistent with previous work suggesting that these neurons are directly wake-promoting and bypass the sleep homeostat[20]. The 44F01 and 60D04 manipulations did not affect food intake during or after activation (Fig. 3b).

Previous studies have found that stimulation of 69F08 neurons in the ellipsoid body accrues sleep pressure either following[21] or bypassing[19] sleep deprivation. In our studies, activation of 69F08 neurons suppressed sleep, and induced both sleep rebound and post-SD hyperphagia while leaving food intake during the heat phase unaffected (Fig. 3c). We confirmed the sleep loss with 69F08 activation in females as well (Fig. S2b). Additionally, 69F08 activation decreased arousal threshold (Fig. S2c), suggesting that 69F08 neurons are wake-promoting. The sleep suppression was not due to the upright placement of the behavioral chambers, since 69F08 stimulation similarly suppressed sleep in ARC chambers oriented horizontally, mimicking the orientation in the widely used Drosophila Activity Monitor (DAM; TriKinetics) system (Fig. S2d).

The DAM system deduces activity levels and sleep based on the number of times an animal crosses an infrared beam at the center of the chamber[22]. This method assumes that any change in activity around the chamber midline is proportional to movement in unsampled areas—i.e., both ends of the chamber. Thus, single-beam DAM systems cannot accurately reflect the activity of manipulations that alter the spatial distributions of the movements[23]. In contrast, the ARC tracks the planar location of the animal, providing a more accurate measure of activity. We hypothesized that the particular assay used might have affected the detection of movement and sleep in 69F08 neuron-stimulated animals, consistent with other reports of artifactual hyperactivity using the DAM system[23,24].

To reconcile the discrepancy between our 69F08 activation results and the previously reported sleep phenotype[19], we ran post hoc analyses on the movement characteristics of the subjects in our original experiment (Fig. 3c). We found that 69F08 activation altered the spatial distribution of fly movements, where the experimental animals spent more time taking shorter strides in the bottom half of the chambers during stimulation (Fig. S2e). We also reanalyzed the activity in a DAM-like manner—i.e., counting the number of times the flies crossed an imaginary line bisecting the chamber. Not surprisingly, sleep estimated using this method failed to capture the sleep loss during 69F08-neuron stimulation (Fig. S2e). Collectively, these results support the conclusion that increased sleep following 69F08 neuron activation is indeed sleep rebound and is not independent of homeostatic sleep debt[19,21].

We hypothesized that the association of hard sleep loss with increased feeding might be due to changes in metabolic demands or energy balance during sleep loss. Thus,

we next measured CO_2 production as a proxy for metabolic rate during various neuronal manipulations. Increased respiration was observed in all manipulations associated with hard sleep loss and increased feeding, as well as with the octopaminergic manipulation (Fig. 4a-d). CO_2 production was not affected by stimulation of 44F01 and 60D04—manipulations that also did not induce sleep rebound or affect feeding behavior (Fig. 4e).

It is unclear whether hyperphagia, observed in all hard sleep loss manipulations, is merely a readout for energy deficit, or if it is more directly influences sleep pressure. The former seems more likely, since manipulations that show increased food intake during stimulation, sufficient to eliminate post-treatment hunger, do not eliminate subsequent sleep rebound (Fig. 3a, Fig. S1b, Fig. S2a). Additionally, preventing food access during stimulation in these manipulations did not further increase sleep rebound compared to controls (Fig. 5). However, these studies are complicated by the possibility of a floor effect with the amount of sleep loss that can be induced. Interestingly, starvation during stimulation did not further increase recovery food intake compared to controls, suggesting that the manipulations tested (TH- and 24C10∩ppk-neuron stimulation) do not directly affect homeostatic feeding (Fig. 5). Taken together, our results suggest that hard sleep loss and energy deficit drive sleep and feeding pressures concomitantly.

Discussion

Our findings expand upon previous work demonstrating that sleep deprivation increases metabolism[8]. While increased metabolic rates have been observed in sleep-deprived animals[25,26], our studies suggest negative energy balance is a critical factor in driving rebound sleep. However, we tested a limited set of sleep-suppressing manipulations. Assessing how other sleep-altering manipulations affect metabolism and feeding behavior could further support our interpretation.

Although hyperphagia is observed in all instances of hard sleep loss, its variable timing (during or after SD), and the lack of an additive effect on sleep rebound with starvation and wake-promoting neuron stimulation (Fig. 5), suggest that feeding pressure is not directly driving sleep pressure. Instead, it may be merely a readout of energy deficit. This may also explain why Tdc2 stimulation increases respiration without inducing sleep rebound—no hyperphagia was observed, suggesting that these flies were not in energy deficit. Since CO_2 production alone does not always accurately reflect energy expenditure[27], it is possible that additional metabolic measurements may reveal a more consistent factor that ties energy balance to sleep rebound. It is also possible that Tdc2 stimulation suppresses the feeding response to metabolic expenditure. Octopaminergic stimulation has been shown to suppress sleep rebound following mechanical sleep deprivation, and octopaminergic MS1 neurons suppress sleep in favor of mating[20,29].

Sleep is critical for normal day-to-day function, and deficient or irregular sleep is associated with numerous adverse effects on health, including on metabolism[30-32]. Sleep homeostasis and rebound sleep serve restorative functions, perhaps only upon signaling from sleep debt. Our findings suggest that one such signal might be negative energy balance. This idea is consistent with the idea that one of the functions of sleep

is to conserve energy, especially since sleep typically decreases metabolic rate[8,35,36]. Overall, our study adds to the collective literature that couples sleep homeostasis to metabolism and energy balance.

While our study provides compelling evidence for the interplay of energy balance and sleep homeostasis, there are limitations. The use of *Drosophila melanogaster*, while enabling powerful genetic manipulations, requires careful consideration when extrapolating findings to mammals with more complex metabolic systems. Additionally, it remains possible that the link between energy deficit and sleep rebound involves not only direct metabolic sensing, but also potential secondary stress responses or circadian misalignment induced by sleep deprivation. Further research using complementary model systems and investigating additional aspects of metabolism could provide a more complete understanding of sleep regulation.

In conclusion, our study establishes a critical link between energy balance and mechanisms of sleep homeostasis. This finding challenges the traditional view of sleep regulation as being primarily driven by the accumulation of sleep debt and suggests a more integrative model where sleep serves a critical role in metabolic restoration. Our work highlights the potential for therapeutic approaches to sleep disorders or metabolic conditions through the manipulation of energy states. Future studies exploring the specific metabolic signals communicating with sleep homeostatic centers will further advance our understanding of this fundamental relationship.

Materials and methods

<u>Fly lines and husbandry</u>. Flies were reared and maintained on a standard cornmeal-sucrose-yeast medium (3.1% active dry yeast, 0.7% agar, 5.8% cornmeal, 1.2% sucrose (all w/v), 1% propionic acid (v/v) and 0.22% Tegosept (w/v, pre-dissolved in ethanol)) in a light-, temperature-, and humidity-controlled incubator (12-/12-h light/dark cycle, 25°C, 60% relative humidity), unless otherwise specified. Young adults (2-4 days old) were separated into single-sex vials (10-15 flies/vial) under CO_2 anesthesia and transferred to fresh food every other day until 7-10 days old. Female flies used in experiments were once-mated. Transgenic flies were typically outcrossed for >10 generations to the indicated strain (a "Cantonized" white-eyed line, wCS; or white-eyed Berlin, w[Berlin]) that was tested to be free of *Wolbachia*.

For thermogenetic experiments, flies were reared and maintained at 18 °C except during testing, during which they were transferred to control (22 °C) or experimental temperatures. 69F08- and ppk∩24C10-neuron stimulation was induced at 29 °C. All other thermogenetic stimulations were performed at 27 °C.

The following lines were acquired from Bloomington Drosophila Stock Center: ChAT-GAL4 (BDSC #6793), TH-GAL4 (BDSC #8848), Tdc2-GAL4 (BDSC #9313), 24C10-GAL4 (BDSC #49075), 44F01-GAL4 (BDSC #45313), 60D04-GAL4 (BDSC #45356), and 69F08-GAL4 (BDSC #39499). The split-GAL4 constructs (ppk∩24C10) were acquired from Dr. William Joiner. UAS-TrpA1 was acquired from Dr. Ulrike Heberlein. All lines except the ppk∩24C10 split-GAL4 were outcrossed.

Diets. Liquid diets were filtered (0.22-μm cellulose acetate sterile syringe filter, VWR) prior to usage. Solid diets were prepared with Bacto™ Agar (BD Diagnostics) and a propionic/phosphoric acid mix was added to prevent bacterial growth. ARC experiments were conducted with a standard liquid diet of 5% sucrose + 5% yeast extract (w/v) except during starvation, during which the feeding capillaries were filled with deionized H_2O. Experiments using the Drosophila Activity Monitor (DAM, TriKinetics) used solid food of 5% sucrose + 5% yeast extract (w/v).

Mechanical sleep deprivation. To mechanically prevent flies from sleeping, we used a custom-built device that agitates fly vials in a 3-dimensional space. The device consisted of a 3D-printed vial carrier secured to two stepper motors (Zhengke Motor #ZGA28RP37.9i) controlled by an Arduino microcontroller. One motor rotated the carrier in a cartwheeling motion and the other delivered short, timed punches to further dislodge and disrupt the flies (Supplemental Video). This agitation cycle was repeated every 5 minutes to effectively prevent sleep. Sleep deprivation with this device was confirmed by subsequent sleep rebound compared to undisturbed controls.

Sleep and feeding behavior measurement (ARC, DAM). We used the Activity Recording CAFE (ARC) system [37], a modified version of CApillary FEeder (CAFE)[38]) [38], to simultaneously measure sleep and food intake. Flies were individually loaded by mouth pipette into chambers with access to liquid food (5% sucrose, 5% yeast extract (w/v) in ddH_2O) in a 5-μL glass capillary. Prior to recording, flies were acclimated to the chambers overnight. Feeding events were detected as sudden, suprathreshold drops in the liquid meniscus level. Arousal threshold was measured as previously described [4,37]. The analyses excluded animals that died during the experiment to avoid artificially high

arousal threshold from non-responsiveness. General malaise or other artifacts were monitored with real-time camera output as well as through post hoc analysis of the animal movements. In selected experiments, we used the Drosophila Activity Monitor (DAM; TriKinetics, Waltham, MA) to validate ARC results.

For horizontally placed ARC experiment, the ARC chambers were loaded with an aliquot of agar on the bottom as in the typical setup. Afterwards, flies were anesthetized by CO_2 and loaded into the chambers.   Before the flies recovered from anesthesia, the top of the chamber was fitted with 200-µL unfiltered pipette tips filled with solid food (5% yeast extract, 5% sucrose, 1% agar, all w/v), instead of capillary tubes holding liquid food. The narrow ends of the pipette tips were trimmed to widen the opening for easier access by flies, and the opposite ends were secured by Parafilm® to reduce evaporation. The chambers were laid flat or horizontally on the incubator shelf and cameras were mounted above. Animal movements were tracked and recorded as in the standard ARC setup, except that food intake was not monitored.

Radiolabeled food intake. To quantify food intake during and following mechanical sleep deprivation, we measured consumption of radiolabeled solid food in standard vials[39]. Flies were divided into four experimental groups: sleep-deprived with radiolabeled ("hot") food, sleep-deprived with non-radiolabeled ("cold") food, undisturbed with hot food, and undisturbed with cold food. Flies in the hot food group had access to food mixed with 1–2 µCi/mL [α-[32]P] during the sleep deprivation period. Following the 12 hours of sleep deprivation, flies in the hot food groups were collected in empty vials and frozen at -80 °C. Flies from the cold food groups were allowed to feed *ad libitum* on hot food for 24 hours and then frozen in empty vials. Accumulated

radiotracer was subsequently quantified in whole flies by liquid scintillation[39]. Pre-weighed aliquots of non-solidified radiolabeled media were used as a conversion factor. We have previously confirmed that the low concentration of radiolabel used (<1 nM) does not affect feeding behavior[39].

Respiratory metabolic expenditure. We used CO_2 generation as a proxy for overall energy expenditure, using the previously described method with slight modifications[40]. A metabolic chamber was fabricated using a non-barrier P1000 micropipette tip (Fisher Scientific cat no. 02-681-172) containing soda lime (Ward's Science #470302-416), separated from the residential area by a piece of low-density packing foam. The narrow end of the tip was fitted with a 50-µL glass capillary (Pyrex Disposable Micro-Sampling Pipets, Corning cat no. 7099S-50) and sealed with hot glue. A 100-µL droplet of 5% sucrose, 5% yeast extract, and 1% agar (w/v) was applied to the wall of each metabolic chamber, including the blank control. Each metabolic chamber was loaded with 4-5 flies using a mouth aspirator and sealed with non-hardening modeling clay (Plastilina, Sargent Art #22-7688). The blank chambers, used as controls for atmospheric pressure, were constructed identically except no flies were loaded. A thin layer chromatography (TLC) box was filled with red-dyed water (Eosin Y, Fisher Chemical cat no. E511-100) and fitted with a 3d-printed rack that suspended the metabolic chambers at an equal height with all glass capillary tips below the surface of the dyed water. The prepared TLC box was sealed with high vacuum grease (Dow Corning) and placed in front of a camera in a temperature-controlled incubator. PhenoCapture (www.phenocapture.com) software was used to capture a series of pictures of the TLC box at 5-minute intervals, and the timelapse images were analyzed using FiJi [41] to

automatically measure the meniscus level of the dyed water inside the glass capillaries. The changes in the meniscus levels were assumed to be directly proportional to respiratory metabolic expenditure.

Data analysis. All analyses were performed using the R statistical package[42]. For one-factor experiments, normally distributed data were first checked for homogeneity of variances with Levene's Test followed by standard analysis of variance (ANOVA) or, in case of unequal variance, Welch's variant of ANOVA. Tukey's multiple comparisons of means test was used for post hoc comparisons between groups and data that were not normally distributed were analyzed using the Kruskal-Wallis test. When comparing between phases of the same experiment, we used repeated measures ANOVA or, if the data were of unequal variances or not normally distributed, the Friedman test. Significant interaction terms were followed up with post hoc within-genotype comparisons between phases with Bonferroni adjustments for multiple comparisons. An α level of .05 and power $(1-\beta)$ of .80 were used for all statistical tests.

Figure 1. Mechanical disturbance and starvation suppress sleep and lead to post-deprivation sleep rebound and hyperphagia.

(a) Mechanical sleep deprivation overnight leads to sleep rebound and hyperphagia the following day. N = 30 Canton-S males per condition. (b) Food intake during mechanical sleep deprivation is unchanged. N = 18 Canton-S males per condition. (c) Sleep rebound following starvation. N = 30 Canton-S males per condition. Box plots represent median and interquartile range. Lines represent mean and the shaded regions SEM. Asterisks denote significant difference between the control and the sleep-deprived group by Welch's t test (a,c) or Tukey multiple comparisons of means (b) (**, $p < 0.01$; ***, $p < 0.001$).

Figure 2. Food intake and sleep are altered during and following broad sleep-suppressing manipulations.

(a) Schematic of the thermogenetic stimulation and references for the terms denoting sleep and feeding behaviors surrounding the stimulation. Baseline sleep is recorded at 22 °C prior to the stimulating heat phase, and thermogenetically stimulated ("heat phase") behaviors are recorded at 27 °C unless specified otherwise. Heat phase Δsleep is defined as the difference between the total sleep during ZT 18-24 of the heat phase and that of the baseline phase. Rebound phase Δsleep is calculated from the difference between sleep during ZT 0-6 (rebound phase) and that of the rebound phase. Heat phase feeding refers to total food intake during the heat stimulation, and recovery

feeding refers to food intake during the 24 hours following the heat stimulation. (b) Cholinergic stimulation elicits sleep loss and rebound, coupled with an increase in food consumption during recovery phase. (c-d) Octopaminergic (c) and dopaminergic (d) stimulation elicit sleep loss but not sleep rebound or increased recovery feeding. (e) Stimulation of cholinergic, octopaminergic, or dopaminergic neurons suppress sleep, but only cholinergic stimulation is followed by sleep rebound. (f) Cholinergic stimulation results in increased recovery feeding. The light pink rectangular background indicates heat phase (b-d). Box plots represent median and interquartile range. All lines represent mean and the shaded regions SEM. N = 17-18 flies per genotype. Following one-way ANOVA, significant differences were compared post-hoc using Tukey pairwise comparisons between the experimental and each of the control lines. Significant differences from the post-hoc comparisons are noted as: #, significantly different from TrpA1 control; *, significantly different from GAL4 control. The number of symbols denote the p value of the difference between the control and the sleep-deprived group (**/##, $p < 0.01$; ***/###, $p < 0.001$).

Supplemental Figure S1. Food intake and sleep are altered during and following broad sleep-suppressing manipulations.

(a) Cholinergic stimulation elicits sleep loss, sleep rebound, and post-SD overfeeding in females (N = 20 per genotype). (b) Prolonged (12 hr) stimulation of octopaminergic neurons suppresses sleep but does not affect post-SD sleep or food intake, whereas prolonged stimulation of dopaminergic neurons suppresses sleep, increases food intake during the heat phase, and induces rebound sleep. The light pink rectangular background indicates heat phase (b-d). Box plots represent median and interquartile range. All lines represent mean and the shaded regions SEM. N = 15 flies per genotype. Following one-way ANOVA, significant differences were compared post-hoc using Games-Howell (if variance not homogeneous by Levene's test) or Tukey pairwise comparisons (if variance homogeneous) between the experimental and each of the control lines. Significant differences from the post-hoc comparisons are noted as: #, significantly different from TrpA1 control; *, significantly different from GAL4 control. The number of symbols denote the p value of the difference between the control and the sleep-deprived group (***/###, $p < 0.001$).

Figure 3. Thermogenetic stimulation of various subsets of cholinergic neurons causes soft or hard sleep loss

(a) 24C10-neuron stimulation elicits sleep loss and increased feeding during stimulation, as well as increased sleep during recovery phase. *N* = 12 males per genotype. (b) Stimulation of 44F01 or 60D04 neurons suppresses sleep without post-SD hypersomnia or hyperphagia. *N* = 6 males per genotype. (c) Stimulation of R2 neurons using 69F08-

GAL4 driver suppresses sleep and food intake during the stimulation, with increased post-SD sleep rebound and food intake. $N = 12$ males per genotype. Box plots represent median and interquartile range. All lines represent mean and the shaded regions SEM. Following one-way ANOVA, significant differences were compared post-hoc using Games-Howell (if variance not homogeneous by Levene's test) or Tukey pairwise comparisons (if variance homogeneous) between the experimental and each of the control lines. Significant differences from the post-hoc comparisons are noted as: #, significantly different from TrpA1 control; *, significantly different from GAL4 control. The number of symbols denote the p value of the difference between the control and the sleep-deprived group (*/#, $p < 0.05$; **/##, $p < 0.01$; ***/###, $p < 0.001$).

Supplemental Figure S2. Thermogenetic stimulation of restricted subsets of cholinergic neurons causes hard sleep loss

(a) Stimulation of a more restricted subset of R2 neurons suppresses sleep and increases feeding during the stimulation, followed by sleep rebound. (b) 69F08 stimulation also elicits sleep loss and rebound sleep in females. N = 20 females per genotype. (c) 69F08 stimulation decreases arousal threshold, consistent with sleep loss. N = 10 males per genotype. (d) Placing the ARC chamber flat against the ground show sleep loss during 69F08 stimulation, suggesting that the sleep loss is not due to the restrictions from vertical orientation of the chamber. N = 20 males per genotype. (e) Spatial map (left) and walk length frequency histogram (middle) of animals within ARC chamber during 69F08 stimulation from Fig. 3c. Simulated bisecting beam crosses from the animal movements show unchanged sleep, contrary to the observation from animal tracking. Box plots represent median and interquartile range. Lines represent mean and the shaded regions SEM. Following one-way ANOVA, significant differences were checked for homogeneity of variance using Levene's test and post-hoc comparisons between the experimental and each of the control lines were performed using Tukey's multiple comparisons test. Significant differences from the post-hoc comparisons are noted as: #, significantly different from TrpA1 control; *, significantly different from GAL4

control. The number of symbols denote the p value of the difference between the control and the sleep-deprived group (*/#, $p < 0.05$; **/##, $p < 0.01$; ***/###, $p < 0.001$).

Figure 4. Increased respiratory metabolism is correlated with presence of hard sleep loss

(a-e) 2-hour time-course (in 30-minute bins) and average CO_2 production during stimulation of the noted neurons at the indicated temperatures. *N* = 6 cohorts per genotype, 4 males per cohort. (f) 4-hour time-course (in 30-minute bins) and average CO_2 production in fed, starved, or starved and refed flies. *N* = 6 cohorts per condition, 4 Canton-S males per cohort. Box plots represent median and interquartile range. Lines represent mean and the shaded regions SEM. Following one-way ANOVA, significant

differences were checked for homogeneity of variance using Levene's test and post-hoc comparisons between the experimental and each of the control lines were performed using Tukey's multiple comparisons test. Significant differences from the post-hoc comparisons are noted as: #, significantly different from TrpA1 control; *, significantly different from GAL4 control. The number of symbols denote the p value of the difference between the control and the sleep-deprived group (*/#, $p < 0.05$; **/##, $p < 0.01$; ***/###, $p < 0.001$).

Figure 5. Starvation during stimulations that accompany increased food intake does not increase rebound sleep or recovery food intake

(a,b) Prohibiting access to food during the TH-(a) or 24C10 ∩ ppk-(b) stimulation-induced SD is insufficient to increase sleep rebound or recovery food intake. Following one-way ANOVA, post-hoc Tukey pairwise comparisons were performed between the experimental (GAL4 > TrpA1) and each of the control lines. Significant differences from the post-hoc comparisons are noted as: #, significantly different from TrpA1 control (wCS × TrpA1); *, significantly different from GAL4 control (GAL4 × w[Berlin]). $N = 10$ males per genotype and treatment.

References

1.   Greer, S.M., Goldstein, A.N., and Walker, M.P. (2013). The impact of sleep deprivation on food desire in the human brain. Nat Commun *4*, 2259. 10.1038/ncomms3259.

2.   St-Onge, M.P., McReynolds, A., Trivedi, Z.B., Roberts, A.L., Sy, M., and Hirsch, J. (2012). Sleep restriction leads to increased activation of brain regions sensitive to food stimuli. Am J Clin Nutr *95*, 818-824. 10.3945/ajcn.111.027383.

3.   Lee, G., and Park, J.H. (2004). Hemolymph sugar homeostasis and starvation-induced hyperactivity affected by genetic manipulations of the adipokinetic hormone-encoding gene in Drosophila melanogaster. Genetics *167*, 311-323. 10.1534/genetics.167.1.311.

4.   Murphy, K.R., Deshpande, S.A., Yurgel, M.E., Quinn, J.P., Weissbach, J.L., Keene, A.C., Dawson-Scully, K., Huber, R., Tomchik, S.M., and Ja, W.W. (2016). Postprandial sleep mechanics in Drosophila. Elife *5*. 10.7554/eLife.19334.

5.   Mansbach, R.S., and Lorenz, D.N. (1983). Cholecystokinin (CCK-8) elicits prandial sleep in rats. Physiology & behavior *30*, 179-183. 10.1016/0031-9384(83)90002-1.

6.   Shemyakin, A., and Kapas, L. (2001). L-364,718, a cholecystokinin-A receptor antagonist, suppresses feeding-induced sleep in rats. Am J Physiol Regul Integr Comp Physiol *280*, R1420-1426. 10.1152/ajpregu.2001.280.5.R1420.

7.   Orr, W.C., Shadid, G., Harnish, M.J., and Elsenbruch, S. (1997). Meal composition and its effect on postprandial sleepiness. Physiology & behavior *62*, 709-712. 10.1016/s0031-9384(97)00012-7.

8.   Stahl, B.A., Slocumb, M.E., Chaitin, H., DiAngelo, J.R., and Keene, A.C. (2017). Sleep-Dependent Modulation of Metabolic Rate in Drosophila. Sleep *40*. 10.1093/sleep/zsx084.

9.   Koban, M., and Swinson, K.L. (2005). Chronic REM-sleep deprivation of rats elevates metabolic rate and increases UCP1 gene expression in brown adipose tissue. Am J Physiol Endocrinol Metab *289*, E68-74. 10.1152/ajpendo.00543.2004.

10.   Schmid, S.M., Hallschmid, M., and Schultes, B. (2015). The metabolic burden of sleep loss. Lancet Diabetes Endocrinol *3*, 52-62. 10.1016/S2213-8587(14)70012-9.

11.   Scharf, M.T., Naidoo, N., Zimmerman, J.E., and Pack, A.I. (2008). The energy hypothesis of sleep revisited. Prog Neurobiol *86*, 264-280. 10.1016/j.pneurobio.2008.08.003.

12.   Reichert, S., Pavon Arocas, O., and Rihel, J. (2019). The Neuropeptide Galanin Is Required for Homeostatic Rebound Sleep following Increased Neuronal Activity. Neuron *104*, 370-384 e375. 10.1016/j.neuron.2019.08.010.

13.   Monsalve, G.C., Van Buskirk, C., and Frand, A.R. (2011). LIN-42/PERIOD controls cyclical and developmental progression of C. elegans molts. Curr Biol *21*, 2033-2045. 10.1016/j.cub.2011.10.054.

14.   Szentirmai, E., and Kapas, L. (2014). Intact brown adipose tissue thermogenesis is required for restorative sleep responses after sleep loss. Eur J Neurosci *39*, 984-998. 10.1111/ejn.12463.

15.   St-Onge, M.P., Roberts, A.L., Chen, J., Kelleman, M., O'Keeffe, M., RoyChoudhury, A., and Jones, P.J. (2011). Short sleep duration increases energy intakes but does not change energy expenditure in normal-weight individuals. Am J Clin Nutr *94*, 410-416. 10.3945/ajcn.111.013904.

16.   Yang, C.L., Schnepp, J., and Tucker, R.M. (2019). Increased Hunger, Food Cravings, Food Reward, and Portion Size Selection after Sleep Curtailment in Women Without Obesity. Nutrients *11*. 10.3390/nu11030663.

17.   Benedict, C., Hallschmid, M., Lassen, A., Mahnke, C., Schultes, B., Schioth, H.B., Born, J., and Lange, T. (2011). Acute sleep deprivation reduces energy expenditure in healthy men. Am J Clin Nutr *93*, 1229-1236. 10.3945/ajcn.110.006460.

18.   Donlea, J.M., Pimentel, D., Talbot, C.B., Kempf, A., Omoto, J.J., Hartenstein, V., and Miesenbock, G. (2018). Recurrent Circuitry for Balancing Sleep Need and Sleep. Neuron *97*, 378-+. 10.1016/j.neuron.2017.12.016.

19.   Liu, S., Liu, Q.L., Tabuchi, M., and Wu, M.N. (2016). Sleep Drive Is Encoded by Neural Plastic Changes in a Dedicated Circuit. Cell *165*, 1347-1360. 10.1016/j.cell.2016.04.013.

20.   Seidner, G., Robinson, J.E., Wu, M., Worden, K., Masek, P., Roberts, S.W., Keene, A.C., and Joiner, W.J. (2015). Identification of Neurons with a Privileged Role in Sleep Homeostasis in Drosophila melanogaster. Curr Biol *25*, 2928-2938. 10.1016/j.cub.2015.10.006.

21.   Satterfield, L.K., De, J., Wu, M., Qiu, T., and Joiner, W.J. (2022). Inputs to the Sleep Homeostat Originate Outside the Brain. J Neurosci *42*, 5695-5704. 10.1523/JNEUROSCI.2113-21.2022.

22.   Chiu, J.C., Low, K.H., Pike, D.H., Yildirim, E., and Edery, I. (2010). Assaying locomotor activity to study circadian rhythms and sleep parameters in Drosophila. Journal of visualized experiments : JoVE. 10.3791/2157.

23.   De, J., Varma, V., Saha, S., Sheeba, V., and Sharma, V.K. (2013). Significance of activity peaks in fruit flies, Drosophila melanogaster, under seminatural conditions. Proc Natl Acad Sci U S A *110*, 8984-8989. 10.1073/pnas.1220960110.

24.   King, L.B., Koch, M., Murphy, K.R., Velazquez, Y., Ja, W.W., and Tomchik, S.M. (2016). Neurofibromin Loss of Function Drives Excessive Grooming in Drosophila. G3 (Bethesda) *6*, 1083-1093. 10.1534/g3.115.026484.

25. Everson, C.A., Bergmann, B.M., and Rechtschaffen, A. (1989). Sleep deprivation in the rat: III. Total sleep deprivation. Sleep *12*, 13-21. 10.1093/sleep/12.1.13.

26. Kushida, C.A., Bergmann, B.M., and Rechtschaffen, A. (1989). Sleep deprivation in the rat: IV. Paradoxical sleep deprivation. Sleep *12*, 22-30. 10.1093/sleep/12.1.22.

27. Oshima, T., Graf, S., Heidegger, C.P., Genton, L., Pugin, J., and Pichard, C. (2017). Can calculation of energy expenditure based on CO_2 measurements replace indirect calorimetry? Crit Care *21*, 13. 10.1186/s13054-016-1595-8.

28. Brown, E.B., Klok, J., and Keene, A.C. (2022). Measuring metabolic rate in single flies during sleep and waking states via indirect calorimetry. J Neurosci Methods *376*, 109606. 10.1016/j.jneumeth.2022.109606.

29. Machado, D.R., Afonso, D.J., Kenny, A.R., Oztu Rk-Colak, A., Moscato, E.H., Mainwaring, B., Kayser, M., and Koh, K. (2017). Identification of octopaminergic neurons that modulate sleep suppression by male sex drive. Elife *6*. 10.7554/eLife.23130.

30. Axelrod, S., Li, X., Sun, Y., Lincoln, S., Terceros, A., O'Neil, J., Wang, Z., Nguyen, A., Vora, A., Spicer, C., et al. (2023). The Drosophila blood-brain barrier regulates sleep via Moody G protein-coupled receptor signaling. Proc Natl Acad Sci U S A *120*, e2309331120. 10.1073/pnas.2309331120.

31. Bedont, J.L., Kolesnik, A., Pivarshev, P., Malik, D., Hsu, C.T., Weljie, A., and Sehgal, A. (2023). Chronic sleep loss sensitizes Drosophila melanogaster to nitrogen stress. Curr Biol *33*, 1613-1623 e1615. 10.1016/j.cub.2023.03.008.

32. Wang, Z., Lincoln, S., Nguyen, A.D., Li, W., and Young, M.W. (2022). Chronic sleep loss disrupts rhythmic gene expression in Drosophila. Front Physiol *13*, 1048751. 10.3389/fphys.2022.1048751.

33. Dissel, S., Angadi, V., Kirszenblat, L., Suzuki, Y., Donlea, J., Klose, M., Koch, Z., English, D., Winsky-Sommerer, R., van Swinderen, B., and Shaw, P.J. (2015). Sleep restores behavioral plasticity to Drosophila mutants. Curr Biol *25*, 1270-1281. 10.1016/j.cub.2015.03.027.

34. Haynes, P.R., Pyfrom, E.S., Li, Y., Stein, C., Cuddapah, V.A., Jacobs, J.A., Yue, Z., and Sehgal, A. (2024). A neuron-glia lipid metabolic cycle couples daily sleep to mitochondrial homeostasis. Nature neuroscience *27*, 666-678. 10.1038/s41593-023-01568-1.

35. Siegel, J.M. (2005). Clues to the functions of mammalian sleep. Nature *437*, 1264-1271. 10.1038/nature04285.

36. Sarikaya, D.P., Cridland, J., Tarakji, A., Sheehy, H., Davis, S., Kochummen, A., Hatmaker, R., Khan, N., Chiu, J., and Begun, D.J. (2020). Phenotypic coupling of sleep and starvation resistance evolves in D. melanogaster. BMC Evol Biol *20*, 126. 10.1186/s12862-020-01691-8.

37. Murphy, K.R., Park, J.H., Huber, R., and Ja, W.W. (2017). Simultaneous measurement of sleep and feeding in individual Drosophila. Nat Protoc *12*, 2355-2366. 10.1038/nprot.2017.096.

38. Ja, W.W., Carvalho, G.B., Mak, E.M., de la Rosa, N.N., Fang, A.Y., Liong, J.C., Brummel, T., and Benzer, S. (2007). Prandiology of Drosophila and the CAFE assay. P Natl Acad Sci USA *104*, 8253-8256. 10.1073/pnas.0702726104.

39. Deshpande, S.A., Carvalho, G.B., Amador, A., Phillips, A.M., Hoxha, S., Lizotte, K.J., and Ja, W.W. (2014). Quantifying Drosophila food intake: comparative analysis of current methodology. Nat Methods *11*, 535-540. 10.1038/nmeth.2899.

40. Yatsenko, A.S., Marrone, A.K., Kucherenko, M.M., and Shcherbata, H.R. (2014). Measurement of metabolic rate in Drosophila using respirometry. Journal of visualized experiments : JoVE, e51681. 10.3791/51681.

41. Schindelin, J., Arganda-Carreras, I., Frise, E., Kaynig, V., Longair, M., Pietzsch, T., Preibisch, S., Rueden, C., Saalfeld, S., Schmid, B., et al. (2012). Fiji: an open-source platform for biological-image analysis. Nat Methods *9*, 676-682. 10.1038/nmeth.2019.

42. R Core Team (2018). R: A language and environment for statistical computing.