CrHCF244 is required for *psbA* translation in *Chlamydomonas*
reinhardtii

Xiaozhuo Wang[1,2], Guannan Wang[1,3], Lexi A. Cheramie[1], Cuiping Zhao[1], Maheshi

Dassanayake[1], James V. Moroney[1] and David J. Vinyard[1,*]

[1] Department of Biological Sciences, Louisiana State University, Baton Rouge, LA 70803

[2] Current address: Department of Biology, Stanford University, Palo Alto, CA 94305

[3] Current address: Division of Biosphere Science and Engineering, Carnegie Institution for

Science, Stanford, CA 94305

* Correspondence: dvinyard@lsu.edu, +1 225 578 0958

Highlight

We identified CrHCF244 as a translation factor of *psbA* in *Chlamydomonas*. *Arabidopsis*

HCF244 partially complements *Chlamydomonas* Δ*CrHCF244* mutant, indicating semi-

conservation of the function of this gene between organisms. Suppressor mutants of Δ*CrHCF244*

suggest the presence of alternative translation factors in *psbA* translation.

Abstract

Translation of *psbA*, the chloroplast gene that encodes the D1 subunit of Photosystem II

(PSII), is important for both PSII biogenesis and repair. The translation of the *psbA* transcript in

the chloroplast is under the control of nuclear gene products. Using a *Chlamydomonas* forward

genetic screen and whole genome sequencing, we found a mutant defective in PSII activity and

mapped the causative gene to be the homolog of *Arabidopsis High Fluorescence* (*HCF244*) gene,

or *CrHCF244*. We then demonstrated that CrHCF244 is required for *psbA* translation in the alga,

consistent with the function of HCF244 in *Arabidopsis*. The *Arabidopsis HCF244* gene also

partially complemented the algal mutant. These results experimentally support the functional

conservation of the homologs in green algae and land plants. Intriguingly, the *CrHCF244* mutant

also exhibited a relatively high rate of suppressor mutants, pointing to the presence of alternative

factor(s)/pathway(s) for D1 translation control. The establishment of *CrHCF244* as a *psbA*

translation factor in *Chlamydomonas* showed the similarities in *psbA* translation regulation in

algae and plants. The future identification of the alternative factor(s) in this alga will provide

insights on *psbA* translation in plants.

Keywords

Photosystem II, *de novo* biogenesis of Photosystem II, Photosystem II repair, *psbA* translation,

regulators of organelle genes, chlorophyll biosynthesis and association, suppressor mutant

Abbreviations

PSII   Photosystem II

27	ROGE	regulators of organelle genes
28	HCF244	high chlorophyll fluorescence 244
29	At	*Arabidopsis thaliana*
30	Cr	*Chlamydomonas reinhardtii*
31	cTP	chloroplast transit peptide
32	cpPDC	chloroplast pyruvate dehydrogenase
33	DLA2	dihydrolipoamide acetyltransferase subunit 2
34	BN-PAGE	Blue native polyacrylamide gel electrophoresis
35	TF	translation factor

Introduction

Introduction

Photosystem II (PSII) is a water-plastoquinone oxidoreductase that enables the capture

and conversion of light energy into storable chemical energy. PSII is the first component in the

photosynthetic electron transport chain that generates ATP and NADPH for CO_2 fixation.

Though much is known about the subunit composition and structure of PSII (Shen, 2015), its

biogenesis is less well understood. As a large protein complex containing over 20 subunits

encoded by both nuclear and chloroplast genomes and containing various cofactors, the

biogenesis of PSII is complicated (Lu, 2016; Nickelsen and Rengstl, 2013; Nixon *et al.*, 2010).

PSII biogenesis is a stepwise and highly coordinated process involving the synthesis of subunits,

integration into the membrane, assembly of protein subunits, and association of cofactors (Lu,

2016; Nickelsen and Rengstl, 2013; Nixon *et al.*, 2010). PSII is also subject to photoinhibition

and a repair process is needed to restore its activity, during which the D1 subunit needs to be

degraded and a new copy of the subunit synthesized *de novo* (Järvi *et al.*, 2015).

Translation of chloroplast genes encoding subunits of photosynthetic complexes

including PSII have been found to be widely regulated by nuclear gene products, which are

termed as regulators of organelle genes (ROGEs). As the control of gene expression in

chloroplasts occurs prevalently at post-transcriptional levels, including mRNA stabilization

and/or its translation, ROGEs are categorized as M and T factors, which refer to protein factors

involved in maturation or stabilization of the target mRNAs and translation regulation of the

mRNAs, respectively. Genetic and biochemical approaches have identified many ROGEs in

regulating PSII in *Chlamydomonas* (e.g. Tba1 (Somanchi *et al.*, 2005), RBP63/DLA2 (Bohne *et*

al., 2013; Ossenbuhl *et al.*, 2002), RBP40/RB38 (Schwarz *et al.*, 2007), MBB1 (Vaistij *et al.*,

2000), MBC1 (Cavaiuolo *et al.*, 2017), TBC2 (Auchincloss *et al.*, 2002)), PSI (e.g. TAA1

(Lefebvre-Legendre *et al.*, 2015)), Cyt b_6f (e.g. MCG1 (Wang *et al.*, 2015)), and ATP synthase

(e.g. TDA1 (Drapier *et al.*, 1992; Eberhard *et al.*, 2011), MDA1 (Viola *et al.*, 2019)). Recently a

systematic screening and characterization discovered several gene specific ROGEs and master

regulators responsible for regulating the expression of multiple ROGEs to be involved in

biogenesis of photosynthetic machinery (Kafri *et al.*, 2023).

Among PSII genes, the translation of *psbA* encoding the D1 subunit is important for both

PSII *de novo* biogenesis and repair. *psbA* gene expression control has been extensively studied in

cyanobacteria, algae, and plants (Mulo *et al.*, 2012). A few protein factors have been identified

as involved in *psbA* translation in *Chlamydomonas*. Mayfield and coworkers biochemically

identified *psbA* mRNA binding proteins RB47, RB60, RB38, and RB55, named based on their

molecular size (Danon and Mayfield, 1991). The gene for RB47 was later cloned as a

polyadenylate (poly (A)) binding protein, thus also named as chloroplast poly (A) binding

protein cPAB1 (Yohn *et al.*, 1998a; Yohn *et al.*, 1998b). Though chloroplast mRNAs generally

do not have poly (A) tails, there are A-rich regions in the UTRs, which could be potential

binding sites. RB60 is a disulfide isomerase protein (Kim and Mayfield, 1997) which may

regulate the redox state of RB47 and thus control its binding to *psbA* mRNA and the translation

(Kim and Mayfield, 2002). Tba1, identified in a genetic screen, is a thioredoxin protein

regulating *psbA* translation (Somanchi *et al.*, 2005). Genetic screens have also found several

Chlamydomonas nuclear mutants defective in *psbA* translation, some of which the causative

genes are not mapped including F35 (Girard-Bascou *et al.*, 1992; Yohn *et al.*, 1996). Besides the

dedicated protein factors involved in *psbA* translation, a "moonlighting" regulator was also

identified. The dihydrolipoamide acetyltransferase subunit 2 (DLA2), a subunit of chloroplast

pyruvate dehydrogenase complex (cpPDC) is involved in D1 translation regulation under

mixotrophic growth condition (Bohne *et al.*, 2013; Neusius *et al.*, 2022; Ossenbuhl *et al.*, 2002).

DLA2 switches its role between acetate metabolism as a subunit of cpPDC, and regulator of D1

translation dissociating from the cpPDC complex and associating with thylakoid membrane

through acetylation of its K197 residue (Neusius *et al.*, 2022).

In land plants, two protein factors have been identified to be involved in *psbA* translation:

High Chlorophyll Fluorescence proteins 173 and 244 (HCF173 and HCF244) (Chotewutmontri

and Barkan, 2020; Link *et al.*, 2012; Schult *et al.*, 2007). Both are members of the short-chain

dehydrogenase superfamily (Kavanagh *et al.*, 2008; Moummou *et al.*, 2012; Oppermann *et al.*,

2001). HCF173 is required for *psbA* translation initiation and mRNA stability (Schult *et al.*,

2007). HCF244, co-expressing with HCF173, is also involved in *psbA* translation, possibly

working together with HCF173, as the double mutant has a more reduced plant size and fewer

numbers of leaves (Link *et al.*, 2012).

In this study, we used forward genetics to identify a PSII defective mutant in

Chlamydomonas in which the causative gene was mapped to be a homolog of *Arabidopsis*

HCF244. By examining the mutant, we provide evidence for its role in D1 translation in this alga.

We show that the *Arabidopsis* homolog gene *HCF244* partially complements the

Chlamydomonas mutant. Intriguingly, we observed a number of suppressor mutants of the

CrHCF244 mutant that showed partial recoveries of PSII activity. In an effort to examine the

relationship of CrHCF244 to the previously identified translation activators RB60 and RB47 in

Chlamydomonas, we generated the corresponding mutants using CRISPR-Cas9 but no effect on

PSII activity was observed in those mutants. This study supports the functional conservation of

the plant ortholog HCF244 in algae. Suppressor mutants suggest an alternative pathway of D1

translation control. Genetic studies add to our understanding of the components and their roles in

D1 translation control.

Materials and methods

Strains and culture conditions

Chlamydomonas strain CC-4533 (Zhang *et al.*, 2014) and *psbA* deletion strain Fud7

(Bennoun *et al.*, 1986) were obtained from the *Chlamydomonas* Resource Center (CRC). CC-

4533 is the background strain for the CLiP library. We refer to CC-4533 as WT here. The *ami6*

mutant and the CRISPR-Cas9 knockout mutants (Δ*CrHCF244-T23*, Δ*RB47*, Δ*RB60-1, 2, 3*)

were generated in the background of CC-4533. Cells were grown in Tris-acetate-phosphate (TAP)

(Harris *et al.*, 2009) or high salt (HS) (Sueoka, 1960) media with shaking at 25°C in an

AlgaeTron incubator (Photon Systems Instruments). Growth tests were done by spotting cells

onto TAP and/or HS plates with 1.5% (w/v) agar under white light at 60 µmol photons m^{-2} s^{-1}

intensity.

Chlamydomonas transformation

Electroporation was used in this study for the delivery of plasmid DNA (for generating

random insertional mutants and complementation studies of mutants) and ribonucleoprotein

(RNP) complexes (for generating CRISPR-Cas9 knockout mutants) into *Chlamydomonas* cells.

Electroporation was done using MAX Efficiency^{TM} transformation reagent (Invitrogen)

according to the manufacturer's instructions. Briefly, cells from a 50 mL TAP culture with OD_{750}

0.3–0.5 were pelleted, washed twice with the transformation reagent and resuspended in 0.5–1

mL transformation reagent. For delivering plasmid, 110 µL of cell suspension was mixed with 1

µg plasmid DNA. For CRISPR Cas9 mutations, the 110 µL cell suspension was mixed with 10

µL Cas9-guide RNA (gRNA) RNP complex and 1 µg donor DNA. The mixture was added into a

cuvette with 0.2 cm gap (Bio-Rad), and electroporated with Gene Pulser® II (Bio-Rad) at a

voltage of 0.35 kV and capacitance of 25 µF. After electroporation, cells were kept at room

temperature (RT) for 10 min, then transferred into 10 mL TAP with 40 mM sucrose and shaken

in the dark overnight. Cells were then pelleted and spread onto selective plates. For

complementation studies of *ami6* and for the Δ*CrHCF244-T23* mutant, plasmids C1 and C2 (see

cloning construct) were electroporated into the mutants, and selected on HS plates.

Whole genome sequencing and analysis

Genomic DNA was extracted from CC-4533 and *ami6* mutant using the phenol-

chloroform-isoamyl method (Newman *et al.*, 1990). The isolated DNA was further treated with

RNase to remove RNA contamination and quantified using a Qubit fluorometer. Library

preparation and Illumina sequencing was performed by Genewiz (NJ, USA). After quality check

with FastQC (v0.11.7, http://www.bioinformatics.babraham.ac.uk/projects/fastqc/), paired-end

reads were aligned to *Chlamydomonas reinhardtii* CC-4532 reference genome (v 5.6 and v6.1)

(Merchant *et al.*, 2007) from Phytozome V13 using bowtie2 (v2.3.5) (Langmead and Salzberg,

2012) with the default settings. Aligned reads were sorted and deduplicated using Picard (v2.25.2,

https://broadinstitute.github.io/picard). We subsequently employed GRIDSS2 structural variant

caller (v2.13.2) (Cameron *et al.*, 2021) to identify rearrangement break points that generally

represent structural variant events using the recommended workflow. Results were organized

into VCF (Variant Call Format) files, and then filtered using StructuralVariantAnnotation

(v1.18.0, DOI: 10.18129/B9.bioc.StructuralVariantAnnotation) in R. The resulting structural

variants were further validated and supplemented with the ones identified by GATK (v4.1)

(O'Connor and van der Auwera, 2020) to generate the final list of variants. Lastly, the identified

variants for the WT and *ami6* mutant were compared to identify these that were only present in

the mutant.

Cloning constructs

Plasmid C1 was used for complementing *ami6* and CRISPR Cas9 knockout Δ*CrHCF244-T23* mutants with *CrHCF244* and the resulting transformants are *ami6-C1* and *T23-C1*. Plasmid C2 was used for complementing *ami6* and CRISPR Cas9 knockout Δ*CrHCF244-T23* mutants by *CrHCF244* with C-terminal tagged Venus and the resulting transformants are *ami6-C2* and *T23-C2*.

To construct the C1 plasmid, the open reading frame (ORF) of *CrHCF244* gene was amplified by Q5® High-Fidelity DNA Polymerase (New England Biolabs) using CC-4533 genomic DNA as template; the linear vector was PCR amplified using pLM006 (Mackinder *et al.*, 2016) as template with primers in Table S1 and the mCherry and His tag sequences were excluded. To construct C2, the above ORF of *CrHCF244* was used and the linear vector was PCR amplified using pLM005 (Mackinder *et al.*, 2016) as template with primers in Table S1 (Venus sequence was included). The PCR products of the ORF of *CrHCF244* and linear vectors were assembled by Gibson assembly (Gibson et al., 2009) using Gibson Assembly® Master Mix (New England Biolabs). For the vector construct to transform the *Arabidopsis HCF244* gene into *Chlamydomonas*, the coding sequence for the HCF244 protein without the transit peptide was codon optimized (IDT Codon Optimization Tool) and then fused to the sequence of the *Chlamydomonas* PsaD transit peptide on the 5' end. The chimeric sequence was synthesized by Twist Bioscience. The synthesized fragment was used as template for PCR using primers in Table S1, after which it was assembled with the PCR product using primers in Table S1 from the backbone plasmid pLM006 (Mackinder *et al.*, 2016), excluding mCherry and His tag sequence. Plasmids were verified by Sanger sequencing before transformation into *Chlamydomonas*.

Generating CRISPR-Cas9 knockout mutants in Chlamydomonas

CRISPR-Cas9 was used for targeted mutagenesis to knock out the *CrHCF244*, *RB47* and *RB60* genes in this study. The gRNA design, *in vitro* assembly with Cas9 to form RNP and delivery of RNP and donor DNA to cells were performed according to a published method (Picariello *et al.*, 2020) with minor modifications. Briefly, the crRNAs targeting the second exon of *CrHCF244* (Cre02.g142146), the fourth exon of *RB47* (Cre01.g039300) and the first exon of *RB60* (Cre02.g088200) were designed on the website of Integrated DNA Technologies (IDT).

The lack of off-targeting was verified by Cas-OFFinder (Bae *et al.*, 2014). The crRNAs and

tracrRNA (sequences were shown in Table S1), synthesized by IDT, were first annealed and then

assembled with Cas9 (Alt-R™ S.p. Cas9 Nuclease V3, IDT) at 37^{o}C for 15 min. Donor DNA for

editing each gene was PCR amplified with primers in Table S1 using the pKS-aphVIII-lox

aphVIII plasmid (Heitzer and Zschoernig, 2007) or the pHyg3 plasmid (Berthold *et al.*, 2002) as

templates to generate antibiotic-resistance cassettes flanked by 40-bp homology arms of the Cas9

cleavage site. The RNP and donor DNA was electroporated into CC-4533 after removal of cell

wall by autolysin. Transformants were selected on TAP plates with 20 µg/mL paromomycin or

hygromycin. Colonies were screened with PCR using primers in Table S1 and/or Sanger

sequencing.

Spotting cells for growth tests

191   For spotting, TAP cultures with OD_{750} around 0.3–0.5 were used. Cells from 1 mL of

192   culture were pelleted, washed with HS, and resuspended in 1 mL HS. OD_{750} was used to

193   normalize the number of cells. Five µL cultures with serial dilutions were spotted onto TAP and

194   HS plates. Plates were kept under 60 µmol photons m^{-2} s^{-1} at 25^{o}C for growth.

Chlorophyll fluorescence measurement

196   Chlorophyll a fluorescence F_v/F_m ((F_m-F_o)/F_m) was measured to indicate PSII activity

197   (Genty *et al.*, 1989). Cells in liquid cultures were measured with a double-modulation

198   fluorometer (FL 6000, Photon Systems Instruments) after two minutes dark adaptation. Cells on

199   plates were measured with a FluorCam (Photon Systems Instruments) after two minutes dark

200   adaptation.

Confocal microscopy for subcellular localization of CrHCF244

Confocal microscopy of *Chlamydomonas* cells was done according to (Mackinder *et al.*,

2016), using a Leica SP8 microscope. The Venus tagged strain (*ami6-C2*) was grown in TAP to

early exponential phase under light intensity of 60 µmol photons m^{-2} s^{-1}. Around 8 µL of culture

was mixed with an equal volume of 0.1% (w/v) low melting agarose on the slide and covered

with a #1.5 cover slip. The observation was done using a 63x water immersion objective. White

light laser at 512 nm was used for excitation of Venus fluorescence and emission between 520-

540 nm was collected. Chlorophyll autofluorescence was excited at 561 nm and emission

between 560-590 nm was collected as chlorophyll fluorescence.

Sodium dodecyl sulfate-polyacrylamide electrophoresis (SDS-PAGE) and immunoblotting

analysis

To analyze the abundance of photosynthetic proteins, *Chlamydomonas* cells were grown

in liquid TAP under light intensity of 20 µmol photons m^{-2} s^{-1}. The low light was used here to

avoid possible secondary effects on the photosynthetic machinery in the mutants from high light

intensity. Total protein was extracted using 2% SDS in 50 mM Tris-HCl (pH 8.0). A BCA assay

was used to measure total protein concentration. Laemmli sample buffer was added to protein

samples followed by incubation at room temperature for 10 min. Around 10 µg of protein was

loaded to home-made 12% Tris-Glycine polyacrylamide gels containing 6 M urea. After

electrophoresis, proteins were transferred to a PVDF membrane with pore size of 0.2 µm

(Amersham™ Hybond™ P) through wet transfer in Tris-Glycine transfer buffer and blocked

with 5% non-fat milk in Tris-buffered saline with 0.05% Tween 20 (TBST). Antibodies of PSII

subunits D1 (Agrisera, AS05 084), D2 (Agrisera, AS06 146), CP43 (Agrisera, AS11 1787),

CP47 (PhytoAB, PHY0319) and PsbO (Agrisera, AS06 142), and antibodies against PSI subunit

PsaA (Agrisera, AS06 172) and Cyt b_6f subunit Cyt f (a gift from Prof. Terry Bricker, LSU) were

used. A rabbit polyclonal antibody against RB60 was made in this study using the RB60 peptide

SEKMPPTIEFNQKNSDKIFN as the antigen by YenZym Antibodies LLC

(https://www.yenzym.com). The PVDF membrane was incubated with diluted primary

antibodies for 1 h at room temperature followed by washing and incubation with 1:20,000

diluted HRP-conjugated secondary antibody (Invitrogen, 31460) at room temperature for 1 h.

ECL reagent (Bio-Rad, Clarity Western ECL Substrate) was applied to the membrane for

chemiluminescence detection using a ChemiDoc XRS+ imaging system.

Crude thylakoid membrane isolation and Blue-native (BN) PAGE

Crude thylakoid membranes from *Chlamydomonas* were prepared as described (Garcia-

Cerdan *et al.*, 2019) with minor modifications. Cells were resuspended in lysis buffer (25 mM

HEPES-NaOH, pH 7.5, 10 mM NaCl, 5 mM $MgCl_2$ and 5% [w/v] glycerol) and passed through

a French press at gauge pressure 700 psi, $4^{o}C$ to lyse cells. The lysate was centrifuged at $3000{\times}g$,

5 min, 4°C to remove unbroken cells and cell debris. The supernatant was then centrifuged at

$30,000 \times g$, 30 min, 4°C. The pellets are used as crude thylakoid membranes. After resuspension

in solubilization buffer (25 mM HEPES-NaOH, pH 7.5, 50 mM NaCl and 5% glycerol) at Chl

concentration of 0.5 µg/µL, the thylakoid membranes were solubilized with 1% n-Dodecyl α-

maltoside (α-DDM) (GLYCON Biochemicals GmbH) on ice for 20 min (Tokutsu *et al.*, 2012),

followed by centrifugation at $20,000 \times g$ for 10 min at 4°C to remove unsolubilized thylakoids.

The supernatant was run on a 5-12% BN-PAGE gel as previously described (Wang *et al.*, 2016).

Purified PSII from the *Chlamydomonas* B-His strain in which a His_6 tag fused to the C-terminus

of PsbB was used as a control (Suzuki *et al.*, 2003).

Chlorophyll measurement

Chlorophyll measurement was done using methanol extraction, as previously described

(Porra *et al.*, 1989). Cells were grown in TAP under light intensity of 35 µmol photons m^{-2} s^{-1}

and in the dark, respectively.

^{14}C-acetate pulse labeling

Pulse labeling of *Chlamydomonas* cells was done according to the literature (Calderon *et*

al., 2023; Chua and Gillham, 1977; Spaniol *et al.*, 2022). Briefly, 5 mL of TAP-grown cells

under 20 µmol photons m^{-2} s^{-1} were pelleted and washed with HS. After resuspension in 0.5 mL

HS, cells were incubated under 20 µmol photons m^{-2} s^{-1} with vigorous shaking for 1.5 h to

deplete the cellular carbon pool. Then, 10 µg/mL cycloheximide was added to the culture to

inhibit cytosolic translation followed by the addition of 20 µCi/mL ^{14}C-acetic acid, sodium salt

(Moravek, MC200) and incubated for 20 min under 20 µmol photons m^{-2} s^{-1}. The culture was

diluted into 10 volumes of cold TAP with 50 mM nonradioactive acetate and cells were pelleted

and total protein was extracted. Protein samples were run on 12-18% Tris-Glycine SDS-PAGE

gels with 6 M urea and then transferred from the gel to a PVDF membrane with pore size of 0.2

µm (Amersham™ Hybond™ P) (Amersham™). The membrane was then exposed to a storage

phosphor screen (Molecular Dynamics) for three weeks and imaged with a Typhoon™ 8600

imager.

Reverse transcription (RT)-PCR

Cell pellets from 10 mL TAP culture with OD750 of 0.3 were suspended in 1 mL TRIzol

TM (Invitrogen). Total RNA was extracted using the Quick-RNA MiniPrep Kit (Zymo

Research). ProtoScript® First Strand cDNA Synthesis Kit (New England Lab) was used to make

cDNA using 500 ng RNA and an oligo-d(T) primer. No reverse transcriptase was added to the

negative control group. Primers for *AtHCF244*(AtHCF244_1F, AtHCF244_2R), *RB47*

(RB47_E4F and RB47_E7R), and *Actin* (Actin_F and Actin_R) are listed in Table S1.

Results

Screen and mapping of a mutant ami6 defective in PSII activity

To look for protein factors involved in PSII biogenesis, random insertional mutagenesis using a paromomycin cassette was used to generate a pool of mutants with resistance to this antibiotic. One mutant was identified which could not grow photoautotrophically and had no PSII activity, which we named *ami6* (Fig. 1A). To identify the candidate causative gene(s) in this mutant, we conducted whole genome sequencing. Analysis of the whole genome sequencing data using *Chlamydomonas reinhardtii* v6.1 (https://phytozome-next.jgi.doe.gov/info/CreinhardtiiCC_4532_v6_1) genome as the reference genome revealed a large deletion (~347 kb) which mapped onto chromosome 15 (Fig. 1B). This genomic variance was verified by PCR and Sanger sequencing (primer sequences are in Supplementary Table S1 and results are in Supplementary Fig. S1). There are 28 genes (Supplementary Dataset S1) located in the deleted genome region. One of the deleted genes, Cre02.g142146, a homolog of *Arabidopsis HCF244*, was able to restore photoautotrophic growth and PSII activity in *ami6* (Fig. 1A). Adding the fluorescent protein Venus tag at the C-terminal of CrHCF244 did not affect the function of the protein (Fig. 1A and 1C). Of note, after we confirmed the causative gene in *ami6* in late May of 2022, we noticed an annotation on this gene in *Chlamydomonas* genome v6.1, the newly released version of the genome, that this gene is mutated in F35, a previously identified nuclear mutant defective in PSII (Girard-Bascou *et al.*, 1992; Yohn *et al.*, 1996) . However, there is no publication regarding on the mapping of the gene in F35. Our identification, mapping and complementation of *ami6* mutant verified the genetic relation of *CrHCF244* with PSII in *Chlamydomonas.*

CRISPR-Cas9 to knock out CrHCF244

We next used CRISPR-Cas9 to generate a *CrHCF244* knockout mutant. The gRNA was designed to target the second exon of the gene, as illustrated in Fig. 2A. Based on PCR screening and Sanger sequencing (Supplementary Fig. S2A and Fig. S2B), transformant number 23 was selected, which is termed as Δ*CrHCF244-T23*. Δ*CrHCF244-T23* has a genomic variation at the Cas9 cleavage site in exon 2 of *CrHCF244*. For sequence alignment see Supplementary Fig. S2B. This mutation would result in the point mutation of F37 to G37 and a frame shift after the 37[th]

residue followed by a premature early stop codon, which abolished the function of the gene.

Δ*CrHCF244-T23* could not grow photoautotrophically and has no PSII activity (Fig. 2B and C),

consistent with the phenotype of *ami6*. Similar to *ami6*, *CrHCF244* restored the photosynthetic

growth and PSII activity of Δ*CrHCF244-T23* (Fig. 2B). The complementation of both the *ami6*

mutant and the CRISPR-Cas9 knockout mutant by *CrHCF244* verified that this gene is required

for PSII function. The function is consistent with its homolog in plants (Chotewutmontri *et al.*,

2020; Link *et al.*, 2012).

Mutation of CrHCF244 specifically affects accumulation of PSII among photosynthetic

complexes

To test the accumulation of photosynthetic complexes in the absence of CrHCF244,

immunoblotting of subunits of PSII, PSI and Cyt b_6f was performed. As shown in Fig. 3A, PSII

core subunits including D1, D2, CP43, CP47 were drastically reduced in the *ami6* mutant and

Δ*CrHCF244-T23* mutant, whereas the abundance of subunits of PSI and Cyt b_6f did not change

in the mutants. The extrinsic subunit PsbO also showed a decrease in abundance, though to a

lesser level than those core subunits. Introducing *CrHCF244* into the two mutants restored the

accumulation of PSII subunits, as shown in the complemented lines (Fig. 3A). With respect of

the PSII complexes, immunoblotting of the protein complexes separated on BN-PAGE showed

that none of the PSII complexes accumulated to a significant level in *ami6* compared to WT (Fig.

3B and C). In the complemented strains, PSII complexes were restored, as shown in Fig. S3. The

results showed absence of CrHCF244 impaired both accumulation of individual subunits and

assembly of PSII complexes.

Mutation of CrHCF244 appears to abolish psbA translation

We used ^{14}C-acetate pulse labeling to look at the synthesis of PSII subunits in *ami6* and

the CRISPR-Cas9 mutants. Fig. 4A showed that in the mutants the synthesis of D1 subunit is not

observable, whereas the synthesis of D2 and CP43 occurred similarly to WT. The synthesis of

CP47 was absent in the mutants. This is due to the assembly-controlled translation which

requires the presence of D1 for translation of CP47 (Minai *et al.*, 2006). The transcription of

psbA gene is slightly decreased in the mutant (Fig. 4B). This indicates that the decrease of D1

synthesis is not caused by defects in transcription but due to defects in translation.

Arabidopsis HCF244 complements the ami6 mutant

To test whether the *Arabidopsis* homolog gene could complement the *Chlamydomonas* mutant, we made a construct in which the sequence of the transit peptide of PsaD from *Chlamydomonas* (CrPsaD) was fused with the sequence of the mature *Arabidopsis* HCF244 protein (Fig. 5A). The construct was then transformed into *ami6* and transformants were selected under photoautotrophic conditions. The results (Fig. 5B) showed that *AtHCF244* can partially complement the *Chlamydomonas* mutant. Transgene expression was verified at the mRNA level by reverse transcription (RT)-PCR, as shown in Fig. 5C.

CrHCF244 localizes to chloroplast

Confocal microscopy was used to examine the subcellular localization of CrHCF244. The complementation strain *ami6-C2* (the same line used in Fig. 1A), which is the *ami6* mutant transformed by C2 construct (depicted in Fig. 1C) was used for microscopy. In this strain, CrHCF244 was tagged with fluorescent protein Venus at the C-terminus. As shown in Fig. 6, the Venus signal mostly overlapped with chlorophyll signal, indicating that CrHCF244 is localized in the chloroplast. The strain photosynthetically restored by the Venus-fused CrHCF244 represents the functional and native localization of the protein. Venus tagged CrHCF244 in the background of WT showed the similar localization (Wang *et al.*, 2023b).

Chlorophyll decreased in the absence of CrHCF244

To test the effect of the absence of *CrHCF244* on the accumulation of chlorophyll, we measured chlorophyll content in the Δ*CrHCF244-T23* mutant grown in the dark and under light, respectively. The total Chl and Chl a/b ratio are shown in Table 1. In the dark, the total Chl decreased around 40% in the mutant and recovered to similar levels of WT in the complemented strains. Under light, the chlorophyll in the mutant also decreased similarly as in the dark. This suggests CrHCF244 could be involved in chlorophyll biosynthesis in *Chlamydomonas*, directly or indirectly through one-helix protein 2 (OHP2), which is involved in chlorophyll binding to apo-D1 and chlorophyll biosynthesis in *Arabidopsis* and *Chlamydomonas* (Hey and Grimm, 2018; Wang *et al.*, 2023a). The change of chlorophyll in the Δ*CrHCF244-T23* mutant is similar to *ohp2* mutant of *Chlamydomonas* (Wang *et al.*, 2023a).

Suppressor mutations suggests an alternative pathway of D1 translational control

Interestingly, we observed colonies originating from Δ*CrHCF244-T23* on minimal
medium (HS) plates. Growth and F_v/F_m measurements (Fig. 7A) showed these strains have
partially restored PSII activity. PCR and Sanger sequencing of *CrHCF244* proved that the
CrHCF244 gene did not undergo secondary mutations in these colonies (Supplementary Fig.
S6A and Fig. S6B), indicating the colonies are likely suppressor mutants that have mutations in
other gene(s). This suggests an alternative pathway for D1 translation control. We were also able
to observe suppressor mutants in *ami6*, in which 27 genes besides *CrHCF244* have been deleted.
We propose the pathway for *psbA* translation control in Fig. S5, in which the alternative pathway
is mediated by an alternative translation factor (TF) that was repressed. The identity of the
suppressor gene(s) is not investigated here and will be of interest for future study.

PSII is not affected in CRISPR-Cas9 mutants of RB47 and RB60

To examine the relationship of CrHCF244 with the previously reported D1 translation
factors RB47 and RB60, we used CRISPR-Cas9 to generate mutants. The gene maps and Cas9-
gRNA RNP cleavage sites are shown in Fig. 8A. We were able to get one mutant for RB47 and
multiple mutants for RB60 from PCR screening of the transformants. Here, we used the one
mutant of RB47, named as *RB47* and three mutants of RB60, named as Δ*RB60-1*, Δ*RB60-2*,
Δ*RB60-3*. PCR identifications of the mutants are shown in Fig. S6A and *RB47* was further
verified by Sanger sequencing (Fig. S6B).

CRISPR-Cas9 editing of RB47 was supposed to introduce a premature stop codon in the
fourth exon of the gene (Fig. S6C). Unexpectedly, sequencing of the transcript in the mutant
revealed a splicing event that partially removed the insertion (Fig. S6E). The final result is a 23
amino acid insertion. *RB60* was successfully knocked out as evidenced at the DNA and protein
levels (Fig. 8C). We note that we were unsuccessful in raising an RB47 antibody.

Unexpectedly, we did not detect significant PSII activity or D1 accumulation defects in
the mutants (Fig. 8C). Our genetic analysis does not support the previously proposed roles of
RB47 and RB60 in D1 translation (Kim and Mayfield, 1997, 2002; Yohn *et al.*, 1998b). This

discrepancy may be due to the fact that previous identifications were mostly based on

biochemical data when targeted gene editings were not possible.

Discussion

The high tendency of *Chlamydomonas* to randomly insert DNA fragments into its

genome makes it very accessible for generating a mutant pool or library. Such genetic screenings

have facilitated the discovery of many protein factors involved in PSII biogenesis and repair

including Tba1 (Somanchi *et al.*, 2005), REP27 (Park *et al.*, 2007), Alb3 (Bellafiore *et al.*, 2002;

Gohre *et al.*, 2006; Ossenbuhl *et al.*, 2004), and RBD1 (Calderon *et al.*, 2013). Recently, high

throughput genetic screening and mapping has resulted in many other potential photosynthesis

genes to be identified (Fauser *et al.*, 2022; Kafri *et al.*, 2023; Li *et al.*, 2019a; Wakao *et al.*,

2021), which includes the components of photosynthetic apparatus, factors for their biogenesis

and maintenance, proteins regulating photosynthesis and other genes whose functions are not yet

characterized. In this study, using forward genetics, we identified a mutant, *ami6*, that is

defective in PSII (Fig. 1A). To understand PSII biogenesis and identify factors involved in this

process, we started by generating a mutant pool and screened it for photosynthetic growth

defective mutants using chlorophyll fluorescence to test the PSII activity and found one mutant

that does not have PSII activity (Fig. 1A). We then used whole genome sequencing of the mutant

and its corresponding WT to map the genomic variation. Analysis of the sequencing data

revealed an approximately 345 kb deletion on chromosome 2 (Supplementary Dataset S1) when

using the *Chlamydomonas reinhardtii* v5.6 genome (https://phytozome-

next.jgi.doe.gov/info/Creinhardtii_v5_6) as the reference genome in early 2022. Recently, we re-

analyzed the whole genome sequencing data using *Chlamydomonas reinhardtii* v6.1

(https://phytozome-next.jgi.doe.gov/info/CreinhardtiiCC_4532_v6_1) genome as the reference

genome and the deletion (~347 kb) was mapped onto chromosome 15 (Fig. 1B), due to the

significant improvements in assembly quality of the v6.1 genome. Analysis of the whole genome

sequencing showed a ~347 kb deletion, where there are 28 genes, on chromosome 15 in the *ami6*

mutant (Fig. 1B). We pinpointed the causative gene to be *Cre02.g142146*, or *CrHCF244*, the

homolog gene of *Arabidopsis HCF244* by complementation (Fig. 1A) and by reproducing the

mutant phenotype using a CRISPR knockout (Δ*CrHCF244-T23*) of the same gene (Fig. 2).

Both *ami6* and Δ*CrHCF244-T23* have no PSII activity (Fig. 1A and Fig. 2B).

Immunoblotting showed that major subunits of PSII are drastically reduced in both mutants,

whereas other photosynthetic complexes are not affected (Fig. 3A). The specific effect of

CrHCF244 on PSII is consistent with its homologs in plants (Chotewutmontri *et al.*, 2020; Link

et al., 2012). PSII biogenesis is a stepwise and highly coordinated process, and the accumulation

of subunits is tightly regulated (Baena-Gonzalez and Aro, 2002; Nickelsen and Rengstl, 2013;

Nixon *et al.*, 2010). It has been established that the translation of PSII subunits is controlled by

an assembly-controlled mode, termed as CES, in which presence of the D2 subunit is

prerequisite for translation of D1, and presence of D1 is prerequisite for translation of CP47

(Wollman *et al.*, 1999). The mechanism of CES is that the unassembled subunit will bind to the 5'

UTR of its own mRNA and repress its translation. For instance, the translated D1 protein, which

hasn't been assembled into any complex will bind to *psbA* mRNA and repress any more

translation of D1 protein. The repression will be released when D1 is assembled into D2-Cyt b_{559}

complex (Minai *et al.*, 2006). To know whether any defects in synthesis caused the reduced

accumulation of PSII subunits, we used ^{14}C-acetate pulse labeling and observed that the

synthesis of D1 is affected in the mutants (Fig. 4A). As expected, in the absence of D1, CP47

translation was abolished as well in the mutants (Fig. 4A). The data presented showed that

CrHCF244 is essential for PSII biogenesis by affecting *psbA* translation, consistent with its role

in land plants (Chotewutmontri *et al.*, 2020; Link *et al.*, 2012).

Due to the large differences in the *psbA* 5' UTR sequences (Shen *et al.*, 2001) and no

homolog of trans-regulating factors has been reported to be involved in *psbA* translation between

Chlamydomonas and *Arabidopsis*, it was not clearly known whether *psbA* translation control is

conserved in algae and plants. In this study, we show that *CrHCF244*, a homolog of *Arabidopsis*

HCF244 function in *psbA* translation, consistent with the function of *HCF244* in *Arabidopsis*

(Link *et al.*, 2012). The *Arabidopsis* ortholog complementation of the *Chlamydomonas* mutant

(Fig. 5B) also demonstrated the functional conservation of this gene from algae to plants,

consistent with the relatively high sequence identity (Fig. S7) of the proteins between the two

organisms. The other *psbA* translation factor identified in *Arabidopsis*, HCF173 (Schult *et al.*,

2007), is also possibly involved in *psbA* translation in *Chlamydomonas*. In a proteomics study, it

was noted that D1 accumulation was reduced in a *CrHCF173* mutant (Kafri *et al.*, 2023) which

suggests it could be a translation factor for *psbA*. Previously RB47 and RB60, were

biochemically identified as possible *psbA* translation factors in *Chlamydomonas*. However,

genetic evidence in this study showed that RB47 and RB60 play no major role in *psbA*

translation (Fig. 8B, 8C, 8D). No *Arabidopsis* homologs of these proteins functioning in *psbA*

translation have been identified. These data also support the hypothesis that similar factors are

being employed in *psbA* translation in the two organisms.

We observed less chlorophyll content in Δ*CrHCF244-T23* compared to the WT and

complemented lines (Table 1), which resembles the result found in the *Chlamydomonas ohp2*

mutant (Wang *et al.*, 2023a). In plants, OHP1 and OHP2 bind and deliver chlorophyll to apo-D1.

OHP1, OHP2, and HCF244 form a complex and there is interdependence of protein

accumulation among the components (Chotewutmontri and Barkan, 2020; Hey and Grimm, 2018;

Li *et al.*, 2019b; Myouga *et al.*, 2018). Future investigation will explore the relationship between

CrHCF244, OHP2, and chlorophyll biosynthesis in *Chlamydomonas*.

Previously, it was proposed that RB60 regulates the redox state of RB47, promoting its

binding to *psbA* mRNA and translation (Kim and Mayfield, 1997). Here we showed that neither

of these proteins is essential for *psbA* translation using genetic evidence (Fig. 8). The previous

conclusions were mostly based on *in vitro* biochemical data (Danon and Mayfield, 1991; Kim

and Mayfield, 1997; Yohn *et al.*, 1998a). There was no genetic study on RB60, and the genetic

analysis on RB47 used mutants with reduced accumulation of RB47 but no mutation in the *RB47*

gene (Bruick and Mayfield, 1999; Yohn *et al.*, 1998b), due to the limitation in targeted

mutagenesis of nuclear genes in *Chlamydomonas*. A previous scenario of unspecific binding of

psbA mRNA was observed in which RB38, originally identified as a *psbA* mRNA associating

protein (Danon and Mayfield, 1991), was later shown to function in *psbD* translation control

(Schwarz *et al.*, 2007).

In this study, we confirmed that CrHCF244 is a *psbA* translation factor. A previous study

identified Tba1 as a *psbA* translation factor (Somanchi *et al.*, 2005) in *Chlamydomonas*. There

was a suggestion that CrHCF173 is also a *psbA* translation factor based on the function of its

Arabidopsis homology and proteomics detection of reduced D1 accumulation in the

Chlamydomonas mutant (Kafri *et al.*, 2023). Therefore, there are at least two or three proteins

required for translation of *psbA in Chlamydomonas*, considering DLA2 is a moonlighting

regulator, which only functions in the presence of acetate (Bohne *et al.*, 2013; Neusius *et al.*,

2022). Questions arise as to why multiple components are required for the translation of one

message and how the absence of each of them would abolish the translation. They could function

in a complex with each of them required for the stability of the complex, like the translation

initiation complex in the cytosol of eukaryotes (Brito Querido *et al.*, 2024; Jackson *et al.*, 2010).

Alternatively, the translation initiation of *psbA* could be a sequential process and each of them is

needed in a stepwise way. Regarding the working mechanism of CrHCF244 in *psbA* translation,

this protein could directly bind to the 5' UTR of *psbA* or bind indirectly as mediated by other

proteins. There are proteins in the SDR superfamily shown to be RNA binding proteins

(Bollenbach and Stern, 2003). We tried to recombinantly express CrHCF244 and test its RNA

binding ability *in vitro* but failed to get soluble expression of the protein in *E. coli*.

Our observation of suppressor mutants of Δ*CrHCF244* suggests the presence of an

alternative complex for D1 translation or perhaps a step in the translation process could be

bypassed in these lines. The inability of the Δ*CrHCF244-T23* mutant to grow

photoautotrophically and the absence of PSII activity indicate this alternative pathway is in

repression. We propose a possible scenario as illustrated in Fig. S5. Another possibility is

mutation in the 5' UTR of the target gene in the background of a translation factor mutant would

(partially) restore the phenotype. The mutation in the 5' UTR led to a secondary structure change

which allowed access for ribosome binding in the absence of the translation factor. A suppressor

mutation of a translation factor of *psbD* was reported and the mutation is in the 5' UTR of the

psbD mRNA (Rochaix *et al.*, 1989). We sequenced the *psbA* 5' UTR of these suppressors and

there is no mutation, suggesting the mutation might involve nuclear gene product(s).

This study shows that CrHCF244 is essential for PSII biogenesis by affecting *psbA*

translation. This identification of a plant homolog involved in *psbA* translation demonstrates the

conservation of *psbA* translation control between algae and plants. Previously, it had been

speculated that *psbA* translation control might be distinct between algae and plants due to the

large difference in the 5' UTR of *psbA* genes, and no shared translation factor proteins had been

identified between the organisms. The establishment of the conservation of *psbA* translation

control will allow the use of *Chlamydomonas*, which has more feasible genetic and systematic

mutant screen tools to facilitate the understanding of this important process in plants. The

observation of partial suppressors of *CrHCF244* mutants suggests an alternative pathway in *psbA*

translation control. Future mapping of the mutation gene(s) in the suppressors and studies of the

plant homolog(s) will improve our understanding in *psbA* translation control.

Supplementary data

Supplementary data for this manuscript is listed as below and available online.

Acknowledgments

We thank Shared Instrumentation Facility at Louisiana State University for confocal microscopy support. We thank Prof. Alice Barkan for helpful discussions. We thank Prof. Jun Minagawa for the gift of the B-Hiss *Chlamydomonas* strain. We thank Prof. Terry Bricker for the Cyt f antibody. We thank Dr. Yuqing Hou at University of Massachusetts Medical School for her advice on CRISPR-Cas9 gene editing in *Chlamydomonas*. BioRender was used for making diagrams (Fig. 1C, Fig.2A, Fig. 5A, Fig. S5 and Fig. 8A)

Author contributions

Conceptualization: XW, JVM, DJV; Experiments: XW, LAC, CZ, JVM, DJV; Data analysis and interpretation: XW, GW, CZ, MD, JVM, DJV; Writing: original draft- XW, JVM, DJV, reviewing and editing- all authors; Supervision of study: DJV; Funding acquisition: DJV and MD.

Conflict of interest

The authors declare that they have no conflict of interests.

Funding

This work is supported by the U.S. Department of Energy, Office of Science, Office of Basic Energy Science, Division of Chemical Sciences, Geosciences, and Biosciences, Photosynthetic Systems through Grant DE-SC0025359. This work was also supported by the US National Science Foundation IOS-EDGE 1923589 and US Department of Energy BER DE-SC0022985 grants for genomic data analysis.

Data availability

All relevant *Chlamydomonas* strains will be archived in Chlamydomonas Resource Center. All relevant data and other materials are available from the authors upon request.

References

Auchincloss AH, Zerges W, Perron K, Girard-Bascou J, Rochaix JD. 2002. Characterization of Tbc2, a nucleus-encoded factor specifically required for translation of the chloroplast psbC mRNA in Chlamydomonas reinhardtii. Journal of Cell Biology 157, 953-962.

Bae S, Park J, Kim JS. 2014. Cas-OFFinder: a fast and versatile algorithm that searches for potential off-target sites of Cas9 RNA-guided endonucleases. Bioinformatics 30, 1473-1475.

Baena-Gonzalez E, Aro EM. 2002. Biogenesis, assembly and turnover of photosystem II units. Philos Trans R Soc Lond B Biol Sci 357, 1451-1459; discussion 1459-1460.

Bellafiore S, Ferris P, Naver H, Gohre V, Rochaix JD. 2002. Loss of Albino3 leads to the specific depletion of the light-harvesting system. The Plant Cell 14, 2303-2314.

Bennoun P, Spierer-Herz M, Erickson J, Girard-Bascou J, Pierre Y, Delosme M, Rochaix JD. 1986. Characterization of photosystem II mutants of Chlamydomonas reinhardii lacking the psbA gene. Plant Molecular Biology 6, 151-160.

Berthold P, Schmitt R, Mages W. 2002. An Engineered Streptomyces hygroscopicus aph 7″ Gene Mediates Dominant Resistance against Hygromycin B in Chlamydomonas reinhardtii. Protist 153, 401-412.

Bohne AV, Schwarz C, Schottkowski M, Lidschreiber M, Piotrowski M, Zerges W, Nickelsen J. 2013. Reciprocal regulation of protein synthesis and carbon metabolism for thylakoid membrane biogenesis. PLoS Biol 11, e1001482.

Bollenbach TJ, Stern DB. 2003. Divalent metal-dependent catalysis and cleavage specificity of CSP41, a chloroplast endoribonuclease belonging to the short chain dehydrogenase/reductase superfamily. Nucleic Acids Research 31, 4317-4325.

Brito Querido J, Diaz-Lopez I, Ramakrishnan V. 2024. The molecular basis of translation initiation and its regulation in eukaryotes. Nature Reviews Molecular Cell Biology 25, 168-186.

Bruick RK, Mayfield SP. 1999. Light-activated translation of chloroplast mRNAs. Trends in Plant Science 4, 190-195.

Calderon RH, de Vitry C, Wollman FA, Niyogi KK. 2023. Rubredoxin 1 promotes the proper folding of D1 and is not required for heme b(559) assembly in Chlamydomonas photosystem II. Journal of Biological Chemistry 299, 102968.

Calderon RH, Garcia-Cerdan JG, Malnoe A, Cook R, Russell JJ, Gaw C, Dent RM, de Vitry C, Niyogi KK. 2013. A conserved rubredoxin is necessary for photosystem II accumulation in diverse oxygenic photoautotrophs. Journal of Biological Chemistry 288, 26688-26696.

Cameron DL, Baber J, Shale C, Valle-Inclan JE, Besselink N, van Hoeck A, Janssen R, Cuppen E, Priestley P, Papenfuss AT. 2021. GRIDSS2: comprehensive characterisation of somatic structural variation using single breakend variants and structural variant phasing. Genome Biology 22, 202.

Cavaiuolo M, Kuras R, Wollman FA, Choquet Y, Vallon O. 2017. Small RNA profiling in Chlamydomonas: insights into chloroplast RNA metabolism. Nucleic Acids Research 45, 10783-10799.

Chotewutmontri P, Barkan A. 2020. Light-induced psbA translation in plants is triggered by photosystem II damage via an assembly-linked autoregulatory circuit. Proc Natl Acad Sci U S A 117, 21775-21784.

Chotewutmontri P, Williams-Carrier R, Barkan A. 2020. Exploring the Link between Photosystem II Assembly and Translation of the Chloroplast psbA mRNA. Plants (Basel, Switzerland) 9.

Chua N-H, Gillham NW. 1977. The sites of synthesis of the principal thylakoid membrane polypeptides in Chlamydomonas reinhardtii. The Journal of cell biology 74, 441-452.

Danon A, Mayfield SP. 1991. Light regulated translational activators: identification of chloroplast gene specific mRNA binding proteins. The EMBO Journal 10, 3993-4001.

Drapier D, Girard-Bascou J, Wollman FA. 1992. Evidence for Nuclear Control of the Expression of the atpA and atpB Chloroplast Genes in Chlamydomonas. The Plant Cell 4, 283-295.

Eberhard S, Loiselay C, Drapier D, Bujaldon S, Girard-Bascou J, Kuras R, Choquet Y, Wollman FA. 2011. Dual functions of the nucleus-encoded factor TDA1 in trapping and translation activation of atpA transcripts in Chlamydomonas reinhardtii chloroplasts. The Plant Journal 67, 1055-1066.

Fauser F, Vilarrasa-Blasi J, Onishi M, Ramundo S, Patena W, Millican M, Osaki J, Philp C, Nemeth M, Salome PA, Li X, Wakao S, Kim RG, Kaye Y, Grossman AR, Niyogi KK, Merchant SS, Cutler SR, Walter P, Dinneny JR, Jonikas MC, Jinkerson RE. 2022. Systematic characterization of gene function in the photosynthetic alga Chlamydomonas reinhardtii. Nature Genetics 54, 705-714.

Garcia-Cerdan JG, Furst AL, McDonald KL, Schunemann D, Francis MB, Niyogi KK. 2019. A thylakoid membrane-bound and redox-active rubredoxin (RBD1) functions in de novo assembly and repair of photosystem II. Proc Natl Acad Sci U S A 116, 16631-16640.

Genty B, Briantais JM, Baker NR. 1989. The Relationship between the Quantum Yield of Photosynthetic Electron-Transport and Quenching of Chlorophyll Fluorescence. Biochim Biophys Acta 990, 87-92.

Gibson DG, Young L, Chuang RY, Venter JC, Hutchison CA, 3rd, Smith HO. 2009. Enzymatic assembly of DNA molecules up to several hundred kilobases. Nature Methods 6, 343-345.

Girard-Bascou J, Pierre Y, Drapier D. 1992. A nuclear mutation affects the synthesis of the chloroplast psbA gene production Chlamydomonas reinhardtii. Current Genetics 22, 47-52.

Gohre V, Ossenbuhl F, Crevecoeur M, Eichacker LA, Rochaix JD. 2006. One of two alb3 proteins is essential for the assembly of the photosystems and for cell survival in Chlamydomonas. The Plant Cell 18, 1454-1466.

Harris EH, Stern DB, Witman G, Harris EH, Harris EH. 2009. *The Chlamydomonas sourcebook*. Amsterdam ; Boston: Academic Press.

Heitzer M, Zschoernig B. 2007. Construction of modular tandem expression vectors for the green alga Chlamydomonas reinhardtii using the Cre/lox-system. BioTechniques 43, 324, 326, 328 passim.

Hey D, Grimm B. 2018. ONE-HELIX PROTEIN2 (OHP2) Is Required for the Stability of OHP1 and Assembly Factor HCF244 and Is Functionally Linked to PSII Biogenesis. Plant Physiol 177, 1453-1472.

Jackson RJ, Hellen CU, Pestova TV. 2010. The mechanism of eukaryotic translation initiation and principles of its regulation. Nature Reviews Molecular Cell Biology 11, 113-127.

Järvi S, Suorsa M, Aro EM. 2015. Photosystem II repair in plant chloroplasts--Regulation, assisting proteins and shared components with photosystem II biogenesis. Biochim Biophys Acta 1847, 900-909.

Kafri M, Patena W, Martin L, Wang L, Gomer G, Ergun SL, Sirkejyan AK, Goh A, Wilson AT, Gavrilenko SE, Breker M, Roichman A, McWhite CD, Rabinowitz JD, Cross FR, Wuhr M, Jonikas MC. 2023. Systematic identification and characterization of genes in the regulation and biogenesis of photosynthetic machinery. Cell 186, 5638-5655 e5625.

Kavanagh KL, Jornvall H, Persson B, Oppermann U. 2008. Medium- and short-chain dehydrogenase/reductase gene and protein families : the SDR superfamily: functional and structural diversity within a family of metabolic and regulatory enzymes. Cellular and Molecular Life Sciences 65, 3895-3906.

Kim J, Mayfield SP. 1997. Protein disulfide isomerase as a regulator of chloroplast translational activation. Science 278, 1954-1957.

Kim J, Mayfield SP. 2002. The active site of the thioredoxin-like domain of chloroplast protein disulfide isomerase, RB60, catalyzes the redox-regulated binding of chloroplast poly(A)-binding protein, RB47, to the 5' untranslated region of psbA mRNA. The Plant Cell Physiol 43, 1238-1243.

Langmead B, Salzberg SL. 2012. Fast gapped-read alignment with Bowtie 2. Nature Methods 9, 357-359.

Lefebvre-Legendre L, Choquet Y, Kuras R, Loubery S, Douchi D, Goldschmidt-Clermont M. 2015. A nucleus-encoded chloroplast protein regulated by iron availability governs expression of the photosystem I subunit PsaA in Chlamydomonas reinhardtii. Plant Physiol 167, 1527-1540.

Li X, Patena W, Fauser F, Jinkerson RE, Saroussi S, Meyer MT, Ivanova N, Robertson JM, Yue R, Zhang R, Vilarrasa-Blasi J, Wittkopp TM, Ramundo S, Blum SR, Goh A, Laudon M, Srikumar T, Lefebvre PA, Grossman AR, Jonikas MC. 2019a. A genome-wide algal mutant library and functional screen identifies genes required for eukaryotic photosynthesis. Nature Genetics 51, 627-635.

Li Y, Liu B, Zhang J, Kong F, Zhang L, Meng H, Li W, Rochaix JD, Li D, Peng L. 2019b. OHP1, OHP2, and HCF244 Form a Transient Functional Complex with the Photosystem II Reaction Center. Plant Physiol 179, 195-208.

Link S, Engelmann K, Meierhoff K, Westhoff P. 2012. The atypical short-chain dehydrogenases HCF173 and HCF244 are jointly involved in translational initiation of the psbA mRNA of Arabidopsis. Plant Physiol 160, 2202-2218.

Lu Y. 2016. Identification and Roles of Photosystem II Assembly, Stability, and Repair Factors in Arabidopsis. Frontiers in Plant Science 7, 168.

Mackinder LC, Meyer MT, Mettler-Altmann T, Chen VK, Mitchell MC, Caspari O, Freeman Rosenzweig ES, Pallesen L, Reeves G, Itakura A, Roth R, Sommer F, Geimer S, Muhlhaus T, Schroda M, Goodenough U, Stitt M, Griffiths H, Jonikas MC. 2016. A repeat protein links Rubisco to form the eukaryotic carbon-concentrating organelle. Proc Natl Acad Sci U S A 113, 5958-5963.

Merchant SS, Prochnik SE, Vallon O, Harris EH, Karpowicz SJ, Witman GB, Terry A, Salamov A, Fritz-Laylin LK, Marechal-Drouard L, Marshall WF, Qu LH, Nelson DR, Sanderfoot AA, Spalding MH, Kapitonov VV, Ren Q, Ferris P, Lindquist E, Shapiro H, Lucas SM, Grimwood J, Schmutz J, Cardol P, Cerutti H, Chanfreau G, Chen CL, Cognat V, Croft MT, Dent R, Dutcher S, Fernandez E, Fukuzawa H, Gonzalez-Ballester D, Gonzalez-Halphen D, Hallmann A, Hanikenne M, Hippler M, Inwood W, Jabbari K, Kalanon M, Kuras R, Lefebvre PA, Lemaire SD, Lobanov AV, Lohr M, Manuell A, Meier I, Mets L, Mittag M, Mittelmeier T, Moroney JV, Moseley J, Napoli C, Nedelcu AM, Niyogi K, Novoselov SV, Paulsen IT, Pazour G, Purton S, Ral JP, Riano-Pachon DM, Riekhof W, Rymarquis L, Schroda M, Stern D, Umen J, Willows R, Wilson N, Zimmer SL, Allmer J, Balk J, Bisova K, Chen CJ, Elias M, Gendler K, Hauser C, Lamb MR, Ledford H, Long JC, Minagawa J, Page MD, Pan J, Pootakham W, Roje S, Rose A, Stahlberg E, Terauchi AM, Yang P, Ball S, Bowler C, Dieckmann CL, Gladyshev VN, Green P, Jorgensen R, Mayfield S, Mueller-Roeber B, Rajamani S, Sayre RT, Brokstein P, Dubchak I, Goodstein D, Hornick L, Huang YW, Jhaveri J, Luo Y, Martinez D, Ngau WC, Otillar B, Poliakov A, Porter A, Szajkowski L, Werner G, Zhou K, Grigoriev IV, Rokhsar DS, Grossman AR. 2007. The *Chlamydomonas* genome reveals the evolution of key animal and plant functions. Science 318, 245-250.

Minai L, Wostrikoff K, Wollman FA, Choquet Y. 2006. Chloroplast biogenesis of photosystem II cores involves a series of assembly-controlled steps that regulate translation. The Plant Cell 18, 159-175.

Moummou H, Kallberg Y, Tonfack LB, Persson B, van der Rest B. 2012. The plant short-chain dehydrogenase (SDR) superfamily: genome-wide inventory and diversification patterns. BMC Plant Biology 12, 219.

Mulo P, Sakurai I, Aro EM. 2012. Strategies for psbA gene expression in cyanobacteria, green algae and higher plants: from transcription to PSII repair. Biochim Biophys Acta 1817, 247-257.

Myouga F, Takahashi K, Tanaka R, Nagata N, Kiss AZ, Funk C, Nomura Y, Nakagami H, Jansson S, Shinozaki K. 2018. Stable Accumulation of Photosystem II Requires ONE-HELIX PROTEIN1 (OHP1) of the Light Harvesting-Like Family. Plant Physiol 176, 2277-2291.

Neusius D, Kleinknecht L, Teh JT, Ostermeier M, Kelterborn S, Eirich J, Hegemann P, Finkemeier I, Bohne AV, Nickelsen J. 2022. Lysine acetylation regulates moonlighting activity of the E2 subunit of the chloroplast pyruvate dehydrogenase complex in Chlamydomonas. The Plant Journal 111, 1780-1800.

Newman SM, Boynton JE, Gillham NW, Randolph-Anderson BL, Johnson AM, Harris EH. 1990. Transformation of chloroplast ribosomal RNA genes in Chlamydomonas: molecular and genetic characterization of integration events. Genetics 126, 875-888.

Nickelsen J, Rengstl B. 2013. Photosystem II assembly: from cyanobacteria to plants. Annual Review of Plant Biology 64, 609-635.

Nixon PJ, Michoux F, Yu J, Boehm M, Komenda J. 2010. Recent advances in understanding the assembly and repair of photosystem II. Annals of Botany 106, 1-16.

O'Connor BD, van der Auwera G. 2020. *Genomics in the Cloud: Using Docker, GATK, and WDL in Terra*: O'Reilly Media, Incorporated.

Oppermann UC, Filling C, Jornvall H. 2001. Forms and functions of human SDR enzymes. Chemico-Biological Interactions 130-132, 699-705.

Ossenbuhl F, Gohre V, Meurer J, Krieger-Liszkay A, Rochaix JD, Eichacker LA. 2004. Efficient assembly of photosystem II in Chlamydomonas reinhardtii requires Alb3.1p, a homolog of Arabidopsis ALBINO3. The Plant Cell 16, 1790-1800.

Ossenbuhl F, Hartmann K, Nickelsen J. 2002. A chloroplast RNA binding protein from stromal thylakoid membranes specifically binds to the 5' untranslated region of the psbA mRNA. Eur J Biochem 269, 3912-3919.

Park S, Khamai P, Garcia-Cerdan JG, Melis A. 2007. REP27, a tetratricopeptide repeat nuclear-encoded and chloroplast-localized protein, functions in D1/32-kD reaction center protein turnover and photosystem II repair from photodamage. Plant Physiol 143, 1547-1560.

Picariello T, Hou Y, Kubo T, McNeill NA, Yanagisawa HA, Oda T, Witman GB. 2020. TIM, a targeted insertional mutagenesis method utilizing CRISPR/Cas9 in *Chlamydomonas reinhardtii*. PLoS One 15, e0232594.

Porra R, Thompson W, Kriedemann P. 1989. Determination of accurate extinction coefficients and simultaneous equations for assaying chlorophylls a and b extracted with four different solvents: verification of the concentration of chlorophyll standards by atomic absorption spectroscopy. Biochimica et Biophysica Acta (BBA)-Bioenergetics 975, 384-394.

Rochaix JD, Kuchka M, Mayfield S, Schirmer-Rahire M, Girard-Bascou J, Bennoun P. 1989. Nuclear and chloroplast mutations affect the synthesis or stability of the chloroplast psbC gene product in Chlamydomonas reinhardtii. The EMBO Journal 8, 1013-1021.

Schult K, Meierhoff K, Paradies S, Toller T, Wolff P, Westhoff P. 2007. The nuclear-encoded factor HCF173 is involved in the initiation of translation of the psbA mRNA in Arabidopsis thaliana. The Plant Cell 19, 1329-1346.

Schwarz C, Elles I, Kortmann J, Piotrowski M, Nickelsen J. 2007. Synthesis of the D2 protein of photosystem II in Chlamydomonas is controlled by a high molecular mass complex containing the RNA stabilization factor Nac2 and the translational activator RBP40. The Plant Cell 19, 3627-3639.

Shen JR. 2015. The Structure of Photosystem II and the Mechanism of Water Oxidation in Photosynthesis. Annual Review of Plant Biology 66, 23-48.

Shen Y, Danon A, Christopher DA. 2001. RNA binding-proteins interact specifically with the Arabidopsis chloroplast psbA mRNA 5' untranslated region in a redox-dependent manner. The Plant Cell Physiol 42, 1071-1078.

Somanchi A, Barnes D, Mayfield SP. 2005. A nuclear gene of Chlamydomonas reinhardtii, Tba1, encodes a putative oxidoreductase required for translation of the chloroplast psbA mRNA. The Plant Journal 42, 341-352.

Spaniol B, Lang J, Venn B, Schake L, Sommer F, Mustas M, Geimer S, Wollman FA, Choquet Y, Muhlhaus T, Schroda M. 2022. Complexome profiling on the Chlamydomonas lpa2 mutant reveals insights into PSII biogenesis and new PSII associated proteins. Journal of Experimental Botany 73, 245-262.

Sueoka N. 1960. Mitotic Replication of Deoxyribonucleic Acid in Chlamydomonas Reinhardi. Proc Natl Acad Sci U S A 46, 83-91.

Suzuki T, Minagawa J, Tomo T, Sonoike K, Ohta H, Enami I. 2003. Binding and Functional Properties of the Extrinsic Proteins in Oxygen-Evolving Photosystem II Particle from a Green Alga, Chlamydomonas reinhardtii having His-tagged CP47. Plant and Cell Physiology 44, 76-84.

Tokutsu R, Kato N, Bui KH, Ishikawa T, Minagawa J. 2012. Revisiting the supramolecular organization of photosystem II in Chlamydomonas reinhardtii. Journal of Biological Chemistry 287, 31574-31581.

Vaistij FE, Boudreau E, Lemaire SD, Goldschmidt-Clermont M, Rochaix JD. 2000. Characterization of Mbb1, a nucleus-encoded tetratricopeptide-like repeat protein required for expression of the chloroplast psbB/psbT/psbH gene cluster in Chlamydomonas reinhardtii. Proc Natl Acad Sci U S A 97, 14813-14818.

Viola S, Cavaiuolo M, Drapier D, Eberhard S, Vallon O, Wollman FA, Choquet Y. 2019. MDA1, a nucleus-encoded factor involved in the stabilization and processing of the atpA transcript in the chloroplast of Chlamydomonas. The Plant Journal 98, 1033-1047.

Wakao S, Shih PM, Guan K, Schackwitz W, Ye J, Patel D, Shih RM, Dent RM, Chovatia M, Sharma A, Martin J, Wei CL, Niyogi KK. 2021. Discovery of photosynthesis genes through whole-genome sequencing of acetate-requiring mutants of *Chlamydomonas reinhardtii*. PLoS Genet 17, e1009725.

Wang F, Dischinger K, Westrich LD, Meindl I, Egidi F, Trösch R, Sommer F, Johnson X, Schroda M, Nickelsen J, Willmund F, Vallon O, Bohne A-V. 2023a. One-helix protein 2 is not required for the synthesis of photosystem II subunit D1 in Chlamydomonas. Plant Physiologyogy 191, 1612-1633.

Wang F, Johnson X, Cavaiuolo M, Bohne AV, Nickelsen J, Vallon O. 2015. Two Chlamydomonas OPR proteins stabilize chloroplast mRNAs encoding small subunits of photosystem II and cytochrome b6 f. The Plant Journal 82, 861-873.

Wang L, Patena W, Van Baalen KA, Xie Y, Singer ER, Gavrilenko S, Warren-Williams M, Han L, Harrigan HR, Hartz LD, Chen V, Ton V, Kyin S, Shwe HH, Cahn MH, Wilson AT, Onishi M, Hu J, Schnell DJ, McWhite CD, Jonikas MC. 2023b. A chloroplast protein atlas reveals punctate structures and spatial organization of biosynthetic pathways. Cell 186, 3499-3518 e3414.

Wang X, Gao F, Zhang J, Zhao J, Ogawa T, Ma W. 2016. A Cytoplasmic Protein Ssl3829 Is Important for NDH-1 Hydrophilic Arm Assembly in Synechocystis sp. Strain PCC 6803. Plant Physiol 171, 864-877.

Wollman F-A, Minai L, Nechushtai R. 1999. The biogenesis and assembly of photosynthetic proteins in thylakoid membranes1This article is dedicated to the memory of our colleague and friend, Alma Gal. Her presence played a major role in our decision to prepare this review article.1. Biochimica et Biophysica Acta (BBA) - Bioenergetics 1411, 21-85.

Yohn CB, Cohen A, Danon A, Mayfield SP. 1996. Altered mRNA binding activity and decreased translational initiation in a nuclear mutant lacking translation of the chloroplast psbA mRNA. Molecular and Cellular Biology 16, 3560-3566.

Yohn CB, Cohen A, Danon A, Mayfield SP. 1998a. A poly(A) binding protein functions in the chloroplast as a message-specific translation factor. Proceedings of the National Academy of Sciences USA 95, 2238-2243.

Yohn CB, Cohen A, Rosch C, Kuchka MR, Mayfield SP. 1998b. Translation of the chloroplast psbA mRNA requires the nuclear-encoded poly(A)-binding protein, RB47. Journal of Cell Biology 142, 435-442.

Zhang R, Patena W, Armbruster U, Gang SS, Blum SR, Jonikas MC. 2014. High-Throughput Genotyping of Green Algal Mutants Reveals Random Distribution of Mutagenic Insertion Sites and Endonucleolytic Cleavage of Transforming DNA. The Plant Cell 26, 1398-1409.

Table 1. Chlorophyll content and Chl a/b ratios in WT, Δ*CrHCF244-T23* and complemented lines. Cells from mixotrophic cultures under light intensity of 35 µmol photons m^{-2} s^{-1} and from heterotrophic cultures in the dark (grey area) in the early exponential phase were used for measurement. Data are presented as means ± standard deviation (n = 3).

		[Chl] µg mL^{-1} OD_{750}^{-1}	Chl-a/b
light	WT	33.5 ± 0.8	3.04 ± 0.01
	Δ*CrHCF244-T23*	21.7 ± 0.6	2.67 ± 0.05
	Δ*CrHCF244-T23-C1*	34.6 ± 1.4	3.08 ± 0.03
	Δ*CrHCF244-T23-C2*	33.2 ± 1.7	3.10 ± 0.03
dark	WT	27.5 ± 0.4	2.75 ± 0.17
	Δ*CrHCF244-T23*	16.9 ± .07	2.67 ± 0.14
	Δ*CrHCF244-T23-C1*	28.0 ± 2.0	2.71 ± 0.11
	Δ*CrHCF244-T23-C2*	28.8 ± 3.0	2.74 ± 0.15

Figure legends:

Figure 1. ami6 growth, PSII activity, and the mapping of genomic variation and the causative gene.

(A) Growth of wild type (WT), *ami6* mutant and two complemented lines of *ami6*, under mixotrophic (TAP) and photoautotrophic (HS) conditions. *ami6-C1* is a representative transformant from transforming *ami6* with the C1 construct (illustrated in **(C)**); *ami6-C2* is a representative transformant from transforming *ami6* with the C2 construct (illustrated in **(C)**). F_v/F_m values are shown for cultures grown on TAP. Spot test results and F_v/F_m values are representative of two replicates. **(B)** Mapping of *ami6* genomic variation by whole genome sequencing revealed a ~347 kb deletion on chromosome 15. **(C)** Illustration of genetic constructs for complementing *ami6* mutant. The C1 and C2 constructs both are under control of the *psaD* promoter (*PpsaD*). The constructs were used for transforming untagged CrHCF244 and Venus-tagged CrHCF244 into the *ami6* mutant.

Figure 2. Growth and PSII activity of a CRISPR-Cas9 knockout CrHCF244 mutant.

(A) *CrHCF244* gene map (5' UTR as green box, exons as blue boxes, 3' UTR as purple box) indicating the gRNA targeting site at exon 2 for CRISPR-Cas9 knockout/gene editing. **(B)** Growth of WT, CRISPR-Cas9 knockout *CrHCF244* mutant Δ*CrHCF244-T23* (shown as *T23* in **(B)** and **(C)**) and the two complemented strains (shown as *T23-C1* and *T23-C2* in **(B)** and **(C)**), which are representative transformants from transforming Δ*CrHCF244-T23* with C1 and C2 construct (illustrated in Fig. 1C), respectively, under mixotrophic (TAP) and photoautotrophic (HS) conditions. **(C)** F_v/F_m values for strains grown on TAP. Spot test results and F_v/F_m values are representative of two replicates.

Figure 3. Accumulation of photosynthetic protein complexes.

(A) Immunoblots of photosynthetic proteins in WT, *ami6* and its two complemented lines (*ami6-C1*, *ami6-C2*), Δ*CrHCF244-T23* and its two complemented lines (*T23-C1*, *T23-C2*) and Fud7 (*psbA* deletion mutant). Coomassie brilliant blue (CBB) stain was shown as a loading control. Data are representative of two biological replicates. **(B)** Blue-Native PAGE (BN-PAGE) separation of photosynthetic protein complexes. Arrows indicate the positions of different PSII

complexes. **(C)** Immunoblotting analysis using D1 antibody of the PSII complexes. The immunoblotting was performed after transferring the protein complexes on the BN-PAGE to PVDF membrane. Arrows indicate PSII complexes, I and II: PSII super-complexes, III: PSII dimer, IV: PSII monomer.

Figure 4. D1 synthesis and psbA transcription.

(A) [14]C-acetate pulse-labeling of WT, *ami6* mutant, Fud7 (*psbA* deletion) and CRISPR-Cas9 knockout mutant Δ*CrHCF244-T23*. Cells were labeled for 20 min in the presence of 10 µg/mL cycloheximide to inhibit cytosol translation. PSII major subunits are shown by the arrows. Data are representative of two biological replicates. **(B)** Quantitative real time PCR (qRT-PCR) to detect *psbA* transcription change in WT, Δ*CrHCF244-T23* and its two complemented lines (*T23-C1*, *T23-C2*). Data are presented as means ± Standard Deviation (n = 4). *** indicates p<0.001 and ns indicates non-significant with One-way ANOVA. The sequences of primers are shown in Table S1.

Figure 5. Complementation of Chlamydomonas ami6 mutant with the Arabidopsis HCF244 gene.

(A) Schematic of the construct for transforming *ami6* with the *Arabidopsis HCF244* gene. Mature AtHCF244 protein (mAtHCF244) in fusion with a chloroplast transit peptide (cTP) derived from PsaD in *Chlamydomonas* was expressed under control of *psaD* promoter (*PpsaD*) **(B)** Growth and PSII activity of WT, *ami6* mutant, and the two complemented strains under mixotrophic (TAP) and photoautotrophic (HS) conditions. F_v/F_m values are shown for strains grown on TAP. Spot test results and F_v/F_m values are representative of two replicates. **(C)** RT-PCR to detect the expression of the transgene *AtHCF244* in the complemented lines. Primers are listed in Table S1. Actin is used as a loading control. PCR of RNA without reverse transcriptase (-RT) is used to confirm there is no genomic DNA contamination.

Figure 6. Chloroplast localization of CrHCF244.

Venus-tagged CrHCF244 plasmid was transformed into the *ami6* mutant. The complemented line *ami6-C2* was grown under photosynthetic condition before imaging. WT strain was imaged under the same conditions.

Figure 7. Suppressor mutants of ΔCrHCF244-T23 suggest an alternative pathway for D1 translation control.

(A) Growth and PSII activity of strains under mixotrophic conditions. WT, CRISPR-Cas9 knockout mutant Δ*CrHCF244-T23* (labeled as *T23* here) and three suppressor mutants (labeled as *T23-Su1*, *T23-Su2*, *T23-Su3*). **(B)** Growth and PSII activity of strains, as in (A), under photoautotrophic conditions. Spot test results and F_v/F_m values are representative of two replicates.

Figure 8. Growth and PSII activity of RB47 and RB60 mutants.

(A) Gene maps of the *RB47* and *RB60* showing the CRISPR-Cas9 cleavage sites. **(B)** Growth of the *RB47* mutant (*RB47*, see description in text) and RB60 (Δ*RB60-1*, Δ*RB60-2*, Δ*RB60-3*) mutants generated by CRISPR-Cas9, under photoautotrophic conditions. PSII activity was measured as F_v/F_m. Spot test results and Fv/Fm values are representative of two replicates. **(C)** Abundance of D1 subunit in the mutants. Immunoblotting of D1 subunit and RB60 was performed in the strains; Coomassie brilliant blue (CBB) stain was shown as a loading control.

Figure 1

Figure 1. ami6 growth, PSII activity, and the mapping of genomic variation and the causative gene.

(A) Growth of wild type (WT), *ami6* mutant and two complemented lines of *ami6*, under mixotrophic (TAP) and photoautotrophic (HS) conditions. *ami6-C1* is a representative transformant from transforming *ami6* with the C1 construct (illustrated in **(C)**); *ami6-C2* is a representative transformant from transforming *ami6* with the C2 construct (illustrated in **(C)**). F_v/F_m values are shown for cultures grown on TAP. Spot test results and F_v/F_m values are representative of two replicates. **(B)** Mapping of *ami6* genomic variation by whole genome sequencing revealed a ~347 kb deletion on chromosome 15. **(C)** Illustration of genetic constructs for complementing *ami6* mutant. The C1 and C2 constructs both are under control of the *psaD* promoter (*PpsaD*). The constructs were used for transforming untagged CrHCF244 and Venus-tagged CrHCF244 into the *ami6* mutant.

Figure 2

Figure 2. Growth and PSII activity of a CRISPR-Cas9 knockout CrHCF244 mutant.

(A) *CrHCF244* gene map (5' UTR as green box, exons as blue boxes, 3' UTR as purple box) indicating the gRNA targeting site at exon 2 for CRISPR-Cas9 knockout/gene editing. **(B)** Growth of WT, CRISPR-Cas9 knockout *CrHCF244* mutant Δ*CrHCF244-T23* (shown as *T23*) and the two complemented strains (shown as *T23-C1* and *T23-C2*), which are representative transformants from transforming Δ*CrHCF244-T23* with C1 and C2 construct (illustrated in Fig. 1C), respectively, under mixotrophic (TAP) and photoautotrophic (HS) conditions. F_v/F_m values were measured for strains grown on TAP. Spot test results and F_v/F_m values are representative of two replicates.

Figure 3

Figure 3. Accumulation of photosynthetic protein complexes.

(A) Immunoblots of photosynthetic proteins in WT, *ami6* and its two complemented lines (*ami6-C1*, *ami6-C2*), Δ*CrHCF244-T23* and its two complemented lines (*T23-C1*, *T23-C2*) and Fud7 (*psbA* deletion mutant). Coomassie brilliant blue (CBB) stain was shown as a loading control. Data are representative of two biological replicates. **(B)** Blue-Native PAGE (BN-PAGE) separation of photosynthetic protein complexes. Arrows indicate the positions of different PSII complexes. **(C)** Immunoblotting analysis using D1 antibody of the PSII complexes. The immunoblotting was performed after transferring the protein complexes on the BN-PAGE to PVDF membrane. Arrows indicate PSII complexes, I and II: PSII super-complexes, III: PSII dimer, IV: PSII monomer.

Figure 4

Figure 4. D1 synthesis and psbA transcription.

(A) ^{14}C-acetate pulse-labeling of WT, *ami6* mutant, Fud7 (*psbA* deletion) and CRISPR-Cas9 knockout mutant Δ*CrHCF244-T23*. Cells were labeled for 20 min in the presence of 10 μg/mL cycloheximide to inhibit cytosol translation. PSII major subunits are shown by the arrows. Data are representative of two biological replicates. **(B)** Quantitative real time PCR (qRT-PCR) to detect *psbA* transcription change in WT, Δ*CrHCF244-T23* and its two complemented lines (*T23-C1*, *T23-C2*). Data are presented as means ± Standard Deviation (n = 4). *** indicates p<0.001 and ns indicates non-significant with One-way ANOVA. The sequences of primers are shown in Table S1.

Figure 5

Figure 5. Complementation of Chlamydomonas ami6 mutant with the Arabidopsis HCF244 gene.

(A) Schematic of the construct for transforming *ami6* with the *Arabidopsis HCF244* gene. Mature AtHCF244 protein (mAtHCF244) in fusion with a chloroplast transit peptide (cTP) derived from PsaD in *Chlamydomonas* was expressed under control of *psaD* promoter (*PpsaD*) **(B)** Growth and PSII activity of WT, *ami6* mutant, and the two complemented strains under mixotrophic (TAP) and photoautotrophic (HS) conditions. F_v/F_m values are shown for strains grown on TAP. Spot test results and F_v/F_m values are representative of two replicates. **(C)** RT-PCR to detect the expression of the transgene *AtHCF244* in the complemented lines. Primers are listed in Table S1. Actin is used as a loading control. PCR of RNA without reverse transcriptase (-RT) is used to confirm there is no genomic DNA contamination.

Figure 6

Figure 6. Chloroplast localization of CrHCF244.

Venus-tagged CrHCF244 plasmid was transformed into the *ami6* mutant. The complemented line *ami6-C2* was grown under photosynthetic condition before imaging. WT strain was imaged under the same conditions.

Figure 7

Figure 7. Growth and PSII activity of suppressor mutants of ΔCrHCF244-T23

(A) Growth and PSII activity of strains under mixotrophic conditions. WT, CRISPR-Cas9 knockout mutant ΔCrHCF244-T23 (labeled as *T23* here) and three suppressor mutants (labeled as *T23-Su1*, *T23-Su2*, *T23-Su3*). **(B)** Growth and PSII activity of strains, as in (A), under photoautotrophic conditions. Spot test results and Fv/Fm values are representative of two replicates.

Figure 8

Figure 8. Growth and PSII activity of RB47 and RB60 mutants.

(A) Gene maps of the *RB47* and *RB60* showing the CRISPR-Cas9 cleavage sites. **(B)** Growth of the *RB47* mutant (*RB47*, see description in text) and RB60 (ΔRB60-1, ΔRB60-2, ΔRB60-3) mutants generated by CRISPR-Cas9, under photoautotrophic conditions. PSII activity was measured as F_v/F_m. Spot test results and Fv/Fm values are representative of two replicates. **(C)** Abundance of D1 subunit in the mutants. Immunoblotting of D1 subunit and RB60 was performed in the strains; Coomassie brilliant blue (CBB) stain was shown as a loading control.