1
2
3
4
5 **Latitudinal clines in floral display associated with adaptive evolution during a**
6 **biological invasion**
7
8
9
10

11 Mia N. Akbar[1], Dale Moskoff[1], Spencer C.H. Barrett[2], Robert I. Colautti[1,3]
12
13
14
15 [1]Department of Biology, Queen's University, Kingston, ON, Canada.
16 [2]Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, ON, Canada.
17
18
19 [3]Corresponding author: robert.colautti@queensu.ca

22
23 **Running head: Floral display clines in *Lythrum salicaria***
24
25

ABSTRACT

Premise: The timing and pattern of flowering -- flowering phenology -- strongly influences

reproductive success in plants. Days to first flower is easy to quantify and widely used to

characterize phenology, but reproductive fitness depends on the full schedule of flower

production over time.

Methods: We examined floral display evolution associated with rapid adaptive evolution and

range expansion among thirteen populations of *Lythrum salicaria* populations, sampled along a

10-degree latitudinal gradient in eastern North America. Growing these collections in a common

garden field experiment at a mid-latitude site, we used Principal Coordinates Analysis (PCoA)

and quantitative metrics analogous to central moments of probability distributions (i.e., mean,

variance, skew, and kurtosis) to quantify flowering schedules..

Key Results: Consistent with adaptation to shorter growing seasons at higher latitudes, we found

that populations from higher latitudes had earlier start and mean flowering day for both

population average (i.e., among individuals) and population aggregate (i.e., pooled flowers).

Clines in population average and aggregate flowering skew increased with latitude while kurtosis

decreased, consistent with a bet-hedging strategy arising in biotic environments with more

herbivores and greater competition for pollinators. Although clines were generally similar at the

average vs aggregate level, we found that individual flowering schedules from more southern

populations tended to flower later than their population aggregate.

Conclusions: Heritable genetic changes in flowering schedule are consistent with biotic and

abiotic responses to natural selection, potentially contributing to rapid evolution and range

expansion of this invasive species.

INTRODUCTION

Phenology affects how organisms interact with their environment, generating important consequences for their survival and reproduction. As targets of natural selection and determinants of population vital rates, the timing of growth, reproduction, and other phenological traits are crucial to understanding eco-evolutionary dynamics. For example, in sexual outcrossing plants, flowering phenology governs interaction with mutualists (Fenster et al., 2004), antagonists (Strauss & Whittall, 2006) and abiotic factors (Jin *et al.*, 2019). Consequently, plants experience spatial and temporal variation in natural selection on flowering phenology (Kudo, 2007), sometimes resulting in the evolution of local adaptation (Anderson et al., 2012; Fitchett et al., 2015; D. W. Inouye, 2022; Panchen, 2022; Panchen & Gorelick, 2017; Prather et al., 2023). As such, flowering phenology is a useful composite trait for investigating evolutionary responses to past and future environmental challenges.

Days to first flower (i.e., flowering time) is a simple yet informative metric with which to study the ecology and evolution of phenology in plant populations. For example, latitudinal and altitudinal clines in days to first flower are common to a variety of plant taxa (Alexander et al., 2009; Colautti & Lau, 2015; Halbritter et al., 2018), and reciprocal transplant experiments have confirmed that these clines are an important component of local adaptation (Ågren & Schemske, 2012; Anderson & Gezon, 2015; Colautti & Barrett, 2013; Ensing & Eckert, 2019; Griffith & Watson, 2005). Flowering time clines are consistent with local adaptation models characterized by stabilizing selection within populations but a shift in optimal flowering time along environmental gradients (de Villemereuil et al., 2020; Gauzere et al., 2020; Kirkpatrick & Barton, 1997). In contrast to the prediction of stabilizing selection, empirical measurements of selection suggest that flowering time is often under directional selection within populations

(Munguía-Rosas et al., 2011). Moreover, long-term phenology records show a variety of

responses to climate change -- from earlier to later to no change in flowering time, depending on

the species (Rafferty et al., 2020; Wolkovich et al., 2012). Thus, there appears to be

disagreement between theoretical models and empirical observations of geographic clines and

local adaptation on the one hand and observed measurements of natural selection and phenotypic

changes in natural populations on the other (Austen et al., 2017). Antagonistic selection on traits

correlated with the onset of flowering offers one potential resolution to this apparent

contradiction.

Day of first flowering may trade off with other aspects of flowering phenology such as

the shape of the flowering schedule. These other aspects of flowering phenology may not be well

represented by flowering time (Newstrom et al. 1994; CaraDonna et al., 2014; B. D. Inouye et

al., 2019) but have important effects on reproductive fitness (Fox, 2003). The shape of the

flowering schedule can be quantified by metrics analogous to central moments of probability

distributions, which include the mean, variance, skew, and kurtosis (Box 1). Not to be confused

with the central moments of a statistical population or sample, the analogous characteristics of a

flowering schedule quantify temporal variation in relative reproductive investment through time.

These characteristics use the same mathematical equations (Table 1) but are not shaped by

probabilistic process and therefore may not correlate with onset or duration of flowering (Fig.

1A). These metrics have rarely been used in studies of flowering phenology, yet they have the

potential to offer a novel perspective on selection and trade-offs affecting the evolution of

phenology.

Central moments of individual flowering schedules describe the timing and intensity of

reproductive investment, and thus are likely to experience strong selection from abiotic or biotic

features of the environment. We are not aware of studies quantifying evolutionary divergence of

entire flowering schedules among naturally occurring populations, but there are reasons to expect

populations to evolve differences. For example, individuals that experience competition for

pollinators later in the growing season may benefit from showy inflorescences that enhance

pollinator attraction (O'Neil, 1997; Thomson, 1980) and be characterized by a flowering

schedule with a more positive skew and longer duration. Herbivores and seed predators can be

unpredictable, which may select for a bet-hedging strategy and possibly delay in the onset of

flowering (Augspurger, 1981; Elzinga et al., 2007; Wright & Meagher, 2003). These conditions

might further result in a longer flowering duration with a higher variance and negative skew in

the flowering schedule if late-season ramp-up in flowering is limited by harsh growing

conditions at the end of a season.

In addition to biotic interactions, flowering schedules are likely to be shaped by abiotic

factors that determine the timing and duration of environmental conditions favorable to the

development of reproductive organs. Shorter growing seasons typically favour an earlier onset of

flowering (Austen & Weis, 2015; Colautti & Barrett, 2013; Griffith & Watson, 2005), and may

favour reproduction only within a shorter growing season; this would result in flowering

schedules with lower duration and variance. In marginal environments, plants may combine

strategies, concentrating flower production into a narrow window but also producing a few

flowers over a long timeframe as a bet-hedging strategy (Forrest & Thomson, 2010; Simons,

2011; Tufto, 2015), which would produce a long-tailed leptokurtic or skewed flowering

schedule. The above examples illustrate the complex ways that biotic interactions and the abiotic

environment could shape selection on individual flowering schedules, and how these differences

can be quantified using equations developed for probability distributions.

Natural selection acts on phenological schedules of individuals, but variation among

individuals within a population affect eco-evolutionary interactions with pollinators, herbivores,

and potential mates. Just as individuals have flowering schedules, the aggregated flower

production of all individuals within a population can be characterized (Box 1), which could yield

emergent properties that are relevant to understanding eco-evolutionary feedbacks. We define

emergent properties here as deviations in the aggregate population from the average of individual

schedules comprising it (Table 1). For example, a population can have a high aggregate variance

but low average individual variance in the flowering schedule if individuals vary in onset and

duration of flowering. In cases like this, the central moments of the aggregate flowering schedule

show characteristics that are emergent and distinct from the typical individual comprising it (Fig.

1). Although emergent properties can arise from adaptive processes acting on individuals, they

can also have important ecological consequences for species communities. For example,

population aggregate flowering schedules are sometimes positively skewed with an abrupt start

and a long tail of flowering (Blionis et al., 2001; Rabinowitz et al., 1981), which may be an

adaptation to enhance pollinator visitation to the population (Stemkovski et al., 2023; Thomson,

1980). Additionally, strong frequency-dependent selection may concentrate flowering within a

narrow but synchronous timeframe to attract more pollinators (Rathcke & Lacey, 1985),

increase mating opportunities (Ison & Weis, 2017), or prevent interspecific cross-pollination

(Fantinato et al., 2018). Therefore, in pollinator or mate-limited environments, population

aggregate flowering schedules may be very similar to the average flowering schedule. Thus,

emergent properties (or a lack thereof) have the potential to provide insight into eco-evolutionary

processes and ecological consequences of aggregate flowering schedules.

Environmental gradients associated with latitude can be useful for studying phenotypic

evolution because selection often varies predictably due to shorter growing seasons and weaker

biotic interactions towards polar latitudes (reviewed in Zvereva & Kozlov, 2021). Adaptation

along gradual environments is commonly manifested as clines in phenotypic traits that are

observable when genetically related individuals are grown in a common garden environment

(Olsson & Ågren, 2002; Panchen, 2022; Sobral et al., 2013). For example, there is good evidence

for the rapid evolution of adaptive flowering time clines in the perennial wetland herb *Lythrum*

salicaria (purple loosestrife) during its invasion of North America. Selection splines (Colautti &

Barrett, 2010), trait correlations (Colautti & Barrett, 2011), reciprocal transplants (Colautti &

Barrett, 2013), and herbarium records (Wu & Colautti, 2022) collectively support a model of

constrained selection in which shorter growing seasons toward the northern range limit favour

earlier flowering. In contrast, longer growing seasons in the southern part of the introduced range

relax selection on flowering time and favors plants that flower later but grow larger in stature.

However, by focusing on the first day of flowering, previous studies have ignored other aspects

of the flowering schedule that may be important for understanding adaptation and constraint

during evolution and invasion of *L. salicaria*.

Here, we investigate clines in flowering schedules among thirteen populations sampled

across ten degrees of latitude in eastern North America and grown in a common garden field

study at the Koffler Scientific Reserve north of Toronto, Ontario, Canada. We quantify both

individual and population aggregate flowering schedules to address the following questions: (1)

How well do common phenological metrics, like the onset and duration of flowering, predict

often overlooked features of the flowering schedule such as the variance, skew and kurtosis? (2)

Are there latitudinal clines in flowering schedules that are consistent with the adaptive

hypotheses outlined above? Specifically, we predicted that longer growing season lengths,

stronger competition for pollinators, and higher rates of herbivory in southern populations will

have favoured genotypes with: (i) a longer flowering duration and (ii) a more negative kurtosis

with a concentrated peak and rare flowering events far from the average flowering day, but also

(iii) an increase in the flowering schedule variance and (iv) a more negative skew with flowering

focused later in the schedule. (3) Are there emergent properties of population aggregate

flowering schedules that differ from the population average of individual flowering schedules?

Because natural selection acts on individual flowering schedules, emergent properties represent

(i) potential biases in evolutionary inference when aggregate characteristics are used in place of

individual or population average characteristics of flowering schedules, and (ii) ecological

phenomena that are shaped by selection and potentially affect herbivore and pollinator

communities.

METHODS

Experimental design --

Details of the field experiment are provided in Colautti & Barrett (2010) and the seeds used were

collected along a 10-degree latitudinal gradient in eastern North America, as described in

Montague *et al.* (2008). Briefly, 13 populations were chosen to represent a latitudinal gradient

from Timmins, Ontario (48.48 °N) to Easton, Maryland (38.75 °N) and were grown at the

Koffler Scientific Reserve at Jokers Hill, located approximately at the geographic center of the

latitudinal gradient (44.03 °N). Prior to inflorescence development, plants were sprayed with

insecticide to reduce incidence of herbivorous insects. The primary purpose of this experiment

was to characterize flowering time clines and genetic trade-offs in a multi-year (2006-2008)

common garden study, as reported in Colautti & Barrett (2010, 2011). In 2008 we additionally

recorded all open flowers on every stem of a subset of 369 individuals of 216 unique seed

families representing the 13 populations (Appendix S1). Our daily census records of this subset

of plants occurred over 125 days on a rotating schedule such that each individual was surveyed

every five days from the onset of flowering in July until September. A 5-day cycle was chosen

because stigma receptivity is nearly zero after 24 h of anthesis (Waites & Ågren, 2006), after

which unpollinated flowers begin to senesce and abscise while pollinated flowers begin to close

and fuse at the calyx, limiting the risk of counting the same flowers on multiple census dates. All

calculations and statistical analyses were conducted in R version 4.2.2 (R Core Team, 2022) in R

Studio v4.2.2 (RStudio Team, 2022), and are available with the original data from the Dryad

database (Dryad DOI: TBD(ROB)).

Characterizing flowering schedules --

We characterized the flowering schedules of each individual plant using the equations shown in

Table 1, standardized to the number of calendar days after the first flower was observed in the

experiment. The number of fruits produced over a growing season is variable in this experiment

because individuals that are locally adapted to mid-latitude balance flowering time with growth

to produce more flowers compared to small, early flowering phenotypes more characteristic of

northern populations or large, late-flowering phenotypes characteristic of more southern

populations. To account for variation in total flower production, we standardized the number of

open flowers every five days for each individual as a proportion of the total number of open

flowers censused over 125 days of observation. We then calculated a weighted census day by

multiplying each day by the proportion of flowers open on that day. The proportion of flowers

and the weighted census days were summed to calculate the weighted mean and other

characteristics of individual flowering schedules analogous to central moments (variance, skew

and kurtosis) (Box 1; Table 1). In addition to calculating the characteristics of every individual's

flowering schedule, we calculated the average of each characteristic for each population.

In addition, the start of flowering, duration, and central moments, we used a Principle

Coordinates Analysis to characterize flowering schedules of individuals as described by Austen

et al., (2014). Briefly, we computed a dissimilarity matrix from raw flower counts using

Kolmogorov-Smirnov (KS) distances (Sokal & Rohlf, 1995) where each matrix cell was a KS

statistic measuring the maximum distance between two flowering schedules. We chose KS

distance because it is robust to differences in flower number and census schedules. However, the

KS statistic is sensitive to differences in scale, so flowering durations were standardized to run

from 0 to 1 by subtracting the shortest duration time from each individual and dividing by the

difference of the longest and shortest durations (Austen et al., 2014). We used the KS matrix as

the input for the **cmdscale** function to generate the PCoA axes. Since KS distances do not meet

the triangle inequality condition of PCoA, we consider only eigenvectors with positive

eigenvalues (Austen et al., 2014; Legendre & Legendre, 1983). It is important to note that the

PCoA axes characterize variation in the shape of flowering schedules among individuals but not

variation in onset and duration of flowering.

In addition to individual flowering schedules, and the average of all individuals within

each population, we calculated characteristics of the aggregate population flowering schedules.

These calculations use the same equations after first pooling the flower counts for each day

across all individuals within a population (Table 1). For example, the aggregate start of flowering

of a population would be the first day that any individual in the population flowered, and

aggregate duration of flowering would be the number of days until the last open flower was

observed in the population.

We used bootstrap models with 999 iterations to compute a bootstrapped mean and 95%

confidence intervals for both population average and population aggregate flowering schedules.

In each iteration, individuals were hierarchically sampled with replacement from within each

population, and then each metric in Table 1 was calculated either from the pooled set of

individuals in each population to calculate the population aggregate characteristics, or for each

individual within the population to calculate the population average characteristics. Original data

with reproducible R code for the bootstrap models are available from the Dryad database (Dryad

<mark>DOI: TBD(ROB))</mark>

Characteristic correlations, clines and emergent properties --

Our first research question addresses whether conventional metrics (i.e., onset and duration of

flowering) correspond to more detailed characteristics of the flowering schedule (i.e., mean,

variance, skew, kurtosis, and PCoA axes). To investigate these relationships, we calculated

pairwise Pearson correlations to compare three scales of analysis, namely (A) average schedules

of each population (n = 13), (B) the deviation of each individual from its population mean (n =

369), and (C) aggregate schedules calculated on pools of individuals for each population (n =

13). In addition to comparing conventional metrics with other flowering schedule characteristics,

the correlational structure of these matrices can also be instructive for testing evolutionary

hypotheses because evolutionary divergence of flowering schedules would produce correlations

among average (A) or aggregate (C) populations that are not evident among individuals within

populations (B).

We used simple linear regression models of bootstrap means to address our second

question concerning whether there were latitudinal clines characteristic of both average and

aggregate flowering schedules. Each univariate linear regression included the bootstrap mean of

a flowering schedule characteristic as the dependent variable, with latitude as a linear and

quadratic predictor, and significance was tested using the **anova** function in R to find the

minimal adequate model.

To quantify emergent properties of population aggregate flowering schedules that differ

from the population average, we subtracted the average value of each population from the

aggregate value for the same population. Therefore, a positive value for an emergent property

means that a population had an aggregate value that was higher than the average individual

comprising it. For example, all populations will have a negative emergent property for onset of

flowering because the average onset of flowering will never be earlier than the day of first flower

in the aggregate population. As with average and aggregate characteristics, we tested for clines in

emergent properties using linear models as described in the previous paragraph.

Finally, we also considered the false discovery rate between 6 and 24 separate statistical

models. We consider a range of degrees of freedom because there are 24 separate statistical

models to test for clines (i.e., six characteristics across three types of comparisons) but the

aggregate, average, and emergent characteristics are each calculated from the same population

data, resulting in no more than 6 independent characteristics.

RESULTS

Flowering schedule characteristics --

Pearson product-moment correlations revealed significant relationships among commonly

measured traits of the flowering schedule (i.e., start and duration) and more detailed

characteristics (i.e., mean, variance, skew, and kurtosis), across three types of comparison (Fig.

2). At the aggregate level, populations that started flowering later also had a later mean flowering

date. Additionally, populations with longer flowering duration had a higher coefficient of

kurtosis indicating more flowering far from the mean flowering date (Fig. 2C). In contrast, there

was no significant correlation between start date and duration among population average

flowering schedules, but there was a significant negative correlation with skew indicating that

populations with a later start date, on average, tend to have individuals with imbalanced

flowering clustered more at later dates (Fig. 2A).

Flowering schedule variance was significantly positively correlated with flowering

duration on average but not among aggregate flowering schedules. Residual, within-population

correlations were generally weaker when compared to correlations among population average

and aggregate flowering schedules (Fig. 2B). In addition, the correlation between onset and

duration of flowing among individuals within populations was in the opposite direction to

population average and aggregate. In other words, plants that started flowering earlier relative to

other individuals within the same population also flowered for a shorter duration (Fig. 2B),

whereas populations with earlier flowering individuals, on average, also tended to have

individuals with longer flowering durations (Fig. 2A).

Principal Coordinates Analysis --

In addition to the above characteristics, we use Principal Coordinates Analysis (PCoA) to fully

characterize similarities and differences among flowering schedules after controlling for

variation in start date and duration. The PCoA of all 369 individuals resulted in 369 independent

eigenvectors, of which 192 had positive eigenvalues. However, of these 192, only the first two

PCoA axes had eigenvectors that explained more than 10% of the variation in flowing schedule

shape, and together they explained 35% of the variation (Table 2). By comparison, the next ten

eigenvectors combined explained another 35% of variation, with the remaining 180 accounting

for 30% of the variation. Thus, we next focus on the first two eigenvectors.

For both the population aggregate and average, the first axis (PCoA1) was positively

correlated with skew and negatively correlated with the mean day of flowering (Fig. 2). PCoA1

was also negatively correlated with duration of flowering when comparing population average

but not population aggregate flowering schedules. In contrast, within-population residual

correlations involving PCoA1 were generally weaker than among population average or

aggregate, with significant correlations for mean, skew and kurtosis (Fig. 2B). Correlations

involving PCoA2 were quite different than PCoA1 within and among all three comparisons. For

example, PCoA2 was not significantly correlated with any other metrics in aggregate

distributions (Fig. 2C) but was negatively correlated with start and kurtosis and positively

correlated with skew and duration among individuals within the same populations (Fig. 2B) and

was positively correlated with variance and duration of average distributions (Fig. 2A). By

mathematical definition of the PCoA, there was no correlation between PCoA1 and PCoA2

among all individuals, nor was there a significant correlation among residuals of individuals

within populations (Fig. 2B). In other words, PCoA1 and PCoA2 change in a correlated way.

This can be seen clearly in Fig. 3, where northern populations cluster at intermediate values of

PCoA1 and PCoA2 compared to mid-latitude (higher PCoA1 but lower PCoA2) and southern

populations (lower PCoA1 and lower PCoA2).

323   ***Latitudinal clines and emergent properties --***

The correlated changes in population aggregate and average flowering schedules are reflected in

significant changes with latitude. For example, the start, mean, and kurtosis of flowering

schedules were each significantly negatively related to latitude for both aggregate and average

schedules (Table 3, Fig. 4). This means that southern populations start flowering later, with a

later average flowering date, and more 'outlier' flowers opening far from the average flowering

date, both in aggregate and on average, when compared to northern populations. However,

aggregate and average schedules vary in two key characteristics – aggregate but not average

schedule duration declined with latitude, whereas average but not aggregate skew was negative

at lower latitudes (Fig. 4). In other words, individuals have similar flowering durations on

average but are more spread out in southern populations, resulting in a longer aggregate duration

towards the south. Similarly, individual schedules are more heavily weighted toward later

flowering dates towards the south, but variation in start dates average out such that aggregate

schedules are more evenly distributed and not significantly related to latitude.

To quantify emergent properties, we compared aggregate to average schedules and this

procedure revealed considerable variation among populations and two latitudinal clines (Table 4;

Fig. 4). Although the aggregate start of flowering must always be earlier than the average,

resulting in a negative emergent property score, we found a significant positive correlation with

latitude. Likewise, the duration of flowering must always be longer in aggregate than on average,

but we found a marginal negative relationship with latitude (Fig. 4, right column). Together,

these clines demonstrate more synchronous flowering schedules in northern populations

compared to those occurring in the south. We also found a nonlinear cline in variance

representing a lower aggregated flowering schedule variance than the average individual

variance at intermediate latitudes.

Overall, we expect about one false positive cline at $\alpha < 0.05$ with 24 independent

statistical tests. But of the 10 significant clines, only five remained significant after a Bonferroni

adjusted $\alpha = 0.0083$ for six independent tests or $\alpha = 0.0021$ for 24 tests. Despite uncertainty in

the detection of clines, bootstrap estimates show a relatively high level of precision for most

population means, as shown by the small error bars in Fig. 4.

DISCUSSION

There is growing evidence for rapid evolution in invasive species (van Kleunen et al., 2018), but

it remains uncertain how often adaptive evolution contributes to their spread (Colautti & Lau,

2015). Rapid evolution of a latitudinal cline in flowering time in response to local growing

conditions likely increased the rate and extent of invasion by *Lythrum salicaria* across North

America (Colautti & Barrett, 2013; Wu & Colautti, 2022). However, previous studies focusing

on the onset and duration of flowering provided only a partial view of how plants allocate

flowering resources (Box 1), raising uncertainty about the extent to which phenological

evolution has been adaptive. We used a common garden study with detailed metrics to

characterize entire flowering schedules and tested hypotheses about adaptive latitudinal clines.

We found that the onset and duration of flowering were not good predictors of other

characteristics of individual flowering schedules (Fig. 2B). In contrast, the average flowering

schedule characteristics were highly correlated among population averages (Fig. 2A) and also

among population aggregate flowering schedules (Fig. 2C). Furthermore, that these among-

population correlations were largely due to correlated evolution along latitudinal clines (Fig. 4).

Emergent properties were highly variable among populations and formed clines indicative of

more synchronized flowering schedules in northern populations. Below we discuss the

implications of these results for understanding the evolutionary causes and ecological

consequences of flowering schedule diversity.

Evidence for correlated evolution --

Date of first flower and duration of flowering are relatively easy to measure and are commonly

reported as focal traits in phenological studies. However, reproductive fitness depends on the full

schedule of flower production, and it has not been clear how well the onset or duration of

flowering correlates with other characteristics of the flowering schedule. In *L. salicaria*, plants

that flowered later also flowered for a longer duration with a higher flowering variance and

slightly lower skew, relative to other individuals sampled from the same population (Fig. 2B).

This result is consistent with a life history trade-off between flowering earlier at a small size or

delaying reproduction to accumulate more resources for later floral investment – a mechanism

that was proposed previously for this species (Colautti & Barrett, 2013; Wu & Colautti, 2022)

but was not validated with floral display data until now.

In contrast to individual flowering schedules, measurements of the flowering schedules of

whole populations provide more limited insights into the evolutionary mechanisms shaping

flowering schedules. The different correlational structure for average (Fig. 2B) versus aggregate

schedules (Fig. 2C) produced significant variation in emergent property characteristics among

populations (Fig. 4C), suggesting that aggregate characteristics are not generally indicative of

individual flowering schedules within a population. Studies that have not considered the full

schedule shape of individual plants may therefore neglect key aspects of phenology that are

crucial for investigating ecological and evolutionary hypotheses. For example, there is

considerable among-population and among-species variability in phenological shifts due to

climate change (Anderson et al., 2012; Fitchett et al., 2015; D. W. Inouye, 2022; Panchen, 2022;

Panchen & Gorelick, 2017; Prather et al., 2023). This variation may be explained by selection on

characteristics of flowering schedules that are not correlated with the onset of flowering.

There were differences in both the strength and direction of correlations among

individuals within *L. salicaria* populations when compared with population average and

aggregate characteristics (Fig. 2). Among-population correlations were caused by either

statistical artifacts or genetic constraints, then we would expect to see similar correlations among

individuals within populations. Instead, the differing correlation matrices are more consistent

with correlated evolution among populations. We cannot rule out the possibility that

environmental effects mask genetic correlations within populations; however, this would require

a much larger experiment with sufficient replication to estimate correlations among relatives

within populations (Price et al., 1988; Rausher, 1992). Nor can we rule out the potential role of

stochastic processes without replicated sampling from independent geographical gradients

(Colautti & Lau, 2015). Nevertheless, our finding that flowering schedules change continuously

with latitude lends additional evidence for the correlated evolution hypothesis. Moreover, the

direction of the observed clines are consistent with adaptive responses to biotic and abiotic

selection, as discussed below.

Latitudinal clines and emergent properties --

Latitudinal clines in the onset of flowering have evolved independently in several species across

North America (McGoey et al., 2020; Montague et al., 2008; Samis et al., 2012; Wu & Colautti,

2022). In our study of *L. salicaria*, we additionally found latitudinal clines in average mean,

skew and kurtosis of flowering schedules (Fig. 4, left column). These types of clines indicate that

individuals from southern environments have floral displays that peak later in the season but also

a few open flowers far from the mean flowering date. These findings correspond to higher

intensity of biotic interactions in southern environments reported in a meta-analysis of terrestrial

ecosystems (Zvereva & Kozlov, 2021).

Lythrum salicaria is a self-incompatible outcrosser (Colautti et al., 2010; Darwin, 1877;

O'Neil, 1997) and therefore individuals in southern populations likely experience strong inter-

specific competition for pollinators. Additionally, several effective biological control insects

have been released into eastern North America, including *Gallerucella* beetles that feed on

meristem tissue as larvae and can drastically reduce flower production (Blossey & Notzold,

1995; Grevstad, 2006). The later average flowering and more negative skew in plants from

southern populations is therefore consistent with selection from higher herbivory and more

competition for pollinators under a longer growing season. We cannot easily parse the specific

effects of biotic and abiotic factors based on clines alone, but we can make the prediction that

clines in herbivore and pollinator activity would be more patchy and less predictable by latitude

compared to season length. It is therefore interesting that we detected strong clines in average

start ($\beta = -0.856$) and mean ($\beta = -0.881$) flowering dates, which we predict to be more sensitive

to season length, in contrast to kurtosis ($\beta = -0.590$), which we predicted as a bet-hedging

strategy for biotic interactions. Skew shows an intermediate pattern ($\beta = 0.745$) and could be an

adaptation to biotic selection but could also be partly an artifact of our mid-latitude common

garden site, as discussed below.

Characteristics of aggregate flowering schedules were generally very different from their

population averages, resulting in high variability in emergent properties among populations (Fig.

4, right column), but these differences did not generally correlate with latitude. Two important

exceptions are the significant and marginally significant clines observed in the onset and

duration of flowering respectively, which characterized more synchronized flowering schedules

in the north. However, the mid-latitude location of our common garden may also be responsible

for these trends. Cooler temperatures that are expected with increasing latitude not only delay the

start of favorable growing conditions, but they also restrict insect pollinator activity. Therefore,

lack of pollinator activity towards the end of the growing season at our mid-latitude field site

may represent an artificial truncation of flowering by southern populations leading to a perceived

difference in flowering schedule synchrony. Overall, variation in emergent properties among *L.*

salicaria populations has important implications for field research on components of floral

display. It has previously been established that individuals with flowering schedules

uncharacteristic of their populations can potentially bias ecological and evolutionary inferences

drawn from aggregate population schedules (Newstrom et al. 1994; B. D. Inouye et al., 2019).

The emergent properties in our study suggest that such cases are likely more common than is

generally appreciated and that they may also be geographically structured. As such, aggregate

schedules should be interpreted cautiously.

Conclusions --

Our study demonstrates that populations of *L. salicaria* in eastern North America have evolved

differences in flowering schedules within less than a century of invasion. This phenological

differentiation among populations includes predictable changes in the timing of resource

allocation to flower development and floral display and adds to the growing list of phenotypic

traits that are targets of selection as invasive species spread along environmental gradients

(MacDougall et al., 2006; McGoey et al., 2020; Molina-Montenegro et al., 2018; Urbanski et al.,

2012). Based on the observed clines in our study, we predict that longer growing seasons

accompanying anthropogenic climate change will alter flowering schedules. These changes are

likely to involve a later start date to flowering, a later mean day of flowering, and a larger

kurtosis, which in turn should increase seed production and promote population growth and

range expansion of *L. salicaria*. The ecological and evolutionary insights drawn from

observations of flowering start dates and population aggregate characteristics should, however,

be interpreted cautiously. In future, more detailed features of individual flowering schedules can

be characterised by multivariate methods and metrics drawn from central moments theory. As we

have shown for *L. salicaria* from eastern North America, these more refined characteristics of

individual flowering schedules might help in developing a deeper understanding of the diverse

ways that species respond to changing environments.

Acknowledgements

The authors thank Z. Burivalova and R. Mackenzie for field assistance, and to E.J. Austen, A.E.

Weis and J. Friedman for feedback on the analysis. Funding was provided from NSERC

Discovery grants to S.C.H.B. and R.I.C. and to M.N.A. from NSERC CGSM.

AUTHOR CONTRIBUTIONS

M.N.A. led the data analysis with assistance from D.M., S.C.H.B., and R.I.C. The

conceptualization and experimental design were developed by S.C.H.B. and R.I.C., and all

authors contributed to the manuscript, based on an original draft written by M.N.A.

DATA AVAILABILITY STATEMENT

Raw data and fully reproducible R code for all figures, tables, and statistical analyses are

available in the Dryad database (DOI:)

LITERATURE CITED

Ågren, J., & Schemske, D. W. (2012). Reciprocal transplants demonstrate strong adaptive

differentiation of the model organism Arabidopsis thaliana in its native range. *New*

Phytologist, *194*(4), 1112–1122. https://doi.org/10.1111/j.1469-8137.2012.04112.x

Alexander, J. M., Edwards, P. J., Poll, M., Parks, C. G., & Dietz, H. (2009). Establishment of

parallel altitudinal clines in traits of native and introduced forbs. *Ecology*, *90*(3),

612–622. https://doi.org/10.1890/08-0453.1

Anderson, J. T., & Gezon, Z. J. (2015). Plasticity in functional traits in the context of climate

change: A case study of the subalpine forb Boechera stricta (Brassicaceae). *Global*

Change Biology, *21*(4), 1689–1703. https://doi.org/10.1111/gcb.12770

Anderson, J. T., Inouye, D. W., McKinney, A. M., Colautti, R. I., & Mitchell-Olds, T. (2012).

Phenotypic plasticity and adaptive evolution contribute to advancing flowering

phenology in response to climate change. *Proceedings of the Royal Society B:*

Biological Sciences, *279*(1743), 3843–3852.

https://doi.org/10.1098/rspb.2012.1051

Augspurger, C. K. (1981). Reproductive synchrony of a tropical shrub: Experimental studies

on effects of pollinators and seed predators in Hybanthus prunifolius (violaceae).

Ecology, *62*(3), 775–788. https://doi.org/10.2307/1937745

Austen, E. J., Jackson, D. A., & Weis, A. E. (2014). Describing flowering schedule shape

through multivariate ordination. *International Journal of Plant Sciences*.

https://doi.org/10.1086/673934

Austen, E. J., Rowe, L., Stinchcombe, J. R., & Forrest, J. R. K. (2017). Explaining the

apparent paradox of persistent selection for early flowering. *New Phytologist*,

215(3), 929–934. https://doi.org/10.1111/nph.14580

Austen, E. J., & Weis, A. E. (2015). What drives selection on flowering time? An

experimental manipulation of the inherent correlation between genotype and

environment: causes of selection on flowering time. *Evolution*, *69*(8), 2018–2033.

https://doi.org/10.1111/evo.12709

Blionis, G. J., Halley, J. M., & Vokou, D. (2001). Flowering phenology of Campanula on Mt

Olympos, Greece. *Ecography*, *24*(6), 696–706. https://doi.org/10.1111/j.1600-

0587.2001.tb00531.x

Blossey, B., & Notzold, R. (1995). Evolution of Increased Competitive Ability in Invasive

Nonindigenous Plants: A Hypothesis. *Journal of Ecology*, *83*(5), 887–889.

https://doi.org/10.2307/2261425

CaraDonna, P. J., Iler, A. M., & Inouye, D. W. (2014). Shifts in flowering phenology reshape a

subalpine plant community. *Proceedings of the National Academy of Sciences*,

111(13), 4916–4921. https://doi.org/10.1073/pnas.1323073111

Colautti, R. I., & Barrett, S. C. H. (2010). Natural selection and genetic constraints on

flowering phenology in an invasive plant. *International Journal of Plant Sciences*,

171(9), 960–971. https://doi.org/10.1086/656444

Colautti, R. I., & Barrett, S. C. H. (2011). Population divergence along lines of genetic

variance and covariance in the invasive plant Lythrum salicaria in eastern North

America. *Evolution*, *65*(9), 2514–2529. https://doi.org/10.1111/j.1558-

5646.2011.01313.x

Colautti, R. I., & Barrett, S. C. H. (2013). Rapid adaptation to climate facilitates range

expansion of an invasive plant. *Science*, *342*(6156), 364–366.

https://doi.org/10.1126/science.1242121

Colautti, R. I., & Lau, J. A. (2015). Contemporary evolution during invasion: Evidence for

differentiation, natural selection, and local adaptation. *Molecular Ecology*, *24*(9),

1999–2017. https://doi.org/10.1111/mec.13162

Colautti, R. I., White, N. A., & Barrett, S. C. H. (2010). Variation of Self-Incompatibility

within Invasive Populations of Purple Loosestrife (*Lythrum salicaria* L.) from Eastern

North America. *International Journal of Plant Sciences*, *171*(2), 158–166.

https://doi.org/10.1086/649023

Darwin, C. (1877). *The Different Forms of Flowers on Plants of the Same Species*. John

Murray, Albermarle Street.

de Villemereuil, P., Charmantier, A., Arlt, D., Bize, P., Brekke, P., Brouwer, L., Cockburn, A.,

Côté, S. D., Dobson, F. S., Evans, S. R., Festa-Bianchet, M., Gamelon, M., Hamel, S.,

Hegelbach, J., Jerstad, K., Kempenaers, B., Kruuk, L. E. B., Kumpula, J., Kvalnes, T.,

… Chevin, L.-M. (2020). Fluctuating optimum and temporally variable selection on

breeding date in birds and mammals. *Proceedings of the National Academy of*

Sciences, *117*(50), 31969–31978. https://doi.org/10.1073/pnas.2009003117

Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E., & Bernasconi, G. (2007). Time after

time: Flowering phenology and biotic interactions. *Trends in Ecology & Evolution*,

22(8), 432–439. https://doi.org/10.1016/j.tree.2007.05.006

Ensing, D. J., & Eckert, C. G. (2019). Interannual variation in season length is linked to

strong co-gradient plasticity of phenology in a montane annual plant. *New*

Phytologist, *224*(3), 1184–1200. https://doi.org/10.1111/nph.16009

Fantinato, E., Del Vecchio, S., Giovanetti, M., Acosta, A. T. R., & Buffa, G. (2018). New

insights into plants co-existence in species-rich communities: The pollination

interaction perspective. *Journal of Vegetation Science*, *29*(1), 6–14.

https://doi.org/10.1111/jvs.12592

Fenster, C. B., Armbruster, W. S., Wilson, P., Dudash, M. R., & Thomson, J. D. (2004).

Pollination syndromes and floral specialization. *Annual Review of Ecology,*

Evolution, and Systematics, *35*, 375–403.

Fitchett, J. M., Grab, S. W., & Thompson, D. I. (2015). Plant phenology and climate change:

Progress in methodological approaches and application. *Progress in Physical*

Geography: Earth and Environment, *39*(4), 460–482.

https://doi.org/10.1177/0309133315578940

Forrest, J., & Thomson, J. D. (2010). Consequences of variation in flowering time within and among individuals of Mertensia fusiformis (Boraginaceae), an early spring wildflower. *American Journal of Botany*, *97*(1), 38–48.

https://doi.org/10.3732/ajb.0900083

Fox, G. A. (2003). Assortative mating and plant phenology: Evolutionary and practical consequences. *Evolutionary Ecology Research*, *5*(1), 18.

Gauzere, J., Teuf, B., Davi, H., Chevin, L.-M., Caignard, T., Leys, B., Delzon, S., Ronce, O., & Chuine, I. (2020). Where is the optimum? Predicting the variation of selection along climatic gradients and the adaptive value of plasticity. A case study on tree phenology. *Evolution Letters*, *4*(2), 109–123. https://doi.org/10.1002/evl3.160

Grevstad, F. S. (2006). Ten-year impacts of the biological control agents Galerucella pusilla and G. calmariensis (Coleoptera: Chrysomelidae) on purple loosestrife (Lythrum salicaria) in Central New York State. *Biological Control*, *39*(1), 1–8.

https://doi.org/10.1016/j.biocontrol.2006.03.007

Griffith, T. M., & Watson, M. A. (2005). Stress avoidance in a common annual: Reproductive timing is important for local adaptation and geographic distribution. *Journal of Evolutionary Biology*, *18*(6), 1601–1612. https://doi.org/10.1111/j.1420-9101.2005.01021.x

Halbritter, A. H., Fior, S., Keller, I., Billeter, R., Edwards, P. J., Holderegger, R., Karrenberg, S., Pluess, A. R., Widmer, A., & Alexander, J. M. (2018). Trait differentiation and

adaptation of plants along elevation gradients. *Journal of Evolutionary Biology*,

31(6), 784–800. https://doi.org/10.1111/jeb.13262

Inouye, B. D., Ehrlén, J., & Underwood, N. (2019). Phenology as a process rather than an

event: From individual reaction norms to community metrics. *Ecological*

Monographs, *89*(2), e01352. https://doi.org/10.1002/ecm.1352

Inouye, D. W. (2022). Climate change and phenology. *WIREs Climate Change*, *13*(3), e764.

https://doi.org/10.1002/wcc.764

Ison, J. L., & Weis, A. E. (2017). Temporal population genetic structure in the pollen pool for

flowering time: A field experiment with Brassica rapa (Brassicaceae). *American*

Journal of Botany, *104*(10), 1569–1580. https://doi.org/10.3732/ajb.1700210

Kirkpatrick, M., & Barton, N. H. (1997). Evolution of a species' range. *The American*

Naturalist, *150*(1), 1–23. https://doi.org/10.1086/286054

Kudo, G. (2007). Flowering phenologies of animal-pollinated plants: Reproductive

strategies and agents of selection. In *Ecology and Evolution of Flowers*. Oxford

University Press, Incorporated.

Legendre, L., & Legendre, P. (1983). *Numerical Ecology*. Elsevier Scientific Publishing

Company.

MacDougall, A. s., Boucher, J., Turkington, R., & Bradfield, G. e. (2006). Patterns of plant

invasion along an environmental stress gradient. *Journal of Vegetation Science*,

17(1), 47–56. https://doi.org/10.1111/j.1654-1103.2006.tb02422.x

McGoey, B. V., Hodgins, K. A., & Stinchcombe, J. R. (2020). Parallel flowering time clines in

native and introduced ragweed populations are likely due to adaptation. *Ecology*

and Evolution, *10*(11), 4595–4608. https://doi.org/10.1002/ece3.6163

Molina-Montenegro, M. A., Acuña-Rodríguez, I. S., Flores, T. S. M., Hereme, R., Lafon, A.,

Atala, C., & Torres-Díaz, C. (2018). Is the Success of Plant Invasions the Result of

Rapid Adaptive Evolution in Seed Traits? Evidence from a Latitudinal Rainfall

Gradient. *Frontiers in Plant Science*, *9*, 208.

https://doi.org/10.3389/fpls.2018.00208

Montague, J. L., Barrett, S. C. H., & Eckert, C. G. (2008). Re-establishment of clinal

variation in flowering time among introduced populations of purple loosestrife

(Lythrum salicaria, Lythraceae). *Journal of Evolutionary Biology*, *21*(1), 234–245.

https://doi.org/10.1111/j.1420-9101.2007.01456.x

Munguía-Rosas, M. A., Ollerton, J., Parra-Tabla, V., & De-Nova, J. A. (2011). Meta-analysis

of phenotypic selection on flowering phenology suggests that early flowering plants

are favoured. *Ecology Letters*, *14*(5), 511–521. https://doi.org/10.1111/j.1461-

0248.2011.01601.x

Newstrom, L. E., Frankie, G. W., & Baker, H. G. (1994). A new classification for plant

phenology based on flowering patterns in lowland tropical rain forest trees at Le

Selva, Costa Rica. Biotropica, 26(2) 151-169.

Olsson, K., & Ågren, J. (2002). Latitudinal population differentiation in phenology, life

history and flower morphology in the perennial herb Lythrum salicaria. *Journal of*

Evolutionary Biology, *15*(6), 983–996. https://doi.org/10.1046/j.1420-

9101.2002.00457.x

O'Neil, P. (1997). Natural selection on genetically correlated phenological characters in

Lythrum salicaria l. (Lythraceae). *Evolution*, *51*(1), 267–274.

https://doi.org/10.2307/2410980

Panchen, Z. A. (2022). Plant reproductive phenology along an elevation gradient in the

extreme environment of the Canadian High Arctic. *Plant Ecology & Diversity*, *15*(5–

6), 213–226. https://doi.org/10.1080/17550874.2022.2147804

Panchen, Z. A., & Gorelick, R. (2017). Prediction of Arctic plant phenological sensitivity to

climate change from historical records. *Ecology and Evolution*, *7*(5), 1325–1338.

https://doi.org/10.1002/ece3.2702

Prather, R. M., Dalton, R. M., barr, billy, Blumstein, D. T., Boggs, C. L., Brody, A. K., Inouye,

D. W., Irwin, R. E., Martin, J. G. A., Smith, R. J., Van Vuren, D. H., Wells, C. P.,

Whiteman, H. H., Inouye, B. D., & Underwood, N. (2023). Current and lagged climate

affects phenology across diverse taxonomic groups. *Proceedings of the Royal*

Society B: Biological Sciences, *290*(1990), 20222181.

https://doi.org/10.1098/rspb.2022.2181

Price, T., Kirkpatrick, M., & Arnold, S. J. (1988). Directional selection and the evolution of

breeding date in birds. *Science*, *240*(4853), 798–799.

https://doi.org/10.1126/science.3363360

R Core Team. (2022). *R: A language and environment for statistical computing.* [Computer

software]. R Foundation for Statistical Computing. https://www.R-project.org/

Rabinowitz, D., Rapp, J. K., Sork, V. L., Rathcke, B. J., Reese, G. A., & Weaver, J. C. (1981).

Phenological properties of wind- and insect-pollinated prairie plants. *Ecology*,

62(1), 49–56. https://doi.org/10.2307/1936667

Rafferty, N. E., Diez, J. M., & Bertelsen, C. D. (2020). Changing climate drives divergent and

nonlinear shifts in flowering phenology across elevations. *Current Biology*, *30*(3),

432-441.e3. https://doi.org/10.1016/j.cub.2019.11.071

Rathcke, B., & Lacey, E. P. (1985). Phenological patterns of terrestrial plants. *Annual*

Review of Ecology and Systematics, *16*(1), 179–214.

https://doi.org/10.1146/annurev.es.16.110185.001143

Rausher, M. D. (1992). The measurement of selection on quantitative traits: Biases due to

environmental covariances between traits and fitness. *Evolution*, *46*(3), 616–626.

https://doi.org/10.1111/j.1558-5646.1992.tb02070.x

RStudio Team. (2022). *RStudio: Integrated Development for R* [Computer software].

RStudio, PBC. http://www.rstudio.com/.

Samis, K. E., Murren, C. J., Bossdorf, O., Donohue, K., Fenster, C. B., Malmberg, R. L.,

Purugganan, M. D., & Stinchcombe, J. R. (2012). Longitudinal trends in climate drive

flowering time clines in North American Arabidopsis thaliana. *Ecology and*

Evolution, *2*(6), 1162–1180. https://doi.org/10.1002/ece3.262

Simons, A. M. (2011). Modes of response to environmental change and the elusive

empirical evidence for bet hedging. *Proceedings of the Royal Society B: Biological*

Sciences, *278*(1712), 1601–1609. https://doi.org/10.1098/rspb.2011.0176

Sobral, M., Guitián, J., Guitián, P., & Larrinaga, A. R. (2013). Selective pressure along a

latitudinal gradient affects subindividual variation in plants. *PLOS ONE*, *8*(9),

e74356. https://doi.org/10.1371/journal.pone.0074356

Sokal, R. R., & Rohlf, F. J. (1995). *Biometry* (3rd ed.). WH Freeman.

Stemkovski, M., Dickson, R. G., Griffin, S. R., Inouye, B. D., Inouye, D. W., Pardee, G. L.,

Underwood, N., & Irwin, R. E. (2023). Skewness in bee and flower phenological

distributions. *Ecology*, *104*(1), e3890. https://doi.org/10.1002/ecy.3890

Strauss, S. Y., & Whittall, J. B. (2006). Non-pollinator agents of selection on floral traits. In

Ecology and Evolution of Flowers (pp. 120–135). Oxford Biology.

Thomson, J. D. (1980). Skewed flowering distributions and pollinator attraction. *Ecology*,

61(3), 572–579. https://doi.org/10.2307/1937423

Tufto, J. (2015). Genetic evolution, plasticity, and bet-hedging as adaptive responses to

temporally autocorrelated fluctuating selection: A quantitative genetic model.

Evolution, 69(8), 2034–2049. https://doi.org/10.1111/evo.12716

Urbanski, J., Mogi, M., O'Donnell, D., DeCotiis, M., Toma, T., & Armbruster, P. (2012). Rapid

Adaptive Evolution of Photoperiodic Response during Invasion and Range

Expansion across a Climatic Gradient. *The American Naturalist*, *179*(4), 490–500.

https://doi.org/10.1086/664709

van Kleunen, M., Bossdorf, O., & Dawson, W. (2018). The ecology and evolution of alien

plants. *Annual Review of Ecology, Evolution, and Systematics*, *49*(1), 25–47.

https://doi.org/10.1146/annurev-ecolsys-110617-062654

Waites, A. R., & Ågren, J. (2006). Stigma receptivity and effects of prior self-pollination on

seed set in tristylous Lythrum salicaria (Lythraceae). *American Journal of Botany*,

93(1), 142–147. https://doi.org/10.3732/ajb.93.1.142

Wolkovich, E. M., Cook, B. I., Allen, J. M., Crimmins, T. M., Betancourt, J. L., Travers, S. E.,

Pau, S., Regetz, J., Davies, T. J., Kraft, N. J. B., Ault, T. R., Bolmgren, K., Mazer, S. J.,

McCabe, G. J., McGill, B. J., Parmesan, C., Salamin, N., Schwartz, M. D., & Cleland,

E. E. (2012). Warming experiments underpredict plant phenological responses to

climate change. *Nature*, *485*(7399), Article 7399.

https://doi.org/10.1038/nature11014

Wright, J. W., & Meagher, T. R. (2003). Pollination and seed predation drive flowering

phenology in Silene latifolia (Caryophyllaceae). *Ecology*, *84*(8), 2062–2073.

https://doi.org/10.1890/02-0676

Wu, Y., & Colautti, R. I. (2022). Evidence for continent-wide convergent evolution and

stasis throughout 150 y of a biological invasion. *Proceedings of the National*

Academy of Sciences, *119*(18), e2107584119.

https://doi.org/10.1073/pnas.2107584119

Zvereva, E. L., & Kozlov, M. V. (2021). Latitudinal gradient in the intensity of biotic

interactions in terrestrial ecosystems: Sources of variation and differences from the

diversity gradient revealed by meta-analysis. *Ecology Letters*, *24*(11), 2506–2520.

https://doi.org/10.1111/ele.13851

Box 1: Summary characteristics of flowering schedules

Central moments (i.e., mean, variance, skew, and kurtosis) arise from probability theory and characterize distributions that can be visualized as histograms with observed values on the x-axis and probability density, frequency, or number of observations on the y-axis. Here, we use the same equations to describe analogous characteristics of flowering schedules that arise from developmental rather than probabilistic process. In contrast to probability histograms, flowering schedules can be visualized as a time series with number or proportion of flowers on the y-axis. Although the underlying processes are distinct, the equations are the same (Table 1). To understand the biological significance of mean, variance, skew and kurtosis, we contrast flowering schedules of different shape (Fig. 1). Standardizing to proportion of open flowers (P_t) over time (t), rather than total flower number (N_t) accounts for variation in total flower number. Moreover, we can use proportions to calculate a weighted mean day of flowering. By analogy to the mean of a probability distribution, the weighted mean of a flowering schedule represents the 'balance point', which better captures the anthesis day of a typical flower.

   As demonstrated in Figure 1A, central moments capture biologically meaningful variation in flowering schedules that may not be correlated with onset or duration of flowering. For example, the variance parameter (σ_i^2) describes how much flowering is spread out, even if the start and end dates are the same. An individual's coefficient of skew (CS_i) accounts for expected differences in variance and thus describes whether flowering is concentrated earlier (i.e., right skew, $CS_i > 0$ as shown by the blue dashed line) or later in the schedule (i.e., left skew, $CS_i < 0$ as shown by the purple dotted line). Finally, the coefficient of kurtosis (CK_i) describes deviations from the Gaussian expectation, contrasting individuals with a concentrated cluster of open flowers and a few flowers over a long-time frame (i.e., leptokurtic distribution, $CK_i > 0$ as shown in the gold dashed line) or individuals that spread out flowering more evenly over the entire schedule (i.e., platykurtic, $CK_i < 0$ as shown by the dashed green line). In addition to calculating these characteristics for individual plants, all flowers (or proportion of flowers) on each day can be pooled to yield a population aggregate flowering schedule (Fig. 1B). The aggregate schedule of a population may not be representative of the individual schedules comprising it, resulting in emergent properties that we define as deviations from the average individual flowering schedule.

Table 1. Equations of flowering schedule characteristics for an average individual within each population and for the aggregate population of pooled individuals.

Summary Statistic	Individual (i)	Population aggregate (j)
Proportion (P) of flowers (F) produced at time t	$P_{i,t} = \dfrac{F_{i,t}}{\sum_t F_{i,t}}$	$P_{t,j}^{\Sigma} = \dfrac{F_{t,j}^{\Sigma}}{\sum_t F_{t,j}^{\Sigma}} = \dfrac{\sum_i F_{i,t,j}}{\sum_t \sum_i F_{i,t,j}}$
Day of first flower	$\alpha_i = \min\{t \mid F_{i,t} > 0\}$	$\alpha_j^{\Sigma} = \min\{t \mid \sum_i F_{i,t,j} > 0\}$
Day of last flower	$\Omega_i = \max\{t \mid F_{i,t} > 0\}$	$\Omega_j^{\Sigma} = \max\{t \mid \sum_i F_{i,t,j} > 0\}$
Duration of flowering	$\Omega_i - \alpha_i$	$\Omega_j^{\Sigma} - \alpha_j^{\Sigma}$
Day of mean flowering	$\mu_i = \sum_t t\, P_{i,t}$	$\mu_j^{\Sigma} = \sum_t t\, P_{t,j}^{\Sigma}$
Flowering schedule variance	$\sigma_i^2 = \sum_t (t - \mu)^2\, P_t$	$\sigma_j^{2\Sigma} = \sum_t \left(t - \mu_j^{\Sigma}\right)^2 P_{t,j}^{\Sigma}$
Flowering schedule coefficient of skew	$CS_i = \dfrac{\sum_t (t - \mu_i)^3\, P_{i,t}}{\sigma_i^3}$	$CS_j = \dfrac{\sum_t \left(t - \mu_j^{\Sigma}\right)^3 P_{t,j}^{\Sigma}}{\sigma_j^{3\Sigma}}$
Flowering schedule coefficient of kurtosis	$CK_i = \dfrac{\sum_t (t - \mu_i)^4\, P_{i,t}}{\sigma_i^4} - 3$	$CK_j = \dfrac{\sum_t \left(t - \mu_j^{\Sigma}\right)^4 P_{t,j}^{\Sigma}}{\sigma_j^{4\Sigma}}$

Table 2. Summary of the first five eigenvector axes of a Principal Coordinates Analysis (PCoA) using Kolmogorov-Smirnov distances between flowering schedules of 369 plants from 13 populations of *Lythrum salicaria* sampled along a latitudinal gradient and grown in a common garden at the Koffler Scientific Reserve in Newmarket, ON. Percent variation is calculated as the eigenvalue over the sum of all positive eigenvalues.

Axis	Eigenvalue	Variation explained (%)
PCoA1	13.4	23.5
PCoA2	6.6	11.5
PCoA3	3.9	6.9
PCoA4	3.2	5.6
PCoA5	2.6	4.7

Table 3. Standardized slope and test statistics of univariate linear regressions testing latitude as a predictor of the bootstrapped means of average and aggregate population flowering schedule traits, and the emergent property (i.e. population aggregate minus average) in populations of *Lythrum salicaria*. Bold text indicates significant values.

Traits		β	$F_{(1,11)}$	P	X^2	P
			Linear term		**Quadratic term**	
Start	Average	-0.856	30.0	**< 0.001**	1.73	0.189
	Aggregate	-0.591	5.92	**< 0.05**	0.01	0.907
	Emergent	0.857	30.52	**< 0.001**	2.50	0.114
Duration	Average	-0.369	1.73	0.215	0.11	0.738
	Aggregate	-0.849	28.37	**< 0.001**	0.01	0.904
	Emergent	-0.470	3.13	0.105	2.96	0.085
Mean	Average	-0.881	38.0	**< 0.001**	0.47	0.494
	Aggregate	-0.845	27.5	**< 0.001**	0.12	0.724
	Emergent	0.368	1.66	0.224	0	0.948
Variance	Average	0.031	0.01	0.920	0.07	0.791
	Aggregate	-0.214	0.53	0.483	0	0.977
	Emergent	0.156	0.276	0.610	6.83	**< 0.01**
Skew	Average	0.745	13.7	**< 0.01**	1.03	0.310
	Aggregate	0.382	1.87	0.198	0.07	0.797
	Emergent	0.227	0.60	0.456	0.12	0.741
Kurtosis	Average	-0.590	5.88	**< 0.05**	0.20	0.659
	Aggregate	-0.592	5.94	**< 0.05**	0.88	0.347
	Emergent	-0.133	0.19	0.664	1.18	0.279

Figure 1. Hypothetical flowering schedules of (A) five individuals with different characteristics and (B) their pooled aggregate flowering schedule. Flowering schedules are scaled to the proportion of total flower production (P_t) over time (t).

Figure 2. Matrices showing pairwise Pearson correlation coefficients among the start, duration, central moments (i.e., variance, skew and kurtosis), and two the first two eigenvectors of a principal coordinates analysis (PCoA) characterizing flowering schedules in populations of *Lythrum salicaria* grown in a common garden. The metrics were calculated using either (A) flowering schedule characteristics *averaged* across individuals within each population ($n = 13$), (B) the *residual* correlations calculated on deviations of individuals from their population means ($n = 369$), or (C) *aggregated* schedules for individuals pooled within populations ($n = 13$). Circles are shown for statistically significant correlations ($p < 0.05$), colour-coded by correlation coefficient (i.e., -1 to 1 as shown on the x-axis).

Figure 3. Principal Coordinates Analysis (PCoA) of Kolmogorov-Smirnov distances between flowering schedules of 369 plants from 13 populations of *Lythrum salicaria* sampled along a latitudinal gradient and grown in a common garden at the Koffler Scientific Reserve in Newmarket, ON. Point colors correspond to latitude of origin, ranging from dark (48° N) to light (38° N). Bar plots show representative flowering schedules of four individual plants. PCoA1 is inversely related to the central moments of variance and mean such that points on the right represent individuals with lower flowering schedule variance and an earlier peak.

Figure 4. Bivariate plots showing relations between latitude and flowering schedule characteristics of 13 populations of *Lythrum salicaria* grown in a common garden field experiment. The bootstrapped estimate and 95% confidence intervals of *average* individual flowering schedule characteristics are shown for each population along the left column. The bootstrapped estimate and 95% confidence intervals of the *aggregate* flowering schedules for all flowers within each population are shown in the middle column. The right column indicates the 'emergent properties' of each population, defined as the aggregate minus the average.

Appendix S1. The distribution of 369 individuals and 216 seed families representing 13 populations of *Lythrum salicaria* sampled along a latitudinal gradient and grown in a common garden at the Koffler Scientific Reserve in Newmarket, ON. Point colors correspond to latitude of origin ranging from dark (48° N) to light (38° N).

Latitude (°)	Number of seed families	Number of individuals
38.75261	15	23
40.34025	15	33
41.31283	18	24
42.23822	20	34
42.52306	12	34
43.69719	19	21
44.49339	19	38
44.59489	19	22
45.26278	16	26
45.43194	17	36
45.49000	16	23
47.69081	14	32
48.47892	16	23

FIGURE LEGENDS

Figure 1. Hypothetical flowering schedules of (A) five individuals with different characteristics and (B) their pooled aggregate flowering schedule. Flowering schedules are scaled to the proportion of total flower production (P_t) over time (t).

Figure 2. Matrices showing pairwise Pearson correlation coefficients among the start, duration, central moments (i.e., variance, skew and kurtosis), and two the first two eigenvectors of a principal coordinates analysis (PCoA) characterizing flowering schedules in populations of *Lythrum salicaria* grown in a common garden. The metrics were calculated using either (A) flowering schedule characteristics *averaged* across individuals within each population ($n = 13$), (B) the *residual* correlations calculated on deviations of individuals from their population means ($n = 369$), or (C) *aggregated* schedules for individuals pooled within populations ($n = 13$). Circles are shown for statistically significant correlations ($p < 0.05$), colour-coded by correlation coefficient (i.e., -1 to 1 as shown on the x-axis).

Figure 3. Principal Coordinates Analysis (PCoA) of Kolmogorov-Smirnov distances between flowering schedules of 369 plants from 13 populations of *Lythrum salicaria* sampled along a latitudinal gradient and grown in a common garden at the Koffler Scientific Reserve in Newmarket, ON. Point colors correspond to latitude of origin, ranging from dark (48° N) to light (38° N). Bar plots show representative flowering schedules of four individual plants. PCoA1 is inversely related to the central moments of variance and mean such that points on the right represent individuals with lower flowering schedule variance and an earlier peak.

Figure 4. Bivariate plots showing relations between latitude and flowering schedule characteristics of 13 populations of *Lythrum salicaria* grown in a common garden field experiment. The bootstrapped estimate and 95% confidence intervals of *average* individual flowering schedule characteristics are shown for each population along the left column. The bootstrapped estimate and 95% confidence intervals of the *aggregate* flowering schedules for all flowers within each population are shown in the middle column. The right column indicates the 'emergent properties' of each population, defined as the aggregate minus the average.