Latitudinal clines in floral display associated with adaptive evolution during a biological invasion

Mia N. Akbar[1], Dale R. Moskoff[1], Spencer C.H. Barrett[2], Robert I. Colautti[1,3]

[1]Department of Biology, Queen's University, Kingston, ON, Canada.
[2]Department of Ecology & Evolutionary Biology, University of Toronto, Toronto, ON, Canada.

[3]Corresponding author: robert.colautti@queensu.ca

Running head: Floral display clines in *Lythrum salicaria*

ABSTRACT

Premise: Flowering phenology strongly influences reproductive success in plants. Days to first

flower is easy to quantify and widely used to characterize phenology, but reproductive fitness

depends on the full schedule of flower production over time.

Methods: We examined floral display traits associated with rapid adaptive evolution and range

expansion among thirteen populations of *Lythrum salicaria*, sampled along a 10-degree

latitudinal gradient in eastern North America. Growing these collections in a common garden

field experiment at a mid-latitude site, we quantified flowering schedule variation using Principal

Coordinates Analysis (PCoA) and quantitative metrics analogous to central moments of

probability distributions (i.e., mean, variance, skew, and kurtosis).

Key Results: Consistent with earlier evidence for adaptation to shorter growing seasons, we

found that populations from higher latitudes had earlier start and mean flowering day, on

average, when compared to populations from southern latitudes. Clines in flowering skew

increased with latitude whereas kurtosis decreased, consistent with a bet-hedging strategy in

biotic environments with more herbivores and greater competition for pollinators.

Conclusions: Heritable genetic clines in flowering schedules are consistent with biotic and

abiotic responses to natural selection, potentially contributing to rapid evolution and range

expansion of this invasive species.

INTRODUCTION

Phenology affects how organisms interact with their environment, with important consequences

for their survival and reproduction. As targets of natural selection and determinants of population

vital rates, the timing of growth, reproduction, and other phenological traits are crucial

components of eco-evolutionary dynamics. Phenology governs interaction with mutualists

(Fenster et al., 2004), antagonists (Strauss & Whittall, 2006) and abiotic factors (Jin *et al.*, 2019)

that can vary over space and time. Consequently, plants experience spatial and temporal

variation in natural selection on flowering phenology (Kudo, 2007), sometimes resulting in the

evolution of local adaptation (Anderson et al., 2012; Colautti & Barrett, 2013; Preite et al.,

2015). As such, flowering phenology is a useful composite trait for investigating evolutionary

responses to past and future environmental challenges (Fitchett et al., 2015; D. W. Inouye, 2022;

Panchen & Gorelick, 2017; Prather et al., 2023).

Days to first flower (i.e., flowering time) is a simple yet informative metric with which to

study the ecology and evolution of phenology in plant populations. For example, latitudinal and

altitudinal clines in days to first flower are commonly reported in a variety of plant taxa

(Alexander et al., 2009; Colautti & Lau, 2015; Ensing & Eckert, 2019; Halbritter et al., 2018;

Olsson & Ågren, 2002; Panchen, 2022), and reciprocal transplant experiments have confirmed

that these clines are an important component of local adaptation (Anderson et al., 2012; Colautti

& Barrett, 2013; Griffith & Watson, 2005). Flowering time clines are consistent with local

adaptation models characterized by stabilizing selection within populations but a shift in optimal

flowering time along environmental gradients (de Villemereuil et al., 2020; Gauzere et al., 2020;

Kirkpatrick & Barton, 1997).

In contrast to the prediction of stabilizing selection, empirical measurements of selection

differentials suggest that flowering time is typically under directional selection within

populations (Munguía-Rosas et al., 2011). Despite evidence for consistent directional selection,

long-term phenology records show a variety of responses to climate change -- from earlier to

later to no change in flowering time, depending on the species (Rafferty et al., 2020; Wolkovich

et al., 2012). Thus, models and empirical observations of geographic clines and local adaptation

appear to disagree with observed measurements of natural selection and variable phenotypic

shifts in natural populations (Austen et al., 2017). Antagonistic selection on unmeasured traits

correlated with the onset of flowering could help to resolve this apparent contradiction.

Day of first flowering is but one aspect of the "flowering schedule" of individual plants --

a time-series of flower production that characterizes reproductive opportunity (CaraDonna et al.,

2014; Fox, 2003; B. D. Inouye et al., 2019; Newstrom et al., 1994). While rarely examined in

studies of flowering phenology evolution, different characteristics of the flowering schedule can

have important effects on reproductive fitness (Forrest & Thomson, 2010; Fox, 2003). In

addition to flowering time, the shape of the flowering schedule can be quantified using

ordination methods like Principal Coordinates Analysis (PCoA; Austen et al., 2014) and metrics

analogous to central moments of probability distributions including the mean, variance, skew,

and kurtosis (Box 1).

Not to be confused with the central moments of a statistical population or sample, the

analogous characteristics of a flowering schedule are not probabilistic, but rather quantify

temporal variation in relative reproductive investment through time. These schedule

characteristics use the same mathematical equations as probability moments (Table 1) but are not

shaped by probabilistic process and therefore may not necessarily correlate with onset or

duration of flowering (Fig. 1). These metrics have rarely been used in studies of flowering

phenology, yet they have the potential to offer a novel perspective on selection and trade-offs

affecting the evolution of phenology.

Central moments of individual flowering schedules describe the timing and intensity of

reproductive opportunity, and thus are likely to experience strong selection from abiotic or biotic

features of the environment. We are not aware of studies quantifying evolutionary divergence of

entire flowering schedules among naturally occurring populations, but there are reasons to expect

populations to evolve differences. For example, individuals that experience competition for

pollinators may benefit from showy inflorescences that enhance pollinator attraction (O'Neil,

1997; Thomson, 1980), which would produce a flowering schedule with a more positive skew

and longer duration. Herbivores and seed predators often select for delayed peak reproduction

(Biere & Antonovics, 1996; Elzinga et al., 2007; Juenger & Bergelson, 1998; Wright &

Meagher, 2003) but may also favour a bet-hedging strategy such that a long flowering duration,

high flowering schedule variance, and positive kurtosis would be advantageous in the presence

of intense herbivory.

In addition to biotic interactions, flowering schedules are likely to be shaped by abiotic

factors that determine the timing and duration of environmental conditions favorable to the

development of reproductive organs. Shorter growing seasons typically favour a more rapid

flowering strategy (Austen & Weis, 2015; Colautti & Barrett, 2013; Griffith & Watson, 2005)

characterized by an earlier onset of flowering, with a shorter duration and smaller variance of the

flowering schedule. In marginal environments with less predictable growing conditions, it may

be advantageous for plants to produce a few flowers over a longer timeframe as a bet-hedging

strategy (Forrest & Thomson, 2010; Simons, 2011; Tufto, 2015), resulting in a long-tailed

leptokurtic or skewed flowering schedule.

Environmental gradients associated with latitude can be useful for studying phenotypic

evolution if natural selection covaries, for example with shorter growing seasons and weaker

biotic interactions towards polar latitudes (reviewed in Zvereva & Kozlov, 2021). Adaptation

along environmental gradients is commonly manifested as clines in phenotypic traits that are

observable when individuals are grown in a common garden environment (Langlet, 1971;

Schwinning et al., 2022). For example, adaptive flowering time clines evolved rapidly in the

perennial wetland herb *Lythrum salicaria* (purple loosestrife) over a century of invasion across

North America. Selection splines (Colautti & Barrett, 2010), trait correlations (Colautti &

Barrett, 2011), reciprocal transplants (Colautti & Barrett, 2013), and herbarium records (Wu &

Colautti, 2022) collectively support a model of selection in which shorter growing seasons

toward the northern range limit favour earlier flowering. In contrast, longer growing seasons in

the southern part of the introduced range relax selection on flowering time and favor plants that

flower later but grow larger in stature. However, by focusing on the first day of flowering,

previous studies of *L. salicaria* overlooked other characteristics of the flowering schedule, which

may be important for understanding adaptation and constraint during its evolution and invasion

in eastern North America.

Here, we investigate clines in flowering schedules among thirteen populations of *L.*

salicaria sampled across ten degrees of latitude in eastern North America and grown in a

common garden field study at the Koffler Scientific Reserve north of Toronto, Ontario, Canada.

We use PCoA and central moments characteristics of individual flowering schedules to address

the following questions: (1) Are common phenological metrics, like the onset and duration of

flowering, associated with overlooked features of the flowering schedule such as the mean,

variance, skew and kurtosis? (2) Are there latitudinal clines in flowering schedules that are

consistent with the adaptive hypotheses outlined above? Specifically, we predicted that the

longer growing season length and more intense biotic interactions at southern latitudes should

create stronger competition for pollinators and higher rates of herbivory in southern populations.

These conditions seem likely to favour genotypes with: (i) a longer flowering duration and (ii) a

later start and mean flowering day, but also (iii) an increase in flowering schedule variance (iv) a

more negative skew, with flowering focused later in the schedule and (v) a more positive

kurtosis, with a concentrated peak and rare flowering events far from the mean flowering day

(see Fig. 2). In line with these predicted adaptive changes, we expect the primary PCoA axes to

reflect population differentiation in flowering schedules based on latitude of origin.

METHODS

Experimental design --

Details of the field experiment are provided in Colautti & Barrett (2010) and the seeds used were

collected along a 10-degree latitudinal gradient in eastern North America, as described in

Montague *et al.* (2008). Briefly, populations were chosen to represent a latitudinal gradient from

Timmins, Ontario (48.48 °N) to Easton, Maryland (38.75 °N) and were surveyed during the

summer of 2003 (see Appendix S1 for a list of locations). When 90% of plants had finished

flowering in the fall, 20-41 stems were randomly harvested from each population. Eight seeds

from each of 20 seed families from each of 13 populations were grown at the Koffler Scientific

Reserve at Jokers Hill, located approximately at the geographic center of the latitudinal gradient

(44.03 °N). Early in the growing season (early May), plants were sprayed with insecticide to

reduce incidence of herbivorous insects. This was sufficient to inhibit colonization of biocontrol

beetles in the area, without impacting pollinator activity later in the season when plants became

reproductive (July through September). The primary purpose of this experiment was to

characterize flowering time clines and genetic trade-offs in a multi-year (2006-2008) common

garden study, as reported in Colautti & Barrett (2010, 2011). In 2007 we additionally recorded

all open flowers on every stem of a subset of 369 individuals of 216 seed families, evenly

represented across the 13 populations (Appendix S1). Our daily census records of this subset of

plants occurred over 125 days on a rotating schedule such that each individual was surveyed

every five days from July 1 until September 30. A 5-day cycle was chosen because stigma

receptivity is nearly zero after 24 h of anthesis (Waites & Ågren, 2006) and flowers are fully

senescent after 5 days. Additionally, unpollinated flowers begin to senesce and abscise while

pollinated flowers begin to close within 48 h of anthesis, limiting the risk of counting the same

flowers on multiple census dates. All calculations and statistical analyses were conducted in R

version 4.2.2 (R Core Team, 2022) in R Studio (Posit team, 2024), and are available with the

original data from the Dryad repository (https://doi.org/10.5061/dryad.jdfn2z3jz).

Characterizing flowering schedules --

We characterized the flowering schedules of each individual plant using the equations shown in

Table 1, where the day that an open flower was first observed in the experiment is $t = 0$. The

number of flowers produced by plants in each population over a growing season is variable in

this experiment because stabilizing selection a our mid-latitude field site favours individuals that

balance flowering time with growth to produce more flowers relative to small, early flowering

phenotypes more characteristic of northern populations or large, late-flowering phenotypes

characteristic of more southern populations. Including flower number as a covariate in our

models would assume linearity between the number of flowers produced with the schedule

variance, skew, and kurtosis. Therefore, we instead standardized the number of observed flowers

for each individual on each census day as a proportion of their total number of open flowers

censused over the experiment. We then calculated a weighted census day by multiplying each

day by the proportion of flowers open on that day. The proportion of flowers and the weighted

census days were summed to calculate the weighted mean and other characteristics of individual

flowering schedules analogous to central moments (variance, skew and kurtosis) (Box 1; Table

1). Specific calculations are provided as custom R functions in the Dryad repository

(https://doi.org/10.5061/dryad.jdfn2z3jz).

In addition to calculating the characteristics of every individual's flowering schedule, we

wrote a bootstrap model in R to compute a bootstrapped mean and 95% confidence intervals

across 999 iterations for each population, and a similar bootstrap model to test for latitudinal

clines. In each iteration, individuals were hierarchically sampled with replacement from within

each population, and then the equations in Table 1 were used to calculate population averages.

The population averages of each iteration were also used to generate 999 estimates of intercept,

linear slope and quadratic regression coefficients to test for latitudinal clines. In this way we

generated bootstrap means and 95% confidence intervals for each population and for the

estimates of latitudinal clines. More details of the bootstrap models are available in R code from

the Dryad repository (https://doi.org/10.5061/dryad,jdfn2z3jz).

In addition to the start of flowering, duration, and central moments of flowering

schedules, we also used a Principle Coordinates Analysis (PCoA) to compare flowering schedule

shape among individuals, as described by Austen et al., (2014). Briefly, we computed a

dissimilarity matrix from raw flower counts using Kolmogorov-Smirnov (KS) distances (Sokal

& Rohlf, 1995) where each matrix cell was a KS statistic measuring the maximum distance

between two flowering schedules. We chose KS distance because it is robust to differences in

flower number and census schedules. However, the KS statistic is sensitive to differences in

scale, so flowering durations were standardized to run from 0 to 1 by subtracting the shortest

duration time from each individual and dividing by the difference of the longest and shortest

durations (Austen et al., 2014). We used the KS matrix as the input for the **cmdscale** function to

generate the PCoA axes. Since KS distances do not meet the triangle inequality condition of

PCoA, any variation that is unable to be converted to Euclidean distance is represented as

negative PCoA axes, which may lead to overestimation of variation explained by the first axes

(Austen et al., 2014; Legendre & Anderson, 1999; Podani & Miklós, 2002). Therefore, we

adjusted our estimates of proportion of variation explained to only include positive eigenvalues

(Austen et al., 2014). We then generated vectors to plot each flowering schedule trait in

multivariate space by computing the standardized covariance of the central moments with all

PCoA axes, filtered for positive eigenvalues (Legendre et al., 1998). It is important to note that

the PCoA axes characterize variation in the shape of flowering schedules among individuals but

not variation in onset and duration of flowering. However, projecting flowering characteristics

onto the first two PCoA axes provides a visualization of how different flowering schedule

characteristics relate to the major axes of variation in schedule shape. Additionally, we visualize

each individual, colour-coded by latitude, to see how schedule shape varies among and within

populations, and along a latitudinal gradient.

Correlations of flowering schedule characteristics --

Our first research question addresses whether conventional metrics (i.e., onset and duration of

flowering) correspond to more detailed characteristics of the flowering schedule (i.e., mean,

variance, skew, kurtosis, and PCoA axes). To investigate these relations, we calculated pairwise

Pearson correlations to compare two scales of analysis, namely (A) average schedules of each

population ($n = 13$), and (B) the deviation of each individual from its population average ($n =$

369). In addition to comparing conventional metrics with other flowering schedule

characteristics, the correlational structure of these matrices can also be instructive for testing

evolutionary hypotheses. For example, the evolutionary divergence of flowering schedules along

biotic and abiotic gradients should produce stronger correlations among population when

compared to correlations among individuals within populations.

RESULTS

Flowering schedule characteristics --

Pearson product-moment correlations revealed significant relations among commonly measured

traits of the flowering schedule (i.e., start and duration) and more detailed characteristics (i.e.,

mean, variance, skew, and kurtosis), both among and within populations (Fig. 3). There was no

significant correlation between days to flower and duration or flowering variance among

population average flowering schedules. However, there was a significant negative correlation

for start date with skew and a positive correlation with kurtosis, indicating that populations with

a later start date, on average, tended to have individuals with flowering clustered at later dates

(Fig. 3A).

For individual flowering schedules, residual within-population correlations were

generally weaker when compared to among-population correlations (Fig. 3B). However, there

were negative correlations between days to flower and duration and variance of flowing among

individuals within populations (Fig. 3B). These correlations were notably weaker and not

statistically significant among population means (Fig. 3A). Skew and kurtosis were strongly

negatively correlated both among (Fig. 3A) and within populations (Fig. 3B).

Principal Coordinates Analysis --

In addition to the above characteristics, we used Principal Coordinates Analysis (PCoA) to

characterize similarities and differences among flowering schedules after controlling for

variation in start date and duration. The PCoA of all 369 individuals resulted in 369 independent

eigenvectors, of which 192 were positive. After adjusting the proportion of variation to include

only positive eigenvalues, only the first two of 192 PCoA axes had eigenvectors that explained

more than 10% of the variation in flowing schedule shape, and together they explained 35% of

the variation (Fig. 4). By comparison, the next ten eigenvectors combined explained another 35%

of variation, with the remaining 180 positive eigenvectors accounting for 30% of the variation.

Thus, we focus here on interpreting the first two eigenvectors. The first axis (PCoA1) was

negatively correlated with mean days to flower, after controlling for variation in start and

duration of flowering (Fig. 4). In contrast, higher values of PCoA2 were primarily associated

with higher flowering variance and lower kurtosis in flowering schedules. Despite standardizing

individual flowering schedules to have the same onset and duration of flowering prior to PCoA,

we found that the start and duration of flowering remained correlated with PCoA1 and PCoA2,

indicating that central moments are correlated with other, unmeasured shape characteristics.

Latitudinal clines --

The significant correlations that we detected among population average flowering schedule traits

in our central moments and PCoA analyses also covaried with latitude. For example, the start,

duration, mean, and kurtosis of flowering schedules were each significantly negatively correlated

with latitude (Fig. 5). This means that on average, individuals from southern populations start

flowering later, have later mean flowering days, and more 'outlier' flowers opening far from the

mean flowering date, when compared to northern populations. Additionally, average skew was

negative for all populations, but more negative at lower latitudes, whereas kurtosis was close to

zero but slightly positive in southern populations (Fig. 5). In other words, individuals from

southern populations tended to produce more flowers far from peak flowering whereas flowering

schedules in individuals from northern populations are more symmetric and concentrated (Fig.

5), even though there were no significant latitudinal clines in schedule variance.

DISCUSSION

There is growing evidence for rapid evolution in invasive species (reviewed in Barrett et al.

2017), but it remains uncertain how often adaptive evolution contributes to their spread. Previous

studies focusing on the onset of flowering provided only a partial view of how plants allocate

flowering resources (Box 1), leaving uncertainty about the extent to which flowering phenology

is adaptive. To address this knowledge gap, we used a common garden study to test for

latitudinal clines in complete flowering schedules in the wetland invader *Lythrum salicaria*. We

found that the onset of flowering was significantly correlated with other aspects of the flowering

schedule including among-population averages (Fig. 2A), and among individuals within

populations (Fig. 2B). Furthermore, the significant among-population correlations were largely

consistent with the evolution of correlated traits along latitudinal clines (Fig. 5). Compared to

northern genotypes, plants from lower latitudes flowered later, for a longer duration, and with a

more negative skew and leptokurtic shape (Fig. 5). Below we discuss the implications of these

results for understanding the evolutionary causes and ecological consequences of flowering

schedule diversity.

Onset and duration of flowering --

Date of first flower and duration of flowering are relatively easy to measure and are commonly

reported as focal traits in phenological studies. However, the reproductive fitness of individuals

depends on the full schedule of flower production, and it has not been clear how well the onset or

duration of flowering represents other flowering schedule characteristics. In our study, the use of

PCoA revealed substantial variation in flowering schedules even after standardizing for the onset

and duration of flowering (Fig. 4). However, correlations of both PCoA1 and PCoA2 with onset

and duration (Fig. 4) indicated that this multi-dimensional variation in schedule appears to

covary with commonly measured parameters. In line with this conclusion, the onset of flowering

was strongly correlated with the mean flowering day, both among population averages (Fig. 3A)

and among individuals within populations (Fig. 3B). Therefore, the onset of flowering appears to

be a good predictor of peak flowering in this study system.

Within populations, one might predict a trade-off between the onset and duration of

flowering. Instead, we found that early flowering plants also flowered for a longer duration with

a higher flowering variance (Fig. 3B). Although counterintuitive, this result is consistent with

studies of other species that have reported that early flowering plants also flower for a longer

duration (Austen et al., 2017; Hendry & Day, 2005). Variation in microhabitat quality (e.g.,

nutrient flux) and/or segregation of deleterious alleles within populations (e.g., inbreeding

depression) could account for this correlation, with early/longer flowering plants representing

microenvironments or genotypes of higher quality. Yet, without evidence for trade-offs it is

difficult to explain how latitudinal clines in flowering time could evolve.

In contrast to the highly significant relations observed between onset, duration, and

variance of flowering within populations (Fig. 3B), the same correlations were non-significant

when calculated among population averages (Fig. 3A). Instead, duration and variance were not

significantly correlated with latitude because populations originating closer to the common

garden latitude site tended to have the longest duration and highest variance. This result is

consistent with previously proposed models of constrained adaptation in this species (Colautti,

Eckert, et al., 2010; Colautti & Barrett, 2011, 2013; Wu & Colautti, 2022). Specifically,

populations that are locally adapted to the latitude of the field experiment balance a trade-off

between early flowering and larger growth to maximize the flowering duration and variance. In

contrast, populations from more distant latitudes have lower flowering duration and variance.

Early flowering northern plants are smaller and run out of resources for flower production earlier

in the season while larger plants from the south have their flowering schedules truncated by the

end of the growing season. Future experiments replicating common gardens at higher and lower

latitudes would be necessary to test for the predicted shift from a positive cline in duration and

variance at southern sites to a negative cline at higher latitudes with shorter growing seasons. In

addition to season length, the presence of biological control beetles are likely to alter flowering

schedules and may have different effects on northern vs southern population, which could be

evaluated with sprayed and unsprayed treatments in a replicated common garden.

Latitudinal clines --

Latitudinal clines in the onset of flowering have evolved independently in several unrelated

angiosperm species across North America (McGoey et al., 2020; Montague et al., 2008; Samis et

al., 2012; Wu & Colautti, 2022). In our study of *L. salicaria*, we additionally found latitudinal

clines in average mean, duration, skew and kurtosis of flowering schedules (Fig. 5). This

latitudinal differentiation is also reflected in Figure 4, where northern populations cluster at

intermediate values of PCoA1 and PCoA2 compared to mid-latitude (higher PCoA1 but lower

PCoA2) and southern populations (lower PCoA1 and lower PCoA2). These types of clines

indicate that individuals from southern environments have floral displays that peak later in the

season but produce a few open flowers far from the mean flowering date. These findings

correspond to predictions based on a bet-hedging strategy with a higher intensity of biotic

interactions predicted in southern environments. Previous meta-analyses have found increasing

biotic interactions at lower latitudes (Zvereva & Kozlov, 2021) and towards warmer range limits

(Paquette & Hargreaves, 2021), but we are not aware of prior studies showing evolved clines in

flowering schedules that correspond to clines in biotic interactions.

Lythrum salicaria is a self-incompatible outcrosser (Colautti et al., 2010; Darwin, 1877;

O'Neil, 1997) and individuals likely experience inter-specific competition for pollinators (King

& Sargent, 2012; Groulx & Sargent, 2018), which may be more intense in southern populations.

Additionally, several effective biological control insects have been released into eastern North

America, including *Gallerucella* beetles that feed on meristem tissue as larvae and can

drastically reduce flower production (Blossey & Notzold, 1995; Grevstad, 2006; Russell-Mercier

& Sargent, 2015). One of the factors along with climatic adaptation that may contribute to the

later flowering time and more negative skew in plants from southern populations is the

possibility of selection from insect herbivores that intensifies at mid-growing season.

Higher values of kurtosis are consistent with the hypothesis that a strategy of producing a

few flowers over a long period of time while also optimizing for an intense flowering peak to

compete for pollinators occurs in southern populations. We cannot easily parse the specific

effects of biotic and abiotic factors based on flowering clines alone, but we can make the

prediction that clines in herbivore and pollinator activity are likely to be more patchy and less

predictable by latitude compared to season length. It is therefore interesting that we detected

stronger clines in average start and mean flowering dates (Fig. 5), which we predicted would be

more sensitive to season length, in contrast to kurtosis, which we predicted as a bet-hedging

strategy associated with biotic interactions. Skew showed an intermediate pattern (Fig. 5) and

could be an adaptation to biotic selection. However, it is also possible that the pattern of skew

may partly be an artifact of our mid-latitude common garden site. Specifically, a lack of

pollinator activity and cooler growing conditions towards the end of the growing season could

act as cues for plants to stop flowering. This would effectively truncate the flowering schedules

of late-flowering plants, creating a more negative skew. Indeed, this could explain the negative

correlation between days to flower and skew among population means (Fig. 3A).

Conclusions --

Our study demonstrated that populations of *L. salicaria* in eastern North America have evolved

differences in flowering schedules within a century of invasion. This phenological differentiation

among populations includes predictable changes in the timing of resource allocation to flower

development and floral display and adds to the growing list of phenotypic traits that are targets of

selection as invasive species spread along environmental gradients (MacDougall et al., 2006;

McGoey et al., 2020; Molina-Montenegro et al., 2018; Urbanski et al., 2012). The ecological

and evolutionary insights drawn from observations of flowering start dates should be interpreted

cautiously. Instead, more detailed features of individual flowering schedules can be characterised

by metrics drawn from central moments theory (Table 1). As we have shown for *L. salicaria*

from eastern North America, these more refined characteristics of individual flowering schedules

help to evaluate adaptive hypotheses for flowering phenology to develop a deeper understanding

of the diverse ways that species respond to changing environments.

ACKNOWLEDGEMENTS

The authors thank Z. Burivalova and R. Mackenzie for field assistance, and to E.J. Austen, A.E.

Weis and J. Friedman for help with the analysis, and to E.J. Austen and an anonymous reviewer

for feedback on the manuscript. Funding was provided from NSERC Discovery grants to

S.C.H.B. and R.I.C. and NSERC CGSM to M.N.A.

AUTHOR CONTRIBUTIONS

402 M.N.A. led the data analysis with assistance from D.R.M., S.C.H.B., and R.I.C. The

403 conceptualization and experimental design were developed by S.C.H.B. and R.I.C., and all

404 authors contributed to the manuscript, based on an original draft written by M.N.A.

DATA AVAILABILITY STATEMENT

407 Raw data and fully reproducible R code for all figures, tables, and statistical analyses are

408 available in the Dryad database (https://doi.org/10.5061/dryad.jdfn2z3jz)

LITERATURE CITED

Alexander, J. M., Edwards, P. J., Poll, M., Parks, C. G., & Dietz, H. (2009). Establishment of parallel altitudinal clines in traits of native and introduced forbs. *Ecology*, *90*(3), 612–622. https://doi.org/10.1890/08-0453.1

Anderson, J. T., Inouye, D. W., McKinney, A. M., Colautti, R. I., & Mitchell-Olds, T. (2012). Phenotypic plasticity and adaptive evolution contribute to advancing flowering phenology in response to climate change. *Proceedings of the Royal Society B: Biological Sciences*, *279*(1743), 3843–3852. https://doi.org/10.1098/rspb.2012.1051

Austen, E. J., Jackson, D. A., & Weis, A. E. (2014). Describing flowering schedule shape through multivariate ordination. *International Journal of Plant Sciences*. https://doi.org/10.1086/673934

Austen, E. J., Rowe, L., Stinchcombe, J. R., & Forrest, J. R. K. (2017). Explaining the apparent paradox of persistent selection for early flowering. *New Phytologist*, *215*(3), 929–934. https://doi.org/10.1111/nph.14580

Austen, E. J., & Weis, A. E. (2015). What drives selection on flowering time? An experimental manipulation of the inherent correlation between genotype and environment: causes of selection on flowering time. *Evolution*, *69*(8), 2018–2033. https://doi.org/10.1111/evo.12709

Barrett, S. C. H., Colautti, R. I., Dlugosch, K. M., & Rieseberg, L. H. (Editors) (2017). *Invasion Genetics: The Baker and Stebbins Legacy*. John Wiley, London.

Biere, A., & Antonovics, J. (1996). Sex-specific costs of resistance to the fungal pathogen Ustilago violacea (*Microbotryum violaceum*) in Silene alba. *Evolution; International*

Journal of Organic Evolution, *50*(3), 1098–1110. https://doi.org/10.1111/j.1558-

5646.1996.tb02350.x

Blossey, B., & Notzold, R. (1995). Evolution of increased competitive ability in invasive

nonindigenous plants: A hypothesis. *Journal of Ecology*, *83*(5), 887–889.

https://doi.org/10.2307/2261425

CaraDonna, P. J., Iler, A. M., & Inouye, D. W. (2014). Shifts in flowering phenology reshape a

subalpine plant community. *Proceedings of the National Academy of Sciences of the*

United States of America, *111*(13), 4916–4921. https://doi.org/10.1073/pnas.1323073111

Colautti, R. I., & Barrett, S. C. H. (2010). Natural selection and genetic constraints on flowering

phenology in an invasive plant. *International Journal of Plant Sciences*, *171*(9), 960–971.

https://doi.org/10.1086/656444

Colautti, R. I., & Barrett, S. C. H. (2011). Population divergence along lines of genetic variance

and covariance in the invasive plant *Lythrum salicaria* in eastern North America.

Evolution, *65*(9), 2514–2529. https://doi.org/10.1111/j.1558-5646.2011.01313.x

Colautti, R. I., & Barrett, S. C. H. (2013). Rapid adaptation to climate facilitates range expansion

of an invasive plant. *Science, 342*(6156), 364–366.

https://doi.org/10.1126/science.1242121

Colautti, R. I., Eckert, C. G., & Barrett, S. C. H. (2010). Evolutionary constraints on adaptive

evolution during range expansion in an invasive plant. *Proceedings of the Royal Society*

B: Biological Sciences, *277*(1689), 1799–1806. https://doi.org/10.1098/rspb.2009.2231

Colautti, R. I., & Lau, J. A. (2015). Contemporary evolution during invasion: Evidence for

differentiation, natural selection, and local adaptation. *Molecular Ecology*, *24*(9), 1999–

2017. https://doi.org/10.1111/mec.13162

Colautti, R. I., White, N. A., & Barrett, S. C. H. (2010). Variation of self-incompatibility within

invasive populations of purple loosestrife (*Lythrum salicaria* L.) from eastern North

America. *International Journal of Plant Sciences*, *171*(2), 158–166.

https://doi.org/10.1086/649023

Darwin, C. (1877). *The Different Forms of Flowers on Plants of the Same Species*. John Murray,

Albermarle Street.

de Villemereuil, P., Charmantier, A., Arlt, D., Bize, P., Brekke, P., Brouwer, L., Cockburn, A.,

Côté, S. D., Dobson, F. S., Evans, S. R., Festa-Bianchet, M., Gamelon, M., Hamel, S.,

Hegelbach, J., Jerstad, K., Kempenaers, B., Kruuk, L. E. B., Kumpula, J., Kvalnes, T., …

Chevin, L.-M. (2020). Fluctuating optimum and temporally variable selection on

breeding date in birds and mammals. *Proceedings of the National Academy of Sciences of*

the United States of America, *117*(50), 31969–31978.

https://doi.org/10.1073/pnas.2009003117

Elzinga, J. A., Atlan, A., Biere, A., Gigord, L., Weis, A. E., & Bernasconi, G. (2007). Time after

time: Flowering phenology and biotic interactions. *Trends in Ecology & Evolution*, *22*(8),

432–439. https://doi.org/10.1016/j.tree.2007.05.006

Ensing, D. J., & Eckert, C. G. (2019). Interannual variation in season length is linked to strong

co-gradient plasticity of phenology in a montane annual plant. *New Phytologist*, *224*(3),

1184–1200. https://doi.org/10.1111/nph.16009

Fenster, C. B., Armbruster, W. S., Wilson, P., Dudash, M. R., & Thomson, J. D. (2004).

Pollination syndromes and floral specialization. *Annual Review of Ecology, Evolution,*

and Systematics, *35*, 375–403.

Fitchett, J. M., Grab, S. W., & Thompson, D. I. (2015). Plant phenology and climate change:

Progress in methodological approaches and application. *Progress in Physical Geography:*

Earth and Environment, *39*(4), 460–482. https://doi.org/10.1177/0309133315578940

Forrest, J., & Thomson, J. D. (2010). Consequences of variation in flowering time within and

among individuals of *Mertensia fusiformis* (Boraginaceae), an early spring wildflower.

American Journal of Botany, *97*(1), 38–48. https://doi.org/10.3732/ajb.0900083

Fox, G. A. (2003). Assortative mating and plant phenology: Evolutionary and practical

consequences. *Evolutionary Ecology Research*, *5*(1), 18.

Gauzere, J., Teuf, B., Davi, H., Chevin, L.-M., Caignard, T., Leys, B., Delzon, S., Ronce, O., &

Chuine, I. (2020). Where is the optimum? Predicting the variation of selection along

climatic gradients and the adaptive value of plasticity. A case study on tree phenology.

Evolution Letters, *4*(2), 109–123. https://doi.org/10.1002/evl3.160

Grevstad, F. S. (2006). Ten-year impacts of the biological control agents *Galerucella pusilla* and

G. calmariensis (Coleoptera: Chrysomelidae) on purple loosestrife (*Lythrum salicaria*) in

Central New York State. *Biological Control*, *39*(1), 1–8.

https://doi.org/10.1016/j.biocontrol.2006.03.007

Griffith, T. M., & Watson, M. A. (2005). Stress avoidance in a common annual: Reproductive

timing is important for local adaptation and geographic distribution. *Journal of*

Evolutionary Biology, *18*(6), 1601–1612. https://doi.org/10.1111/j.1420-

9101.2005.01021.x

Groulx, A. F., & Sargent, R. D. (2018). Purple loosestrife provides long-distance pollinator

attraction to a coflowering native species. *International Journal of Plant Sciences*, 179

(8), 593-602.

Halbritter, A. H., Fior, S., Keller, I., Billeter, R., Edwards, P. J., Holderegger, R., Karrenberg, S.,

Pluess, A. R., Widmer, A., & Alexander, J. M. (2018). Trait differentiation and

adaptation of plants along elevation gradients. *Journal of Evolutionary Biology*, *31*(6),

784–800. https://doi.org/10.1111/jeb.13262

Hendry, A. P., & Day, T. (2005). Population structure attributable to reproductive time: Isolation

by time and adaptation by time. *Molecular Ecology*, *14*(4), 901–916.

https://doi.org/10.1111/j.1365-294X.2005.02480.x

Inouye, B. D., Ehrlén, J., & Underwood, N. (2019). Phenology as a process rather than an event:

From individual reaction norms to community metrics. *Ecological Monographs*, *89*(2),

e01352. https://doi.org/10.1002/ecm.1352

Inouye, D. W. (2022). Climate change and phenology. *WIREs Climate Change*, *13*(3), e764.

https://doi.org/10.1002/wcc.764

Juenger, T., & Bergelson, J. (1998). Pairwise versus diffuse natural selection and the multiple

herbivores of scarlet gilia, *Ipomopsis aggregata*. *Evolution; International Journal of*

Organic Evolution, *52*(6), 1583–1592. https://doi.org/10.1111/j.1558-

5646.1998.tb02239.x

King, V. M., & Sargent, R. D. (2012). Presence of an invasive plant species alters pollinator

visitation to a native. *Biological Invasions*, 14, 1809-1818.

Kirkpatrick, M., & Barton, N. H. (1997). Evolution of a species' range. *The American Naturalist*,

150(1), 1–23. https://doi.org/10.1086/286054

Kudo, G. (2007). Flowering phenologies of animal-pollinated plants: Reproductive strategies and

agents of selection. In *Ecology and Evolution of Flowers*. Oxford University Press,

Incorporated.

Langlet, O. (1971). Two hundred years of genecology. *Taxon*, *20*(5–6), 653–721. https://doi.org/10.2307/1218596

Legendre, P., & Anderson, M. J. (1999). Distance-based redundancy analysis: Testing multispecies responses in multifactorial ecological experiments. *Ecological Monographs*, *69*(1), 1–24.

Legendre, P., Legendre, L., & others. (1998). Numerical ecology: Developments in environmental modelling. *Developments in Environmental Modelling*, *20*(1).

MacDougall, A. s., Boucher, J., Turkington, R., & Bradfield, G. e. (2006). Patterns of plant invasion along an environmental stress gradient. *Journal of Vegetation Science*, *17*(1), 47–56. https://doi.org/10.1111/j.1654-1103.2006.tb02422.x

McGoey, B. V., Hodgins, K. A., & Stinchcombe, J. R. (2020). Parallel flowering time clines in native and introduced ragweed populations are likely due to adaptation. *Ecology and Evolution*, *10*(11), 4595–4608. https://doi.org/10.1002/ece3.6163

Molina-Montenegro, M. A., Acuña-Rodríguez, I. S., Flores, T. S. M., Hereme, R., Lafon, A., Atala, C., & Torres-Díaz, C. (2018). Is the success of plant invasions the result of rapid adaptive evolution in seed traits? Evidence from a latitudinal rainfall gradient. *Frontiers in Plant Science*, *9*, 208. https://doi.org/10.3389/fpls.2018.00208

Montague, J. L., Barrett, S. C. H., & Eckert, C. G. (2008). Re-establishment of clinal variation in flowering time among introduced populations of purple loosestrife (*Lythrum salicaria*, Lythraceae). *Journal of Evolutionary Biology*, *21*(1), 234–245. https://doi.org/10.1111/j.1420-9101.2007.01456.x

Munguía-Rosas, M. A., Ollerton, J., Parra-Tabla, V., & De-Nova, J. A. (2011). Meta-analysis of phenotypic selection on flowering phenology suggests that early flowering plants are

favoured. *Ecology Letters*, *14*(5), 511–521. https://doi.org/10.1111/j.1461-

0248.2011.01601.x

Newstrom, L. E., Frankie, G. W., & Baker, H. G. (1994). A new classification for plant

phenology based on flowering patterns in lowland tropical rain forest trees at La Selva,

Costa Rica. *Biotropica*, *26*(2), 141–159. https://doi.org/10.2307/2388804

Olsson, K., & Ågren, J. (2002). Latitudinal population differentiation in phenology, life history

and flower morphology in the perennial herb *Lythrum salicaria*. *Journal of Evolutionary*

Biology, *15*(6), 983–996. https://doi.org/10.1046/j.1420-9101.2002.00457.x

O'Neil, P. (1997). Natural selection on genetically correlated phenological characters in *Lythrum*

salicaria L. (Lythraceae). *Evolution*, *51*(1), 267–274. https://doi.org/10.2307/2410980

Panchen, Z. A. (2022). Plant reproductive phenology along an elevation gradient in the extreme

environment of the Canadian High Arctic. *Plant Ecology & Diversity*, *15*(5–6), 213–226.

https://doi.org/10.1080/17550874.2022.2147804

Panchen, Z. A., & Gorelick, R. (2017). Prediction of Arctic plant phenological sensitivity to

climate change from historical records. *Ecology and Evolution*, *7*(5), 1325–1338.

https://doi.org/10.1002/ece3.2702

Paquette, A., & Hargreaves, A. L. (2021). Biotic interactions are more often important at

species' warm versus cool range edges. *Ecology Letters*, *24*(11), 2427–2438.

https://doi.org/10.1111/ele.13864

Podani, J., & Miklós, I. (2002). Resemblance coefficients and the horseshoe effect in principal

coordinates analysis. *Ecology*, *83*(12), 3331–3343.

Posit team. (2024). *RStudio: Integrated Development Environment for R*. Posit Software, PBC.

http://www.posit.co/

Prather, R. M., Dalton, R. M., barr, B., Blumstein, D. T., Boggs, C. L., Brody, A. K., Inouye, D.

W., Irwin, R. E., Martin, J. G. A., Smith, R. J., Van Vuren, D. H., Wells, C. P.,

Whiteman, H. H., Inouye, B. D., & Underwood, N. (2023). Current and lagged climate

affects phenology across diverse taxonomic groups. *Proceedings of the Royal Society B:*

Biological Sciences, *290*(1990), 20222181. https://doi.org/10.1098/rspb.2022.2181

Preite, V., Stöcklin, J., Armbruster, G. F. J., & Scheepens, J. F. (2015). Adaptation of flowering

phenology and fitness-related traits across environmental gradients in the widespread

Campanula rotundifolia. *Evolutionary Ecology*, *29*(2), 249–267.

https://doi.org/10.1007/s10682-015-9754-y

R Core Team. (2022). *R: A language and environment for statistical computing.* [Computer

software]. R Foundation for Statistical Computing. https://www.R-project.org/

Rafferty, N. E., Diez, J. M., & Bertelsen, C. D. (2020). Changing climate drives divergent and

nonlinear shifts in flowering phenology across elevations. *Current Biology*, *30*(3), 432-

441.e3. https://doi.org/10.1016/j.cub.2019.11.071

Russell-Mercier, J. L., & Sargent, R. D. (2015). Indirect effects of herbivory on plant–pollinator

interactions in invasive Lythrum salicaria. *American Journal of Botany*, *102*(5), 661–668.

https://doi.org/10.3732/ajb.1500043

Samis, K. E., Murren, C. J., Bossdorf, O., Donohue, K., Fenster, C. B., Malmberg, R. L.,

Purugganan, M. D., & Stinchcombe, J. R. (2012). Longitudinal trends in climate drive

flowering time clines in North American Arabidopsis thaliana. *Ecology and Evolution*,

2(6), 1162–1180. https://doi.org/10.1002/ece3.262

Schwinning, S., Lortie, C. J., Esque, T. C., & DeFalco, L. A. (2022). What common-garden

experiments tell us about climate responses in plants. *Journal of Ecology*, *110*(5), 986–

996. https://doi.org/10.1111/1365-2745.13887

Simons, A. M. (2011). Modes of response to environmental change and the elusive empirical

evidence for bet hedging. *Proceedings of the Royal Society B: Biological Sciences*,

278(1712), 1601–1609. https://doi.org/10.1098/rspb.2011.0176

Sokal, R. R., & Rohlf, F. J. (1995). *Biometry* (3rd ed.). WH Freeman.

Strauss, S. Y., & Whittall, J. B. (2006). Non-pollinator agents of selection on floral traits. In

Ecology and Evolution of Flowers (pp. 120–135). Oxford Biology.

Thomson, J. D. (1980). Skewed flowering distributions and pollinator attraction. *Ecology*, *61*(3),

572–579. https://doi.org/10.2307/1937423

Tufto, J. (2015). Genetic evolution, plasticity, and bet-hedging as adaptive responses to

temporally autocorrelated fluctuating selection: A quantitative genetic model. *Evolution*,

69(8), 2034–2049. https://doi.org/10.1111/evo.12716

Urbanski, J., Mogi, M., O'Donnell, D., DeCotiis, M., Toma, T., & Armbruster, P. (2012). Rapid

Adaptive Evolution of Photoperiodic Response during Invasion and Range Expansion

across a Climatic Gradient. *The American Naturalist*, *179*(4), 490–500.

https://doi.org/10.1086/664709

Waites, A. R., & Ågren, J. (2006). Stigma receptivity and effects of prior self-pollination on seed

set in tristylous *Lythrum salicaria* (Lythraceae). *American Journal of Botany*, *93*(1), 142–

147. https://doi.org/10.3732/ajb.93.1.142

Wolkovich, E. M., Cook, B. I., Allen, J. M., Crimmins, T. M., Betancourt, J. L., Travers, S. E.,

Pau, S., Regetz, J., Davies, T. J., Kraft, N. J. B., Ault, T. R., Bolmgren, K., Mazer, S. J.,

McCabe, G. J., McGill, B. J., Parmesan, C., Salamin, N., Schwartz, M. D., & Cleland, E.

E. (2012). Warming experiments underpredict plant phenological responses to climate

change. *Nature*, *485*(7399), Article 7399. https://doi.org/10.1038/nature11014

Wright, J. W., & Meagher, T. R. (2003). Pollination and seed predation drive flowering

phenology in *Silene latifolia* (Caryophyllaceae). *Ecology*, *84*(8), 2062–2073.

https://doi.org/10.1890/02-0676

Wu, Y., & Colautti, R. I. (2022). Evidence for continent-wide convergent evolution and stasis

throughout 150 y of a biological invasion. *Proceedings of the National Academy of*

Sciences of the United States of America, *119*(18), e2107584119.

https://doi.org/10.1073/pnas.2107584119

Zvereva, E. L., & Kozlov, M. V. (2021). Latitudinal gradient in the intensity of biotic

interactions in terrestrial ecosystems: Sources of variation and differences from the

diversity gradient revealed by meta-analysis. *Ecology Letters*, *24*(11), 2506–2520.

https://doi.org/10.1111/ele.13851

Box 1: Summary characteristics of flowering schedules

Central moments (i.e., mean, variance, skew, and kurtosis) arise from probability theory and characterize distributions that can be visualized as histograms with observed values on the x-axis and probability density, frequency, or number of observations on the y-axis. Here, we use the same equations to describe analogous characteristics of flowering schedules that arise from developmental rather than probabilistic processes. In contrast to probability histograms, flowering schedules can be visualized as a time series with number or proportion of flowers on the y-axis. Although the underlying processes are distinct, the equations are the same (Table 1). To understand the biological significance of mean, variance, skew and kurtosis, we contrast flowering schedules of different shape (Fig. 1). Standardizing to proportion of open flowers (P_t) over time (t), rather than total flower number (N_t) accounts for variation in total flower number per plant. Moreover, we can use proportions to calculate a weighted mean day of flowering. By analogy to the mean of a probability distribution, the weighted mean of a flowering schedule represents the 'balance point', which better captures the start of anthesis of a typical flower.

　　As described in Table 1 and visualized in Figure 1, central moments capture biologically meaningful variation in flowering schedules that may not be correlated with onset or duration of flowering. For example, the variance parameter (σ_i^2) describes how much flowering is spread out, even if the start and end dates are the same. An individual's coefficient of skew (CS_i) accounts for expected differences in symmetry and thus describes whether flowering is concentrated earlier (i.e., right skew, $CS_i > 0$ as shown by the blue dashed line) or later in the schedule (i.e., left skew, $CS_i < 0$ as shown by the purple dotted line), relative to the mean flowering date. Finally, the coefficient of kurtosis (CK_i) describes deviations from the Gaussian expectation, contrasting individuals with a concentrated cluster of open flowers and a few flowers over a longer timeframe (i.e., leptokurtic distribution, $CK_i > 0$ as shown in the gold dashed line) to individuals that spread out flowering more evenly over the entire schedule (i.e., platykurtic, $CK_i < 0$ as shown by the dashed green line).

Table 1. Equations of average flowering schedule characteristics for a population, with subscripts i for individual and t for census time point.

Summary Statistic	Calculation
Proportion (P) of flowers (F) produced at time t	$$P_{i,t} = \frac{F_{i,t}}{\sum_t F_{i,t}}$$
Day of first flower	$$\alpha_i = \min\{t \mid F_{i,t} > 0\}$$
Day of last flower	$$\Omega_i = \max\{t \mid F_{i,t} > 0\}$$
Duration of flowering	$$\Omega_i - \alpha_i$$
Day of mean flowering	$$\mu_i = \sum_t t \, P_{i,t}$$
Flowering schedule variance	$$\sigma_i^2 = \sum_t (t - \mu)^2 \, P_t$$
Flowering schedule coefficient of skew	$$CS_i = \frac{\sum_t (t - \mu_i)^3 \, P_{i,t}}{\sigma_i^3}$$
Flowering schedule coefficient of kurtosis	$$CK_i = \frac{\sum_t (t - \mu_i)^4 \, P_{i,t}}{\sigma_i^4} - 3$$

FIGURE LEGENDS

Figure 1. Hypothetical flowering schedules of five individuals with different characteristics, as described in Box 1. Flowering schedules are scaled to the proportion of total flower production per plant (P_t) over time (t).

Figure 2. Three hypothetical flowering schedules demonstrating predicted clines in central moments from northern (blue dotted line) to southern (red dashed line) genotypes, relative to a standard normal schedule at mid-latitude (solid pink line). Each schedule is scaled as the proportion of total flowers (P_t) over time (t). The northern schedule is defined by an earlier mean, lower variance and positive skew. The southern schedule is defined by a later mean, higher variance, negative skew, and higher kurtosis.

Figure 3. Matrices showing pairwise Pearson correlation coefficients among the start, duration, central moments (i.e., variance, skew and kurtosis), using either: (A) flowering schedule characteristics averaged across individuals within each population ($n = 13$), or (B) the residual correlations calculated on deviations of individuals from their population means ($n = 369$). Circles are shown for statistically significant correlations ($p < 0.05$), colour-coded by correlation coefficient (i.e., -1 to 1 as shown on the x-axis).

Figure 4. Principal Coordinates Analysis (PCoA) of Kolmogorov-Smirnov distances between flowering schedules of 369 plants from 13 populations sampled along a latitudinal gradient and grown in a common garden at the Koffler Scientific Reserve in Newmarket, ON. Point colours correspond to latitude of origin ranging from dark (48° N) to light (38° N). Bar plots show representative flowering schedules of four individual plants. Vector directions correlate to loadings on PCoA1 and PCoA2.

Figure 5. Bivariate plots showing bootstrap means and statistical relations between latitude and average flowering schedule characteristics of 13 populations of *Lythrum salicaria*, grown in a common garden field experiment at mid-latitude (44.03ºN). The bootstrapped average population means (black dots) and quadratic regression lines (red curves) are based on 999 bootstrap iterations. To visualize variation among bootstrap iterations, individual iteration averages (grey dots) and regression lines (grey curves) are also shown for a random sample of 99 bootstrap iterations.

Figure 1. Hypothetical flowering schedules of five individuals with different characteristics, as described in Box 1. Flowering schedules are scaled to the proportion of total flower production per plant (P_t) over time (t).

Figure 2. Three hypothetical flowering schedules demonstrating predicted clines in central moments from northern (blue dotted line) to southern (red dashed line) genotypes, relative to a standard normal schedule at mid-latitude (solid pink line). Each schedule is scaled as the proportion of total flowers (P_t) over time (t). The northern schedule is defined by an earlier mean, lower variance and positive skew. The southern schedule is defined by a later mean, higher variance, negative skew, and higher kurtosis.

Figure 3. Matrices showing pairwise Pearson correlation coefficients among the start, duration, central moments (i.e., variance, skew and kurtosis), using either (A) flowering schedule characteristics averaged across individuals within each population ($n = 13$), or (B) the residual correlations calculated on deviations of individuals from their population means ($n = 369$). Circles are shown for statistically significant correlations ($p < 0.05$), colour-coded by correlation coefficient (i.e., -1 to 1 as shown on the x-axis).

Figure 4. Principal Coordinates Analysis (PCoA) of Kolmogorov-Smirnov distances between flowering schedules of 369 plants from 13 populations sampled along a latitudinal gradient and grown in a common garden at the Koffler Scientific Reserve in Newmarket, ON. Point colours correspond to latitude of origin ranging from dark (48° N) to light (38° N). Bar plots show representative flowering schedules of four individual plants. Vector directions correlate to loadings on PCoA1 and PCoA2.

Figure 5. Bivariate plots showing bootstrap means and statistical relations between latitude and average flowering schedule characteristics of 13 populations of *Lythrum salicaria*, grown in a common garden field experiment at mid-latitude (44.03ºN). The bootstrapped average population means (black dots) and quadratic regression lines (red curves) are based on 999 bootstrap iterations. To visualize variation among bootstrap iterations, individual iteration averages (grey dots) and regression lines (grey curves) are also shown for a random sample of 99 bootstrap iterations.

Appendix S1. The distribution of 369 individuals from 216 seed families representing 13 populations of *Lythrum salicaria* sampled along a latitudinal gradient and grown in a common garden at the Koffler Scientific Reserve in Newmarket, ON.

Population Latitude (°)	*N* Seed families	*N* Individuals
38.75261	15	23
40.34025	15	33
41.31283	18	24
42.23822	20	34
42.52306	12	34
43.69719	19	21
44.49339	19	38
44.59489	19	22
45.26278	16	26
45.43194	17	36
45.49000	16	23
47.69081	14	32
48.47892	16	23