THE DUAL-SPECIFICITY KINASE DYRK1A INTERACTS WITH THE HEPATITIS B VIRUS

GENOME AND REGULATES THE PRODUCTION OF VIRAL RNA

Florentin Pastor[1], Emilie Charles[1], Chiara Di Vona[3], Maëlys Chapelle[1], Michel Rivoire[4],

Guillaume Passot[5], Benoit Chabot[2], , Susana de la Luna[3,6], Julie Lucifora[1], David Durantel[1],

Anna Salvetti[1*]

[1] International Center for Research in Infectiology (CIRI), INSERM U1111, Université Claude

Bernard Lyon 1, CNRS UMR5308, ENS de Lyon, Lyon, France

[2] Department of Microbiology and Infectious Diseases, Faculty of Medicine and Health

Sciences, Université de Sherbrooke, Sherbrooke, Quebec, Canada.

[3] Genome Biology Program, Centre for Genomic Regulation (CRG), and CIBER of Rare

Diseases, Barcelona, Spain.

[4] INSERM U1032, Centre Léon Bérard (CLB), Lyon, France.

[5] Service de chirurgie générale et oncologique, Hôpital Lyon Sud, Hospices Civils de Lyon Et

CICLY, EA3738, université Lyon 1

[6] Institució Catalana de Recerca i Estudis Avançats (ICREA), Barcelona, Spain.

* corresponding author: anna.salvetti@inserm.fr

Keywords: HBV, DYRK1A, cccDNA, transcription, virus-host interaction,

ABSTRACT

The genome of Hepatitis B virus (HBV) persists in infected hepatocytes as a nuclear episome (cccDNA) that is responsible for the transcription of viral genes and viral rebound, following antiviral treatment arrest in chronically infected patients. Beside pegylated interferon-alpha, there is currently no clinically approved therapeutic protocol able to target cccDNA (1). The development of alternative strategies aiming at permanently abrogating HBV RNA production requires a thorough understanding of cccDNA transcriptional and post-transcriptional regulation. In a previous study, we discovered that 1C8, a compound that inhibits the phosphorylation of some cellular RNA-binding proteins (RBPs), could decrease the level of HBV RNAs. Here, we aimed at identifying kinases responsible for this effect. Among the kinases targeted by 1C8, we focused on DYRK1A, a dual-specificity kinase that controls the transcription of cellular genes by phosphorylating transcription factors, histones, chromatin regulators as well as RNA polymerase II. The results of a combination of genetic and chemical approaches using HBV-infected hepatocytes, indicated that DYRK1A positively regulates the production of HBV RNAs. In addition, we found that DYRK1A associates with cccDNA, and stimulates the production of HBV nascent RNAs. Finally, reporter gene assays showed that DYRK1A up-regulates the activity of the HBV enhancer 1/X promoter in a sequence-dependent manner. Altogether, these results strongly suggest that DYRK1A participates in the early onset of the HBV life cycle by stimulating the production of HBx, a viral factor absolutely required to trigger the complete cccDNA transcriptional program.

INTRODUCTION

Chronic infection by Hepatitis B virus (HBV) is due to the stable persistence and activity of the viral genome, called cccDNA (for covalently-closed circular DNA), within the nucleus of infected hepatocytes. Clinically accepted treatments are represented mostly by nucleoside analogs, which target viral reverse-transcription and reduce viremia, but rarely lead to a stable functional cure. Indeed, these compounds do not affect cccDNA stability and transcriptional activity. As a consequence, the continuous production of viral RNAs and proteins allows viral replication to resume off-treatment, and, more importantly, contributes to immune escape mechanisms and liver disease progression (2, 3). Despite some experimental proof-of-concept approaches (1, 4), *in vivo* therapies aiming at eradicating HBV cccDNA and other replication intermediates from infected hepatocytes are unrealistic to date (1). Therefore, strategies directed to permanently silence the viral genome and/or destabilize the viral RNAs, requiring an in-depth understanding of cccDNA transcriptional and post-transcriptional regulation, are currently investigated.

After binding to its main receptor and co-receptors on the hepatocyte membrane, HBV particles are internalized within endosomes from which, the viral nucleocapsids are released later on. Subsequently, these nucleocapsids are directed to the nuclear membrane, translocate through the nuclear pore, and finally disassemble within the inner nuclear basket (5, 6). This process results in the release into the nucleoplasm of the viral relaxed circular DNA genome (rcDNA), with a covalently-attached polymerase, and of 240 copies of the HBc (Core) protein, the unique capsid component. Once in the nucleus, the rcDNA is repaired by cellular factors, chromatinized, and loaded with cellular and viral proteins to form cccDNA, the viral double-stranded DNA episome responsible for the production of viral RNAs (7). This 3.2 Kb viral genome contains 2 enhancers (enhancer 1 and 2) and 4 promoters (preCore/Core, preS1, preS2, and X) that harbor binding sites for several ubiquitous and hepatocyte-specific transcription factors, and drive the transcription of 5 unspliced and several spliced RNAs (8-10). Seven viral proteins are translated from unspliced transcripts and are sufficient to initiate productive replication. Among them, HBx is a 154 amino-acid

protein translated from mRNAs produced from enhancer 1/X promoter with diverse

transcription initiation sites (11). This viral protein is absolutely required to initiate and

maintain cccDNA in a transcriptionally active state (12, 13). The reason for this requirement

remained obscure for a long time until the discovery that, in the absence of HBx, cccDNA is

bound and silenced by the Smc5/6 cohesin complex. The transcriptional repression of

cccDNA is relieved by HBx by hijacking the CUL4A E3 ubiquin ligase to induce the

degradation of the Smc5/6 complex (14-16).

In addition to HBx, the viral HBc protein, translated from the pregenomic RNA (pgRNA)

transcribed from the enhancer 2/preCore/Core promoter, is also suspected of playing

important regulatory roles in the production of viral RNAs (17). Indeed, besides its essential

role to form capsids in the cytoplasm, HBc is also present in the nucleus where, as HBx, it

associates with cccDNA (18, 19). The consequences of this interaction on viral transcription

are, however, still a matter of debate (20-23). The recent observations that capsid-derived

HBc stably associates with the viral genome once released into the nucleus (24), and that it

interacts with Hira, a histone variant deposited on cccDNA during the chromatinization

process (25), additionally suggests that it may be involved during cccDNA repair and/or

chromatinization.

In a previous study, we found that HBc interacts with several cellular RBPs within the

hepatocyte's nucleus raising the hypothesis that it may hijack their activities to control viral

replication (26). In the same study, we discovered that a drug named 1C8, capable of

inhibiting the phosphorylation of some RBPs associated with HBc, could reduce the level of

nascent HBV RNAs (26). These results suggested that one or more protein kinases targeted

by this 1C8 might control the synthesis of viral RNAs by phosphorylating cellular and/or viral

proteins.

The objective of the current study was to identify the kinase(s) targeted by 1C8 responsible

for its (their) effect on cccDNA transcriptional activity. Our results pointed to DYRK1A as one

of these kinases. DYRK1A is a pleiotropic serine/threonine kinase that, among other

activities, regulates the transcription of cellular genes by associating with chromatin and

phosphorylating histones, chromatin regulators as well as the carboxy-terminal domain of the

RNA polymerase II (RNA Pol II) (27-29). DYRK1A also participates in nuclear condensates

associated to transcription elongation (30). We show that DYRK1A interacts with cccDNA

and up-regulates the transcription of viral RNAs. In addition, our data strongly suggest that

DYRK1A transcriptional activity is mediated by the recognition of a consensus motif within

enhancer I/X promoter region that drives HBx production. Therefore, we propose that

DYRK1A is a novel cellular proviral factor that participates in the early onset of HBV

transcription by firing the initial transcription of HBx, a viral factor required to unleash the

complete HBV transcriptional program leading to productive replication.

MATERIALS AND METHODS

Cell culture and infection

HepaRG cells were cultured, differentiated, and infected by HBV as previously described (31). Primary human hepatocytes (PHHs) were freshly prepared from human liver resection obtained from the Centre Léon Bérard and Hôpital Lyon Sud (Lyon) with French ministerial authorizations (AC 2013-1871, DC 2013 – 1870, AFNOR NF 96 900 sept 2011) as previously described (32). To generate HepaRG cells lines over-expressing HA-tagged human DYRK1A and DYRK1B (wild-type and kinase-dead versions (KR), in which the ATP binding Lys179 was replaced by Arg),) and FLAG-tagged SRPK1 (33), the open reading frame of each kinase was inserted into the pLenti4/TO lentiviral vector plasmid, under the control of the minimal CMV/TetOn promoter. Lentiviral particles generated from these plasmids were used to transduce HepaRG-TR cells (13). Transduced cells were selected using blasticidin (10 μg/mL) and zeocyn (100 μg/mL), amplified and frozen as polyclonal lines. HuH7-NTCP cells were cultivated in 10% fetal calf serum supplemented DMEM (4.5 g/L glucose) (34). HBV genotype D inoculum (ayw subtype) was prepared from HepAD38 (35) cell supernatant by polyethylene-glycol-MW-8000 (PEG8000, Sigma) precipitation (8% final) as previously described (36). The titer of endotoxin free viral stocks was determined by qPCR. Cells were infected overnight in a media supplemented with 4% final of PEG, at indicated multiplicity of infections (MOIs), as previously described (31). Measure of secreted HBs and HBe antigens was performed by chemo-luminescent immune assay (AutoBio, China), following manufacturer's instructions and expressed as international units/mL (IU/mL) and Paul Erlich Internation units/mL (PEIU/mL), respectively.

Chemical compounds

Unless otherwise specified, chemical reagents, drugs, antibiotics were purchased from Sigma Aldrich. Tenofovir (TFV) was a kind gift of Gilead Sciences (Foster city, USA). The core assembly modulator (CAM) used in the experiments was previously described (37), and resynthesized by AI-Biopharma (Montpellier, France). 1C8, *i.e.* 1-Methyl-N-(5-

nitrobenzo[d]isothiazol-3-yl)-4-oxo-1,4-dihydropyridine-3-carboxamide, was synthesized and

purified at 99% by AGV (Montpellier, France). RG7834 (38), a molecule that destabilizes

HBV RNAs, was synthetized by AI-Biopharma (Montpellier, France). AZ191 was purchased

from SelleckChem, IFN-α (Roferon) from Roche, SRPIN340 and TG003 from MedChem

Express. DB18 was a kind gift from René Grée (University of Rennes, France) ((39).

siRNA Transfection

Differentiated HepaRG (dHepaRG) cells or PHHs seeded into a 24-well plate were

transfected with 25 nM of siRNA using Lipofectamine RNAiMax (Life Technologies), following

manufacturer's instructions: siDYRK1A (Dharmacon SmartPool L-004805-00), siDYRK1B

(Dharmacon SmartPool L-004806-00), siCLK2 (Dharmacon J-004801-10), siSRPK1

(Dharmacon SmartPool: L-003982-00) and siControl (siCTL) (Dharmacon D-001810).

Viability/cytoxicity assays.

Neutral red uptake and sulforhodamine B assays were performed to estimate cell

viability/cytotoxicity as previously described (40). Puromycin (Puro), was used as a positive

control at a concentration of 10 µM.

Nucleic acid extraction and analysis

HBV RNAs and DNA were extracted from cells with the Nucleospin RNA (Macherey-Nagel)

and MasterPureTM Complexe Purification Kit (BioSearch Technologies) kit without digestion

with proteinase K (41), respectively, according to the manufacturer's instruction. RNA

reverse transcription was performed using Maxima First Strand cDNA Synthesis kit

(Thermofisher). Quantitative PCR for HBV were performed using HBV specific primers and

normalized to PRNP or RPLP0 housekeeping genes as previously described (42).

Pregenomic RNA (pgRNA) was quantified using the TaqMan Fast Advanced Master Mix (Life

Technologies) and normalized to GUS B cDNA levels. HBV cccDNA was quantified from

total DNA by TaqMan qPCR analyses and normalized to β-globin DNA level, as previously

described (43). Unless otherwise stated, results are expressed as the mean normalized ratio

+/- SD, of 3 independent experiments, each performed in triplicate.

Western blot analyses

Proteins were resolved by SDS-PAGE and then transferred onto a nitrocellulose membrane.

Membranes were incubated with the primary antibodies corresponding to the indicated

proteins. Proteins were revealed by chemiluminescence (Super Signal West Dura Substrate,

Pierce) using a secondary peroxydase-conjugated antibody (Dako) at a dilution of 1:10000.

Primary antibodies used were CLK1 (Clinisciences, ARP52021, 1:1000), CLK2 (Abcam,

Ab65082, 1:1000), SRPK1 (Abcam, Ab131160, 1:1000), DYRK1A (Abcam Ab69811, 1:1000

and SantaCruz sc-100376, 1:1000), DYRK1B (Cell Signaling 5672, 1:1000), HBc (provided

by A. Zlotnick (Indiana University, USA), 1:40000), β-tubulin (Abcam 6046,1:10000), β-actin

(Sigma A2228, 1:2000), cyclin D1 (Cell Signaling 55506, 1:2000), HA (Abcam 130275;

1:1000), and FLAG (Sigma F1804; 1:1000).

Native cccDNA-immunoprecipitation (cccDNA-IP) and ChIP

179 cccDNA was immunoprecipitated from frozen cell pellets following a modified native

180 chromatin-immuno-precipitation (ChIP) protocol as previously described (24). Antibodies (2

181 μg) used were: DYRK1A (Abcam 69811 and Sigma D1694), HBc (Life Invitrogen MA1-7609),

182 total RNA Pol II (Diagenode C15200004), RNA Pol II phosphor-Ser5 (Abcam Ab5131), RNA

183 Pol II phosphor-Ser2 (Abcam Ab5095) and control IgG (Cell Signaling 2729).

Luciferase assays

186 Huh7.NTCP cells seeded in a 48-well plates were transfected using TransIT-LT1 transfection

187 reagent (Mirus) and lysed 72 hours later using the Bright-Glo Luciferase reagent (Promega),

188 according to the manufacturer's instructions. Luciferase activity was quantified on a

189 Luminoskan (Life Technologies). Plasmids expressing luciferase under the control of HBV

promoters were purchased from Addgene: preC/C_Luc (pHBV_Luc; ref:71414); pX_Luc (pHBV-X/EnhI-Luc; ref:71418). Mutations M1 (<u>CAGGTG</u>cAac) and M2 (<u>CAGGTGTACA</u>) in pX_Luc were introduced using Q5 site-directed mutagenesis kit (New England Biolabs). Plasmids expressing DYRK1A or DYRK1A_KR under the control of the CMV promoter were previously described (29).

Run-ON

HBV nascent RNAs were quantified as previously described (44, 45). Briefly, cells were incubated for 2h at 37°C with 5-bromouridine (BrU, Sigma), before RNA extraction and immunoprecipitation using an anti-BrdU antibody (BD Pharmingen) and Dynabeads coated with anti-mouse IgG (Life Technologies), overnight at 4°C. Captured RNAs were then eluted, concentrated and quantified by RT-qPCR. Total RNAs, extracted before IP were similarly quantified by RT-qPCR. Controls were provided by cells treated with actinomycin D (Invitrogen, 1 μM; 1 h before labeling), RG7834 (0.05 μM, added twice 5 and 3 days before labeling) or cells incubated in the absence of BrU.

Statistical analysis

Statistical analyses were performed using the GraphPad Prism 9 software and a two-tailed Mann-Whitney non-parametric tests. In graph bars, error bars represent standard deviation (SD) unless otherwise indicated. A p value ≤ 0.05 was considered as significant: *, p value ≤ 0.05; **, p value ≤ 0.01; ***, p value ≤ 0.001.

RESULTS

1C8 targets five CMGC kinases expressed in human hepatocytes

Our previous results indicated that the compound 1C8, an inhibitor of the phosphorylation of some cellular RBPs (46, 47), downregulated the synthesis of HBV RNAs without affecting cccDNA levels (26). This observation suggested that one or more kinases involved in RBPs phosphorylation had a proviral function on HBV transcription. A kinase profiler assay conducted using 1C8 (Eurofins Pharma Discovery) identified 18 different kinases highly inhibited by the compound (Figure S1) (48). Among them, the members of the Cdc2-like kinases (CLK) family CLK1, CLK2 and CLK4, and those of the dual-specificity tyrosine phosphorylation-regulated kinases (DYRKs) family DYRK1A and DYRK1B, were found. All these kinases that belong to the CMGC group of the eukaryotic kinome (49, 50), phosphorylate RBPs, in particular SR proteins, and their inhibition has been proposed to have therapeutic potential (51, 52). Interestingly, the serine-arginine protein kinases (SPRKs) family that also belongs to the CMGC group and are well-known regulators of SR proteins, were not found as 1C8 targets.

One first requirement for the involvement of these kinases in HBV life cycle is their expression in hepatocytes. Thus, we analyzed the level of expression of these kinases in the human hepatocyte cell line HepaRG, using either proliferating or differentiated (dHepaRG) cells, and in primary human hepatocytes (PHHs), whether infected or not with HBV. We detected the expression of CLK1 and CLK2 mRNAs (CLK4 was not measured as it is highly similar to CLK1) in HepaRG cells, irrespectively of their differentiation or infectious status. In comparison, the mRNA levels of DYRK1A and DYRK1B were much lower (Figure S2A). Similar results were observed in PHHs and in human liver biopsies (Figure S2B and C). Moreover, the four kinases were detected at the protein level in both cell types (Figure 1). We conclude that the four kinases targeted by 1C8 are expressed in human hepatocytes.

Inhibitors targeting DYRK1A/B, but not CLK kinases, reduce HBV replication

In order to identify which CLK/DYRK kinase(s) targeted by 1C8 potentially regulates HBV

replication, the effect of inhibitors of these kinases was analyzed on the production of HBV

RNAs and the secretion of HBs and HBe antigens in HBV-infected PHHs, two traits

commonly used to measure HBV productive infection (Figure 2). As positive controls, we

used IFN-α, an inhibitor of HBV transcription (53), which reduces HBV RNAs, and a class I

CAM, which prevents capsid assembly (54), and also reduces HBe secretion (Figure 2B)

(37). Treatment of cells with either SRPIN340 (55), an inhibitor of the kinase SRPK1, not

targeted by 1C8, or with TG003, a well-known CLK inhibitor (56), had no effect on HBV RNA

production or HBs and HBe secretion (Figure 2B).(55) The absence of effect of inhibitors

targeting CLKs was further confirmed using DB18 (Figure S3), a compound that selectively

targets CLK1/4 and CLK2 but not DYRK1A (39). In contrast, both 1C8 and AZ191, a DYRK1

kinases inhibitor but with no effect on CLKs (57), induced a decrease in HBV RNA levels

(Figure 2B and S3). The effect of AZ191 was stronger than that of 1C8 (Figure 2B), with no

evidence of cell toxicity at the dose of 10-20 µM used (Figure S4). The lower effect of 1C8

could be due to the poor solubility of 1C8 and/or to a stronger susceptibility to the hepatocyte

detoxifying activity. Altogether, these results pointed towards DYRK1A/1B as potential

kinases able to directly or indirectly regulate HBV replication.

256   Production of HBV RNAs responds to DYRK1A/DYRK1B levels

To provide further support to the results with the kinase inhibitors, we set up knock-down

(KD) experiments in HBV-infected dHepaRG cells, using a well-established protocol in our

lab, consisting in two successive siRNAs transfections followed by downstream analyses 10

days post-transfection (dpt) (Figure S5A). Unfortunately, despite several attempts using

different sources of siRNA, efficient KD of CLK1 mRNA was never achieved in hepatocytes,

precluding further analysis with this kinase. It is possible that a previously described stress-

responsive mechanism, able to rapidly compensate for the loss of CLK1 (58), may be the

cause of this failure (58). In contrast, CLK2 and SRPK1 expression was reduced in HBV-

infected dHepaRG cells upon siRNA treatment, but no effect on HBV RNAs was observed

(Figure S5B and C). Transfection of siRNAs targeting either DYRK1A or DYRK1B resulted in

the visible death of most cells between 7- and 10-dpt. This cytopathic effect led us to modify

the depletion protocol to investigate effects at earlier time-points (Figure S6A), during which

no toxic effects were observed (Figure S6B and C), but with KD efficiency gradually

increasing (Figure S6D). In these conditions, total viral RNA levels remained unaffected

when targeting DYRK1B, but they were clearly reduced when targeting DYRK1A (Figure

S6E). Importantly, DYRK1A KD induced not only a significant decrease in total HBV RNAs,

but also in pgRNA and downstream parameters (HBe and HBs secretion), without affecting

cccDNA levels (Figure 3).

To confirm this result, the effect of DYRK1A KD was also assessed in HBV-infected PHHs, in

which the effect of DYRK1A KD could be evaluated at longer times after transfection due to

the lack of toxicity of the siRNA treatment with good depletion efficiency (Figure S7A and B).

Surprisingly, discordant results were obtained using this cell type. For 3 out of 7 total

experiments, each performed with PHHs from independent donors, KD of DYRK1A resulted

in a decrease of viral RNAs without affecting cccDNA levels (Figure S7C and D), in

agreement with the results in dHepaRG cells. However, in four of these experiments, the

opposite result was observed despite similar levels of DYRK1A KD (Figure S7E and F). One

possibility for these discordant results could be related to compensatory effects between

DYRK1A and its close paralog DYRK1B (49, 50), effects that might have variable penetrance

given that PHHs are from different donors. In fact, an increase in DYRK1B levels in response

to silencing DYRK1A has been shown in a different physiological context (59). To test this

possibility, we quantified the levels of DYRK1A and DYRK1B mRNAs in control and KD cells

by RT-qPCR. The analyses revealed that DYRK1B mRNA was up-regulated in PHHs

transfected with siRNA against DYRK1A. In addition, the up-regulation of DYRK1B mRNA

was significantly higher in PHHs where HBV RNAs increased following DYRK1A KD,

compared to PHHs where the opposite effect was observed (Figure S8A), suggesting a

possible positive correlation between the level of DYRK1B compensation and the effect of

DYRK1A KD on HBV RNAs. The increase in DYRK1B upon DYRK1A KD was also observed

in dHepaRG cells but at a relatively lower level compared to PHHs, despite a similar

efficiency in DYRK1A KD (Figure S8B), highlighting a layer of complexity linked to the cell

line used. In contrast, DYRK1A mRNA levels were not upregulated upon DYRK1B KD in

either dHepaRG cells or PHH (Figure S8B and S9B). Importantly, the variations in DYRK1B

mRNA were confirmed at the protein level (Figure S9A). Interestingly, in those PHHs where

DYRK1B compensation upon DYRK1A KD was higher, DYRK1B KD, either alone or in

combination with DYRK1A, induced a decrease in HBV RNA (Figure S9C). Altogether, these

results indicated a regulatory cross-talk between DYRK1B and DYRK1A whereby DYRK1A

depletion is compensated by an increase in DYRK1B mRNA, either due to the existence of

DYRK1A-responsive elements within the DYRK1B promoter region or to DYRK1A-

dependent cellular events that impact on DYRK1B transcription and/or mRNA stability. More

importantly, the level of DYRK1B compensation provided a rational explanation for the

divergent results observed upon DYRK1A KD in PHHs.

Next, we wondered whether the observed effect on HBV transcription depends on DYRK1A

dosage. To this end, HepaRG cells over-expressing either a wild-type (wt) DYRK1A or a

kinase-dead version (DYRK1A_KR), under the control of a tetracyclin (Tet)-inducible

promoter were developed (Figure 4A and B). Notably, over-expression of DYRK1A_WT, but

not of SRPK1, led to an increase in HBV parameters, either intracellular viral RNAs or

extracellular HBs and HBe antigens (Figure 4C-F). This effect was partially dependent on

DYRK1A catalytic activity, as overexpression of the kinase-dead version induced a much

lower increase (Figure 4C-F), further supporting the kinase activity dependence showed by

the experiments with the inhibitors. Interestingly, overexpression of DYRK1B induced only a

modest increase in HBV pgRNA but not in total RNAs, indicating a lower capacity to

modulate viral RNAs compared to DYRK1A (Figure S10).

Altogether, the loss-of-function and gain-of-function studies indicate that DYRK1A could

exert a proviral function by increasing the levels of HBV RNAs, and that its catalytic activity is

involved in this function. They also highlight a complex cell-specific compensatory

mechanism between DYRK1A and DYRK1B expression, and, at least partially, redundant

functions on HBV replication.

323 DYRK1A associates to HBV cccDNA and up-regulates transcriptional activity from

324 enhancer 1/HBx promoter region

Previous studies indicated that DYRK1A promotes the transcription of a set of cellular genes

by associating to chromatin at RNA Pol II promoter regions and close to transcriptional start

sites (TSS) (29). We therefore investigated whether DYRK1A could associate to HBV

cccDNA using our recently optimized native ChIP protocol (24). This protocol is efficient in

detecting the presence of HBc bound to cccDNA (Figure 5A), and showed that DYRK1A was

associated to cccDNA in HBV-infected dHepaRG cells (Figure 5A). The same result was

obtained in HBV-infected PHHs with two different antibodies to DYRK1A (Figure 5B and C).

These results indicated that DYRK1A interacts with cccDNA, either directly or indirectly via

other cellular and/or viral factors.

DYRK1A recruitment to chromatin is mediated by a 10 nucleotides sequence

TCTCGCG(A/G)(G/T)(A/G) located near TSSs (29, 60) (Figure 6A). Association of DYRK1A

to such sites regulates the transcription of downstream genes by phosphorylating the C-

terminal domain (CTD) of RNA Pol II (29). Interestingly, a sequence partially matching the

DYRK1A motif is present in the enhancer 1/X promoter region of the HBV genome, upstream

of the HBx TSSs (Figure 6A). To investigate whether the enhancer 1/X promoter could be a

target of DYRK1A, reporter assays were conducted with constructs containing the luciferase

cDNA under the control of either the enhancer 1/X promoter (pX_Luc) regions or enhancer

2/preCore/Core (preC/C_Luc) as a negative control, co-transfected together with constructs

expressing either DYRK1A wt or DYRK1A-KR. As shown in Figure 6B, DYRK1A expression

significantly increased the activity of the enhancer 1/X promoter reporter, but not that of the

enhancer 2/preCore/Core reporter. Interestingly, a lower but still significant enhancement

was observed following transection of the plasmid expressing the DYRK1A kinase-dead

version. In this regard, it is worth noting that a transcriptional effect independent of the

DYRK1A kinase activity was observed in one-hybrid assays (29), suggesting a possible

scaffolding activity. To further determine if the DYRK1A transactivating activity was mediated by the putative DYRK1A motif within enhancer 1/X promoter, two different mutant motifs were generated (Figure 6A). Both pX_M1_Luc and pX_M2_Luc constructs no longer responded to DYRK1A (Figure 6C), indicating that the enhancing activity of DYRK1A did depend on the presence of the motif characteristic of its recruitment to cellular chromatin. Altogether, these results indicated that DYRK1A associates to cccDNA and strongly suggested that its effect on HBV RNAs was, at least in part, linked to its capacity to recognize a sequence matching the DYRK1A motif present on the enhancer 1/HBx promoter region.

Over-expression of DYRK1A up-regulates the transcription of HBV RNAs

To investigate whether DYRK1A acts at the transcriptional level, RUN-On assays were performed. For this, BrU-labeled RNAs were extracted from HBV-infected dHepaRG cells, with DYRK1A (WT or KR versions) expression induced by Tet treatment. To proof that the experimental set up does detect nascent RNAs, we show that treatment of cells with RG7834, a compound that degrades HBV RNAs (38, 44), had no effect on nascent RNAs, whereas actinomycin D, an inhibitor of RNA Pol II, blocked their synthesis (Figure 7). As shown above, over-expression of DYRK1A wt increased total HBV RNAs, whereas its kinase inactive version had no effect (Figure 7, left panel). Importantly, up-regulation of nascent HBV RNAs was observed only when the expression of active DYRK1A was induced (Figure 7, right panel), strongly suggesting that the DYRK1A-dependent effect takes place at the transcriptional level.

DISCUSSION

So far, only few cellular kinases were reported to modulate the HBV life cycle. In addition, most if not all of them, exert their effects at late steps of the infectious cycle, during capsid assembly and pgRNA packaging (17). This is the case of CDK2, PKC and PLK1, which phosphorylate the HBc protein during capsid assembly in the cytoplasm (61-64). In this study, we asked whether cellular kinases could also regulate the HBV life cycle by impacting on the production of HBV RNAs. This question arose from the observation that 1C8, a compound able to dephosphorylate some RBPs, including SR proteins involved in the HBV life cycle, could down-regulate the production of nascent HBV RNAs (26, 47).

Among the kinases identified as the main 1C8 targets *in vitro,* we focused on DYRK1A. This kinase belongs to the DYRK family, within the CMGC group of kinases, a family of constitutively active serine/threonine kinases that participate in a wide variety of processes involved in cell growth, differentiation and transcription (49, 50). DYRK1A is an essential gene in mammals, whose alterations in expression have been associated to different human diseases including Down syndrome when present in trisomy or a neurodevelopmental syndrome when in haploinsufficency (65). It is a pleiotropic kinase regulating a variety of cellular processes including transcription and splicing. DYRK1B is the closest paralog of DYRK1A within the family, sharing a high degree of homology at the primary and secondary structure, with activities connected mostly to cancer development (66). DYRK1A and DYRK1B have cytoplasmic and nuclear localization but, only DYRK1A localizes to nuclear speckles, the major storage sites of RBPs, and its over-expression induces speckle disassembly (33). Not surprisingly, DYRK1A can phosphorylate RBPs, among which several SR proteins (52).

Our results show that the genetic manipulation of DYRK1A levels in HBV-infected hepatocytes modulates the production of HBV RNAs without affecting cccDNA levels, suggesting a positive, stimulatory role. This role is less evident for DYRK1B, though our results indicate that the cellular context might be relevant for this kinase. In fact, we found that depletion of DYRK1A induces a compensatory effect on DYRK1B levels, which could

partially compensate for DYRK1A loss, thereby affecting the final impact on HBV RNA

production. The kinase activity of DYRK1A is involved in the process based on the

concurring results of the inhibitory role of small molecule inhibitors and the lack of activity of

a kinase-dead version of DYRK1A on HBV parameters.

Previous studies have shown that DYRK1A is recruited to the proximal promoter of several

cellular genes characterized by the presence of a common 10 nucleotides motif (29). We

found that DYRK1A associates to the viral genome, cccDNA, in infected hepatocytes.

Recruitment to cccDNA could be dependent on the presence of a DYRK1A motif within the

enhancer 1/X promoter as shown by the results with reporter assays mutated for the

consensus sequence. This potential binding site is adjacent to a previously described

interferon-stimulated response element (ISRE) (53), opening the possibility of a functional

cross-talk between the activity of DYRK1A and the interferon pathway. The alignment of

more than ten thousand HBV sequences derived from all genotypes

(https://hbvdb.lyon.inserm.fr/HBVdb/) indicates that this site is conserved among all

genotypes, and further support its role as a general regulatory element. These findings do no

exclude that DYRK1A may associate to cccDNA via other target sequences. Additional

techniques to precisely identify the DYRK1A association site within the cccDNA, such as the

recently developed Cut&Run method (67), could confirm our finding. Additionally, targeted

mutations in cccDNA should also be performed to determine whether the effect of DYRK1A

on HBV life cycle is dependent on its association to the viral genome.

The results showed a good correlation between the activity of DYRK1A and the production of

nascent HBV RNAs or the stimulation of reporters containing the DYRK1A motif, supporting

a role for the enzymatic activity of the kinase. However, we detected a small stimulatory

activity independent of the kinase activity, which has been also observed previously (29).

This could be due to the capacity of DYRK1A to function as a scaffold protein to recruit other

factors involved in the processes.

In any case, the Run-On experiments indicated that DYRK1A stimulates HBV transcription.

We envision several scenarios that are not mutually exclusive. First, DYRK1A may regulate

the phosphorylation status of the RNA Pol II CTD on cccDNA, thereby contributing to

increase transcription initiation and/or elongation. Second, transcriptional regulation by

DYRK1A could result from its interaction with chromatin associated factors shown to be

involved in cccDNA transcriptional regulation, such as the histone acetyl transferase

p300/CBP or the transcriptional repressor HP1 (12, 27, 28, 68, 69). Third, DYRK1A may

exert its effect on HBV by targeting some cellular RBPs, as it does phosphorylate several

splicing-related proteins. Finally, the DYRK1A effect might require some viral factors. In this

regard, HBc preferentially associates to CpG island II within cccDNA, a region which partially

covers enhancer 1/X promoter, an association that correlates with a permissive epigenetic

state (20, 21). Future experiments should investigate whether HBc and DYRK1A can interact

in infected hepatocytes as well as consequences of this interaction on HBc phosphorylation

and association to viral DNA and RNAs.

A major question raised this study concerns the role of DYRK1A during the HBV life cycle.

We are aware that the effect of DYRK1A on the level of HBV RNAs was moderate (always

lower than 2-fold), arguing against a major function during the productive viral cycle.

However, this level of enhancement may be important for the onset of HBV transcription

once the cccDNA is formed in the nucleus, in particular to fire transcription of HBx mRNAs,

an event that is required for the initiation of the complete HBV transcriptional program (12,

13). Indeed, since the HBx protein is not present within the viral capsids, some basal level of

HBx mRNA synthesis has to occur, early after infection, in order to induce Smc5/6

degradation and de-repress the HBV genome (14-16). While some studies have reported the

presence of HBx mRNA within secreted particles, no evidence has been provided so far to

show that these mRNA molecules are transferred into cells and produce HBx (11, 70, 71).

Therefore, it is possible that a low and transient stimulation of HBx mRNA transcription early

upon cccDNA formation may lead to the synthesis of a basal level of HBx sufficient to induce

Smc5/6 degradation (72). In this regard, it is interesting to note that in a recent phospho-

proteomic analysis we found that DYRK1A was among the top list of kinases whose activity

is predicted to be up-regulated upon HBV infection (73).

In conclusion, this study describes for the first time the involvement of DYRK1A, a kinase

which controls the expression of several cellular genes, in the regulation of HBV RNAs

production and strongly suggests that this nuclear kinase may be involved in the early control

of HBV transcription. Further deciphering the exact role of DYRK1A during the HBV life cycle

as well as its molecular targets will be important to further understand the factors which are

critical to unleash the HBV transcriptional program and to develop strategies to counteract it.

ACKNOWLEDGMENTS

We would like to thank Maud Michelet and Anëlle Dubois for their help for PHH isolation, as

well as the staff from Pr. Michel Rivoire's and Dr. Guillaume Passot's surgery rooms for

providing us with liver resections.

FUNDING

This work was funded by Institut National de la Santé et de la Recherche Médicale

(INSERM), the Centre National de la Recherche Scientifique (CNRS), and Université Claude

Bernard Lyon 1 (UCBL). It was also supported by grants from the Ministry of Science and

Innovation (PID2022-139904NB to SdlL], Generalitat de Catalunya (2021SGR01229 and

CERCA Programme to SdlL), Agence Nationale de Recherche sur le Sida et les hépatites

virales (ANRS, ECTZ11892 to AS) and fellowship to FP (ANRS, ECTZ119385). BC is

supported by grant from CIHR.

REFERENCES

1. Lucifora J, Protzer U. 2016. Attacking hepatitis B virus cccDNA--The holy grail to hepatitis B cure. J Hepatol 64:S41-8.

2. Zeisel MB, Lucifora J, Mason WS, Sureau C, Beck J, Levrero M, Kann M, Knolle PA, Benkirane M, Durantel D, Michel ML, Autran B, Cosset FL, Strick-Marchand H, Trepo C, Kao JH, Carrat F, Lacombe K, Schinazi RF, Barre-Sinoussi F, Delfraissy JF, Zoulim F. 2015. Towards an HBV cure: state-of-the-art and unresolved questions--report of the ANRS workshop on HBV cure. Gut 64:1314-26.

3. Fanning GC, Zoulim F, Hou J, Bertoletti A. 2019. Therapeutic strategies for hepatitis B virus infection: towards a cure. Nat Rev Drug Discov doi:10.1038/s41573-019-0037-0.

4. Martinez MG, Smekalova E, Combe E, Gregoire F, Zoulim F, Testoni B. 2022. Gene Editing Technologies to Target HBV cccDNA. Viruses 14.

5. Tsukuda S, Watashi K. 2020. Hepatitis B virus biology and life cycle. Antiviral Res 182:104925.

6. Blondot ML, Bruss V, Kann M. 2016. Intracellular transport and egress of hepatitis B virus. J Hepatol 64:S49-59.

7. Diogo Dias J, Sarica N, Neuveut C. 2021. Early Steps of Hepatitis B Life Cycle: From Capsid Nuclear Import to cccDNA Formation. Viruses 13.

8. Quasdorff M, Protzer U. 2010. Control of hepatitis B virus at the level of transcription. J Viral Hepat 17:527-36.

9. Xia Y, Guo H. 2020. Hepatitis B virus cccDNA: Formation, regulation and therapeutic potential. Antiviral Res 180:104824.

10. Sommer G, Heise T. 2008. Posttranscriptional control of HBV gene expression. Front Biosci 13:5533-47.

11. Stadelmayer B, Diederichs A, Chapus F, Rivoire M, Neveu G, Alam A, Fraisse L, Carter K, Testoni B, Zoulim F. 2020. Full-length 5'RACE identifies all major HBV transcripts in HBV-infected hepatocytes and patient serum. J Hepatol 73:40-51.

12.   Belloni L, Pollicino T, De Nicola F, Guerrieri F, Raffa G, Fanciulli M, Raimondo G,

Levrero M. 2009. Nuclear HBx binds the HBV minichromosome and modifies the

epigenetic regulation of cccDNA function. Proc Natl Acad Sci U S A 106:19975-9.

13.   Lucifora J, Arzberger S, Durantel D, Belloni L, Strubin M, Levrero M, Zoulim F, Hantz

O, Protzer U. 2011. Hepatitis B virus X protein is essential to initiate and maintain

virus replication after infection. J Hepatol 55:996-1003.

14.   Abdul F, Diman A, Baechler B, Ramakrishnan D, Kornyeyev D, Beran RK, Fletcher

SP, Strubin M. 2022. Smc5/6 silences episomal transcription by a three-step function.

Nat Struct Mol Biol 29:922-931.

15.   Decorsiere A, Mueller H, van Breugel PC, Abdul F, Gerossier L, Beran RK, Livingston

CM, Niu C, Fletcher SP, Hantz O, Strubin M. 2016. Hepatitis B virus X protein

identifies the Smc5/6 complex as a host restriction factor. Nature 531:386-9.

16.   Murphy CM, Xu Y, Li F, Nio K, Reszka-Blanco N, Li X, Wu Y, Yu Y, Xiong Y, Su L.

2016. Hepatitis B Virus X Protein Promotes Degradation of SMC5/6 to Enhance HBV

Replication. Cell Rep 16:2846-2854.

17.   Diab A, Foca A, Zoulim F, Durantel D, Andrisani O. 2018. The diverse functions of the

hepatitis B core/capsid protein (HBc) in the viral life cycle: Implications for the

development of HBc-targeting antivirals. Antiviral Res 149:211-220.

18.   Bock CT, Schranz P, Schroder CH, Zentgraf H. 1994. Hepatitis B virus genome is

organized into nucleosomes in the nucleus of the infected cell. Virus Genes 8:215-29.

19.   Bock CT, Schwinn S, Locarnini S, Fyfe J, Manns MP, Trautwein C, Zentgraf H. 2001.

Structural organization of the hepatitis B virus minichromosome. J Mol Biol 307:183-

96.

20.   Chong CK, Cheng CYS, Tsoi SYJ, Huang FY, Liu F, Seto WK, Lai CL, Yuen MF,

Wong DK. 2017. Role of hepatitis B core protein in HBV transcription and recruitment

of histone acetyltransferases to cccDNA minichromosome. Antiviral Res 144:1-7.

21.   Guo YH, Li YN, Zhao JR, Zhang J, Yan Z. 2011. HBc binds to the CpG islands of

HBV cccDNA and promotes an epigenetic permissive state. Epigenetics 6:720-6.

22.   Tu T, Zehnder B, Qu B, Urban S. 2021. D e novo synthesis of hepatitis B virus

532   nucleocapsids is dispensable for the maintenance and transcriptional regulation of

533   cccDNA. JHEP Rep 3:100195.

534   23.   Zhong Y, Wu C, Xu Z, Teng Y, Zhao L, Zhao K, Wang J, Wang W, Zhan Q, Zhu C,

535   Chen X, Liang K, Cheng X, Xia Y. 2022. Hepatitis B Virus Core Protein Is Not

536   Required for Covalently Closed Circular DNA Transcriptional Regulation. J Virol

537   96:e0136222.

538   24.   Lucifora J, Pastor F, Charles E, Pons C, Auclair H, Fusil F, Rivoire M, Cosset FL,

539   Durantel D, Salvetti A. 2021. Evidence for long-term association of virion-delivered

540   HBV core protein with cccDNA independently of viral protein production. JHEP Rep

541   3:100330.

542   25.   Locatelli M, Quivy JP, Chapus F, Michelet M, Fresquet J, Maadadi S, Aberkane AN,

543   Diederichs A, Lucifora J, Rivoire M, Almouzni G, Testoni B, Zoulim F. 2022. HIRA

544   Supports Hepatitis B Virus Minichromosome Establishment and Transcriptional

545   Activity in Infected Hepatocytes. Cell Mol Gastroenterol Hepatol 14:527-551.

546   26.   Chabrolles H, Auclair H, Vegna S, Lahlali T, Pons C, Michelet M, Coute Y, Belmudes

547   L, Chadeuf G, Kim Y, Di Bernardo A, Jalaguier P, Cosset FL, Fusil F, Rivoire M,

548   Arnold LD, Lopatin U, Combet C, Zoulim F, Grierson D, Chabot B, Lucifora J,

549   Durantel D, Salvetti A. 2020. Hepatitis B virus Core protein nuclear interactome

550   identifies SRSF10 as a host RNA-binding protein restricting HBV RNA production.

551   PLoS Pathog 16:e1008593.

552   27.   Li S, Xu C, Fu Y, Lei PJ, Yao Y, Yang W, Zhang Y, Washburn MP, Florens L, Jaiswal

553   M, Wu M, Mohan M. 2018. DYRK1A interacts with histone acetyl transferase p300

554   and CBP and localizes to enhancers. Nucleic Acids Res 46:11202-11213.

555   28.   Jang SM, Azebi S, Soubigou G, Muchardt C. 2014. DYRK1A phoshorylates histone

556   H3 to differentially regulate the binding of HP1 isoforms and antagonize HP1-

557   mediated transcriptional repression. EMBO Rep 15:686-94.

29.   Di Vona C, Bezdan D, Islam AB, Salichs E, Lopez-Bigas N, Ossowski S, de la Luna

S. 2015. Chromatin-wide profiling of DYRK1A reveals a role as a gene-specific RNA

polymerase II CTD kinase. Mol Cell 57:506-20.

30.   Lu H, Yu D, Hansen AS, Ganguly S, Liu R, Heckert A, Darzacq X, Zhou Q. 2018.

Phase-separation mechanism for C-terminal hyperphosphorylation of RNA

polymerase II. Nature 558:318-323.

31.   Gripon P, Rumin S, Urban S, Le Seyec J, Glaise D, Cannie I, Guyomard C, Lucas J,

Trepo C, Guguen-Guillouzo C. 2002. Infection of a human hepatoma cell line by

hepatitis B virus. Proc Natl Acad Sci U S A 99:15655-60.

32.   Lecluyse EL, Alexandre E. 2010. Isolation and culture of primary hepatocytes from

resected human liver tissue. Methods Mol Biol 640:57-82.

33.   Alvarez M, Estivill X, de la Luna S. 2003. DYRK1A accumulates in splicing speckles

through a novel targeting signal and induces speckle disassembly. J Cell Sci

116:3099-107.

34.   Michelet M, Alfaiate D, Chardes B, Pons C, Faure-Dupuy S, Engleitner T, Farhat R,

Riedl T, Legrand AF, Rad R, Rivoire M, Zoulim F, Heikenwalder M, Salvetti A,

Durantel D, Lucifora J. 2022. Inducers of the NF-kappaB pathways impair hepatitis

delta virus replication and strongly decrease progeny infectivity in vitro. JHEP Rep

4:100415.

35.   Ladner SK, Otto MJ, Barker CS, Zaifert K, Wang GH, Guo JT, Seeger C, King RW.

1997. Inducible expression of human hepatitis B virus (HBV) in stably transfected

hepatoblastoma cells: a novel system for screening potential inhibitors of HBV

replication. Antimicrob Agents Chemother 41:1715-20.

36.   Luangsay S, Gruffaz M, Isorce N, Testoni B, Michelet M, Faure-Dupuy S, Maadadi S,

Ait-Goughoulte M, Parent R, Rivoire M, Javanbakht H, Lucifora J, Durantel D, Zoulim

F. 2015. Early inhibition of hepatocyte innate responses by hepatitis B virus. J

Hepatol 63:1314-22.

37.   Lahlali T, Berke JM, Vergauwen K, Foca A, Vandyck K, Pauwels F, Zoulim F,

586   Durantel D. 2018. Novel Potent Capsid Assembly Modulators Regulate Multiple Steps

587   of the Hepatitis B Virus Life Cycle. Antimicrob Agents Chemother 62.

588   38.   Mueller H, Wildum S, Luangsay S, Walther J, Lopez A, Tropberger P, Ottaviani G, Lu

589   W, Parrott NJ, Zhang JD, Schmucki R, Racek T, Hoflack JC, Kueng E, Point F, Zhou

590   X, Steiner G, Lutgehetmann M, Rapp G, Volz T, Dandri M, Yang S, Young JAT,

591   Javanbakht H. 2018. A novel orally available small molecule that inhibits hepatitis B

592   virus expression. J Hepatol 68:412-420.

593   39.   Brahmaiah D, Kanaka Durga Bhavani A, Aparna P, Sampath Kumar N, Solhi H, Le

594   Guevel R, Baratte B, Ruchaud S, Bach S, Singh Jadav S, Raji Reddy C, Roisnel T,

595   Mosset P, Levoin N, Gree R. 2021. Discovery of DB18, a potent inhibitor of CLK

596   kinases with a high selectivity against DYRK1A kinase. Bioorg Med Chem 31:115962.

597   40.   Isorce N, Lucifora J, Zoulim F, Durantel D. 2015. Immune-modulators to combat

598   hepatitis B virus infection: From IFN-alpha to novel investigational immunotherapeutic

599   strategies. Antiviral Res 122:69-81.

600   41.   Allweiss L, Testoni B, Yu M, Lucifora J, Ko C, Qu B, Lutgehetmann M, Guo H, Urban

601   S, Fletcher SP, Protzer U, Levrero M, Zoulim F, Dandri M. 2023. Quantification of the

602   hepatitis B virus cccDNA: evidence-based guidelines for monitoring the key obstacle

603   of HBV cure. Gut doi:10.1136/gutjnl-2022-328380.

604   42.   Lucifora J, Xia Y, Reisinger F, Zhang K, Stadler D, Cheng X, Sprinzl MF,

605   Koppensteiner H, Makowska Z, Volz T, Remouchamps C, Chou WM, Thasler WE,

606   Huser N, Durantel D, Liang TJ, Munk C, Heim MH, Browning JL, Dejardin E, Dandri

607   M, Schindler M, Heikenwalder M, Protzer U. 2014. Specific and nonhepatotoxic

608   degradation of nuclear hepatitis B virus cccDNA. Science 343:1221-8.

609   43.   Werle-Lapostolle B, Bowden S, Locarnini S, Wursthorn K, Petersen J, Lau G, Trepo

610   C, Marcellin P, Goodman Z, Delaney WEt, Xiong S, Brosgart CL, Chen SS, Gibbs

611   CS, Zoulim F. 2004. Persistence of cccDNA during the natural history of chronic

hepatitis B and decline during adefovir dipivoxil therapy. Gastroenterology 126:1750-8.

44. Mueller H, Lopez A, Tropberger P, Wildum S, Schmaler J, Pedersen L, Han X, Wang Y, Ottosen S, Yang S, Young JAT, Javanbakht H. 2019. PAPD5/7 Are Host Factors That Are Required for Hepatitis B Virus RNA Stabilization. Hepatology 69:1398-1411.

45. Paulsen MT, Veloso A, Prasad J, Bedi K, Ljungman EA, Magnuson B, Wilson TE, Ljungman M. 2014. Use of Bru-Seq and BruChase-Seq for genome-wide assessment of the synthesis and stability of RNA. Methods 67:45-54.

46. Cheung PK, Horhant D, Bandy LE, Zamiri M, Rabea SM, Karagiosov SK, Matloobi M, McArthur S, Harrigan PR, Chabot B, Grierson DS. 2016. A Parallel Synthesis Approach to the Identification of Novel Diheteroarylamide-Based Compounds Blocking HIV Replication: Potential Inhibitors of HIV-1 Pre-mRNA Alternative Splicing. J Med Chem 59:1869-79.

47. Shkreta L, Blanchette M, Toutant J, Wilhelm E, Bell B, Story BA, Balachandran A, Cochrane A, Cheung PK, Harrigan PR, Grierson DS, Chabot B. 2017. Modulation of the splicing regulatory function of SRSF10 by a novel compound that impairs HIV-1 replication. Nucleic Acids Res 45:4051-4067.

48. Shkreta L, Toutant J, Delannoy A, Durantel D, Salvetti A, Ehresmann S, Sauvageau M, Delbrouck JA, Gravel-Trudeau A, Comeau C, Huard C, Coulombe-Huntington J, Tyers M, Grierson D, Boudreault PL, Chabot B. 2024. The anticancer potential of the CLK kinases inhibitors 1C8 and GPS167 revealed by their impact on the epithelial-mesenchymal transition and the antiviral immune response. Oncotarget 15:313-325.

49. Aranda S, Laguna A, de la Luna S. 2011. DYRK family of protein kinases: evolutionary relationships, biochemical properties, and functional roles. FASEB J 25:449-62.

50. Lindberg MF, Meijer L. 2021. Dual-Specificity, Tyrosine Phosphorylation-Regulated Kinases (DYRKs) and cdc2-Like Kinases (CLKs) in Human Disease, an Overview. Int J Mol Sci 22.

51.   Wang E, Pineda JMB, Kim WJ, Chen S, Bourcier J, Stahl M, Hogg SJ, Bewersdorf

JP, Han C, Singer ME, Cui D, Erickson CE, Tittley SM, Penson AV, Knorr K, Stanley

RF, Rahman J, Krishnamoorthy G, Fagin JA, Creger E, McMillan E, Mak CC, Jarvis

M, Bossard C, Beaupre DM, Bradley RK, Abdel-Wahab O. 2023. Modulation of RNA

splicing enhances response to BCL2 inhibition in leukemia. Cancer Cell 41:164-180

e8.

52.   Pastor F, Shkreta L, Chabot B, Durantel D, Salvetti A. 2021. Interplay Between

CMGC Kinases Targeting SR Proteins and Viral Replication: Splicing and Beyond.

Front Microbiol 12:658721.

53.   Belloni L, Allweiss L, Guerrieri F, Pediconi N, Volz T, Pollicino T, Petersen J,

Raimondo G, Dandri M, Levrero M. 2012. IFN-alpha inhibits HBV transcription and

replication in cell culture and in humanized mice by targeting the epigenetic regulation

of the nuclear cccDNA minichromosome. J Clin Invest 122:529-37.

54.   Schlicksup CJ, Wang JC, Francis S, Venkatakrishnan B, Turner WW,

VanNieuwenhze M, Zlotnick A. 2018. Hepatitis B virus core protein allosteric

modulators can distort and disrupt intact capsids. Elife 7.

55.   Fukuhara T, Hosoya T, Shimizu S, Sumi K, Oshiro T, Yoshinaka Y, Suzuki M,

Yamamoto N, Herzenberg LA, Herzenberg LA, Hagiwara M. 2006. Utilization of host

SR protein kinases and RNA-splicing machinery during viral replication. Proc Natl

Acad Sci U S A 103:11329-33.

56.   Muraki M, Ohkawara B, Hosoya T, Onogi H, Koizumi J, Koizumi T, Sumi K, Yomoda

J, Murray MV, Kimura H, Furuichi K, Shibuya H, Krainer AR, Suzuki M, Hagiwara M.

2004. Manipulation of alternative splicing by a newly developed inhibitor of Clks. J

Biol Chem 279:24246-54.

57.   Ashford AL, Oxley D, Kettle J, Hudson K, Guichard S, Cook SJ, Lochhead PA. 2014.

A novel DYRK1B inhibitor AZ191 demonstrates that DYRK1B acts independently of

GSK3beta to phosphorylate cyclin D1 at Thr(286), not Thr(288). Biochem J 457:43-

56.

58. Ninomiya K, Kataoka N, Hagiwara M. 2011. Stress-responsive maturation of Clk1/4
pre-mRNAs promotes phosphorylation of SR splicing factor. J Cell Biol 195:27-40.

59. Ackeifi C, Swartz E, Kumar K, Liu H, Chalada S, Karakose E, Scott DK, Garcia-
Ocana A, Sanchez R, DeVita RJ, Stewart AF, Wang P. 2020. Pharmacologic and
genetic approaches define human pancreatic beta cell mitogenic targets of DYRK1A
inhibitors. JCI Insight 5.

60. Di Vona C, Barba L, Ferrari R, de la Luna S. 2023. Loss of the DYRK1A Protein
Kinase Results in the Reduction in Ribosomal Protein Gene Expression, Ribosome
Mass and Reduced Translation. Biomolecules 14.

61. Ludgate L, Ning X, Nguyen DH, Adams C, Mentzer L, Hu J. 2012. Cyclin-dependent
kinase 2 phosphorylates s/t-p sites in the hepadnavirus core protein C-terminal
domain and is incorporated into viral capsids. J Virol 86:12237-50.

62. Luo J, Xi J, Gao L, Hu J. 2020. Role of Hepatitis B virus capsid phosphorylation in
nucleocapsid disassembly and covalently closed circular DNA formation. PLoS
Pathog 16:e1008459.

63. Diab A, Foca A, Fusil F, Lahlali T, Jalaguier P, Amirache F, N'Guyen L, Isorce N,
Cosset FL, Zoulim F, Andrisani O, Durantel D. 2017. Polo-like-kinase 1 is a proviral
host factor for hepatitis B virus replication. Hepatology 66:1750-1765.

64. Wittkop L, Schwarz A, Cassany A, Grun-Bernhard S, Delaleau M, Rabe B, Cazenave
C, Gerlich W, Glebe D, Kann M. 2010. Inhibition of protein kinase C phosphorylation
of hepatitis B virus capsids inhibits virion formation and causes intracellular capsid
accumulation. Cell Microbiol 12:962-75.

65. Deboever E, Fistrovich A, Hulme C, Dunckley T. 2022. The Omnipresence of
DYRK1A in Human Diseases. Int J Mol Sci 23.

66. Becker W. 2018. A wake-up call to quiescent cancer cells - potential use of DYRK1B
inhibitors in cancer therapy. FEBS J 285:1203-1211.

67. Meers MP, Bryson TD, Henikoff JG, Henikoff S. 2019. Improved CUT&RUN
chromatin profiling tools. Elife 8.

696 68. Cougot D, Wu Y, Cairo S, Caramel J, Renard CA, Levy L, Buendia MA, Neuveut C.
697   2007. The hepatitis B virus X protein functionally interacts with CREB-binding
698   protein/p300 in the regulation of CREB-mediated transcription. J Biol Chem
699   282:4277-4287.

700 69. Riviere L, Gerossier L, Ducroux A, Dion S, Deng Q, Michel ML, Buendia MA, Hantz
701   O, Neuveut C. 2015. HBx relieves chromatin-mediated transcriptional repression of
702   hepatitis B viral cccDNA involving SETDB1 histone methyltransferase. J Hepatol
703   63:1093-102.

704 70. Bai L, Zhang X, Kozlowski M, Li W, Wu M, Liu J, Chen L, Zhang J, Huang Y, Yuan Z.
705   2018. Extracellular Hepatitis B Virus RNAs Are Heterogeneous in Length and
706   Circulate as Capsid-Antibody Complexes in Addition to Virions in Chronic Hepatitis B
707   Patients. J Virol 92.

708 71. Niu C, Livingston CM, Li L, Beran RK, Daffis S, Ramakrishnan D, Burdette D, Peiser
709   L, Salas E, Ramos H, Yu M, Cheng G, Strubin M, Delaney WI, Fletcher SP. 2017.
710   The Smc5/6 Complex Restricts HBV when Localized to ND10 without Inducing an
711   Innate Immune Response and Is Counteracted by the HBV X Protein Shortly after
712   Infection. PLoS One 12:e0169648.

713 72. Doitsh G, Shaul Y. 2004. Enhancer I predominance in hepatitis B virus gene
714   expression. Mol Cell Biol 24:1799-808.

715 73. Pastor F, Charles E, Belmudes L, Chabrolles H, Cescato M, Rivoire M, Burger T,
716   Passot G, Durantel D, Lucifora J, Couté Y, Salvetti A. 2024. Deciphering the
717   phospho-signature induced by hepatitis B virus in primary human hepatocytes.
718   bioRxiv doi:https://doi.org/10.1101/2024.04.10.588822 2024.04.10.588822.

719 74. Yoo S, Wang W, Wang Q, Fiel MI, Lee E, Hiotis SP, Zhu J. 2017. A pilot systematic
720   genomic comparison of recurrence risks of hepatitis B virus-associated hepatocellular
721   carcinoma with low- and high-degree liver fibrosis. BMC Med 15:214.

722   75.   Chen JY, Lin JR, Tsai FC, Meyer T. 2013. Dosage of Dyrk1a shifts cells within a p21-

723   cyclin D1 signaling map to control the decision to enter the cell cycle. Mol Cell 52:87-

724   100.

FIGURE CAPTIONS

Figure 1. Expression levels of selected CMGC kinases in human hepatocytes. Western

blot analysis of CLK1, CLK2, SRPK1, DYRK1A, DYRK1B expression in human hepatocytes

(HepaRG cells or PHHs), mock- or HBV-infected. HepaRG: non-differentiated cells;

dHepaRG: differentiated cells; dHepaRG_HBV: HBV-infected dHepaRG; PHH: mock-

infected cells; PHH-HBV: HBV-infected cells. #1 and #2 refers to PHH purified from two

different donors. The detection of HBc is used as a control for HBV infection. β-Actin is used

as loading control.

Figure 2. Chemical inhibitors targeting DYRK1A and DYRK1B reduce HBV RNAs

production. A. Experimental outline. PHHs derived from different donors were infected with

HBV (MOI: 500 vge/cell) and seven days later, treated three times with the indicated

compounds. Viral parameters were measured 2 days after the last treatment. IFN: Interferon-

α (500 IU/mL), CAM: capsid assembly modulator (class I, 10 µM), 1C8 TG003, AZ191, and

SRPIN340 were used at 20 µM. **B.** Relative quantification of total intracellular viral RNAs,

and of secreted HBs and HBe antigen levels, as indicated. Statistical values are from

comparisons between the treated and mock-treated cells for each of the measurements and

expressed as the mean +/- SD of 3 independent experiments, each performed in triplicate.

.

Figure 3. Knock-down of DYRK1A decreases HBV RNAs levels in HBV-infected

dHepaRG cells. A. Experimental outline. Cells were infected and 7 days later transfected

with siRNAs targeting DYRK1A. DYRK1A protein levels (**B**) and the viral parameters

indicated (**C**) were quantified 4-dpt, a time point at which cell toxicity was not observed

(Figure S6). Statistical values are from comparisons between siDYRK1A and siCTL for each

of the measurements and expressed as the mean +/- SD of 3 independent experiments,

each performed in triplicate.

Figure 4. Over-expression of DYRK1A increases HBV RNA levels in HBV-infected dHepaRG cells. A. Experimental outline. dHepaRG cells engineered to express either HA-DYRK1A, HA-DYRK1A_KR or FLAG-SRPK1, under the control of a tetracyclin (Tet)-inducible promoter, were infected with HBV (500 vge/cell) with additions of Tet (5 µg/mL) or mock-treated every 2-3 days for ten days. Western blot analyses were performed to detect the over-expressed kinases with anti-tag antibodies (**B**) and viral parameters (**C** to **F**). Results are normalized to the mock situation and expressed as the mean +/- SD of 4 independent experiments, each performed in triplicate.

Figure 5. DYRK1A interacts with HBV cccDNA. Native ChIP analyses were performed on HBV-infected dHepaRG (**A**) or PHH (**B** and **C**) using antibodies against HBc or DYRK1A. The results obtained with two different antibodies against DYRK1A are shown in panels **B** and **C**. cccDNA was quantified from immuno-precipitated material by TaqMan qPCR. Results represent the mean +/- the standard error of the mean of 3 (dHepaRG) or 5 (PHHs) independent experiments.

Figure 6. DYRK1A activity on HBV transcription regulatory regions. A. Schematic view of a linearized HBV genome (genotype D, ayw subtype), showing the position of the four main ORFs (the HBx ORF is indicated by a large green arrow) and of the two enhancer/promoter regions. The enhancer 1/X promoter region contains a putative DYRK1A recognition site that partially matches the consensus motif found in DYRK1A responsive cellular promoters. The two mutation M1 and M2 introduced within enhancer 1/X promoter are shown below. Nucleotide positions are numbered from the unique EcoRI site. **B.** Plasmid containing the luciferase cDNA under the control of the enhancer 1/HBx (pX_Luc) promoter or the enhancer 2/preCore/Core (preC/C-Luc) regions were cotransfected together with a plasmid coding for DYRK1A, DYRK1A_KR, or a control plasmid into Huh7-NTCP cells. Luciferase assays were performed three days later. **C.** Luciferase assays were performed by transfecting plasmids containing either a wt or two different mutated putative DYRK1A motifs.

Results are normalized to the cells co-transfected with a control plasmid (Mock) and

expressed as the mean +/- SD of 4 independent experiments, each performed in triplicate.

Figure 7. DYRK1A over-expression increases the level of HBV nascent RNAs. Run-On

analysis. dHepaRG-TR-DYRK1A or -DYRK1A_KR cells were infected with HBV and treated

with Tet or mock-treated as indicated in the legend of Figure 6A. Controls were provided by

cells incubated in the absence of BrU (- BrU) or in the presence of actinomycin D (Act D) or

RG7834. Labeled RNAs were immunoprecipitated using an anti-BrdU antibody. Input HBV

RNAs (left panel) and immunoprecipitated RNAs representing nascent RNAs (right panel)

were quantified by RT-qPCR. Results are expressed as the mean+/- SD of 2 independent

experiments, each performed in triplicate. ND: value under the detection limit.

Supplemental Figure 1. Kinase profiler assay of 1C8 (10 µM). Only the kinases inhibited by

1C8 with a residual activity<20% are shown. Grey bars indicate the CLK and DYRK kinases

targeted by the compound.

Supplemental Figure 2. RNA levels of selected CMGC kinases in dHepaRG, PHHs and

liver biopsies. RNA levels of CMGC kinases indicated measured by RT-qPCR. Relative

RNA levels are expressed by the 2∆Ct value calculated using the PRNP mRNA as control in

HepaRG cells (undifferentiated or differentiated, **A**) or PHHs (**B**), HBV-infected or mock-

infected. PHHs used were purified from two different donors, each analyzed in triplicate. **C.**

CLK1, CLK2, DYRK1A, DYRK1B and SRPK1 mRNA expression in human liver using RNA-

Seq data retrieved from Yoo *et al.* (74) expressed as Reads Per Kilobase Million (RPKM).

Supplemental Figure 3. Comparative effects of AZ191 and DB18 in HBV-infected PHH.

A. Experimental outline. B. PHH were infected and total viral RNAs and pgRNA were

quantified by RTqPCR after treatment. IFN: Interferon-α (500 IU/mL). Doses of AZ191 and

DB18 were 10 µM and 20 µM. Results are expressed as the mean +/- SD of two independent

experiments, each performed in triplicate.

Supplemental Figure 4. Toxicity assays of AZ191 in PHHs. Cells infected and treated

with various concentrations of AZ191, as indicated in Figure 2A, were analyzed for signs of

toxicity using either a neutral red (**A**) or a sulforhodamine B (**B**) assay. Puromycin (Puro) was

used as a positive control at 10 µM. Results are expressed as the mean +/- SD of 3

independent experiments, each performed in triplicate.

Supplemental Figure 5. Effect of CLK2 and SRPK1 KD in HBV-infected dHepaRG cells.

A. Experimental outline. **B.** KD efficiency of the siRNA treatments measured by RT-qPCR at

15-dpt. Results are shown relative to siCTL. **C.** Effect of the kinases KD on the production

HBV intracellular RNAs and secreted HBs/HBe antigens, expressed as relative to siCTL

(mean +/- SD, n>5, each performed in triplicate).

Supplemental Figure 6. DYRK1A and DYRK1B KD in HBV-infected dHepaRG cells. A.

Experimental outline. HBV-infected cells were transfected once with siRNA and then

harvested at 1-, 2-, and 4-dpt. **B** and **C**. Toxicity assays in siRNA transfected dHepaRG cells.

Infected and transfected cells were analyzed for signs of toxicity using either a neutral red

(**B**) or a sulforhodamine B (**C**) assay. Puromycin (Puro) was used as positive control at 10

µM. **D.** KD efficiency of the siRNA treatments measured by RT-qPCR at the indicated time

points. Results are shown relative to siCTL (mean +/- SD, n=2-3). **E**. Effect of the KD for the

kinases indicated on HBV RNA production. Intracellular RNA extracted at each time point

was analyzed by RT-qPCR to quantify HBV total RNAs. In all cases, results are shown

relative to siCTL (mean +/- SD, n=2).

Supplemental Figure 7. Divergent effects of DYRK1A KD in HBV-infected PHH. A.

Experimental outline. PHH were infected with HBV and 7 days later transfected with siRNA

targeting DYRK1A or with a control siRNA (siCTL). **B**. Representative Western blot analysis

of DYRK1A KD levels. **C** to **D**. DYRK1A and HBV RNA analysis of a set of 3 independent

experiments, each performed in triplicate, in which a decrease of HBV RNAs was observed.

E and **F** represent another set of 4 experiments in which HBV RNAs increased upon

DYRK1A KD. Results are expressed as the mean +/- SD. Statistical analysis is from

comparisons between siDYRK1A and control siRNA for each of the measurements.

Supplemental Figure 8. Compensatory effect of DYRK1B/DYRK1A levels in PHHs and

dHepaRG cells. **A.** mRNA levels were quantified in HBV-infected PHHs transfected with

siRNA targeting DYRK1A as shown in Figure S7A. Results were clustered according to their

effect on HBV RNA levels (see Figure S7D and F). **B**. Analysis of DYRK1A/DYRK1B

compensatory mRNAs variations in HBV-infected dHepaRG cells transfected with siRNA

targeting either of the kinases. Statistical analysis is from comparisons between siRNA

targeting one of the kinases and control siRNA for DYRK1A or DYRK1B mRNA levels.

Supplemental Figure 9. Compensatory effect of DYRK1B/DYRK1A levels in HBV-

infected PHHs. A. DYRK1A and DYRK1B protein levels in HBV-infected PHHs with KD for

each kinase. Cells were infected with HBV and then transfected with indicated siRNA.

Western blot analysis was performed at 7-dpt. Cyclin D1 levels were analyzed as a marker of

functional DYRK1 KD, since its accumulation has been shown to depend on both DYRK1A

and DYRK1B (57, 75). PHHs (+) and (-) refers to experiments in which an increase (+) or a

decrease (-) of HBV RNAs levels was observed following DYRK1A KD (see Figure S7). The

bands were quantified using ImageJ (numbers relative to control siRNA, below the blots). **B.**

and **C.** Effect of single DYRK1B or double DYRK1A/1B KD in HBV-infected PHHs. Cells

were infected and transfected as indicated in Figure S7A. DYRK1A, DYRK1B (**B**), and HBV

(**C**) RNAs were quantified at 7-dpt. Results are normalized versus control siRNA and

expressed as the mean +/- SD of 2 independent experiments, each performed in triplicate.

Supplemental Figure 10. Effect of DYRK1B and DYRK1B_KR over-expression on HBV

parameters. dHepaRG cells engineered to express either HA-DYRK1B or HA-DYRK1B_KR

under the control of a tetracyclin (Tet)-inducible promoter were infected with HBV and then

treated with Tet as indicated in Figure 4A. **A.** Western blot analysis to detect over-expressed

HA-DYRK1B or HA-DYRK1B_KR. **B** and **C.** Quantification of HBV RNAs. Results are

normalized to the mock situation and expressed as the mean +/- SD, of 2 independent

experiments, each performed in triplicate.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7