Direct observation of fluorescent proteins in gels: a rapid cost-efficient, and quantitative alternative to immunoblotting

Matthieu Sanial[1], Ryan Miled[1], Marine Alves[1], Sandra Claret[1], Nicolas Joly[1], Véronique Proux-Gillardeaux[1], Anne Plessis[1], Sébastien Léon[1,2]

Affiliations:

[1] : Université Paris Cité, CNRS, Institut Jacques Monod, F-75013, Paris, France.

[2] : Author for correspondence:

Sébastien Léon, PhD

Institut Jacques Monod

15 Rue Hélène Brion

75205 Paris cedex 13, France.

email: sebastien.leon@ijm.fr ; tel. +33 (0)1 57 27 80 57.

Running title:

In-gel fluorescence detection of fluorescent proteins as an alternative to immunoblotting

1 Abstract

The discovery of Green Fluorescent Protein (GFP) and its derivatives has revolutionized cell
biology. These fluorescent proteins (FPs) have enabled the real-time observation of protein
localization and dynamics within live cells. Applications of FP vary from monitoring gene/protein
expression patterns, visualizing protein-protein interactions, measuring protein stability, assessing
protein mobility and creating biosensors. The utility of FPs also extends to biochemical approaches
through immunoblotting and proteomic analyses, aided by anti-FP antibodies and nanobodies. FPs
are notoriously robust proteins with a tightly folded domain that confers a strong stability and a
relative resistance to degradation and denaturation. In this study, we report that various green and
red FPs can be maintained in a native, fluorescent form during the entire process of protein sample
extraction, incubation with sample buffer, loading and migration on SDS-PAGE with only minor
adaptations of traditional protocols. This protocol results in the ability to detect and quantify in-gel
fluorescence (IGF) of endogenously-expressed proteins tagged with FPs directly after migration,
using standard fluorescence-imaging devices. This approach eliminates the need for antibodies and
chemiluminescent reagents, as well as the time-consuming steps inherent to immunoblotting such
as transfer onto a membrane and antibody incubations. Overall, IGF detection provides clearer data
with less background interference, a sensitivity comparable or better to antibody-based detection, a
better quantification and a broader dynamic range. After fluorescence imaging, gels can still be used
for other applications such as total protein staining or immunoblotting if needed. It also expands
possibilities by allowing the detection of FPs for which antibodies are not available. Our study
explores the feasibility, limitations, and applications of IGF for detecting endogenously expressed
proteins in cell extracts, providing insights into sample preparation, imaging conditions, and
sensitivity optimizations, and potential applications such as co-immunoprecipitation experiments.

Introduction

The development of GFP as a tool to study protein localization and dynamics in live cells has
been a revolution in the field of cell biology, as recognized by the Nobel prize in Chemistry awarded
in 2008 to Osamu Shimomura, Martin Chalfie and Roger Tsien (Chalfie et al., 1994; Tsien, 1998).
GFP and its derivatives, as well as other fluorescent proteins (FPs) isolated since then from other
organisms with various properties (Lambert, 2019; Shaner et al., 2005), have become instrumental
in many fields of biology. They allow to monitor protein expression and localization in live cells, and
can be used as reporters of gene/protein expression patterns in organisms or tissues. Various
techniques have been implemented using GFP and its variants to visualize protein-protein
interactions in cells, through bimolecular fluorescence complementation (Hu and Kerppola, 2003;
Romei and Boxer, 2019), or fluorescence resonance energy transfer (FRET, BRET) (Hochreiter et
al., 2015). This also opened the way to the development of various probes and biosensors to report
on the quantity and/or localization of ions, metabolites and other organic molecules in cells and
tissues (Chandris et al., 2021; Wang et al., 2023; Zacharias et al., 2002). The various folding kinetics
of FPs have been exploited to report on protein stability and half-life (fluorescent timers) (Khmelinskii
et al., 2012; Subach et al., 2009; Terskikh et al., 2000). The fact that fluorescence bleaches upon
intense illumination also provides a way to evaluate protein mobility and dynamics in live cells
(Fluorescence Recovery after Photobleaching [FRAP] and Fluorescence Loss in Photobleaching
[FLIP]) (White and Stelzer, 1999). Similarly, photo-activatable and photo-switchable FPs allow the
study of distinct protein pools over time by activating or converting their fluorescence upon
illumination (Wang et al., 2023). Following the development of robust anti-GFP antibodies and in
particular of nanobodies (Rothbauer et al., 2006), GFP can also be used like any other epitope for
protein detection by techniques ranging from immunoblotting to immunoprecipitation and proteomic
analyses (Cristea et al., 2005; Rothbauer et al., 2008). Finally, the ability to express a GFP-binding
protein (derived from a llama single chain antibody) in cells fused with cellular proteins provides a
way to modify the subcellular localization and interactions of a GFP-tagged protein at will (Rothbauer
et al., 2008).

Because of these many applications, GFP and other FPs are frequently used as protein tags in
cell biology studies. An early step in the workflow consists in verifying the expression of an intact
fusion protein in cells by immunoblotting. Immunoblotting is also required when GFP is used as any
epitope to monitor the fusion protein's integrity, stability or post-translational modifications, or in
interaction studies based on co-immunoprecipitations. Protein detection by immunoblotting involves
primary antibodies and most generally a chemiluminescent reaction (ECL) using secondary
antibodies coupled to an enzyme (horseradish peroxidase). Whereas this approach is considered to
be more sensitive, enzyme kinetics may interfere with the linearity of the signal vis-à-vis the amount
of detected protein, a problem that may hinder the quantification of signals.

Over the past 20 years, fluorescent alternatives to chemiluminescence-based protein detection
were introduced, and the development of fluorescent dyes and the amelioration of fluorescence-
based imaging systems have allowed for a better sensitivity and better quantification of signals
(Eaton et al., 2014). Although initial systems focused on fluorescence detection at high wavelengths
(far-red and infrared, i.e. around 700 nm and above) to minimize background autofluorescence,
commercial imaging systems now allow the visualization of fluorescence at additional wavelengths
compatible with usual commercial fluorophores, eg. those used for immunofluorescence.

The versatility of these imaging systems is such that, in principle, the endogenous fluorescence
of the widely used FPs (eg. GFP) in extracts could be monitored directly in gels without the need of
transfer to membrane or the use of antibodies. SDS-PAGE involves denaturation of protein samples
at high temperatures in the presence of SDS and reducing agents prior to loading onto a gel,
procedures that are usually considered incompatible with the maintenance of FP fluorescence in gel
(Chew et al., 2009). A few studies have reported the observation of GFP-fluorescence in gels, but
these approaches require the use of native PAGE (Nemec et al., 2017), which has drawbacks
compared to regular, denaturing SDS-PAGE. On the other hand, GFP and FPs in general are known
to be tightly folded and more resistant to denaturation than most proteins (Saeed and Ashraf, 2009;
Ward, 2005). For example, a few studies exploited the ability of GFP to resist the adverse conditions
of SDS-PAGE to run protein overexpression screens in *Escherichia coli* or other cell types (Bird et
al., 2015; Bomholt et al., 2013; Geertsma et al., 2008; Krasnoselska et al., 2021; Madani et al., 2021;
Muller-Lucks et al., 2012; Newstead et al., 2007), or on recombinantly expressed or purified proteins
in vitro (Campbell et al., 2002; Donate-Macian et al., 2019; Nakatani et al., 2019; Weinberger Ii and
Lennon, 2021).

The development and popularity of fluorescence imaging devices is likely to lead to a
generalization of the use of FPs for direct visualization of in-gel fluorescence (IGF). Accordingly, a
recent study used this method for easy detection of FP-tagged proteins in protein extracts of cell
culture (Ruan et al., 2023).

In this study, we document that in-gel fluorescence (IGF) detection is a rapid and cost-effective
alternative to immunoblotting to visualize FP-tagged proteins expressed at endogenous levels in cell
extracts. IGF avoids the time-consuming process of transfer to a membrane, incubations with
antibodies and washes, with substantial improvements of the cost of the experiment as this does not
require primary nor secondary antibodies, membranes, or chemiluminescence substrates, and of
the quality of the data (less background and higher dynamic range). We describe the conditions of
use and limitations of various GFPs and RFPs for IGF, as well as sample preparation, imaging
conditions, sensitivity and possible applications.

Results

The fluorescence of endogenously yeGFP-tagged proteins can be visualized from cell extracts in SDS-PAGE gel after migration

To know whether we can detect GFP fluorescence from endogenously expressed proteins in total protein extracts, we tagged two yeast genes (*BMH1*, encoding a 14-3-3 protein) or HXK1, encoding hexokinase 1) at their endogenous locus with yeGFP, a yeast codon-optimized GFP (Cormack et al., 1997) which also harbors two mutations (S65G, S72A - a.k.a. "GFPmut 3", Cormack et al., 1996) selected to increase GFP fluorescence (ID on FPBase https://www.fpbase.org : A2OWC, Lambert, 2019). yeGFP is often used for fluorescent tagging as it is incorporated into a toolbox for PCR-based tagging of yeast genes at their endogenous genetic locus (Janke et al., 2004). Exponentially growing cells (5 OD equivalents) were lysed with glass beads at 4°C in 100 µL native lysis buffer (containing Tris-HCl, Triton X-100 and NaCl; see Material and Methods for more details). 4X Laemmli sample buffer (Bio-Rad, containing LDS and DTT) was then added to the lysates (1X final), and samples were then incubated at 30°C or 95°C for 5 min, before loading on commercial precast TGX gels, and run in Tris-Glycine-SDS (TGS) buffer. After migration, the gel was imaged with gel-imaging devices whose excitation/emission wavelengths were compatible with the observation of green fluorescent proteins (ChemiDoc MP, Bio-Rad: 460-490/518-546 nm; and Typhoon, Cytiva: 488/505-545 nm) (**Figure S1**). The gels displayed a green fluorescent signal at the size of the tagged proteins, as revealed by immunoblotting with anti-GFP antibodies after transfer of the same gel to a membrane (**Figure 1A**). Thus, at least some of the GFP molecules remained fluorescent throughout the sample preparation and gel migration process. Noteworthy, other lysis buffers were tested and also allowed the maintenance of GFP fluorescence (see Material and Methods for more details). On a side note, we observed that gel exposure to UV-light (such as during total protein labeling with a trihalo compound, eg. "Stain-Free" technology from Bio-Rad: 45 sec exposure at 300-400 nm) partially bleached the GFP fluorescence signal (**Figure S2**). Thus, this labeling should preferentially be performed after visualizing the GFP signal, especially if the signal is weak. Finally, as expected, GFP was no longer fluorescent when samples were denatured at 95°C instead of 30°C, or when extracts were prepared from cells precipitated with TCA in which proteins are fully denatured (**Figure 1A**).

Whereas these mild denaturing conditions allow the detection of in-gel fluorescence, two important points should be noted. First, the lower level of denaturation can impact on the migration of the GFP-tagged protein. At 30°C, Bmh1-yeGFP and Hxk1-yeGFP migrated faster than when samples were heated at 95°C or precipitated with TCA, as determined after transfer and immunoblotting with anti-GFP antibodies (**Figure 1A**). In-gel fluorescence detection systematically correlated with the presence of the fast-migrating band, suggesting it may correspond to a folded species of yeGFP. To address this point, proteins were transferred onto a nitrocellulose membrane

in classical conditions (liquid transfer), and immunoblotted with anti-GFP antibodies. Bmh1-yeGFP

fluorescence signal could be detected on the nitrocellulose membrane even after transfer and

incubation with antibodies, allowing the sequential observation of yeGFP fluorescence and antibody-

based labeling during the same acquisition. This revealed that the yeGFP fluorescence signal

precisely overlaid with the fast-migrating band detected with the anti-GFP antibody (**Figure 1B**).

Thus, yeGFP remains mostly folded in these mild denaturation conditions allowing in-gel

fluorescence observation, but can also cause a variation in the apparent molecular weight of yeGFP-

tagged proteins as compared to denatured proteins.

A second aspect that should be considered when using low denaturing conditions is that various

antibodies may recognize folded and denatured fluorescent proteins with different affinities. For

instance, anti-GFP antibodies may display a greater affinity towards folded or unfolded GFP, leading

to potential artifacts during quantification of the signal (**Figure S3A**). This could be circumvented by

denaturing proteins directly on the membrane after transfer (**Figure S3B**, see Material and Methods).

Comparison of GFP(S65T), yeGFP and EGFP for in-gel fluorescence

visualization

Following the development of tools for tagging of yeast genes with various tags by homologous

recombination (Longtine et al., 1998; Wach et al., 1997), Huh et al. generated a collection of yeast

strains which is widely used by the community and in which most genes are individually tagged with

GFP(S65T) (Huh et al., 2003) (FPBase ID: B6J33). To evaluate whether strains from this collection

are suitable for in-gel fluorescence detection, we retrieved the Bmh1-GFP(S65T) strain and

subjected it to the same analysis. A fluorescent signal was detected in the gel when samples were

incubated at 30°C, but not 95°C (**Figure 2A**). When compared to the signal obtained when tagging

the same protein with yeGFP, the fluorescent signal observed with GFP(S65T) was weaker,

probably because GFP(S65T) was less robust and more prone to denaturation. Indeed, a large

fraction of the Bmh1-GFP(S65T) protein pool was denatured even when incubated at 30°C prior to

gel loading, as judged by the migration pattern revealed with anti-GFP antibodies and antibodies

directed to the proteins of interest (**Figure 2A**). Thus, GFP(S65T) is subject to a partial denaturation

even in conditions which otherwise preserve yeGFP fluorescence. This was confirmed when

comparing the staining of Hxk1-GFP(S65T) with that of Hxk1-yeGFP (**Figure 2A**). In conclusion,

fusions with GFP(S65T) (including those from the GFP collection, Huh et al., 2003) are not the best-

suited to detect in-gel fluorescence in our conditions.

We also tested another commonly used GFP variant, EGFP, which derives from the original

avGFP and carries mutations F64L and S65T (a.k.a. "GFPmut 1", FPBase ID: R9NL8, Cormack et

al., 1996). This variant is also commonly used for the tagging of yeast genes at their endogenous

genetic locus (Janke et al., 2004) and is widely used in other eukaryotic organisms, notably through

their availability in the pEGFP-N1/pEGFP-C1 plasmids for expression in mammalian cells. The

signals obtained with EGFP were comparable to those obtained for yeGFP, making it appropriate
for visualization of IGF (**Fig 2B**).

In-gel fluorescence detection of proteins tagged with mNeonGreen and superfolder GFP

We next tested the suitability of more recently developed green-fluorescent proteins for IGF
detection (**Figure 3**). For comparative purposes, the same protein that we used for the initial variants
tested, Bmh1, was tagged with the brighter dLanYFP-derivative, mNeonGreen (FPBase ID: ZRKRV,
Shaner et al., 2013) or with superfolder GFP (sfGFP), a tightly folded GFP which is even more
resistant to denaturation than traditional GFP (FPBase ID: B4SOW, Pedelacq et al., 2006).

In all cases, a fluorescent signal was detected in the gel when samples were incubated at 30°C
prior to loading (**Figure 3**). Various fluorescence intensities were observed, with the strongest signal
obtained for Bmh1-yeGFP. However, these variations could be due to (i) the spectral properties of
these fluorescent proteins which may not appropriately match the imaging systems used for their
detection, (ii) the ability of these fluorescent proteins to endure the sample preparation conditions
and SDS-PAGE protocol, and (iii) an effect of the tag on the expression level of the tagged protein
(**Figure 3**). Indeed, in our experiments, Bmh1-yeGFP was more highly expressed than Bmh1-
mNeonGreen or Bmh1-sfGFP, as determined by immunoblotting these proteins with anti-Bmh1
antibodies. We hypothesized that this lower level of expression might be due to the fact that only
Bmh1-yeGFP is codon-optimized for expression in yeast. However, optimizing codon usage for
mNeonGreen expression in yeast (ymNeonGreen) neither led to an increase in protein expression
nor in IGF signal (**Figure S4**). This hypothesis remains open for sfGFP, as we did not test
optimization of codon usage for this FP.

The loss of IGF signal upon denaturation at 95°C was accompanied by a change in the migration
pattern as revealed by western blotting using anti-Bmh1 antibodies, with only one band present at a
higher apparent molecular weight, corresponding to the non-fluorescent species. An intermediate
situation was observed upon denaturation at 50°C, with a decrease in Bmh1-mNeonGreen IGF,
unlike what was observed for Bmh1-yeGFP or sfGFP (**Figure 3**). Accordingly, by immunoblotting,
treatment of samples at 50°C led to an almost full conversion of Bmh1-mNeonGreen from the native
to the denatured form, which was less pronounced for Bmh1-yeGFP. Thus, mNeonGreen is
amenable to IGF detection but is less heat-stable than other green FPs. In contrast, Bmh1-sfGFP
migration was unaffected at 50°C compared to 30°C, highlighting the stronger heat-stability of this
fluorescent variant. Overall, we conclude that sfGFP and EGFP (see **Figure 2B**) are more resistant
to heat-induced denaturation than yeGFP, GFP(S65T) and mNeonGreen.

These observations led us to quantify the heat-stability of fluorescence of the most robust green
FPs, which could be helpful for applications in which samples must be incubated at temperatures
higher than 30°C. We determined in-gel fluorescence as a function of the denaturation temperature
(**Figure 4**). In-gel fluorescence of Bmh1-sfGFP was unaltered until reaching 59°C, whereas Bmh1-

yeGFP fluorescence decreased starting at 54°C (**Figure 4A, C-D**). The temperature at which 50%

of the signal is still present was 67°C for sfGFP and 59°C for yeGFP. Interestingly, EGFP behaved

similarly to sfGFP (**Figure 4B, D**). The only additional gain of tagging with sfGFP over the widely

used EGFP is that it migrates almost exclusively as a single band at low denaturing temperatures

(30°C). Again, denaturation of all green FP-tagged Bmh1 was accompanied by a decrease in its

mobility on gels, which could clearly be visualized using anti-Bmh1 antibodies (**Figure 4, A-C**).

Altogether, these results allow us to reach the following conclusions. First, Bmh1-yeGFP and

Bmh1-EGFP give a strong IGF signal that, in our experiments, could be attributed to a higher

expression level coupled to a relatively high resistance to denaturation, especially in the case of

EGFP. Indeed, EGFP was as resistant as sfGFP to heat-induced denaturation in our conditions.

Second, despite being more sensitive to the conditions of extraction and imaging, mNeonGreen in-

gel fluorescence can still be used at low denaturation temperatures, which can avoid the purchase

of specific antibodies – indeed, mNeonGreen shares only 25% identity with the traditional avGFP

and is not recognized by anti-GFP antibodies. Finally, care should be taken regarding the expression

level of the tagged protein which, despite tagging at the endogenous genetic locus, can lead to

strong variations in expression depending on which FP was used for tagging (see **Figure 3**).

In-gel fluorescence of red fluorescent proteins

We aimed at expanding our observations to other, non-green fluorescent proteins. We could

not test blue-fluorescent proteins because none of the commonly-used imaging devices (Chemidoc

MP and Typhoon) provide filters compatible with the excitation/emission spectra of blue-fluorescent

proteins. However, these experiments were possible for red-fluorescent proteins. We tagged Bmh1

with the widely used fluorescent protein mCherry, a monomeric DsRed derivative with many

advantages such as high photostability, faster maturation and high pH-stability (FPBase ID: ZERB6,

Shaner et al., 2004). We also fused Bmh1 with mRuby2, a bright derivative of eqFP611 (FPBase ID:

8MJ78, Lam et al., 2012) , and with TagRFP-T, which derives from eqFP578 (Merzlyak et al., 2007)

with enhanced photostability (FPBase ID: LF3LJ, Shaner et al., 2008). Protein extracts were run on

SDS-PAGE and in-gel fluorescence was observed at excitation/emission wavelengths of 532/550-

800nm (Typhoon) or 520-545/577-613nm (ChemiDoc MP). All samples gave a red fluorescent signal

in gel (**Figure 5A**). Fluorescence signals were globally stronger when using a long-pass filter on the

Typhoon (550-800nm), which collects more fluorescence signals than the narrower filter present on

the Chemidoc MP (577-613nm) (**Figure S5**). The strongest fluorescence signal was obtained for

Bmh1-mCherry, whereas Bmh1-TagRFP-T gave the weakest signal. Immunoblotting using anti-

Bmh1 antibodies revealed migration patterns comparable to those obtained for green fluorescent

proteins: a faster-migrating band corresponded to the fluorescent species, whereas the slower

migrating form corresponded to the denatured pool. The ratio of fluorescent over denatured proteins

was higher for mCherry, although a minor denatured pool was already present in mild denaturing

conditions. Notably, the fluorescent pool of Bmh1-mRuby2 migrated much faster than any other FP-

tagged Bmh1 protein, with a decrease of about 15 kDa in apparent MW compared to the denatured
version. Bmh1-mCh was also the best detected red fluorescent protein on the Chemidoc MP (**Figure**
5A). Altogether, our conclusions are that (i) mild denaturation conditions are compatible with IGF of
RFPs, with mCherry being the best tag in our hands, with a stronger signal and resistance to heat-
induced denaturation; (ii) IGF allows the visualization of proteins tagged with RFPs originating from
various organisms without the need to purchase specific antibodies; (iii) tagging of proteins with red
FPs can lead to significant changes in the apparent molecular weight of the fusion proteins in mild
denaturing conditions.

Because mCherry is a widely used FP and is the most fluorescent of all RFPs tested, we then
evaluated the resistance of Bmh1-mCh to temperature-induced denaturation (**Figure 5, B-C**).
mCherry fluorescence was maintained at up to 70°C, with 50% of the signal still being present at
79°C. Thus, of all FPs tested (including green FPs), mCherry is the most resistant to denaturation
and is therefore perfectly appropriate for IGF signal detection.

Linearity of the signal and sensitivity of IGF compared to immunoblotting-
based chemiluminescence detection

Having established the capacity to detect fluorescence of numerous FPs in gels, we then turned
to studying the linearity of the signal and the sensitivity of this approach compared to the traditional
immunoblotting approach. When serial dilutions of Bmh1-yeGFP samples were initially assayed, we
discovered that the simple dilution of the protein lysate in Laemmli buffer (containing SDS and DTT)
was sufficient to promote FP denaturation, as determined by examining the migration profile using
anti–Bmh1 antibodies. This was likely a consequence of an increase in the detergent/protein ratio,
as previously observed (Xu and Keiderling, 2004) (**Figure S6**). To circumvent this problem, we
repeated the experiment using a non-denaturing sample buffer for dilutions of the sample (Laemmli
buffer without LDS). Under these conditions, protein fluorescence was maintained in spite of the
dilution (**Figure 6**).

In-gel fluorescence was compared to the signal obtained using primary antibodies followed by
secondary antibodies coupled to horseradish peroxidase (HRP) for chemiluminescent detection. We
used either antibodies raised against Bmh1 (**Figure 6A**) or GFP (**Figure 6B**). Overall, IGF detection
was comparable or better in sensitivity to chemiluminescent detection. This was particularly true
when comparing IGF with anti-GFP-based detection. Overall, we found the fluorescent signal to
more closely follow a linear plot than the chemiluminescent signal, over a range of 320-fold (**Figure**
6B). This result was confirmed for all FPs we tested, including mCherry (**Figure S7**). Thus, we
conclude that IGF detection is more reliable than chemiluminescent detection.

Increasing the pH protects against FP denaturation at low protein concentrations

Our observations above (**Figure S6**) suggested that sample dilution in Laemmli buffer compromises the integrity of GFP. Although we observed that this could be circumvented by the use of detergent-free Laemmli buffer (**Figure 6**), we sought to find conditions in which sample dilution would maintain FP integrity despite the presence of detergent (**Figure 7A**). First, we observed that dilution of the protein lysate in native lysis buffer (Tris-HCl 100 mM pH 8.0, NaCl 0.15 M, Glycerol 5%, Triton X-100 0.5 %) instead of water (**Figure 7A**) led to a protection of fluorescence, as determined both by IGF detection and the observation of native/denatured species by immunoblotting (**Figure 7B**). This result could partially be attributed to the presence of NaCl in the native lysis buffer, which preserved fluorescence in itself (**Figure 7B**). Moreover, increasing the pH to 8.0 or above was sufficient to protect fluorescence at low protein concentrations (**Figure 7C**), allowing IGF detection even for diluted protein samples. We conclude that increasing the sample pH to pH 8.0 protects GFP denaturation in diluted samples.

Use of IGF for the detection of FP-tagged proteins interaction by co-immunoprecipitation

We sought to exploit the power of IGF detection in a situation in which two fluorescent proteins must be detected. Protein-protein interactions can be studied by the co-expression of tagged proteins and immunoprecipitation of one partner to reveal the presence of the potential partner in the immunoprecipitate. As a case study, we studied the interaction between the yeast 14-3-3 proteins Bmh1 and Bmh2, which are known to heterodimerize *in vivo* (Chaudhri et al., 2003). Lysates of yeast expressing Bmh1-yeGFP and Bmh2-ymCh (a yeast codon–optimized version of mCherry, see Material and Methods) were subjected to immunoprecipitation using nanobodies-based traps. Similarly to our observations on diluted samples (section above), we realized that immunoprecipitates of FP proteins were denatured in the presence of Laemmli buffer, likely because of the low protein concentration in these samples. Based on our findings above (**Figure 7**), immunoprecipitates were resuspended in Laemmli buffer at pH 8.0 before loading, and samples were migrated on SDS-PAGE and imaged for IGF. In both cases (GFP-trap: **Figure 8A**; RFP-trap: **Figure 8B**), IGF detection allowed detection of both the bait and the prey on the same gel after exposure to the appropriate wavelengths, within minutes after uncasting the gel. One limitation was a bleed-through of intense green fluorescence signals in the red channel, which was notably observed using the Chemidoc (**Figure S8**) but also using the Typhoon to a limited extent (**Figure 8**). Thus, when looking at proteins tagged in both channels, the size of the green fluorescence-tagged protein should be different from that of the mCherry-tagged protein, and appropriate negative controls (samples without the GFP-tagged protein) should be included to determine the specificity of the signal.

Use of IGF in cell extracts from other organisms

Although IGF detection is based on endogenous FP fluorescence and thus should not be
affected by the cellular context in which these proteins are expressed, we tested the use of this
technique in other eukaryotic systems, including Drosophila cells or tissues (**Figure 9, A-B**), MDCK
cells (**Figure 9, C**) and *Caenorhabditis elegans* whole organisms (**Figure 9D**). Protein extracts were
prepared from FP-expressing cells in native conditions, resuspended in Laemmli buffer and samples
were loaded on SDS-PAGE gels (see Material and Methods). Overall, only minor adaptations to
current protein extraction protocols were made, and in all cases, IGF was observed, indicating the
applicability of IGF detection to the analysis of protein lysates from other organisms. In particular,
detection of the EGFP-tagged plasma membrane transducer Smoothened (SMO), which is
phosphorylated in response to Hedgehog (Jia et al., 2004) shows that IGF detection can be used to
study membrane proteins and post-translational modifications. In the case of total extracts from *C.*
elegans worms, we could detect GFP and mCherry fluorescence signals, however, we observed the
presence of non-specific fluorescent signals in the green channel that may complicate the readout
of IGF. First, autofluorescence of *C. elegans* animals was previously reported (Pincus et al., 2016),
and using red fluorescent proteins would avoid this problem. Second, we propose additional ways
to circumvent this problem. A band migrating just below the 70-kDa marker appeared in lysates from
WT animals which did not express any FP construct (**Figure 9D**), but this band disappeared upon
denaturation at 50°C. A fluorescent smear of unknown origin was also observed at around 250 kDa
(**Supp Fig 9A**), which resisted the denaturation at 50°C. However, this smear was not visible when
running the samples on homemade gels compared to commercial, pre-cast gels (**Supp Fig 9B**),
probably because this fluorescent material did not migrate into the resolving gel. Whereas these
observations provide ways to circumvent the non-specific green fluorescence observed in *C.*
elegans, it remains that care should be taken when setting up IGF detection in other material,
including the use of negative, non-FP-tagged extracts to ensure the specificity of the signal.
335

Discussion

Our study introduces in-gel fluorescence (IGF) detection as a rapid and cost-effective
alternative to traditional immunoblotting for visualizing fluorescent protein (FPs) signals in cell
extracts from yeast and other organisms, with minor adaptations of current protocols. The key points
resulting from our observations and optimizations are summarized in **Figure 10.**

Our work builds on the widespread use of FPs, particularly green and red fluorescent proteins
and their derivatives, as powerful tools in cell biology. GFP, originally developed as a live-cell
imaging marker, has since become key for various applications, including the monitoring of protein
expression and as a proteomic tool. Immunoblotting has traditionally been the method of choice to
visualize and quantify the expression of FP-tagged proteins. However, it comes with drawbacks, as
this is a time-consuming experiment with a substantial cost due to the reagents required, and is not
optimal for quanatifications compared to fluorescence-based approaches. Here, we propose IGF
detection as a fast, easy and efficient alternative to this technique.

Our study offers a comprehensive examination of IGF detection as an innovative tool for
protein detection after SDS-PAGE. We provide a systematic comparison of a number of widely used
FPs. Our evaluation considered key parameters such as fluorescence intensity, heat-stability, and
their overall suitability for IGF detection in the context of standard SDS-PAGE protocols and
accessible imaging devices. Importantly, this technique can be performed without major adaptation
of classical protocols except that protein denaturation should be carried out at low temperatures.

Overall, IGF detection is also a cost-effective method. This technique eliminates the need for
specific antibodies, which may even not be available in the case of less conventional FPs, as well
as nitrocellulose membrane or ECL reagents. This makes it an attractive option for labs that operate
under budget constraints. Because the observation is done directly from the gel after migration with
no additional steps, problems frequently observed with immunoblotting can be avoided, such as
heterogeneity of protein transfer, uneven antibody deposition, stains, and uneven ECL spreading on
the membrane.

Moreover, IGF is a time-saving technique, bypassing the need for protein transfer and
blotting/washing steps, as proteins can be visualized immediately after migration. For example, we
were able to use this technique to quickly screen for GFP-containing clones directly on colonies after
transformation (See Material and Methods). This complementary approach to PCR-based screening
allows to quickly monitor protein expression and size in multiple clones. The advantages of IGF
detection are even more appealing in the case of co-immunoprecipitations of two fluorescent
proteins, since both the prey and the bait can be visualized within minutes after migration, without
interference of primary antibody/secondary antibody recognition which can sometimes hinder signal
analysis and quantification.

Another key advantage of IGF detection is its versatility. All fluorescent proteins tested gave
a fluorescent signal in gels, which is advantageous notably in the case of FPs for which antibodies

are not yet available. However, not all FPs perform equally under all conditions. The results show
variations in the efficiency of IGF detection, emphasizing the need to consider both the FP's
fluorescence properties and its resistance to denaturation. Our findings demonstrate that EGFP and
sfGFP, in particular, display strong in-gel signals and good resistance to denaturation. These
features make them prime candidates for IGF applications. Moreover, mCherry stands out for its
exceptional heat-stability, maintaining its fluorescence even at relatively high denaturing
temperatures. This expands the possibilities of IGF detection to proteins that require higher
denaturation temperatures (such as proteins forming dimers or assembled in tight complexes). In all
cases, though, the intensity of the signal and the ability to specifically detect a fluorophore depend
on the available filters on the imaging systems, which were sometimes limiting (especially in the
case of mCherry) and led to bleed-through of signals when visualizing green- and red-tagged
proteins in the same sample.

Since fluorescent signals originate from the protein itself, the IGF detection method reduces
potential biases that can be encountered after using primary and secondary antibodies. Like all
fluorescence detection protocols, signal detection does not depend on a chemiluminescence
enzymatic reaction, which imposes kinetic constraints and whose activity can vary depending on
substrate accessibility. Depending on the antibody used, and thanks to the constant improvement in
the sensitivity of imaging devices, we found that IGF detection could outperform antibody-based
chemiluminescent detection, and the signal was globally more linear with respect to protein
concentration. Moreover, changing acquisition settings may increase sensitivity, with less
background than observed with chemiluminescence.

However, like all techniques, IGF has its limitations. In particular, this technique relies on a
partial denaturation of proteins at lower temperatures than usually performed in classical protocols.
Depending on the FP used for tagging, proteins treated in mild denaturation conditions can migrate
as 2 discrete bands corresponding to the fluorescent and denatured forms of the FP. The apparent
molecular weight of FP-tagged proteins is usually lower than under classical denaturation conditions,
with up to 15-kDa difference with the denatured form in the case of mRuby2. Furthermore, antibodies
may recognize denatured and non-denatured forms of the FPs with different affinities. Although IGF
was compatible with a range of lysis buffers, it is also possible that the composition of the lysis buffer
in which proteins are resuspended may alter overall fluorescence, depending on the abilities of FPs
to remain folded and fluorescent in these conditions. While these factors do not diminish the value
of IGF, they highlight the need for appropriate controls when setting up IGF for protein detection.

The method's simplicity, cost-effectiveness, reliability and compatibility with existing FPs and
current protocols make IGF detection an attractive alternative to immunoblotting which can greatly
benefit research in a range of fields. From basic molecular biology to more applied biomedical
research, IGF offers a robust and economical solution to detect fluorescent proteins in gels. Future
studies could further explore the method's robustness across different cell types and conditions,
ensuring its broad applicability in diverse experimental settings.

Material and Methods

Yeast strains and cultures.

Strains are derivatives of the BY4741 background and are listed in Supplementary Table 1. Proteins were tagged with various FPs at their endogenous loci by homologous recombination using plasmids listed in Supplementary Table 2, or originated from the collection of GFP-tagged strains (Huh et al., 2003). All yeast strains were constructed by transformation with the standard lithium acetate–polyethylene glycol protocol using homologous recombination and verified by polymerase chain reaction (PCR) on genomic DNA. Yeast cells were grown in YPD medium (2% w/v) or in SC medium [containing yeast nitrogen base (1.7 g/L; MP Biomedicals), ammonium sulfate (5 g/liter, Sigma-Aldrich), the appropriate drop-out amino acid preparations (MP Biomedicals), and 2% (w/v) glucose, unless otherwise indicated]. Precultures were incubated at 30°C for 8 hours and diluted in the evening to 20-mL cultures to reach mid-log phase the next morning.

Preparation of cell lysates from yeast

Exponentially growing cells (5 OD_{600} equivalents) were washed twice in water and resuspended in 100µL of cold native lysis buffer [Tris-HCl pH 8.0, 100 mM, NaCl, 0.15 M, Glycerol 5% v/v, Triton X-100 0.5% v/v, PMSF 0.5 mM, Complete antiprotease EDTA free (Roche, #11836170001)], and incubated for 5 min on ice-cold water with glass beads (Sigma-Aldrich, #G8772) to cool samples down prior to lysis. Cells were then lysed in a cold room on a vortex (4 x 30 sec, with 1 min incubation on ice-cold water between each lysis). Cell lysates were retrieved by piercing the bottom of the 1.5-mL tubes and brief centrifugation in a minispin centrifuge. Lysates were cleared by centrifugation at 4°C for 5 min at 3,000 xg. Supernatants were collected and 4X Laemmli sample buffer (Bio-Rad, #161-0747) containing DTT (12.5 µM final concentration) was added to the lysates (1X final concentration). Samples were denatured as indicated for each experiment, and loaded on a SDS-PAGE gel.

To use IGF from colonies growing on plates for screening purposes, 3 days-old colonies were harvested and resuspended in 100 µL cold native lysis buffer [Tris-HCl pH 8.0, 100 mM, NaCl, 0.15 M, Glycerol 5% v/v, Triton X-100 0.5% v/v, PMSF 0.5 mM, Complete antiprotease EDTA free (Roche, #11836170001)] and lysed as described above for exponentially growing cells. 15 µL of lysate were collected and 4X Laemmli sample buffer (Bio-Rad, #161-0747) containing DTT (12.5 µM final concentration) was added to the lysates (1X final concentration). Samples were denatured at 30°C for 5 min, and 20 µL were loaded on SDS-PAGE gel.

Relative quantitations were carried out with a detergent-compatible Bradford reagent (Abcam, #ab119216), and samples were then incubated for 5 min. Denaturation temperatures varied from 30°C to 95°C and are indicated in the Results section and the Figures. Equivalent protein amounts

446 were loaded on commercial precast 4-20% TGX gels (Bio-Rad, #4561094), and run in Tris-glycine-
447 SDS (TGS) buffer at 150 V for 45 min unless otherwise indicated.

Preparation of yeast protein extracts with TCA

A volume of culture corresponding to 1 OD_{600} equivalent of exponentially growing yeast cells
was collected, to which TCA (100%, w/v; Sigma-Aldrich) was added (10% final concentration).
Whole cells were precipitated on ice for 10 min, samples were then centrifuged at 16,000 xg at 4°C
for 10 min, the pellet was resuspended in 100 µL of a 10% (w/v) TCA solution and broken for 10 min
with glass beads (Sigma-Aldrich, #G8772) at room temperature. Lysates were transferred to another
1.5-mL tube to remove glass beads and centrifuged for 5 min at 16,000 xg at 4°C. Protein pellets
were resuspended in 50 µL of 1X Laemmli sample buffer, which included 50 mM Tris-base (final
concentration) to buffer for the TCA present in the pellet.

Preparation of cell lysates from Drosophila cultured cells

Drosophila wing imaginal disc cultured cells [Clone 8 (Cl-8) cells] responsive to Hedgehog (HH)
were cultured and transiently transfected using transitory insect transfection reagent (Mirusbio, #
MIR 6100) using pAct-EGFP, pAct-SMO-EGFP, and pAct-HH (Malpel et al., 2007; Sanial et al.,
2017) as described previously (Sanial et al., 2017). Forty-eight hours after transfection, cells were
washed in PBS and lysed in 1% Triton X-100, 150 mM NaCl, 50 mM Tris-HCl pH 8.0, 1 mM
dithiothreitol with complete EDTA-free antiprotease mix (Roche, #11836170001) and PhosSTOP
(Roche, #04906837001). Lysates were centrifuged (12,000 xg) for 10 min at 4°C and the
supernatant was mixed with 4X Laemmli sample buffer (Bio-Rad) containing DTT (12.5 µM final
concentration). Samples were incubated for 5 minutes at 25°C before loading on a homemade 10%
acrylamide gel without SDS. Gels were run for 90 min at 150 V in TGS buffer on a MiniProtean setup
(Bio-Rad).

Preparation of cell lysates from fly ovary tissue

Ovary extracts were obtained from Ptub64c-GAL4 and Ptub64c-GAL4; UbiEGFP-PAR3
transgenic flies (Kullmann and Krahn, 2018) by dissecting ovaries (20 flies per genotype) into 1X
PBS. Ovaries were placed on ice in lysis buffer [10 mM Tris-HCl pH 7.5, 150 mM NaCl, Complete
Protease Inhibitor cocktail (Roche)] and mechanically homogenized using micro pestles in matching
tubes. Ovaries lysates were spun at 3000 xg for 20 min at 4°C to eliminate debris. Supernatants
were incubated for 5 min at 72°C in LDS sample buffer with Sample Reducing Agent (ThermoFisher,
#NP0007). Samples were loaded on a NuPage, 4–12% Bis-Tris gels (ThermoFisher, #NP0322) and
run in 1X MOPS buffer (ThermoFisher, #NP0001) for 90 min at 150v.

Preparation of cell lysates from MDCK cells

MDCK cells (ATCC CCL-34) were transfected with Lipofectamine 2000 (Thermofisher #11668027) as described (Walch et al., 2018). Arf1-EGFP and Arf1-mCherry plasmids were obtained by subcloning Arf1 (Wessels et al., 2006) into pEGFP-N3 (Clontech) and pmCherry-N1 (Clontech), respectively. Cells were lysed in 6-well plates at confluency with 200 µL of lysis buffer [Tris-HCl pH 8.0 50 mM, NaCl 150mM, EDTA 10 mM, Triton X-100 0.5% (v/v) with complete antiprotease mix (Roche #11836145001)]. After 20 min incubation on ice, cells were scraped from the plate, collected in tubes and vortexed 3 x 5 sec, with 30 sec-incubation on ice between each step. Lysates were cleared by centrifugation 30 min, 11000 xg at 4°C. After protein quantitation using a Bradford reagent (Bio-Rad #5000205), samples were mixed with Laemmli sample buffer (Bio-Rad #161-0747) (1X final concentration). 50 µg of proteins were loaded per well on a precast 4-20%TGX gel (Bio-Rad, #4561094), and run in TGS buffer at 200 V for 45 min.

Preparation of cell lysates from *C. elegans*

Worms [*C. elegans* strain N2 (wildtype ancestral, Bristol) and JDU233: *ijmSi63 [pJD520; mosII_5'mex-5_GFP::tba-2; mCherry::his-11; cb-unc-119(+)] II; unc-119(ed3) III* (Lacroix et al., 2024)] were cultured on MGM++ plate and fed on OP50 bacterial strain at 23C. Worm lysate was obtained from an asynchronous population. A total volume equivalent of 50 µL worm pellet was washed twice in bacterial M9 minimal medium containing 0.05% (v/v) Tween-20, resuspended into 200 µL Lysis buffer [50mM Tris-HCl pH 8.0, 200 mM NaCl, 5% glycerol and Complete protease inhibitor cocktail (Roche #11836145001)] and sonicated for 1 min on ice. Total extract was incubated for 5 min at 50°C with 1x final pH8 Laemmli sample buffer containing DTT (12.5 µM final concentration). 40 µg of worm extract were loaded per well onto a precast 4-20%TGX gel (Bio-Rad, #4561094), and run in TGS buffer at 200 V for 45 min.

In-gel fluorescence imaging

After migration, the gel was imaged with gel-imaging systems whose excitation/emission wavelengths were compatible with the observation of green and red fluorescent proteins. Imaging with the Amersham Typhoon 5 (Cytiva) was carried out using the 488 nm laser with the Cy2 filter 525BP20 (for green FPs) or the 532 nm laser with the Cy3 filter LPG550 (for red FPs). The scanning resolution was 25 µm/pixel. PMT voltage was adjusted so that collected signals do not saturate. Imaging on the Chemidoc MP (Bio-Rad) was carried out using standard excitation/emission wavelengths recommended for Cy2/Cy3 filters (green FP: Epi-blue, excitation: 460−490 nm, emission: 518–546 nm; red FP: Epi-green, excitation: 520−545 nm, emission: 577–613 nm).

Total proteins were visualized by in-gel fluorescence using a trihalo compound incorporated in the commercial pre-cast SDS–PAGE gels (stain-free TGX gels, 4–20%; Bio-Rad) after 45 sec UV-

512　induced photoactivation using a ChemiDoc MP imager (Bio-Rad), serving as a loading control. Note
513　that UV irradiation should preferentially be performed after FP imaging (see **Figure S2**).

Immunoblotting

After fluorescence imaging, gels were equilibrated in transfer buffer [25 mM Tris, 190 mM
glycine, 20% ethanol (v/v), 0.02% SDS (w/v)] for 15 min and transferred at 100 V onto a nitrocellulose
membrane in a liquid transfer system (MiniProtean, Bio-Rad). Membranes were blocked in Tris-
buffered saline solution (50 mM Tris pH 7.6, 150 mM NaCl) containing 0.1 % Tween-20 (v/v) (TBS-
T) and 2% fat-free milk (w/v) for 30 min and incubated for at least 2 hours with the appropriate
primary antibodies. Membranes were washed 3 x 10 min in TBS-T and incubated for at least two
hours with the corresponding secondary antibody (coupled with horseradish peroxidase or
Alexa680). Membranes were then washed again 3 x 10 min in TBS-T and incubated with ECL Select
Western Blotting Detection Reagent (Cytiva #RPN2235) except in **Figure S3B** in which Clarity
Western ECL Substrate (Bio-Rad #1705060) was used because of the strong signal obtained with
the anti-GFP antibody 3H9. Luminescence signals were acquired using a ChemiDoc MP (Bio-Rad).
For denaturation of proteins on the membrane (**Figure S3B**), the membrane treatment was adapted
from Xu et al (2019) (Xu et al., 2019): after transfer, the membrane was washed with water, incubated
with cold 50% methanol (v/v) on ice for 30 min. The membrane was then washed with water, quickly
dried on lint-free wipes and incubated in a sandwich between two sheets of Whatman paper and two
glass slides at 75°C in a preheated thermocycler for 30 min.

Antibodies

The primary antibodies used are: anti-GFP (clones 7.1 and 13.1; Roche #11814460001; 1/5,000
dilution), Anti-GFP (Chromotek #3H9; 1/5,000), anti-GFP-DyLight800 (Rockland, #600-145-215;
1/2,000), anti-Bmh1 (kind gift of S. Lemmon, Univ. Miami, USA; 1/15,000) (Gelperin et al., 1995);
anti-Hkx2 (Rockland, #100-4159; 1/3,000). Secondary antibodies are anti-Mouse IgG-HRP (Sigma-
Aldrich, #A5278; 1/5000), anti-Rabbit IgG-HRP (Sigma-Aldrich, #A6154; 1/5,000), anti-Rat IgG-HRP
(Jackson Immuno, #112-035-143; 1/5,000), and anti-Mouse IgG Alexa Fluor 680 (ThermoFisher,
#21057; 1/5,000).

Test of the linearity of the signal by sample dilution

Extraction was performed in native conditions using 15 OD_{600} equivalent of yeast lysed in 300
µL of cold native lysis buffer [100 mM Tris-HCl pH 8.0, NaCl 0.15 M, Glycerol 5% v/v, Triton X-100
0.5% v/v, PMSF 0.5 mM, Complete antiprotease EDTA free (Roche, #11836170001)]. 4X Laemmli
sample buffer (Biorad) was added to the lysates (1X final, 12.5 µM DTT final concentration), and
samples were then incubated for 5 min at 30°C. Samples were serial-diluted with 1X native sample
buffer (62.5 mM Tris-HCl pH 6.8, 10% glycerol (v/v), 0.005% bromophenol blue (w/v)], as follows: 9

times at 0.75X dilution, followed by 5 times at 0.5X dilution (final dilution >300 fold). Fluorescence
levels were quantified using ImageQuant TL v8.2 (Cytiva).

Sample denaturation with a range of temperatures

Extraction was performed in native conditions using $15OD_{600}$ equivalent of yeast lysed in 300µL
of cold native lysis buffer [100 mM Tris-HCl pH 8.0, NaCl 0.15 M, Glycerol 5% v/v, Triton X-100 0.5%
v/v, PMSF 0.5 mM, Complete antiprotease EDTA free (Roche, #11836170001)]. 4X Laemmli sample
buffer was added to the lysates (1X final, DTT 12.5 µM final concentration). Samples were distributed
in 0.5 mL Eppendorf tube and kept on ice until heat treatment. Samples were heated at various
temperatures in an Eppendorf Gradient Mastercycler for 5 min, and kept at 22°C after incubation.
The equivalent of 0.5 OD_{600} was loaded in each well. Fluorescence levels were quantified using
ImageQuant TL v8.2 (Cytiva).

Co-immunoprecipitations of fluorescent proteins from yeast lysates

Exponentially growing cells (40 OD_{600} equivalents) were washed twice in water and
resuspended in 400 µL of cold IP lysis buffer (100 mM Tris-HCl pH8, NaCl 200 mM, Glycerol 5%,
Triton X-100 0.1%, EDTA 1 mM) containing protease inhibitors [PMSF 0.5 mM, Complete
antiprotease EDTA free (Roche, #11836170001), NaF 10 mM], and incubated for 5 min on ice-cold
water with glass beads. Cells were then lysed in a cold room on a vortex (4 x 30 sec, with 1 min
incubation on ice-cold water between each lysis). Relative quantitation was made with Bradford
reagent to use equivalent amounts of proteins for co-immunoprecipitation. Samples were diluted
4:10 as per the manufacturer's instructions with dilution buffer (10 mM Tris-HCl pH 8.0, NaCl 200
mM, EDTA 1 mM) containing protease inhibitors [PMSF 0.5 mM, Complete antiprotease EDTA free
(Roche, #11836170001), NaF 10 mM]. A fraction (3%) was taken as the "Input" fraction. 10 µL GFP-
Trap Magnetic Particles M-270 (Proteintech, #gtd) or RFP-Trap Magnetic Agarose (Proteintech,
#rtma) were added to the remaining of the sample and incubated on a rotating wheel for 1h at 4°C
and washed 3 times with cold IP lysis buffer. Proteins were eluted in 40 µL of FP trap sample buffer
[66 mM Tris-HCl pH 8.0, 2% SDS, 100 mM NaCl, 12.5 µM DTT, 10% (v/v) glycerol, 0.002%
bromophenol blue (w/v)] by incubating at 50°C for 20 min with soft shaking in a dry bath incubator.
Half of the IP and 1% of the diluted lysate were loaded on the gel.

Construction of yeast-optimized mCherry and mNeonGreen

The mCherry and mNeonGreen sequences were codon-optimized for *Saccharomyces cerevisiae* using GENEius tool (giving ymCherry and ymNeonGreen, respectively) and synthetized (Eurofins Genomics), with a GSAGAGAGAGAG or GSGAGAGAGAGAGA linker respectively, flanked by the traditional S1/S3 oligo sequences used for amplification (Janke et al., 2004). Linkers were also codon-optimized but sequences are different to avoid recombination. DNA was provided cloned into pEX-A128 containing a polylinker site to allow subcloning into the pYM vector series (Janke et al., 2004) with various selection markers. ymNeonGreen was cloned in pYM16 at BsiW1/AscI sites, giving pYM-ymCherry-NAT (pSL778), and ymCherry was cloned in pYM17 at HindIII/AscI sites, giving pYM-ymNeonGreen-Hph (pSL766).

Acknowledgments

The authors wish to thank Dr Silke Hauf (Virginia Tech, Blacksburg, VA, USA), Dr Sandra Lemmon (University of Miami, Coral Gables, FL, USA), Dr Myriam Ruault (Institut Curie, Paris, France), Dr Julien Dumont (Institut Jacques Monod, Paris, FR) and Dr Alenka Copic (CRBM, Montpellier, FR) for sharing material, and Dr Catherine L. Jackson (Institut Jacques Monod, Paris, FR) for discussions and critical reading of the manuscript. This work was supported by grants from Université Paris-Cité IDEx (https://u-paris.fr; ANR-18-IDEX-0001, "Emergence en Recherche" RS30J20IDXA1_EMERLEON) and Agence Nationale pour la Recherche ("AMPKILL", ANR-23-CE13-0012-01) to S. Léon. N. Joly is supported by a grant from the Fondation ARC pour la recherche sur le cancer (ARCPJA2022050005002) and a grant from Université Paris-Cité IDEx (https://u-paris.fr; ANR-18-IDEX-0001, "Emergence en Recherche" RS30J23IDX64_KATAREP).

The authors declare no competing interests.

References

Bird, L.E., H. Rada, A. Verma, R. Gasper, J. Birch, M. Jennions, J. Löwe, I. Moraes, and R.J. Owens. 2015. Green fluorescent protein-based expression screening of membrane proteins in Escherichia coli. *J Vis Exp*:e52357.

Bomholt, J., C. Helix-Nielsen, P. Scharff-Poulsen, and P.A. Pedersen. 2013. Recombinant production of human Aquaporin-1 to an exceptional high membrane density in Saccharomyces cerevisiae. *PLoS One*. 8:e56431.

Brachmann, C.B., A. Davies, G.J. Cost, E. Caputo, J. Li, P. Hieter, and J.D. Boeke. 1998. Designer deletion strains derived from Saccharomyces cerevisiae S288C: a useful set of strains and plasmids for PCR-mediated gene disruption and other applications. *Yeast*. 14:115-132.

Campbell, R.E., O. Tour, A.E. Palmer, P.A. Steinbach, G.S. Baird, D.A. Zacharias, and R.Y. Tsien. 2002. A monomeric red fluorescent protein. *Proc Natl Acad Sci U S A*. 99:7877-7882.

Chalfie, M., Y. Tu, G. Euskirchen, W.W. Ward, and D.C. Prasher. 1994. Green fluorescent protein as a marker for gene expression. *Science*. 263:802-805.

Chandris, P., C.C. Giannouli, and G. Panayotou. 2021. Imaging Approaches for the Study of Metabolism in Real Time Using Genetically Encoded Reporters. *Front Cell Dev Biol*. 9:725114.

Chaudhri, M., M. Scarabel, and A. Aitken. 2003. Mammalian and yeast 14-3-3 isoforms form distinct patterns of dimers in vivo. *Biochemical and Biophysical Research Communications*. 300:679-685.

Chew, F.N., W.S. Tan, T.C. Ling, C.S. Tan, and B.T. Tey. 2009. Quantitation of green fluorescent protein using a gel-based imaging method. *Anal Biochem*. 384:353-355.

Cormack, B.P., G. Bertram, M. Egerton, N.A.R. Gow, S. Falkow, and A.J.P. Brown. 1997. Yeast-enhanced green fluorescent protein (yEGFP): a reporter of gene expression in Candida albicans. *Microbiology (Reading)*. 143 (Pt 2):303-311.

Cormack, B.P., R.H. Valdivia, and S. Falkow. 1996. FACS-optimized mutants of the green fluorescent protein (GFP). *Gene*. 173:33-38.

Cristea, I.M., R. Williams, B.T. Chait, and M.P. Rout. 2005. Fluorescent proteins as proteomic probes. *Mol Cell Proteomics*. 4:1933-1941.

Donate-Macian, P., E. Alvarez-Marimon, F. Sepulcre, J.L. Vazquez-Ibar, and A. Peralvarez-Marin. 2019. The Membrane Proximal Domain of TRPV1 and TRPV2 Channels Mediates Protein(-)Protein Interactions and Lipid Binding In Vitro. *Int J Mol Sci*. 20.

Eaton, S.L., M.L. Hurtado, K.J. Oldknow, L.C. Graham, T.W. Marchant, T.H. Gillingwater, C. Farquharson, and T.M. Wishart. 2014. A guide to modern quantitative fluorescent western blotting with troubleshooting strategies. *J Vis Exp*:e52099.

Geertsma, E.R., M. Groeneveld, D.J. Slotboom, and B. Poolman. 2008. Quality control of overexpressed membrane proteins. *Proc Natl Acad Sci U S A*. 105:5722-5727.

Gelperin, D., J. Weigle, K. Nelson, P. Roseboom, K. Irie, K. Matsumoto, and S. Lemmon. 1995. 14-3-3 proteins: potential roles in vesicular transport and Ras signaling in *Saccharomyces cerevisiae*. *Proc Natl Acad Sci U S A*. 92:11539-11543.

Hochreiter, B., A.P. Garcia, and J.A. Schmid. 2015. Fluorescent proteins as genetically encoded FRET biosensors in life sciences. *Sensors (Basel)*. 15:26281-26314.

Hu, C.D., and T.K. Kerppola. 2003. Simultaneous visualization of multiple protein interactions in living cells using multicolor fluorescence complementation analysis. *Nat Biotechnol*. 21:539-545.

Huh, W.K., J.V. Falvo, L.C. Gerke, A.S. Carroll, R.W. Howson, J.S. Weissman, and E.K. O'Shea. 2003. Global analysis of protein localization in budding yeast. *Nature*. 425:686-691.

Janke, C., M.M. Magiera, N. Rathfelder, C. Taxis, S. Reber, H. Maekawa, A. Moreno-Borchart, G. Doenges, E. Schwob, E. Schiebel, and M. Knop. 2004. A versatile toolbox for PCR-based tagging of yeast genes: new fluorescent proteins, more markers and promoter substitution cassettes. *Yeast*. 21:947-962.

Jia, J., C. Tong, B. Wang, L. Luo, and J. Jiang. 2004. Hedgehog signalling activity of Smoothened requires phosphorylation by protein kinase A and casein kinase I. *Nature*. 432:1045-1050.

Khmelinskii, A., P.J. Keller, A. Bartosik, M. Meurer, J.D. Barry, B.R. Mardin, A. Kaufmann, S. Trautmann, M. Wachsmuth, G. Pereira, W. Huber, E. Schiebel, and M. Knop. 2012. Tandem

fluorescent protein timers for in vivo analysis of protein dynamics. *Nat Biotechnol*. 30:708-
714.

Krasnoselska, G.O., M. Dumoux, N. Gamage, H. Cheruvara, J. Birch, A. Quigley, and R.J. Owens.
2021. Transient Transfection and Expression of Eukaryotic Membrane Proteins in Expi293F
Cells and Their Screening on a Small Scale: Application for Structural Studies. *Methods Mol*
Biol. 2305:105-128.

Kullmann, L., and M.P. Krahn. 2018. Redundant regulation of localization and protein stability of
DmPar3. *Cell Mol Life Sci*. 75:3269-3282.

Lacroix, B., S. Vigneron, J.C. Labbe, L. Pintard, C. Lionne, G. Labesse, A. Castro, and T. Lorca.
2024. Increases in cyclin A/Cdk activity and in PP2A-B55 inhibition by FAM122A are key
mitosis-inducing events. *EMBO J*. 43:993-1014.

Lam, A.J., F. St-Pierre, Y. Gong, J.D. Marshall, P.J. Cranfill, M.A. Baird, M.R. McKeown, J.
Wiedenmann, M.W. Davidson, M.J. Schnitzer, R.Y. Tsien, and M.Z. Lin. 2012. Improving
FRET dynamic range with bright green and red fluorescent proteins. *Nat Methods*. 9:1005-
1012.

Lambert, T.J. 2019. FPbase: a community-editable fluorescent protein database. *Nat Methods*.
16:277-278.

Lee, S., W.A. Lim, and K.S. Thorn. 2013. Improved blue, green, and red fluorescent protein tagging
vectors for S. cerevisiae. *PLoS One*. 8:e67902.

Leon, S., Z. Erpapazoglou, and R. Haguenauer-Tsapis. 2008. Ear1p and Ssh4p are new adaptors
of the ubiquitin ligase Rsp5p for cargo ubiquitylation and sorting at multivesicular bodies. *Mol*
Biol Cell. 19:2379-2388.

Longtine, M.S., A. McKenzie, 3rd, D.J. Demarini, N.G. Shah, A. Wach, A. Brachat, P. Philippsen,
and J.R. Pringle. 1998. Additional modules for versatile and economical PCR-based gene
deletion and modification in Saccharomyces cerevisiae. *Yeast*. 14:953-961.

Madani, G., E. Lamping, H.J. Lee, M. Niimi, A.K. Mitra, and R.D. Cannon. 2021. Small-Scale Plasma
Membrane Preparation for the Analysis of Candida albicans Cdr1-mGFPHis. *J Vis Exp*.

Malpel, S., S. Claret, M. Sanial, A. Brigui, T. Piolot, L. Daviet, S. Martin-Lanneree, and A. Plessis.
2007. The last 59 amino acids of Smoothened cytoplasmic tail directly bind the protein kinase
Fused and negatively regulate the Hedgehog pathway. *Dev Biol*. 303:121-133.

Merzlyak, E.M., J. Goedhart, D. Shcherbo, M.E. Bulina, A.S. Shcheglov, A.F. Fradkov, A. Gaintzeva,
K.A. Lukyanov, S. Lukyanov, T.W. Gadella, and D.M. Chudakov. 2007. Bright monomeric
red fluorescent protein with an extended fluorescence lifetime. *Nat Methods*. 4:555-557.

Muller-Lucks, A., S. Bock, B. Wu, and E. Beitz. 2012. Fluorescent in situ folding control for rapid
optimization of cell-free membrane protein synthesis. *PLoS One*. 7:e42186.

Nakatani, T., N. Yasui, I. Tamura, and A. Yamashita. 2019. Specific modification at the C-terminal
lysine residue of the green fluorescent protein variant, GFPuv, expressed in Escherichia coli.
Scientific reports. 9:4722.

Nemec, A.A., L.A. Howell, A.K. Peterson, M.A. Murray, and R.J. Tomko, Jr. 2017. Autophagic
clearance of proteasomes in yeast requires the conserved sorting nexin Snx4. *J Biol Chem*.
292:21466-21480.

Newstead, S., H. Kim, G. von Heijne, S. Iwata, and D. Drew. 2007. High-throughput fluorescent-
based optimization of eukaryotic membrane protein overexpression and purification in
Saccharomyces cerevisiae. *Proc Natl Acad Sci U S A*. 104:13936-13941.

Pedelacq, J.D., S. Cabantous, T. Tran, T.C. Terwilliger, and G.S. Waldo. 2006. Engineering and
characterization of a superfolder green fluorescent protein. *Nat Biotechnol*. 24:79-88.

Pincus, Z., T.C. Mazer, and F.J. Slack. 2016. Autofluorescence as a measure of senescence in C.
elegans: look to red, not blue or green. *Aging (Albany NY)*. 8:889-898.

Romei, M.G., and S.G. Boxer. 2019. Split Green Fluorescent Proteins: Scope, Limitations, and
Outlook. *Annu Rev Biophys*. 48:19-44.

Rothbauer, U., K. Zolghadr, S. Muyldermans, A. Schepers, M.C. Cardoso, and H. Leonhardt. 2008.
A versatile nanotrap for biochemical and functional studies with fluorescent fusion proteins.
Mol Cell Proteomics. 7:282-289.

Rothbauer, U., K. Zolghadr, S. Tillib, D. Nowak, L. Schermelleh, A. Gahl, N. Backmann, K. Conrath,
S. Muyldermans, M.C. Cardoso, and H. Leonhardt. 2006. Targeting and tracing antigens in
live cells with fluorescent nanobodies. *Nat Methods*. 3:887-889.

Ruan, Z., J. Lee, Y. Li, J. Du, and W. Lu. 2023. Human Pannexin 1 Channel is NOT Phosphorylated
by Src Tyrosine Kinase at Tyr199 and Tyr309. *bioRxiv*.
Saeed, I.A., and S.S. Ashraf. 2009. Denaturation studies reveal significant differences between GFP
and blue fluorescent protein. *Int J Biol Macromol*. 45:236-241.
Sanial, M., I. Becam, L. Hofmann, J. Behague, C. Arguelles, V. Gourhand, L. Bruzzone, R.A.
Holmgren, and A. Plessis. 2017. Dose-dependent transduction of Hedgehog relies on
phosphorylation-based feedback between the G-protein-coupled receptor Smoothened and
the kinase Fused. *Development*. 144:1841-1850.
Shaner, N.C., R.E. Campbell, P.A. Steinbach, B.N. Giepmans, A.E. Palmer, and R.Y. Tsien. 2004.
Improved monomeric red, orange and yellow fluorescent proteins derived from Discosoma
sp. red fluorescent protein. *Nat Biotechnol*. 22:1567-1572.
Shaner, N.C., G.G. Lambert, A. Chammas, Y. Ni, P.J. Cranfill, M.A. Baird, B.R. Sell, J.R. Allen, R.N.
Day, M. Israelsson, M.W. Davidson, and J. Wang. 2013. A bright monomeric green
fluorescent protein derived from Branchiostoma lanceolatum. *Nat Methods*. 10:407-409.
Shaner, N.C., M.Z. Lin, M.R. McKeown, P.A. Steinbach, K.L. Hazelwood, M.W. Davidson, and R.Y.
Tsien. 2008. Improving the photostability of bright monomeric orange and red fluorescent
proteins. *Nat Methods*. 5:545-551.
Shaner, N.C., P.A. Steinbach, and R.Y. Tsien. 2005. A guide to choosing fluorescent proteins. *Nat*
Methods. 2:905-909.
Subach, F.V., O.M. Subach, I.S. Gundorov, K.S. Morozova, K.D. Piatkevich, A.M. Cuervo, and V.V.
Verkhusha. 2009. Monomeric fluorescent timers that change color from blue to red report on
cellular trafficking. *Nat Chem Biol*. 5:118-126.
Terskikh, A., A. Fradkov, G. Ermakova, A. Zaraisky, P. Tan, A.V. Kajava, X. Zhao, S. Lukyanov, M.
Matz, S. Kim, I. Weissman, and P. Siebert. 2000. "Fluorescent timer": protein that changes
color with time. *Science*. 290:1585-1588.
Tsien, R.Y. 1998. The green fluorescent protein. *Annu Rev Biochem*. 67:509-544.
Wach, A., A. Brachat, C. Alberti-Segui, C. Rebischung, and P. Philippsen. 1997. Heterologous HIS3
marker and GFP reporter modules for PCR-targeting in Saccharomyces cerevisiae. *Yeast*.
13:1065-1075.
Walch, L., E. Pellier, W. Leng, G. Lakisic, A. Gautreau, V. Contremoulins, J.M. Verbavatz, and C.L.
Jackson. 2018. GBF1 and Arf1 interact with Miro and regulate mitochondrial positioning
within cells. *Scientific reports*. 8:17121.
Wang, M., Y. Da, and Y. Tian. 2023. Fluorescent proteins and genetically encoded biosensors.
Chem Soc Rev. 52:1189-1214.
Ward, W.W. 2005. Biochemical and Physical Properties of Green Fluorescent Protein. *In* Green
Fluorescent Protein. 39-65.
Weinberger Ii, J., and C.W. Lennon. 2021. Monitoring Protein Splicing Using In-gel Fluorescence
Immediately Following SDS-PAGE. *Bio Protoc*. 11:e4121.
Wessels, E., D. Duijsings, T.K. Niu, S. Neumann, V.M. Oorschot, F. de Lange, K.H. Lanke, J.
Klumperman, A. Henke, C.L. Jackson, W.J. Melchers, and F.J. van Kuppeveld. 2006. A viral
protein that blocks Arf1-mediated COP-I assembly by inhibiting the guanine nucleotide
exchange factor GBF1. *Dev Cell*. 11:191-201.
White, J., and E. Stelzer. 1999. Photobleaching GFP reveals protein dynamics inside live cells.
Trends Cell Biol. 9:61-65.
Wosika, V., E. Durandau, C. Varidel, D. Aymoz, M. Schmitt, and S. Pelet. 2016. New families of
single integration vectors and gene tagging plasmids for genetic manipulations in budding
yeast. *Mol Genet Genomics*. 291:2231-2240.
Xu, J., H. Sun, G. Huang, G. Liu, Z. Li, H. Yang, L. Jin, X. Cui, L. Shi, T. Ma, A. Kameyama, and W.
Dong. 2019. A fixation method for the optimisation of western blotting. *Scientific reports*.
9:6649.
Xu, Q., and T.A. Keiderling. 2004. Effect of sodium dodecyl sulfate on folding and thermal stability
of acid-denatured cytochrome c: a spectroscopic approach. *Protein Sci*. 13:2949-2959.
Zacharias, D.A., J.D. Violin, A.C. Newton, and R.Y. Tsien. 2002. Partitioning of lipid-modified
monomeric GFPs into membrane microdomains of live cells. *Science*. 296:913-916.
763
764

Figure legends

Figure 1. Detection of in-gel fluorescence from endogenously expressed proteins
tagged with yeGFP.

A. Control (WT) yeast or yeast expressing Bmh1-yeGFP, Hxk1-yeGFP were lysed in the
indicated conditions (Nat.: native conditions; TCA: lysis in denaturing conditions with TCA),
samples were resuspended in LDS sample buffer (1X final, Bio-Rad) and incubated for 5 min
at the indicated temperatures. After migration on a commercial precast 4-20% TGX gels (Bio-
Rad), gels were imaged for green fluorescence using a Typhoon and a Chemidoc MP, and
total proteins were visualized by the stain-free technology on a Chemidoc MP. After transfer
to a nitrocellulose membrane, proteins were immunoblotted with anti-GFP antibodies
(Roche) and detected by chemiluminescence using anti-mouse antibodies coupled to HRP.

B. Protein lysates were prepared as in A. Proteins were incubated at 30°C or 95°C for 5 min,
with and without prior incubation at 30°C (5 min). After migration on a commercial precast 4-
20%TGX gel (Bio-Rad), gels were imaged for green fluorescence using a Chemidoc MP, and
total proteins were visualized by the stain-free technology on a Chemidoc MP. Proteins were
then transferred to a nitrocellulose membrane and imaged again for yeGFP fluorescence,
which was maintained during transfer. Proteins were then immunoblotted with anti-GFP
antibodies (Roche) and anti-mouse antibodies coupled to Alexa Fluor 680. yeGFP
fluorescence was then visualized again before detecting the anti-GFP antibodies by
fluorescence on a Chemidoc MP.

Figure 2. Comparison of in-gel fluorescence of yeGFP-, GFP(S65T)- and EGFP-tagged
proteins

A. Yeast expressing Bmh1-yeGFP, Bmh1-GFP(S65T), Hxk1-yeGFP and Hxk1-GFP(S65T)
were lysed in native conditions, samples were resuspended in LDS sample buffer and
incubated for 5 min at the indicated temperatures. After migration on a commercial precast
4-20% TGX gels (Bio-Rad), gels were imaged for green fluorescence using a Chemidoc MP,
and total proteins were visualized by the stain-free technology on a Chemidoc MP. Proteins
were then transferred to a nitrocellulose membrane and imaged again for green
fluorescence, which was maintained during transfer. Proteins were then immunoblotted with
anti-GFP antibodies (Roche) and anti-mouse antibodies coupled to Alexa Fluor 680. Green
fluorescence was then visualized again before detecting the anti-GFP antibodies by
fluorescence. The membranes were then stripped and incubated with anti-Bmh1 or anti-
Hxk1/2 antibodies and then with anti-rabbit antibodies coupled to HRP, and revealed by
chemiluminescence on a Chemidoc MP.

B. Yeast expressing Bmh1-yeGFP and Bmh1-EGFP were lysed in native conditions, samples
were resuspended in LDS sample buffer and incubated for 5 min at the indicated
temperatures. After migration on a commercial precast 4-20% TGX gel (Bio-Rad), gels were
imaged for green fluorescence using a Typhoon or a Chemidoc MP, and total proteins were
visualized by the stain-free technology on a Chemidoc MP. Proteins were then transferred to
a nitrocellulose membrane and immunoblotted with anti-Bmh1 antibodies and then with anti-
rabbit antibodies coupled to HRP, and revealed by chemiluminescence on a Chemidoc MP.
* indicates the fluorescent species.

Figure 3. Comparison of in-gel fluorescence of yeGFP-, sfGFP- and mNeonGreen-
tagged proteins

Yeast expressing Bmh1-yeGFP, Bmh1-sfGFP or Bmh1-mNeonGreen were lysed in native
conditions, samples were resuspended in LDS sample buffer and incubated for 5 min at the
indicated temperatures. After migration on a commercial precast 4-20%TGX gel (Bio-Rad), gels
were imaged for green fluorescence using a Typhoon or a Chemidoc MP, and total proteins were
visualized by the stain-free technology on a Chemidoc MP. Proteins were then transferred to a
nitrocellulose membrane and immunoblotted with anti-Bmh1 antibodies and then with anti-rabbit
antibodies coupled to HRP, and revealed by chemiluminescence on a Chemidoc MP. * indicates
the fluorescent species.

Figure 4. Temperature-sensitivity of in-gel fluorescence of yeGFP-, EGFP- and sfGFP-
tagged proteins

A. Yeast expressing Bmh1-yeGFP were lysed in native conditions, samples were resuspended
in LDS sample buffer and incubated for 5 min at the indicated temperatures in a gradient
thermocycler (note that the temperature range is different for sfGFP). After migration on a
commercial precast 4-20%TGX gel (Bio-Rad), gels were imaged for green fluorescence
using a Typhoon, and total proteins were visualized by the stain-free technology on a
Chemidoc MP. Proteins were then transferred to a nitrocellulose membrane and
immunoblotted with anti-Bmh1 antibodies and then with anti-rabbit antibodies coupled to
HRP, and revealed by chemiluminescence on a Chemidoc MP.
B. Same as A. using yeast expressing Bmh1-EGFP.
C. Same as A. using yeast expressing Bmh1-sfGFP.
D. Quantification of the green fluorescence signal as a function of the denaturation temperature
for various green FP-tagged Bmh1 (n=3; ± SD). Solid line: sigmoidal fit of the data (GraphPad
Prism), dotted line: 95% confidence interval of the fit.

Figure 5. Comparison of in-gel fluorescence of tagRFP-T-, mRuby2- and mCherry-
tagged proteins

A. Yeast expressing Bmh1-tagRFP-T, Bmh1-mRuby2 or Bmh1-mCherry were lysed in native
conditions, samples were resuspended in LDS sample buffer and incubated for 5 min at the
indicated temperatures. After migration on a commercial precast 4-20%TGX gel (Bio-Rad),
gels were imaged for red fluorescence using a Typhoon or a Chemidoc MP, and total proteins
were visualized by the stain-free technology on a Chemidoc MP. Proteins were then
transferred to a nitrocellulose membrane and immunoblotted with anti-Bmh1 antibodies and
then with anti-rabbit antibodies coupled to HRP, and revealed by chemiluminescence on a
Chemidoc MP. * indicates the fluorescent species.

B. Yeast expressing Bmh1-mCherry were lysed in native conditions, samples were
resuspended in LDS sample buffer and incubated for 5 min at the indicated temperatures in
a gradient thermocycler. After migration on a commercial precast 4-20%TGX gel (Bio-Rad),
gels were imaged for red fluorescence using a Typhoon, and total proteins were visualized
by the stain-free technology on a Chemidoc. Proteins were then transferred to a nitrocellulose
membrane and immunoblotted with anti-Bmh1 antibodies and then with anti-rabbit antibodies
coupled to HRP, and revealed by chemiluminescence on a Chemidoc MP.

C. Quantification of the red fluorescence signal as a function of the denaturation temperature
for mCherry-tagged Bmh1 (n=3; ±SD). Solid line: sigmoidal fit of the data (GraphPad Prism),
dotted line: 95% confidence interval of the fit.

Figure 6. Sensitivity and linearity of in-gel fluorescence detection of EGFP

A. *Left.* Yeast expressing Bmh1-EGFP were lysed in native conditions, samples were
resuspended in LDS sample buffer and incubated for 5 min at 30°C. Samples were serially
diluted (right to left) into sample buffer without LDS (see Material and Methods), and loaded
onto on a commercial precast 4-20%TGX gel (Bio-Rad). After migration, gels were imaged
for green fluorescence using a Typhoon, and proteins were then transferred to a
nitrocellulose membrane and immunoblotted with anti-Bmh1 antibodies and then with anti-
rabbit antibodies coupled to HRP, and revealed by chemiluminescence on a Chemidoc MP.
Right. Quantification of the signals obtained for green fluorescence (black) and
chemiluminescence (red) as a function of sample dilution. Linear-scaled and log-plot scaled
graphs are shown.

B. *Left.* Same experiment as in A, but using anti-GFP antibodies and anti-mouse antibodies
coupled to HRP to reveal proteins by chemiluminescence. *Right.* Quantification of the signals
obtained for fluorescence (black) and chemiluminescence (red) as a function of sample
dilution. Linear-scaled and log-plot scaled graphs are shown.

Figure 7. High pH preserves green fluorescence in diluted protein samples

A. Design of the experiment. Protein lysates prepared in native conditions from yeast
expressing Bmh1-EGFP are diluted 1:5 with dilution buffer (water, lysis buffer, NaCl at
various concentrations, or 100 mM Tris-HCl solution at various pH) before mixing with 4X
sample buffer pH6.8 (1X final concentration). Samples are then incubated at 50°C for 10 min
before being loaded on SDS-PAGE.

B. Effect of lysis buffer and NaCl concentration on in-gel fluorescence. Samples were diluted
with water, native lysis buffer (1X, 0.75X, 0.5X or 0.25X in water) or NaCl solution (at the
indicated concentration) before being incubated at 50°C for 10 min. After migration on a
commercial precast 4-20%TGX gel (Bio-Rad), gels were imaged for green fluorescence
using a Typhoon, and total proteins were visualized by the stain-free technology on a
Chemidoc MP. Proteins were then transferred to a nitrocellulose membrane and
immunoblotted with anti-GFP antibodies and then with anti-mouse antibodies coupled to
HRP, and revealed by chemiluminescence on a Chemidoc MP.

C. Effect of pH on in-gel fluorescence. Samples were diluted with native lysis buffer (1X), water
or Tris-HCl 100 mM solution (final concentration) at the indicated pH before being incubated
at 50°C for 10 min. After migration on a commercial precast 4-20%TGX gel (Bio-Rad), gels
were imaged for green fluorescence using a Typhoon, and total proteins were visualized by
the stain-free technology on a Chemidoc MP.

Figure 8. IGF detection in the context of co-immunoprecipitation

A. Protein lysates prepared in native conditions with native IP lysis buffer from yeast expressing
Bmh2-ymCherry (a yeast codon-optimized mCherry) with or without the co-expression of
Bmh1-yeGFP were subjected to co-immunoprecipitation using GFP-trap beads.
Immunoprecipitates were incubated at 50°C for 20 min in 1X Laemmli sample buffer at pH
8.0 which allows to maintain protein fluorescence at low protein concentrations. Input
samples and immunoprecipitates (IP) are shown. After migration on a commercial precast 4-
20%TGX gel (Bio-Rad), gels were imaged for green and red fluorescence using a Typhoon,
and total proteins were visualized by the stain-free technology on a Chemidoc MP. * indicates
bleed-through of the green fluorescence into the red channel.

B. Protein lysates prepared in native conditions from yeast expressing Bmh1-yeGFP with or
without the co-expression of Bmh2-ymCh were subjected to co-immunoprecipitation using
RFP-trap beads and treated as in A. Input samples and immunoprecipitates (IP) are shown.
After migration on a commercial precast 4-20%TGX gel (Bio-Rad), gels were imaged for
green and red fluorescence using a Typhoon, and total proteins were visualized by the stain-
free technology on a Chemidoc MP.

Figure 9. IGF detection in cells from various organisms

A. Drosophila wing-imaginal-disc cultured cells [Clone 8 (Cl-8) cells] were transfected with
plasmids encoding the Hedgehog (HH) transducer Smoothened (SMO) fused to EGFP and
EGFP alone. They were also transfected (+) or not (-) with a construct allowing the
expression of HH. Cells were lysed in native conditions (see Material and Methods), lysates
were mixed with 4X Laemmli sample buffer and incubated for 5 minutes at 25°C before
loading on a homemade PAGE gel. IGF was detected on a Typhoon. SMO-EGFP is
phosphorylated in the presence of HH, causing a slower migration (red arrowhead).

B. Transgenic *Drosophila* flies expressing or not an EGFP-tagged version of the polarity protein
PAR-3 were dissected and ovaries were lysed in native conditions (see Material and
Methods). Samples were incubated for 5 min at 72°C in LDS sample buffer, loaded on a
NuPage Bis-Tris gel (ThermoFisher) and IGF was detected on a Typhoon after migration.
After fluorescence imaging, total proteins were stained with Coomassie stain and imaged on
the infrared channel on a Chemidoc MP.

C. MDCK cells expressing Arf1 fused to EGFP or mCherry were lysed in native conditions (see
Material and Methods). Samples were incubated for 10 min at 50°C in LDS sample buffer
and loaded on a gel. After migration on a commercial precast 4-20%TGX gel (Bio-Rad), gels
were imaged for green and red fluorescence using a Chemidoc MP, and total proteins were
visualized by the stain-free technology on a Chemidoc MP. Note that some bleed-through of
green fluorescence is observed in the red channel (see also **Fig. 8A** and **Fig. S8**).

D. Transgenic *C. elegans* expressing or not GFP-tagged alpha-tubulin and mCherry-tagged
histone H2B were lysed in native conditions (see Material and Methods) by sonication.
Samples were incubated for 5 min at 50°C in Laemmli sample buffer (pH 8.0). After migration
on a commercial precast 4-20%TGX gel (Bio-Rad), gels were imaged for green and red
fluorescence using a Typhoon, and total proteins were visualized by the stain-free technology
on a Chemidoc MP. * denotes a non-specific fluorescence protein present in extracts of WT
C. elegans whose fluorescence disappears when heating the sample at 50°C.

Figure 10. Overview of the workflow and technical considerations when using IGF
detection to visualize proteins in gel.

White boxes refer to figure panels in which specific points are addressed.

Supplementary material

Supplementary tables.

Table S1: Yeast strains used in this study

Name	Genotype & Description	Origin & Reference
ySL0066: **BY4741 (WT)**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0*	(Brachmann et al., 1998)
ySL2719: **Bmh1-yeGFP**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-yeGFP::KANMX*	This study
ySL3509: **Hxk1-yeGFP**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; HXK1-yeGFP::HPHNT1*	This study
ySL0435: **Bmh1-GFP(S65T)**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1:GFP(S65T)::HIS3MX6*	(Huh et al., 2003)
ySL3674: **Bmh1-EGFP**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-EGFP::HIS3MX6*	This study
ySL3422: **Bmh1-sfGFP**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-sfGFP::HIS3MX6*	This study
ySL3423: **Bmh1-mNeongreen**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-mNeonGreen::HPHNT1*	This study
ySL3595: **Bmh1-ymNeongreen**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-ymNeonGreen::HPHNT1*	This study
ySL3421: **Bmh1-TagRFP-T**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-TagRFP-T::CaURA3*	This study
ySL3420: **Bmh1-mRuby2**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-mRUBY2::HIS3MX6*	This study
ySL3419: **Bmh1-mCherry**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-mCherry::KANMX4*	This study
ySL3593: **Bmh2-ymCherry**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH2-ymCherry::NATNT2*	This study
ySL3595: **Bmh2-ymCherry Bmh1-yeGFP**	*MAT a; ura3Δ0, his3Δ1, leu2Δ0, met15Δ0; BMH1-GFP::KanMx4; BMH2-ymCherry::NATNT2*	This study

939 **Table S2: Plasmids for yeast tagging used in this study**

Name	Description	Origin & Reference
pYM12	Plasmid to tag genes with yeGFP at their endogenous locus (*KANMX4*)	(Janke et al., 2004)
pYM25	Plasmid to tag genes with EGFP at their endogenous locus (*HPHNT1*)	(Janke et al., 2004)
pYM28	Plasmid to tag genes with EGFP at their endogenous locus (*HIS3MX6*)	(Janke et al., 2004)
GTH-g (pSL648)	Plasmid to tag genes with sfGFP at their endogenous locus (*HIS3MX6*)	Addgene #81104 (Wosika et al., 2016)
pFA6-mNeongreen::Hph (pSL643)	Plasmid to tag genes with mNeonGreen at their endogenous locus (*HPHNT1*)	Silke Hauf's lab, Virginia Tech
pFa6-link-yoTag-RFP-T-CaURA3 (pSL558)	Plasmid to tag genes with yeast codon-optimized Tag-RFP-T at their endogenous locus (*CaURA3*)	Addgene #44877 (Lee et al., 2013)
pFA6a-link-yomRuby2-SpHIS5 (pS649)	Plasmid to tag genes with yeast codon-optimized Tag-RFP-T at their endogenous locus (*SpHIS5*)	Addgene #44843 (Lee et al., 2013)
pYM-mCherry (pSL1)	Plasmid to tag genes with mCherry at their endogenous locus (*KANMX4*)	(Leon et al., 2008)
pYM-ymCherry-NAT (pSL778)	Plasmid to tag genes with yeast codon-optimized mCherry (*NATNT2*)	This study
pYM-ymNeonGreen-Hph (pSL766)	Plasmid to tag genes with yeast codon-optimized mNeonGreen (*HPHNT1*)	This study

Supplementary Figure legends.

Supplementary Figure 1. Spectral properties of green fluorescent proteins used in this
study.

A. Excitation wavelengths of yeGFP (= GFPmut3), EGFP, sfGFP and mNeonGreen. Data were
retrieved from FPBase (Lambert, 2019). The window of excitation provided by the Chemidoc
MP imaging system is indicated in gray (460-490nm), the wavelength of the excitation laser
of the Typhoon is indicated as a dotted line (488 nm), as per the manufacturer's indications.
B. Emission wavelengths of yeGFP (=GFPmut3), EGFP, sfGFP, and mNeonGreen. Data were
retrieved from FPBase (Lambert, 2019) and have been divided by the quantum yield to
evaluate fluorescence efficiency for each FP. The window of emission collected by the
Chemidoc MP system is indicated in gray (518-546 nm), that of the Typhoon is indicated as
shaded (515-535 nm), as per the manufacturer's indications.

Supplementary Figure 2. Effect of UV irradiation and freezing on the fluorescence of
various fluorescent proteins.

A. Native lysates of cells expressing Bmh1 tagged with yeGFP, EGFP, sfGFP, mNeonGreen or
mCherry were kept on ice or frozen (-20°C, -80°C, or in liquid N_2) for one hour, thawed on
ice and incubated at 30°C for 5 min, and loaded onto SDS-PAGE. IGF was detected on a
Chemidoc MP before (top) and after (bottom) UV irradiation, which is required for the
"StainFree" labeling (Bio-Rad) of total proteins with a trihalo compound (see Material and
Methods).
B. Quantification of the effect of UV exposure on IGF (ratio of signal obtained after UV exposure
to that obtained before UV exposure, for each fluorophore; n=4; ± SD).

Supplementary Figure 3. Effect of denaturation temperature on the recognition of GFP-
tagged proteins by various anti-GFP antibodies.

A. A native lysate of cells expressing Bmh1-yeGFP was resuspended in LDS sample buffer and
incubated at the indicated temperatures. After migration, gels were imaged for green
fluorescence using a Chemidoc MP, and total proteins were visualized by the stain-free
technology on a Chemidoc MP. Proteins were then transferred to a nitrocellulose membrane,
immunoblotted with the indicated antibodies and revealed by chemiluminescence on a
Chemidoc MP. * indicates the fluorescent species.
B. A native lysate of cells expressing Bmh1-yeGFP was resuspended in LDS sample buffer and
incubated at the indicated temperatures. After migration, gels were imaged for green
fluorescence using a Chemidoc MP, and total proteins were visualized by the stain-free
technology on a Chemidoc MP. Proteins were then transferred to a nitrocellulose membrane

and immunoblotted with the indicated antibodies and revealed by chemiluminescence on a
Chemidoc MP. Membranes on the right were treated with methanol and high temperature
(see Material and Methods) to denature proteins and ameliorate the detection of GFP-tagged
proteins treated at low temperatures. * indicates the fluorescent species.

Supplementary Figure 4. Comparison of IGF signals of mNeonGreen and a
corresponding yeast-optimized version.

Yeast expressing Bmh1-mNG and Bmh1-ymNG were lysed in native conditions, samples
were resuspended in LDS sample buffer and incubated for 5 min at the indicated temperatures. After
migration, gels were imaged for green fluorescence using a Typhoon or a Chemidoc MP, and total
proteins were visualized by the stain-free technology on a Chemidoc MP. Proteins were then
transferred to a nitrocellulose membrane and immunoblotted with anti-Bmh1 antibodies and then
with anti-rabbit antibodies coupled to HRP, and revealed by chemiluminescence on a Chemidoc MP.
The size difference is due to a longer linker present in the ymNeonGreen construct.* indicates the
fluorescent species.

Supplementary Figure 5. Spectral properties of red fluorescent proteins used in this
study.

A. Excitation wavelengths of mCherry, TagRFP-T and mRuby2. Data were retrieved from
FPBase (Lambert, 2019). The window of excitation provided by the Chemidoc MP imaging
system is indicated in gray (520-545nm), the wavelength of the excitation laser of the
Typhoon is indicated as a dotted line (532 nm), as per the manufacturer's indications.
B. Emission wavelengths of mCherry, TagRFP-T and mRuby2. Data were retrieved from
FPBase (Lambert, 2019). The window of emission collected by the Chemidoc MP system is
indicated in gray (577-613nm), that of the Typhoon is indicated as shaded (≥550 nm), as per
the manufacturer's indications.

Supplementary Figure 6. Effect of sample dilution on FP denaturation.

Top, Yeast expressing Bmh1-yeGFP were lysed in native conditions, samples were
resuspended in LDS sample buffer, incubated for 5 min at 30°C and serially diluted (1:2) in LDS
sample buffer. After migration, gels were imaged for green fluorescence using a Typhoon. Proteins
were then transferred to a nitrocellulose membrane and immunoblotted with anti-Bmh1 antibodies
and then with anti-rabbit antibodies coupled to HRP, and revealed by chemiluminescence on a
Chemidoc MP. Red arrowhead: size of the native (fluorescent) protein; blue arrowhead, size of the
denatured protein.

Bottom, Quantification of the signals obtained for fluorescence (black) and chemiluminescence
(red: native protein, blue: denatured protein) as a function of sample dilution. Linear-scaled and log-

plot scaled graphs are shown. yeGFP denaturation increases with sample dilution, causing a bias

when quantifying fluorescence.

Supplementary Figure 7. Sensitivity and linearity of in-gel fluorescence detection of

various FP

A.   Yeast expressing Bmh1-yeGFP were lysed in native conditions, samples were resuspended

in LDS sample buffer and incubated for 5 min at 30°C. Samples were serially diluted (right

to left) into sample buffer without LDS (see Material and Methods), and loaded onto SDS-

PAGE. After migration, gels were imaged for green fluorescence using a Typhoon, and

proteins were then transferred to a nitrocellulose membrane and immunoblotted with anti-

Bmh1 antibodies and then with anti-rabbit antibodies coupled to HRP, and revealed by

chemiluminescence on a Chemidoc MP. *Right.* Quantification of the signals obtained for

green fluorescence (black) and chemiluminescence (red) as a function of sample dilution.

Linear-scaled and log-plot scaled graphs are shown.

B.   Same as A, with Bmh1-sfGFP.

C.   Same as A, with Bmh1-mNeonGreen.

D.   Same as A, with Bmh1-mCherry.

1027

1028   Supplementary Figure 8. Bleedthrough of green fluorescence in the red fluorescence

1029   channel when using a Chemidoc MP.

1030   The same gel as that presented in **Fig 8A** was imaged for red fluorescence in a Chemidoc MP.

1031   * indicates the green fluorescent signal that is detected in the red channel.

1032

Supplementary Figure 9. Endogenous fluorescent material in *C. elegans* worm extracts.

A.   Extended version of the gel presented in **Figure 9D**. Total protein extracts of *C. elegans*

worms were prepared as detailed in *Material and Methods* and loaded onto precast TGX 4-

20% gels (Bio-Rad). After migration, gels were imaged for fluorescence using a Typhoon

at the indicated excitation/emission wavelengths. A fluorescent smear of unknown origin is

visible in the green channel at around 250 kDa.

B.   The same samples as in A were loaded onto a precast TGX 4-20% gel (Bio-Rad) (left) or

homemade 10% SDS-PAGE gel (right). After migration, gels were imaged for fluorescence

using a Typhoon at the indicated excitation/emission wavelengths. The fluorescent smear

in the green channel at around 250 kDa was no longer observed in homemade gels.

1043

1044

1045

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5

Figure 6

Figure 7

Figure 8

Figure 9

Lysis: in **non-denaturing** conditions (eg. native lysis buffer), to preserve fluorescence.
Compatible with (max. tested concentration):
triton x-100 (1%) [9A], EDTA (10 mM) [9C], DTT (2 mM) [9C], antiprotease and antiphosphatase cocktails [9A], PMSF (0.5 mM), NaF (10 mM) [8].

Freezing

Sample Buffer

Co-IP: use regular IP/TRAP protocols. [8]

stable in sample buffer under classical freezing conditions:
- -20°C,
- -80°C, [S2A]
- -LN$_2$.

Classical 1X Laemmli buffer (containing SDS or LDS), 12.5 µM DTT.

1x IP sample buffer: Tris-HCl **pH8** 66 mM, SDS 2%, NaCl 0.1 M, DTT 12.5 µM. [7] [8]

Denaturation
°C

Start with **5 min at 30°C.**
Max temp:
- yeGFP 50°C, sfGFP 60°C, EGFP 60°C. [4]
- mCh 70° [5]
- mNG 30° [3]

20 min at 50°C with soft shaking. [8]

Migration
- Prestained protein ladders are fluorescent. Use unstained ladder or load only few microliters. [S9]
- In regular TGS buffer, **150V** is recommended to prevent overheating. Optional: electrophoresis tank can be placed in ice-cold water. Use pre-chilled buffer.
- Fill outside chamber with buffer to cover the whole gel.
- Uncast, cut blue migration front to avoid fluorescence signal.
- Keep gel in water (no transfer buffer).

Gel imaging
- Fluorescence imaging first then total protein acquisition (U.V. exposure) if applicable. [S2B]
- ⚠ Green fluorescence may bleed through in the red channel. [8A] [S8] [9C]
- Set up exposure time / PMT power to avoid saturation.

Protein staining
- Coomassie blue [9B]
- Silver nitrate

Protein transfer and immunoblotting
- **Equilibrate gel** in transfer buffer for 15Min.
- Liquid transfer works great (60 min at 100v).
- If Ponceau staining is required, image fluorescence on membrane before (**fluorescence is not maintained in Ponceau acidic solution**).
- To detect fluorescence on the membrane after antibody incubation, fluorescence should be detected before applying chemiluminescent reagent (**fluorescence is not maintained in hydrogen peroxide solution**).
- **Protein denaturation on membrane** can increase antibody detection of low temperature-treated samples. [S3B]

Figure 10

Figure S1

Figure S2

Figure S3

Figure S4

Figure S5

Figure S6

Figure S7

Figure S8

A

Denaturation temp. (°C):	30	50	30	50	30	50	30	50	30	50
Wild-type	+ −	+ −	+ −	+ −	+ −	+ −	+ −	+ −	+ −	+ −
GFP::tba-2; mCherry::his-11	− +	− +	− +	− +	− +	− +	− +	− +	− +	− +

ex:488, em:515-535
GFP

ex:532, em:550-800
mCherry

Overlay

ex:635, em655-685

Total Proteins

250kd
130kd
100kd
70kd
55kd
35kd
25kd
15kd
10kd

Typhoon   Chemidoc

B

Precast TGX 4-20%   Homemade 10%

| Protein Ladder | + − | + − | + − | + − |
| GFP::tba-2; mCherry::his-11 | − + | − + | − + | − + |

Fluorescent smear

Stacking

250kd
130kd
100kd
70kd
55kd
35kd
25kd
15kd
10kd

250kd
130kd
100kd
70kd
55kd
35kd
25kd

Resolving

ex:488
em:515-535
GFP

ex:532
em:550-800
mCherry

ex:488
em:515-535
GFP

ex:532
em:550-800
mCherry

Figure S9