Title: The proteotranscriptomic characterization of venom in the white seafan *Eunicella singularis* elucidates

the evolution of Octocorallia arsenal

Authors

Maria Vittoria Modica[1]*, Serena Leone[1]*, Marco Gerdol[2], Samuele Greco[2], Didier Aurelle[3], Marco Oliverio[4],

Giulia Fassio[4], Khadija El Koulali[5], Célia Barrachina[6], Sebastien Dutertre[7]

* equal contribution

1. Department of Biology and Evolution of Marine Organisms (BEOM), Stazione Zoologica Anton Dohrn, Italy

2. Department of Life Sciences, University of Trieste, Trieste, Italy

3. Aix Marseille Univ, Université de Toulon, CNRS, IRD, MIO, Marseille, France; Institut de Systématique, Evolution,

Biodiversité (ISYEB), Muséum National d'Histoire Naturelle, CNRS, Sorbonne Université, EPHE, Paris, France

4. Department of Biology and Biotechnology "Charles Darwin", Sapienza University of Roma, Italy

5. BCM, Univ. Montpellier, CNRS, INSERM, Montpellier, France

6. Platform MGX - Montpellier GenomiX, IGF, Univ. Montpellier, France.

7. Institut des Biomolécules Max Mousseron, UMR 5247, Université Montpellier – CNRS, Montpellier, France

Corresponding author email mariavittoria.modica@szn.it.
20

21 **Keywords**: venom, Cnidaria, Octocorallia, proteomics, transcriptomics, toxins

Abstract

All the members of the phylum Cnidaria are characterized by the production of venom in specialized structures, the nematocysts. Venom of jellyfish (Medusozoa) and sea anemones (Anthozoa) has been investigated since the 1970s, revealing a remarkable molecular diversity. Specifically, sea anemones harbour a rich repertoire of neurotoxic peptides, some of which have been developed in drug leads. However, venoms of the vast majority of Anthozoa species remain uncharacterized, particularly in the class Octocorallia. To fill this gap, we applied a proteo-transcriptomic approach to investigate the venom composition in *Eunicella singularis*, a gorgonian species common in Mediterranean hard-bottom benthic communities. Our results highlighted the peculiarities of the venom of *E. singularis* with respect to sea anemones, which is reflected in the presence of several toxins with novel folds, worthy of functional characterization. A comparative genomic survey across the octocoral radiation allowed us to generalize these findings and provided insights into the evolutionary history, molecular diversification patterns and putative adaptive roles of venom toxins. A comparison of whole-body and nematocyst proteomes revealed the presence of different cytolytic toxins inside and outside the nematocysts. Two instances of differential maturation patterns of toxin precursors were also identified, highlighting the intricate regulatory pathways underlying toxin expression.

Introduction

Cnidaria is the most ancient venomous lineage among the extant Metazoa, having originated 600-800 million years ago [1,2]. Venom is produced in the nematocysts, specialized cells with high morphological diversity [3,4] and is involved in different functions including prey capture, defence from predators, and possibly also *intra-* and *inter*-specific competition [5]. In addition, in sea anemones the production of venom occurs also in ectodermal gland cells, which may represent an ancestral secretory pathway for these animals [6].

Within Cnidaria, the Anthozoa is a large, well-supported clade comprising over 6,000 species, and it includes both hexacorals (e.g.: sea anemones and reef-building corals) and octocorals (e.g.: soft corals, sea fans and sea pens) [7,8]. Anthozoa are the subject of active investigation on venoms, but traditionally most studies have been focused on sea anemones (Hexacorallia, Actiniaria) [9–15]. Despite a reduced taxon sampling, these studies highlighted Actiniaria produce complex venoms, including over 15 different protein families with neurotoxic or cytolytic action [14–16]. Comparative analyses have demonstrated that sea anemones produce a neurotoxin-rich venom, while the venom of jellyfish (Scyphozoa) and hydroids (Hydrozoa) is abundant in cytolysins and enzymes [17]. These neurotoxins have been used since the 1970s to investigate nervous system functioning, and some of them possess a remarkable biomedical potential as drug leads. For instance, a synthetic analogue of the ShK toxin from *Stichodactyla helianthus* (ShK 186, or dalazatide) is in phase II trial for the treatment of several autoimmune diseases [18].

Much less is known about the venoms of Hexacorallia other than sea anemones (*e.g.*, reef-building corals and black corals) and of the other major lineages within Anthozoa: tube anemones (Ceriantharia) and Octocorallia [19]. Venom composition was investigated in four Ceriantharia species only and found predominantly cytolytic [20]. Octocorals, instead, have been completely neglected to date by venom studies, probably since they were supposed to possess few, small, and poorly diversified nematocysts [21]. Only their secondary metabolites have been characterized: these molecules, mostly produced by bacteria of the coral microbiomes, play key roles in defence [22]. Octocorallia include some of the most conspicuous organisms of rocky subtidal communities, both in tropical and temperate seas. These ecosystem engineers play important ecological roles and contribute substantially to benthic communities' diversity: in the

Mediterranean Sea, gorgonians constitute up to 40% of the biomass in rocky sublittoral habitats [23].

The white gorgonian *Eunicella singularis* (Esper, 1791) is widespread and abundant in the north-western

Mediterranean [24,25]. Despite hosting *Symbiodinium* zooxanthellae, *E. singularis* displays low levels of

primary productivity and relies on heterotrophic nutrition to meet its metabolic needs, feeding mainly on

zooplankton integrated with particulated organic matter (POM) [26]. Predation on zooplankton,

intraspecific competition and defence from predators (mostly snails, crabs and fish) [27] are physiological

activities likely supported, in *E. singularis* and allies, by the production of a specialized venom. Indeed,

preliminary evidence have demonstrated that haemolytic peptides are present in different species of

octocorals [28], and the diversity and abundance of nematocysts could be higher than previously held [21].

However, toxin diversity has not been addressed in Octocorallia so far, hampering our complete

understanding of a crucial eco-evolutionary question concerning Cnidaria biology and evolution: to what

extent are main toxin families conserved across Anthozoa? To start answering this question, we provided a

first overview of the venom produced by an octocoral, applying to *E. singularis* an integrated proteo-

transcriptomic approach, which employed whole colony transcriptomic data to match proteomic data

obtained from chemical-induced nematocysts discharge or from the whole body of polyps. Our analyses

unveiled several previously unreported putative toxin families, providing the first comprehensive

description of the venom of an octocoral species. Leveraging these findings, we aimed to assess the degree

of conservation of specific toxin folds of Octocorallia in comparison to other Anthozoa, and to elucidate how

venom toxin families diversified throughout the evolutionary radiation of octocorals. Employing a

comparative approach on available Octocorallia genomes, we reconstructed the evolutionary patterns of

the main toxin families, highlighting distinctive evolutionary innovations specific to Octocorallia. Eventually,

by complementing our analyses with untemplated, deep-learning assisted structural modelling of the most

relevant toxins, we attempted to infer their function and adaptive value.

Methods

(a) Specimens collection, venom sample preparation and RNA and proteins extraction

Colonies of *Eunicella singularis* were collected by scuba diving at a depth of 25-30 m around Giannutri Island,

in the Tuscany Archipelago (Italy) and kept alive in seawater until further processing. Polyps were dissected

from the colony under a stereomicroscope and pooled in three aliquots of about 20 each (about 15 mg of

tissue) that were preserved in Trizol at -80 °C until further processing. Total RNA was extracted using Trizol

(Thermo Fisher Scientific), following the manufacturer's instructions, obtaining 5-10,000 ng with RQN values

between 6.7 and 8.8. Proteins from 10 additional polyps were extracted through homogenization in milliQ

water, followed by centrifugation and analysis of the supernatant to produce the whole body proteome.

Additionally, 10 live colony fragments about 25 mm long were immersed for 30 seconds into 1 mL of EtOH

in a 1.5 mL tube to induce nematocyst discharge [29]. Samples were freeze-dried and resuspended in 100

µL milliQ water for proteomic analysis.

(b) RNASeq

To account for variations in venom expression among samples, three cDNA libraries for RNASeq were

prepared using an Illumina TruSeq Stranded mRNA Sample Preparation kit according to manufacturer's

instructions and validated qualitatively and quantitatively on a Fragment Analyser, as well as by qPCR on a

ROCHE LightCycler 480. The three libraries were then processed for paired-end sequencing-by-synthesis in

an Illumina HiSeq 2500 at the MGX facility (Montpellier, France), and quality controlled using FastQC. After

trimming, we obtained a number of reads longer than 30 bp and quality above Q20 comprised between 21

and 30 million.

(c) Bioinformatic analyses of transcriptomic data

Paired-end raw trimmed sequencing data obtained from the three libraries were pooled and *de novo*

assembled using Trinity v.2.5.1 [30] with default settings, allowing a minimum contig length of 200

nucleotides. The assembled transcriptome was virtually translated using TransDecoder v.5.01

(https://github.com/TransDecoder/), setting the minimum allowed ORF length to 100 codons.

Expression levels of toxin transcripts were calculated based on unique gene counts as Transcripts Per Million

(TPM), as this normalised unit allows an efficient and reliable comparison of gene expression levels both

within and between samples [31]. Briefly, trimmed reads from each library were separately mapped against

the transcriptome assembly, previously filtered to remove sequence redundancy by considering only the

longest transcript for each gene model.

(d) Protein digestion

Protein extracts were denatured, reduced and alkylated. Briefly, the volume of each sample was adjusted

to 100 µL with 100 mM triethylammonium bicarbonate (TEABC). 1M DTT (1 µL) was added to each sample,

which was then incubated 30 min at 60 °C before adding 10 µL 0.5 M Iodoacetoamide (IAA). Alkylation lasted

30 min in the dark. Enzymatic digestion was performed by adding 2 µg trypsin (Gold, Promega, Madison

USA) in TEABC 100 mM and incubating overnight at 30 °C. Then, peptides were purified and concentrated

using OMIX Tips C18 reverse-phase resin (Agilent Technologies Inc.) according to the manufacturer's

specifications. Peptides samples were finally dehydrated in a vacuum centrifuge.

(e) Shotgun proteomics

Samples were resuspended in 20 µL of 0.1% formic acid (buffer A) and 1 µL was loaded onto an analytical

50 cm reversed-phase column (0.75 mm i.d., Acclaim Pepmap 100® C18, Thermo Fisher Scientific) and

separated with an Ultimate 3000 RSLC system (Thermo Fisher Scientific) coupled to a Q Exactive HF (Thermo

Fisher Scientific) *via* a nano-electrospray source, using a 120-min gradient of 2 to 40% of buffer B (80% ACN,

0.1% formic acid) and a flow rate of 300 nL/min. MS/MS analyses were performed in a data-dependent

mode. Full scans (375 – 1,500 m/z) were acquired in the Orbitrap mass analyser with a 60,000 resolution at

200 m/z. For the full scans, 3×10^6 ions were accumulated within a maximum injection time of 60 ms and

detected in the Orbitrap analyser. The twelve most intense ions with charge states ≥ 2 were sequentially

isolated to a target value of 1×10^5 with a maximum injection time of 45 ms and fragmented by HCD (Higher-

energy collisional dissociation) in the collision cell (normalised collision energy of 28%) and detected in the

Orbitrap analyser at a resolution of 30,000.

(f) Bioinformatic analyses of proteomic data

PEAKS Studio 8.5 (Bioinformatics solutions, Waterloo, ON, Canada), a de novo assisted database software

[32-34] was used to analyse MS/MS data from *E. singularis* venom, with parameters set as in [35]. The

MS/MS spectra obtained from shotgun proteomics were matched to a custom database resulting from the

whole-body transcriptome of the same colony, translated in the six open reading frames.

Protein isoforms were considered only when their existence was supported by MS/MS fragmentation data,

whereas translated transcripts identified by PEAKS Studio as belonging to the same protein group were

clustered with CD-HIT [36] using an identity threshold of 90%. Only the most representative sequence was

retained for the subsequent annotation.

Protein annotation was performed as detailed in the Suppl. Met. and Fig.1. Toxin nomenclature of mature

polypeptides followed Oliveira et al. [37]

Structure predictions of mature toxins were performed with the colabfold implementation of AlphaFold2,

using MMSeqs2 for multiple sequence alignment generation [38,39]. To increase model confidence for the

most challenging targets, custom multiple sequence alignments with orthologue octocorallian sequences

were used as inputs and each model was built using 12 recycles. All predictions were performed without the

use of templates and energy minimization. The structures that were modelled with reasonable confidence

(average pLDDT ≥ 70, which is generally recognised as the threshold for reliable models) [40] were submitted

to the DALI [41] and RUPEE [42] servers for structural homology search against the Protein Data Bank.

Homology modelling of the saposin-like toxins in the closed conformation was performed with Modeller

v9.24 [43]. Electrostatic surfaces were calculated with APBS [44]. UCSF ChimeraX was used for model

analysis, visualization and graphics [45].

(g) Comparative genomic analyses

Toxins identified in *Eunicella singularis* were used as queries for sequence homology searches against a

selection of available anthozoan genomes, representative of the diversity of Octocorallia [46] and major

Hexacorallia lineages summarized in Suppl. Tab. 1.

Sequence homology searches were carried out with tBLAStn [47], based on an initial E-value significance

threshold arbitrarily set to 10. Potential hits were manually curated, by annotating the inferred CDS,

detecting putative splicing donor and acceptor sites with the aid of Genie [48]. The presence of *bona fide*

orthologues between *E. singularis* and the other species was assessed through combined observations, i.e.

(i) the conservation of the main topological feature of the encoded precursor proteins, such as the

identification of a signal peptide and the cysteine residues involved in the formation of the disulphide bonds

(whenever present); (ii) the shared presence of conserved protein domains, detected with a HMM profile

approach, based on the Pfam-A database; (iii) the conservation of gene architecture (e.g. intron number,

phase and position), with respect to the orthologous gene from the congeneric species *E. verrucosa*.

Whenever needed (i.e., U-GRTX-Esi20 and 21), a fine-scale classification of orthologous sequences was aided

by Maximum Likelihood phylogenetic inference analyses, carried out with IQ-Tree [49] with 1000 ultrafast

bootstrap replicates based on the best-fitting model of molecular evolution detected by ModelFinder [50].

RESULTS AND DISCUSSION

(a) *E. singularis* **produces a complex venom**

The extraction of venom from cnidarians, like *E. singularis*, is performed by stimulating the animals to fire

their nematocysts and then purifying the toxins. Traditional methods use electrical or mechanical

stimulation followed by toxin isolation from seawater. Alternatively, nematocysts can be isolated by density

gradient centrifugation and mechanically disrupted to release their contents. Electrical stimulation of *E.*

singularis colonies proved insufficient to recover enough crude venom for subsequent analysis. Hence, we

attempted an ethanol-induced discharge method. While this has been validated for the box jellyfish

Chironex fleckeri [29], its use for Anthozoan venoms has never been reported. Our results showed that the

protein fraction recovered from ethanol-immersed *E. singularis* colonies had a composition comparable to

the contents of nematocysts: proteomic analysis confirmed 162 translated transcripts (NEM-P, Suppl. Tab.

2), reduced to 108 non-redundant precursor protein sequences. Based on an extensive bioinformatic

annotation (see Methods, Suppl. Met. and Fig. 1), 33 of these sequences were associated to structural

components or basal physiology functions, 15 were enzymes, 28 were classified as unknown and 32 were

identified as putative toxins (Fig. 2B).

 Figure 1. Schematic representation of proteotranscriptomics and bioinformatic annotation pipelines employed in this
 work.

 Only 17 of them could be also detected also in the whole-body proteome (WB-P, Suppl. Tab. 3) of *E.*

 singularis, which, by comparison, comprised ca. 1400 translated transcripts. As the NEM-P constitutes a

 subset of the WB-P, its enrichment in putative venom components served as indirect confirmation of

 nematocysts firing due to the ethanol treatment. To identify additional putative toxins from other tissues

 (*e.g.*, ectodermal glands), we also performed a targeted HMM search in the WB-P, looking for known

 cnidarian toxin families, as defined by VenomZone (https://venomzone.expasy.org). This search produced

 only four sequences previously undetected in the NEM-P: a protein with homology to the Small Cysteine

 Rich Proteins (SCRiPs) family and three cytolysins (see below). The absence of these proteins in the NEM-P

 could be due either to incomplete ethanol-induced nematocyst firing due to differential reactivity of diverse

 types of nematocysts, or to the existence of additional venom delivery systems, such the ectodermal glands,

 as previously demonstrated for sea anemones [51], that were not affected by ethanol exposure. In

alternative, these proteins may lack a venom function despite having toxin-like folds (e.g. they might be

involved in neurotransmission or digestion). The NEM-P comprised some transcripts encoding precursors

of multiple mature toxins, which were further classified by predicted fold (Fig. 1C and Suppl. Tab. 4). In total,

we identified 44 mature toxins, including a calglandulin, a protein characterized by multiple EF-hand motifs

with an accessory role in venom production and processing, omitted from the detailed discussion below.

Figure 2. Structural and functional annotation of the venom from *E. singularis.* (A) Total Ion Chromatogram (TIC) of the
ethanol extract; (B) Functional annotation of the translated transcripts confirmed at the protein level in the NEM-P; (C)
Fold distribution of the mature toxins produced by *E. singularis* based on HMM profile search; the x axis represents the
number of proteomics-validated sequences.

Following [37], the newly characterized toxins were collectively termed "Gorgotoxins" (GRTX-Esi), according

to the most conservative placement of *Eunicella* in the family Gorgoniidae, although recent work has

proposed a separate family for this genus (Eunicellidae) [46].

(b) **Established toxin folds are present in the venom of *E. singularis***

Small Cystein Rich Proteins-like (octo-SCRiPs)

We retrieved four toxins with homology to hexacorallian SCRiPs, which are known to induce severe

neurotoxicity in zebrafish larvae, possess antimicrobial activity and potentiate the transient receptor

potential ankyrin 1 (TRPA1) in mouse [52-54]. These comprised U-GRTX-Esi1 (particularly abundant at the

transcript level, TPM 1585), 2a-b and 3 (Fig. 3). These proteins exhibited a peptide coverage of the predicted

mature sequence exceeding 95%, shared a significant homology and up to 68% pairwise sequence similarity

with hexacoral SCRiPs. Despite this similarity, each toxin showcased distinct features.

In particular, U-GRTX-Esi2a-b derived from a precursor containing two consecutive highly conserved SCRiP-

like domains (pairwise identity 78.9%), separated by a protease cleavage site. Interestingly, while both forms

could be detected in the WB-P, only U-GRTX-Esi2b was detectable in the nematocyst discharge (Fig. 3B and

Suppl. Fig. 1).

U-GRTX-Esi3, solely detectable in the WB-P, featured a more complex architecture, comprising two SCRiP-

like domains in the mature polypeptide, confirmed by the proteomic coverage (Fig. 3C). The absence of

predicted cleavage sites between these domains would qualify U-GRTX-Esi3 as a Secreted Cysteine-Rich

Repeat Protein (SCREP), an emerging typology of venom proteins found in various organisms, often with

high affinity and avidity towards their biological target and a multi-modal action [55].

248
Figure 3. Tissue-specific maturation patterns of the SCRiP-like toxins- from *E. singularis.* The sequences of the
precursors of U-GRTX-Esi1 (A), 2a-b (B) and 3 (C) are reported. Red residues correspond to predicted signal peptide,
green boxes define the SCRiP domains, the red rectangle defines a PQM protease recognition site [56], orange and blue
lines indicate the peptide coverage of the precursors in the WB-P and NEM-P, respectively.

Although SCRiPs-encoding genes were identified, often in large numbers and associated with tandemly

duplicated gene clusters, in the genomes of all the octocoral species investigated in this study, they have a

patchy occurrence in Hexacorallia, supporting a complex evolutionary history characterized by lineage-

specific expansions and rapid diversification [55]. It is unclear whether SCREPs with two consecutive SCRiP

domains were already present in the latest common ancestor of all Anthozoans, or independently acquired

in different lineages. Indeed, clear orthologues of U-GRTX-Esi3 could be identified only in *P. clavata,*

Trachytela sp. and *R. reniformis*: this architecture had been previously identified also in the hexacoral

Orbicella faveolata [53]. However, while the only SCRiP so far reported in Octocorallia (in *Scleronephthya*

gracillima) presented a conventional cysteine framework [55], the domains here identified display a shift in

the cysteine pattern, which nonetheless did not affect the overall protein fold (Suppl. Fig. 2).

ShK-containing toxins

The ShK toxins, first identified in the venom of the sea anemone *Stichodactyla helianthus*, are potent K_V1

channels blockers [57]. Our search for ShK domains in the proteomes of *E. singularis* led to the identification

of U-GRTX-Esi4 in the NEM-P (TPM 103), with three consecutive ShK domains separated by short linker

regions. Only the C-terminal domain has the typical ShKT cysteine pattern, whereas the first two domains

present an unusual shift of the C-terminal cysteine. None of the domains of U-GRTX-Esi4 presents the key

Lys residue necessary for binding $K_V1.2$ and $K_V1.3$, while the subsequent Tyr residue, also important for

binding K_V1, is extremely conserved (Suppl. Fig. S3). This architecture resembles the precursors of the NEP3

protein family from *N. vectensis* [58]. However, only the first ShK domain of NEP3 is retained in the mature

NEP3 toxin, while U-GRTX-Esi4 lacks putative cleavage sites and we found proteomic evidence for the first

and the last ShK domain, suggesting the occurrence of a multidomain protein in *E. singularis*.

Protease inhibitors

Toxins containing domains associated with protease inhibition are widespread across the tree of life. Among

them, Kunitz-like peptides are frequently found in venoms [59] where they can act as inhibitors of several

endogenous and exogenous proteases [60] but also show neurotoxic and neuromodulatory activity on a

variety of receptors and ion channels [61,62], sometimes with multimodal action [57,59]. Kunitz-like toxins

that bind ion channels can be used to paralyze prey and are being extensively studied for their analgesic and

neuroprotective properties [63-67]. Our HMM search revealed the occurrence of two Kunitz-like toxins. PI-

GRTX-Esi5, detectable in the NEM-P only, presented two Kunitz domains separated only by an Asp-Ile

dipeptide, an architecture previously detected in salivary proteins of blood-feeding animals [67,68]. Since

orthologous genes sharing the same 3 exons/2 introns organization of PI-GRTX-Esi5 were not detected in

Scleralcyonacea, this domain combination most likely represents a recent innovation of the

Malacalcyonacea. The second toxin, PI-GRTX-Esi6, with a single Kunitz domain, was detected both in the

NEM-P and in the WB-P, and derived from a precursor with a complex architecture, with the Kunitz-like

domain followed by three Kazal-type domains, all separated by propeptidase cleavage sites. Also in this case,

proteomic evidence indicated a differential maturation pattern of the precursor, with the Kazal-like peptides

from this precursor only retrieved in the WB-P (Suppl. Tab. 2). The Kunitz domain of PI-GRTX-Esi5 presented

the Arg residue (conventionally "P1") associated with the trypsin inhibition, whereas its replacement with a

Met in both domains of PI-GRTX-Esi6 suggests a chymotrypsin specificity [69]. In addition, PI-GRTX-Esi6

presented a Leu residue aligning with the crucial position for K_V1 binding of LmKTT, a scorpion Kunitz-like

toxin [70]. These similarities may hint to multimodal activity for PI-GRTX-Esi5, while the lack of key residue

conservation in PI-GRTX-Esi6 does not allow inferring any putative function (Suppl. Fig. 4)

Other peptides present in *E. singularis* nematocysts displaying protease inhibitory domains (Kazal-type,

cystatines, antistasins, and macins) were detected but did not present novelty elements; they are briefly

described in Suppl. Data.

Colipases

Finally, a striking feature within the NEM-P of *E. singularis* was the abundance of toxins with a predicted

colipase fold (U-GRTX-Esi14-18). Colipases are knottins containing five disulphide bridges, found in spider,

snake and tick venoms [71], but to date not in Cnidaria. The prototypical example, MIT-1 [72] interacts

through its N-terminal pentapeptide AVITG with prokineticin receptors, stimulating smooth muscle

contractions, and exhibiting hyperalgesic effects [73,74]. *E. singularis* colipases lacked the AVITG

pentapeptide, a feature shared with the atracotoxins of the spider *Hadronyche versuta,* whose

pharmacological target is unknown to date [75].

Apart from the most abundantly expressed protein, U-GRTX-Esi14, all colipases derived from modular

precursors containing multiple domains separated by propeptidase cleavage sites. Both U-GRTX-Esi15a-c

and U-GRTX-Esi16 were encoded by genes displaying a five exons/four introns architecture (Fig. 4), well-

conserved in several orthologues from multiple Octocorallia and Hexacorallia species, pointing out a shared

ancient origin with secondary losses in some lineages. U-GRTX-Esi14 and U-GRTX-Esi17 most likely derive

from paralogous gene copies, which underwent structural reorganization during evolution. In detail, U-

GRTX-Esi17 entirely lost exon 2, which resulted in a shorter precursor protein with an incomplete disulphide

pattern. U-GRTX-Esi14 further lost the colipase-fold encoded by exon 3, which was likely reduced and later

fused with exon 1. This scenario is supported by the observations that exon 1 in U-GRTX-Esi14 is significantly

longer than its paralogues and includes a dibasic propeptidase cleavage site usually present at the 3'end of

exon 3. Nevertheless, such remarkable modifications occurred in the latest common ancestor of all

Malacalcyonacea, as evidenced by the identification of U-GRTX-Esi14 and U-GRTX-Esi17 orthologues in *D.*

gigantea, *M. muricata*, *P. clavata*, *Trachythela* sp. and *P. subtilis* (the latter limited to orthologues of U-

GRTX-Esi14).

The loss of exon 2 in the gene encoding for U-GRTX-Esi17 results in a shorter precursor protein, with an N-

terminal incomplete colipase domain. The maturation of this precursor leads to the release of a polypeptide

lacking six out of the ten cysteines (U-GRTX-Esi18), whose presence was also confirmed in the NEM-P. The

lack of most of the cysteine scaffold has a major impact on the protein structure: the model predicted with

good confidence by AlphaFold2 shows in fact only a long β-hairpin flanked by disulphide stabilized tails.

Notably, the transcripts for all these toxins had exceptionally high TPM values (1806, 569, 826 and 429,

respectively for the U-GRTX-Esi14 to 17/18), compared to other proteins detected in the NEM-P (cfr. Suppl.

Tab. 4).

Figure 4. Schematic representation of the evolution of the colipase genes in Anthozoa, with references to specific *E. singularis* genes. Exons are in blue, cysteine residues are indicated by the yellow bars, scissors indicate cleavage sites.

336
337
338

(c) **Novel putative toxins were retrieved in the NEM-P of *Eunicella singularis***

We identified in the NEM-P 10 additional proteins possessing the typical sequence features of toxins (small

size, abundance of Cys, presence of a signal peptide), lacking sequence similarity with any known protein

and often encoded by highly expressed transcripts. Their genomic organization is summarized in Suppl. Fig.

5. High-confidence structural models obtained with Alphafold2 were used to retrieve structural homologues

in the Brookheaven Protein Data Bank, in the attempt to infer their biological functions (Fig.5).

Figure 5. Overview of the structures predicted by AlphaFold2 for the novel toxins from *E. singularis.* Models are coloured according to pLDDT, with higher values (>70) indicating reliable prediction. (A) The putative cytolytic U-GRTX-Esi19 and, on the right, PulS from *K. oxytoca* (PDB 4A56); Overlay of the structures of the two disulphide stabilized helical hairpin toxins U-GRTX-Esi20 (B) and U-GRTX-Esi21 (C) with acrorhagin-1 (PDB 6UX5, right); The predicted structures of the two novel αβ toxins U-GRTX-Esi22 (D) and U-GRTX-Esi23 (E); (F) The SCRiP-like arrangement of U-GRTX-Esi24; (G) The two-domain organization of U-GRTX-Esi25 (left) and its surface charge distribution (right). The ShK domain is adjacent to an all-helical domain. The extended blue patch on the protein surface is common in cytolytic proteins; (H) U-GRTX-Esi26 (left) compared to the byssal protein Pvfp-5β from the mussel *Perna viridis* (PDB 7QAB, right); (I) The structure of U-GRTX-Esi28

Among them, U-GRTX-Esi19 was encoded by an extremely abundant transcript (TPM 2898). The mature

protein (100 amino acids, 6 Cys) was detected in both the NEM-P and WB-P and had a weak similarity at the

C-terminus with Amoebapore A, the major pore-forming protein of *Entamoeba histolytica*, suggestive of a

possible cytolytic function [76]. Most of the protein structure was modeled with high confidence (pLDDT >

80, Fig. 5A), highlighting a compact fold characterized by a knotted four-helix bundle stabilized by three

inter-helices disulphide bridges. A similar arrangement, although devoid of disulphide stabilization, is typical

of bacterial pilotins PulS and GspS, chaperone lipoproteins that play a role in the assembly of the T2SS

secretion system in pathogenic Gram-negative bacteria [77,78]. This putative toxin appears as an

evolutionary innovation of Malacalcyonacea, although orthologues are missing in *Phenganax subtilis* and in

Xenia sp. (Fig. 7, Suppl. Fig. 6). The cysteine framework is always conserved, while the length of some loops

varies across species. The abundance of this protein suggests a strong functional relevance, but activity

assays will be necessary to clarify its function.

The two putative toxins U-GRTX-Esi20 and 21 did not show any obvious sequence similarity, with a pairwise

identity of only 21%. U-GRTX-Esi20 is 48 amino acids long and derives from a longer propeptide

proteolytically cleaved at the C-terminus, whereas U-GRTX-Esi21 displays a shorter sequence with a very

high cysteine density (8 out of 36 amino acids). The models obtained (Fig. 5B and C) highlighted a structural

relationship, as both presented a disulphide-stabilized helical hairpin motif, recently characterized in

acrorhagin-1 (U-AITX-Aeq5a) [79], a toxin from the acrorhagi of *A. equina* used in conspecific aggressive

encounters [80]. Acrorhagin-1 can induce local tissue necrosis and displays toxicity on crabs, but its

mechanism of action has not been clarified, although it has been demonstrated that the toxin does not

interact with either potassium or sodium ion channels and it does not possess membranolytic activity

[79,80]. U-GRTX-Esi20 and 21 are stabilized by three and four disulphide bridges, respectively, which overlap

with those of acrorhagin-1, despite a sequence identity below 20% in both cases.

Figure 6. A) multiple sequence alignment of the complete precursor amino acid sequences of U-GRTX-Esi20, U-GRTX-Esi21, their paralogs, and their orthologs in *E. verrucosa* and *D. gigantea*. **B)** Maximum likelihood phylogeny of the sequences belonging to the same superfamily of U-GRTX-Esi20 and U-GRTX-Esi21 (indicated with orange arrows), identified in different Malacalcyoneacea and Scleralcyonacea species. All Malacalcyonacea sequences (excluding those from *Xenia*) can be classified within six highly supported monophyletic clades. Bootstrap support values are only shown for the major nodes of the tree, and those denoting the basal node of the six aforementioned clades are highlighted with a yellow background.

Our genomic survey indicates that these two toxins belong to two distinct monophyletic orthogroups within

a very large superfamily of cysteine-rich peptides, encoded by ancestrally duplicated paralogous genes with

intronless structures, that also include other members in *E. singularis* transcriptome, not detected in the

NEM-P (Fig. 6). High-confidence orthologues were exclusively found in Malacalcyonacea, except for *P.*

subtilis and *Xenia*, forming 6 well-supported clades (Type I to Type VI) (Fig.6). This suggests a rather recent

origin of these orthogroups, which derive from a large group of similar peptides also found in other

cnidarians, worthy of more extensive investigation. In fact, an ortholog of acrorhagin has been detected in

the sensory neurons of *N. vectensis* [81], suggesting an ancestral recruitment from the nervous to the venom

system for this class of peptides, followed by lineage-specific diversification in the Octocorallia (Fig. 7).

398
399 **Figure 7.** Distribution of orthologues of a subset of *E. singularis* toxins across the Octocorallia genomes here
400 investigated. Phylogenetic relationships of the Octocorallia after [46].
401

The next two putative toxins, U-GRTX-Esi22 (60 amino acids, 10 cysteines, Fig. 5D) and 23 (45 amino acids,

8 cysteines, Fig. 5E) did not present any sequence similarity with known proteins. In both cases, the models

highlighted an unusual disulphide-stabilized αβ fold, suggesting the existence of peculiar and to date

undiscovered cnidarian toxin folds. The gene encoding U-GRTX-Esi23 is present with two similar paralogous

genes in *E. singularis* and has clearly detectable orthologs in all Malacalcyonacea except *P. subtilis* and *Xenia*

sp. (Suppl. Fig. 6, Fig. 7) A divergent paralogue, with a slightly different cysteine pattern, is found in

Scleralcyonacea. This toxin most likely constitutes an evolutionary innovation of Malacalcyonacea but could

derive from an ancestral sequence with a similar cysteine connectivity present in the latest common

ancestor of all Octocorallia. Conversely, U-GRTX-Esi22 presents a single-copy gene, with a detectable

orthologue only in *P. clavata* and *M. muricata,* and is probably a recent innovation, only shared by a few

related orders within Malacalcyonacea (Suppl. Fig. 7, Fig. 7).

The structure of U-GRTX-Esi24, a 45 amino acids protein with 8 cysteines and a 50% sequence identity with

Conotoxin Cltx-2 from *Conus californicus,* could also be modelled with acceptable confidence (Fig. 5F).

Despite undetected by HMM profile analysis, the structure presented a SCRiP-like W-shaped fold. This

domain has a broad but scattered distribution in Octocorallia, suggesting an evolutionary origin slightly older

than most other toxins described in this work. Nevertheless, it is most often found repeated multiple times

within the same protein (also in *E. singularis*), with several independent occurrences of single-domain

peptides scattered across the tree of life (Fig. 7; Suppl. Fig. 8).

Unique case among the novel toxins from *E. singularis,* U-GRTX-Esi25 was predicted to possess a modular

architecture: a recognizable N-terminal ShK domain followed by a region lacking any significant homology.

The two domains were clearly distinguishable in the three-dimensional prediction (Fig. 5G), where they

appeared as directly connected, without spacer sequences or putative cleavage sites. The unknown C-

terminal domain was modelled as an asymmetrical three-helices arrangement, with three disulphide bridges

connecting and stabilizing the secondary structure elements. Interestingly, despite the complex architecture

predicted, several basic residues from both domains in the model aligned perfectly, forming an extended

positively charged patch (Fig. 5G), a feature that suggests a possible membranolytic and/or antimicrobial

action [82]. This peculiar combination of domains, paired with a shared 3 exons/2 introns gene architecture,

was detected in Malacalcyonacea (except *P. subtilis* and *Xenia* sp). A single orthologous gene was found in

Paragorgia papillata (Scleralcyonacea), suggesting that the appearance of this protein could predate the

split between Malacolcyonacea and Scleralcyonacea (Fig. 7; Suppl. Fig. 9).

The model predicted for U-GRTX-Esi26 (77 residues, 12 Cys) presented an elongated structure made of two

domains: an N-terminal region in which two antiparallel beta strands are connected by three disulphide

bridges to a proline-rich stretch, in a defensin-like fold found in other cnidarian toxins, and a C-terminal EGF-

like domain (Fig. 5H). A similar modular structure was recently described in the byssal adhesive protein Pvfp-

5β from the mussel *Perna viridis* [83]; despite the absence of sequence similarity, this structural similarity

might indicate for this protein a structural or adhesive function. However, orthologues of the gene were

found only in Malacalcyonacea (apart from *Dendronephthya* and *Xenia)*, all sharing a common 3 exons/2

introns gene structure. Three divergent paralogues were found In *P. clavata*, probably originated through a

species- or genus-specific gene duplication process (Suppl. Fig. 10).

The last two novel toxins identified in the NEM-P of *E. singularis,* U-GRTX-Esi27 and 28, were short peptides

associated with relatively low transcription levels but a high proteomic coverage. In particular, U-GRTX-Esi27

displayed a weak sequence similarity to both SCRiPs and BDS toxins from *A. sulcata*, which are potent ligands

of K_v3 channels [84]. The mature sequence of U-GRTX-Esi27, a 39 amino acids long peptide, contained 8

cysteines, 6 of which correspond to the BDS framework. An additional cysteine pair occupied the position

of two extremely conserved residues in sea anemone BDS toxins, *i.e.*, a tryptophan and a tyrosine residue,

otherwise present in all other members of the family (Suppl. Fig. 11). Only low confidence models could be

predicted by AlphaFold2, thus preventing any function inference. Despite the similarity with BDS,

orthologous genes encoding for similar proteins were not detected in the genomes of the other Octocorallia

considered in this study except from *E. verrucosa.* U-GRTX-Esi27 might therefore constitute an evolutionary

innovation of the genus *Eunicella* or the family Eunicellidae (Fig. 7). Finally, U-GRTX-Esi28 possessed a weak

sequence homology with the ω-hexatoxin-Hi1g. The modelled structure confirmed this similarity, revealing

a cystine knotted core (Fig. 5I), with the addition of an extended and disordered C-terminus, possibly

proteolitically cleaved. Orthologues coding for this protein are shared by nearly all Malacalcyonacea but are

lacking in *Xenia* (Suppl. Fig. S12). This finding, along with the conservation of the gene structure, in which

two exons are spaced by an intron with an equal placement in all species, the signal peptide sequence, and

the cleavage sites, strongly indicated that this protein represents an evolutionary innovation of

Malacalcyonacea (Fig. 7).

(d) A diverse range of cytolytic proteins are produced both inside and outside the nematocysts

Our analysis highlighted the absence, in the NEM-P of *E. singularis*, of the common cnidarian cytolytic

proteins. A targeted hmmscan search in the WB-P yielded instead positive hits for profiles associated with

both the aerolysin (PF01117) and the full anemone cytotoxin (PF06369) families. Coherently with the

approach adopted for the NEM-P annotation, we considered only the proteins encoded by complete

transcripts presenting a signal peptide; this approach resulted in the identification of three pore forming

toxins (PFTs), characterized by diverse structural features. The first one, Δ-GRTX-Esi29, was an aerolysin-like

protein, structurally related to the β-PFT Hydralysin [85]; the other two, Δ-GRTX-Esi30 and 31, which shared

~50% sequence similarity, belonged to the actinoporins family, the most abundant family of cnidarian porins

[86]. Interestingly, we observed relevant non-conservative substitutions in *E. singularis* actinoporins, which

may indicate selectivity towards different lipids (Suppl. Fig. 13).

Finally, the highest scoring match in the proteomic analysis of the nematocysts content of *E. singularis*

corresponded to a 699 amino-acid translated transcript with homology to prosaposin (pSAP) [87]. In

humans, the maturation of the orthologous pSAP produces four globular saposins (SAP-A/D), which are co-

factors of sphingolipid degrading enzymes. The saposin fold is very ancient and characterizes all the saposin-

like proteins (SAPLIPs), which include the amoebapores [88-90]. SAPs and SAPLIPs have a distinctive cysteine

framework but display minimal conservation (<25% sequence identity). They are often released in several

copies from the same precursor gene [91], assuming a closed, monomeric conformation at lysosomal pH

[88,92]. All these proteins are involved in membrane destabilization/solubilization, acting through different

mechanisms: lysosomal SAPs undergo a conformational change upon pH shift or in the presence of lipids or

detergents, exposing the residues of the hydrophobic core and encapsulating a large number of lipids [93-

95]. Other SAPLIPs, like the amoebapores from *E. histolytica* and the caenopores from *C. elegan*s, form

oligomeric pores [96,97]. Finally, basic SAPLIPs, like NK-lysin and granulysin, act by "molecular

electroporation", inserting within the membranes [98,99]. In rare cases, the saposin fold has also been

detected in non-cytolytic neurotoxins, like the macro-conotoxins con-ikot-ikot and Mu8.1 [100,101], but

these proteins are highly dissimilar from SAPLIPs in terms of sequence and cysteine pattern and are encoded

by single domain (non pSAP-like) transcripts. The six sequences deriving from the maturation of the pSAP

precursor of *E. singularis* (Δ-GRTX-Esi32a-f) were all identified at the proteomic level in the NEM-P. These

proteins shared a 24%-65% similarity, higher than what generally observed for homologous SAPLIPs, and

were more similar to human saposins SAP-A and C (similarity 26-40%) than to antimicrobial porogenic

peptides like amoebapore A (16-25%) or caenopore-5 (13-27%).

We examined the surface potential distribution of the SAPLIPs from *E. singularis*, since the activity of SAPLIPs

correlates to their surface electrostatics [102,103]. Except Δ-GRTX-Esi32a and c, they all shared with human

SAP-A extended patches of negative charge (Fig. 8), which hints to a pH-dependent mechanism for activation

and membrane interaction [104]. In contrast, Δ-GRTX-Esi32a and c presented a more balanced distribution

of basic and acidic regions and extended neutral patches, like caenopore-5, suggesting a different, possibly

non-monomeric, native arrangement and a pH-independent mechanism of action, possibly through pore

formation.

Δ-GRTX-Esi32d   Δ-GRTX-Esi32e   Δ-GRTX-Esi32f

Figure 8. Comparison of the surface electrostatic potentials of the three model saposin-like proteins SAP-A (4UEX),
Amoebapore A (1OF9) and Caenopore-5 (2JSA) with the six SAPLIPs identified in the venom of *E. singularis.* The
extended patches of negative charge (red) observed in Δ-GRTX-Esi32b, d, e, f are observed in human SAP-A and are
indicative of a pH-dependent conformational switch. The lower charge density of Δ-GRTX-Esi32a and c may indicate a
different mode of action*.

Conclusions

We have provided the first preliminary depiction of the venomous arsenal of a widespread octocoral.

Although experimental limitations did not allow us to conduct our analysis on an isolated venom fraction,

by exposing whole polyps of the gorgonian *Eunicella singularis* to ethanol [29] we isolated a sample enriched

in proteins presenting typical toxin features. Among them, we detected several folds previously identified

in cnidarian venoms, some proteins with structural similarities to venom components of unrelated

organisms, and a few with new folds. Even in the absence of functional characterization, the NEM-P seems

to provide a good snapshot of the nematocysts content in *E. singularis,* although we could not rule out the

possibility that these proteins derive from other tissues*.* The scarce similarity observed between octocoral

and hexacoral putative toxins repertoire is not surprising, given the ancient divergence between these two

lineages that recent works dated about 500-700 MYA [2]. In comparison, Elapidae and Viperidae snakes,

which present remarkable differences in their venom composition, have diverged only 50 MYA [105].

However, our results clearly indicate, within the NEM-P of *E. singularis,* an elevated content of putative

neurotoxic peptides, a feature previously reported for sea anemones. Cytolytic toxins with different

structures (and, likely, specificity/function) seem distributed both inside and outside the nematocysts.

The transcriptionally more abundant components of *E. singularis* venom are the potentially cytolytic U-

GRTX-Esi19 and the acrorhagin-like U-GRTX-Esi20, the six colipases, the SCRiP-like U-GRTX-Esi1a and the

SCREP U-GRTX-Esi4 with three ShK domains. With the exception of the "octo-SCRiPs", which, however, are

characterized by sequence peculiarities, these toxins represent evolutionary innovations of Octocorallia,

either shared by the entire lineage (*e.g.,* some colipases), or restricted to the Malacalcyonacea (*e.g.*, U-

GRTX-Esi19), or even exclusive of a group of sublineages within the Malacalcyonacea (*e.g.*, U-GRTX-Esi20)

(Fig.7).

Putative neurotoxins in *E. singularis* included both novel and established cnidarian folds, the latter often

detected within multidomain SCREPs [106]. This is the case, *e.g.,* of the highly expressed U-GRTX-Esi4, with

three ShK domains: multidomain ShK proteins have been so far reported in the transcriptomes of parasitic

roundworms, in *Hydra* and *Nematostella* [58], and in the vampire snails [107]. The toxin U-GRTX-Esi3

represents another case of SCREP, presenting a double SCRiP domain in the mature polypeptide: this

architecture had not been documented in Cnidaria mature toxins to date, although it has been recognized

at the transcript level (e.g. in *Orbicella faveolata*). Two domains were also retrieved in the Kunitz BPTI-like

PI-GRTX-Esi5, an architecture so far investigated in detail in hematophagous invertebrates only, but

extremely abundant among the putative ion-channel impairing SCREP toxins [67,106]. While the abundance

of the SCREPs is most likely underestimated across venomous Metazoa, also due to the paucity of proteomic

studies confirming the architecture of mature toxins, their abundance in the venom of *Eunicella* might

indeed possess some functional significance.

Interestingly, conventional pore forming toxins (PFTs) of Cnidaria, including the aerolysin-like Δ-GRTX-Esi29

and the two actinoporins Δ-GRTX-Esi30 and 31 were not retrieved in the NEM-P. As anticipated, this could

depend on the intrinsic limitations of the venom collection procedure employed or reflect an extra-

nematocystic production of these proteins. Indeed, finding PFTs outside the nematocysts is not unusual:

e.g., hydralysin is an exclusively non-nematocystic PFT involved in prey digestion [84]. Actinoporins, on the

other hand, have been to date detected both in the nematocysts and in other tissues. The extra-

nematocystic localization retrieved in *E. singularis* might suggest that its actinoporins share the digestive

function of the hydralysin-like Δ-GRTX-Esi29. Further histological and functional analysis are needed to

confirm this hypothesis, and to evaluate the impact of the observed mutations at critical sites on chemical

selectivity and pore formation mechanism of *Eunicella* actinoporins.

Among the proteins of the NEM-P, likely representing the nematocyst content, the cytolytic activity seems

to be ascribable mostly to the six saposins Δ-GRTX-Esi32a-f, abundant at the proteomic level. The analysis

of their surface charge hints to a considerable diversification of targets and modes of action for this novel

group of cytolytic polypeptides: most of them might be inactive at the low pH typical of mature nematocysts

but could be activated by the abrupt pH change which follows discharge [108]. On the contrary, the neutral

surface electrostatic of Δ-GRTX-Esi32a and c might suggest a pH-independent activation. The novel toxin U-

GRTX-Esi19, transcribed at high levels and with a fold compatible with a cytolytic activity, might complement

saposins.

While some of these putative toxins are shared by all the investigated octocorals, others had a taxonomic

range restricted to Malacalcoynacea and were not retrieved in the analysed Scleralcyonacea, suggesting

they might constitute evolutionary innovations arisen after the split between the two major Octocorallia

lineages. This is the case of the novel toxins U-GRTX-Esi19 and U-GRTX-Esi28 (Fig. 7). In addition, several

toxins were present in a subset of Malacalcyonacea species and are frequently lacking in the representatives

of two specific families, Xeniidae (or pumping corals) and Acrossotiidae (i.e., tube corals) (Fig. 7). This peculiar taxonomic distribution might suggest a key role in predation for these toxins, since they are absent in species with a mixotrophic habit, in which autotrophy sustained by symbiotic dinoflagellates is the most relevant source of energy. In contrast, the other Malacalcyonacea included in our analysis (e.g., *Paramuricea*, *Trachytela*) are suspension feeders that largely rely on zooplankton intake [109]. Novel toxins, such as U-GRTX-Esi26, U-GRTX-Esi22, the acrorhagin-like U-GRTX-Esi21 and U-GRTX-Esi27 might thus constitute a predation-related group of putative toxins. In particular, U-GRTX-Esi27 appears to be restricted to *Eunicella* as a genus-specific adaptation (Fig. 7).

The inference on the potential function of octocoral putative toxins is, however, heavily affected by the sparse information available on the biology of these organisms. For example, some Octocorallia species possess sweeper and thread-like tentacles, nematocysts-rich structures that have been hypothesized to mediate interspecific competition, similarly to sea anemones acrorhagi [110]. The occurrence of these structures across the Octocorallia radiation has been poorly investigated to date: this lack of comprehensive information hampers the evaluation of a potential role of specific toxins in competition. Nevertheless, although our comparative genomic approach has revealed a powerful tool to restrict the range of potential functions, further functional characterization studies are needed to unequivocally establish the physiological role of these toxins.

Interestingly, our proteomic analysis pointed towards two cases of differential toxin maturation pathways in the nematocyst vs whole body. The first concerns the two SCRiP isotoxins U-GRTX-Esi2a-b, derived from a double-domain precursor, with both isoforms present in the WB-P while only U-GRTX-Esi2b represented in the NEM-P (Suppl. Fig. 1). The second corresponds to the toxin PI-GRTX-Esi6 and the three isotoxins PI-GRTX-Esi7a-c, encoded by a single transcript in which a Kunitz-like domain (retrieved both in the whole body and in the nematocyst proteomes) is followed by three Kazal-type PI domains only identified in the whole-body proteome. The contribution of alternative cleavage sites to venom diversity has been demonstrated in conopeptides [111], but cell- or tissue-specific differential maturation patterns have not been so far documented for venom toxins, suggesting that the mechanisms regulating venom toxin expression could be more complex than previously assessed.

Through our multidisciplinary approach, we have demonstrated that Octocorallia produce a complex

neurotoxin-rich venom, showcasing an overall scarce resemblance to the hexacoral venom repertoire. This

distinctiveness is underscored by the presence of novel folds with a complex evolutionary history. These

findings call for extensive work, focusing both on the biology of this understudied group of organisms and

on the functional characterization of their toxins, to unravel the adaptive significance of Octocorallia arsenal.

Data Availability

The Transcriptome Shotgun Assembly project has been deposited at DDBJ/EMBL/GenBank under the

accession GKVI00000000. The version described in this paper is the first version, GKVI01000000. Proteomic

data have been submitted to PRIDE with reference 1-20240529-113849-2368759. Other data underlying this

article are available in the online supplementary material.

Acknowledgements

This work has received funding from the European Union's Horizon 2020 research and innovation

programme under the Marie Skłodowska-Curie grant agreement No 748902. The MGX acknowledges

financial support from France Génomique National infrastructure, funded as part of "Investissement

d'Avenir" program managed by Agence Nationale pour la Recherche (contract ANR-10-INBS-09). Mass

spectrometry experiments were carried out using the facilities of the Montpellier Proteomics Platform

(PPM, BioCampus Montpellier). Dr. Dany Dominguez-Perez is gratefully acknowledged for his help in data

handling and curation.

References

1. Van Iten H, Marques AC, Leme J, Pacheco MLA, Simões MG. 2014 Origin and early diversification of

the phylum Cnidaria Verrill: Major developments in the analysis of the taxon's proterozoic-cambrian

history. Palaeontology 57, 677–690. (doi:10.1111/pala.12116)

2. McFadden CS, Quattrini AM, Brugler MR, Cowman PF, Dueñas LF, Kitahara M V., Paz-García DA,

Reimer JD, Rodríguez E. 2021 Phylogenomics, Origin, and Diversification of Anthozoans (Phylum

Cnidaria). *Syst Biol* **70**, 635–647. (doi:10.1093/sysbio/syaa103)

3. David CN, Özbek S, Adamczyk P, Meier S, Pauly B, Chapman J, Hwang JS, Gojobori T, Holstein TW.

2008 Evolution of complex structures: minicollagens shape the cnidarian nematocyst. *Trends in*

Genetics **24**, 431–438. (doi:10.1016/j.tig.2008.07.001)

4. Fautin DG. 2009 Structural diversity, systematics, and evolution of cnidae. *Toxicon* **54**, 1054–1064.

(doi:10.1016/j.toxicon.2009.02.024)

5. Kass-Simon G, Scappaticci, Jr. AA. 2002 The behavioral and developmental physiology of

nematocysts. *Can J Zool* **80**, 1772–1794. (doi:10.1139/z02-135)

6. Moran Y, Genikhovich G, Gordon D, Wienkoop S, Zenkert C, Özbek S, Technau U, Gurevitz M. 2012

Neurotoxin localization to ectodermal gland cells uncovers an alternative mechanism of venom

delivery in sea anemones. *Proc. Royal Soc B* **279**, 1351–1358. (doi:10.1098/rspb.2011.1731)

7. Marques AC, Collins AG. 2004 Cladistic analysis of Medusozoa and cnidarian evolution. *Inv Biol* **123**,

23–42. (doi:10.1111/j.1744-7410.2004.tb00139.x)

8. Kayal E, Bentlage B, Sabrina Pankey M, Ohdera AH, Medina M, Plachetzki DC, Collins AG, Ryan JF.

2018 Phylogenomics provides a robust topology of the major cnidarian lineages and insights on the

origins of key organismal traits. *BMC Evol Biol* **18**, 68. (doi:10.1186/s12862-018-1142-0)

9. Norton RS. 1991 Structure and structure-function relationships of sea anemone proteins that

interact with the sodium channel. *Toxicon* **29**, 1051–1084. (doi:10.1016/0041-0101(91)90205-6)

10. Norton RS. 2009 Structures of sea anemone toxins. *Toxicon* **54**, 1075–1088.

(doi:10.1016/j.toxicon.2009.02.035)

11. Honma T, Shiomi K. 2006 Peptide toxins in sea anemones: Structural and functional aspects. *Mar*

Biotech **8**, 1–10. (doi:10.1007/s10126-005-5093-2)

12. Shiomi K. 2009 Novel peptide toxins recently isolated from sea anemones. *Toxicon* **54**, 1112–1118.

(doi:10.1016/j.toxicon.2009.02.031)

13. Jouiaei M, Yanagihara AA, Madio B, Nevalainen TJ, Alewood PF, Fry BG. 2015 Ancient venom

systems: A review on Cnidaria toxins. *Toxins* **7**, 2251–2271. (doi:10.3390/toxins7062251)

14. Madio B, King GF, Undheim EAB. 2019 Sea anemone toxins: A structural overview. *Mar Drugs* **17**,

325. (doi:10.3390/md17060325)

15. Menezes C, Thakur NL. 2022 Sea anemone venom: Ecological interactions and bioactive potential.

Toxicon **208**, 31–46. (doi:10.1016/j.toxicon.2022.01.004)

16. Prentis PJ, Pavasovic A, Norton RS. 2018 Sea anemones: Quiet achievers in the field of peptide

toxins. *Toxins* **10**, 36. (doi:10.3390/toxins10010036)

17. Rachamim T, Morgenstern D, Aharonovich D, Brekhman V, Lotan T, Sher D. 2015 The dynamically

evolving nematocyst content of an Anthozoan, a scyphozoan, and a hydrozoan. *Mol Biol Evol* **32**,

740–753. (doi:10.1093/molbev/msu335)

18. Pennington MW, Czerwinski A, Norton RS. 2018 Peptide therapeutics from venom: Current status

and potential. *Bioorg Med Chem* **26**, 2738–2758. (doi:10.1016/j.bmc.2017.09.029)

19. Schmidt CA, Daly NL, Wilson DT. 2019 Coral venom toxins. *Front Ecol Evol* **7**.

(doi:10.3389/fevo.2019.00320)

20. Klompen AML, Macrander J, Reitzel AM, Stampar SN. 2020 Transcriptomic analysis of four cerianthid

(Cnidaria, Ceriantharia) venoms. *Mar Drugs* **18**, 413. (doi:10.3390/MD18080413)

21. Yoffe C, Lotan T, Benayhau Y. 2012 A modified view on octocorals: *Heteroxenia fuscescens*

nematocysts are diverse, featuring both an ancestral and a novel type. *PLoS One* **7**, 1–5.

(doi:10.1371/journal.pone.0031902)

22. Raimundo I, Silva SG, Costa R, Keller-Costa T. 2018 Bioactive secondary metabolites from octocoral-

associated microbes. New chances for blue growth. *Mar Drugs* **16**, 485. (doi:10.3390/md16120485)

23. Ballestreros E. 2006 Mediterranean coralligenous assemblages: A synthesis of present knowledge.

In *Oceanography and marine biology: An annual review*, pp. 123–195.

24. Linares C, Coma R, Garrabou J, Díaz D, Zabala M. 2008 Size distribution, density and disturbance in

two Mediterranean gorgonians: *Paramuricea clavata* and *Eunicella singularis*. *J Appl Ecol* **45**, 688–

699. (doi:10.1111/j.1365-2664.2007.01419.x)

25. Gori A, Rossi S, Berganzo E, Pretus JL, Dale MRT, Gili JM. 2011 Spatial distribution patterns of the

gorgonians *Eunicella singularis*, *Paramuricea clavata*, and *Leptogorgia sarmentosa* (Cape of Creus,

Northwestern Mediterranean Sea). *Mar Biol* **158**, 143–158. (doi:10.1007/s00227-010-1548-8)

26. Coma R, Llorente-Llurba E, Serrano E, Gili JM, Ribes M. 2015 Natural heterotrophic feeding by a

temperate octocoral with symbiotic zooxanthellae: a contribution to understanding the

mechanisms of die-off events. *Coral Reefs* **34**, 549–560. (doi:10.1007/s00338-015-1281-3)

27. Renzi JJ, Shaver EC, Burkepile DE, Silliman BR. 2022 The role of predators in coral disease dynamics.

Coral Reefs **41**, 405–422. (doi:10.1007/s00338-022-02219-w)

28. Radwan FFY, Aboul-Dahab HM, Burnett JW. 2002 Some toxicological characteristics of three

venomous soft corals from the Red Sea. *Comp Biochem Physiol C* **132**, 25–35. (doi:10.1016/S1532-

0456(02)00045-5)

29. Jouiaei M *et al.* 2015 Firing the sting: Chemically induced discharge of cnidae reveals novel proteins

and peptides from box jellyfish (*Chironex fleckeri*) Venom. *Toxins* **7**, 936–950.

(doi:10.3390/toxins7030936)

30. Grabherr MG *et al.* 2011 Full-length transcriptome assembly from RNA-Seq data without a reference

genome. *Nat Biotechnol* **29**, 644–652. (doi:10.1038/nbt.1883)

31. Wagner GP, Kin K, Lynch VJ. 2012 Measurement of mRNA abundance using RNA-seq data: RPKM

measure is inconsistent among samples. *Theor Biosci* **131**, 281–285. (doi:10.1007/s12064-012-0162-

3)

32. Ma B, Zhang K, Hendrie C, Liang C, Li M, Doherty-Kirby A, Lajoie G. 2003 PEAKS: Powerful software

for peptide de novo sequencing by tandem mass spectrometry. *Rapid Commun Mass Spectr* **17**,

2337–2342. (doi:10.1002/rcm.1196)

33. Zhang J *et al.* 2012 PEAKS DB: de novo sequencing assisted database search for sensitive and

accurate peptide identification. *Mol Cell Proteomics* **11**, M111.010587.

(doi:10.1074/mcp.M111.010587)

34. Han Y, Ma B, Zhang K. 2005 SPIDER: software for protein identification from sequence tags with de

novo sequencing error. *J Bioinform Comput Biol* **3**, 697–716. (doi:10.1142/s0219720005001247)

35. Degueldre M et al. 2017 Diversity in sequences, post-translational modifications and expected

pharmacological activities of toxins from four *Conus* species revealed by the combination of cutting-

edge proteomics, transcriptomics and bioinformatics. *Toxicon* **130**, 116–125.

(doi.org/10.1016/j.toxicon.2017.02.014)

36. Huang Y, Niu B, Gao Y, Fu L, Li W. 2010 CD-HIT Suite: A web server for clustering and comparing

biological sequences. *Bioinformatics* **26**, 680–682. (doi:10.1093/bioinformatics/btq003)

37. Oliveira JS, Fuentes-Silva D, King GF. 2012 Development of a rational nomenclature for naming

peptide and protein toxins from sea anemones. *Toxicon* **60**, 539–550.

(doi:10.1016/j.toxicon.2012.05.020)

38. Jumper J *et al.* 2021 Highly accurate protein structure prediction with AlphaFold. *Nature* **596**, 583–

589. (doi:10.1038/s41586-021-03819-2)

39. Mirdita M, Schütze K, Moriwaki Y, Heo L, Ovchinnikov S, Steinegger M. 2022 ColabFold: making

protein folding accessible to all. *Nat Methods* **19**, 679–682. (doi:10.1038/s41592-022-01488-1)

40. Tunyasuvunakool K *et al.* 2021 Highly accurate protein structure prediction for the human

proteome. *Nature* **596**, 590–596. (doi:10.1038/s41586-021-03828-1)

41. Holm L. 2022 Dali server: structural unification of protein families. *Nucleic Acids Res* **50**, W210–

W215. (doi:10.1093/nar/gkac387)

42. Ayoub R, Lee Y. 2019 Rupee: A fast and accurate purely geometric protein structure search. *PLoS*

One **14**, e0213712. (doi:10.1371/journal.pone.0213712)

43. Webb B, Sali A. 2016 Comparative protein structure modeling using MODELLER. *Curr Protoc*

Bioinformatics **2016**, 5.6.1-5.6.37. (doi:10.1002/cpbi.3)

44. Jurrus E *et al.* 2018 Improvements to the APBS biomolecular solvation software suite. *Protein*

Science **27**, 112–128. (doi:10.1002/pro.3280)

45. Meng EC, Goddard TD, Pettersen EF, Couch GS, Pearson ZJ, Morris JH, Ferrin TE. 2023 UCSF

ChimeraX: Tools for structure building and analysis. *Protein Science* **32**, e4792.

(doi:10.1002/pro.4792)

46. McFadden CS, Van Ofwegen LP, Quattrini AM. 2022 Revisionary systematics of Octocorallia

(Cnidaria: Anthozoa) guided by phylogenomics. *Bull Soc Syst Biol* **1**. (doi:10.18061/bssb.v1i3.8735)

47. Gertz EM, Yu Y-K, Agarwala R, Schäffer AA, Altschul SF. 2006 Composition-based statistics and

translated nucleotide searches: Improving the TBLASTN module of BLAST. *BMC Biol* **4**, 41.

(doi:10.1186/1741-7007-4-41)

48. Reese MG. 1997 Improved splice site detection in Genie. *J Comput Biol* **4**, 311–323.
(doi:10.1089/cmb.1997.4.311)

49. Minh BQ, Schmidt HA, Chernomor O, Schrempf D, Woodhams MD, von Haeseler A, Lanfear R. IQ-
TREE 2: New Models and Efficient Methods for Phylogenetic Inference in the Genomic Era. 2020.
Mol Biol Evol. **37**, 1530-1534 (doi: 10.1093/molbev/msaa015).

50. Kalyaanamoorthy S, Minh BQ, Wong TKF, Von Haeseler A, Jermiin LS. 2017 ModelFinder: Fast model
selection   for   accurate   phylogenetic   estimates.   *Nat   Methods*   **14**,   587–589.
(doi:10.1038/nmeth.4285)

51. Sachkova MY, Singer SA, Macrander J, Reitzel AM, Peigneur S, Tytgat J, et al. 2019. The Birth and
Death of Toxins with Distinct Functions: A Case Study in the Sea Anemone Nematostella. *Mol Biol*
Evol **36**, 2001. (doi: 10.1093/molbev/msz132)

52. Jouiaei M, Sunagar K, Federman Gross A, Scheib H, Alewood PF, Moran Y, Fry BG. 2015. Evolution of
an ancient venom: recognition of a novel family of cnidarian toxins and the common evolutionary
origin of sodium and potassium neurotoxins in sea anemone. *Mol Biol Evol* **32**, 1598-610. (doi:
10.1093/molbev/msv050)

53. Sunagawa S, DeSalvo MK, Voolstra CR, Reyes-Bermudez A, Medina M. 2009 Identification and gene
expression analysis of a taxonomically restricted cysteine-rich protein family in reef-building corals.
PLoS One **4**, e4865. (doi:10.1371/journal.pone.0004865)

54. Logashina YA *et al.* 2017 New disulfide-stabilized fold provides sea anemone peptide to exhibit both
antimicrobial and TRPA1 potentiating properties. *Toxins* **9**, 154. (doi:10.3390/toxins9050154)

55. Barroso RA, Ramos L, Moreno H, Antunes A. 2024 Evolutionary Analysis of Cnidaria Small Cysteine-
Rich Proteins (SCRiPs), an Enigmatic Neurotoxin Family from Stony Corals and Sea Anemones
(Anthozoa: Hexacorallia). *Toxins* **16**, 75. (doi:10.3390/toxins16020075)

56. Kozlov SA, Grishin EV. 2007. The universal algorithm of maturation for secretory and excretory
protein precursors. *Toxicon* **49**, 721–6.

57. Pennington MW, Czerwinski A, Norton RS. 2018. Peptide therapeutics from venom: Current status

and potential. *Bioorg. Med. Chem.* 26:2738–2758.

58. Columbus-Shenkar YY, Sachkova MY, Macrander J, Fridrich A, Modepalli V, Reitzel AM, Sunagar K, Moran Y. 2018. Dynamics of venom composition across a complex life cycle**. *Elife* **9,** e35014. (doi: 10.7554/eLife.35014)

59. Sintsova O *et al.* 2021 Sea anemone kunitz-type peptides demonstrate neuroprotective activity in the 6-hydroxydopamine induced neurotoxicity model. *Biomedicines* **9**, 283. (doi:10.3390/biomedicines9030283)

60. Peigneur S, Billen B, Derua R, Waelkens E, Debaveye S, Béress L, Tytgat J. 2011 A bifunctional sea anemone peptide with Kunitz type protease and potassium channel inhibiting properties. *Biochem Pharmacol* **82**, 81–90. (doi:10.1016/j.bcp.2011.03.023)

61. Nikolaev M V., Dorofeeva NA, Komarova MS, Korolkova Y V., Andreev YA, Mosharova I V., Grishin E V., Tikhonov DB, Kozlov SA. 2017 TRPV1 activation power can switch an action mode for its polypeptide ligands. *PLoS One* **12**, e0177077. (doi:10.1371/journal.pone.0177077)

62. García-Fernández R, Peigneur S, Pons T, Alvarez C, González L, Chávez MA, Tytgat J. 2016 The kunitz-type protein ShPI-1 inhibits serine proteases and voltage-gated potassium channels. *Toxins* **8**, 110. (doi:10.3390/toxins8040110)

63. Andreev YA *et al.* 2013 Polypeptide modulators of TRPV1 produce analgesia without hyperthermia. *Mar Drugs* **11**, 5100–5115. (doi:10.3390/md11125100)

64. Wang X *et al.* 2021 Toxic Peptide From *Palythoa caribaeorum* Acting on the TRPV1 Channel Prevents Pentylenetetrazol-Induced Epilepsy in Zebrafish Larvae. *Front Pharmacol* **12**, 763089. (doi:10.3389/fphar.2021.763089)

65. Hwang SM, Jo YY, Cohen CF, Kim YH, Berta T, Park CK. 2022 Venom Peptide Toxins Targeting the Outer Pore Region of Transient Receptor Potential Vanilloid 1 in Pain: Implications for Analgesic Drug Development. *Int J Mol Sci* **23**, 5772. (doi:10.3390/ijms23105772)

66. Kvetkina A *et al.* 2022 Kunitz-Type Peptides from Sea Anemones Protect Neuronal Cells against Parkinson's Disease Inductors via Inhibition of ROS Production and ATP-Induced P2X7 Receptor Activation. *Int J Mol Sci* **23**, 5115. (doi:10.3390/ijms23095115)

67. Dai SX, Zhang A Di, Huang JF. 2012 Evolution, expansion and expression of the Kunitz/BPTI gene

family associated with long-term blood feeding in *Ixodes scapularis*. *BMC Evol Biol* **12**, 4.

(doi:10.1186/1471-2148-12-4)

68. Modica MV, Lombardo F, Franchini P, Oliverio M. 2015 The venomous cocktail of the vampire snail

Colubraria reticulata (Mollusca, Gastropoda). *BMC Genomics* **16**, 441. (doi:10.1186/s12864-015-

1648-4)

69. Mishra M. 2020 Evolutionary Aspects of the Structural Convergence and Functional Diversification

of Kunitz-Domain Inhibitors. *J Mol Evol* **88**, 537–548. (doi:10.1007/s00239-020-09959-9)

70. Chen Z *et al.* 2013 Genomic and Structural Characterization of Kunitz-Type Peptide LmKTT-1a

Highlights Diversity and Evolution of Scorpion Potassium Channel Toxins. *PLoS One* **8**, e60201.

(doi:10.1371/journal.pone.0060201)

71. Fry BG *et al.* 2009 The Toxicogenomic Multiverse: Convergent Recruitment of Proteins Into Animal

Venoms.   *Annu   Rev   Genomics   Hum   Genet*   **10**,   483–511.

(doi:10.1146/annurev.genom.9.081307.164356)

72. Boisbouvier J, Albrand J-P, Blackledge M, Jaquinod M, Schweitz H, Lazdunski M, Marion D. 1998 A

structural homologue of colipase in black mamba venom revealed by NMR floating disulphide bridge

analysis. *J Mol Biol* **283**, 205–219. (doi:10.1006/jmbi.1998.2057)

73. Kaser A, Winklmayr M, Lepperdinger G, Kreil G. 2003 The AVIT protein family. *EMBO Rep* **4**, 469–

473. (doi:10.1038/sj.embor.embor830)

74. Negri L, Lattanzi R, Giannini E, Melchiorri P. 2007 Bv8/Prokineticin proteins and their receptors. *Life

Sci* **81**, 1103–1116. (doi:10.1016/j.lfs.2007.08.011)

75. Wen S *et al.* 2005 Discovery of an MIT-like atracotoxin family: Spider venom peptides that share

sequence homology but not pharmacological properties with AVIT family proteins. *Peptides* **26**,

2412–2426. (doi:10.1016/j.peptides.2005.05.012)

76. Hecht O, van Nuland NA, Schleinkofer K, Dingley AJ, Bruhn H, Leippe M, Grötzinger J. 2004 Solution

Structure of the Pore-forming Protein of *Entamoeba histolytica*. *J Biol Chem* **279**, 17834–17841.

(doi:10.1074/jbc.M312978200)

77. Tosi T, Nickerson NN, Mollica L, Jensen MR, Blackledge M, Baron B, England P, Pugsley AP, Dessen

A. 2011 Pilotin-secretin recognition in the type II secretion system of *Klebsiella oxytoca*. *Mol*

Microbiol **82**, 1422–1432. (doi:10.1111/j.1365-2958.2011.07896.x)

78. Korotkov K V., Hol WGJ. 2013 Crystal structure of the pilotin from the enterohemorrhagic

Escherichia coli type II secretion system. *J Struct Biol* **182**, 186–191. (doi:10.1016/j.jsb.2013.02.013)

79. Krishnarjuna B, Sunanda P, Villegas-Moreno J, Csoti A, A.V. Morales R, Wai DCC, Panyi G, Prentis P,

Norton RS. 2021 A disulfide-stabilised helical hairpin fold in acrorhagin I: An emerging structural

motif in peptide toxins. *J Struct Biol* **213**, 107692. (doi:10.1016/j.jsb.2020.107692)

80. Honma T, Minagawa S, Nagai H, Ishida M, Nagashima Y, Shiomi K. 2005 Novel peptide toxins from

acrorhagi, aggressive organs of the sea anemone *Actinia equina*. *Toxicon* **46**, 768–774.

(doi:10.1016/j.toxicon.2005.08.003)

81. Sachkova MY, Landau M, Surm JM, Macrander J, Singer SA, Reitzel AM, et al. 2020. Toxin-like

neuropeptides in the sea anemone *Nematostella* unravel recruitment from the nervous system to

venom. *PNAS* **117**, 27481–92. (doi: 10.1073/pnas.2011120117)

82. Anderluh G, Lakey JH. 2008 Disparate proteins use similar architectures to damage membranes.

Trends Biochem Sci **33**, 482–490. (doi:10.1016/j.tibs.2008.07.004)

83. Morando MA, Venturella F, Sollazzo M, Monaca E, Sabbatella R, Vetri V, Passantino R, Pastore A,

Alfano C. 2022 Solution structure of recombinant Pvfp-5β reveals insights into mussel adhesion.

Commun Biol **5**, 1–10. (doi:10.1038/s42003-022-03699-w)

84. Yeung SYM, Thompson D, Wang Z, Fedida D, Robertson B. 2005 Modulation of Kv3 subfamily

potassium currents by the sea anemone toxin BDS: Significance for CNS and biophysical studies. *J*

Neurosci **25**, 8735–8745. (doi:10.1523/JNEUROSCI.2119-05.2005)

85. Sher D, Fishman Y, Melamed-Book N, Zhang M, Zlotkin E. 2008 Osmotically driven prey

disintegration in the gastrovascular cavity of the green hydra by a pore-forming protein. *The FASEB*

Journal **22**, 207–214. (doi:10.1096/fj.07-9133com)

86. Macrander J, Daly M. 2016 Evolution of the cytolytic pore-forming proteins (Actinoporins) in sea

anemones. *Toxins* **8**, 1–16. (doi:10.3390/toxins8120368)

87. Darmoise A, Maschmeyer P, Winau F. 2010 The Immunological Functions of Saposins. *Adv Immunol*
105, 25–62. (doi:10.1016/S0065-2776(10)05002-9)

88. Ahn VE, Leyko P, Alattia J-R, Chen L, Privé GG. 2006 Crystal structures of saposins A and C. *Protein*
Sci **15**, 1849–1857. (doi:10.1110/ps.062256606)

89. Munford RS, Sheppard PO, O'Hara PJ. 1995 Saposin-like proteins (SAPLIP) carry out diverse functions
on a common backbone structure. *J Lipid Res* **36**, 1653–1663.

90. Bruhn H. 2005 A short guided tour through functional and structural features of saposin-like
proteins. *Biochem J* **389**, 249–257. (doi:10.1042/BJ20050051)

91. Herbst R, Marciano-Cabral F, Leippe M. 2004 Antimicrobial and pore-forming peptides of free-living
and potentially highly pathogenic *Naegleria fowleri* are released from the same precursor molecule.
Journal of Biological Chemistry **279**, 25955–25958. (doi:10.1074/jbc.M401965200)

92. Hill CH, Read RJ, Deane JE. 2015 Structure of human saposin A at lysosomal pH. *Acta Crystallogr.*
Sect F **71**, 895–900. (doi:10.1107/S2053230X15008584)

93. Hawkins CA, De Alba E, Tjandra N. 2005 Solution structure of human saposin C in a detergent
environment. *J Mol Biol* **346**, 1381–1392. (doi:10.1016/j.jmb.2004.12.045)

94. Alattia JR, Shaw JE, Yip CM, Privé GG. 2006 Direct Visualization of Saposin Remodelling of Lipid
Bilayers. *J Mol Biol* **362**, 943–953. (doi:10.1016/j.jmb.2006.08.009)

95. Popovic K, Holyoake J, Pomès R, Privé GG. 2012 Structure of saposin A lipoprotein discs. *PNAS* **109**,
2908–2912. (doi:10.1073/pnas.1115743109)

96. Leippe M, Bruhn H, Hecht O, Grotzinger J. 2005 Ancient weapons: the three-dimensional structure
of amoebapore A. *Trends Parasitol* **21**, 5–7. (doi:10.1016/j.pt.2004.10.009)

97. Mysliwy J, Dingley AJ, Stanisak M, Jung S, Lorenzen I, Roeder T, Leippe M. 2010 Caenopore-5: The
three-dimensional structure of an antimicrobial protein from. *Dev Comp Immunol* **34**, 323-30 (doi:
10.1016/j.dci.2009.11.00

98. Liepinsh E, Andersson M, Ruysschaert JM, Otting G. 1997 Saposin fold revealed by the NMR
structure of NK-lysin. *Nat Struct Biol* **4**, 793–795. (doi:10.1038/nsb1097-793)

99. Anderson DH, Sawaya MR, Cascio D, Ernst W, Modlin R, Krensky A, Eisenberg D. 2003 Granulysin

Crystal Structure and a Structure-derived Lytic Mechanism. *J Mol Biol* **325**, 355–365.

(doi:10.1016/S0022-2836(02)01234-2)

100.   Chen L, Dürr KL, Gouaux E. 2014 X-ray structures of AMPA receptor-cone snail toxin

complexes illuminate activation mechanism. *Science* **345**, 1021–1026.

(doi:10.1126/science.1258409)

101.   Hackney CM *et al.* 2023 A previously unrecognized superfamily of macro-conotoxins

includes an inhibitor of the sensory neuron calcium channel Cav2.3. *PLoS Biol* **21**, e3002217.

(doi:10.1371/journal.pbio.3002217)

102. Olmeda B, García-Álvarez B, Pérez-Gil J. 2013 Structure-function correlations of pulmonary

surfactant protein SP-B and the saposin-like family of proteins. EBJ 42, 209–222.

(doi:10.1007/s00249-012-0858-9)

103. Olmeda B, García-Álvarez B, Gómez MJ, Martínez-Calle M, Cruz A, Pérez-Gil J. 2015 A model for

the structure and mechanism of action of pulmonary surfactant protein B. *FASEB Journal* **29**, 4236–

4247. (doi:10.1096/fj.15-273458)

104. Garrido-Arandia M, Cuevas-Zuviría B, Díaz-Perales A, Pacios LF. 2018 A comparative study of

human saposins. *Molecules* **23**, 422. (doi:10.3390/molecules23020422)

105. Yin W *et al.* 2016 Evolutionary trajectories of snake genes and genomes revealed by comparative

analyses of five-pacer viper. *Nat Commun* **7**, 13107. (doi:10.1038/ncomms13107)

106. Maxwell M, Undheim EAB, Mobli M. 2018 Secreted cysteine-rich repeat proteins 'SCREPs': A novel

multi-domain architecture. *Front Pharmacol* **9**, 1333. (doi:10.3389/fphar.2018.01333)

107. Gerdol M, Cervelli M, Mariottini P, Oliverio M, Dutertre S, Modica MV. 2019 A recurrent motif:

Diversity and evolution of ShKT domain containing proteins in the vampire snail *Cumia reticulata*.

Toxins **11**. (doi:10.3390/toxins11020106)

108. Berking S, Herrmann K. 2006 Formation and discharge of nematocysts is controlled by a proton

gradient across the cyst membrane. *Helgol Mar Res* **60**, 180–188. (doi:10.1007/s10152-005-0019-

y)

109. Fabricius KE, Dommisse M. 2000. Depletion of suspended particulated matter over coastal reef

communities dominated by zooxanthellate soft corals. *MEPS* **196**, 157-167

(doi:10.3354/meps196157)

110. Sebens KP, Miles JS. 1988 Sweeper Tentacles in a Gorgonian Octocoral: Morphological

Modifications for Interference Competition. *Biol Bull* **175**, 378–387. (doi:10.2307/1541729)

111. Dutertre S, Jin AH, Kaas Q, Jones A, Alewood PF, Lewis RJ. 2013 Deep venomics reveals the

mechanism for expanded peptide diversity in cone snail venom. *Mol Cell Proteomics* **12**, 312–329.

(doi:10.1074/mcp.M112.021469)