Early development of the glucocorticoid stress response in dyeing poison frog tadpoles

Lisa L. Surber-Cunningham [a*], Lucas S. Jimenez [a], Lauren W. Mobo [b], Sarah E. Westrick [a,c], Eva

K. Fischer [a,d]

[a] Department of Evolution, Ecology, and Behavior; School of Integrative Biology, University

of Illinois Urbana-Champaign, Urbana IL USA

[b] Department of Natural Sciences, Parkland College, Champaign IL USA

[c] Department of Biology, Wake Forest University, Winston-Salem, NC USA

[d] Department of Neurobiology, Physiology, and Behavior, University of California Davis, Davis

CA USA

*Corresponding author:

Lisa L. Surber-Cunningham

Department of Evolution, Ecology, and Behavior

School of Integrative Biology

University of Illinois Urbana-Champaign

505 S Goodwin Ave

Urbana, IL 61801

lsurber2@illinois.edu

+1 (707) 508-5083

Abstract

In vertebrates, the glucocorticoid "stress" response (corticosterone or cortisol) through the hypothalamic-pituitary-adrenal (HPA) axis influences many essential functions, including behavior, metabolism, immunity, and ontogenetic transitions. During development, stress responses can be adaptive if they facilitate antipredator behavior and modulate developmental speed to adjust to environmental conditions; however, these same responses can be maladaptive when energetic costs become too high and developmental speed trades-off with size and health at maturity. Thus, the timing of HPA-axis development may be aligned with specific developmental challenges and opportunities presented by a species' life history strategy. In anurans (frogs and toads), corticosterone plays critical roles in development and behavior, and concentrations can fluctuate in response to environmental stressors. Given the role of corticosterone in ontogenetic changes and behaviors, we studied the development of the HPA axis in tadpoles of the dyeing poison frog (*Dendrobates tinctorius*), a species with a unique life history that includes transport to water after hatching on land and aggressive and cannibalistic behavior. We measured the excretion rate and whole-body concentration of corticosterone and the corticosterone response to adrenocorticotropic hormone (ACTH) in free-swimming tadpoles after transport and throughout metamorphosis. We found no significant differences across development in excretion rates or whole-body concentration of corticosterone, nor corticosterone response to ACTH, indicating that that the glucocorticoid response develops early in ontogeny. This pattern differs from those in other species of tadpoles, suggesting the unique ecological pressures faced by *D. tinctorius* have shaped the development of its HPA axis. More broadly, this study illustrates how life history strategies and tradeoffs impact the timing of the HPA axis development.

Keywords: development, ACTH challenge, anurans, larvae, HPA

Introduction

Across development an essential challenge for organisms is to detect and respond to environmental challenges. In vertebrates, these responses are mediated in part via glucocorticoid "stress" hormones that regulate many essential functions, including behavior, metabolism, immunity and ontogenetic transitions (Wada 2008, reviewed in Carr, 2011 and Landys et al., 2006). Given their wide-spread effects, glucocorticoids are also essential in coordinating phenotypic responses to stressors (Dantzer, 2023; Hau et al., 2016). Though the stress response is essential, chronically elevated glucocorticoids can cause problems including decreased immune function (Charmandari et al., 2005), elevated heart rate and blood pressure (Charmandari et al., 2005), negatively impacted growth rate (Glennemeier & Denver, 2002) and compromised cognitive ability later in life (Kitaysky et al., 2003). At an evolutionary timescale, the need to balance essential functions with negative effects leads to variation in when physiological stress responses develop (Wada, 2008).

Selection may mitigate tradeoffs between stress responsivity and hyperactivity by changing when the stress response develops (Wada, 2008). Many altricial species have minimal glucocorticoid reactivity early in life, a pattern thought to facilitate faster growth rates. For example, mice undergo a Stress Hypo-Responsive Period (SHRP) at four to fourteen days post birth, in which individuals have low basal stress hormone levels and an inability to mount a stress response to a mild stressor (Schmidt et al., 2002). However, in species with independent and active young, an early developed stress response may be crucial despite potential negative

side effects. For example, precocial birds like turkeys show significant glucocorticoid activity

from the day they hatch (Davis & Siopes, 1985), likely to facilitate immediate antipredator

behaviors from their ground nests (Wada, 2008). In general, there is a balance between the

timing and abundance of glucocorticoid production associated with life-history characteristics,

developmental timing, and survival strategies across species.

In vertebrates, the neuroendocrine stress response is primarily mediated by the

hypothalamic-pituitary adrenal (HPA) axis (Carr, 2011; Dantzer, 2023) or hypothalamic-

pituitary-interrenal axis (HPI) in amphibians and fish (Denver, 2009a). When the HPA axis is

activated, the hypothalamus releases corticotropin releasing hormone (CRH) that induces the

pituitary gland (or anterior pituitary (Carr, 2011)) to secrete adrenocorticotropic hormone

(ACTH). ACTH is transported to the adrenal cortex, which then releases glucocorticoid steroid

hormones (Sapolsky et al., 1986). The glucocorticoid hormones cortisol and corticosterone are

well-studied and – despite their many functions – are commonly referred to simply as 'stress

hormones.' Though cortisol and corticosterone are generally considered functionally

interchangeable, which glucocorticoid is dominant varies across species (Sapolsky et al., 2000).

The role of glucocorticoids in development is particularly interesting in the unique life-

history of amphibians where they play a key role in metamorphosis (Denver, 1993, 1998,

2009b), and developmental trade-offs in response to environmental stressors are well-studied

(McDiarmid & Altig, 1999). In many species of larval anurans (tadpoles of frogs and toads),

corticosterone levels increase throughout development, peaking at metamorphosis (Jaudet &

Hatey, 1984; Krug et al., 1983). These developmental shifts in corticosterone are in addition to

fluctuations in response to a variety of well-studied environmental stressors that tadpoles face,

including predation, competition, and habitat desiccation (Bókony et al., 2021; Denver, 1998;

Forsburg et al., 2019a; Ledón-Rettig et al., 2023; Ledón-Rettig & Lagon, 2021). Acute changes

in corticosterone impact tadpole antipredator behavior (Fraker et al., 2009; Kulkarni &

Gramapurohit, 2017), morphology (Glennemeier & Denver, 2002; Hossie et al., 2010), and

increase developmental speed (Denver, 1993, 1998, 2009b). Elevated corticosterone can increase

developmental rate to rapidly escape a drying or dangerous pond (Pfennig et al., 1991), but rapid

metamorphosis can have a negative impact on growth and survivorship later in life (Francesco

Ficetola & De Bernardi, 2006). Thus, the unique life-histories and diverse environments of

tadpoles highlight the potential for glucocorticoid mediated developmental tradeoffs.

We studied HPA axis activity in tadpoles of the dyeing poison frog (*Dendrobates*

tinctorius), a neotropical anuran species with several unique life history traits. Despite the

variation in life histories, environmental pressures, and developmental rates across tadpole

species, glucocorticoids and HPA responsivity have largely been studied in temperate, rapidly

developing species. Compared to most temperature frog species that lay hundreds to thousands

of eggs in the water and provide no care to eggs or offspring, adult *D. tinctorius* lay three to eight

terrestrial eggs and provide parental care. Males clean, hydrate, and defend their eggs, and after

hatching carry tadpoles "piggyback" to pools of water in fallen trees, trees with buttresses, and

fallen palm bracts (Rojas, 2014; Rojas & Pašukonis, 2019). Once this transition from land to

water is complete, tadpoles are independent from their parents and faced with predators,

competition for resources, disease, habitat desiccation, and aggressive and cannibalistic

conspecifics (Fischer et al., 2020; Rojas & Pašukonis, 2019). *D. tinctorius* tadpoles are able to

survive in highly variable pool conditions (Fouilloux et al., 2021) and exhibit distinct behavioral

responses to different environmental stressors from an early stage (Surber-Cunningham et al.,

2024). Their developmental period is relatively slow, lasting a reported two months in the wild

(Rojas, 2014; Rojas & Pašukonis, 2019) and three to four months the lab (personal observation).

Our study aimed to determine if HPA axis functionality shifts during the ontogeny of *D.*

tinctorius tadpoles, potentially facilitating developmental transitions and supporting key

physiological and behavioral adaptations. Given the shared and distinct life history features

between *D. tinctorius* tadpoles and more temperate species, we had two alternative predictions.

On the one hand, given the constraints of metamorphosis and environmental challenges shared

with other tadpoles (predation, desiccation, etc.), *D. tinctorius* tadpoles' stress response could

have a developmental trajectory similar to other species, with a peak in HPA axis activity close

to metamorphosis. Alternatively, we predicted *D. tinctorius* might have an early-developed HPA

axis to facilitate the behavioral dexterity involved in coordinated transport, the ontogenetic

transition from land to water, responses to environmental challenges immediately upon arrival in

pools, and an extended developmental period. To test these alternatives, we compared (1) the

excretion rate and whole-body concentration of corticosterone and (2) the corticosterone

response to ACTH across developmental stages. We additionally confirmed corticosterone as the

dominant glucocorticoid and validated waterborne hormone sampling methods previously used

in adult *D. tinctorius* (Rodríguez López, 2015; Westrick et al., 2023) in tadpoles.

Methods

Animal husbandry

D. tinctorius tadpoles used in this study were hatched in the Fischer Lab frog colony at the

University of Illinois Urbana-Champaign (UIUC). Once parents deposited a tadpole into a pool,

we moved the tadpole to a transparent plastic cup (6cm diameter, 7.5cm height) with mesh

bottom within a larger plexiglass aquarium. We separated tadpoles into three aquaria with

tadpole densities of 12 – 20 individuals per aquarium in approximately 20 liters of reverse

osmosis (RO) water. In each aquarium, tadpoles shared water through the mesh bottoms of each

plastic cup but were unable to physically interact. We kept the water temperature constant at 22.8

$\pm1°C$ and kept the room under a 12:12-hour light/dark cycle. Tadpoles were also given 15

minutes of UV light per day to support development (Pahkala et al., 2003). We fed tadpoles a

diet of shrimp flakes and rabbit food pellets three times per week. We performed daily health

checks, weekly partial water changes, and weekly water quality checks (pH, salinity,

temperature). All animal care and experimental procedures were approved by the Animal Care

and Use Committee of the University of Illinois at Champaign-Urbana (IACUC Protocol number

20147).

Staging & morphological measurements

We measured tadpoles' total body mass, body length, tail length, head width, and Gosner stage.

To measure wet mass, we tared a portable scale (Weightman model B200) with a small cup of

water, and a tadpole was gently placed into the cup via a dry spoon. We recorded mass in grams

to the nearest 0.01 g. To measure total body length, tail length, and head width, we placed

tadpoles on a millimeter grid, and straightened body position gently with a fine paint brush.

Because tadpole developmental speed is highly plastic (Pfennig et al., 1991), it is standard

practice to use developmental stage rather than age. We categorized tadpole development via

Gosner stage, a numeric system based on key morphological characteristics (Gosner, 1960). We

examined tadpoles under a stereo microscope (Leica S9E) for accurate Gosner staging. We

separated tadpoles into three developmental categories: early stage (no external limb

development, Gosner stages 25-29, n=10), middle stage (subtle external leg development,

Gosner stages 30-34, n=10), and late stage (prominent external leg development, Gosner stages

35-40, n=10). Tadpoles were out of the water for no longer than five minutes during

morphometric data collection, and no tadpoles were harmed in the process.

ACTH hormone challenge

To test for differences in the physiological stress response, we manipulated the HPA axis via an

ACTH challenge. ACTH challenges are used across vertebrate taxa to stimulate the HPA axis

and assess stress responsivity (Hammond et al., 2019; McClelland & Woodley, 2021; Murray et

al., 2020). ACTH injections took place one or two days after morphological measurements. We

utilized a within-individual repeated measures design in which tadpoles were injected with both

an intraperitoneal injection of 0.5 ug ACTH per gram body weight (Sigma-Aldrich®, product no.

A7075) (Forsburg et al., 2019) diluted in 0.9% saline and a control injection of 0.9% saline (with

volume matched to the ACTH injection), with six days between injections. We used a 50μL

Hamilton syringe (Gastight® #1705) with single use 30G hypodermic needle (Excel Int, model

30Gx1) for injections. During injection, we placed tadpoles on a clean wet sponge and held them

in place with a wet strip of paper towel.

Waterborne hormone collection

We used waterborne hormone collection as a minimally invasive method for repeated hormone

sampling in small-bodied amphibians. To collect each hormone sample, we soaked the tadpole in

50mL RO water in a ten-centimeter diameter cylindrical glass container for 1 hour. To measure

pre-injection glucocorticoids, we collected a sample immediately prior to injection. To measure

how glucocorticoids respond to an ACTH and control injections over time, we collected water

samples at four time points post-injection: one hour, two hours, four hours, and twenty-four

hours. Pre-injection hormone collection started between 9am and10am, with injections one hour

after (between 10am and 11am). All individuals completed an ACTH trial and a saline trial one

week apart; thus, we collected ten waterborne hormone samples per individual (five per trial).

The order of the injections (ACTH/control) was counter-balanced to control for order effects.

After collection, we filtered water samples through filter paper (Whatman 1001-125) to remove

any debris and stored the filtered samples in 50mL conical tubes at -20°C until steroid hormone

extraction (see below). We collected a total of n=295 waterborne hormone samples (n = 30

tadpoles). One individual died between week one and two, but death believed unrelated to

experiment and data points still included.

Whole body hormone extraction

We euthanized tadpoles immediately following the 24-hour hormone collection using an

overdose of sodium bicarbonate buffered tricaine methane sulfonate (MS-222), a commonly-

used anesthetic for fish and amphibians (Ramlochansingh et al., 2014; Topic Popovic et al.,

2012). We rinsed bodies thoroughly with RO water, and stored bodies individually in

microcentrifuge tubes at -80°C for two weeks. Prior to whole-body hormone extraction, we

weighed bodies to the nearest 0.001 g using a microbalance (Mettler Toledo ML203T/00).

Whole bodies were homogenized in liquid nitrogen using a mortar and pestle. We transferred

homogenized tissue to a pre-weighed 15 mL centrifuge tube and recorded the powder-only mass.

Following weighing, we added 5 mL of methanol and vortexed for about 30 seconds. Samples

were stored overnight at -20 °C. The next day we centrifuged tubes with methanol and tissue (4

°C, 2000 rpm) for fifteen minutes (Eppendorf centrifuge 5810R). We carefully pipetted the

supernatant (~ 5 mL) into a 50 mL conical tube and added 45 mL of RO water. Samples were

then processed through the solid phase extraction process via the vacuum manifold procedure

(see below). We collected N=29 body samples for hormonal analysis (early=10, middle=9,

late=10).

Steroid hormone extraction

We extracted hormones from water samples and methanol-extracted whole-body samples using

solid-phase extraction via C18 cartridges (Waters Corporation) on a vacuum manifold. We

primed the cartridges with 6 mL of 100% methanol, rinsed with 6 mL RO water, and then passed

the 50mL sample through. We used 4 mL of 100% methanol to elute the collected total steroid

hormones into 13x100 mm borosilicate glass vials. We dried samples on a 37°C heating block

under a stream of nitrogen gas. After completely dry, we covered tubes with tight fitting caps and

stored at -20°C until quantification.

Enzyme linked immunosorbent assay

Methods for quantification of waterborne hormone samples using enzyme immunoassay

(EIA) have been utilized and validated in various amphibians (Baugh et al., 2018; Forsburg et

al., 2019; Gabor et al., 2016). These methods have been successfully applied in *D. tinctorius* by

our research group (Westrick et al., 2023) and others (Rodríguez López, 2015). We previously

confirmed parallelism between standards provided by the manufacturer and water samples from

D. tinctorius for corticosterone and cortisol (Westrick et al., 2023).

Commented [SW1]: If you say enzyme-linked immunosorbent assay, I would use ELISA. If you just say enzyme immunoassay, you can use EIA. Since you changed everything to EIA below, I changed the words up here. (same thing tho)

To quantify the concentration of cortisol and corticosterone in each sample, we used

DetectX® kits from Arbor Assays™. We reconstituted samples by adding assay buffer (catalog

no. X065) to the dried waterborne (500 μL) and whole-body samples (750 μL) in borosilicate

glass vials. We ran each sample in duplicate on both a corticosterone and a cortisol EIA.

Individuals that completed all sample collections had a total of ten waterborne samples and one

whole-body sample. Because we were interested in within-individual differences, we set up

plates so that all the samples taken from an individual were run on the assay to reduce potential

variation caused by plate-to-plate variation. We followed manufacturer guidelines for running all

EIA plates. We prepared the standard curve and samples using the high sensitivity 100 μL assay

format, such that the lowest point of the curve measured 19.531 pg/mL. The lowest point of the

cortisol curve measured 50 pg/mL. We measured the optical density at 450 nm for all assays.

The manufacturer reported cross reactivity of the cortisol EIA with corticosterone as 1.2%, and

cross reactivity of the corticosterone EIA with cortisol as 0.38%.

To generate a standard curve for each plate, we used MyAssay software

(www.MyAssay.com) to fit a four-parameter logistic curve (4PLC) regression. We used this

software to calculate the concentration of glucocorticoids (pg/mL of assay buffer) for each

sample based on the standard curve and averaged across duplicates. To calculate the excretion

rate for water samples, we multiplied the concentration by the volume of assay buffer used to

reconstitute the sample (0.5 mL) to get mass (pg) of the hormone in the sample. We then divided

by the collection time (one hour) to get the release rate (pg of hormone per hour of collection).

For whole-body samples concentrations, we corrected for the milliliters of assay buffer used

(0.75 mL) and the mass of powder collected after grinding (pg of hormone per gram of dry body

mass).

We removed samples with a coefficient of variation (CV) greater than 20% (n = 17 out of

328 total corticosterone measurements, n=233 out of 328 total cortisol measurements). Most

cortisol samples fell at the low end of the plate measurement range, and we believe high CVs for

cortisol arose primarily from measurement noise at the extremes of the detectable range. The

remaining average intra-assay CV across all samples was 6.85% for corticosterone and 10.01%

for cortisol.

To test the extraction efficiency of our hormone analysis, we collected additional samples

(N=12) by soaking tadpoles in RO water for one hour. We divided each sample equally to

measure the concentration of a control and a spike. For the spiked samples, we added either a

cortisol standard (N=6) or a corticosterone standard (N=6). These samples were then processed

through the extraction, drying, and plating methods described previously. To determine the

"observed" concentration of the spiked standard, we subtracted the control sample's value from

the corresponding spiked sample's value. The recovery rate was calculated by dividing the

observed concentration by the expected concentration (i.e., the amount added to the spike) and

multiplying by 100. Our results showed average recovery rates of 108% (range: 96.7%–127%)

for cortisol and 87.5% (range: 78.5%–102%) for corticosterone.

Statistical analysis

All statistical analyses were performed with R (v. 2022.07.1 + 554 R Core Development Team,

2022) in Rstudio (v. 2022.12.0 Rstudio Team, 2022). We log transformed corticosterone and

cortisol release rates for all tests to achieve normality. For all analyses described below, we ran

linear mixed models using the lmer() function in 'lme4' package (Bates et al., 2015), and

calculated test stats and p-values using the package 'lmertest'(Kuznetsova et al., 2017). To

analyze if there were significant inter-plate differences, we ran separate models for

corticosterone and cortisol with the glucocorticoid concentration as the dependent variable, plate

number as a fixed effect and individual as a random effect. The variation between EIA plates

was non-significant for corticosterone ($F_{1,30.9}=0.367$, p=0.716) and cortisol ($F_{1,17.08}=0.017$,

p=0.898).

To analyze which glucocorticoid (cortisol/corticosterone) was released at a higher rate,

we ran linear mixed effects models with the pre-injection glucocorticoid release rate (pg/hr) as

the dependent variable, and glucocorticoid type (cortisol/corticosterone), developmental stage

(early/middle/late), and the interaction between glucocorticoid type and developmental stage as

fixed effects, and individual as a random effect. We ran linear models using the whole-body

hormone concentrations to understand the abundance within the body, with concentration of

glucocorticoids (pg/g) as the dependent variable, and glucocorticoid type, treatment

(ACTH/saline control), developmental stage, and all interactions as fixed effects.

To assess pre-injection differences in corticosterone across developmental stages, we ran

linear mixed effects models with the pre-injection glucocorticoid release rate (pg/hr) as the

dependent variable, and developmental stage (early/middle/late) as a fixed effect, and individual

as a random effect. Tadpoles underwent two trials (one ACTH and one saline injection) one

week apart, thus we originally included week in this model, but removed it when nonsignificant

($F_{1,\,26.1}=0.218$, p=0.645). We ran a linear model using the whole-body hormone concentrations to

understand the abundance within the body, with concentration of glucocorticoids (pg/g) as the

dependent variable, and developmental stage as a fixed effect. We did not include body-size as a

covariate in our models as it was confounded with developmental stage (later stage tadpoles are

larger) and created problems with collinearity in the models.

We analyzed if whole-body corticosterone concentration predicted waterborne

corticosterone release rate via linear model, with week two 24-hour waterborne release rate (the

sample time point immediately preceding the body collection) as the dependent variable and the

whole-body corticosterone concentration, developmental stage (early/middle/late), and their

interaction as fixed effects. We did not include treatment in this model, because there were no

differences between treatments at this time point.

To analyze if corticosterone increased in response to ACTH and if this response changed

across developmental stages, we ran a linear mixed effects models with corticosterone

concentration as the dependent variable. We included all water-borne sampling time points and

fit models with treatment, timepoint, developmental stage, and all interactions between as fixed

effects and individual ID as a random effect. We included sampling week as a fixed effect in

original models but removed it to prevent over-fitting because it was not statistically significant

($F_{1,\ 238.26}$=0.074, p=0.786). We ran a post-hoc analyses using the package 'emmeans' (Lenth et

al., 2022) with pairwise tests to compare across timepoints, to compare the timepoints within

each treatment group (timepoint x treatment), to compare the treatment groups within timepoint

(timepoint x treatment), and the developmental groups between treatments (treatment x

development).

Results

Corticosterone release rate was significantly higher than cortisol release rate pre-injection

($F_{1,59.201}$=189.475, p<0.0001, Figure 1A) and in whole-bodies ($F_{1,34}$=46.546, p<0.0001, Figure

1B) in all developmental stages. Given the approximately seven-fold greater abundance of

corticosterone and the high number of cortisol samples dropped due to low concentration (see

Methods), we report differences across development and in response to ACTH challenge for

corticosterone only.

Corticosterone release rate did not differ across developmental stages in pre-injection

waterborne samples ($F_{2,47.04}$=0.361 p=0.699) (Figure 2A) or whole-body samples ($F_{2,25}$=1.043,

p=0.370) (Figure 2B). Whole-body corticosterone predicted waterborne corticosterone

($F_{1,18}$=21.95, p<0.005), and this relationship did not differ across developmental stages

($F_{2,18}$=0.506, p=0.611) (Figure 2C).

When we compared ACTH and saline treatment groups, we found an overall higher

corticosterone excretion rate in the ACTH treatment group (treatment: $F_{1,230.6}$=4.45, p<0.05)

(Figure 3A). Sampling time point also had a significant effect on corticosterone excretion rate

(time: $F_{4,224.94}$ = 24.85, p <0.0001), though the pattern of change was the same in both treatment

groups (treatment*time: $F_{4,224.84}$ = 1.43, p=0.2243) (Table 1). Comparing between treatment

groups, corticosterone release rate was higher in ACTH treated than saline treated tadpoles at

one hour post injection (t_{226} = 2.138, p<0.05) and was marginally higher at two hours post

injection ($t_{1,227}$ = 1.932, p=0.055) (Table 2).

We found no effect of developmental stage on treatment response

(treatment*time*development: $F_{8,224.84}$=0.759, p=0.639) (Figure 3B, 3C, 3D).

Discussion

A major way vertebrates coordinate responses to environmental stressors is through

activation of the hypothalamic-pituitary adrenal (HPA) axis (Carr, 2011; Dantzer, 2023, Denver,

2009a). However, what constitutes an appropriate stress response can change across

development as morphological and energetic changes shift the potential tradeoffs imposed by

excessive or insufficient HPA axis activity. This shifting landscape may in turn shape evolved

differences in the timing of the HPA axis development based on a species' life history. In this

study we investigated HPA axis development in dyeing poison frog (*Dendrobates tinctorius*)

tadpoles, a species with a unique life history that includes transport to water after hatching on

land, as well as aggressive and cannibalistic behavior. We specifically tested the alternatives that

D. tinctorius would (1) exhibit a developmental pattern similar to most tadpoles, with increased

GC abundance just before metamorphosis (Glennemeier & Denver, 2002; Jaudet & Hatey, 1984;

Krug et al., 1983), or (2) have an early-developed HPA axis given the importance of

glucocorticoids in ontogenetic transitions (Wada, 2008) and for rapid phenotypic responses to

short-term environmental stressors (Dantzer, 2023; Hau et al., 2016).

We found no differences across tadpole developmental stages in corticosterone excretion

rate in waterborne samples, whole-body concentration of corticosterone, nor ACTH response,

supporting the prediction that *D. tinctorius* tadpoles have an early developed and consistently

active HPA axis. This result contrasts with findings in other species, where late-stage tadpoles

have corticosterone levels two to three times higher than early or middle-stage tadpoles

(Glennemeier & Denver, 2002; Jaudet & Hatey, 1984; Krug et al., 1983). Notably, other species

that have been studied tend to be explosive breeders that produce 10s to 1000s of aquatic eggs

and provide no parental care. We suggest these key life history differences could contribute to

our findings here.

Our study also adds to the mounting evidence that there is species variation in HPA axis

reactivity across development in tadpoles. In American bullfrogs (*R. catesbeiana*), late-stage

tadpoles mount a response greater than early stage tadpoles (Krug et al., 1983), while Northern

leopard frogs (*Rana pipiens*) and African clawed frogs (*Xenopus laevis*) have a larger ACTH

response as early-stage tadpoles than late-stage (Glennemeier, 2002). In *D. tinctoirus*, we found

equal reactiveness to ACTH across development. Below we discuss multiple, non-mutually

exclusive reasons why an early-developed and consistent HPA axis could be important in this

species.

We posit that the early development of the HPA axis in *D. tinctorius* tadpoles may

facilitate transportation and transition from terrestrial hatching sites to developmental pool, and

once there, enable plastic responses to highly variable environmental conditions and conflicts

with conspecifics. In *D. tinctorius*, tadpoles hatch terrestrially, then crawl onto the backs of their

parents and hold on as they are transported for minutes to hours to their new aquatic habitat. This

action takes collaboration between parents and offspring, as well as water balance while in the

open air and during transition to a fully aquatic environment. An early developed stress response

may facilitate these metabolically demanding behaviors (Magomedova & Cummins, 2016). We

did not compare glucocorticoid abundance or responsivity pre-transport in this study, as out non-

invasive waterborne sampling method would have inherently confounded measuring the effects

of the terrestrial to aquatic transition. However, this is an exciting avenue for future research as

this land-to-water transition is fundamentally different from the majority of frogs that lay aquatic

eggs and do not provide care.

After arriving at their pond, *D. tinctorius* face challenges similar to those of most tadpoles

(predation, competition, desiccation) as well as some additional challenges. The pools *D.*

tinctorius tadpoles inhabit vary considerable in water quality, predator and conspecific

abundance, and pool stability, especially compared to other neotropical species (Fouilloux et al.,

2021). These variables are known to cause plastic developmental shifts in other tadpoles

(McDiarmid & Altig, 1999), and such plasticity could be facilitated by an early developed HPA

axis in *D. tinctorius*. Additionally, *D. tinctorius* tadpoles are aggressive and cannibalistic

(Fouilloux et al., 2022; Fischer et al., 2020), which makes interactions with conspecifics a

potential danger or a potential meal. Indeed, we previously demonstrated that *D. tinctorius*

exhibit behavioral responses to injured conspecific cues that are distinct from responses to food

or predator cues (CITE LISA'S PAPER!). We suggest that corticosterone may be mediating

behavioral responses to these unique cues as glucocorticoids are known mediators of behavioral

plasticity generally (Dantzer, 2023), and are positively correlated with aggression (Summers &

Winberg, 2006) and antipredator behavior (Fraker et al., 2009) in other tadpole species. Our

findings lay the foundation for future studies to investigate the role of corticosterone as a

mediator of phenotypic plasticity and juvenile aggression.

In general, tadpole responses to ACTH challenge were as expected: treatment groups did not

differ pre-injection, corticosterone excretion was relatively higher following ACTH treatment,

and corticosterone levels recovered to pre-injections concentrations 24 hours post-injection.

While we expected this general response, we further asked whether the shape or magnitude of

the response differed across development. We found no significant differences in ACTH

response across developmental stages. This observation has two important implications. First, it

demonstrates that *D. tinctorius* tadpoles can mount a physiological stress response even early in

development, confirming our prediction that this species has an early developed HPA axis.

Second, these results indicate that developmental effects of corticosterone do not interfere with

stress responsivity in this species. Taken together, these observations demonstrate that *D.*

tinctorius can respond rapidly and consistently to stressors throughout their aquatic

developmental period.

Although the response to ACTH was as we expected, one unanticipated result in our study

was that corticosterone excretion pre-injection did not differ significantly from excretion one-

hour post-injection in either the ACTH or saline group. This suggests both groups exhibited

some handling stress associated with waterborne hormone collection (moving from their home

cup to the hormone collection cup), as has been observed in other species (Wong et al., 2008).

Though waterborne hormone sampling requires some handling (Wong et al., 2008) and is

inherently an integrative measure across time, we expect any effects of sampling procedure to be

consistent across experimental groups. Similarly, any stress resulting from injection is shared in

ACTH and control treatments. Importantly, we nonetheless found that corticosterone was

elevated in the ACTH group as compared to saline, indicating that the HPA axis was activated

by our pharmacological manipulation above responses to handling and/or injection. Indeed,

when treated with ACTH, individuals had elevated corticosterone and corticosterone decreased

more slowly than when treated with saline. At present, we cannot distinguish whether the lower

corticosterone at 2-hours and 4-hours post injection results from short-term habituation to sample

collection or negative feedback from the HPA-axis, and either alternative suggests that all

tadpoles experienced at least a slight activation in the sampling and injection process. We note

that we have no indication for habituation at longer timescales, as corticosterone levels at 24

hours post injection and in the second week of sampling were similar to those pre-injection and

in the first week. While these observations underscore the importance of considering habituation

procedures when conducting experiments involving glucocorticoid manipulation in tadpoles and

other organisms, they do not negate the results and interpretations discussed here.

We found that whole-body corticosterone concentration predicted the release rate of

waterborne corticosterone, and that this relationship did not change across development. From a

technical perspective, our findings validate the use of waterborne hormone measurements in

tadpoles of *D. tinctorius*. Waterborne hormone sampling is minimally invasive, allowing for

repeated sampling in small-bodied individuals and non-destructive sampling of wild populations,

thereby providing many potential avenues for future study. While correlations between body and

water samples are expected, confirmation of this pattern is also biologically interesting, as

previous studies have documented changes in body/water correlations across tadpole

development (McClelland & Woodley, 2021). McClelland & Woodley (2021) report a lack of

correlation between waterborne and plasma hormone concentrations in early-stage tadpoles and

those undergoing metamorphic climax, suggesting excretion rate can be plastic across

development, at least in some species. Data from additional species are needed to understand

whether and how these patterns may be related to developmental changes in HPA-axis activity.

Finally, our study confirmed that corticosterone is the dominant and most abundant

glucocorticoid in *D. tinctorius*. Across tadpole development, corticosterone was approximately

seven times more abundant than cortisol. Corticosterone is generally considered the primary

glucocorticoid in amphibians (Denver, 2009a; Glennemeier & Denver, 2002; Hossie et al., 2010;

Jaudet & Hatey, 1984), however, there are exceptions. Indeed, previous studies in adult *D.*

tinctorius demonstrate that, although corticosterone is the ACTH responsive glucocorticoid,

adults release equal amounts of corticosterone and cortisol (Westrick et al. 2023), and increased

cortisol in parents is associated with tadpole transport (Fischer & O'Connell, 2020). Taken

together, these previous observations and our data here suggest that the production and relative

abundance of corticosterone and cortisol may shift post-metamorphosis or as adults become

sexually mature. Further exploration of these patterns and their potential functional consequences

is an area of ongoing research.

Conclusions

We found no significant differences in glucocorticoid excretion rates, whole body

concentrations, or physiological responses to ACTH injection across tadpole developmental

stages in *D. tinctorius*. Taken together, these results suggest that the HPA axis is functional early

in poison frog tadpole ontogeny. Our findings provide a foundation for exploring how *D.*

tinctorius tadpoles physiologically respond to a habitat transitions and environmental stressors,

as well as prepare for and undergo metamorphosis. Additionally, this pattern contrasts with other

tadpole species, providing inroads for exciting comparative work to further build our

understanding of how life-history shapes adaptive differences in the developmental timing of

stress responses.

Acknowledgements

We would like to thank Drs. Britt Carlson and Nathan Schroeder from the Phenotypic Plasticity

Research Experience for Community College Students (PRECS) program for adding Lauren

Mobo to the team. We also thank Drs. Chris Holmes and Ken Paige, program directors of the

Access and Achievement Program for adding Lucas Jimenez to the project. We appreciate the

members of the Fischer Lab for feedback on previous versions of the manuscript and help with

tadpole husbandry. Our work would not have been possible without the animal care support of

the Division of Animal Resources (DAR) at UIUC.

Declarations of interest

None.

Author contributions

Lisa L. Surber-Cunningham: Conceptualization, Methodology, Formal Analysis,

Investigation, Data Curation, Writing – Original Draft, Writing – Review & Editing,

Visualization, Supervision, Project Administration. **Lucas S. Jimenez**: Investigation. **Lauren**

W. Mobo: Investigation. **Sarah E. Westrick**: Conceptualization, Methodology, Formal

Analysis, Writing – Review & Editing. **Eva K. Fischer**: Conceptualization, Methodology,

Formal Analysis, Funding Acquisition, Resources, Supervision, Writing – Review & Editing.

Funding

Financial support was provided by the Edward Banks Award for the Study of Behavior from the

UIUC EEB Department (to L.L.S.), the Francis M. and Harlie M. Clark Research Fellowship

from UIUC School of integrative biology (to L.L.S.), Access and Achievement Program Summer

Research Fellowship through the College of Liberal Arts & Sciences and the UIUC School of

Integrative Biology (to L.S.J.), NSF REU through the Phenotypic Plasticity Research Experience

for Community College Students through the Institute for Genomic Biology at UIUC and

Parkland College (REU 1950819/1950786 to L.W.M), NSF PRFB funding (#2010714 to

S.E.W.), NIH R35 (#R35GM147207 to E.K.F.), Research Board funding from UIUC (RB21025

to E.K.F), and UIUC Laboratory Start-up funds (to E.K.F).

References

Bates D, Mächler M, Bolker B, Walker S (2015). "Fitting Linear Mixed-Effects Models Using

lme4." *Journal of Statistical Software*, 67(1), 1–48. doi:10.18637/jss.v067.i01.

Baugh, A. T., Bastien, B., Still, M. B., & Stowell, N. (2018). Validation of water-borne steroid

hormones in a tropical frog (Physalaemus pustulosus). *General and Comparative*

Endocrinology, *261*, 67–80. https://doi.org/10.1016/j.ygcen.2018.01.025

Bókony, V., Ujhegyi, N., Hamow, K. Á., Bosch, J., Thumsová, B., Vörös, J., Aspbury, A. S., &

Gabor, C. R. (2021). Stressed tadpoles mount more efficient glucocorticoid negative

feedback in anthropogenic habitats due to phenotypic plasticity. *Science of The Total*

Environment, *753*, 141896. https://doi.org/10.1016/j.scitotenv.2020.141896

Carr, J. A. (2011). Stress and Reproduction in Amphibians. In *Hormones and Reproduction of*

Vertebrates (pp. 99–116). Elsevier. https://doi.org/10.1016/B978-0-12-374931-4.10006-9

Charmandari, E., Tsigos, C., & Chrousos, G. (2005). Endocrinology of the stress reponse.

Annual Review of Physiology, *67*(1), 259–284.

https://doi.org/10.1146/annurev.physiol.67.040403.120816

Chloe A Fouilloux, Lutz Fromhage, Janne K Valkonen, & Bibiana Rojas. (2022). Size-

dependent aggression towards kin in a cannibalistic species. *Behavioral Ecology*.

https://doi.org/10.1093/beheco/arac020

Dantzer, B. (2023). Frank Beach Award Winner: The centrality of the hypothalamic-pituitary-

adrenal axis in dealing with environmental change across temporal scales. *Hormones and*

Behavior, 105311. https://doi.org/10.1016/j.yhbeh.2023.105311

Davis, G. S., & Siopes, T. D. (1985). Adrenal Cortical Response of Tom Poults. *Poultry Science*,

64(11), 2189–2194. https://doi.org/10.3382/ps.0642189

Denver, R. J. (1998). *Hormonal Correlates of Environmentally Induced Metamorphosis in the*

Western Spadefoot Toad, Scaphiopus hammondii.

Denver, R. J. (2009). Endocrinology of complex life cycles: Amphibians. *Hormones, Brain and*

Behavior Online, *2*, 707–745. https://doi.org/10.1016/B978-008088783-8.00021-8

Fischer, E. K., Alvarez, H., Lagerstrom, K. M., McKinney, J. E., Petrillo, R., Ellis, G., &

O'Connell, L. A. (2020). Neural correlates of winning and losing fights in poison frog

tadpoles. *Physiology and Behavior*, *223*. https://doi.org/10.1016/j.physbeh.2020.112973

Fischer, E. K., & O'Connell, L. A. (2020). Hormonal and neural correlates of care in active

versus observing poison frog parents. *Hormones and Behavior*, *120*(January), 104696.

https://doi.org/10.1016/j.yhbeh.2020.104696

Forsburg, Z. R., Goff, C. B., Perkins, H. R., Robicheaux, J. A., Almond, G. F., & Gabor, C. R.

(2019). Validation of water-borne cortisol and corticosterone in tadpoles: Recovery rate

from an acute stressor, repeatability, and evaluating rearing methods. *General and*

Comparative Endocrinology, *281*, 145–152. https://doi.org/10.1016/j.ygcen.2019.06.007

Fouilloux, C. A., Serrano Rojas, S. J., Carvajal-Castro, J. D., Valkonen, J. K., Gaucher, P.,

Fischer, M., Pašukonis, A., & Rojas, B. (2021). Pool choice in a vertical landscape:

Tadpole-rearing site flexibility in phytotelm-breeding frogs. *Ecology and Evolution*,

ece3.7741. https://doi.org/10.1002/ece3.7741

Fraker, M. E., Hu, F., Cuddapah, V., McCollum, S. A., Relyea, R. A., Hempel, J., & Denver, R.

J. (2009). Characterization of an alarm pheromone secreted by amphibian tadpoles that

induces behavioral inhibition and suppression of the neuroendocrine stress axis.

Hormones and Behavior, *55*(4), 520–529. https://doi.org/10.1016/j.yhbeh.2009.01.007

Francesco Ficetola, G., & De Bernardi, F. (2006). Trade-off between larval development rate and

Post-metamorphic Traits in the Frog Rana latastei. *Evolutionary Ecology*, *20*(2), 143–

158. https://doi.org/10.1007/s10682-005-5508-6

Gabor, C. R., Zabierek, K. C., Kim, D. S., Da Barbiano, L. A., Mondelli, M. J., Bendik, N. F., &

Davis, D. R. (2016). A Non-Invasive Water-Borne Assay of Stress Hormones in Aquatic

Salamanders. *Copeia*, *104*(1), 172–181. https://doi.org/10.1643/OT-14-207

Glennemeier, K. A. (2002). Developmental Changes in Interrenal Responsiveness in Anuran

Amphibians. *Integrative and Comparative Biology*, *42*(3), 565–573.

https://doi.org/10.1093/icb/42.3.565

Glennemeier, K. A., & Denver, R. J. (2002). Small changes in whole-body corticosterone

content affect larval Rana pipiens fitness components. *General and Comparative*

Endocrinology, *127*(1), 16–25. https://doi.org/10.1016/S0016-6480(02)00015-1

Gosner, K. L. (1960). A Simplified Table for Staging Anuran Embryos and Larvae with Notes

on Identification. *Herpetologica*, *16*(3), 183–190.

Hammond, T. T., Vo, M., Burton, C. T., Surber, L. L., Lacey, E. A., & Smith, J. E. (2019).

Physiological and behavioral responses to anthropogenic stressors in a human-tolerant

mammal. *Journal of Mammalogy*, *100*(6), 1928–1940.

https://doi.org/10.1093/jmammal/gyz134

Hau, M., Casagrande, S., Ouyang, J. Q., & Baugh, A. T. (2016). Glucocorticoid-Mediated

Phenotypes in Vertebrates. In *Advances in the Study of Behavior* (Vol. 48, pp. 41–115).

Elsevier. https://doi.org/10.1016/bs.asb.2016.01.002

Hossie, T. J., Ferland-Raymond, B., Burness, G., & Murray, D. L. (2010). Morphological and

behavioural responses of frog tadpoles to perceived predation risk: A possible role for

corticosterone mediation? *Ecoscience*, *17*(1), 100–108. https://doi.org/10.2980/17-1-3312

Jaudet, G. J., & Hatey, J. L. (1984). Variations in aldosterone and corticosterone plasma levels

during metamorphosis in Xenopus laevis tadpoles. *General and Comparative*

Endocrinology, *56*(1), 59–65. https://doi.org/10.1016/0016-6480(84)90061-3

Kitaysky, A. S., Kitaiskaia, E. V., Piatt, J. F., & Wingfield, J. C. (2003). Benefits and costs of

increased levels of corticosterone in seabird chicks. *Hormones and Behavior*, *43*(1), 140–

149. https://doi.org/10.1016/S0018-506X(02)00030-2

Krug, E. C., Honn, V., Battista, J., & Nicoll, C. S. (1983). Corticosteroids in Serum of Rana

catesbeiana during Development and Metamorphosis. *General and Comparative*

Endocrinology, *52*(2), 232–241.

Kulkarni, P. S., & Gramapurohit, N. P. (2017). Effect of corticosterone on larval growth,

antipredator behaviour and metamorphosis of Hylarana indica. *General and Comparative*

Endocrinology, *251*, 21–29. https://doi.org/10.1016/j.ygcen.2016.09.001

Kuznetsova, A., Brockhoff, P. B., & Christensen, R. H. B. (2017). **lmerTest** Package: Tests in

Linear Mixed Effects Models. *Journal of Statistical Software*, *82*(13).

https://doi.org/10.18637/jss.v082.i13

Ledón-Rettig, C. C., Lo, K. M., & Lagon, S. R. (2023). Baseline corticosterone levels in

spadefoot toads reflect alternate larval diets one year later. *General and Comparative*

Endocrinology, *339*, 114291. https://doi.org/10.1016/j.ygcen.2023.114291

Ledón-Rettig, C., & Lagon, S. (2021). A novel larval diet interacts with nutritional stress to

modify juvenile behaviors and glucocorticoid responses. *Evology and Evolution*, *11*(16),

10880–10897.

Magomedova, L., & Cummins, C. L. (2016). *Glucocorticoids and Metabolic Control* (S. Herzig,

Ed.; Vol. 233). Springer International Publishing. https://doi.org/10.1007/978-3-319-

29806-1

McClelland, S. J., & Woodley, S. K. (2021). Water-borne corticosterone assay is a valid method

in some but not all life-history stages in Northern Leopard Frogs. *General and*

Comparative Endocrinology, *312*, 113858. https://doi.org/10.1016/j.ygcen.2021.113858

McDiarmid, R. W., & Altig, R. (Eds.). (1999). *Tadpoles: The biology of anuran larvae.*

University of Chicago Press.

Murray, M. J., Young, M. A., & Santymire, R. M. (2020). Use of the ACTH challenge test to

identify the predominant glucocorticoid in the southern sea otter (Enhydra lutris nereis).

Conservation Physiology, *8*(1), coz116. https://doi.org/10.1093/conphys/coz116

Pahkala, M., Laurila, A., & Merilä, J. (2003). Effects of ultraviolet-B radiation on behaviour and

growth of three species of amphibian larvae. *Chemosphere*, *51*(3), 197–204.

https://doi.org/10.1016/S0045-6535(02)00813-5

Pfennig, D. W., Mabry, A., & Orange, D. (1991). Environmental Causes of Correlations between

Age and Size at Metamorphosis in Scaphiopus Multiplicatus. *Ecology*, *72*(6), 2240–2248.

https://doi.org/10.2307/1941574

Ramlochansingh, C., Branoner, F., Chagnaud, B. P., & Straka, H. (2014). Efficacy of Tricaine

Methanesulfonate (MS-222) as an Anesthetic Agent for Blocking Sensory-Motor

Responses in Xenopus laevis Tadpoles. *PLoS ONE*, *9*(7), e101606.

https://doi.org/10.1371/journal.pone.0101606

Rodríguez López, C. (2015). Does testosterone correlate with the aposematic syndrome in

Amazonian poison. *Universidad de Los Andes*.

Rojas, B. (2014). Strange parental decisions: Fathers of the dyeing poison frog deposit their

tadpoles in pools occupied by large cannibals. *Behavioral Ecology and Sociobiology*,

68(4), 551–559. https://doi.org/10.1007/s00265-013-1670-y

Rojas, B., & Pašukonis, A. (2019). From habitat use to social behavior: Natural history of a

voiceless poison frog, Dendrobates tinctorius. *PeerJ*, *2019*(9).

https://doi.org/10.7717/peerj.7648

Sapolsky, R. M., Krey, L. C., & McEwen, B. S. (1986). *The Neuroendocrinology of Stress and*

Aging: The Glucocorticoid Cascade Hypothesis. *7*(3).

Sapolsky, R. M., Romero, L. M., & Munck, A. U. (2000). *How Do Glucocorticoids Influence*

Stress Responses? Integrating Permissive, Suppressive, Stimulatory, and Preparative

Actions. *21*(1).

Schmidt, M., Oitzl, M. S., Levine, S., & De Kloet, E. R. (2002). The HPA system during the

postnatal development of CD1 mice and the effects of maternal deprivation.

Developmental Brain Research, *139*(1), 39–49. https://doi.org/10.1016/S0165-

3806(02)00519-9

Summers, C. H., & Winberg, S. (2006). Interactions between the neural regulation of stress and

aggression. *Journal of Experimental Biology*, *209*(23), 4581–4589.

https://doi.org/10.1242/jeb.02565

Surber-Cunningham, L. L., Oza, S. S., & Fischer, E. K. (2024). Chemical cues of conspecific

predation elicit distinct behavioural responses in cannibalistic poison frog tadpoles.

Animal Behaviour, *208*, 79–89. https://doi.org/10.1016/j.anbehav.2023.12.003

Topic Popovic, N., Strunjak-Perovic, I., Coz-Rakovac, R., Barisic, J., Jadan, M., Persin

Berakovic, A., & Sauerborn Klobucar, R. (2012). Tricaine methane-sulfonate (MS-222)

application in fish anaesthesia. *Journal of Applied Ichthyology*, *28*(4), 553–564.

Wada, H. (2008). Glucocorticoids: Mediators of vertebrate ontogenetic transitions. *General and*

Comparative Endocrinology, *156*(3), 441–453.

https://doi.org/10.1016/j.ygcen.2008.02.004

Westrick, S. E., Paitz, R. T., & Fischer, E. K. (2023). *Why not both? A case study measuring*

cortisol and corticosterone in poison frogs. https://doi.org/10.1101/2023.06.19.545597

Wong, S., Dykstra, M., Campbell, J., & Earley, R. (2008). Measuring water-borne cortisol in

convict cichlids (Amatitlania nigrofasciata): Is the procedure a stressor? *Behaviour*,

145(10), 1283–1305. https://doi.org/10.1163/156853908785765863

Tables

Table 1. Post-hoc results comparing corticosterone excretion rates pre and post injection samples, separated for ACTH and saline treatments.

Collection hours	ACTH		Saline	
	t	p	t	p
Pre - 1	t_{225}= -2.362	0.1297	t_{224}=-0.169	0.998
Pre - 2	t_{225}= 1.926	0.3065	**t_{225}= 3.990**	**<0.001**
Pre - 4	**t_{225}= 4.367**	**<0.0005**	**t_{224}= 5.848**	**<0.0001**
Pre - 24	t_{225}= 1.546	0.5337	t_{225} =1.033	0.8396
1 - 2	**t_{225}= 4.245**	**<0.0005**	**t_{225} = 4.157**	**<0.0005**
1 - 4	**t_{225}= 6.686**	**<0.0001**	**t_{224}= 6.016**	**<0.0001**
1 - 24	**t_{225}= 3.874**	**<0.005**	t_{225} =1.197	0.7530
2 - 4	t_{225}= 2.369	0.1276	t_{225} =1.788	0.3833
2 - 24	t_{225}= -0.394	0.9949	**t_{225} = -2.861**	**<0.05**
4 - 24	**t_{224}= -2.795**	**<0.05**	**t_{225} = -4.653**	**<0.0005**

Table 2. Post hoc comparisons of corticosterone excretion of ACTH and saline across the hormone time collections.

Collection hour	corticosterone	
	t value	p value
Pre-Injection	T_{226} = -0.050	0.9600
Post 1 hour	**$T_{1,226}$ = 2.138**	**0.0336**
Post 2 hour	**$T_{1,227}$ = 1.932**	**0.0546**
Post 4 hour	$T_{1,226}$ = 1.335	0.1832
Post 24 hour	$T_{1,226}$ = -0.534	0.5937

648 **Figures**

Figure 1. Tadpoles excrete more corticosterone (orange) than cortisol (blue) (A) pre-injection (B) whole-body. We collapsed developmental groups together because there were no differences between them. Boxplots show the median, the first and third quartiles, and whiskers as 1.5x the interquartile range. Black dots show individual data points. Red diamonds show group means.

Figure 2. Early (dark blue), middle (blue), and late-stage (light blue) tadpoles do not differ in corticosterone in (A) pre-injection excretion rate or (B) whole-body concentration. Boxplots show the median, the first and third quartiles, and whiskers as 1.5x the interquartile range. Black dots show individual data points. Red diamonds show group means. (C) Whole-body corticosterone predicted corticosterone excreted, with no significant differences in the water-body relationship between developmental groups.

Figure 3. When (A) averaged across all developmental groups, tadpoles injected with ACTH (red) excreted significantly more corticosterone one hour and marginally more corticosterone two hours post injection compared to the saline control (gray). There was no significant difference in ACTH response between (B) early, (C) middle, or (D) late-stage tadpoles. Data are shown as means +/- standard error.