Low-invasive, wide-field, and cellular resolution two-photon imaging
of neural population activity in brainstem and nucleus tractus solitarii

Authors:

Masakazu Agetsuma[1,2,3,*], Azumi Hatakeyama[3], Daisuke Yamada[3,*], Hiroshi Kuniishi[4],

Eri Takeuchi[5], Shinji Tsuji[1], Chihiro Ito[2], Tomoko Kobayashi[1], Atsushi Noritake[6],

Yoshitsugu Aoki[5], Hiroshi Yukawa[2], Akiyoshi Saitoh[3], Junichi Nabekura[1], Masayuki

Sekiguchi[3,5]

Affiliations:

[1]Division of Homeostatic Development, National Institute for Physiological Sciences, 38
Nishigohnaka Myodaiji-cho, Okazaki, Aichi, 444-8585, Japan

[2] Quantum Regenerative and Biomedical Engineering Team, Institute for Quantum Life
Science, National Institutes for Quantum Science and Technology (QST), Anagawa 4-9-
1, Chiba Inage-ku, Chiba 263-8555, Japan

[3] Laboratory of Pharmacology, Faculty of Pharmaceutical Sciences, Tokyo University of
Science, 2641 Yamazaki, Noda, Chiba 278-8510, Japan

[4]Division of Development of Mental Functions, Research Center for Child Mental
Development, University of Fukui, 23-3, Matsuokashimoaizuki, Eiheiji-cho, Yoshida-
gun, Fukui 910-1193, Japan

[5]Department of Molecular Therapy, National Institute of Neuroscience, National Center
of Neurology and Psychiatry, 4-1-1 Ogawahigashi, Kodaira, Tokyo 187-8502, Japan.

[6]Division of Behavioral Development, Department of System Neuroscience, National
Institute for Physiological Sciences, 38 Nishigohnaka Myodaiji-cho, Okazaki, Aichi,
444-8585, Japan

26
27 *Corresponding author. Email: age@nips.ac.jp, yamadada@rs.tus.ac.jp
28

Keywords:
Brain-viscera communication
Emotion
Two photon live brain imaging
Nucleus tractus solitarii
Vagus nerve stimulation
Brainstem
36

Summary: Brain-viscera communication plays a crucial role in regulating mental health,
with the vagus nerve being a key structure mediating this interaction. Clinically, artificial
vagus nerve stimulation (VNS) is used to treat various neuropsychiatric disorders,
highlighting the importance of vagal afferent fibers in regulating emotion. The nucleus
tractus solitarii (NTS) is a brainstem structure proposed to receive signals from vagal
afferents and relay them to brain networks for emotion regulation. However, due to the
anatomical complexity and difficulty in accessing the deep-brain NTS region in living
animals, the mechanisms remain unclear. Here, we developed a wide-field and deep-brain
two-photon imaging method using a double-prism based optical interface. This approach
enables the identification of cellular-resolution neural activities in the NTS while
preserving the cerebellum, which covers the NTS and is important for emotion regulation,
intact. We systematically evaluated how NTS neurons respond to VNS and a
gastrointestinal hormone, suggesting the usefulness of this method for investigating the
role of the vagus-NTS pathway *in vivo*.

Introduction
Optical techniques are powerful for measuring and artificially manipulating neuronal
activity to elucidate the function of neuronal circuits and dynamic information processing
in the brain. These techniques have been widely used in rodent brain research. However,
technical limitations remain in accessing deep brain regions. For example, the brainstem,
a key deep-brain region for brain-viscera communication, has been difficult to optically
investigate its physiological properties.

Regulating emotion through brain-viscera communication has received significant
attention, and the abnormal communication is linked to neuropsychiatric disorders[1-6].
Afferent vagus fibers originating in the nodose ganglion in the cervical cord play a crucial
role in brain-viscera communication, receiving information from various internal organs
and relaying it to the central nervous system through the brainstem area[5,6]. Artificial vagus
nerve stimulation (VNS), which is clinically applied to treat various neurological and
psychiatric disorders[7-9], mainly target the afferent vagus nerve. Initially reported in the
19th century, the FDA approved VNS for treating drug-resistant seizures in 1997 and
depression in 2005[10,11]. Currently, several neuropsychiatric and related disorders are
considered potentially treatable with VNS[12]. For safe and effective VNS, the parameter

settings must be carefully optimized[13,14]. Research using rodent models is crucial for this

optimization by elucidating the mechanisms underlying VNS and its therapeutic effects.

Previous studies in rodents demonstrated that VNS effectively suppresses seizures[15,16],

mitigates cerebral ischemia[17], and treats chronic pain[18]. Furthermore, VNS significantly

reduces depressive symptoms[19,20] and facilitates the extinction of associative fear

memory[21]. These findings suggest that the insights gained from animal models are critical

to advancing VNS-based clinical treatments and understanding the mechanisms of

emotion regulation through the vagus nerve.

Previous studies have indicated that the nucleus tractus solitarii (NTS) is likely a

critical gateway for regulating emotions through brain-viscera communication and for

VNS-based clinical treatments[6,8,22,23]. Afferent vagus fibers receive information from

various internal organs and are proposed to relay it to the NTS[6,23-26]. The NTS projects to

various brain regions involved in emotion regulation, such as the amygdala, lateral

hypothalamic area, bed nucleus of the stria terminalis, and periaqueductal gray[6,8], which

suggest the involvement of the vagus-NTS pathway in regulating emotion. VNS has also

been shown to modulate the activity of various brain regions[27-29], including the locus

coeruleus, whose activity is regulated by the NTS. Lesions of the locus coeruleus suppress

the anti-epileptic effects of VNS[15]. The actual role of the NTS and the detailed

mechanisms underlying the therapeutic effects of VNS, however, remain unclear. It is

also unclear how information from various organs relayed to the NTS for regulating

emotion is processed, and whether the information from various organs is segregated or

integrated within the NTS.

Addressing these questions has been challenging because the NTS is located deep in

the brain and is surrounded by regions critically involved in vital functions, making direct

observation difficult. Removing the cerebellum to expose the brainstem allows for

observations of a wide area of the brainstem including the NTS, which is useful to study

the role of the NTS in regulating the information from the digestive systems[30]. However,

the cerebellum is also an essential brain structure for emotional regulation[31-34] and is

modulated by the VNS-based treatment[28]. Therefore, the development of a novel

technique for *in vivo* recordings of brainstem neurons without removing the cerebellum

is in high demand. A technique to optically access the brainstem and NTS in the deep

brain area while keeping the cerebellum intact would expand the possibilities for neural

activity imaging and optogenetic neural activity manipulation, thereby profoundly advancing our understanding of the function of the NTS and its contribution to emotional regulation.

This aim motivated us to develop an *in vivo* two-photon imaging technique using a double-prism based optical interface (<u>D</u>ouble-<u>P</u>rism based brain<u>S</u>tem imaging under <u>C</u>erebellar <u>A</u>rchitecture and <u>N</u>eural circuits, "D-PSCAN"). The D-PSCAN method enables the observation of neural population activity with single-cell resolution in the brainstem and NTS situated deeply beneath the cerebellum while keeping the cerebellum and surrounding neural connections intact. We demonstrated the capability of this method to distinguish NTS activity from that of descending vagus neurons whose cell bodies lie ventrally to the NTS. The recorded neural activity data were used to evaluate the physiological characteristics of NTS neurons, i.e., their responsiveness to VNS. We also demonstrated the application of this imaging technique for detecting responsiveness to a gastrointestinal hormone cholecystokinin-8 (CCK), a more physiological stimulus than VNS. Our results suggest the applicability of this method for detailed understanding of the physiological mechanism of the NTS and future research on emotional regulation through the vagus-NTS pathway.

Results:

<u>Double microprism for optical access to the brainstem area deep below the cerebellum</u>

We developed a novel *in vivo* two-photon calcium imaging method for observing the deep brain area in mice that enables neural population activity recordings in the brainstem and NTS. Using a genetically encoded Ca^{2+} indicator, GCaMP, expressed by an adeno-associated virus (AAV), we aimed to detect neural activities in the NTS (Figure 1A–B) to observe the responsiveness of NTS neurons to various stimuli. Viruses encoding GCaMP were injected into the NTS at approximately postnatal day (P) 50–60 for *in vivo* imaging experiments, at least 1 month before the microprism implantation described below. Although primarily targeted at the NTS, viruses often spread to adjacent areas, beneficially inducing additional expression in surrounding regions and allowing us to visualize wide areas of the brainstem.

The brainstem is located deeply below (2.0–2.5 mm) the dorsal surface of the cerebellum. One potential problem in observing neural activities in the brainstem is that

even when the mouse head is fixed under the objective lens for two-photon imaging by
tightly attaching the head plate to the skull, the brainstem can still move somewhat
independently in response to the mouse's body movements[30]. Although standard skull
reference points (bregma and lambda) are typically referred to when targeting specific
brain areas, it is also crucial to visually capture the brain structure under observation in
each mouse. In a previous study, the cerebellum was removed to address these issues[30],
but the cerebellum is also involved in emotion control[31]; therefore, to accurately elucidate
the role of the vagus-NTS pathway in regulating emotion, it is essential to have a wide
field of view of the brainstem while keeping the cerebellum and neural connections intact.
To achieve this aim, we developed a technique to optically access the deep brainstem
area using a double prism (Figure 1C–E). We designed an implantable double microprism
assembly comprising a 2-mm right-angle glass microprism bonded to another 2-mm
right-angle glass microprism and a 4 mm × 4 mm or 3 mm × 4 mm cover glass window at
the surface-end position of the imaging window (Figure 1C). Dental cement was used to
fix this assembly at the correct position for NTS imaging. The cover glass enhanced
adhesion to the dental cement, thereby securing the position of the double microprism
assembly and preventing dental cement from obstructing its optical path. We implanted
the assembly (simply termed "double prism") between the cerebellum and the brainstem
using a custom-made vacuum-based manipulator (Figure 1D and Figure S1A–C). To
implant the double prism, the skull over the dorsal surface of the cerebellum was carefully
removed using a dental drill and fine forceps (Figure S1D–E). A micro-knife was used to
incise the thick dura mater between the posterior edge of the cerebellum and the spinal
cord. Removing the dura mater over the cerebellum not only allows for smooth insertion
of the prism but also adequately relieves the pressure that might occur when the prism is
inserted between the cerebellum and brainstem.
Next, the newly developed vacuum-based prism holder was used to insert and implant
the double prism between the cerebellum and brainstem (Figure 1D–E, Figure S1C, F–
G). Sufficient space and no neural fibers remained between the cerebellum and brainstem
around the NTS area, enabling smooth insertion of the prism without cutting neural fibers.
Using a stereotaxic frame and vacuum-based prism holder attached to an equipped
manipulator, the prism can be precisely operated during the implantation while being
maintained in a horizontal position. The vacuum-based prism holder has a hook on one

side (Figure 1D and Figure S1B–C), which exerts a pushing force when sliding the sharp

corner of the double prism between the cerebellum and brainstem from the caudal side to

the rostral side (Figure 1D–E). After slowly inserting the double prism to approximately

1.6 mm deep, we waited a few minutes to ensure that breathing and other physical

conditions were not disturbed (Figure 1E). The double prism was slid further down at 0.1-

mm increments until reaching a maximum depth of 2.0 mm. Any physical reactions such

as disrupted breathing occurring during the insertion led us to promptly retract the prism.

Setting a stereomicroscope at an angle such that the brainstem can be visualized through

the double prism allows for clear observation of the arrangement of blood vessels in the

brainstem and fourth ventricle while inserting the prism.

After the double prism was fully inserted, UV craft resin was used to fix the double

prism and coat the surface of the targeted nerve structures (Figure S1G), followed by tight

fixation with dental cement. A resin coating was applied with the double prism remaining

in the holder and maintaining sufficient contact with the brainstem target area (including

the NTS) to prevent the resin from getting between them. After the UV resin had cured,

the holder was carefully released, and the double prism was firmly fixed to the

surrounding areas, including the remaining skull bone rostral to the cerebellum, using

dental cement. For further head fixation for two-photon imaging under the microscope

objective, the head plate was attached to the skull with dental cement (Figure 1F).

The double prism allows for optical access to the brainstem located on the ventral side

of the cerebellum, which exceeds 2.0 mm in thickness. Using the wide field of view and

horizontal coordinates, the position of the fourth ventricle serves as a key reference for

locating the NTS. Two-photon imaging through the double prism enabled us to detect

neural activities in the NTS at single-cell resolution, visualized by GCaMP (Figure 1G).

We named this method "D-PSCAN" as explained above.

We confirmed the specificity of the method for detecting NTS neural activities. A

primary purpose for developing this imaging method was to investigate how information

from afferent vagus nerve inputs is processed in the NTS. On the other hand, the dorsal

motor nucleus of the vagus nerve (10N, also known as DMNV or DMV), where the cell

bodies of the vagal efferent fibers originate, is located just ventrally to the NTS[6],

suggesting the importance of confirming specificity in the NTS and distinguishing NTS

neurons from 10N neurons. Importantly, the two-photon imaging technique offers

superior spatial resolution at depth. Based on this advantage, we confirmed that the

obtained signals originated from the brainstem surface area, i.e., the NTS, rather than

deeper regions such as 10N. To further test the specificity, we prepared fixed brain slices

after the *in vivo* imaging experiments and checked the expression of GCaMP within the

NTS. We used ChAT, an immunohistochemical marker, to identify 10N neurons[35] and

distinguish NTS neurons histologically. Expression of GCaMP was not observed in

ChAT-positive cells within 10N (Figure 1B), indicating that NTS neuron activities can

be detected separately from efferent vagus neuron activities using this method. We also

checked for unanticipated retrograde labeling of the vagal afferent terminals onto the NTS.

Cell bodies of vagus neurons projecting to the NTS are located in the nodose ganglion[6,36].

In mice that showed GCaMP expression in the NTS neurons, we observed no GCaMP-

expressing cell bodies in the nodose ganglion, i.e., vagus sensory neurons (Figure S2).

These findings demonstrated that, when imaging neural activities around the NTS region

using this method, we can distinguish activities of NTS neurons from the signals of vagal

afferent nerve terminals. Thus, the D-PSCAN method enables us to specifically evaluate

the activities of the NTS neurons separately from the activities of vagal efferent neurons

and vagal afferent fiber terminals.

In vivo imaging of neural population activities at single-cell resolution in the brainstem

(NTS) of live mice

Using the D-PSCAN method, we performed *in vivo* two-photon imaging and recorded

the neural population activities of the NTS and surrounding brainstem areas. Spontaneous

activities and stimulus-evoked responses of the brainstem neurons were recorded by

imaging GCaMP fluorescence changes ($\Delta F/F0$). Brain motion artifacts interfered with

imaging in awake mice but were minimized under anesthesia, enabling stable detection

of the neural activities at a single-cell resolution both inside and outside of the NTS area

(Figure 2 and Supplementary Movie 1, 2). Time-lapse imaging at various depths was

performed by adjusting the imaging focal plane using a piezo to move up and down in

depth (Figure 2A and Figure S3). Volumetric live imaging (2 mm x 2 mm horizontally,

or smaller for zooming, 200–480 μm dorsoventrally, at 0.4–1.0 Hz temporal resolution)

of neural activities in the wide brainstem area (including the NTS) of the head-fixed

mouse was performed through the implanted double prism. Importantly, this wide field
of view and piezo-based volumetric imaging allowed us to detect GCaMP fluorescence
and neural activities not only from inside but also from outside of the NTS areas and to
confirm the specificity of the stimulus-evoked responses in the NTS, as explained below.
233

<u>Detection of single cell activities in the NTS as the direct efferent target of the vagus</u>
<u>nerve</u>

To demonstrate the usefulness of the D-PSCAN for investigating the physiological
characteristics of the NTS neurons, we applied the method to observe the neural responses
to artificial electrical stimulation of the afferent vagus nerve under various conditions
(vagus nerve stimulation, or VNS). We tested whether NTS neurons respond to the stimuli
and further evaluated the diversity of their responsiveness. To perform VNS, the bipolar
platinum hook electrode that has been used in rodents[37] was employed (Figure 2B). After
isolating the vagus nerve from the carotid artery, a bipolar platinum hook electrode was
wrapped around the left vagus nerve and fixed in place without contacting other tissues
using the insulating silicone adhesive (Kwik-Sil)[38,39]. The electrode was positioned to
keep the positive and negative poles apart. This setup allows for stable activation of the
vagus nerve while preventing irritation to the surrounding tissue. To create an anodal
block for the efferent direction, the cathode and anode were located on the rostral and
caudal sides, respectively, of the vagus nerve. The mechanism of the anodal block is
proposed to selectively bias activation toward vagal afferent fibers[40-44]. We selectively
targeted the left vagus nerve, which is usually chosen for safety reasons, as stimulation
of the right vagus nerve has a greater impact on heart rate[45,46]. We conducted *in vivo* NTS
imaging experiments after closing the surgical wound.

We imaged the *in vivo* neural activity in the brainstem as VNS was repeatedly applied.
The VNS parameters were established based on previous studies, and various conditions
(i.e., various current intensities and stimulus frequencies) within the ranges reported in
prior rodent and human studies[13,42,43,47] were used to investigate the neural response
properties. Figure 2C presents an example in which the stimulation intensity was set to
1.0 mA and the stimulation frequency was 10 Hz. A pulse width of 100 microseconds
(μsec) was consistently selected across all experiments in the present study. At a stimulus

frequency of 10 Hz, stimulation (100-µsec current) occurred 10 times per second. The

term "trial" is defined as each 10-sec VNS duration, while the term "session" is defined

as a series of VNS trials at each condition (e.g., 1.0 mA at 10 Hz, 0.5 mA at 10 Hz, 1.0

mA at 5 Hz, etc.). In each session, the inter-trial interval was set at 50 sec. In the example

cases shown in Figure 2, 10-sec VNS was repeated multiple times within a session to

assess the neural responses to each VNS condition.

Taking advantage of the 2 mm x 2 mm large field of view provided by the double

prism, we conducted volumetric observations of a wide brainstem area, including the NTS,

and compared neural responses to VNS inside and outside of the NTS (Figure 2D). In an

example case (Figure 2D), we scanned a total of 16 z-planes at each time point. We

stacked every 4 consecutive plane from top to bottom to create an average image for each

time point with carefully correcting the motion artifact (see Methods for details). This

stacking was done with a two-plane overlap; for instance, the first stack included the 1st

to 4th z-planes, the second stack included the 3rd to 6th z-planes, and subsequent stacks

followed this pattern. As a result, we obtained 7 stacked planes in total in this experiment,

labeled Zp1 through Zp7, respectively. This procedure is advantageous to enhance the

signal to noise ratio in each movie, which is critical for further automatic detection of the

active neurons[48]. The motion-corrected movies were analyzed to automatically detect the

active neurons using a constrained nonnegative matrix factorization algorithm, as

described previously[48,49]. This detection method is advantageous in isolating the activity

of the individual neuron, even when other active neurons are nearby or overlapped within

the field of view[48] (see Methods for details). After automatic detection of the active

neurons, we checked the existence of overlapped detection of the regions of interest (ROIs,

i.e., active neurons) between adjacent Zps, and removed such overlaps by manual

inspection.

Temporal dynamics of neural activity were clearly observed within the field of view

at multiple depths (Figure 2D and Supplementary Movie 1). Here, we present the GCaMP

signals at Zp1, Zp2, and Zp4. Neurons exhibiting stimulus-evoked and/or spontaneous

neural activity were automatically identified, and representative neurons are shown in the

graph. In the case of neurons outside of the NTS, we observed no clear responses to VNS

and detected only spontaneous (stimulus-independent) activities (neurons #1 and #2 in

Figure 2D). On the other hand, we detected a large number of neurons responding to VNS

in the NTS area (neurons #3–#21).

Two-photon imaging utilizes a scanning-type observation system, enabling seamless

transition from wide-field to high-magnification imaging using a single objective lens.

Depending on the point spread function of the objective lens, this approach facilitates the

acquisition of images with enhanced spatial resolution compared with maximum wide-

field imaging. Using this mechanism, after confirming the position of the NTS in the wide

field of view, we narrowed down the imaging area to the NTS and observed the neural

activities at high magnification (Figure 2E and Supplementary Movie 2). We detected

individual neuron cell bodies more clearly through visual inspection at higher

magnifications compared with lower magnifications, confirming that most neurons

identified at lower magnification were indeed NTS neuron cell bodies, which exhibited

clear responses to VNS.

Note that there was variability in VNS responsiveness among neurons detected within

the NTS (Figure 2D and E), prompting us to further examine their characteristics more

thoroughly by varying the stimulus conditions.

<u>Diverse neuronal responses to VNS in the NTS</u>

In vivo brainstem imaging was performed during VNS administered at various

intensities and frequencies to explore how NTS neurons respond to different intensities

and frequencies. Additionally, we assessed the responsiveness of the NTS neurons to

repetitive stimulation across different intensities and frequencies.

First, we set the stimulus frequency at 10 Hz and varied the stimulus intensities

(Figure 3A–E and Figures S4A, S5). Most of the stimulus-responsive NTS neurons

exhibited enhanced responses to higher stimulus intensities. On the other hand, one

neuron detected outside of the NTS showed only a spontaneous activity change and no

clear response to VNS (neuron #2 in Figure 3 and Figures S4, S5). NTS neurons showed

minimal responses to VNS at 0.1 mA and 0.25 mA and there was no clear difference

between these intensities, while there was a substantial increase in the neural response at

0.5 mA. The difference between 0.5 mA and lower intensities suggests a possible

threshold between them for recorded NTS neurons. At 1.0 mA, the responses increased

further. We also calculated various parameters for evaluating the neural responses and
observed consistent tendencies (Figure 3E).

We next evaluated the effect of repetitive stimulation on the neural responses at
various stimulus intensities (Figure 3C–D and Figure S4A). At 0.1 mA and 0.25 mA, we
did not observe a change in the responses to repeated stimulation (50-sec intervals) at
each stimulus intensity. On the other hand, at higher intensities of 0.5 mA and 1.0 mA,
some neurons exhibited gradually enhanced responses during the session with repeated
stimulations.

Next, using the same recorded neurons, we set the stimulus intensity at 1.0 mA and
varied the stimulus frequencies (Figure 3F–H and Figure S4B). The neuronal responses
to VNS increased gradually as the stimulus frequency was increased from 1 Hz to 10 Hz,
similar to when the stimulus intensity was increased. Interestingly, increasing the
frequency to 15 Hz resulted in a reduction in the neural response compared with that at
10 Hz. We also investigated the effects of repetitive stimulation during each session at
various frequencies. At 1 Hz, 2.5 Hz, and 5 Hz, we did not observe clear changes in
response to repeated stimulation during each session. At 10 Hz, however, most of the
NTS neurons showed a gradually increased response. Notably, at a higher frequency of
15 Hz, despite similar initial responses to those at 10 Hz, neurons exhibited decreased
responses to subsequent stimuli during the session.

These results indicate that repeated stimulation may uniquely affect synaptic
interactions and signal transmission between the vagal afferent nerve and NTS neurons.
While we consider the physiological mechanisms behind the observed acceleration and
deceleration below in the discussion section, our findings also suggest that the frequency-
dependent increase and decrease in responses to VNS are not merely artifacts caused by
the slow off-kinetics of GCaMP. Changes in neural activities and associated calcium
concentrations occur more rapidly than the signal changes detected by calcium indicators
like GCaMP. This discrepancy can theoretically lead to an artificial amplification of
baseline signals during repetitive stimuli, which may account for the increase in
fluorescent signals at 10 Hz. Our imaging system, however, reliably detected an opposing
decrease in the GCaMP signal in response to repetitive stimulation at 15 Hz, despite
similar initial response levels to those observed at 10 Hz. Thus, the enhanced responses

at 10 Hz are also likely to reflect the actual responses of the neurons, rather than being

artifacts.

Together, these findings indicate that the D-PSCAN method effectively detects

various types of neurons in the NTS under different stimulus conditions. This method

demonstrated that specific VNS conditions can directly and effectively modulate NTS

neuron activities, potentially influencing whole-brain neural networks involved in the

regulation of emotion.

<u>Chemical stimulation triggered both immediate and long-lasting responses in NTS</u>

<u>neurons</u>

Up to this point, we examined the response of NTS neurons to artificial electrical

stimulation of the vagal afferent nerve. Emotional responses are also governed by more

physiological stimuli and conditions. To assess the suitability of the D-PSCAN method

for capturing responses under physiological conditions, we measured the activity of NTS

neurons following intravenous administration of CCK (Figure 4A). CCK is a hormone

released from the gastrointestinal tract after meals[50]. It directly influences the afferent

fibers of the vagus nerve[51] and is proposed to transmit signals via the NTS[51,52] to various

brain regions, including the feeding center.

To assess whether the intravenous administration process itself affects neural activity,

we also monitored responses during saline administration as a negative control. The

timing of the intravenous administration was determined by including SR101 in the

injected solution and observing the signal detected in the blood vessels surrounding the

NTS using *in vivo* two-photon microscopy (Supplementary Movie 3).

With this method, we first demonstrated that saline administration did not induce a

clear change in neural activity (Figure 4B–C and Supplementary Movie 4). In contrast,

CCK administration induced a change in NTS neuron activities (Figure 4B–C and

Supplementary Movie 5); upon administering CCK, we immediately observed clear and

significant signal changes in NTS neurons (Figure 4B–E). This immediate response of

the NTS neuron is consistent with a previous study demonstrating the responsiveness of

vagal afferents to CCK[53]. Additionally, for more than 5 minutes after CCK administration,

we observed a higher frequency of spontaneous activity compared with before CCK

administration or after saline administration (Figure 4C, F and Figure S6). This prolonged

activation in the NTS neuron is also consistent with the previous observation in vagal

afferent fibers[53]. These results demonstrate that the D-PSCAN method is applicable to

the detection of changes in NTS neuron activity induced under physiological conditions.

Discussion:

In the present study, to understand how the information processing mechanisms of

brain-viscera pathways mediated by the NTS regulate emotions, we developed a new

imaging technique named D-PSCAN based on *in vivo* two-photon imaging and double

prism-based optical access. This method enables wide-field observation of deep-brain

areas in the brainstem, including the NTS, while keeping the cerebellum, an important

region for emotional regulation, intact. We confirmed that the D-PSCAN method allows

for specific identification of NTS neurons, distinguishing them from surrounding regions

based on horizontal coordinates in its wide field of view. Additionally, based on the

advantages of two-photon imaging, we confirmed that the obtained signals originated

from the surface brainstem area, i.e., the NTS, rather than deeper regions such as 10N.

As we observed no GCaMP expression in 10N, it appears unlikely that vagal afferent

terminals were labeled by GCaMP, ensuring almost no or undetectable contamination of

their signal in the observed NTS areas. These findings together indicate that our method

provides specific identification of NTS activities and assesses their responsiveness to

VNS and CCK.

Using this method, we observed the responses of mouse NTS neurons to VNS, a

technique clinically employed to treat neuropsychiatric disorders in humans. We

confirmed that responses to electrical stimulation depended on the intensity and

frequency, observed a threshold-like effect, and found that repeated excessive stimulation

could reduce the responses. In addition to responses to artificial stimuli like VNS, we

detected significant immediate responses and smaller but sustained activities following

physiological stimuli such as CCK. These results demonstrate the usefulness of the D-

PSCAN method for investigating how the NTS neurons process information from visceral

organs for emotional regulation.

Recent studies have advanced our understanding of the complex mechanisms by

which the NTS processes various types of visceral information[22,23,30,36,54]. How these

mechanisms underlie the therapeutic effects of VNS and regulate emotion, however,

remains unclear. Our imaging technique, which allows for detailed observation of NTS

neuron activities while maintaining the integrity of cerebellar circuits crucial for

emotional regulation[31-34], is expected to be instrumental in deepening our knowledge of

these complex mechanisms.

Repeated VNS induced both enhanced and diminished NTS responses, depending on

the stimulus conditions. As discussed above, the observed NTS responses are unlikely to

be artifacts caused by the calcium indicator (GCaMP). These results indicate that repeated

stimulation may uniquely affect synaptic interactions and signal transmission between

the vagal afferent nerve and NTS neurons. Inhibitory interneurons in the NTS are

involved in information processing during mechanical stimulation of digestive organs[30,55],

suggesting the possible contribution of such complex circuitries within the NTS for

adjusting the VNS signal. The activity of NTS neurons during VNS might also be

modulated through interactions with other brain regions, including supramedullary

regions[56]. Therefore, our imaging method will be useful for future investigations into

these detailed regulatory mechanisms of the NTS. While an anodal block is likely to

enhance specificity in targeting vagal afferent fibers, VNS might also weakly stimulate

efferent fibers, which in turn might stimulate visceral organs and further activate the vagal

afferent fibers. This may also influence the rates of the neuron responses. Overall, as

optimizing VNS parameters for human patients is critical[13,14], our NTS imaging method

in mice could help reveal the underlying mechanisms of VNS, leading to more effective

and personalized therapy settings.

Previous electrophysiological studies suggest that CCK induces prolonged

activation of the vagal afferent nerve[53,57]. The CCK-evoked enhancement of the

spontaneous fluctuations in the GCaMP signal lasting for more than 5 minutes is

consistent with these electrophysiological observations. These results demonstrate that

the D-PSCAN method has significant potential to detect changes in NTS neuron activity

under physiological conditions.

Overall, our development of the wide-field and deep-brain two-photon imaging

method utilizing a prism-based optical interface marks a significant advancement in the

ability to study the NTS *in vivo* while maintaining an intact cerebellum. Future

applications of this imaging system, particularly in awake animals, will be crucial for

understanding the dynamic interactions within the NTS and its role in neuropsychiatric

disorders. By enabling detailed observation of NTS activities, this method offers a

powerful tool for advancing our understanding of the vagus nerve's impact on brain
function and emotional health, potentially leading to improved clinical interventions for
neuropsychiatric conditions.
453

Acknowledgements
We thank Okito Hashimoto for technical advice and discussion. We also thank the
members of the laboratory for their help, especially Ikuko Takeda, Yuki Ichihashi, Ryo
Fujikawa, and Fumiko Kito for experiments and data analyses. This study was supported
by the JSPS KAKENHI Grant (grant number JP18K06536, JP20H05076, JP21H02801,
JP22H05081, JP22H05519, JP24H01260 to M.A.), the Cooperative Study Program of
National Institute for Physiological Sciences (19-233, 20-140 to M.S.; 21-244,
22NIPS218 to D.Y.), the practical research project for rare/intractable diseases from
Agency for Medical Research and Development (AMED; Grant #: 20ek0109315h003 to
M.S.), and the Research Grant for Neurological and Psychiatric Disorders from National
Center of Neurology and Psychiatry, Japan (NCNP; Grant #: 30-3, 1-1, and 2-6 to M.S.).
465

Author Contributions
M.A., D.Y. and M.S. conceived and coordinated the whole project. M.A. designed and
developed a method for double-prism-based *in vivo* two photon imaging with the support
of T.K. and J.N.; M.A., D.Y., H.K., E.T. and M.S. performed imaging experiments during
the VNS and CCK-administration; M.A., A.H., D.Y., and S.T. performed data analyses
with the support of C.I., A.N., H.Y., A.S. and M.S.; M.A., A.H., D.Y., A.N., Y.A., and
M.S. wrote the paper, with contributions from all authors.
473

474 **Data availability**
475 The data that support the findings of this study are available from the corresponding
476 author upon reasonable request. The anatomical information in the Allen Brain Atlas
477 (https://atlas.brain-map.org/) was used for the anatomical description, determination of
478 the virus injection area, and evaluation of the recorded brain regions.
479
480

481 **<u>Methods</u>**
482

Animals. All animal experiments were carried out in accordance with National Institutes
of Health guidelines and approved by the National Institute for Physiological Sciences
Animal Care and Use Committee (approval number 18A102), or in accordance with the
National Institute of Neuroscience, National Center of Neurology and Psychiatry (Tokyo,
Japan) and were approved by the Institutional Animal Investigation Committee (approval
number: 2020004). Male C57BL/6 mice housed under a 12-h light/dark cycle with free
access to food and water were used for all experiments. Mice at 3–10 months of age were
used for the experiments.
491
Virus injection. Nucleus tractus solitarii (NTS) is a structure distributing in the dorsal
medulla near the area postrema. To express GCaMP, a genetically encoded calcium
indicator for monitoring neural activity, we used a gene expression system based on the
AAV vector. We tested AAV1/Syn.GCaMP6s, AAV1/Syn.GCaMP6f, and
AAVdj/Syn.GCaMP7f, which were produced as described previously[58,59] using plasmids
obtained from Addgene. As the GCaMP was expressed under the regulation of the Syn
(Synapsin) promoter, all of the recording targets were assumed to be neuron-specific. All
of the viruses allowed us to visualize the neural activity in the NTS, and type of virus
used in each experiment is indicated in each corresponding figure legend. Viruses were
injected into mice at approximately postnatal day (P) 50–60 for experiments at least 1
month before the microprism implantation, which was followed by *in vivo* imaging.
Injection procedures were performed as described previously[60], with some modifications
to target the NTS efficiently. During surgery, the mice were anesthetized with isoflurane
(initially 2% [partial pressure in air] and then reduced to 1%). A small circle (~1 mm in
diameter) of the skull was thinned over the cerebellum (around the injection site) using a
dental drill to mark the site for a small craniotomy. Viruses were injected into the bilateral
NTS regions at two sites (depth 4.7 and 3.7 mm from the dorsal surface of the cerebellum,
volume 250 nl/site, at 100 nl/min) to sufficiently cover the NTS, over a 5-min period at
each depth using a UMP3 microsyringe pump (World Precision Instruments). The X-Y
coordinates for the injection site were usually 0.5 mm lateral to the midline and 3.3 mm
caudal to lambda. The beveled side of the injection needle was faced to the caudal so that
the needle could be smoothly inserted into and passed through the cerebellum, and the
AAV would cover the NTS similarly in both hemispheres. We designed our injection
protocol (especially the volume and depth) carefully to widely cover the area around the

NTS, while the anatomical coordinates of the field of view for the two-photon imaging

were precisely targeted using visual guide through the microprism implanted as described

below (see the section "*In vivo* two-photon imaging in the wide brainstem area"). Also,

as shown in Figure 1B, the expression of GCaMP was excluded in the 10N area, which

placed next to and ventrally to the NTS as marked by the ChAT immunostaining. The

reason for the exclusion is unclear, but it may be because the promoters and/or serotypes

used in the present studies do not match for inducing the GCaMP expression in these

ChAT-positive neurons. In any case, since neurons in the 10N area receive synaptic inputs

from NTS neurons[6,61] and are a source of vagal efferents, the lack of expression in the

neurons of this area rather positively aids in the interpretation of the GCaMP signals

detected by *in vivo* NTS imaging. In some mice, a deeper area (more than 0.5 mm from

the dorsal surface of the brainstem, i.e., the surface of the microprism) also expressed the

GCaMP when the fluorescence was checked in the brain slices (Figure 1B), but this

deeper area was beyond the deepest limit of our two-photon imaging system for the

detection of the sufficient signal intensity through the double prism *in vivo*. These results

indicates that our recording system simply allows specific targeting of the neurons within

the NTS and together with laterally-surrounding wide areas.

***In vivo* two-photon imaging in the wide brainstem area**. We developed a novel,

minimally invasive method based on the double prism, which allows *in vivo* two-photon

imaging of deep brain regions, specifically around the NTS. This innovative approach

also enables simultaneous vagus nerve stimulation (VNS) or intravenous administration

of cholecystokinin-8 (CCK).

In vivo two-photon imaging was performed as described previously[49,60], with some

special development for the optical access to the brainstem and NTS as explained below.

Before the surgery, we designed an implantable double microprism assembly, comprising

a 2-mm right-angle glass microprism (TS N-BK7, 2 mm AL+MgF2, Edmund) bonded to

another 2-mm right-angle glass microprism and a 3 mm x 3 mm, 3 mm × 4 mm or

4 mm × 4 mm cover glass window (No.1; Matsunami) at the surface-end position of the

imaging window, was prepared. Dental cement was used to fix this assembly at the correct

position for NTS imaging. The cover glass was used to enhance adhesion to the dental

cement, securing the position of the double microprism assembly and preventing dental

cement from obstructing its optical path. We implanted the assembly (or simply termed

"double prism") between the cerebellum and the brainstem using a custom-made vacuum-

based manipulator (Figure 1D and Figures S1A–C).

At 1–8 months after the virus injection, the mice were anesthetized with isoflurane

(initially 2% [partial pressure in air] and reduced to 1% during the surgery). For the

implantation of the double prism, the skull over the dorsal surface of the cerebellum was

carefully removed using a dental drill and fine forceps (Figure S1D–E). A micro-knife

was used to incise the thick dura mater between the posterior edge of the cerebellum and

the spinal cord. Removing the dura mater over the cerebellum not only allows for smooth

insertion of the prism but also adequately relieves the pressure that might occur when the

prism is inserted between the cerebellum and the brainstem.

Next, the newly developed vacuum-based prism holder was used to insert and implant

the double prism between the cerebellum and the brainstem (Figure 1D–E, Figure S1B–

C, F). Sufficient space and no neural fibers remained between the cerebellum and the

brainstem around the NTS area, enabling smooth insertion of the prism without cutting

neural fibers. Using a stereotaxic frame and vacuum-based prism holder attached to an

equipped manipulator, the prism can be precisely operated during the implantation while

being maintained in a horizontal position. The vacuum-based prism holder has a hook on

one side (Figure 1D and Figure S1B–C), which exerts a pushing force when sliding the

sharp corner of the double prism between the cerebellum and the brainstem from the

caudal side to the rostral side (Figure 1D–E). After slowly inserting the double prism to

approximately 1.6 mm deep, we waited a few minutes to ensure that breathing and other

physical conditions were not disturbed (Figure 1E). The double prism was slid further

down at 0.1-mm increments until reaching a maximum depth of 2.0 mm. Any physical

reactions such as disrupted breathing occurring during the insertion led us to promptly

retract the prism. When retracting the double prism assembly, it is crucial to ensure that

the suction force by the vacuum is sufficient, as insufficient suction may lead to failure

of the retraction. The area directly beneath the microprism was minimally compressed

but remained intact, minimizing motion artifacts caused by the animal's breathing. Setting

a stereomicroscope at an angle such that the brainstem can be visualized through the

double prism allows for clear observation of the arrangement of blood vessels in the

brainstem and fourth ventricle while inserting the prism. Bleeding was rarely observed

during the implantation procedure.

After the double prism was fully inserted, UV craft resin was used to fix the double
prism and coat the surface of the targeted nerve structures (Figure S1G), followed by tight
fixation with dental cement. A resin coating was applied with the double prism remaining
in the holder and maintaining sufficient contact with the brainstem target area (including
the NTS) to prevent the resin from getting between them. After the UV resin had cured,
the holder was carefully released, and the double prism was firmly fixed to the
surrounding areas, including the remaining skull bone rostral to the cerebellum, using
dental cement. For the further head fixation for two-photon imaging under the microscope
objective, the head plate was attached to the skull with dental cement (Figure 1F).

The double prism allows for optical access to the brainstem located on the ventral side
of the cerebellum, which exceeds 2.0 mm in thickness. Using the wide field of view and
horizontal coordinates, the position of the fourth ventricle serves as a key reference for
locating the NTS. Two-photon imaging through the double prism enabled us to detect
neural activities in the NTS at the single-cell resolution, visualized by GCaMP (Figure
1G). We named this method "D-PSCAN", or Double-Prism based brainStem imaging
under Cerebellar Architecture and Neural circuits.

During the consecutive imaging sessions, mice were anesthetized with isoflurane
(1%) or a ketamine (70 mg/kg) and xylazine (10.5 mg/kg) mixture (see the sections below
for more details). Experiments were performed in a completely dark environment. The
activity of NTS neurons was recorded by imaging fluorescence changes with a
multiphoton microscope (Nikon A1R MP, Nikon) and a Mai Tai DeepSee Ti:sapphire
laser (Spectra-Physics) at 920 nm, through a 4x dry objective, 0.28 N.A. (Olympus).
Scanning and image acquisition were controlled by NIS-elements software (Nikon).
Brain motion artifacts interfered with imaging in awake mice but were minimized under
anesthesia, enabling the stable detection of the neural activities at a single cellular
resolution both in and out of the NTS area (Figure 2, Figure S3 and Supplementary Movie
1 and 2). Time-lapse imaging at various depths was performed by adjusting the imaging
focal plane using a piezo to move up and down in depth (Figure 2A and Figure S3).
Volumetric live imaging (2 mm x 2 mm horizontally, or smaller for zooming, 200–480
μm dorsoventrally, at around 0.4–1.0 Hz temporal resolution) of neural activities in the
wide brainstem area (including the NTS) of the head-fixed mouse was performed through
the implanted double prism. Importantly, this wide field of view and piezo-based

volumetric imaging allowed us to detect the GCaMP fluorescence and neural activities

not only from inside but also from outside of the NTS areas and confirm the specificity

of the stimulus-evoked responses in the NTS, as explained below.

Vagus nerve stimulation during NTS imaging. For the NTS imaging experiments with

vagus nerve stimulation (VNS), a bipolar platinum hook electrode, connected to a

programmable stimulator (SEN-8203, NIHON KODEN, Tokyo, Japan) and isolator (SS-

202J, NIHON KODEN), was wrapped around the cervical vagal trunk, and was fixed and

insulated from surrounding tissue using the quick-setting silicone adhesive Kwik-Sil

(World Precision Instruments, FL, USA). After separating the vagus nerve from the

carotid artery, a bipolar platinum hook electrode was wrapped around the left vagus nerve

and fixed in place without contacting other tissues using the insulating silicone adhesive,

Kwik-Sil[38,39]. The electrode was positioned to keep the positive and negative poles apart.

This setup allows for stable activation of the vagus nerve while preventing irritation to

the surrounding tissue. To create an anodal block for the efferent direction, the cathode

and anode were located on the rostral and caudal sides, respectively, of the vagus nerve.

The mechanism of the anodal block is proposed to selectively bias activation toward vagal

afferent fibers[40-44]. We selectively targeted the left vagus nerve, which is usually chosen

for safety reasons, as stimulation of the right vagus nerve has a greater impact on heart

rate[45]. We conducted *in vivo* NTS imaging experiments after closing the surgical wound.

The mouse's head was fixed under the objective lens by using the head plate tightly

attached to the skull under anesthesia (a ketamine [70 mg/kg] and xylazine [10.5 mg/kg]

mixture). VNS was triggered using custom Labview software. To determine optimal

stimulation parameters VNS was delivered in blocked trials for 10 s at the frequencies of

1.0, 2.5, 5.0, 10, or 15 Hz, at the intensities of 0.1, 0.25, 0.5, or 1.0 mA, with a pulse width

of 100 µsec. The maximum current intensity and frequency of stimulation were set at 1.0

mA and 10 Hz according to the previous study[37,47]. The first VNS train was delivered at

least 10 s after the start of imaging, and VNS was repeated three times in total (Figure 3

and Figures S4, S5), or more (Figure 2), with a 50-sec interval. Neither facial nor muscle

twitching (an indication of off-target stimulation of musculature surrounding the vagus

nerve) was observed with any parameters used in any of the mice included in the present

study. VNS and NTS imaging were performed with the mouse head fixed under the

microscope objective lens using the head plate. The timings of the imaging and VNS were

synchronously recorded with image acquisition (GCaMP imaging in NTS) using NI

software (Labview; National Instruments) and NI-DAQ (National Instruments).

Intravenous administration of CCK during NTS imaging. To examine the neural

activity change in response to more physiological stimulation than VNS, we employed

cholecystokinin octapeptide (CCK); it has been demonstrated that intravenous

administration of CCK activates vagal afferents and increases firing rate of hepatic branch

of the vagus nerve[53] (Cox and Randich, 1997). On the day of the experiment, mouse's

head was fixed under the objective lens by using the head plate tightly attached to the

skull under anesthesia (1% isoflurane). During the Ca^{2+} imaging, CCK solution (2.0

μg/kg) or saline was mixed with sulforhodamine 101 (SR101; 0.1 mM) and intravenously

administered through the tail vein. 60 sec after the start of the imaging, saline with SR101

was intravenously administered and performed for 10 minutes. Using the same animal,

60 sec after the beginning of the imaging, CCK with SR101 was intravenously

administered and performed for 15 minutes. Intravenous administration and NTS imaging

were performed with the mouse head fixed under the microscope objective lens using the

head plate.

Imaging data analyses and statistics.

The activity of NTS neurons was recorded by imaging fluorescence changes with a

multiphoton microscope (Nikon A1R MP, Nikon). Time-lapse imaging at various depths

was performed by adjusting the imaging focal plane using a piezo to move up and down

in depth (Figure 2A and Figure S3). The volumetric live imaging (at around 0.4–1.0 Hz

temporal resolution) of NTS neural activities was performed through the implanted

double prism. We recorded a total of 10–16 focal planes (z-planes) at each time point for

each mouse. In an example case (Figure 2D), we recorded a total of 16 z-planes at each

time point. We stacked every 4 consecutive plane from top to bottom to create an average

image for each time point. This stacking was done with a two-plane overlap; for instance,

the first stack included the 1st to 4th z-planes, the second stack included the 3rd to 6th z-

planes, and subsequent stacks followed this pattern. As a result, we obtained 7 stacked

planes in total in this experiment, labeled as Zp1 through Zp7, respectively. This

procedure is advantageous to enhance the signal to noise ratio in each movie, which is

critical for further automatic detection of the active neurons[48]. After the automatic

detection of the active neurons (details for this detection method are described below),

we checked the existence of the overlapped detection of the regions of interest (ROIs, i.e.,

active neurons) between adjacent Zps next each other, and removed such overlaps by the

manual inspection.

Importantly, before stacking 4 consecutive z-planes at each time point as explained

above, they were processed to correct for brain motion artifacts using the enhanced

correlation coefficient image alignment algorithm[62]. For this z-axis correction, first (most

dorsal surface frame) was used as a reference for correcting the second z-plane. Then, an

average image of these two aligned z-planes was further used for the correction of the

third z-plane. Eventually an average image of these three aligned z-planes was used for

the motion correction of the fourth z-plane. Through this procedure, four z-planes were

motion corrected and used to accurately obtain a stacked (averaged) image for each Zp at

each time point. Next, the time-series of stacked GCaMP images at each Zp was further

processed to correct for brain motion artifacts using the enhanced correlation coefficient

image alignment algorithm[62]. Temporal dynamics of neural activity were clearly observed

at multiple depths after this motion correction procedure (Supplementary Movies 1 and

2).

These motion-corrected movies were used for automatic ROI detection. The ROIs for

the detection of neural activity were automatically selected using a constrained

nonnegative matrix factorization algorithm in MATLAB R2014a/R2019b as described

previously[48,49], with some manual adjustments as described above. This detection method

is advantageous in isolating the activity of the individual neuron, even when other active

neurons are nearby or overlapped in the field of view[48]. The same constrained

nonnegative matrix factorization package for ROI detection also provides an option for

signal processing and measurement of the signal change ($\Delta F/F$) of each neuron. We used

this method to identify neurons and calculate $\Delta F/F$ for them. While neurons for the

analyses were initially automatically detected, neurons responding to noisy signals with

no apparent calcium transient at any time during the experimental days were identified

by visual inspection and excluded from further analysis. To apply the same regions of

interest (ROIs) for analyzing images obtained across multiple conditions (i.e., various

VNS conditions or saline vs CCK experiments), the movies from the same mouse and the

same field of view were precisely aligned with each other using the same enhanced

correlation coefficient algorithm as above, and these multiple movies of the same field of

view were tentatively concatenated for the ΔF/F measurement. The obtained ΔF/F data

were eventually separated to evaluate the neural responses under each condition.

For the statistical analysis, we used MATLAB R2022a/R2023b. The Wilcoxon

signed-rank test for a paired comparison and the Kolmogorov-Smirnov test for a two-

group comparison with pooled data (from all neurons) were used to determine statistical

significance ($p < 0.05$) in Figure 4. Two-tailed tests were selected for all statistical

analyses. Graphs were produced by MATLAB R2022a/R2023b.

As for the imaging data analysis, one concern was that the off kinetics of the GcaMP6f

signal might not be sufficiently fast to consider the responses to 3 trails of VNS as

independent data, and thus statistical evaluation based on these responses might not be

reliable to evaluate the neural responses. As shown in Fig. 3G, however, neurons showed

a reduced response to repeated stimulation, indicating that calcium signaling was not

enhanced by delayed off-kinetics.

After the CCK administration, we observed prolonged neural activation, which was

detected as continuous calcium transients (Figure 4 and Figure S6). To evaluate the event

frequency of the transients, we identified the timing of the calcium transients by analyzing

the first derivative of ΔF/F (Figure S6). The first derivatives of each neuron, from the

entire imaging session (including both the saline and CCK experiments), were

thresholded at 0.5 times the standard deviation to detect active time points. The onset of

each event was then used to determine the timing of each transient. The intervals between

calcium transient events were then calculated and used to compare the event frequencies

between the saline and CCK experiments (Figure 4F).

Drugs

CCK was purchased from PEPTIDE INSTITUTE, INC. and was dissolved in saline (2.0

μg/kg). In addition, sulforhodamine (SR101), a red fluorescent dye, is diluted to 0.1 mM

in saline or CCK solution to confirm the successful intravenous administration and the

injection timing during NTS imaging.

References

1. Khalsa, S.S., Adolphs, R., Cameron, O.G., Critchley, H.D., Davenport, P.W.,
Feinstein, J.S., Feusner, J.D., Garfinkel, S.N., Lane, R.D., Mehling, W.E., et al.
(2018). Interoception and Mental Health: A Roadmap. Biol Psychiatry Cogn
Neurosci Neuroimaging *3*, 501-513. 10.1016/j.bpsc.2017.12.004.
2. Chen, W.G., Schloesser, D., Arensdorf, A.M., Simmons, J.M., Cui, C.,
Valentino, R., Gnadt, J.W., Nielsen, L., Hillaire-Clarke, C.S., Spruance, V., et al.
(2021). The Emerging Science of Interoception: Sensing, Integrating,
Interpreting, and Regulating Signals within the Self. Trends Neurosci *44*, 3-16.
10.1016/j.tins.2020.10.007.
3. Dantzer, R., O'Connor, J.C., Freund, G.G., Johnson, R.W., and Kelley, K.W.
(2008). From inflammation to sickness and depression: when the immune
system subjugates the brain. Nat Rev Neurosci *9*, 46-56. 10.1038/nrn2297.
4. Yamada, T., and Katagiri, H. (2007). Avenues of communication between the
brain and tissues/organs involved in energy homeostasis. Endocr J *54*, 497-505.
10.1507/endocrj.kr-106.
5. D'Mello, C., and Swain, M.G. (2011). Liver-brain inflammation axis. Am J
Physiol Gastrointest Liver Physiol *301*, G749-761. 10.1152/ajpgi.00184.2011.
6. Berthoud, H.R., and Neuhuber, W.L. (2000). Functional and chemical anatomy
of the afferent vagal system. Auton Neurosci *85*, 1-17. 10.1016/S1566-
0702(00)00215-0.
7. Beekwilder, J.P., and Beems, T. (2010). Overview of the clinical applications of
vagus nerve stimulation. J Clin Neurophysiol *27*, 130-138.
10.1097/WNP.0b013e3181d64d8a.
8. Bonaz, B., Picq, C., Sinniger, V., Mayol, J.F., and Clarencon, D. (2013). Vagus
nerve stimulation: from epilepsy to the cholinergic anti-inflammatory pathway.
Neurogastroenterol Motil *25*, 208-221. 10.1111/nmo.12076.
9. Reid, S.A. (1990). Surgical technique for implantation of the neurocybernetic
prosthesis. Epilepsia *31 Suppl 2*, S38-39. 10.1111/j.1528-1157.1990.tb05847.x.
10. Penry, J.K., and Dean, J.C. (1990). Prevention of intractable partial seizures by
intermittent vagal stimulation in humans: preliminary results. Epilepsia *31 Suppl
2*, S40-43. 10.1111/j.1528-1157.1990.tb05848.x.

11.   Groves, D.A., and Brown, V.J. (2005). Vagal nerve stimulation: a review of its

774   applications and potential mechanisms that mediate its clinical effects. Neurosci

775   Biobehav Rev *29*, 493-500. 10.1016/j.neubiorev.2005.01.004.

776   12.   Wang, Y., Zhan, G., Cai, Z., Jiao, B., Zhao, Y., Li, S., and Luo, A. (2021). Vagus

777   nerve stimulation in brain diseases: Therapeutic applications and biological

778   mechanisms. Neurosci Biobehav Rev *127*, 37-53.

779   10.1016/j.neubiorev.2021.04.018.

780   13.   Thompson, S.L., O'Leary, G.H., Austelle, C.W., Gruber, E., Kahn, A.T., Manett,

781   A.J., Short, B., and Badran, B.W. (2021). A Review of Parameter Settings for

782   Invasive and Non-invasive Vagus Nerve Stimulation (VNS) Applied in

783   Neurological and Psychiatric Disorders. Front Neurosci *15*, 709436.

784   10.3389/fnins.2021.709436.

785   14.   Heck, C., Helmers, S.L., and DeGiorgio, C.M. (2002). Vagus nerve stimulation

786   therapy, epilepsy, and device parameters: scientific basis and recommendations

787   for use. Neurology *59*, S31-37. 10.1212/wnl.59.6_suppl_4.s31.

788   15.   Krahl, S.E., Clark, K.B., Smith, D.C., and Browning, R.A. (1998). Locus

789   coeruleus lesions suppress the seizure-attenuating effects of vagus nerve

790   stimulation. Epilepsia *39*, 709-714. 10.1111/j.1528-1157.1998.tb01155.x.

791   16.   Krahl, S.E., Senanayake, S.S., and Handforth, A. (2001). Destruction of

792   peripheral C-fibers does not alter subsequent vagus nerve stimulation-induced

793   seizure suppression in rats. Epilepsia *42*, 586-589. 10.1046/j.1528-

794   1157.2001.09700.x.

795   17.   Jiang, Y., Li, L., Liu, B., Zhang, Y., Chen, Q., and Li, C. (2014). Vagus nerve

796   stimulation attenuates cerebral ischemia and reperfusion injury via endogenous

797   cholinergic pathway in rat. PLoS One *9*, e102342.

798   10.1371/journal.pone.0102342.

799   18.   Shao, P., Li, H., Jiang, J., Guan, Y., Chen, X., and Wang, Y. (2023). Role of

800   Vagus Nerve Stimulation in the Treatment of Chronic Pain.

801   Neuroimmunomodulation *30*, 167-183. 10.1159/000531626.

802   19.   Furmaga, H., Shah, A., and Frazer, A. (2011). Serotonergic and noradrenergic

803   pathways are required for the anxiolytic-like and antidepressant-like behavioral

804   effects of repeated vagal nerve stimulation in rats. Biol Psychiatry *70*, 937-945.

805   10.1016/j.biopsych.2011.07.020.

20. Krahl, S.E., Senanayake, S.S., Pekary, A.E., and Sattin, A. (2004). Vagus nerve
stimulation (VNS) is effective in a rat model of antidepressant action. J
Psychiatr Res *38*, 237-240. 10.1016/j.jpsychires.2003.11.005.
21. Noble, L.J., Souza, R.R., and McIntyre, C.K. (2019). Vagus nerve stimulation as
a tool for enhancing extinction in exposure-based therapies.
Psychopharmacology (Berl) *236*, 355-367. 10.1007/s00213-018-4994-5.
22. Andresen, M.C., and Kunze, D.L. (1994). Nucleus tractus solitarius--gateway to
neural circulatory control. Annu Rev Physiol *56*, 93-116.
10.1146/annurev.ph.56.030194.000521.
23. Paton, J.F. (1999). The Sharpey-Schafer prize lecture: nucleus tractus solitarii:
integrating structures. Exp Physiol *84*, 815-833.
24. Appleyard, S.M., Marks, D., Kobayashi, K., Okano, H., Low, M.J., and
Andresen, M.C. (2007). Visceral afferents directly activate catecholamine
neurons in the solitary tract nucleus. J Neurosci *27*, 13292-13302.
10.1523/JNEUROSCI.3502-07.2007.
25. Doyle, M.W., and Andresen, M.C. (2001). Reliability of monosynaptic sensory
transmission in brain stem neurons in vitro. J Neurophysiol *85*, 2213-2223.
10.1152/jn.2001.85.5.2213.
26. Traub, R.J., Sengupta, J.N., and Gebhart, G.F. (1996). Differential c-fos
expression in the nucleus of the solitary tract and spinal cord following noxious
gastric distention in the rat. Neuroscience *74*, 873-884. 10.1016/0306-
4522(96)00173-x.
27. Naritoku, D.K., Terry, W.J., and Helfert, R.H. (1995). Regional induction of fos
immunoreactivity in the brain by anticonvulsant stimulation of the vagus nerve.
Epilepsy Res *22*, 53-62. 10.1016/0920-1211(95)00035-9.
28. Chae, J.H., Nahas, Z., Lomarev, M., Denslow, S., Lorberbaum, J.P., Bohning,
D.E., and George, M.S. (2003). A review of functional neuroimaging studies of
vagus nerve stimulation (VNS). J Psychiatr Res *37*, 443-455. 10.1016/s0022-
3956(03)00074-8.
29. Lomarev, M., Denslow, S., Nahas, Z., Chae, J.H., George, M.S., and Bohning,
D.E. (2002). Vagus nerve stimulation (VNS) synchronized BOLD fMRI
suggests that VNS in depressed adults has frequency/dose dependent effects. J
Psychiatr Res *36*, 219-227. 10.1016/s0022-3956(02)00013-4.

30.   Ran, C., Boettcher, J.C., Kaye, J.A., Gallori, C.E., and Liberles, S.D. (2022). A
brainstem map for visceral sensations. Nature *609*, 320-326. 10.1038/s41586-
022-05139-5.

31.   Adamaszek, M., Manto, M., and Schutter, D.J.L.G. (2022). The Emotional
Cerebellum https://doi.org/10.1007/978-3-030-99550-8.

32.   Supple, W.F., Jr., and Leaton, R.N. (1990). Cerebellar vermis: essential for
classically conditioned bradycardia in the rat. Brain Res *509*, 17-23.
10.1016/0006-8993(90)90303-s.

33.   Supple, W.F., Jr., and Kapp, B.S. (1993). The anterior cerebellar vermis:
essential involvement in classically conditioned bradycardia in the rabbit. J
Neurosci *13*, 3705-3711. 10.1523/JNEUROSCI.13-09-03705.1993.

34.   Chen, C.H., Williams, J.L., and Lutherer, L.O. (1994). Cerebellar lesions alter
autonomic responses to transient isovolaemic changes in arterial pressure in
anaesthetized cats. Clin Auton Res *4*, 263-272. 10.1007/BF01827432.

35.   Armstrong, D.M., Rotler, A., Hersh, L.B., and Pickel, V.M. (1988). Localization
of choline acetyltransferase in perikarya and dendrites within the nuclei of the
solitary tracts. J Neurosci Res *20*, 279-290. 10.1002/jnr.490200302.

36.   Zhao, Q., Yu, C.D., Wang, R., Xu, Q.J., Dai Pra, R., Zhang, L., and Chang, R.B.
(2022). A multidimensional coding architecture of the vagal interoceptive
system. Nature *603*, 878-884. 10.1038/s41586-022-04515-5.

37.   Caravaca, A.S., Gallina, A.L., Tarnawski, L., Tracey, K.J., Pavlov, V.A., Levine,
Y.A., and Olofsson, P.S. (2019). An Effective Method for Acute Vagus Nerve
Stimulation in Experimental Inflammation. Front Neurosci *13*, 877.
10.3389/fnins.2019.00877.

38.   Altidor, L.K., Bruner, M.M., Deslauriers, J.F., Garman, T.S., Ramirez, S., Dirr,
E.W., Olczak, K.P., Maurer, A.P., Lamb, D.G., Otto, K.J., et al. (2021). Acute
vagus nerve stimulation enhances reversal learning in rats. Neurobiol Learn
Mem *184*, 107498. 10.1016/j.nlm.2021.107498.

39.   Kawada, T., Sonobe, T., Nishikawa, T., Hayama, Y., Li, M., Zheng, C., Uemura,
K., Akiyama, T., Pearson, J.T., and Sugimachi, M. (2020). Contribution of
afferent pathway to vagal nerve stimulation-induced myocardial interstitial
acetylcholine release in rats. Am J Physiol Regul Integr Comp Physiol *319*,
R517-R525. 10.1152/ajpregu.00080.2020.

40.   Vuckovic, A., Tosato, M., and Struijk, J.J. (2008). A comparative study of three

techniques for diameter selective fiber activation in the vagal nerve: anodal

block, depolarizing prepulses and slowly rising pulses. J Neural Eng *5*, 275-286.

10.1088/1741-2560/5/3/002.

41.   Brindley, G.S., and Craggs, M.D. (1980). A technique for anodally blocking

large nerve fibres through chronically implanted electrodes. J Neurol Neurosurg

Psychiatry *43*, 1083-1090. 10.1136/jnnp.43.12.1083.

42.   Ahmed, U., Chang, Y.C., Cracchiolo, M., Lopez, M.F., Tomaio, J.N., Datta-

Chaudhuri, T., Zanos, T.P., Rieth, L., Al-Abed, Y., and Zanos, S. (2020). Anodal

block permits directional vagus nerve stimulation. Sci Rep *10*, 9221.

10.1038/s41598-020-66332-y.

43.   Lu, K.H., Cao, J., Phillips, R., Powley, T.L., and Liu, Z. (2020). Acute effects of

vagus nerve stimulation parameters on gastric motility assessed with magnetic

resonance imaging. Neurogastroenterol Motil *32*, e13853. 10.1111/nmo.13853.

44.   Cunningham, J.T., Mifflin, S.W., Gould, G.G., and Frazer, A. (2008). Induction

of c-Fos and DeltaFosB immunoreactivity in rat brain by Vagal nerve

stimulation. Neuropsychopharmacology *33*, 1884-1895.

10.1038/sj.npp.1301570.

45.   Yap, J.Y.Y., Keatch, C., Lambert, E., Woods, W., Stoddart, P.R., and Kameneva,

T. (2020). Critical Review of Transcutaneous Vagus Nerve Stimulation:

Challenges for Translation to Clinical Practice. Front Neurosci *14*, 284.

10.3389/fnins.2020.00284.

46.   Yuan, H., and Silberstein, S.D. (2016). Vagus Nerve and Vagus Nerve

Stimulation, a Comprehensive Review: Part II. Headache *56*, 259-266.

10.1111/head.12650.

47.   Cao, J., Lu, K.H., Powley, T.L., and Liu, Z. (2017). Vagal nerve stimulation

triggers widespread responses and alters large-scale functional connectivity in

the rat brain. PLoS One *12*, e0189518. 10.1371/journal.pone.0189518.

48.   Pnevmatikakis, E.A., Soudry, D., Gao, Y., Machado, T.A., Merel, J., Pfau, D.,

Reardon, T., Mu, Y., Lacefield, C., Yang, W., et al. (2016). Simultaneous

Denoising, Deconvolution, and Demixing of Calcium Imaging Data. Neuron *89*,

285-299. 10.1016/j.neuron.2015.11.037.

49.   Agetsuma, M., Sato, I., Tanaka, Y.R., Carrillo-Reid, L., Kasai, A., Noritake, A.,
905   Arai, Y., Yoshitomo, M., Inagaki, T., Yukawa, H., et al. (2023). Activity-
906   dependent organization of prefrontal hub-networks for associative learning and
907   signal transformation. Nat Commun *14*, 5996. 10.1038/s41467-023-41547-5.

908   50.   Moran, T.H., and Kinzig, K.P. (2004). Gastrointestinal satiety signals II.
909   Cholecystokinin. Am J Physiol Gastrointest Liver Physiol *286*, G183-188.
910   10.1152/ajpgi.00434.2003.

911   51.   Rogers, R.C., and Hermann, G.E. (2008). Mechanisms of action of CCK to
912   activate central vagal afferent terminals. Peptides *29*, 1716-1725.
913   10.1016/j.peptides.2008.06.023.

914   52.   Yamada, D., Koppensteiner, P., Odagiri, S., Eguchi, M., Yamaguchi, S., Yamada,
915   T., Katagiri, H., Wada, K., and Sekiguchi, M. (2017). Common Hepatic Branch
916   of Vagus Nerve-Dependent Expression of Immediate Early Genes in the Mouse
917   Brain by Intraportal L-Arginine: Comparison with Cholecystokinin-8. Front
918   Neurosci *11*, 366. 10.3389/fnins.2017.00366.

919   53.   Cox, J.E., and Randich, A. (1997). CCK-8 activates hepatic vagal C-fiber
920   afferents. Brain Res *776*, 189-194. 10.1016/s0006-8993(97)01036-6.

921   54.   Altschuler, S.M., Bao, X.M., Bieger, D., Hopkins, D.A., and Miselis, R.R.
922   (1989). Viscerotopic representation of the upper alimentary tract in the rat:
923   sensory ganglia and nuclei of the solitary and spinal trigeminal tracts. J Comp
924   Neurol *283*, 248-268. 10.1002/cne.902830207.

925   55.   Bailey, T.W., Appleyard, S.M., Jin, Y.H., and Andresen, M.C. (2008).
926   Organization and properties of GABAergic neurons in solitary tract nucleus
927   (NTS). J Neurophysiol *99*, 1712-1722. 10.1152/jn.00038.2008.

928   56.   Cooper, C.M., Farrand, A.Q., Andresen, M.C., and Beaumont, E. (2021). Vagus
929   nerve stimulation activates nucleus of solitary tract neurons via supramedullary
930   pathways. J Physiol *599*, 5261-5279. 10.1113/JP282064.

931   57.   Blackshaw, L.A., and Grundy, D. (1990). Effects of cholecystokinin (CCK-8) on
932   two classes of gastroduodenal vagal afferent fibre. J Auton Nerv Syst *31*, 191-
933   201. 10.1016/0165-1838(90)90185-l.

934   58.   Lock, M., Alvira, M., Vandenberghe, L.H., Samanta, A., Toelen, J., Debyser, Z.,
935   and Wilson, J.M. (2010). Rapid, simple, and versatile manufacturing of

recombinant adeno-associated viral vectors at scale. Hum Gene Ther *21*, 1259-

1271. 10.1089/hum.2010.055.

59.   Shibata, A.C.E., Ueda, H.H., Eto, K., Onda, M., Sato, A., Ohba, T., Nabekura, J.,

and Murakoshi, H. (2021). Photoactivatable CaMKII induces synaptic plasticity

in single synapses. Nat Commun *12*, 751. 10.1038/s41467-021-21025-6.

60.   Agetsuma, M., Hamm, J.P., Tao, K., Fujisawa, S., and Yuste, R. (2018).

Parvalbumin-Positive Interneurons Regulate Neuronal Ensembles in Visual

Cortex. Cereb Cortex *28*, 1831-1845. 10.1093/cercor/bhx169.

61.   Davis, S.F., Derbenev, A.V., Williams, K.W., Glatzer, N.R., and Smith, B.N.

(2004). Excitatory and inhibitory local circuit input to the rat dorsal motor

nucleus of the vagus originating from the nucleus tractus solitarius. Brain Res

1017, 208-217. 10.1016/j.brainres.2004.05.049.

62.   Evangelidis, G.D., and Psarakis, E.Z. (2008). Parametric image alignment using

enhanced correlation coefficient maximization. IEEE Trans Pattern Anal Mach

Intell *30*, 1858-1865. 10.1109/TPAMI.2008.113.

Figures (from next page)

Figure. 1

Figure 1. Schematic overview of *in vivo* NTS imaging with a double prism

(A) Sagittal view of a mouse brain illustrating how GCaMP was expressed in NTS neurons by

AAV injection.

(B) GCaMP expression in the NTS (coronal view). (left) Schematic illustration of the injection

site modified from the Allen Brain Atlas. (right) Expression of GCaMP (green) in the NTS,

exclusive to the 10N area (marked by ChAT immunohistochemical staining in magenta). Blue,

DAPI.

(C) Two 2-mm right-angled prisms are attached to each other using UV-curable optical adhesive

and bonded to a 3 mm × 3 mm square cover glass. In this illustration, the letter "m" at the

bottom of the double prism appears through the double prism and the objective lens. Similarly,

in the picture at the right bottom corner, the letter "m" in the word "prism" appears through

the double prism.

(D) A vacuum-type prism holder is used to implant the double prism. Black arrows indicate the

directions in which the double prisms are pushed forward. Thick blue arrows indicate

directions in which the double prisms are vacuumed up to be held by the holder. A thin blue

arrow indicates the protrusion to push the double prism during the implantation.

(E) Implantation of the double prism. The double prism is carefully inserted between the

cerebellum and the brainstem approximately 1.6−2.0 mm from the posterior edge of the

cerebellum. The area containing the NTS is observed through the double prism.

(F) A head plate is attached to the mouse skull using dental cement after the microprism

implantation.

(G) Example of *in vivo* brainstem imaging showing neural activity in a live mouse. Left:

Representative two-photon images showing GCaMP fluorescence in a large field of view

covering the wide brainstem. Each number in the image indicates a region of interest (ROI),

each corresponding to a putative single neuron. Median fluorescence over time is presented in

green-fire-blue to show a baseline fluorescence level. Middle: Magnified view of a section

from the left panel. Right: Representative traces showing changes in GCaMP signal over

baseline ($\Delta F/F0$) in selected ROIs. Gray indicates the timing of VNS. Zp4, 4th z-stacked plane

obtained by the volumetric scanning (see also Figure 2A and Figure S3).

D, dorsal; R, right; P, posterior; A, anterior.

Figure. 2

Figure 2. *In vivo* volumetric imaging of NTS activity during repeated VNS

(A) Volumetric live imaging of neural activities in the wide brainstem area, including the NTS. Two-photon imaging was performed through the objective lens (black) and the implanted double prism in the head-fixed mouse under anesthesia. The wide field of view and piezo-based z-scanning enabled volumetric imaging of the GCaMP fluorescence. Colored circles illustrate examples of detected regions of interest (ROIs) at Zp1, Zp2, and Zp3.

(B) *In vivo* NTS imaging was conducted during electrical stimulation of the vagus nerve through a bipolar platinum electrode, which wrapped the cervical vagal trunk in a mouse. The electrode was implanted and fixed with the insulating silicone adhesive. To create anodal block for efferent direction, cathode (black) and anode (red) were located on rostral and caudal site of vagus nerve, respectively.

(C) Schematic of VNS protocol. In this example, a 1.0 mA, 10 Hz, 100 μsec train of electrical stimulation was delivered to the left vagus nerve via an electrode. The term "trial" refers to a

VNS with a pulse width of 100 μsec for 10 sec, regardless of frequency and current intensity.

The term "session" is defined as a series of 10-sec VNS presentations at a fixed frequency

and current intensity, with a 50-sec interval, and this example represents a session at 1.0 mA

and 10 Hz.

(D) An example of *in vivo* volumetric imaging of the NTS during VNS at 2.0 mA and 10 Hz.

AAVdj/Syn.GCaMP7f was used for this experiment. (Left) As examples, Zp1, 2, and 4,

which contain automatically detected ROIs in this mouse, are shown (Zp4 here is the same

data as Figure 1G). Median fluorescence over time for each panel is presented in green-fire-

blue to show a baseline fluorescence level. (Right) Detection of spontaneous and/or VNS-

evoked neural activities ($\Delta F/F0$) from 21 example neurons outside (red) and inside (orange)

the NTS. See also Supplementary Movie1.

(E) Higher-magnification imaging provides clear single-cellular precision. NTS imaging was

performed during VNS at 0.5 mA and 10 Hz. AAV1/Syn.GCaMP6f was used for this

experiment. (Left) As examples, Zp1, and 5, which contain automatically detected ROIs in

this mouse, are shown. To clearly distinguish the ROIs by color, median GCaMP fluorescence

in the NTS is shown in grayscale. (Right) Eleven ROIs, each corresponding to a putative

single neuron, are sorted by the magnitude of their responses to the VNS. See also

Supplementary Movie2.

Zp, z-stacked horizontal plane obtained by the volumetric scanning. Gray indicates the timing of

VNS. D, dorsal; P, posterior; A, anterior; R, right.

Figure. 3

Figure 3. Variable responsiveness of NTS neurons characterized by various VNS conditions

(A) Schematic of VNS protocol. In this example, a 1.0 mA, 10 Hz, 100 μsec train of electrical

stimulation was delivered. Each experiment in this figure includes a session consisting of

three trials of 10-sec VNS presentations, with a 50-sec interval.

(B) Various parameters for evaluating neural responses. The neural response ($\Delta F/F0$) to VNS

(gray shaded area), with the horizontal axis representing the time from the VNS onset.

Average signal (0–10, 0–20, 10–20 sec), peak time, and amplitude, which are calculated for

evaluating neural responses are indicated by green arrows.

(C) Neural responses of an example neuron to VNS at a fixed frequency of 10 Hz, with various

stimulus intensities. Colored lines indicate the neural response to the first, second, and third

VNS. Dashed line indicates the mean of the responses to three VNSs.

(D) Summary of responses of simultaneously recorded neurons (n = 8 neurons in a mouse) to

VNS at a fixed frequency of 10 Hz with various stimulus intensities. The mean $\Delta F/F0$ of each

neuron for 10 sec from the onset of the VNS at various stimulus intensities are summarized.

(E) Summary of neuronal responses to VNS at a fixed frequency of 10 Hz, with various stimulus

intensities. Mean values for three trials, for each neuron (gray line) or for all neurons (black

line), were calculated. Mean $\Delta F/F0$ of 0–10 sec: the mean of $\Delta F/F0$ for 10 sec from the onset

of the VNS. Mean $\Delta F/F0$ of 0–20 sec: the mean of $\Delta F/F0$ for 20 sec from the onset of the

VNS. Mean $\Delta F/F0$ of 10–20 sec: the mean of $\Delta F/F0$ for 10 sec from the offset of the VNS.

Peak time: time from the VNS onset to the $\Delta F/F0$ peak. Amplitude: $\Delta F/F0$ peak amplitude at

each VNS presentation.

(F) An example of $\Delta F/F0$ of neuronal responses to VNS at a fixed stimulus intensity of 1.0 mA,

with various frequencies. Colored lines indicate the neural response to the first, second, and

third VNS. Dashed line indicates mean of the responses to three VNSs.

(G) Summary of simultaneously recorded neurons (n = 8 neurons in a mouse) to VNS at a fixed

stimulus intensity of 1.0 mA, with various frequencies. The mean $\Delta F/F0$ of each neuron for

10 sec from the onset of the VNS at various stimulus frequencies are summarized.

(H) Summary of neuronal responses to VNS at a fixed stimulus intensity of 1.0 mA, with various

frequencies. Mean values for three trials, for each neuron (gray line) or for all neurons (black

line), were calculated. Mean $\Delta F/F0$ of 0–10 sec, mean $\Delta F/F0$ of 0–20 sec, mean $\Delta F/F0$ of 10–

20 sec, peak time, and amplitude were calculated as described in the legend of E.

AAV1/Syn.GCaMP6s was used for this experiment. Gray shaded area indicates the timing

of VNS. Boxes in D and G, 25th–75th percentile; Red bars in D and G, median of all neurons.

Error bars in E and H, standard deviation.

Figure. 4

Figure 4.   Imaging NTS neural activity in response to intravenous CCK administration

(A) Schematic of *in vivo* NTS imaging while Saline or CCK was administered into the tail vein

of an anesthetized mouse (co-administered with red fluorescent dye, SR101). Note that CCK

administration was performed more than 10 min after saline administration.

(B) Averaged fluorescence of GCaMP signals during pre- and post-administration (60 sec

duration) of Saline or CCK. (Top) Baseline-subtracted signal is shown in green over a

background of the baseline image shown in gray, (middle) baseline fluorescence image,

(bottom) GCaMP signal. Dashed circles indicates the NTS area. 4V, fourth ventricle. See also

Supplementary Movies 4 and 5.

(C) Traces of GCaMP fluorescence in simultaneously recorded NTS neurons (same neurons on

the left and right). The timings of the injections (saline, left; CCK, right) are indicated by a

gray dotted line. Traces regarding more than five minutes after the injection are expanded to

visualize the higher frequency of spontaneous activity (i.e., frequent calcium transients)

uniquely observed after CCK administration.

(D) The mean $\Delta F/F0$ for each 60-sec bin along the time course was calculated for saline (left) or

CCK (right) administration. Each data point represents the mean $\Delta F/F$ for each neuron in each

bin.

(E) The mean $\Delta F/F$ for 60 sec immediately after the administration of saline or CCK. Horizontal

red bars, median of all neurons at each condition (saline or CCK). **$p < 0.01$, two-tailed

Wilcoxon signed-rank test ($p = 0.0011$).

(F) The cumulative distribution of intervals between calcium transient events from 60 to 240

seconds after the administration of saline or CCK. Neural activation was evaluated by

detecting the timing of the calcium transients as described in Figure S6 (see also the Methods

for more details). Data from all neurons (6 neurons shown in C) were pooled for each group

for statistical comparison. **$p < 0.01$, Kolmogorov-Smirnov test ($p = 0.0075$).

AAVdj/Syn.GCaMP7f was used for this experiment.

Supplementary Materials

Supplementary Materials (Figures. S1 to S6, and caption for Supplementary Movies S1 to S5) are

explained in a separate document.

1088

A

B

NTS

4V

NTS   NTS
10N   10N
12N   12N

Merge   DAPI

GCaMP   ChAT

300 µm

C

Objective lens

m

Cover glass

Double prism

pris m

2 mm   2 mm

D

Protrusion to push prism in

Implant

E

1.6 ~ 2.0 mm

Edge

NTS

Double prism

F

A

R

G

4V   Zp 4

NTS

500 µm

Zp 4

Intensity F [a.u.]
2000
1494
988
481
-25

200 µm

1.0 (ΔF/F0)

100 sec

A

D ↑ → P
NTS
Zp1
Zp2
Zp3

Zp1 Zp2 Zp3

B

Vagus nerve

C

Example : 1.0 mA and 10 Hz VNS

50 sec interval

10 sec trial

100 msec

Pulse: 100 μsec, 1.0 mA

10 times/sec ⇒ 100 times/trial

D

Zp1 Zp2 Zp4

21
19 17
18 20
15 13 14
16 7 6
11 5 9
10 8
12

Intensity F [a.u.]
2000
1494
988
481
-25

A ↑ → R
500 μm

1
3

1.0 (ΔF/F0)
100 sec

E

Zp1 Zp5

9
10 2 6
8
11 5

1
4 3

A ↑ → R
100 μm

1.0 (ΔF/F0)
100 sec

A Example : 1.0 mA and 10 Hz VNS
Session: 3 trials

B Data explanation

C Various intensities, fixed frequency (10 Hz) : An example neuron (Neuron #5)

D Various intensities, fixed frequency (10 Hz) : Summary of mean ΔF/F0 of 0–10 sec (n = 8 neurons)

E Mean ΔF/F0 of 0–10 sec   Mean ΔF/F0 of 0–20 sec   Mean ΔF/F0 of 10–20 sec   Peak time   Amplitude

F Various frequencies, fixed intensity (1.0 mA) : An example neuron (Neuron #5)

G Various frequencies, fixed intensity (1.0 mA) : Summary of mean ΔF/F0 of 0–10 sec (n = 8 neurons)

H Mean ΔF/F0 of 0–10 sec   Mean ΔF/F0 of 0–20 sec   Mean ΔF/F0 of 10–20 sec   Peak time   Amplitude

A

Saline or CCK with SR101

B

	Saline		CCK	
	Pre	Post	Pre	Post

Merge

Baseline

Intensity F [a.u.]

Activation

4V

A

R

NTS

200 μm

Intensity F [a.u.]

C An example mouse

Neuron #1
Neuron #2
Neuron #3
Neuron #4
Neuron #5
Neuron #6

0.5 (ΔF/F0)

50 sec

0.5 (ΔF/F0)

50 sec

Saline

1.0 (ΔF/F0)

100 sec

CCK

1.0 (ΔF/F0)

100 sec

D

Saline administration

Mean ΔF/F0 of each bin

0.3

0.2

0.1

Neuron #1
Neuron #2
Neuron #3
Neuron #4
Neuron #5
Neuron #6

−60–0   0–60   60–120   120–180   180–240   240–300 (sec)

CCK administration

Mean ΔF/F0 of each bin

0.3

0.2

0.1

−60–0   0–60   60–120   120–180   180–240   240–300 (sec)

E Mean ΔF/F0 of 0–60 sec after administration

ΔF/F0

0.3

0.2

0.1

**

Saline   CCK

F During 60–240 sec after administration

Cumulative Probability

0.5

Saline vs CCK
$p < 0.01$

Saline
CCK

1   10   100

Inter-calcium-transient intervals (sec)