Non-disruptive inducible labeling of ER-membrane contact sites using the Lamin B Receptor

Laura Downie [1], Nuria Ferrandiz [1,2], Megan Jones[1], and Stephen J. Royle [1,✉]

[1]Centre for Mechanochemical Cell Biology, Warwick Medical School, Gibbet Hill Road, Coventry, CV4 7AL, UK; [2]Present address: Centro de Investigación del Cáncer, Salamanca, Spain.

Membrane contact sites (MCSs) are areas of close proximity between organelles that allow the exchange of material, among other roles. The endoplasmic reticulum (ER) has MCSs with a variety of organelles in the cell. MCSs are dynamic, responding to changes in cell state, and are therefore best visualized through inducible labeling methods. However, existing methods typically distort ER-MCSs, by expanding contacts or creating artificial ones. Here we describe a new method for inducible labeling of ER-MCSs using the Lamin B receptor (LBR) and a generic anchor protein on the partner organelle. Termed *LaBeRling*, this versatile, one-to-many approach allows labeling of different types of ER-MCSs (mitochondria, plasma membrane, lysosomes, early endosomes, lipid droplets and Golgi), on-demand, in interphase or mitotic cells. LaBeRling is non-disruptive and does not change ER-MCSs in terms of the contact number, extent or distance measured; as determined by light microscopy or a deep-learning volume electron microscopy approach. We applied this method to study the changes in ER-MCSs during mitosis and to label novel ER-Golgi contact sites at different mitotic stages in live cells.

membrane contact sites | ER-Golgi contacts | inducible labeling
Correspondence: *s.j.royle@warwick.ac.uk*

Introduction

In eukaryotic cells, membrane contact sites (MCSs) are areas of close proximity between two membranes of different identity. MCSs allow the exchange of material between the two membranes without fusion, and they also regulate organelle positioning, dynamics, and number (Scorrano et al., 2019). The ER occupies a large volume of the cell and forms contacts with multiple membranes of different identity (Wu et al., 2018). ER-MCSs therefore play a central role in cellular communication across organelles; they are also dynamic and must adapt in response to changes in cell state. The ER and other membrane compartments remodel upon entry into mitosis (Carlton et al., 2020). For example, the nuclear envelope breaks down and the Golgi fragments (Ungricht and Kutay, 2017; Shorter and Warren, 2002). How ER-MCSs change in response to this remodeling is not well understood. Recent work indicates that changes in ER-MCSs may be coordinated with other mitotic processes through regulation of tethering proteins in mitosis (James et al., 2024). The study of MCSs is hampered by the lack of live cell labeling methods that i) specifically label MCSs, ii) do not interfere with contacts, and iii) allow visualization of contacts during specific cell cycle stages.

ER-MCSs were first observed in electron microscopy (EM) studies in fixed cells (Scorrano et al., 2019). Other visualization methods include those based on measures of organelle membrane proximity from live cell multispectral imaging (Valm et al., 2017) or fixed super-resolution 3D images (Cardoen et al., 2024). Ideally, MCS labeling methods in live cells must distinguish MCS from regions where membranes are in close proximity by chance, as recently reviewed by Nakatsu and Tsukiji (2023). Examples for visualizing ER-MCSs include proximity-dependent methods, where fluorescence is produced when protein tags on each membrane are in close proximity: fluorescence resonance energy transfer (FRET) (Venditti et al., 2019), split fluorescent proteins (FPs) (Yang et al., 2018), and dimerisation-dependent FPs (ddFPs) (Miner et al., 2024). Synthetic constructs based on tethering proteins have also been used to visualize ER-MCSs (Chang et al., 2013). A major limitation of these methods is that labeling is not inducible. To study MCSs during mitosis, labeling that persists and potentially stabilizes interphase contacts is undesirable.

Inducible methods allow temporal control of MCS labeling, and typically use tagged proteins on each apposing membrane that heterodimerize in response to light or a chemical (Sittewelle et al., 2023; Nakatsu and Tsukiji, 2023). However, the initial expression and/or the induced heterodimerization of these proteins usually increase the contact between membranes, through expanding the area of pre-existing MCSs or by creating artificial tethers between the ER and the apposing membrane. Expansion of ER-mitochondria MCSs over time after the inducing labeling has been observed in live cells and by EM (Csordás et al., 2010; Komatsu et al., 2010). Here the ER could be seen wrapping around the mitochondria surface; increasing ER-mitochondria coverage from ~10 % to ~30–90 % after induction (Csordás et al., 2010). Moreover, an optogenetic approach also showed an increase of ~20 % in ER/mitochondria signal overlap in live HeLa cells; and a larger increase in ER-lysosome contacts using this approach in live

COS7 cells (Benedetti et al., 2020). ER-plasma membrane (PM) contacts were also increased after labeling using rapamycin-induced heterodimerization (Várnai et al., 2007). In fact, we previously exploited this property to artificially "glue" the ER to the PM during mitosis in order to free chromosomes trapped by the ER (Ferrandiz et al., 2022). This limits the use of inducible methods for the study of MCSs, as labeling manipulates the ER-MCSs.

Our aim was to develop an inducible labeling system which allows fast and specific labeling of ER-membrane contacts, without disrupting the contacts themselves. Using chemical-induced heterodimerization with a tagged anchor protein at the target membrane, we serendipitously found that Lamin B receptor (LBR), which localizes to the ER and inner nuclear envelope, specifically labeled ER-MCSs upon relocalization. We called this method *LaBeRling*. We found that LaBeRling caused no detectable change in the contact number, extent or distance measured from high resolution 3D EM datasets. It can be used to label multiple different ER-MCS types and we applied this method to study MCSs in mitosis and use it to reveal novel ER-Golgi MCSs in live mitotic cells.

Results

LBR forms clusters after relocalization to the plasma membrane

In principle, induced heterodimerization of proteins tagged with FKBP and FRB domains can be used to label membrane contact sites on-demand. The advantage to this method is that the moment of labeling is controlled by the investigator, a major disadvantage is that the heterodimerization may distort existing contact sites or even induce new, or artificial contacts (Figure 1A). We began by comparing the relocalization of two different proteins: Sec61β and the lamin B receptor, LBR; to the plasma membrane (PM) using induced heterodimerization. Both Sec61β and LBR are found in the ER, with LBR being additionally present on the inner nuclear envelope in interphase. Using Stargazin-dCherry-FRB as a plasma membrane anchor, we found that FKBP-GFP-Sec61β relocalization causes the ER to become glued to the plasma membrane in interphase or during mitosis, as reported previously (Ferrandiz et al., 2022). By contrast, LBR-FKBP-GFP was found in discrete clusters at the plasma membrane following rapamycin addition (Figure 1B).

Similar results were seen when using SH4-FRB-EBFP2, a peripheral membrane protein, in place of the multipass Stargazin plasma membrane anchor (Supplementary Figure S1A,B). Suggesting that the identity of the anchor was unimportant and that the difference in behavior could be attributed solely to the ER-resident protein. Interestingly, there was minimal clustering of the plasma membrane anchor upon relocalization of LBR-FKBP-GFP (Figure S1A and Figure 1C). We also found no evidence for co-clustering of mCherry-Sec61β or LBR-mCherry when LBR-FKBP-GFP was relocalized to the plasma membrane which suggested that the clusters did not represent non-specific aggregates of ER or LBR protein itself (Figure S1A,B). Indeed because the ER stayed intact while LBR-FKBP-GFP clustered in the ER at discrete sites on the plasma membrane, it suggested that this manipulation may be labeling ER-PM contact sites.

Properties of LBR-FKBP-GFP clusters at the plasma membrane following relocalization

We next investigated the properties of the clusters that form after inducing the relocalization of LBR-FKBP-GFP to the plasma membrane. To avoid the possibility that overexpression of LBR-FKBP-GFP contributes to cluster formation, we generated a knock-in cell line where LBR was tagged with FKBP-GFP at its endogenous locus (Supplementary Figure S2). We characterized the properties of the LBR-FKBP-GFP clusters in cells in interphase or at metaphase using 3D segmentation of confocal z-stacks of HCT116 LBR-FKBP-GFP knock-in cells expressing Stargazin-mCherry-FRB treated with rapamycin (200 nM, 20 min) (Figure 1C). The surface area of the clusters (see Methods for definition) was variable but the median cluster area per cell was (0.50 ± 0.07, interphase; 0.55 ± 0.06 μm^2, metaphase), with no significant difference between interphase and metaphase cells (Figure 1D,E). Hundreds of clusters were detected in each cell, but to normalize for differences in cell size and understand the density of LBR-FKBP-GFP clusters at the plasma membrane a cell surface approximation was generated and the density of clusters per unit area was determined (Figure 1F,G). Again the density of clusters was similar in interphase and metaphase cells (0.26 ± 0.12, interphase; 0.21 ± 0.08 μm^{-2}, metaphase). Given the average size of the clusters and their density, the coverage of the plasma membrane was ∼10 %, which is similar to published estimates of PM coverage with ER-PM contact sites (Wu et al., 2017).

To investigate the formation and dynamics of LBR-FKBP-GFP clusters, we imaged mitotic cells during the induced relocalization of LBR-FKBP-GFP to the plasma membrane (Figure 1H). These movies revealed that multiple clusters that were distributed around the cell, formed simultaneously and with similar kinetics (Supplementary Video SV1). The clusters began to form ∼90 s after rapamycin addition, with typically less than 20 clusters in a single confocal slice, and persisted throughout the duration of the movie (up to 5 min). The clusters that formed did so at the expense of fluorescence in the ER which "drained away" with similar kinetics.

Using a spot-tracking procedure we could monitor the behavior of each cluster. Analysis on this time scale revealed that once formed, the total number, size, and brightness of the clusters was essentially constant on

Figure 1. Properties of LBR-FKBP-GFP clusters at the plasma membrane following induced relocalization.
(**A**) Schematic diagram to show heterodimerization of FKBP and FRB domains with rapamycin and how this may be used to label ER-membrane contact sites, but also how this method may distort contact sites. Membrane contact site (MCS) is labeled by a blue box and a green shading is used indicate membranes within close distance of each other. (**B**) Example micrographs of HCT116 cells co-expressing LBR-FKBP-GFP or FKBP-GFP-Sec61β (green) and optionally Stargazin-dCherry-FRB, each treated with rapamycin (200 nм) for 30 min before fixation and DNA staining (blue). Scale bars, 10 µm; Insets, 3× expansion of ROI. (**C**) Micrographs of typical HCT116 LBR-FKBP-GFP (green) knock-in cells expressing Stargazin-mCherry-FRB (red), stained with SiR-DNA (blue), treated with rapamycin (200 nм). Single confocal slices or z-projections for cells in interphase or mitosis are shown. (**D**) Raincloud plot to show size distribution of LBR-FKBP-GFP clusters analyzed in 3D. (**E**) Plot of the median contact area for each cell (dots). Mean and ± sd are indicated by crossbar. (**F**) 3D cell surface approximation generated using the location of all segmented LBR-FKBP-GFP clusters (see Methods). (**G**) Plot of the density of clusters (total clusters divided by cell surface). Dots show cells, mean and ± sd are indicated by crossbar. (**H**) Stills from a movie of LBR-FKBP-GFP cluster formation upon rapamycin addition (200 nм, orange bar). Cells were as described in A. Timescale, mm:ss. Scale bars, 10 µm. (**I**) Plots to show the number, size, and intensity of LBR-FKBP-GFP clusters in a single slice over time. Thin green lines and dark green line, clusters from and average for the cell shown in F; black dotted line, average of 14 different cells.

this time scale (Figure 1I). There were very few splitting or merging of clusters, and any appearance or disappearance of clusters could be attributed to movement into or out of the imaging plane. Overall, the mobility of the clusters was very low ($0.016\,\mu m\,s^{-1}$, median, n = 16 cells).

Together these observations indicate that LBR-FKBP-GFP clusters do not form at random locations at the plasma membrane. Instead the coordinated appearance and stability of fluorescence suggests that relocalized LBR-FKBP-GFP labels pre-existing ER-PM contact sites.

Relocalized LBR-FKBP-GFP labels pre-existing ER-PM contact sites

Are the LBR-FKBP-GFP clusters that form after relocalization, pre-existing ER-PM contact sites? To answer this question we took confocal z-stacks of interphase and mitotic cells before and after LBR-FKBP-GFP relocalization to Stargazin-EBFP2-FRB using rapamycin (200 nM). To identify the ER-PM contact sites, we used a modified MAPPER construct (mScarlet-I3-6DG5-MAPPER) which is an established marker of ER-PM contact sites (Chang et al., 2013). From these images, we could clearly see colocalization between the contact sites marked by MAPPER and the clusters of relocalized LBR-FKBP-GFP in interphase and mitotic cells (Figure 2A). This was also true of the relocalization captured in live HCT116 or HeLa cells (Supplementary Video SV2, SV3). The contact sites where colocalization occurred were present before LBR-FKBP-GFP was relocalized, indicating that labeling is of pre-existing contact sites.

3D segmentation and quantification of the contact sites marked by MAPPER and the clusters of LBR-FKBP-GFP confirmed that the formation of LBR-FKBP-GFP clusters by inducing relocalization was significant. There was no change in the number of MAPPER clusters per cell in mitotic cells, and a small but significant decrease in interphase which was probably attributable to photobleaching (Figure 2B). Importantly, we found no evidence for increases in MAPPER clusters after LBR-FKBP-GFP relocalization which would have indicated the artificial formation of new contact sites. When the colocalization of the two fluorescence channels was measured we found in most cells LBR-FKBP-GFP clusters were contact sites (Figure 2C). The fraction of contact sites that were labeled by LBR-FKBP-GFP was lower, a result which may have been influenced by photobleaching of GFP vs mScarlet-I3.

We found no influence of MAPPER expression on the number of LBR-FKBP-GFP clusters that form upon relocalization to the plasma membrane. For example, there were 213.4 ± 89.4 LBR-FKBP-GFP clusters in MAPPER expressing interphase cells compared with 277.2 ± 112.6 in those not co-expressing MAPPER (p = 0.7, Tukey's post-hoc test). This indicates that there is no interference between the two labeling types.

Finally, this dataset also revealed that there are fewer ER-PM contact sites at metaphase than there are in interphase (Figure 2B). for example, the number of MAPPER clusters in interphase and metaphase cells, before rapamycin treatment was significantly lower (429.3 ± 74.5, interphase; 258.5 ± 45.3, metaphase; p = 0.002). A pattern repeated for post-rapamycin treatment (p = 0.002) and for LBR-FKBP-GFP clusters in MAPPER expressing cells post-rapamycin (p = 0.04). In summary, we could confirm that LBR-FKBP-GFP relocalization to the plasma membrane labels pre-existing ER-PM contact sites and that there were no additional sites created by this relocalization. We also documented a decrease in the number of contacts in mitotic cells compared with non-dividing cells.

Relocalization of LBR-FKBP-GFP to mitochondria highlights ER-mitochondria contact sites

Having established that LBR can be used to inducibly label ER-PM contact sites, we next tested if it could be used to label ER-mitochondria contact sites. HCT116 cells transiently co-expressing MitoTrap (pMito-mCherry-FRB) and either LBR-FKBP-GFP or FKBP-GFP-Sec61β were imaged in interphase or mitosis (Figure 3A). Following relocalization with rapamycin (200 nM), LBR-FKBP-GFP was clustered at discrete sites on each mitochondrion with typically one contact per mitochondrion visible by light microscopy. By contrast, FKBP-GFP-Sec61β fluorescence completely surrounded each mitochondrion suggesting that the relocalization of this protein caused the ER to wrap around the mitochondria (Figure 3A). This pattern was similar in interphase or mitotic cells, and mirrored the observations previously using a plasma membrane anchor.

The clusters of LBR-FKBP-GFP on mitochondria following relocalization were also observed with the endogenously tagged protein in live cells, again suggesting that the cluster formation was not an artifact of overexpression or fixation (Figure 3B). Live-cell imaging of the relocalization revealed that cluster formation was rapid (∼20 s) and came at the expense of fluorescence in the ER (Supplementary Video SV4). The appearance of discrete clusters of LBR-FKBP-GFP upon relocalization to mitochondria suggested that these clusters most likely represent mitochondria-ER contact sites.

Inducible labeling of ER-mitochondria contacts using LBR does not affect ER-mitochondria contacts

Are ER-mitochondria contacts altered by the relocalization of LBR-FKBP-GFP to MitoTrap? As there was no obvious choice of marker to assess this, we used a 3D-EM approach to examine all ER-mitochondria contacts (Figure 4). We first imaged live HCT116 LBR-FKBP-GFP knock-in cells co-expressing MitoTrap and confirmed the relocalization of LBR-FKBP-GFP to mitochondria or not in the case of the control (no rapamycin, Figure 4A). The same cell was then processed for SBF-SEM and relocated for imaging by 3D-EM. We used a

Figure 2. Relocalized LBR-FKBP-GFP labels pre-existing ER-PM contact sites.
(**A**) Example micrographs of HCT116 cells co-expressing LBR-FKBP-GFP (green) and Stargazin-EBFP2-FRB (not shown) and optionally MAPPER (mScarlet-I3-6DG5-MAPPER, red) as indicated. A single slice of a z-stack of the same cell is shown before (pre) or after (post) rapamycin (200 nM, 20 min) addition. Scale bars, 10 µm. (**B**) Comparison of total 3D clusters per cell detected in mScarlet-I3 (MAPPER) or GFP (LBR-FKBP-GFP) channels. Thin lines indicate the pre and post values for each cell. Thick lines and dots indicate the average per experimental repeat. Color indicates experimental repeat. Paired t-tests with Holm-Bonferroni correction for multiple testing: ns, not significant; ***, $p < 0.001$; **, $p < 0.01$. (**C**) Colocalization analysis. For cells co-expressing MAPPER, the number of MAPPER or LBR clusters is shown and the extent that these clusters colocalize with LBR or MAPPER clusters is indicated by the colorscale.

Figure 3. Relocalization of LBR-FKBP-GFP to mitochondria highlights ER-mitochondria contact sites.
(**A**) Example micrographs of HCT116 cells expressing either LBR-FKBP-GFP or FKBP-GFP-Sec61β (green) and MitoTrap (pMito-mCherry-FRB, red). Relocalized samples were treated with rapamycin (200 nM) for 30 min before fixation. Control samples were not treated with rapamycin. A single slice from a z-stack of an interphase or metaphase cell are shown. Scale bars, 10 μm; Insets, 4× expansion of ROI. (**B**) Stills from a live cell imaging experiment with HCT116 LBR-FKBP-GFP (green) knock-in cells co-expressing MitoTrap (red), treated with rapamycin (200 nM) as indicated. Time, mm; Scale bars, 10 μm; Insets, 6.5× expansion of ROI. (**C**) Line profiles measured around the mitochondrial perimeter from cells in A, as represented in the schematic. Plots show the intensity of LBR-FKBP-GFP or FKBP-GFP-Sec61bβ (green) and MitoTrap (magenta) signal measured around an individual mitochondria. Insets show the line profile images.

Figure 4. Inducible labeling of ER-mitochondria contacts using LBR does not affect ER-mitochondria contacts.
(**A**) Example micrographs of HCT116 LBR-FKBP-GFP (green) knock-in cells expressing MitoTrap (magenta) treated with Rapamycin (200 nM) or not (Control). Mitotic stage was confirmed by imaging SiR-DNA (not shown). Following fixation and processing, the same cell was imaged again by serial block face-scanning electron microscopy (SBF-SEM). Finally, SBF-SEM datasets were used to infer the location of ER (green) and mitochondria (magenta), a Z-projection is shown. Scale bar, 10 μm. (**B**) Large-scale machine learning segmentation of ER and mitochondria from SBF-SEM data. (i) Supervized training of nnU-Net using a subvolume of one SBF-SEM dataset in 3dfullres mode. The resulting model is then used to infer the location of ER and mitochondria in the whole volume of multiple datasets. (ii) An image analysis pipeline (see Methods) detects the ER-mitochondria contact areas that are equal or less than the search distance (10–80 nm) from the nearest mitochondrion. (iii) The contacts may be visualized in 3D: orange contact is shown at a highlighted mitochondrion (arrow), ER is represented by green contour lines for clarity. (iv) Contacts detected can be mapped back to the original data for verification. A random selection of contacts from the 50 nm search distance collection are shown. (**C**) Plots of the total ER-mitochondria contact surface area per mitochondrion vs the surface area of the mitochondrion, for each search distance (indicated in grey box, nm). (**D**) Mean ER-mitochondria contact surface area per cell for each search distance. Each cell is represented as a dot, the mean ± sd is shown by a crossbar; n_{cell} = 3 (control), 4 (rapamycin); p values from Student's t-test with Welch's correction.

machine learning approach to infer in 3D the mitochondria and ER in all of the resulting datasets (control, 3; rapamycin, 4) using a manually segmented subvolume from one dataset (Figure 4B). Next, using the inferred maps of mitochondria and ER, we used an automated procedure to detect the regions of ER that were within a defined distance of the mitochondria, and then segment those to measure the size of the ER region that is in contact with the mitochondrion (see Methods). We verified that the procedure identified genuine mitochondrion-ER contacts by sampling a number of contacts and by visualizing them in 3D (Figure 4B). Using the resulting data we could measure the total ER contacts per mitochondrion and plot the total contact area as a function of mitochondrion surface area (Figure 4C). At all defined distances assessed, from 10–80 nm, we found that the contacts were similar between the two experimental groups (Figure 4D). These data indicate that relocalization of LBR-FKBP-GFP can be used to discretely label ER-mitochondria contact sites, without affecting the morphology of those contacts. Since LBR labeling of ER-PM contacts was shown to be similarly non-invasive, we propose that our method is an innocuous way to label contact sites inducibly. We call this method, LaBeRling.

Relocalized LBR-FKBP-GFP can mark several different ER-membrane contact types

As LaBeRling can be used to highlight ER-PM and ER-mitochondria contact sites, we next asked if it could be applied to similarly label other ER-membrane contact sites. To do this we tested three additional anchor proteins: perilipin-3 (PLIN3), early endosome antigen 1 (EEA1), and lysosome-associated membrane glycoprotein 1 (LAMP1); that mark lipid droplets, early endosomes, and lysosomes, respectively. We compared the relocalization of LBR-FKBP-GFP with that of FKBP-GFP-Sec61β to differentiate genuine contact site labeling from non-specific recruitment of ER to the target membrane.

In each case, whether in interphase or mitotic cells, LBR-FKBP-GFP was relocalized to clusters on the surface of the lipid droplet, endosome or lysosome. By contrast, relocalized FKBP-GFP-Sec61β matched the fluorescence of the anchor suggesting that this manipulation had engulfed the target organelle and distorted any pre-existing ER-membrane contact sites (Figure 5). However, the difference between the two relocalized proteins was hard to distinguish due to the small size of each organelle, relative to the size of a MCS.

Together these data indicate that LBR-FKBP-GFP can be used to label ER-membrane contact sites between ER and lipid droplets, endosomes, or lysosomes; and that these contacts are maintained during mitosis. Given that the relocalization of LBR-FKBP-GFP can be used generically to mark five different types of ER-MCS, we suggest that LaBeRling can be used as a multi-purpose label of ER-MCSs.

LBR sterol reductase activity is not required for ER-MCS labeling

We wondered if LBR was unique in being able to be used in this way to label ER-MCSs. To look at this we investigated the relocalization three other proteins: emerin, LAP2β and BAF. Each protein was tagged at the N-terminus with FKBP-GFP and expressed in HCT116 cells alone (control) or with Stargazin-mCherry-FRB and all cells treated with rapamycin (200 nM). We found that each of the relocalized proteins completely coated the surface of the plasma membrane or mitochondria in interphase or mitosis (Supplementary Figure S3). This suggests that something unique to LBR meant that it can be used as an inducible ER-MCS marker.

LBR transmembrane regions are important for sterol reductase activity, essential in the cholesterol biosynthesis pathway (Tsai et al., 2016). To determine if the sterol reductase function is required to label MCSs, we generated two cholesterol synthesis point mutants (N547D or R583Q) associated with disease and tested their ability to inducibly label ER-PM MCSs in mitosis (Clayton et al., 2010). Both mutants were relocalized to discrete puncta that were indistinguishable from those formed after relocalization of the WT LBR-FKBP-GFP to Stargazin-mCherry-FRB (Figure 6A,B). We made a series of truncation constructs to try to isolate the minimal region needed for inducible labeling of ER-MCSs. The two smallest constructs of this series contained either the first two transmembrane (TM) domains or the first TM domain only (GFP-FKBP-LBR(1-288) or FKBP-LBR(1-245)-GFP, respectively). Surprisingly, both of these constructs formed discrete clusters upon relocalisation, similar to the full-length protein (GFP-FKBP-LBR) (Figure 6B,C). These results suggest that the property of MCS targeting is contained in the N-terminal region and first TM domain. However, the truncated constructs all had reduced expression compared to the full-length, so these minimal versions did not supercede the wild-type LBR as an inducible labeler of ER-MCSs. Like the point mutants, the truncated constructs are not predicted to have any sterol reductase activity, so we can rule out cholesterol biosynthesis as the reason why LBR can be used to label ER-MCS.

Using LaBeRling to investigate novel contact sites: ER-Golgi contact sites in mitosis

Contact sites between the ER and trans-Golgi network (TGN) have been described (Venditti et al., 2019). The Golgi undergoes massive remodeling during mitosis (Carlton et al., 2020) but due to the lack of inducible labeling techniques, the status of ER-Golgi MCSs during mitosis has not been studied. Having established LaBeRling we sought to establish whether we could label ER-Golgi MCSs and if so, to test if they persist during mitosis. HTC116 cells transiently co-expressing LBR-FKBP-GFP and FRB-mCherry-Giantin(3131-3259) were imaged follow-

Figure 5. Using LBR-FKBP-GFP relocalization to mark lipid droplet-ER, endosome-ER or lysosome-ER contact sites.
(**A**) Example micrographs of HCT116 cells expressing either LBR-FKBP-GFP or FKBP-GFP-Sec61β (green) together with the indicated mCherry-FRB tagged protein anchor localizing at the target membrane (red), either lipid droplets (FRB-mCherry-PLIN3), early endosomes (FRB-mCherry-EEA1), or lysosomes (Lamp1-mCherry-FRB), and stained with DAPI (blue). Lipid droplet number was increased by incubation with oleic acid (200 μM) for 17 h. Similar treatment reported to make no significant change to LD-ER contacts (Valm et al., 2017). Relocalized samples were treated with rapamycin (200 nM) for 30 min before fixation. Control samples were not treated with rapamycin.' A single slice from a z-stack of an interphase or metaphase cell are shown. Scale bars, 10 μm; insets, 4× (lipid droplet) or 6× (early endosome and lysosome) expansion of ROI. (**B**) Line profiles corresponding to the above cells. Plots show the intensity of LBR-FKBP-GFP or FKBP-GFP-Sec61bβ (green) and anchor protein (magenta) signal measured around the perimeter of the structure. Insets show the line profile images.

ing relocalization with rapamycin (200 nM) (Figure 7A). We compared these to similarly prepared samples co-expressing FKBP-GFP-Sec61β and FRB-mCherry-Giantin(3131-3259). In interphase cells, LBR-FKBP-GFP formed distinct puncta when relocalized to FRB-mCherry-Giantin(3131-3259) Golgi structures. By contrast, FKBP-GFP-Sec61β coated much of the interphase Golgi structures after relocalization (Figure 7A). These results suggest that LBR can be used to selectively label ER-Golgi MCSs.

In mitotic cells, relocalized LBR-FKBP-GFP was observed to be at puncta coinciding with Golgi fragments at prometaphase and at metaphase (Figure 7A). This relocalization pattern could be observed in live mitotic cells (SV6) forming with dynamics similar to that of other ER-MCS labeling events. Moreover, in mitotic HCT116 LBR-FKBP-GFP knock-in cells the same labeling was present confirming that the labeling of ER-Golgi MCSs

in mitosis was not due to overexpression of the LBR protein (Figure 7B). Again, by contrast, large patches of FKBP-GFP-Sec61β signal were observed at relocalised at FRB-mCherry-Giantin(3131-3259) Golgi fragments in prometaphase and metaphase cells (Figure 7A). These results suggest that selective labeling of ER-Golgi MCSs by LBR was possible and was distinguishable from non-specific heterodimerization between ER and Golgi.

The persistence of LaBeRling in mitotic cells suggest that ER-Golgi MCSs are maintained during mitosis. To examine these contacts in further detail, we examined ER-MCSs in mitotic HCT116 cells by SBF-SEM (Figure 7C and Supplementary Figure S4). Small clusters of vesicles were readily observable within 30 nm of ER in metaphase and telophase cells. These clusters match the mitotic Golgi clusters described in EM images of HeLa cells in prometaphase, metaphase and telophase

Figure 6. LBR cholesterol synthesis function is not required for labeling/truncated LBR has similar labeling to the full-length protein/LBR cholesterol synthesis mutants and truncated LBR form relocalised clusters.

(**A**) Schematic of GFP- and FKBP-tagged LBR, sterol reductase point mutants (N547D or R583Q) and truncated proteins (expressing LBR amino acids 1-288 or 1-245). Example micrographs of HCT116 cells expressing FKBP- and GFP-tagged LBR sterol reductase mutants (**B**) or truncated LBR (**C**) (green) alongside Stargazin-mCherry-FRB (red), stained with DAPI (blue). Relocalized samples were treated with rapamycin (200 nᴍ) for 30 min before fixation. Control samples were not treated with rapamycin.' Scale bars, 5 μm; insets, 2.5× expansion of ROI.

Figure 7. Using LBR-FKBP-GFP relocalization to identify novel ER contact sites.

(**A**) Example micrographs of synchronised HCT116 cells expressing either LBR-FKBP-GFP or FKBP-GFP-Sec61β (green) together with pmScarlet-Giantin-C1 or FRB-mCherry-Giantin(3131-3259) (red), and stained with DAPI (blue). Similarly synchronised HCT116 LBR-FKBP-GFP knock-in cells expressing either pmScarlet-Giantin-C1 or FRB-mCherry-Giantin(3131-3259) are shown in (**B**). All samples were treated with rapamycin (200 nM) for 30 min before fixation. Single slices from z-stacks of an interphase, early mitotic or metaphase cell are shown. Scale bars, 10 μm; insets, 4× expansion of ROI. (**C**) Single slices of SBF-SEM dataset of a metaphase HCT116 cell are shown. Depth of each slice within the dataset (nm) is indicated. Example Golgi clusters are shown by yellow arrows on the full slice image. Three sequential slices of these regions (3× expansion) are shown beside. Scale bars, 2 μm and 0.5 μm on zoom region. (**D**) Schematic representation of ER-Golgi contacts in interphase (top) and metaphase (bottom) cells, with example contact sites indicated in the expanded region (blue box).

(Lucocq et al., 1987), and more recently in NRK cells in prophase, metaphase and late anaphase (Jokitalo et al., 2001). Together these data suggest that ER-Golgi MCSs are maintained in mitotic cells and can be labeled by the relocalization of LBR-FKBP-GFP to Golgi membranes using FRB-mCherry-Giantin(3131-3259) (Figure 7D).

Discussion

In this study, we developed a new method, LaBeRling, for inducible labeling of ER-MCSs using the ER protein LBR. Unlike previous inducible labeling approaches, LaBeRling does not alter existing ER-plasma membrane or ER-mitochondria MCSs and it does not induce artificial contacts. Moreover, labeling is fast (<2 min) and persists over many minutes without distortion (up to 30 min measured here) of MCSs, so it is ideal for labeling MCSs at discrete stages of the cell cycle. Finally, we used this method to demonstrate the presence of ER-Golgi MCSs in mitosis, at a time of mitotic Golgi dispersal.

LaBeRling uses heterodimerization of LBR with a generic anchor protein on the target membrane. This anchor protein does not need to localize to MCSs, and can be readily interchanged, so that LBR can be used to label a range of ER-MCSs (between plasma membrane, mitochondria, early endosomes, lysosomes, lipid droplets, and Golgi) in interphase and mitotic cells. In the past, other approaches have used MCS tethering proteins for inducible labeling (Chung et al., 2015; Lees et al., 2017), with the logic that this will increase specificity. This is problematic because both the overexpression of tethering proteins themselves and their subsequent heterodimerization can distort MCSs and induce artificial contacts (Chung et al., 2015; He et al., 2017), as similarly seen using membrane targeting anchors (Csordás et al., 2010; Komatsu et al., 2010; Miner et al., 2024). LaBeRling delivers more universal labeling of contact sites since the initial coverage of the target membrane and the ER is diffuse and homogeneous yet specific to the respective compartments. We envisage that LaBeRling could also be applied to label other ER-membrane contacts, for example, ER-peroxisome and ER-autophagosome MCSs. One drawback of LaBeRling is that resolving MCSs at small organelles by conventional light microscopy is challenging. We found that ER-endosome MCSs labeled by LBR and the non-specific ER-endosome attachment by Sec61β were hard to distinguish. However, it would be possible to adapt LaBeRling for super-resolution imaging to better resolve MCSs at small organelles.

Our serendipitous discovery that LBR can be used to label MCSs is intriguing: what makes LBR so special? We saw that the anchor protein remains homogeneously distributed after LBR relocalization and so the MCS specificity of the labeling seems to be driven by LBR, from the ER side. This observation could mean that ER-MCSs are not truly bipartite and are perhaps governed by the ER. Why LBR behaves in this way, whereas other ER proteins do not, is unresolved. Cholesterol synthesis activity was not essential for labeling because point mutants and truncated LBR proteins that lack sterol reductase activity all labeled EM-plasma membrane MCSs similarly to full-length WT LBR. It is possible that the intermembrane distance at the ER-MCSs is such that the size of LBR-FKBP-GFP is in a "sweet spot" to be immobilized by the heterodimerization procedure, but not so large as to crosslink membranes outside of the MCS. In support of this possibility there is evidence that the altering the spacing in optogenetic heterodimerization of ER-plasma membrane linkages affects labeling efficiency (He et al., 2017). Adjusting linkage distance of ER-plasma membrane contacts has also been indicated to affect protein translocation into the MCS (Várnai et al., 2007; Chang et al., 2013). However, the intermembrane distances of MCSs are reported to be rather variable and we observed that various anchor proteins can be used successfully, which argues against the idea that physical spacing can explain the labeling behavior. Whatever the mechanism, the observation that sterol reductase activity is not required means that the LBR point mutants can be used to ensure the local sterol environment at the MCS is not modified by labeling. This may be important because MCSs are reported to be enriched with cholesterol (Hayashi and Fujimoto, 2010; Fujimoto et al., 2012; Zung et al., 2024), and an ideal labeling method would not perturb the lipids at the endogenous MCS. Conversely, additional protein domains could be added to LBR to deliver new enzymatic activities to ER-MCSs in order to experimentally manipulate these areas and answer new biological questions.

In interphase, ER-TGN MCSs have been described (Wu et al., 2018; Venditti et al., 2019) and we used a generic Golgi anchor protein and LBR-FKBP-GFP to detect ER-Golgi MCSs, which were presumably ER-TGN contacts. We used our inducible method to show that ER-Golgi MCSs are maintained in mitosis, a time when the Golgi has been disassembled (Carlton et al., 2020). To our knowledge this is the first labeling of these MCSs during mitosis but is corroborated by evidence from electron microscopy studies. Briefly, the Golgi is disassembled by severing the Golgi ribbons into stacks and then dispersing the stacks into Golgi "blobs" and "haze" (Misteli and Warren, 1995), which correspond to vesicular clusters and single vesicles, respectively. Each Golgi cluster likely contains a mixture of vesicles from TGN, cis- and medial-Golgi. For example, a subset of the vesicles within the mitotic Golgi clusters had TGN markers in HeLa cells by EM (Lucocq et al., 1987) and an incomplete overlap of different Golgi markers stained within mitotic Golgi clusters was detected by light microscopy in NRK cells (Puri et al., 2004). Therefore ER-TGN contacts are an efficient way for the ER to remain in contact with Golgi clusters. The function of ER-Golgi contacts

maintained during mitosis is unclear. The spindle operates in an "exclusion zone" which is largely membrane free (Nixon et al., 2017). Maintained ER-Golgi contacts could serve simply to exclude clusters from the spindle area and to prevent these membrane fragments from interfering with chromosome segregation. Another possibility is that the ER-Golgi contacts may coordinate reassembly by allowing the clusters to surf on the ER towards the spindle pole and the midbody where the Golgi twins coalesce and begin to reassemble (Gaietta et al., 2006).

We used LaBeRling to probe other ER-MCSs in mitosis. For example, we saw that the total number of ER-plasma membrane MCSs was reduced at metaphase compared with interphase cells. This observation is seemingly at odds with our observation that the density of contacts remained constant. However, our density measurement uses a surface constructed from the contact sites rather than the plasma membrane surface area for normalization. So while the net number of contacts decreases, the spacing between them is maintained as the plasma membrane is drawn up and away from the underlying ER (Erickson and Trinkaus, 1976). How this process is regulated and how MCS are modified during mitosis are interesting questions for the future.

To conclude, the method we describe to label ER-MCS using LBR can be applied to a wide variety of questions at many different types of contact in cells. It can be further developed to tweak the properties of ER-MCSs or to improve resolution of ER-MCSs by light microscopy.

Methods

Molecular biology

The following plasmids were available from Addgene or previous work: pEBFP2-N1 (Addgene #54595); EGFP-BAF (Addgene #101772); Emerin pEGFP-C1 (637) (Addgene #61993); FKBP-alpha(740-977)-GFP (Addgene #100731); FKBP-GFP-Sec61β (Addgene #172442); FRB-mCherry-Giantin (Addgene #186575); GFP-EEA1 (Addgene #42307); Lamp1-mCherry-FRB (Addgene #186576); LAP2 Full I pAcGFP-N1 monomeric GFP (1317) (Addgene #62044); LBR pEGFP-N2 (646) (Addgene #61996); pFKBP-GFP-C1 (Clarke and Royle, 2018); pMaCTag-P05 (Addgene #120016); pMito-mCherry-FRB (Addgene #59352); pMito-dCherry-FRB (Addgene #186573); pmScarlet-Giantin-C1 (Addgene #85048); SH4-FRB-mRFP (Addgene #100741); Stargazin-dCherry-FRB (Addgene #172444); Stargazin-GFP-LOVpep (Addgene #80406); Stargazin-mCherry-FRB (Addgene #172443) (Ferrandiz et al., 2022; Wood et al., 2017; Küey et al., 2022).

To generate the pFRB-mCherry-C1 vector, FRB was amplified from pMito-mCherry-FRB plasmid (using CG CGGCTAGCGGCCACCATGATCCTCTGGCATGAGA TGTGGCATGAAGGC and TCGCaccggtggGCCGGCC TGCTTTGAGATTCGTCGGAACAC) and inserted into pmCherry-C1 vector (using NheI-AgeI). pmCherry-C1 vector was made by substituting mCherry for EGFP in pEGFP-C1 [Clontech] by AgeI-XhoI digestion).

LBR-FKBP-GFP was generated by cutting LBR from LBR-mCherry using BamHI-KpnI sites and ligating into pFKBP-GFP-N1 plasmid, where FKBP was inserted into pEGFP-N1 at BamHI-AgeI sites. Similarly, FKBP-GFP-emerin was made by digestion of Emerin pEGFP-C1 (637) and ligation into FKBP-GFP-C1 plasmid using XhoI and BamHI sites. To clone FKBP-GFP-LAP2β, BglII-SalI sites were introduced at either end and a C-terminal stop codon to LAP2β amplified by PCR from LAP2 Full I pAcGFP-N1 monomeric GFP (1317) plasmid template (primers aagcttAGATCTATGCCGGAGT TCCTAGAGG and tcgagGTCGACCTAgCAGTTGGA TATTTTAGTATCTTGAAG), and ligating into pFKBP-GFP-C1. FKBP-GFP-BAF was generated by digestion of mCherry-BAF plasmid and ligation into pFKBP-GFP-C1 at BglII and Acc65I sites. The mCherry-BAF plasmid used in this cloning was made by PCR amplification of the BAF-encoding region from EGFP-BAF (primers aag cttAGATCTATGACAACCTCCCAAAAGC and tcgagAA GCTTCTACAAGAAGGCATCACACC) and ligation into pmCherry-C1 vector (BglII-HindIII)

LBR point mutations (N547D or R583Q) were introduced by site-directed mutagenesis using LBR-FKBP-GFP plasmid template. Primer with mismatch were designed to introduce the mutation N547D (GTTCGCCAC CCCGATTACTTGGGTGATCTCATC and GATGAGAT CACCCAAGTAATCGGGGTGGCGAAC) or R583Q (C ATGTTGCTTGTCCACCAAGAAGCTCGTGACG and C GTCACGAGCTTCTTGGTGGACAAGCAACATG).

The LBR truncations were generated by PCR amplication of the LBR region from LBR pEGFP-N2 (646) template. To generate the GFP-FKBP-LBR(1-288) plasmid, the primer set aagacaGGATCCATGCCAAGTAGGAAA TTTGCCG and tcgagCAATTGTTAATCAATAAGAGGC GTTCCTTCTACAAC was used. The PCR product was ligated into pEGFP-FKBP-C1 using BamHI-MfeI sites. FKBP-LBR(1-245)-GFP truncation was made by amplifying the region encoding amino acids 1-245 (using aag acaGGTACCGTGAACCGTCAGATCCGCTAG and tcg agACCGGTagAGGAGGGAAATTCAGAAGACTGGGA TCTTTC). The PCR product was ligated in substitution of the AP2A1 encoding region in FKBP-alpha(740-977)-GFP (using KpnI-AgeI).

The plasma membrane anchor Stargazin-EBFP2-FRB was made by PCR of Stargazin encoding region from Stargazin-GFP-LOVpep (primers gcggctagcATGGGG CTGTTTGATCGAGGTGTTCA and TTTACTCATGG ATCCttTACGGGCGTGGTCCGG) and then ligating into pEBFP2-N1. Stargazin-EBFP2 was amplified from the resulting plasmid (primers aagcttGCTAGCcATGG GGCTGTTTGATCGAGG and tcgagGGTACCccCTTG TACAGCTCGTCCATGC), excluding the stop codon, and ligated N-terminal to FRB in a plasmid encoding Stargazin-dCherry-FRB. SH4-FRB-EBFP2 was made

by substituting mRFP of SH4-FRB-mRFP with EBFP2 from pEBFP2-N1 (using AgeI-NotI).

Endosomal anchor FRB-mCherry-EEA1 was made by cutting full-length EEA1 from GFP-EEA1 using XhoI-BamHI and ligating into pFRB-mCherry-C1. Lipid droplet anchor FRB-mCherry-PLIN3 was made from a mGFP-PLIN3 full-length plasmid that was generated by DNA synthesis and Gibson Assembly, cutting FRB-mCherry from FRB-mCherry plasmid (NheI-BsrGI) and substituting for the mGFP of mGFP-PLIN3.

The template plasmid for the C-terminal PCR tagging CRISPR method (Fueller et al., 2020) encoding FKBP-GFP tag (pMaCTag-P05-FKBP-GFP) was generated by amplifying the region encoding FKBP-GFP from pFKBP-GFP-N1 and introducing BamHI-SpeI restriction sites (primers aagcttGGATCCCCGCCACCAATGGGAGTGCAGGTGG and tcgagACTAGTTTACTTGTACAGCTCGTCCATGCCGAGAGT), and then ligating in place of the GFP tag in available pMaCTag-P05 to give pMaCTag-P05 FKBP-GFP. To generate the PCR product for editing, LBR-specific tagging oligos were designed using the online design tool (M1, CGTGACGAGTACCACTGTAAGAAGAAATACGGCGTGGCTTGGGAAAAGTACTGTCAGCGTGTGCCCTACCGTATATTTCCATACATCTACTCAGGTGGAGGAGGTAGTG and M2, TTTGCAAATGGCAGCTGGAATTGCAGGAGTATTTTGTAGAAAAGCCAGAAGAGCAAAAAAAGAGCATTAGTAGATGTATGATCTACACTTAGTAGAAATTAGCTAGCTGCATCGGTACC). Primers for genotyping PCR to confirm LBR-FKBP-GFP knock-in were AGAATTGGGGGAAAGCAGG and CATCCTTACTTGTATTTTTCCTATGTTAACTG or AAGACAATAGCAGGCATGCT and CAGTGGCACCATAGGCATAA.

The mScarlet-I3-6DG5-MAPPER plasmid was designed based on the available GFP-MAPPER sequence and generated by DNA synthesis (Chang et al., 2013). Briefly, the GFP-encoding sequence of GFP-MAPPER was replaced with mScarlet-I3 and the unwanted FRB sequence was substituted with a codon-optimized sequence of Neoleukin-2/15 (PDB code, 6DG5), which is structurally similar and close in mass to FRB. Linker sequence lengths were maintained similar to that in the GFP-MAPPER sequence.

Cell biology

HCT116 (CCL-247; ATCC) cells and lines derived from HCT116 were maintained in DMEM supplemented with 10 % FBS and 100 U mL^{-1} penicillin/streptomycin. All cell lines were kept in a humidified incubator at 37 °C and 5 % CO_2. Cells were routinely tested for mycoplasma contamination by a PCR-based method.

HCT116 LBR-FKBP-GFP CRISPR knock-in cells were generated using a C-terminal PCR tagging CRISPR method (Fueller et al., 2020). HCT116 cells were transfected with the M1-M2 PCR cassette and a plasmid encoding Cas12a (pVE13300). Cells were selected and maintained in media supplemented with puromycin dihydrochloride (Gibco) at 1.84 µM. Populations of cells

positive for GFP signal were selected by FACS and the positive pools of cells were characterized by genotyping PCR, western blot and microscopy.

For transient transfection of HCT116 and derived cells, Fugene-HD (Promega) was used according to the manufacturer's instructions.

Heterodimerization of FKBP and FRB tags was induced through addition of rapamycin to media at a final concentration of 200 nM. For fixed cell experiments, 200 nM rapamycin (J62473, Alfa Aesar) was prepared in complete media 2 mL per well of a 6 well plate; growth media was removed, rapamycin-containing media added, and then the plate was returned to the incubator until fixation. To apply rapamycin to live cells on the microscope, rapamycin solution (400 nM) in imaging media (Leibovitz's L-15 Medium, no phenol red, supplemented with 10 % FBS) was diluted (1:2) to final concentration 200 nM using media in the dish. To visualize DNA in live cells, dishes were incubated for 15–30 min with 0.1 µM SiR-DNA (Spirochrome) prepared in complete media.

Cells were synchronised by treatment with thymidine (2 mM, 16 h), followed by washout, 7–8 h incubation and then RO-3306 treatment(9 µM, 16 h). Cells were released from synchronisation and incubated for 15 min or 30 min before applying rapamycin solution (final concentration 200 nM) and incubating a further 30 min before fixation.

To increase the number of lipid droplets, the media on cells was supplemented with oleic acid (O3008, Sigma) at 200 µM) for around 17 h before fixation.

Fluorescence methods

For fixed-cell imaging, cells were seeded onto glass cover slips (16 mm diameter and thickness number 1.5, 0.16–0.19 mm). Cells were fixed using 3 % PFA/4 % sucrose in PBS for 15 min. After fixation, cells were washed with PBS and then incubated in permeabilization buffer (0.5 % v/v Triton X-100 in 1xPBS) for 10 min. Cells were washed twice with PBS, before 45–60 min blocking (3 % BSA, 5 % goat serum in 1xPBS). Antibody dilutions were prepared in blocking solution. After blocking, cells were incubated for 1 h with primary antibody, PBS washed (3 washes, 5 min each), 1 h secondary antibody incubation, 5 min each PBS wash (3 washes), mounting with Vectashield containing DAPI (Vector Labs Inc.) and then sealed.

Western blotting

For Western blotting, cells were harvested, and lysates were prepared by sonication of cells in UTB buffer (8 M urea, 50 mM Tris, and 150 mM 2-mercaptoethanol). Lysates were incubated on ice for 30 min, clarified in a benchtop centrifuge (20 800 g) for 15 min at 4 °C, boiled in Laemmli buffer for 10 min, and resolved on a precast 4–15 % polyacrylamide gel (Bio-Rad). Proteins were transferred to nitrocellulose using a Trans-Blot Turbo Transfer System (Bio-Rad). The following antibodies were used: rat monoclonal anti-GFP (3h9, Chromotek)

at 1:1000 in 3 % BSA TBST; mouse anti-LBR antibody (polyclonal) (SAB1400151, Sigma-Aldrich) at 1:500 in 2 % milk TBST; goat anti-rat IgG-Peroxidase antibody (A9037, Sigma) at 1:5000 in 5 % milk TBST; sheep anti-mouse IgG-HRP (NXA931, Cytiva) at 1:5000 in 5 % milk TBST; loading control HRP-conjugated mouse anti-β-actin (C4) (sc-47778, Santa Cruz Biotechnology) at 1:20000 in 2 % milk TBST.

Microscopy

As described previously, images were captured using a Nikon CSU-W1 spinning disc confocal system with SoRa upgrade (Yokogawa) with either a Nikon, 100×, 1.49 NA, oil, CFI SR HP Apo TIRF or a 63×, 1.40 NA, oil, CFI Plan Apo objective (Nikon) with optional 2.8× intermediate magnification and 95B Prime camera (Photometrics) (Ferrandiz et al., 2022). The system has a CSU-W1 (Yokogawa) spinning disk unit with 50 μm and SoRa disks (SoRa disk used), Nikon Perfect Focus autofocus, Okolab microscope incubator, Nikon motorized xy stage and Nikon 200 μm z-piezo. Excitation was via 405 nm, 488 nm, 561 nm and 638 nm lasers with 405/488/561/640 nm dichroic and Blue, 446/60; Green, 525/50; Red, 600/52; FRed, 708/75 emission filters. Acquisition and image capture was via NiS Elements (Nikon). All microscopy data was stored in an OMERO database in native file formats.

Serial block face-scanning electron microscopy

Preparation of samples for serial block face-scanning electron microscopy (SBF-SEM) was performed as described previously (Ferrandiz and Royle, 2023; Ferrandiz et al., 2022). Briefly, HCT116 LBR-FKBP-GFP knock-in cells expressing MitoTrap were plated onto gridded dishes and prior to imaging, incubated for around 30 min with 0.5 μM SiR-DNA (Spirochrome) to visualize DNA. Using light microscopy, live cells were imaged to confirm the induced labeling following rapamycin (200 nM) treatment. Control cells not treated with rapamycin were imaged in parallel and the coordinate position of the cell of interest recorded for correlation by SBF-SEM. Cells were washed twice with PB (phosphate buffer) before fixing (2.5 % glutaraldehyde, 2 % paraformaldehyde, 0.1 % tannic acid (low molecular weight) in 0.1 M phosphate buffer, pH 7.4) for 1 h at room temperature. Samples were washed three times with PB and then post-fixed in 2 % reduced osmium (equal volume of 4 % OsO$_4$ prepared in water and 3 % potassium ferrocyanide in 0.1 M PB solution) for 1 h at room temperature, followed by a further three washes with PB. Cells were then incubated for 5 min at room temperature in 1 % (w/v) thiocarbohydrazide solution, followed by three PB washes. A second osmium staining step was then included, incubating cells in a 2 % OsO$_4$ solution prepared in water for 30 min at room temperature, followed by three washes with PB. Cells were then incubated in 1 % uranyl acetate solution at 4 ℃ overnight. This was followed by a further three washes

with PB. Walton's lead aspartate was prepared adding 66 mg lead nitrate (TAAB) to 9 mL 0.03 M aspartic acid solution at pH 4.5, and then adjusting to final volume of 10 mL with 0.03 M aspartic acid solution and to pH 5.5 (pH adjustments with KOH). Cells were incubated in Walton's lead aspartate for 30 min at room temperature and then washed three times in PB. Samples were dehydrated in an ethanol dilution series (30 %, 50 %, 70 %, 90 %, and 100 % ethanol, 5 min incubation in each solution) on ice, then incubated for a further 10 min in 100 % ethanol at room temperature. Finally, samples were embedded in an agar resin (AGAR 100 R1140, Agar Scientific). SBF-SEM imaging was carried out by the Biomedical Electron Microscopy Unit at University of Liverpool, UK.

Data analysis

Analysis of confocal z-stacks of LBR-FKBP-GFP clusters at the plasma membrane was by 3D Spot Finder in 3D Image Suite plugin in Fiji. Briefly, outputs were fed into R where the size and number of clusters was stored. Contact area was defined as half of the surface area of a cluster. The location of clusters was used to find the surface of the cell using alphashape3d. The total number of clusters divided by the surface area of the alpha shape was used to determine the density of clusters per cell. For analysis of cluster formation in movies of LBR-FKBP-GFP relocalization to the plasma membrane, a weka segmentation method was used in Track-Mate/Fiji to define the clusters that formed and track individual clusters over time. The outputs of these Track-Mate XML files was analyzed using TrackMateR (Sittewelle and Royle, 2024). Tracks shorter than 4 frames or those that terminated before 100 s were removed from analysis and the remainder analyzed for shape, intensity, and number over time.

Analysis of ER-PM contacts labelled by LBR-FKBP-GFP/MAPPER was done using a semi-automatated procedure to segment each channel in 3D using 3D spot finder in 3D Image Suite in Fiji. Output were processed in R, where the total numbers of clusters per cell was analyzed per cell. Comparisons between pre and post rapamycin was done using paired t-tests on the individual cell data, with Holm-Bonferroni correction for multiple testing. Comparison between conditions was done using the experimental means using ANOVA with Tukey's post hoc test.

For ER-mitochondria contact analysis, each SBF-SEM dataset was aligned using SIFT and cropped. A subvolume of one dataset (48 slices) was manually segmented for ER and mitochondria using IMOD. The segmentation and corresponding raw data were used to train nnU-Net v2 (Isensee et al., 2021) running on a GPU workstation (Intel Core i9-7900X, 128 GB, with TITAN Xp GPU). The resulting model was used to infer the ER and mitochondria in all datasets. Visualization of the inference maps or the manual segmented IMOD models was done using ChimeraX. A series of Fiji/ImageJ scripts were used

to process the output. Briefly, an exact euclidean distance transform (EDT) was generated using the mitochondria channel to give a 3D volume of distances from each voxel to the nearest mitochondrion. Then the overlap between ER channel and EDT at distances of 10–80 nm was calculated in 10 nm increments, to give the ER regions (contacts) within the appropriate distance from the mitochondrion. This result was segmented in 3D to classify all of the contacts. In addition, the mitochondria channel was also segmented in 3D. These outputs were processed in R to match each contact with its corresponding mitochondrion, which allowed the comparison of total contact surface area per mitochondrion with the mitochondrion surface area.

Data and software availability

ACKNOWLEDGEMENTS

We thank Maelle Lorvellec and Laura Cooper from the Computing and Advanced Microscopy Unit (CAMDU) for their help and support. Alex Moore provided technical assistance to make the LBR-FKBP-GFP knock-in cells and to manually segment SBF-SEM data for nnU-Net training. Alison Beckett at the Liverpool Biomedical EM Unit carried out SBF-SEM imaging. We are grateful to members of the Royle lab for feedback on the project and manuscript. This work was supported by a Programme Award from Cancer Research UK (C25425/A27718). LD and MJ were supported by studentships from BBSRC Midlands Integrative Biosciences Training Partnership (MIBTP2, BB/M01116X/1 and MIBTP2020, BB/T00746X/1).

AUTHOR CONTRIBUTIONS

L. Downie carried out all experimental work with the exception of SBF-SEM experiments (performed by N. Ferrandiz), CRISPR knock-in cell line generation (N. Ferrandiz), LBR point mutation and truncation work (M. Jones). Data analysis was done by L. Downie and S.J. Royle.

References

Benedetti, L., Marvin, J. S., Falahati, H., Guillén-Samander, A., Looger, L. L., and De Camilli, P. (2020). Optimized Vivid-derived Magnets photodimerizers for subcellular optogenetics in mammalian cells. *eLife*, 9:e63230. doi: 10.7554/eLife.63230.

Cardoen, B., Vandevoorde, K. R., Gao, G., Ortiz-Silva, M., Alan, P., Liu, W., Tiliakou, E., Vogl, A. W., Hamarneh, G., and Nabi, I. R. (2024). Membrane contact site detection (MCS-DETECT) reveals dual control of rough mitochondria-ER contacts. *The Journal of Cell Biology*, 223(1):e202206109. doi: 10.1083/jcb.202206109.

Carlton, J. G., Jones, H., and Eggert, U. S. (2020). Membrane and organelle dynamics during cell division. *Nature Reviews. Molecular Cell Biology*, 21(3):151–166. doi: 10.1038/s41580-019-0208-1.

Chang, C.-L., Hsieh, T.-S., Yang, T. T., Rothberg, K. G., Azizoglu, D. B., Volk, E., Liao, J.-C., and Liou, J. (2013). Feedback regulation of receptor-induced Ca2+ signaling mediated by E-Syt1 and Nir2 at endoplasmic reticulum-plasma membrane junctions. *Cell Reports*, 5(3):813–825. doi: 10.1016/j.celrep.2013.09.038.

Chung, J., Torta, F., Masai, K., Lucast, L., Czapla, H., Tanner, L. B., Narayanaswamy, P., Wenk, M. R., Nakatsu, F., and De Camilli, P. (2015). PI4P/phosphatidylserine countertransport at ORP5- and ORP8-mediated ER–plasma membrane contacts. *Science (New York, N.Y.)*, 349(6246):428–432. doi: 10.1126/science.aab1370.

Clarke, N. I. and Royle, S. J. (2018). FerriTag is a new genetically-encoded inducible tag for correlative light-electron microscopy. *Nature Communications*, 9(1):2604. doi: 10.1038/s41467-018-04993-0.

Clayton, P., Fischer, B., Mann, A., Mansour, S., Rossier, E., Veen, M., Lang, C., Baasanjav, S., Kieslich, M., Brossuleit, K., Gravemann, S., Schnipper, N., Karbasyian, M., Demuth, I., Zwerger, M., Vaya, A., Utermann, G., Mundlos, S., Stricker, S., Sperling, K., and Hoffmann, K. (2010). Mutations causing Greenberg dysplasia but not Pelger anomaly uncouple enzymatic from structural functions of a nuclear membrane protein. *Nucleus (Austin, Tex.)*, 1(4):354–366. doi: 10.4161/nucl.1.4.12435.

Csordás, G., Várnai, P., Golenár, T., Roy, S., Purkins, G., Schneider, T. G., Balla, T., and Hajnóczky, G. (2010). Imaging interorganelle contacts and local calcium dynamics at the ER-mitochondrial interface. *Molecular Cell*, 39(1):121–132. doi: 10.1016/j.molcel.2010.06.029.

Erickson, C. A. and Trinkaus, J. P. (1976). Microvilli and blebs as sources of reserve surface membrane during cell spreading. *Experimental Cell Research*, 99(2):375–384. doi: 10.1016/0014-4827(76)90595-4.

Ferrandiz, N. and Royle, S. J. (2023). 3D Ultrastructural Visualization of Mitosis Fidelity in Human Cells Using Serial Block Face Scanning Electron Microscopy (SBF-SEM). *Bio-Protocol*, 13(13):e4708. doi: 10.21769/BioProtoc.4708.

Ferrandiz, N., Downie, L., Starling, G. P., and Royle, S. J. (2022). Endomembranes promote chromosome missegregation by ensheathing misaligned chromosomes. *The Journal of Cell Biology*, 221(6):e202203021. doi: 10.1083/jcb.202203021.

Fueller, J., Herbst, K., Meurer, M., Gubicza, K., Kurtulmus, B., Knopf, J. D., Kirrmaier, D., Buchmuller, B. C., Pereira, G., Lemberg, M. K., and Knop, M. (2020). CRISPR-Cas12a-assisted PCR tagging of mammalian genes. *The Journal of Cell Biology*, 219(6): e201910210. doi: 10.1083/jcb.201910210.

Fujimoto, M., Hayashi, T., and Su, T.-P. (2012). The role of cholesterol in the association of endoplasmic reticulum membranes with mitochondria. *Biochemical and Biophysical Research Communications*, 417(1):635–639. doi: 10.1016/j.bbrc.2011.12.022.

Gaietta, G. M., Giepmans, B. N. G., Deerinck, T. J., Smith, W. B., Ngan, L., Llopis, J., Adams, S. R., Tsien, R. Y., and Ellisman, M. H. (2006). Golgi twins in late mitosis revealed by genetically encoded tags for live cell imaging and correlated electron microscopy. *Proceedings of the National Academy of Sciences of the United States of America*, 103(47): 17777–17782. doi: 10.1073/pnas.0608509103.

Hayashi, T. and Fujimoto, M. (2010). Detergent-resistant microdomains determine the localization of sigma-1 receptors to the endoplasmic reticulum-mitochondria junction. *Molecular Pharmacology*, 77(4):517–528. doi: 10.1124/mol.109.062539.

He, L., Jing, J., Zhu, L., Tan, P., Ma, G., Zhang, Q., Nguyen, N. T., Wang, J., Zhou, Y., and Huang, Y. (2017). Optical control of membrane tethering and interorganellar communication at nanoscales †Electronic supplementary information (ESI) available: Experimental procedures, supplemental sequences, figures referred in the text, supplemental movies. See DOI: 10.1039/c7sc01115f Click here for additional data file. Click here for additional data file. Click here for additional data file. Click here for additional data file. *Chemical Science*, 8(8):5275–5281. doi: 10.1039/c7sc01115f.

Isensee, F., Jaeger, P. F., Kohl, S. A. A., Petersen, J., and Maier-Hein, K. H. (2021). nnU-Net: a self-configuring method for deep learning-based biomedical image segmentation. *Nature Methods*, 18(2):203–211. doi: 10.1038/s41592-020-01008-z.

James, C., Möller, U., Spillner, C., König, S., Dybkov, O., Urlaub, H., Lenz, C., and Kehlenbach, R. H. (2024). Phosphorylation of ELYS promotes its interaction with VAPB at decondensing chromosomes during mitosis. *EMBO reports*. doi: 10.1038/s44319-024-00125-6.

Jokitalo, E., Cabrera-Poch, N., Warren, G., and Shima, D. T. (2001). Golgi clusters and vesicles mediate mitotic inheritance independently of the endoplasmic reticulum. *The Journal of Cell Biology*, 154(2):317–330. doi: 10.1083/jcb.200104073.

Komatsu, T., Kukelyansky, I., McCaffery, J. M., Ueno, T., Varela, L. C., and Inoue, T. (2010). Organelle-Specific, Rapid Induction of Molecular Activities and Membrane Tethering. *Nature methods*, 7(3):206–208. doi: 10.1038/nmeth.1428.

Küey, C., Sittewelle, M., Larocque, G., Hernández-González, M., and Royle, S. J. (2022). Recruitment of clathrin to intracellular membranes is sufficient for vesicle formation. *eLife*, 11:e78929. doi: 10.7554/eLife.78929.

Lees, J. A., Messa, M., Sun, E. W., Wheeler, H., Torta, F., Wenk, M. R., De Camilli, P., and Reinisch, K. M. (2017). Lipid transport by TMEM24 at ER-plasma membrane contacts regulates pulsatile insulin secretion. *Science (New York, N.Y.)*, 355(6326):eaah6171. doi: 10.1126/science.aah6171.

Lucocq, J. M., Pryde, J. G., Berger, E. G., and Warren, G. (1987). A mitotic form of the Golgi apparatus in HeLa cells. *The Journal of cell biology*, 104(4):865–874. doi: 10.1083/jcb.104.4.865.

Miner, G. E., Smith, S. Y., Showalter, W. K., So, C. M., Ragusa, J. V., Powers, A. E., Zanellati, M. C., Hsu, C. H., Marchan, M. F., and Cohen, S. (2024). Contact-FP: A Dimerization-Dependent Fluorescent Protein Toolkit for Visualizing Membrane Contact Site Dynamics. *Contact (Thousand Oaks (Ventura County, Calif.))*, 7:25152564241228911. doi: 10.1177/25152564241228911.

Misteli, T. and Warren, G. (1995). Mitotic disassembly of the Golgi apparatus in vivo. *Journal of Cell Science*, 108 (Pt 7):2715–2727. doi: 10.1242/jcs.108.7.2715.

Nakatsu, F. and Tsukiji, S. (2023). Chemo- and opto-genetic tools for dissecting the role of membrane contact sites in living cells: Recent advances and limitations. *Current Opinion in Chemical Biology*, 73:102262. doi: 10.1016/j.cbpa.2022.102262.

Nixon, F. M., Honnor, T. R., Clarke, N. I., Starling, G. P., Beckett, A. J., Johansen, A. M., Brettschneider, J. A., Prior, I. A., and Royle, S. J. (2017). Microtubule organization within mitotic spindles revealed by serial block face scanning electron microscopy and image analysis. *J Cell Sci*, 130(10):1845–1855. doi: 10.1242/jcs.203877.

Puri, S., Telfer, H., Velliste, M., Murphy, R. F., and Linstedt, A. D. (2004). Dispersal of Golgi matrix proteins during mitotic Golgi disassembly. *Journal of Cell Science*, 117(Pt 3): 451–456. doi: 10.1242/jcs.00863.

Scorrano, L., De Matteis, M. A., Emr, S., Giordano, F., Hajnóczky, G., Kornmann, B., Lackner, L. L., Levine, T. P., Pellegrini, L., Reinisch, K., Rizzuto, R., Simmen, T., Stenmark, H., Ungermann, C., and Schuldiner, M. (2019). Coming together to define membrane contact sites. *Nature Communications*, 10(1):1287. doi: 10.1038/s41467-019-09253-3.

Shorter, J. and Warren, G. (2002). Golgi architecture and inheritance. *Annual Review of Cell and Developmental Biology*, 18:379–420. doi: 10.1146/annurev.cellbio.18.030602.133733.

Sittewelle, M. and Royle, S. J. (2024). Passive diffusion accounts for the majority of intracellular nanovesicle transport. *Life Science Alliance*, 7(1):e202302406. doi: 10.26508/lsa.202302406.

Sittewelle, M., Ferrandiz, N., Fesenko, M., and Royle, S. J. (2023). Genetically encoded imaging tools for investigating cell dynamics at a glance. *Journal of Cell Science*, 136(7): jcs260783. doi: 10.1242/jcs.260783.

Tsai, P.-L., Zhao, C., Turner, E., and Schlieker, C. (2016). The Lamin B receptor is essential for cholesterol synthesis and perturbed by disease-causing mutations. *eLife*, 5:e16011. doi: 10.7554/eLife.16011.

Ungricht, R. and Kutay, U. (2017). Mechanisms and functions of nuclear envelope remodelling. *Nature Reviews. Molecular Cell Biology*, 18(4):229–245. doi: 10.1038/nrm.

2016.153.

Valm, A. M., Cohen, S., Legant, W. R., Melunis, J., Hershberg, U., Wait, E., Cohen, A. R., Davidson, M. W., Betzig, E., and Lippincott-Schwartz, J. (2017). Applying systems-level spectral imaging and analysis to reveal the organelle interactome. *Nature*, 546(7656): 162–167. doi: $10.1038/\text{nature}22369$.

Venditti, R., Rega, L. R., Masone, M. C., Santoro, M., Polishchuk, E., Sarnataro, D., Paladino, S., D'Auria, S., Varriale, A., Olkkonen, V. M., Di Tullio, G., Polishchuk, R., and De Matteis, M. A. (2019). Molecular determinants of ER-Golgi contacts identified through a new FRET-FLIM system. *The Journal of Cell Biology*, 218(3):1055–1065. doi: $10.1083/\text{jcb}.201812020$.

Várnai, P., Tóth, B., Tóth, D. J., Hunyady, L., and Balla, T. (2007). Visualization and manipulation of plasma membrane-endoplasmic reticulum contact sites indicates the presence of additional molecular components within the STIM1-Orai1 Complex. *The Journal of Biological Chemistry*, 282(40):29678–29690. doi: $10.1074/\text{jbc}.M704339200$.

Wood, L. A., Larocque, G., Clarke, N. I., Sarkar, S., and Royle, S. J. (2017). New tools for "hot-wiring" clathrin-mediated endocytosis with temporal and spatial precision. *The Journal of Cell Biology*, 216(12):4351–4365. doi: $10.1083/\text{jcb}.201702188$.

Wu, H., Carvalho, P., and Voeltz, G. K. (2018). Here, there, and everywhere: The importance of ER membrane contact sites. *Science (New York, N.Y.)*, 361(6401):eaan5835. doi: $10.1126/\text{science}.aan5835$.

Wu, Y., Whiteus, C., Xu, C. S., Hayworth, K. J., Weinberg, R. J., Hess, H. F., and De Camilli, P. (2017). Contacts between the endoplasmic reticulum and other membranes in neurons. *Proceedings of the National Academy of Sciences of the United States of America*, 114 (24):E4859–E4867. doi: $10.1073/\text{pnas}.1701078114$.

Yang, Z., Zhao, X., Xu, J., Shang, W., and Tong, C. (2018). A novel fluorescent reporter detects plastic remodeling of mitochondria-ER contact sites. *Journal of Cell Science*, 131(1):jcs208686. doi: $10.1242/\text{jcs}.208686$.

Zung, N., Aravindan, N., Boshnakovska, A., Valenti, R., Preminger, N., Jonas, F., Yaakov, G., Willoughby, M., Homberg, B., Keller, J., Kupervaser, M., Dezorella, N., Dadosh, T., Wolf, S. G., Itkin, M., Malitsky, S., Brandis, A., Barkai, N., Fernandez-Busnadiego, R., Reddi, A., Rehling, P., Rapaport, D., and Schuldiner, M. The molecular mechanism of on-demand sterol biosynthesis at organelle contact sites, (2024). URL https://www.biorxiv.org/content/10.1101/2024.05.09.593285v2. Pages: 2024.05.09.593285 Section: New Results.

Supplementary Information

Figure S1. LBR-FKBP-GFP relocalisation does not cluster the plasma membrane anchor or other ER proteins.
Single slices from z-stacks of live HCT116 cells co-expressing LBR-FKBP-GFP (green), LBR-mCherry (**A**) or mCherry-Sec61β (**B**) (red) and SH4-FRB-EBFP2 (not shown in merge), treated with rapamycin (200 nM). Scale bars, 10 μm; Insets, 4× expansion of ROI.

Figure S2. HCT116 LBR-FKBP-GFP knock-in cells pool

(**A**) Schematic of C-terminal PCR tagging of LBR with FKBP-GFP. (**B**) FACS plots to show the collection of a mixed population of GFP positive cells. (**C**) Agarose gel to show genotyping PCR results using indicated primers. (**D**) Western blot of lysates collected from parental cells or the edited cell pool. Proteins were detected by anti-GFP or anti-LBR along with an actin loading control. Expected mass of tagged gene product (LBR-FKBP-GFP) and the untagged endogenous LBR are indicated by green and black arrows respectively. (**E**) Stills of single slices from a z-stack of live HCT116 LBR-FKBP-GFP (green) knock-in cells with SiR-DNA staining (magenta) progressing through mitosis. Time, mm:ss; scale bar, 10 μm; insets, 2× zoom.

Figure S3. Other proteins tested do not form clusters when relocalised to the plasma membrane or mitochondria.
Example micrographs of HCT116 cells co-expressing emerin, LAP2β or BAF construct tagged with FKBP-GFP at the N-terminus (green), with Stargazin-mCherry-FRB, pMito-mCherry-FRB (**C**) or a control with no FRB construct expressed (as indicated), stained with DAPI (blue). All samples were treated with rapamycin (200 nM) for 30 min prior to fixation. Shown are single slices from z-stacks of an interphase and metaphase cell. Scale bars, 10 μm; insets, 3× expansion of ROI.

Figure S4. Further examples of mitotic ER-Golgi contacts by SBF-SEM.
Single slices of metaphase (**A**) and telophase (**B**) HCT116 cell SBF-SEM datasets are shown. Depth of each slice within the dataset (nm) is indicated. Example Golgi clusters are shown by yellow arrows on the full slice image. Three sequential slices of these regions (3× expansion) are shown beside. Scale bars, 2 µm and 0.5 µm on zoom region.

Supplementary Videos

Figure SV1. Example of induced relocalization of LBR-FKBP-GFP to the plasma membrane.
Movie of a mitotic HCT116 LBR-FKBP-GFP knock-in cell co-expressing Stargazin-mCherry-FRB (not shown). Rapamycin (200 nM) is added between the first and second frame. Time, mm:ss. Playback, 10 fps. Scale bar, 10 µm.

Figure SV2. Example of LBR-FKBP-GFP relocalisation to the plasma membrane in HCT116 cells expressing MAPPER.
Movie of a mitotic HCT116 cell co-expressing LBR-FKBP-GFP (green, left), mScarlet-I3-6DG5-MAPPER (red, second channel), Stargazin-EBFP2-FRB (third channel) and stained with SiR-DNA (blue). Rapamycin (200 nM) is added between the first and second frame. Time, mm:ss. Playback, 2 fps. Insets, 3× expansion of ROI. Scale bar, 10 µm.

Figure SV3. Example of LBR-FKBP-GFP relocalisation to the plasma membrane in HeLa cells expressing MAPPER.
Movie of a mitotic HeLa cell co-expressing LBR-FKBP-GFP (green, left), mScarlet-I3-6DG5-MAPPER (red, second channel), Stargazin-EBFP2-FRB (third channel) and stained with SiR-DNA (blue). Rapamycin (200 nM) is added between the first and second frame. Time, mm:ss. Playback, 2 fps. Insets, 3× expansion of ROI. Scale bar, 10 µm.

Figure SV4. Example of induced relocalization of LBR-FKBP-GFP to mitochondria.
Movie of a mitotic HCT116 LBR-FKBP-GFP (green, left) knock-in cell co-expressing MitoTrap (pMito-mCherry-FRB, red, middle), stained with SiR-DNA (blue). Rapamycin (200 nM) is added during frame 2. Time, mm:ss. Playback, 10 fps. Scale bar, 10 µm.

Figure SV5. Example of induced relocalization of FKBP-GFP-Sec61β to mitochondria.
Movie of a mitotic HCT116 cell co-expressing FKBP-GFP-Sec61β (green, left), MitoTrap (pMito-mCherry-FRB, red, middle), stained with SiR-DNA (blue). Rapamycin (200 nM) is added between the first and second frame. Time, mm:ss. Playback, 1 fps. Scale bar, 10 µm.

Figure SV6. Golgi HCT116 video.
Movie of a mitotic HCT116 cell co-expressing LBR-FKBP-GFP (green, left), FRB-mCherry-Giantin3131-3259 (red, middle) and stained with SiR-DNA (blue). Rapamycin (200 nM) is added between the first and second frame. Time, mm:ss. Playback, 2 fps. Insets, 3× expansion of ROI. Scale bar, 10 µm.