Sonal Gupta[1], Simon C. Groen[1,2,3], Maricris L. Zaidem[1,4], Andres Godwin C. Sajise[5], Irina

Calic[6,7], Mignon A. Natividad[5], Kenneth L. McNally[5], Georgina V. Vergara[5,8], Rahul Satija[1,9],

Steven J. Franks[6], Rakesh K. Singh[5,10], Zoé Joly-Lopez[1,11*], Michael D. Purugganan[1*]

[1]Center for Genomics and Systems Biology, New York University, New York, NY USA;

[2]Department of Nematology and Department of Botany & Plant Sciences, University of

California, Riverside, CA USA; [3]Center for Plant Cell Biology, Institute for Integrative Genome

Biology, University of California, Riverside, CA USA; [4]Department of Biology, University of

Oxford, Oxford, England; [5]International Rice Research Institute, Los Baños, Philippines;

[6]Department of Biological Sciences, Fordham University, Bronx, NY USA; [7]Inari Agriculture

Nv, Gent, Belgium; [8]Institute of Crop Science, University of the Philippines, Los Baños,

Philippines; [9]New York Genome Center, New York, NY USA; [10]International Center for

Biosaline Agriculture, Dubai, UAE (current affiliation); [11]Département de Chimie, Université du

Quebéc à Montréal, Montreal, Quebec, Canada.

*Corresponding authors: Z.J.-L. (joly-lopez.zoe@uqam.ca); M.D.P. (mp132@nyu.edu)

Title: **Systems genomics of salinity stress response in rice**

Short Title: **Systems genomics of salinity stress response**

ABSTRACT

Populations can adapt to stressful environments through changes in gene expression. However, the role of gene regulation in mediating stress response and adaptation remains largely unexplored. Here, we use an integrative field dataset obtained from 780 plants of *Oryza sativa* ssp. *indica* (rice) grown in a field experiment under normal or moderate salt stress conditions to examine selection and evolution of gene expression variation under salinity stress conditions. We find that salinity stress induces increased selective pressure on gene expression. Further, we show that *trans*-eQTLs rather than *cis*-eQTLs are primarily associated with rice's gene expression under salinity stress, potentially via a few master-regulators. Importantly, and contrary to the expectations, we find that *cis-trans* reinforcement is more common than *cis-trans* compensation which may be reflective of rice diversification subsequent to domestication. We further identify genetic fixation as the likely mechanism underlying this compensation/reinforcement. Additionally, we show that *cis-* and *trans*-eQTLs are under different selection regimes, giving us insights into the evolutionary dynamics of gene expression variation. By examining genomic, transcriptomic, and phenotypic variation across a rice population, we gain insights into the molecular and genetic landscape underlying adaptive salinity stress responses, which is relevant for other crops and other stresses.

Keywords: phenotypic selection, linear selection, stabilizing selection, disruptive selection, eQTL, decoherence

INTRODUCTION

Plants face numerous stresses that reduce their growth and fitness, and they have a variety of adaptations to help them deal with environmental challenges (Mareri et al., 2022). For crop plants, stresses such as drought, heat and salinity can greatly limit productivity and agricultural sustainability worldwide (Fahad et al., 2017; Kopecká et al., 2023); indeed, as a direct result of various abiotic stresses, an estimated ~ 50-70% of crop yields are lost (Francini and Sebastiani, 2019). While there is a wealth of information on the physiological responses of plants, including crops, to stress (H. Zhang et al., 2022), we are still developing an understanding of the underlying genetic responses to stress and the complex interactions between stresses and environmental cues, networks of gene expression regulation, physiological responses and ultimately plant fitness (Cramer et al., 2011; Siddiqui et al., 2021).

Much of the research on the genetic basis of stress responses in crops has focused on identifying individual genes associated with specific responses and/or tolerance traits (Cui et al., 2020; Jiang et al., 2019; Kumar and Wigge, 2010; Laohavisit et al., 2013; Yuan et al., 2014). Stress responses, however, are complex traits that can be influenced by multiple genetic pathways with numerous interconnected genes. Expression levels of genes are controlled by regulatory elements and are often environmentally induced, suggesting a critical role of regulatory divergence in adaptive evolution (De Clercq et al., 2021; Wilkins et al., 2016). Furthermore, changes in gene expression can lead to changes in the transcript network correlational structure, reshaping regulatory pathways (Lea et al., 2019) and influencing the way in which gene expression variation can influence adaptation to stressful environments (Signor and Nuzhdin, 2018; Wagner and Lynch, 2008). As such, there is an increasing need to understand how evolutionary dynamics of gene expression and transcript abundance relate to the genetic underpinnings of stress responses and adaptation in crops (Groen et al., 2020; Ruffley et al., 2023; Siddiqui et al., 2021).

One of the most important stresses for many plants is salinity. Soil salinity causes osmotic imbalance between the plant and the soil, which impedes the uptake of water and other key nutrients (Hakim et al., 2014; Munns, 2002), possibly leading to acute ion toxicity (Liang et al., 2018). While some plants are considered halophytic and can thrive in saline environments, other plants are highly sensitive to salts and experience negative effects of salinity even at low concentrations (Carillo et al., 2011). *Oryza sativa* (Asian rice) is one such salt-sensitive crop,

facing significant yield loss due to soil salinity (Hussain et al., 2017; Melino and Tester, 2023).

Rice responds to salinity stress by adjusting physiological and biochemical processes involved in

osmotic and ion homeostasis, nutritional balances and oxidative stress (Castillo et al., 2007;

Miller et al., 2010; Munns and Tester, 2008; Qin and Huang, 2020; Wang et al., 2012). Although

studies have identified multiple genes associated with these processes (for reviews, see Ponce et

al., (2021) and Liu et al., (2022)), we still lack a systems-level understanding of how gene

expression variation mediates responses to salinity stress and evolves at the molecular level.

Here, we use an integrative system genomics approach to comprehensively dissect the

genome-wide molecular and phenotypic response to salinity stress in rice. This study builds on

prior work by our group that examined selection on gene expression in rice in response to normal

and dry conditions (Ćalić et al., 2022; Groen et al., 2021, 2020). Using genomic, transcriptomic,

and phenotypic datasets obtained from 130 diverse accessions of rice subjected to moderate

levels of salinity stress, we (i) explore the selection on gene expression variation under salinity

stress, (ii) dissect the genetic architecture of gene expression variation under saline conditions,

and (iii) identify genes, molecular pathways, and salt stress response traits as well as associated

trade-offs in the saline environment. We demonstrate that salinity stress induces increased

selective pressure on gene expression, and we identify variation in biological processes and

physiological traits that is beneficial and detrimental to plants in a saline environment, providing

novel insights into the molecular landscape underlying an adaptive response to excess salt. We

integrate these datasets with genomic sequence information to elucidate the genetic architecture

and regulatory networks governing rice's response to salinity stress.

RESULTS

Variation in transcript abundance. To understand the microevolutionary dynamics of gene

expression variation under salinity stress, we first investigated variation in transcript abundance

in 130 accessions of *O. sativa* ssp. *indica*. We conducted a field experiment in the dry season of

2017 (January-May) at the International Rice Research Institute in Los Baños, Laguna,

Philippines. Three replicates of each accession were planted separately in a normal wet paddy

field as well as a similar field in which plants were exposed to moderate salinity stress (salt

levels maintained at 6 dSm^{-1}) maintained until maturity. Average fecundity was significantly

lower in the saline field than in normal conditions (Fig. 1a; two-tailed paired t-test *P* = 1.658 x

10^{-8}) with most of genotypes having significantly lower fecundity in the saline field (n = 94; one-

sample proportion test $P = 3.639 \times 10^{-7}$).

To examine gene expression in the field, mRNA levels were measured in leaf blades

from 780 plants (130 accessions in triplicates for each environment) via 3'-end-biased mRNA

sequencing. Leaf blades were sampled 38 days after sowing (DAS), corresponding to 7 days of

plants being exposed to stress in the saline field. Population variance in gene expression of

18,141 widely expressed transcripts was partitioned into genotype (G), environment (E), and

genotype × environment (G×E) effects (Supplementary Table 1). At a conservative false

discovery rate (FDR) of 0.001, all but 3 transcripts displayed a significant genotype effect,

indicating that expression levels of most transcripts are heritable. This effect of genotype is

reflected in the broad-sense heritability (H^2) distribution of gene expression levels (Fig. 1b),

which had a median value $H^2 = 0.53$ (range of 0.012 - 0.987; Supplementary Table 1). In

addition to the high levels of heritability, 16,371 transcripts had a significant G×E term,

indicating that for many transcripts, genotypes showed heritably different levels of expression in

different environments. The significant G×E indicates genetic variation for plasticity, indicating

that this plasticity can evolve. Although we see a widespread heritable plastic response in gene

expression, we found little evidence of genotype-dependent plasticity in fitness (G×E for fitness

$P = 0.49$; Fig. 1a), indicating that the G×E of transcripts does not translate to the complex trait of

fitness. This could be due to a combination of factors, like gene interactions (pleiotropy and

epistasis) leading to little to no effect on fitness, or environment specific genotype-dependent

gene regulation (environment specific eQTLs). Furthermore, only a relatively small number of

transcripts (254) showed significant variation due to E, indicating genotype-independent

plasticity in gene expression for only a small fraction of genes.

Selection on gene expression. To identify transcripts associated with high fitness in normal and

saline environments, we measured the strength of selection on gene transcript levels. We did this

using phenotypic selection analysis (Lande and Arnold, 1983), taking the total number of filled

rice seeds/grains (total fecundity) as a proxy for fitness (Groen et al., 2020). We estimated the

linear (S) and quadratic (C) selection differentials, which are estimates for directional (negative

or positive S) selection, and stabilizing (negative C) or disruptive (positive C) selection. We

calculated the raw (S and C), variance-standardized (S_σ and C_σ), and mean-standardized

differentials (S_μ and C_μ). We found that both mean and variance of transcript expression vary

significantly between conditions (Mann-Whitney U-test, mean and standard deviation: $P < 2.2$ x

10^{-16}), which means that the interpretation of the strength of selection based on either of these

differentials could be misleading (Supplementary Table 2). To overcome this, and given that we

were interested in selection on gene expression, which were all measured in the same units, we

used the non-standardized raw selection differentials for all downstream analyses. We also report

the mean- and variance-standardized selection coefficients (Supplementary Table 3).

161   Most transcripts were (nearly) neutral ($|S| < 0.1$) in both the normal and saline

environments (Fig. 1c). Additionally, within the normal environment, there was a general trend

towards stronger positive compared to negative selection on gene expression, indicating that

higher expression of transcribed genes is associated with greater fitness (Table 1, Mann-Whitney

U-test, normal: $P = 3.95$ x 10^{-15}). In comparison, though the strength of positive selection was

higher than that of negative selection in the saline field, higher expression of most of the

transcribed genes is associated with lower fitness (Table 1, Mann-Whitney U-test, saline: $P =$

0.0068). Although no transcripts cleared the Bonferroni correction threshold in saline conditions,

under normal conditions seventeen transcripts cleared the threshold, thirteen of which were

under positive directional selection (Supplementary Table 3).

171   We also found that a high proportion of transcripts experienced stabilizing selection ($C <$

0) in both normal and saline environments. This effect was pronounced under normal conditions,

with the strength of stabilizing selection being stronger than that of disruptive selection (Table 1;

Mann-Whitney U-test, normal: $P < 2.2$ x 10^{-16}). In contrast, within the saline environment, there

was no detectable difference between stabilizing and disruptive selection (Mann-Whitney U-test,

normal: $P = 0.514$), due to a higher proportion of transcripts experiencing stronger disruptive

selection. This indicates that in the saline environment, as compared to normal conditions, an

increase in fitness is associated with extremes in transcript abundance.

179   Comparing the distribution of selection differentials between environmental conditions,

180   we found that selection was stronger in the saline field compared to the normal wet paddy field

181   (Table 1). This was true for the overall strength of directional selection (Mann-Whitney U-test,

182     $P=0.012$), as well as for negative ($P=8.27$ x $10^{-6}0.009$), stabilizing ($P < 2.2$ x 10^{-16}) and

183   disruptive ($P < 2.2$ x 10^{-16}) selection, but not positive selection ($P=0.735$).

Gene expression trade-offs. Next we compared the proportions of genes showing an opposite direction of selection in across environments (antagonistic pleiotropy- AP) and those showing selection in only one environment (conditional neutrality- CN) (Anderson et al., 2011). To account for the inherent bias associated with the detection of CN, we used a more stringent P-value cutoff to define CN ($P < 0.025$) in comparison to AP ($P < 0.05$) (Groen et al., 2020; Siddiqui et al., 2021). We found that 10.80% of the transcripts showed selection patterns consistent with CN, while only 0.28% of transcripts showed AP (Fig. 1f; Supplementary Table 4). The proportion of transcripts showing CN was greater than expected by chance and greater than the proportion showing AP (two-tailed proportion z-test $P < 2.2 \times 10^{-16}$), even accounting for the fact that there is an inherent bias favoring detection of CN (Anderson et al., 2013). These results are consistent with a lack of trade-offs in which the expression of a gene is favored in one environment and disfavored in another.

Among the 51 AP transcripts, increased expression of 38 was beneficial only in normal conditions (higher expression associated with higher fitness in normal conditions, but lower fitness under saline conditions). Gene ontology (GO) term analyses of these 38 transcripts indicated that many of these genes were involved in photosynthesis and metabolic processes (Supplementary Fig. S1; Supplementary Table 4). This is consistent with an observed reduction of photosynthesis in rice under salinity stress (Radanielson et al., 2018; Tsai et al., 2019).

We then wondered whether any difference underlie gene regulation of these AP transcripts relative to the non-AP transcripts, potentially indicating the genetic basis for this trade-off. We identified single nucleotide polymorphisms (SNPs) associated with transcript expression levels in each environment separately using whole-genome polymorphism data (expression quantitative trait loci [eQTL] analyses; see below). We found SNPs regulating expression of two photosynthesis related AP transcripts (*PSAN* and *CRR7*) only in the normal environment (Supplementary Fig. S2). We further looked at the 13 AP transcripts that were beneficial only in the saline environment (Supplementary Table 4). Although there was no significant GO enrichment for these transcripts, among them we identified a cyclophilin-encoding transcript (*OsCYP2*), which has been shown to confer salt tolerance in rice (Lee et al., 2015; Roy et al., 2022; Ruan et al., 2011). However, no SNP was associated with the expression of this gene in either normal or saline conditions.

Biological processes under selection. To investigate the broader biological processes associated with differential selection (strong selection in only one environment), we ranked all GO biological processes by their median selection strength in each environment and identified the processes with significantly stronger selection relative to their respective environment. We identified 13 and 18 processes that were under strong differential selection in normal and saline conditions, respectively (Supplementary Table 5; Fig. 2a). Processes primarily involved in various aspects of growth and defense were under stronger selection in normal conditions, whereas processes associated with regulation of flowering, cell cycle control and reproduction showed stronger selection under saline conditions (Fig. 2a). This provides insight into specific biological processes involved in flowering time regulation and reduced yield associated with salinity stress, as has been observed in multiple species (Chang et al., 2019; Li et al., 2007; Van Zandt and Mopper, 2002; R. Zhang et al., 2022). Furthermore, studies have found that the osmotic stress induced by salinity causes a reduction in the cyclin-dependent kinases (CDKs) responsible for cell-cycle transitions (G1/S and G2/M) (Ma et al., 2015; Schuppler et al., 1998; West et al., 2004). Aligned with this, our study also supports the notion that salinity stress affects cell cycle regulation, and leads to reduced growth and reproduction.

Gene expression usually operates within the context of robust gene interaction/regulatory networks (Amiri et al., 2018; Israel et al., 2016; Ko and Brandizzi, 2020). The fact that genes interact in networks means that selection on one gene can lead to indirect selection on interacting genes, which can constrain the response of a population to selection on gene expression (Agrawal and Whitlock, 2010; Groen et al., 2020; Kondrashov and Houle, 1997). To examine this phenomenon, we identified suites of correlated transcripts using principal component (PC) analysis on genome-wide gene expression levels. We then estimated the linear (β) and quadratic selection (γ) gradients for PCs explaining over 0.5% of variance in each environment (Supplementary Table 6). Although quadratic selection was generally weak, linear selection on some PCs showed significant directional selection (Supplementary Table 7).

Using the breeder's equation (Falconer and Mackay, 1996), we predicted the response to selection on these PCs to examine the constraints on a population's microevolutionary response to selection on gene expression. We found that over half of PCs (7 of 11 and 7 of 12 PCs in normal and saline conditions, respectively) displayed opposite signs of direct and indirect responses to selection (Supplementary Table 7). This finding contrasts with prior work in rice

under dry and wet conditions that found a lack of constraint in response to selection for most

traits except seed size because direct and indirect responses to selection were largely similar

(Ćalić et al., 2022), indicating that different types of stress may either constrain or facilitate the

response to selection.

We found PCs enriched for metabolic pathways and biosynthesis of phenylpropanoids

and secondary metabolites to be under selection in the normal environment (Fig. 2a;

Supplementary Fig. S3; Supplementary Table 7). Different transcripts involved in these

pathways were under positive and negative selection which may act to keep these pathways in a

steady-state. Interestingly, circadian rhythm was found to be under positive selection, with an

overall positive response to selection, in the saline environment (Fig. 2a; Supplementary Fig. S3;

Supplementary Table 7). This is in alignment with the role of circadian clock genes in conferring

salt tolerance (Kim et al., 2013; Wei et al., 2021; Xu et al., 2022), and indicates a tentative

increase in expression of circadian clock genes with continuous exposure to soil salinity.

Salinity stress induces decoherence. Gene expression levels are generally correlated, but these

correlations can be perturbed by environmental stresses, a phenomenon that is termed

decoherence (Lea et al., 2019). Such decoherence has been demonstrated in humans and primates

(Lea et al., 2019; Pu et al., 2022; Watowich et al., 2022), but little is known about how stress

alters the correlation structure of transcript levels in plants. To examine decoherence in rice, we

utilized the recently developed CILP (Correlation by Individual Level Product) method (Lea et

al., 2019), which detects the systematic loss of correlation in gene expression among individuals.

Since CILP calculates product correlations for all possible pairs of genes, we used transcripts

with selection strengths greater than 0.1 ($|S| > 0.1$) in at least one environment with expression

greater than 0 in at least 50% of individuals (2,051 transcripts; Supplementary Table 8) to reduce

data dimensionality.

We found that the correlation structure for gene expression was broadly similar across the

2.10 M unique transcript pairs, but > 29,000 transcript pairs (involving 1,742 unique transcripts)

showed significantly different correlations between normal and saline conditions (FDR < 5%)

(Fig. 3a; Supplementary Table 8). This change in correlation structure indicates possible

divergence in gene interactions between environments, which may presage a restructuring of

gene networks. GO term enrichment analysis of the transcripts that show decoherence with

significant pairs greater than the median (median significant pair per transcript = 12, n = 853)

highlighted important pathways related to plant responses and potentially tolerance to excess salt

(Fig. 3b). For instance, circadian rhythm genes like *OsPRR37* and GIGANTA (*GI*), which have

been shown to confer salt tolerance in rice and Arabidopsis (Kim et al., 2013; Wei et al., 2021),

differed significantly in their interactions between the two environments. Similarly, to manage

the energy requirements of salt stress, the relative abundance of metabolites involved in energy

producing pathways like glycolysis, tricarboxylic acid (TCA) cycle and various other metabolic

processes have been shown to be altered as an early response to salt stress (Che-Othman et al.,

2017; Jacoby et al., 2011; Yang and Guo, 2018). Additionally, aligned with our results, multiple

studies have reported a positive correlation between salt tolerance and levels of secondary

metabolites and amino acids with osmoprotectant properties associated with lowering osmotic

stress (Krishnamurthy and Bhagwat, 1990; Petrusa and Winicov, 1997; Tari et al., 2010). Our

results suggest that salt exposure induces decoherence of gene expression in some transcripts,

that this decoherence results from a restructuring of the gene expression network, and this

restructuring can allow salt stress tolerance, providing a potential molecular mechanism

underlying this tolerance.

Selection on traits. In addition to gene expression data, we also collected phenotypic data for 13

organismal traits in normal and saline conditions, which provides us with the opportunity to

examine how salt stress influences complex phenotypes and identify connections between

variation in traits with fitness consequences and underlying patterns of gene expression. Since

these traits were measured on different scales, we estimated variance-standardized selection

gradients on these traits, focusing only on seven traits that were not strongly correlated to limit

the contribution of indirect selection (Pearson correlation coefficient < 0.6; Supplementary Fig.

S4). We identified three traits – leaf osmotic potential (LOP), chlorophyll a content (Chl_a), and

flowering time (FT) – that displayed different selection patterns in normal versus saline

environments (Fig. 4; Supplementary Table 8).

Leaf osmotic potential is a key trait associated with water transport in plants, and which

decreases with increasing salinity, leading to low water uptake by the plant. This trait was under

positive selection in the control environment but under stabilizing selection under saline

conditions, which suggests that an optimal LOP is important under salt stress (Fig. 4).

Furthermore, we found Chl_a content at the reproductive stage to be under negative selection in

saline conditions. Although ion toxicity has previously been shown to reduce chlorophyll content

(Ashraf and Bhatti, 2000; Taïbi et al., 2016), our results showed that this reduction is associated

with increased survival and reproductive fitness, consistent with the general trend for reduced

photosynthesis under salinity stress. We also found that FT was under positive selection

(selection for later flowering) in normal conditions but under stabilizing selection in saline

conditions (Fig. 4). Moreover, FT was significantly reduced under saline conditions (two-tailed

paired t-test $P = 0.0012$; Supplementary Fig. S5), implying that salt stress selects for an optimal

flowering time, with the trend shifted towards earlier flowering compared to that in the normal

wet paddy.

Genetic architecture of gene expression variation. To dissect the genetic architecture of gene

expression variation, we identified expression quantitative trait loci (eQTLs) and examined

whether and how selection acts on these eQTLs. We identified *cis-* and *trans*-eQTLs (FDR <

0.001) regulating expression of 3,065 and 3,277 genes in normal and salinity stress conditions,

respectively (Supplementary Table 9). We observed that 49.64% (29,623 of a total 59,669) of

cis-eQTLs were common in both environments, as compared to 18.62% (25,528 of a total

137,100) of *trans*-eQTLs; this result was robust to FDR cutoff (FDR of 0.01 and 0.05).

Moreover, this is consistent with previous observations from studies on other species that have

found 48%-77% overlapping *cis*-eQTLs and 9%-60% common *trans*-eQTLs across

environments (Smith and Kruglyak, 2008; Snoek et al., 2012; Sterken et al., 2023). It further

indicates that *trans*-eQTLs might be more environment-specific than *cis*-eQTLs. We tested this

explicitly by identifying loci showing gene-environment interaction (G × eQTL) and found that

at FDR of 0.05, *cis*-eQTLs constituted merely 0.28% (142 of 50718) of the total identified G ×

eQTLs.

We identified eQTL hotspots, which are regions of the genome that are associated with

expression variation of a large number of genes (Qu et al., 2018). These regions can occur either

due to low amounts of recombination (high linkage disequilibrium- LD) or because they contain

master-regulators that pleiotropically control expression of multiple functionally-associated

genes (Hammond et al., 2011; Kliebenstein, 2009; West et al., 2007). To account for LD and

identify regions likely to contain master-regulators, we chose to focus on the subset of genes

regulated by the lead-SNPs (SNP with the most significant association) in a given 100-kb region.

Through this approach, we identified 11 *trans*-eQTL hotspots (number of unique genes > 30) in

saline conditions, but none in normal conditions (Supplementary Fig. S6; Supplementary Table

10). These results indicate that gene regulation under stress conditions may be dependent on a

few master-regulators, as has been shown for drought stress in rice and maize (Kuroha et al.,

2017; Liu et al., 2020). Interestingly, one of the hotspots, Chr7: 25.9-26.0Mb, influenced the

expression of a disproportionately high number of genes (191 genes) and contains the gene

OsFLP, a R2R3 myb-like transcription factor that regulates stomatal development (Wu et al.,

2019). *OsFLP* has recently been associated with salt tolerance in rice (Qu et al., 2022).

It has been shown previously that when a gene is regulated via both *cis* and *trans* factors,

their effects tend to drive target gene expression in opposite directions, canceling the combined

effect on expression. This is commonly referred to as *cis-trans* compensation, which is thought

to arise due to stabilizing selection that maintains similar gene expression over evolutionary

timescales in the face of new mutations (Coolon et al., 2014; Goncalves et al., 2012; Landry et

al., 2005; Lovell et al., 2018). Contrary to this expectation, we found that most genes appear to

have a reinforcing (*cis* and *trans* effect in the same direction) rather than a compensatory pattern

(two-tailed proportion z-test $P = 1.65 \times 10^{-11}$; Fig. 5a; Supplementary Table 11). Among the 524

cis-trans co-occurring genes in saline conditions, 317 were reinforcing (~60.5%) and 207 were

compensatory (39.5%). This pattern held for normal conditions as well (Supplementary Fig. S7a;

Supplementary Table 11). Thus, expression patterns of these genes are expected to either

increase or decrease expression over evolutionary timescales, indicating the presence of

directional selection.

We hypothesized that this excess of reinforcing rather than compensatory effects could be

driven by the extensive diversity among rice landraces that arose from crop diversification

following domestication (Meyer and Purugganan, 2013). To test this hypothesis, we examined

whether genes under *cis-trans* reinforcement showed evidence of higher inter-varietal variation

in gene expression (population-wide variance between mean accession expression levels) as

compared to genes under *cis-trans* compensation. Supporting our hypothesis, we found

significantly higher inter-varietal expression variation for genes under *cis-trans* reinforcement in

saline (Fig. 5b; Mann-Whitney *U*-test = 0.002; mean-inter-varietal expression variation

compensating = 1.12, reinforcing = 1.27) and in normal (Supplementary Fig. S7b) conditions.

Furthermore, it has been suggested that *cis-trans* compensation/reinforcement can arise due to

the genetic fixation of compensating/reinforcing *trans*-regulatory variants (McManus et al.,

2014; Signor and Nuzhdin, 2018), which could lead to elevated linkage disequilibrium (LD)

between the *cis*- and *trans*-regulatory variants acting on a gene. This was indeed the case:

estimated LD between all pairs of *cis*- and *trans*-regulatory variants for the identified

compensating and reinforcing genes was significantly higher as compared to the background LD

(mean r^2 compensating: 0.33, reinforcing: 0.36, background = 0.056; two-tailed permutation test

$P = 0.0019$).

To examine the pattern of past selection acting on *cis*- and *trans*-eQTLs (before, during

and after rice domestication, including the crop diversification phase) we plotted the folded site-

frequency spectrum (SFS) of the minor allele frequencies and inferred the relative strength of

historic selection (Joly-Lopez et al., 2020) We found that *trans*-eQTLs for both environments

had been under strong purifying selection, with the SFS being significantly left-shifted compared

to background SNPs (two-tailed t-test $P < 10^{-16}$; mean MAF background = 0.20, *trans* normal =

0.0936, *trans* saline = 0.0910). Not surprisingly, given their highly pleiotropic nature, this effect

was more pronounced for *trans*-eQTLs regulating multiple genes (two-tailed t-test $P < 10^{-16}$;

normal mean MAF in unique *trans*-eQTLs = 0.099, multiple *trans*-eQTL = 0.073; saline mean

MAF in unique *trans*-eQTLs = 0.097, multiple *trans*-eQTL = 0.071). In contrast to *trans*-eQTLs,

we found that *cis*-eQTLs in both normal and saline conditions had potentially been under

balancing selection (Fig. 5c, Supplementary Fig. S8) with the SFS being significantly right-

shifted for *cis*-eQTLs relative to background SNPs (two-tailed t-test $P < 10^{-16}$; mean MAF

background = 0.20, *cis* normal = 0.238, *cis* saline = 0.237). Moreover, for both conditions, the

estimated nucleotide diversity (π) for 100-kb regions flanking *cis*-eQTLs was also significantly

higher as compared to genome-wide 100-kb blocks (two-tailed t-test $P < 10^{-11}$; mean π

background = 0.286, *cis* normal = 0.297, *cis* saline = 0.297).

Comparing patterns of past selection for *cis*-eQTLs specific to each condition revealed no

significant difference between the normal and saline environments. However, we did observe

trans-eQTLs in saline conditions to be under stronger purifying selection (two-tailed t-test $P = $

0.017). Taken together, this indicates that *cis*- and *trans*-eQTLs are under different selection

regimes in rice, and that purifying selection is stronger on *trans*-eQTLs regulating gene

expression under salt stress conditions, giving us insights into the macroevolutionary dynamics

of gene-expression variation.

DISCUSSION

Gene expression is a key link in the chain of adaptive organismal responses of organisms to

environmental challenges. In this study, we used a systems genomics approach to examine

genome-wide transcript levels in rice in both normal and moderately saline field conditions to

examine the evolutionary response and genetic architecture of gene expression under salinity

stress. We found selection on expression of a set of genes, and for some genes, selection on

expression differed among environments, indicating that salinity can select for changes in gene

expression. We saw that the genetic regulatory pathways can be modified by salinity, as

indicated by decoherence, which provides a potential genetic mechanism underlying salinity

tolerance. We found limited evidence for trade-offs given a lack of antagonistic pleiotropy in the

fitness effects of expression compared across environments, and we also found that there did not

appear to be strong cis-trans compensation in gene regulation. These results provide us with

insights into the fitness consequences and genetic architecture of gene expression under salt

stress in rice, as we discuss below.

418   One of the primary goals of this study was to characterize selection on gene expression

and how this selection varies between saline and normal conditions. Before discussing our

results, it is important to note that our conclusions are influenced by the way in which selection

differentials are standardized. Selection differentials are estimated through regression

coefficients in the relationship between trait values and fitness (Lande 1979). Because traits are

often measured on different scales, selection differentials are usually standardized to allow for

meaningful comparisons and interpretation. The most common method of standardization is

variance standardization, in which traits are standardized to a mean of zero and a standard

deviation of one (Lande and Arnold, 1983). With this approach, the selection differential,

multiplied by the heritability, gives the expected response to selection in standard deviation

units. So while this is a valid approach, there is concern that when variance-standardized

selection differentials are interpreted as reflecting the "strength" of selection based on their

magnitude, they can be misleading because trait variance can vary among traits and

environments, such that the trait variance and the magnitude of selection are conflated (Hereford

et al., 2004). Instead, the use of mean-standardized selection estimates have been suggested

(Hereford et al., 2004; Matsumura et al., 2012). However, it has not been widely recognized that

mean-standardized selection estimates can also be biased if mean trait values vary consistently

across conditions, such that magnitude of selection is conflated with trait means. We show here

that this was indeed the case: the mean gene expression values differed consistently between

saline and normal conditions, leading to biases when comparing the strength of selection across

treatments when mean-standardized selection differentials were used. Specifically, using mean-

standardized selection coefficients, we found selection on gene expression to be stronger under

normal conditions, whereas unstandardized and variance-standardized selection coefficients

indicated stronger selection under saline conditions. This result indicates that it is important for

researchers to carefully consider the advantages and potential drawbacks of both variance-

standardized and mean-standardized, as well as unstandardized, selection differentials in making

decisions about which to use in any particular system and depending of the goals of the study,

but also to consider these ideas in interpreting selection differentials as reflecting the strength of

selection.

Because we found that both variance-standardized and mean-standardized selection

gradients were biased, as explained above, we draw our conclusions about patterns of selection

using unstandardized selection gradients. We believe this approach to be appropriate given that

gene expression values are measured on the same scale and normalized for every transcript, and

our goal is to identify transcripts under selection and compare selection across treatments. Using

unstandardized selection coefficients, our study shows that most genes appear to have nearly

neutral levels ($|S| < 0.1$) of selection on transcript levels in both normal and saline environments.

This is consistent with previous studies conducted in various plant and non-plant species, which

have shown that most traits (including transcript abundance) are under very weak selection, and

only a few traits experience strong selection at microevolutionary timescales (Ahmad et al.,

2021; Hoekstra et al., 2001; Kingsolver et al., 2001). There does seem to be a tendency,

however, towards stronger positive selection over negative selection on gene expression,

meaning that higher expression of a majority of genes is generally associated with higher fitness.

Looking at quadratic selection, we found evidence for both stabilizing and disruptive selection

on gene expression. Our results are consistent with the expectation of stabilizing selection being

more common than disruptive selection in normal conditions (Kingsolver et al., 2001), and

reinforce the idea that disruptive selection is more widespread under stress conditions (Ahmad et

al., 2021), potentially reflecting the prevalence of frequency- and density-dependent competition

for resources under stress (Kingsolver and Pfennig, 2007).

Comparing the distribution of selection coefficients, we further found that selection was

stronger in moderate stress saline field conditions compared to normal wet paddy field

conditions, indicating that salinity stress induced increased selective pressure on gene expression

at microevolutionary timescales. Although exposure to salinity stress increased the strength of

selection on gene expression, this increase was relatively low compared to what has been

reported for drought stress in rice (Agrawal and Whitlock, 2010; Groen et al., 2020; Kondrashov

and Houle, 1997). This could potentially be attributed to the levels of stress experienced by the

plants – the drought stress treatment was more severe in terms of fitness after stress exposure

(Agrawal and Whitlock, 2010; Groen et al., 2020; Kondrashov and Houle, 1997) in comparison

to the moderate salinity stress in this study – but further work is needed to examine whether there

is indeed a relationship between stress intensity and selection strength. While this work provides

insight into the direction of selection and the predicted degree of evolutionary change under the

specific conditions of this study, more work would need to be done over multiple generations

and under additional conditions to make more realistic predictions of microevolutionary change

in the field.

Our work contributes to an ongoing debate regarding whether the intensity of selection

on gene expression increases under stress. We found in this study of rice and salinity, and in a

prior study on drought (Groen et al. 2020), that directional selection was greater under stress

compared to normal conditions. This finding is consistent with prior studies on other systems

like fruit flies (*Drosophila melanogaster*) (Groen et al., 2020; Jasnos et al., 2008; Kondrashov

and Houle, 1997) and yeast (Groen et al., 2020; Jasnos et al., 2008; Kondrashov and Houle,

1997), but contrasts with a recent study in *D. melanogaster* reporting no change in selection due

to increasing levels of nutritional stress (Arbuthnott and Whitlock, 2018). Our results also

support prior findings in rice (Ćalić et al., 2022; Groen et al., 2020) and in *Boechera stricta*

(Anderson et al., 2013) that antagonistic pleiotropy occurs and can be important for local

adaptation, while conditional neutrality in gene expression may be much more common. Our

finding indicates that it may be feasible to breed salinity tolerant rice varieties without a yield

penalty under non-saline conditions, though further work is needed to verify this.

We examined the genetic architecture of gene expression variation as well, and found that

trans-eQTLs rather than *cis*-eQTLs are primarily associated with rice gene expression under

salinity stress, potentially via a few master-regulators. *Trans*-eQTLs may be important in

environment-dependent gene expression changes, while *cis*-eQTLs may be more robust to

environmental changes as has been observed in other species (Smith and Kruglyak, 2008; Snoek

et al., 2012; Sterken et al., 2023). This can be attributed to *trans*-eQTLs' larger mutational target

along with a larger effect of drift than of positive selection in fixing them. These results further

corroborates with other studies that have found a more dominant effect of *trans*-eQTLs on

within-species variation than between-species variation (Emerson et al., 2010; Metzger et al.,

2016; Wittkopp et al., 2008).

We found that *cis*- and *trans*-eQTLs show different patterns of selection, with *cis*-eQTLs

showing evidence for balancing selection and *trans*-eQTLs showing purifying selection. We see

this pattern for rice, a largely selfing species, in contrast to outcrossing species where both *cis*-

and *trans-eQTLs* have been found to be under negative selection (Hernandez et al., 2019;

Josephs et al., 2015). Future studies will be needed to investigate whether the mating system has

an influence on the pattern of natural selection acting on variants regulating gene expression.

Finally, and contrary to expectations, we show that under saline conditions, *cis-trans*

reinforcement is more prevalent than *cis-trans* compensation. This result may be driven by rice

domestication and subsequent population diversification. Additionally, we find significantly

elevated levels of LD among the *cis*- and *trans*-regulatory variants for compensating and

reinforcing genes, indicating the role of genetic fixation as an underlying mechanism for *cis*-

trans compensation/reinforcement.

Systems genomic approaches provide insights into large-scale patterns across genome-

wide data over multiple scales. This approach can also identify possible genes or genetic

pathways that may prove critical in various processes. Our analysis, for example, shows that

many of the potential antagonistically pleiotropic genes are involved in photosynthesis and

metabolic processes. Importantly, we found that the circadian rhythm pathway was under

positive selection (selection for increased expression) in saline conditions and that the genes

involved in circadian rhythm showed significant decoherence between the two environments.

These findings make sense in light of the fact that circadian rhythm has been shown to regulate

salt tolerance and flowering time in rice (Liang et al., 2021; Wei et al., 2021), so responses to salt

stress may alter the expression patterns of genes in the circadian rhythm network, leading to

decoherence. In further support of this idea of a link between salt tolerance and circadian rhythm,

our selection analyses on physiological traits shows that salt stress leads to selection for earlier

flowering. Additionally, our decoherence analyses identified transcripts in important pathways

related to plant responses and potentially tolerance to excess salts, including the carbon

metabolic pathways (glycolysis and tricarboxylic acid cycle), and accumulation of secondary

metabolites and sugar moieties with osmoprotectant properties (Jacoby et al., 2011;

Krishnamurthy and Bhagwat, 1990; Petrusa and Winicov, 1997; Tari et al., 2010; Yang and Guo,

2018).

Other genes that are beneficial only in the saline environment include a cyclophilin-

encoding transcript (*OsCYP2*), which has been shown to confer salt tolerance in rice (Ruan et al.,

2011). Moreover, we also identified eQTL variants regulating expression of two photosynthesis

related antagonistically pleiotropic transcripts (*PSAN* and *CRR7*) in the normal environment.

Finally, our *trans*-eQTL analysis identified a hotspot on chromosome 7 that contains *OsFLP*, a

R2R3 myb-like transcription factor that regulates stomatal development (Wu et al., 2019), and

appears to be involved in salt tolerance in rice (Qu et al., 2022). Together, these loci are possible

new targets for functional and translational studies of salinity stress in rice. Coupled with the

insights gained by a systems genomic approach in inferring large scale patterns from genome-

wide information, this integration of data across multiple scales across a rice population has

allowed us to provide an integrated examination of the molecular and genetic landscape

underlying adaptive plant salinity stress responses.

MATERIALS AND METHODS

Plant Material. Domesticated rice is primarily classified into two distinct genetic subgroups, *O.*

sativa ssp. *indica* and *O. sativa* ssp. *japonica*. These subgroups are grown in sympatry and are

often recognized as subspecies, given the reproductive barriers between them (Nadir et al.,

2018). Further analyses have identified a widely accepted classification consisting of genetically

distinct varietal groups namely, *indica, aus/circum-aus, aromatic/circum-basmati, tropical*

japonica, and *temperate japonica* (Garris et al., 2005). For this study, a total of 130 *O. sativa* ssp.

indica (including *indica* and *circum-aus* groups) and 65 *O. sativa* ssp. *japonica* (including

circum-basmati, tropical japonica, and *temperate japonica*) accessions were selected, including

traditional varieties/landraces and three additionally replicated salt-sensitive and -tolerant test

varieties. We focused our analyses on *O. sativa* ssp. *indica* since it is the predominant global

varietal group (Supplementary Table 12). Seeds were obtained from the International Rice

Genebank Collection (IRGC) at the International Rice Research Institute (IRRI) in the

Philippines, and from a 2016 bulk seed collection obtained from plants grown under normal (wet

and non-saline) conditions at IRRI during the course of a previous study (Groen et al., 2020).

Field Experiment. The field experiment was conducted in the dry season of 2017 at IRRI in Los

Baños, Laguna, Philippines. Seeds from each accession were sown on December 16, 2016, and

seedlings were then transplanted into the experimental fields at 17 days after sowing (DAS), on

January 5, 2017. The field experiment was conducted across two locations: site L4

(14°09'34.6"N 121°15'42.4"E) was prepared as the non-salinized "normal" environment and site

L5 (14°09'35.2"N 121°15'42.5"E) as the salinized environment, following Groen et al., (2020).

Within each field environment, there were three blocks and three replicates of each genotype

(accession), with each genotype planted once per block in a random location. Each plant was

planted in a single-row with 0.2-m × 0.2-m spacing between them for a total of one focal plant

and seven neighboring plants (included in the experiment) per plot. Each experimental plot

included the accessions NSIC Rc 222 and NSIC Rc 182 that served as border rows

(Supplemental Fig. S9; Supplemental Table S13). The application of salt in site L5 started on

January 19, 2017, when the plants were 31 days old. The salinity level was monitored by

recording electrical conductivity (EC), using EC meters installed in each of the parcels at a depth

of 30 cm. The EC levels were recorded twice per day until reaching an EC=6 dSm^{-1} and then

recorded daily. The salinity levels were then maintained at 6 dSm^{-1} (considered mild to moderate

salinity stress) until maturity. Management and maintenance of the fields included the

application of basal fertilizer, spraying of insecticides against thrips and removal of plants

potentially infected with rice tungro virus.

Tissue collection for transcriptome sequencing. Leaf sampling was performed as previously

described (Groen et al., 2020). Briefly, leaf collection in the non-saline and saline field was done

from 10:00h to 12:00h at 38 DAS (8 days after the beginning of the salt treatment) in the non-

saline and saline field from 10:00h to 12:00h. Both fields were sampled simultaneously and with

individuals within a block collected in the same order. For each sample, about 10 cm of leaf

length were cut into small pieces and placed in chilled 5-mL tubes containing 4 mL of RNALater

(Fisher Scientific) solution for RNA stabilization and storage. Leaf samples from each of

the 5-ml tubes were then transferred into pairs of 2-mL tubes (one for processing and

one for backup), then stored at − 80 °C.

Yield harvesting and panicle trait phenotyping. A total of 780 plants were harvested

individually and labeled such that the yield of all plants used for each type of measurement

(mRNA sequencing, phenotypic measurements) was known. Individual seeds collected were

further categorized as filled, partially filled, and unfilled, using manual assessment and a seed

counter (Hoffman Manufacturing). Panicle length measurements and panicle trait phenotyping

(PTRAP) of 30 seeds were also performed.

Functional trait phenotyping. In addition to yield-related measurements, we collected data on a

number of physiological, morphological, and phenological traits to assess differences between

rice accessions in response to soil salinity. In both the non-saline and saline fields, we recorded

leaf osmotic potential (LOP) in the vegetative stage, performed chlorophyll analysis based on 1-

mg leaf samples and measured ion content (analysis of sodium, Na^+, and potassium, K^+,

analysis) based on 20-mg of leaf samples in both the vegetative and reproductive stages. We also

measured plant height for growth rate (measured once a week until maturity). Flowering time

was recorded as the day on which 50% of plants in a plot flowered; these plants included the

focal plant and its seven neighboring plants. Whole plants were both harvested at the vegetative

stage and at maturity to measure wet and dry biomass.

Extraction of total RNA for library construction. Leaf samples stored at −80 °C were thawed

at room temperature briefly and excess RNALater was removed. Tissue samples were then flash-

frozen in liquid nitrogen and ground using a TissueLyser II (Qiagen). After this, total RNA was

extracted using the RNeasy Plant Mini Kit according to manufacturer's protocol (Qiagen) and

eluted in nuclease-free water. The integrity of total RNA was assessed by agarose gel

electrophoresis, and RNA from a random subset of samples was further assessed by Agilent

TapeStation (Agilent Technologies). RNA concentration was quantified on a Qubit (Invitrogen).

Samples were stored at −80 °C until library preparation.

RNA-seq library preparation and sequencing. Library preparation for 780 samples was

performed as described in Groen et al. (2020) and followed a plate-based 3′-end mRNA

sequencing (3′ mRNA-seq) protocol. Briefly, total RNA from each sample was transferred

individually into 96-well plates and normalized to a concentration of 10 ng in 50 μL nuclease-

free water. Then, mRNA samples were reverse-transcribed using Superscript II Reverse

Transcriptase (Thermo Fisher Scientific) and cDNAs were amplified using the Smart-seq2

protocol (Picelli et al., 2013) with modifications (Groen et al., 2020). This resulted in

multiplexed pools of 96 samples each that were used for library preparation with the Nextera XT

DNA sample prep kit (Illumina), returning 3′-biased cDNA fragments, similar to the Drop-seq

protocol (Macosko et al., 2015). The resulting cDNA libraries were then quantified on an Agilent

BioAnalyzer and sequenced at the NYU Genomics Core on an Illumina NextSeq 500 with the

configuration HighOutput 1 x 75 base pairs (bp) and the settings: Read 1 of 20 bp (bases 1–12,

well barcode; bases 13–20, unique molecular identifier (UMI)) and Read 2 of 50 bp. Raw

sequence reads have been submitted to the SRA (BioProject PRJNA1010833).

RNA-seq data processing and data normalization. 3′ mRNA-seq read data were processed as

previously described in Groen et al. (2020). Briefly, Drop-seq tools v1.12

(https://github.com/broadinstitute/Drop-seq) and Picard tools v2.9.0

(https://broadinstitute.github.io/picard/) were used to generate the metadata. The reference

genome, Nipponbare IRGSP 1.0

(ftp://ftp.ncbi.nlm.nih.gov/genomes/all/GCF/001/433/935/GCF_001433935.1_IRGSP-1.0), and

annotations were indexed with STAR v020201 (Dobin et al., 2013). Prior to generating read

counts, raw reads were converted from FASTQ to unaligned BAM format using Picard tools

FastqToSam before being processed using the unified script for a FASTQ starting format. After

this, digital gene-expression matrices displaying either UMI or raw read counts with transcripts

as rows and samples as columns containing counts from reads with 96 expected sample barcodes

were produced using the DigitalExpression utility. Sample barcodes corresponding to beads

never exposed to rice total RNA were filtered out based on low numbers of transcribed elements

as described previously (Groen et al., 2020; Macosko et al., 2015). Rice individuals that ended

up being discarded due to low numbers of transcribed elements, were sequenced again using

another library.

UMI counts per sample were normalized through dividing by the total number of

detected UMIs in that sample and multiplying by 1×10^6 to obtain transcripts per million. The

resulting data matrices were then merged into one digital gene-expression super-matrix,

containing transcripts-per-million expression data for all samples. Elements with very low

transcription levels (transcript models with a sigma signal < 20) were discarded, after which a

robust normalization was conducted using an invariant set normalization protocol within the

DChip utility v2010.01 (Li and Wong, 2001). All downstream analyses were done in log-space,

using normalized expression levels ($\log_2$[normalized transcripts-per-million value $+ 1$]) of

transcribed elements estimated using R v3.4.3 (Computing, 2013; Robinson et al., 2010). In a

final step of filtering, transcripts that were not detected in at least 10% of individuals across our

populations and did not derive from protein-coding genes on nuclear chromosomes were

removed prior to performing subsequent analyses.

Quantitative genetics of fecundity and gene expression. The effect of genotype (G),

environment (E) and genotype-by-environment (G×E) on fecundity (number of filled grains) was

assessed using two-way analysis of variance (ANOVA) with E as a fixed effect, and G and G×E

as random effects. The significance of each term was determined using the F-tests (fixed effect)

and likelihood ratio test (random effects). Fecundity, averaged by genotype, was further

compared between the environments using a two-tailed paired t-test. Variation in gene

expression was partitioned similarly using the same model as above, and significance of each

term was tested using F-tests via a mixed-model ANOVA (Howell, 1997). Multiple testing was

controlled using a False-Discovery Rate (FDR) of 0.001. Broad-sense heritabilities were

estimated as $H^2 = \sigma^2_G / (\sigma^2_G + \sigma^2_{GE}/e + \sigma^2_E/re)$, with σ^2_G, σ^2_E and σ^2_{GE} as the variance explained

due to G, E, and G×E; e and r represent the number of environments and number of replicates

per environment, respectively. Inter-varietal differences in gene expression was estimated for

each environmental condition as the population-wide variance between accession mean

expression levels.

Univariate and multivariate selection analyses. Univariate selection differentials consider each trait separately and represent the total strength of selection acting on a trait (Lande, 1979; Lande and Arnold, 1983). Fitness was estimated as fecundity, which was the total number of filled rice grains per individual. Unstandardized linear selection differentials (S) on gene expression were estimated as coefficients of linear regression with fecundity as the dependent variable in each regression and transcript abundance and block as the independent variables. Unstandardized quadratic selection differentials (C) were estimated similarly as twice the coefficients of quadratic regression for transcript abundance and fecundity (Conner and Hartl, 2004; Stinchcombe et al., 2008). Using these we then estimated the variance-standardized, and mean-standardized selection differentials (Hereford et al., 2004; Lande and Arnold, 1983). Data preparation included filtering out individuals with zero fecundity followed by normalizing fecundity fitness by mean fitness. Further, transcript abundances were standardized by subtracting population mean abundance of the transcript and dividing by the standard deviation (SD) of the same transcript. To satisfy normality and remove noise inherent in expression data, transcripts with expression values more than 3 standard deviations from the mean were removed, which affected fewer than 1% of individuals. Selection differentials were estimated for transcripts that were expressed in at least 20 individuals and were estimated separately in each of the two environments. Multiple testing was controlled using Bonferroni correction (Bland and Altman, 1995).

  Multivariate selection gradients represent the strength of direct selection acting on each trait, after removing indirect selection caused by correlations with other traits (Lande and Arnold, 1983). Principal component analysis (PCA) was performed on transcript abundance using the prcomp function in R (Computing, 2013; Kassambara, 2017) and PCs explaining over 0.5% of variance in each environment were chosen for multivariate selection analyses. Linear (β) and quadratic (γ) selection gradients were estimated as coefficients of multiple regression with the normalized fecundity fitness as the dependent variable and the PCs as the independent variables (Conner and Hartl, 2004). We then chose the top one percent of transcripts showing the highest loadings of the PCs (n=182) and counted the number of transcripts showing evidence for positive selection (same directionality between loading of the transcript on the PC and univariate selection differential (S) acting on the transcript estimated above) versus negative selection (opposite directionality between loading of the transcript on the PC and univariate selection

differential (*S*) acting on the transcript estimated above) and based on the majority assigned a

directionality to the selection gradients on PCs. Variance-standardized multivariate selection

gradients were estimated for functional traits without strong correlation to avoid collinearity

among traits (Lande and Arnold, 1983; (Presotto et al., 2019), using a Pearson correlation

coefficient < 0.6 as threshold, which was estimated using the cor function in R (Computing,

2013).

Selection analyses on gene ontology biological processes. Gene Ontology (GO) term

annotations for rice genes/transcripts were downloaded from Monocots PLAZA 5.0 (Van Bel et

al., 2022). All fourth-level biological-process terms were downloaded using GO.db v3.15.0

(Carlson, 2022) and only these terms were considered for further analyses. Next, terms with

fewer than 20 transcripts in our dataset were filtered out to minimize redundancy, leaving a total

of 670 terms and 10,235 associated transcripts. The selection strength on a biological-process

term was estimated as the median selection strength of all transcripts annotated with that term. A

term was considered to be under significantly stronger selection compared to the transcriptome-

wide median if the median strength of selection for a term was over the transcriptome-wide

median selection strength by at least the 95% confidence interval for the selection strength of

that term. GO enrichments were done using ShinyGO (Ge et al., 2020).

Regulatory decoherence analyses. To examine regulatory decoherence in rice, a recently

developed method, CILP (Correlation by Individual Level Product), was used (Lea et al., 2019).

Since CILP calculates product correlations for all possible pairs of genes, only transcripts with a

selection strength greater than 0.1 ($|S| > 0.1$) in at least one environment with expression greater

than 0 in at least 50% individuals were included to reduce the dimensionality (leaving 2,318

transcripts). Multiple testing was controlled using a False-Discovery Rate (FDR) of 0.05.

Genotype data and SNP calling. Raw FASTQ files were downloaded from the Sequence Read

Archive (SRA) website under BioProject PRJEB6180 (Wang et al., 2018) and under bioprojects

PRJNA422249 and PRJNA557122 for 92 accessions (Gutaker et al., 2020) (Supplementary

Table 12). Further, genomes of 19 accessions were re-sequenced, and submitted to the SRA

(BioProject PRJNA1012700), leading to a total of 125 accessions for which genomic data was

available (Supplementary Table 12).

Raw reads were processed for quality control and adapter trimming using the bbduk

program of BBTools version 37.66 (https://jgi.doe.gov/data-and-tools/bbtools/) using the

options: minlen = 25 qtrim = rl trimq = 10 ktrim = r k = 25 mink = 11 hdist = 1 tpe tbo. The

output from this program was mapped to the reference genome *O. sativa* Nipponbare IRGSP 1.0

genome that was downloaded from NCBI Genome

(https://www.ncbi.nlm.nih.gov/genome/?term=txid4530[orgn]*)* using bwa-mem2 v2.1

(Vasimuddin et al., 2019). PCR duplicates were marked and removed using the Picard tools

version 2.9.0. SNPs were called using GATK HaplotypeCaller v4.2.0.0 to obtain a multi-

accession joint SNP file. Only SNPs that were above 5bp distance from an indel variant were

taken. Next, SNPs were filtered using the recommended GATK hard filtering (Van der Auwera

et al., 2013). Further, using vcftools v0.1.16 (Danecek et al., 2011), SNPs with at least 80%

genotype calls and a minor allele frequency of 0.05 were retained (--max-missing 0.8 --maf

0.05). Since rice is a inbred species, we also removed any SNPs that displayed heterozygosity of

over 5% identified using vcftools v0.1.16 –hardy (Danecek et al., 2011). Next, missing genotype

calls were imputed and phased using Beagle v4.1 (Browning and Browning, 2016), and using

vcftools v0.1.16 -m2 -M2 (Danecek et al., 2011) only biallelic SNPs were retained for further

analyses. Finally, SNPs were randomly pruned such that one SNP per 1000bp was retained using

vcftools v0.1.16 –thin (Danecek et al., 2011), leaving a SNP dataset of 246,714 markers.

G-matrix estimation and prediction of short-term phenotypic evolution. A G-matrix (G)

representing the additive genetic variance and covariance was estimated for the principal

component axes (PCs) by taking the eigengene. This was done by deploying GREML v1.94

(Yang et al., 2011). Although the principal components are by definition uncorrelated at the level

of the individual replicate plants, they start showing genetic covariances when loading values of

replicates from each genotype are averaged. Next, using the multivariate breeder's equation (Δz

$= G \beta$), we predicted the response to short-term phenotypic selection (Δz) on the PCs.

Association mapping. Association mapping was performed between the SNP markers and gene

expression values recorded in the normal and saline environments. For this, the linear model in

Matrix eQTL was used (Shabalin, 2012). The normalized gene expression values were averaged

over the replicates in each environment separately and these averages were subsequently used to

test for associations. The first five principal components (PCs) of the kinship matrix were

estimated using GAPIT v3 (Wang and Zhang, 2021) and added as covariates to control for

population structure. Associations were considered significant at a false-discovery rate (FDR) <

0.001, and when significant were included in downstream analyses. Due to long stretches of

homozygosity attributed to the highly inbred nature of rice, *trans*-eQTLs were defined as being

on a different chromosome or at least 1 Mb away from a gene under its influence on the same

chromosome; *cis*-eQTLs were defined as <100 kb away from an associated gene.

To identify significant G × eQTLs, we ran Matrix eQTL on the difference of expression in the

normal and saline conditions (normal – saline) at FDR 0.05 (Smith and Kruglyak, 2008).

For each gene regulated by a SNP in *cis* or *trans*, a lead SNP was identified as the SNP

with the most significant association within a 100-kb region. Furthermore, *trans*-eQTL hotspots

were identified through analyzing the number of unique genes regulated by lead SNPs in a given

100-kb region. To detect genes under *cis-trans* compensation or reinforcement, the effect sizes of

all lead *trans*-eQTLs were averaged for each gene and compared to the lead *cis*-eQTL for the

same gene. Next, for these genes we estimated the mean proportion of individuals with opposite

and same direction *cis-trans* allelic configuration. Genes were defined as compensating and

reinforcing if they had at least 60% of individuals with opposite and same *cis-trans* allelic

configuration, respectively. To examine the LD structure, we estimated r^2 using plink v1.9

(Purcell et al., 2007) between (1) all pairs of *cis-* and *trans*-variants for the identified

compensating and reinforcing genes and (2) 1,000 datasets of randomly selected SNP pairs (with

equal numbers of variant pairs and a similar distribution of distances between the *cis-* and *trans-*

variants as in 1). Next, we compared r^2 between (1) and (2) using a two-tailed permutation test.

To examine the patterns of past selection on eQTLs, we used the minor allele frequency

(MAF) of the 246,714 SNP markers and compared these using the t.test function in R

(Computing, 2013). Furthermore, we estimated the site-wise nucleotide diversity (π) and

averaged it over 50-kb flanking regions around each *cis*-eQTLs (100-kb region total). We

compared this π to the background nucleotide diversity, estimated as π averaged over 100-kb

blocks throughout the genome minus the 100-kb *cis*-eQTL region above. The difference in mean

of nucleotide diversity was tested using the t.test function in R (Computing, 2013).

ACKNOWLEDGEMENTS

We thank the New York University Center for Genomics and Systems Biology GenCore Facility for sequencing support, and the New York University High Performance Computing for supplying computational resources. We would also like to thank William Mauck and Nicholas Rogers for technical support, as well as Adrian E. Platts and Jae Young Choi, for their help with post-sequencing data processing. We are grateful to current members of the Purugganan laboratory (particularly J. Flowers, A. Kurbidaeva, O. Alam) and Elena Hamann at Fordham University (currently Heinrich Heine University Düsseldorf) for insightful discussions. S.C.G. is supported by a grant from the Gordon and Betty Moore Foundation/Life Sciences Research Foundation (award no. GBMF2550.06 to S.C.G.), startup funds from the University of California Riverside, and a grant from the National Institute of General Medical Sciences of the National Institutes of Health (award no. R35GM151194 to S.C.G.). This work was funded in part by grants from the US National Science Foundation Plant Genome Research Program (IOS 1546218 and 2204374) and the Zegar Family Foundation.

AUTHOR CONTRIBUTIONS

M.D.P. conceived and directed the project; Z.J.-L., R.K.S., G.V.V., A.G.C.S., K.L.M. designed and coordinated field experiments; Z.J.-L., S.C.G., M.A.N., A.G.C.S. performed field experiments; Z.J.-L., M.L.Z., S.C.G. processed samples and extracted RNA for sequencing; Z.J.-L. and R.S. made RNA-seq libraries; M.L.Z., S.C.G. made whole-genome re-sequencing libraries; I.C., S.C.G., Z.J.-L., S.J.F., K.L.M. designed and ran the bioinformatics workflow for RNA-seq; S.G. performed selection, statistical and genetic-marker-based analyses with assistance from S.J.F. and M.D.P.; S.C.G., Z.J.-L., S.G., M.A.N. processed fitness, higher-level-trait and gene expression data; and S.G., Z.J.-L. and M.D.P. wrote the manuscript. All authors contributed towards the editing and final review of the manuscript.

TABLES

Table 1: Summary statistics of selection on gene expression.

		Control	Salt
Median $|S|$		0.0501	0.0507
$S > 0$	# Transcripts	9654	8885
	Median S	0.053	0.051
$S < 0$	# Transcripts	8415	9133
	Median S	-0.047	-0.050
$C > 0$	# Transcripts	7175	8713
	Median C	0.077	0.115
$C < 0$	# Transcripts	10894	9304
	Median C	-0.096	-0.111

844   **Figures**

846
Figure 1: **The strength and pattern of selection on heritable gene expression**. a, The *O.*

sativa ssp. *indica* populations showed higher average fitness in the normal (blue) and saline

(pink) field (two-tailed paired t-test $P = 1.658 \times 10^{-8}$) and fitness further showed a significant

effects of genotype (G) and environment (E); genotype × environment (G×E) was not

significant. Analysis of variance (ANOVA) [G and E ($P < 0.001$), G×E ($P = 0.49$)]; n=130

accessions. b, Broad-sense heritability (H^2) distribution of *Oryza sativa* spp. *indica* transcripts.

Two-way ANOVA, genotype FDR-adjusted q < 0.001, n = 130 accessions. c-e, The strength of

linear selection |*S*|, linear selection differentials (*S*), and quadratic selection differentials (*C*) for

genome-wide gene expression in normal (blue) and saline (pink) conditions. f, Conditionally

neutral (light gray), and antagonistically pleiotropic transcripts (blue and magenta represent

beneficial expression in normal and saline conditions, respectively). Black represents transcripts

experiencing selection in the same direction in both environments (expression is beneficial or

detrimental in both environments).

Figure 2: Biological processes and pathways with differential responses to selection under saline conditions. a, GO biological processes under stronger selection in normal (blue) and saline conditions (pink). b, Linear selection gradients *(β)*, along with direct (D), indirect (I) and total (T) responses to selection on suites of transcripts in normal (blue) and saline conditions (pink).

Figure 3: Salinity stress induces regulatory decoherence. a, Pearson correlation coefficients between pairs of transcripts ($|S| > 0.1$ and expression greater than 0 in at least 50% individuals) in normal (x-axis) and saline conditions (y-axis). Pink and blue represent pairs with correlation stronger in saline and normal conditions, respectively; gray represents correlation that is not significantly different between conditions. b, Enrichment of transcripts with significant pairs greater than the median (median significant pair per transcript = 12, n = 853) involved in regulatory decoherence post salt exposure.

902 **Figure 4: Traits with different selection profiles under salt stress.** Linear *(β)* and quadratic *(γ)*

903 selection gradients on the traits LOP (leaf osmotic potential), Chl_a (chlorophyll a content), and

904 FT (flowering time). Error bars represent mean ± SE; dots and asterisks indicate significance of

905 selection-gradient at two-sided unadjusted $P < 0.1$ and $P < 0.05$.

Figure 5: Genetic architecture of gene expression variation during salt stress. a, Effect sizes of genes with both *cis* and *trans* factors under saline conditions showing excess of reinforcing cis-trans (teal) in comparison to compensating cis-trans (salmon). b, Inter-varietal variation in gene expression for genes under compensating control is significantly lower than for those under reinforcing control; one-sided Mann-Whitney $P = 0.0028$. c, Frequency distribution of MAF (minor allele frequency) for *cis*-eQTLs (blue) and *trans*-eQTLs (yellow) in saline conditions against the genome-wide background (gray).

References

Agrawal AF, Whitlock MC. 2010. Environmental duress and epistasis: how does stress affect the strength
of selection on new mutations? *Trends Ecol Evol* **25**:450–458. doi:10.1016/j.tree.2010.05.003

Ahmad F, Debes PV, Nousiainen I, Kahar S, Pukk L, Gross R, Ozerov M, Vasemägi A. 2021. The
strength and form of natural selection on transcript abundance in the wild. *Mol Ecol* **30**:2724–2737.
doi:10.1111/mec.15743

Amiri A, Coppola G, Scuderi S, Wu F, Roychowdhury T, Liu F, Pochareddy S, Shin Y, Safi A, Song L,
Zhu Y, Sousa AMM, PsychENCODE Consortium, Gerstein M, Crawford GE, Sestan N, Abyzov A,
Vaccarino FM. 2018. Transcriptome and epigenome landscape of human cortical development
modeled in organoids. *Science* **362**. doi:10.1126/science.aat6720

Anderson JT, Lee C-R, Rushworth CA, Colautti RI, Mitchell-Olds T. 2013. Genetic trade-offs and
conditional neutrality contribute to local adaptation. *Mol Ecol* **22**:699–708. doi:10.1111/j.1365-
294X.2012.05522.x

Anderson JT, Willis JH, Mitchell-Olds T. 2011. Evolutionary genetics of plant adaptation. *Trends Genet*
27:258–266. doi:10.1016/j.tig.2011.04.001

Arbuthnott D, Whitlock MC. 2018. Environmental stress does not increase the mean strength of selection.
J Evol Biol **31**:1599–1606. doi:10.1111/jeb.13351

Ashraf MY, Bhatti AS. 2000. Effect of salinity on growth and chlorophyll content in rice. *Pakistan*
journal of scientific and industrial research **43**:130–132.

Bland JM, Altman DG. 1995. Multiple significance tests: the Bonferroni method. *BMJ* **310**:170.
doi:10.1136/bmj.310.6973.170

Browning BL, Browning SR. 2016. Genotype Imputation with Millions of Reference Samples. *Am J Hum*
Genet **98**:116–126. doi:10.1016/j.ajhg.2015.11.020

Ćalić I, Groen SC, Choi JY, Joly-Lopez Z, Hamann E, Natividad MA, Dorph K, Cabral CLU, Torres RO,
Vergara GV, Henry A, Purugganan MD, Franks SJ. 2022. The influence of genetic architecture on
responses to selection under drought in rice. *Evol Appl* **15**:1670–1690. doi:10.1111/eva.13419

Carillo P, Grazia M, Pontecorvo G, Fuggi A, Woodrow P. 2011. Salinity stress and salt tolerance In:
Shanker A, Venkateswarlu B, editors. Abiotic Stress in Plants - Mechanisms and Adaptations.
London, England: InTech. doi:10.5772/22331

Carlson M. 2022. GO. db: A set of annotation maps describing the entire Gene Ontology.

Castillo EG, Tuong TP, Ismail AM, Inubushi K. 2007. Response to Salinity in Rice: Comparative Effects
of Osmotic and Ionic Stresses. *Plant Prod Sci* **10**:159–170. doi:10.1626/pps.10.159

Chang J, Cheong BE, Natera S, Roessner U. 2019. Morphological and metabolic responses to salt stress

of rice (Oryza sativa L.) cultivars which differ in salinity tolerance. *Plant Physiol Biochem* **144**:427–

435. doi:10.1016/j.plaphy.2019.10.017

Che-Othman MH, Millar AH, Taylor NL. 2017. Connecting salt stress signalling pathways with salinity-

induced changes in mitochondrial metabolic processes in C3 plants. *Plant Cell Environ* **40**:2875–

2905. doi:10.1111/pce.13034

Computing RFS. 2013. R: A language and environment for statistical computing. *Vienna: R Core Team*.

Conner JK, Hartl DL. 2004. A primer of ecological genetics. New York, NY: Oxford University Press.

Coolon JD, McManus CJ, Stevenson KR, Graveley BR, Wittkopp PJ. 2014. Tempo and mode of

regulatory evolution in Drosophila. *Genome Res* **24**:797–808. doi:10.1101/gr.163014.113

Cramer GR, Urano K, Delrot S, Pezzotti M, Shinozaki K. 2011. Effects of abiotic stress on plants: a

systems biology perspective. *BMC Plant Biol* **11**:163. doi:10.1186/1471-2229-11-163

Cui Y, Lu S, Li Z, Cheng J, Hu P, Zhu T, Wang X, Jin M, Wang X, Li L, Huang S, Zou B, Hua J. 2020.

CYCLIC NUCLEOTIDE-GATED ION CHANNELs 14 and 16 Promote Tolerance to Heat and

Chilling in Rice. *Plant Physiol* **183**:1794–1808. doi:10.1104/pp.20.00591

Danecek P, Auton A, Abecasis G, Albers CA, Banks E, DePristo MA, Handsaker RE, Lunter G, Marth

GT, Sherry ST, McVean G, Durbin R, 1000 Genomes Project Analysis Group. 2011. The variant call

format and VCFtools. *Bioinformatics* **27**:2156–2158. doi:10.1093/bioinformatics/btr330

De Clercq I, Van de Velde J, Luo X, Liu L, Storme V, Van Bel M, Pottie R, Vaneechoutte D, Van

Breusegem F, Vandepoele K. 2021. Integrative inference of transcriptional networks in Arabidopsis

yields novel ROS signalling regulators. *Nat Plants* **7**:500–513. doi:10.1038/s41477-021-00894-1

Dobin A, Davis CA, Schlesinger F, Drenkow J, Zaleski C, Jha S, Batut P, Chaisson M, Gingeras TR.

2013. STAR: ultrafast universal RNA-seq aligner. *Bioinformatics* **29**:15–21.

doi:10.1093/bioinformatics/bts635

Emerson JJ, Hsieh L-C, Sung H-M, Wang T-Y, Huang C-J, Lu HH-S, Lu M-YJ, Wu S-H, Li W-H. 2010.

Natural selection on cis and trans regulation in yeasts. *Genome Res* **20**:826–836.

doi:10.1101/gr.101576.109

Fahad S, Bajwa AA, Nazir U, Anjum SA, Farooq A, Zohaib A, Sadia S, Nasim W, Adkins S, Saud S,

Ihsan MZ, Alharby H, Wu C, Wang D, Huang J. 2017. Crop Production under Drought and Heat

Stress: Plant Responses and Management Options. *Front Plant Sci* **8**:1147.

doi:10.3389/fpls.2017.01147

Falconer DS, Mackay TFC. 1996. Introduction to quantitative genetics. Essex, England: Prentice Hall.

Francini A, Sebastiani L. 2019. Abiotic Stress Effects on Performance of Horticultural Crops.

Horticulturae **5**:67. doi:10.3390/horticulturae5040067

Garris AJ, Tai TH, Coburn J, Kresovich S, McCouch S. 2005. Genetic structure and diversity in Oryza

sativa L. *Genetics* **169**:1631–1638. doi:10.1534/genetics.104.035642

Ge SX, Jung D, Yao R. 2020. ShinyGO: a graphical gene-set enrichment tool for animals and plants.

Bioinformatics **36**:2628–2629. doi:10.1093/bioinformatics/btz931

Goncalves A, Leigh-Brown S, Thybert D, Stefflova K, Turro E, Flicek P, Brazma A, Odom DT, Marioni

JC. 2012. Extensive compensatory cis-trans regulation in the evolution of mouse gene expression.

Genome Res **22**:2376–2384. doi:10.1101/gr.142281.112

Groen SC, Ćalić I, Joly-Lopez Z, Platts AE, Choi JY, Natividad M, Dorph K, Mauck WM, Bracken B,

Cabral CLU, Kumar A, Torres RO, Satija R, Vergara G, Henry A, Franks SJ, Purugganan MD.

2020. The strength and pattern of natural selection on gene expression in rice. *Nature* **578**:572–576.

doi:10.1038/s41586-020-1997-2

Groen SC, Hamann E, Ćalić I, Cochran C, Konshok R, Purugganan MD, Franks SJ. 2021. Fitness costs

and benefits of gene expression plasticity in rice under drought. *bioRxiv*.

doi:10.1101/2021.03.16.435597

Gutaker RM, Groen SC, Bellis ES, Choi JY, Pires IS, Bocinsky RK, Slayton ER, Wilkins O, Castillo CC,

Negrão S, Oliveira MM, Fuller DQ, Guedes JAD, Lasky JR, Purugganan MD. 2020. Genomic

history and ecology of the geographic spread of rice. *Nat Plants* **6**:492–502. doi:10.1038/s41477-

020-0659-6

Hakim MA, Juraimi AS, Hanafi MM, Ismail MR, Selamat A, Rafii MY, Latif MA. 2014. Biochemical

and anatomical changes and yield reduction in rice (Oryza sativa L.) under varied salinity regimes.

Biomed Res Int **2014**:208584. doi:10.1155/2014/208584

Hammond JP, Mayes S, Bowen HC, Graham NS, Hayden RM, Love CG, Spracklen WP, Wang J,

Welham SJ, White PJ, King GJ, Broadley MR. 2011. Regulatory hotspots are associated with plant

gene expression under varying soil phosphorus supply in Brassica rapa. *Plant Physiol* **156**:1230–

1241. doi:10.1104/pp.111.175612

Hereford J, Hansen TF, Houle D. 2004. Comparing strengths of directional selection: how strong is

strong? *Evolution* **58**:2133–2143. doi:10.1111/j.0014-3820.2004.tb01592.x

Hernandez RD, Uricchio LH, Hartman K, Ye C, Dahl A, Zaitlen N. 2019. Ultrarare variants drive

substantial cis heritability of human gene expression. *Nat Genet* **51**:1349–1355. doi:10.1038/s41588-

019-0487-7

Hoekstra HE, Hoekstra JM, Berrigan D, Vignieri SN, Hoang A, Hill CE, Beerli P, Kingsolver JG. 2001.

Strength and tempo of directional selection in the wild. *Proc Natl Acad Sci U S A* **98**:9157–9160.

doi:10.1073/pnas.161281098

Howell DC. 1997. Statistical Methods for Psychology. Cengage Learning.

Hussain S, Zhang J-H, Zhong C, Zhu L-F, Cao X-C, Yu S-M, Allen Bohr J, Hu J-J, Jin Q-Y. 2017.

Effects of salt stress on rice growth, development characteristics, and the regulating ways: A review.

J Integr Agric **16**:2357–2374. doi:10.1016/S2095-3119(16)61608-8

Israel JW, Martik ML, Byrne M, Raff EC, Raff RA, McClay DR, Wray GA. 2016. Comparative

Developmental Transcriptomics Reveals Rewiring of a Highly Conserved Gene Regulatory Network

during a Major Life History Switch in the Sea Urchin Genus Heliocidaris. *PLoS Biol* **14**:e1002391.

doi:10.1371/journal.pbio.1002391

Jacoby RP, Taylor NL, Millar AH. 2011. The role of mitochondrial respiration in salinity tolerance.

Trends Plant Sci **16**:614–623. doi:10.1016/j.tplants.2011.08.002

Jasnos L, Tomala K, Paczesniak D, Korona R. 2008. Interactions between stressful environment and gene

deletions alleviate the expected average loss of fitness in yeast. *Genetics* **178**:2105–2111.

doi:10.1534/genetics.107.084533

Jiang Z, Zhou X, Tao M, Yuan F, Liu L, Wu F, Wu X, Xiang Y, Niu Y, Liu F, Li C, Ye R, Byeon B, Xue

Y, Zhao H, Wang H-N, Crawford BM, Johnson DM, Hu C, Pei C, Zhou W, Swift GB, Zhang H, Vo-

Dinh T, Hu Z, Siedow JN, Pei Z-M. 2019. Plant cell-surface GIPC sphingolipids sense salt to trigger

Ca2+ influx. *Nature* **572**:341–346. doi:10.1038/s41586-019-1449-z

Joly-Lopez Z, Platts AE, Gulko B, Choi JY, Groen SC, Zhong X, Siepel A, Purugganan MD. 2020. An

inferred fitness consequence map of the rice genome. *Nat Plants* **6**:119–130. doi:10.1038/s41477-

019-0589-3

Josephs EB, Lee YW, Stinchcombe JR, Wright SI. 2015. Association mapping reveals the role of

purifying selection in the maintenance of genomic variation in gene expression. *Proceedings of the*

National Academy of Sciences **112**:15390–15395. doi:10.1073/pnas.1503027112

Kassambara A. 2017. Practical Guide To Principal Component Methods in R: PCA, M(CA), FAMD,

MFA, HCPC, factoextra. STHDA.

Kim W-Y, Ali Z, Park HJ, Park SJ, Cha J-Y, Perez-Hormaeche J, Quintero FJ, Shin G, Kim MR, Qiang

Z, Ning L, Park HC, Lee SY, Bressan RA, Pardo JM, Bohnert HJ, Yun D-J. 2013. Release of SOS2

kinase from sequestration with GIGANTEA determines salt tolerance in Arabidopsis. *Nat Commun*

4:1352. doi:10.1038/ncomms2357

Kingsolver JG, Hoekstra HE, Hoekstra JM, Berrigan D, Vignieri SN, Hill CE, Hoang A, Gibert P, Beerli

P. 2001. The strength of phenotypic selection in natural populations. *Am Nat* **157**:245–261.

doi:10.1086/319193

Kingsolver JG, Pfennig DW. 2007. Patterns and Power of Phenotypic Selection in Nature. *Bioscience*

57:561–572. doi:10.1641/B570706

Kliebenstein D. 2009. Quantitative genomics: analyzing intraspecific variation using global gene

expression polymorphisms or eQTLs. *Annu Rev Plant Biol* **60**:93–114.

1060   doi:10.1146/annurev.arplant.043008.092114

1061 Ko DK, Brandizzi F. 2020. Network-based approaches for understanding gene regulation and function in
1062   plants. *Plant J* **104**:302–317. doi:10.1111/tpj.14940

1063 Kondrashov AS, Houle D. 1997. Genotype—environment interactions and the estimation of the genomic
1064   mutation rate in Drosophila melanogaster. *Proceedings of the Royal Society of London Series B:*
1065   *Biological Sciences* **258**:221–227. doi:10.1098/rspb.1994.0166

1066 Kopecká R, Kameniarová M, Černý M, Brzobohatý B, Novák J. 2023. Abiotic Stress in Crop Production.
1067   *Int J Mol Sci* **24**. doi:10.3390/ijms24076603

1068 Krishnamurthy R, Bhagwat KA. 1990. Accumulation of choline and glycinebetaine in salt-stressed wheat
1069   seedlings. *Curr Sci* **59**:111–112.

1070 Kumar SV, Wigge PA. 2010. H2A.Z-containing nucleosomes mediate the thermosensory response in
1071   Arabidopsis. *Cell* **140**:136–147. doi:10.1016/j.cell.2009.11.006

1072 Kuroha T, Nagai K, Kurokawa Y, Nagamura Y, Kusano M, Yasui H, Ashikari M, Fukushima A. 2017.
1073   eQTLs Regulating Transcript Variations Associated with Rapid Internode Elongation in Deepwater
1074   Rice. *Front Plant Sci* **8**:1753. doi:10.3389/fpls.2017.01753

1075 Lande R. 1979. Quantitative Genetic Analysis of Multivariate Evolution, Applied to Brain: Body Size
1076   Allometry. *Evolution* **33**:402–416. doi:10.2307/2407630

1077 Lande R, Arnold SJ. 1983. The measurement of selection correlated characters. *Evolution* **37**:1210–1226.
1078   doi:10.1111/j.1558-5646.1983.tb00236.x

1079 Landry CR, Wittkopp PJ, Taubes CH, Ranz JM, Clark AG, Hartl DL. 2005. Compensatory cis-trans
1080   evolution and the dysregulation of gene expression in interspecific hybrids of Drosophila. *Genetics*
1081   **171**:1813–1822. doi:10.1534/genetics.105.047449

1082 Laohavisit A, Richards SL, Shabala L, Chen C, Colaço RDDR, Swarbreck SM, Shaw E, Dark A, Shabala
1083   S, Shang Z, Davies JM. 2013. Salinity-induced calcium signaling and root adaptation in Arabidopsis
1084   require the calcium regulatory protein annexin1. *Plant Physiol* **163**:253–262.
1085   doi:10.1104/pp.113.217810

1086 Lea A, Subramaniam M, Ko A, Lehtimäki T, Raitoharju E, Kähönen M, Seppälä I, Mononen N, Raitakari
1087   OT, Ala-Korpela M, Pajukanta P, Zaitlen N, Ayroles JF. 2019. Genetic and environmental
1088   perturbations lead to regulatory decoherence. *Elife* **8**. doi:10.7554/eLife.40538

1089 Lee SS, Park HJ, Jung WY, Lee A, Yoon DH, You YN, Kim H-S, Kim B-G, Ahn JC, Cho HS. 2015.
1090   OsCYP21-4, a novel Golgi-resident cyclophilin, increases oxidative stress tolerance in rice. *Front*
1091   *Plant Sci* **6**:797. doi:10.3389/fpls.2015.00797

1092 Liang L, Zhang Z, Cheng N, Liu H, Song S, Hu Y, Zhou X, Zhang J, Xing Y. 2021. The transcriptional
1093   repressor OsPRR73 links circadian clock and photoperiod pathway to control heading date in rice.

Plant Cell Environ **44**:842–855. doi:10.1111/pce.13987

Liang W, Ma X, Wan P, Liu L. 2018. Plant salt-tolerance mechanism: A review. *Biochem Biophys Res*
Commun **495**:286–291. doi:10.1016/j.bbrc.2017.11.043

Li C, Wong WH. 2001. Model-based analysis of oligonucleotide arrays: expression index computation
and outlier detection. *Proc Natl Acad Sci U S A* **98**:31–36. doi:10.1073/pnas.98.1.31

Li K, Wang Y, Han C, Zhang W, Jia H, Li X. 2007. GA signaling and CO/FT regulatory module mediate
salt-induced late flowering in Arabidopsis thaliana. *Plant Growth Regul* **53**:195–206.
doi:10.1007/s10725-007-9218-7

Liu C, Mao B, Yuan D, Chu C, Duan M. 2022. Salt tolerance in rice: Physiological responses and
molecular mechanisms. *The Crop Journal* **10**:13–25. doi:10.1016/j.cj.2021.02.010

Liu S, Li C, Wang H, Wang S, Yang S, Liu X, Yan J, Li B, Beatty M, Zastrow-Hayes G, Song S, Qin F.
2020. Mapping regulatory variants controlling gene expression in drought response and tolerance in
maize. *Genome Biol* **21**:163. doi:10.1186/s13059-020-02069-1

Lovell JT, Jenkins J, Lowry DB, Mamidi S, Sreedasyam A, Weng X, Barry K, Bonnette J, Campitelli B,
Daum C, Gordon SP, Gould BA, Khasanova A, Lipzen A, MacQueen A, Palacio-Mejía JD, Plott C,
Shakirov EV, Shu S, Yoshinaga Y, Zane M, Kudrna D, Talag JD, Rokhsar D, Grimwood J, Schmutz
J, Juenger TE. 2018. The genomic landscape of molecular responses to natural drought stress in
Panicum hallii. *Nat Commun* **9**:5213. doi:10.1038/s41467-018-07669-x

Macosko EZ, Basu A, Satija R, Nemesh J, Shekhar K, Goldman M, Tirosh I, Bialas AR, Kamitaki N,
Martersteck EM, Trombetta JJ, Weitz DA, Sanes JR, Shalek AK, Regev A, McCarroll SA. 2015.
Highly Parallel Genome-wide Expression Profiling of Individual Cells Using Nanoliter Droplets.
Cell **161**:1202–1214. doi:10.1016/j.cell.2015.05.002

Mareri L, Parrotta L, Cai G. 2022. Environmental Stress and Plants. *Int J Mol Sci* **23**.
doi:10.3390/ijms23105416

Matsumura S, Arlinghaus R, Dieckmann U. 2012. Standardizing Selection Strengths to Study Selection in
the Wild: A Critical Comparison and Suggestions for the Future. *Bioscience* **62**:1039–1054.
doi:10.1525/bio.2012.62.12.6

Ma X, Qiao Z, Chen D, Yang W, Zhou R, Zhang W, Wang M. 2015. CYCLIN-DEPENDENT KINASE
G2 regulates salinity stress response and salt mediated flowering in Arabidopsis thaliana. *Plant Mol*
Biol **88**:287–299. doi:10.1007/s11103-015-0324-z

McManus CJ, May GE, Spealman P, Shteyman A. 2014. Ribosome profiling reveals post-transcriptional
buffering of divergent gene expression in yeast. *Genome Res* **24**:422–430.
doi:10.1101/gr.164996.113

Melino V, Tester M. 2023. Salt-Tolerant Crops: Time to Deliver. *Annu Rev Plant Biol* **74**:671–696.

doi:10.1146/annurev-arplant-061422-104322

Metzger BPH, Duveau F, Yuan DC, Tryban S, Yang B, Wittkopp PJ. 2016. Contrasting Frequencies and

Effects of cis- and trans-Regulatory Mutations Affecting Gene Expression. *Mol Biol Evol* **33**:1131–

1146. doi:10.1093/molbev/msw011

Meyer RS, Purugganan MD. 2013. Evolution of crop species: Genetics of domestication and

diversification. *Nature Reviews Genetics*. doi:10.1038/nrg3605

Miller G, Suzuki N, Ciftci-Yilmaz S, Mittler R. 2010. Reactive oxygen species homeostasis and

signalling during drought and salinity stresses. *Plant Cell Environ* **33**:453–467. doi:10.1111/j.1365-

3040.2009.02041.x

Munns R. 2002. Comparative physiology of salt and water stress. *Plant Cell Environ* **25**:239–250.

doi:10.1046/j.0016-8025.2001.00808.x

Munns R, Tester M. 2008. Mechanisms of salinity tolerance. *Annu Rev Plant Biol* **59**:651–681.

doi:10.1146/annurev.arplant.59.032607.092911

Nadir S, Khan S, Zhu Q, Henry D, Wei L, Lee DS, Chen L. 2018. An overview on reproductive isolation

in Oryza sativa complex. *AoB Plants* **10**:ly060. doi:10.1093/aobpla/ply060

Petrusa LM, Winicov I. 1997. Proline status in salt-tolerant and salt-sensitive alfalfa cell lines and plants

in response to NaCl. *Plant Physiol Biochem*.

Picelli S, Björklund ÅK, Faridani OR, Sagasser S, Winberg G, Sandberg R. 2013. Smart-seq2 for

sensitive full-length transcriptome profiling in single cells. *Nat Methods* **10**:1096–1098.

doi:10.1038/nmeth.2639

Ponce KS, Meng L, Guo L, Leng Y, Ye G. 2021. Advances in Sensing, Response and Regulation

Mechanism of Salt Tolerance in Rice. *Int J Mol Sci* **22**. doi:10.3390/ijms22052254

Presotto A, Hernández F, Mercer KL. 2019. Phenotypic selection under two contrasting environments in

wild sunflower and its crop-wild hybrid. *Evol Appl* **12**:1703–1717. doi:10.1111/eva.12828

Pu J, Yu H, Guo Y. 2022. A Novel Strategy to Identify Prognosis-Relevant Gene Sets in Cancers. *Genes*

13:862. doi:10.3390/genes13050862

Purcell S, Neale B, Todd-Brown K, Thomas L, Ferreira MAR, Bender D, Maller J, Sklar P, de Bakker

PIW, Daly MJ, Sham PC. 2007. PLINK: a tool set for whole-genome association and population-

based linkage analyses. *Am J Hum Genet* **81**:559–575. doi:10.1086/519795

Qin H, Huang R. 2020. The phytohormonal regulation of Na+/K+ and reactive oxygen species

homeostasis in rice salt response. *Mol Breed* **40**:47. doi:10.1007/s11032-020-1100-6

Qu W, Gurdziel K, Pique-Regi R, Ruden DM. 2018. Lead Modulates trans- and cis-Expression

Quantitative Trait Loci (eQTLs) in Drosophila melanogaster Heads. *Front Genet* **9**:395.

doi:10.3389/fgene.2018.00395

Qu X, Zou J, Wang J, Yang K, Wang X, Le J. 2022. A Rice R2R3-Type MYB Transcription Factor OsFLP Positively Regulates Drought Stress Response via OsNAC. *Int J Mol Sci* **23**. doi:10.3390/ijms23115873

Radanielson AM, Angeles O, Li T, Ismail AM, Gaydon DS. 2018. Describing the physiological responses of different rice genotypes to salt stress using sigmoid and piecewise linear functions. *Field Crops Res* **220**:46–56. doi:10.1016/j.fcr.2017.05.001

Robinson MD, McCarthy DJ, Smyth GK. 2010. edgeR: a Bioconductor package for differential expression analysis of digital gene expression data. *Bioinformatics* **26**:139–140. doi:10.1093/bioinformatics/btp616

Roy S, Mishra M, Kaur G, Singh S, Rawat N, Singh P, Singla-Pareek SL, Pareek A. 2022. OsCyp2-P, an auxin-responsive cyclophilin, regulates Ca2+ calmodulin interaction for an ion-mediated stress response in rice. *Physiol Plant* **174**:e13631. doi:10.1111/ppl.13631

Ruan S-L, Ma H-S, Wang S-H, Fu Y-P, Xin Y, Liu W-Z, Wang F, Tong J-X, Wang S-Z, Chen H-Z. 2011. Proteomic identification of OsCYP2, a rice cyclophilin that confers salt tolerance in rice (Oryza sativa L.) seedlings when overexpressed. *BMC Plant Biol* **11**:34. doi:10.1186/1471-2229-11-34

Ruffley M, Leventhal L, Hateley S, Rhee SY, Exposito-Alonso M. 2023. Conflicts in natural selection constrain adaptation to climate change in Arabidopsis thaliana. *bioRxiv*. doi:10.1101/2023.10.16.562583

Schuppler U, He PH, John PC, Munns R. 1998. Effect of water stress on cell division and cell-division-cycle 2-like cell-cycle kinase activity in wheat leaves. *Plant Physiol* **117**:667–678. doi:10.1104/pp.117.2.667

Shabalin AA. 2012. Matrix eQTL: ultra fast eQTL analysis via large matrix operations. *Bioinformatics* **28**:1353–1358. doi:10.1093/bioinformatics/bts163

Siddiqui MN, Léon J, Naz AA, Ballvora A. 2021. Genetics and genomics of root system variation in adaptation to drought stress in cereal crops. *J Exp Bot* **72**:1007–1019. doi:10.1093/jxb/eraa487

Signor SA, Nuzhdin SV. 2018. The Evolution of Gene Expression in cis and trans. *Trends Genet* **34**:532–544. doi:10.1016/j.tig.2018.03.007

Smith EN, Kruglyak L. 2008. Gene-environment interaction in yeast gene expression. *PLoS Biol* **6**:e83. doi:10.1371/journal.pbio.0060083

Snoek LB, Terpstra IR, Dekter R, Van den Ackerveken G, Peeters AJM. 2012. Genetical Genomics Reveals Large Scale Genotype-By-Environment Interactions in Arabidopsis thaliana. *Front Genet* **3**:317. doi:10.3389/fgene.2012.00317

Sterken MG, Nijveen H, van Zanten M, Jiménez-Gómez JM, Geshnizjani N, Willems LAJ, Rienstra J,

1196   Hilhorst HWM, Ligterink W, Snoek BL. 2023. Plasticity of maternal environment-dependent
1197   expression-QTLs of tomato seeds. *Theor Appl Genet* **136**:28. doi:10.1007/s00122-023-04322-0
1198 Stinchcombe JR, Agrawal AF, Hohenlohe PA, Arnold SJ, Blows MW. 2008. Estimating nonlinear
1199   selection gradients using quadratic regression coefficients: double or nothing? *Evolution* **62**:2435–
1200   2440. doi:10.1111/j.1558-5646.2008.00449.x
1201 Taïbi K, Taïbi F, Ait Abderrahim L, Ennajah A, Belkhodja M, Mulet JM. 2016. Effect of salt stress on
1202   growth, chlorophyll content, lipid peroxidation and antioxidant defence systems in Phaseolus
1203   vulgaris L. *S Afr J Bot* **105**:306–312. doi:10.1016/j.sajb.2016.03.011
1204 Tari I, Kiss G, Deér AK, Csiszár J, Erdei L, Gallé Á, Gémes K, Horváth F, Poór P, Szepesi Á, Simon
1205   LM. 2010. Salicylic acid increased aldose reductase activity and sorbitol accumulation in tomato
1206   plants under salt stress. *Biol Plant* **54**:677–683. doi:10.1007/s10535-010-0120-1
1207 Tsai Y-C, Chen K-C, Cheng T-S, Lee C, Lin S-H, Tung C-W. 2019. Chlorophyll fluorescence analysis in
1208   diverse rice varieties reveals the positive correlation between the seedlings salt tolerance and
1209   photosynthetic efficiency. *BMC Plant Biol* **19**:403. doi:10.1186/s12870-019-1983-8
1210 Van Bel M, Silvestri F, Weitz EM, Kreft L, Botzki A, Coppens F, Vandepoele K. 2022. PLAZA 5.0:
1211   extending the scope and power of comparative and functional genomics in plants. *Nucleic Acids Res*
1212   **50**:D1468–D1474. doi:10.1093/nar/gkab1024
1213 Van der Auwera GA, Carneiro MO, Hartl C, Poplin R, Del Angel G, Levy-Moonshine A, Jordan T,
1214   Shakir K, Roazen D, Thibault J, Banks E, Garimella KV, Altshuler D, Gabriel S, DePristo MA.
1215   2013. From FastQ data to high confidence variant calls: the Genome Analysis Toolkit best practices
1216   pipeline. *Curr Protoc Bioinformatics* **43**:11.10.1–11.10.33. doi:10.1002/0471250953.bi1110s43
1217 Van Zandt PA, Mopper S. 2002. Delayed and carryover effects of salinity on flowering in Iris hexagona
1218   (Iridaceae). *Am J Bot* **89**:1847–1851. doi:10.3732/ajb.89.11.1847
1219 Vasimuddin M, Misra S, Li H, Aluru S. 2019. Efficient Architecture-Aware Acceleration of BWA-MEM
1220   for Multicore Systems2019 IEEE International Parallel and Distributed Processing Symposium
1221   (IPDPS). pp. 314–324. doi:10.1109/IPDPS.2019.00041
1222 Wagner GP, Lynch VJ. 2008. The gene regulatory logic of transcription factor evolution. *Trends Ecol*
1223   *Evol* **23**:377–385. doi:10.1016/j.tree.2008.03.006
1224 Wang H, Zhang M, Guo R, Shi D, Liu B, Lin X, Yang C. 2012. Effects of salt stress on ion balance and
1225   nitrogen metabolism of old and young leaves in rice (Oryza sativa L.). *BMC Plant Biol* **12**:194.
1226   doi:10.1186/1471-2229-12-194
1227 Wang J, Zhang Z. 2021. GAPIT Version 3: Boosting Power and Accuracy for Genomic Association and
1228   Prediction. *Genomics Proteomics Bioinformatics* **19**:629–640. doi:10.1016/j.gpb.2021.08.005
1229 Wang W, Mauleon R, Hu Z, Chebotarov D, Tai S, Wu Z, Li M, Zheng T, Fuentes RR, Zhang F,

Mansueto L, Copetti D, Sanciangco M, Palis KC, Xu J, Sun C, Fu B, Zhang H, Gao Y, Zhao X,
Shen F, Cui X, Yu H, Li Z, Chen M, Detras J, Zhou Y, Zhang X, Zhao Y, Kudrna D, Wang C, Li R,
Jia B, Lu J, He X, Dong Z, Xu J, Li Y, Wang M, Shi J, Li J, Zhang D, Lee S, Hu W, Poliakov A,
Dubchak I, Ulat VJ, Borja FN, Mendoza JR, Ali J, Li J, Gao Q, Niu Y, Yue Z, Naredo MEB, Talag
J, Wang X, Li J, Fang X, Yin Y, Glaszmann J-C, Zhang J, Li J, Hamilton RS, Wing RA, Ruan J,
Zhang G, Wei C, Alexandrov N, McNally KL, Li Z, Leung H. 2018. Genomic variation in 3,010
diverse accessions of Asian cultivated rice. *Nature* **557**:43–49. doi:10.1038/s41586-018-0063-9

Watowich MM, Chiou KL, Montague MJ, Cayo Biobank Research Unit, Simons ND, Horvath JE, Ruiz-
Lambides AV, Martínez MI, Higham JP, Brent LJN, Platt ML, Snyder-Mackler N. 2022. Natural
disaster and immunological aging in a nonhuman primate. *Proc Natl Acad Sci U S A* **119**.
doi:10.1073/pnas.2121663119

Wei H, Wang X, He Y, Xu H, Wang L. 2021. Clock component OsPRR73 positively regulates rice salt
tolerance by modulating OsHKT2;1-mediated sodium homeostasis. *EMBO J* **40**:e105086.
doi:10.15252/embj.2020105086

West G, Inzé D, Beemster GTS. 2004. Cell cycle modulation in the response of the primary root of
Arabidopsis to salt stress. *Plant Physiol* **135**:1050–1058. doi:10.1104/pp.104.040022

West MAL, Kim K, Kliebenstein DJ, van Leeuwen H, Michelmore RW, Doerge RW, St Clair DA. 2007.
Global eQTL mapping reveals the complex genetic architecture of transcript-level variation in
Arabidopsis. *Genetics* **175**:1441–1450. doi:10.1534/genetics.106.064972

Wilkins O, Hafemeister C, Plessis A, Holloway-Phillips M-M, Pham GM, Nicotra AB, Gregorio GB,
Jagadish SVK, Septiningsih EM, Bonneau R, Purugganan M. 2016. EGRINs (Environmental Gene
Regulatory Influence Networks) in Rice That Function in the Response to Water Deficit, High
Temperature, and Agricultural Environments. *Plant Cell* **28**:2365–2384. doi:10.1105/tpc.16.00158

Wittkopp PJ, Haerum BK, Clark AG. 2008. Regulatory changes underlying expression differences within
and between Drosophila species. *Nat Genet* **40**:346–350. doi:10.1038/ng.77

Wu Z, Chen L, Yu Q, Zhou W, Gou X, Li J, Hou S. 2019. Multiple transcriptional factors control stomata
development in rice. *New Phytol* **223**:220–232. doi:10.1111/nph.15766

Xu X, Yuan L, Xie Q. 2022. The circadian clock ticks in plant stress responses. *Stress Biol* **2**:15.
doi:10.1007/s44154-022-00040-7

Yang J, Lee SH, Goddard ME, Visscher PM. 2011. GCTA: a tool for genome-wide complex trait
analysis. *Am J Hum Genet* **88**:76–82. doi:10.1016/j.ajhg.2010.11.011

Yang Y, Guo Y. 2018. Unraveling salt stress signaling in plants. *J Integr Plant Biol* **60**:796–804.
doi:10.1111/jipb.12689

Yuan F, Yang H, Xue Y, Kong D, Ye R, Li C, Zhang J, Theprungsirikul L, Shrift T, Krichilsky B,

Johnson DM, Swift GB, He Y, Siedow JN, Pei Z-M. 2014. OSCA1 mediates osmotic-stress-evoked

Ca2+ increases vital for osmosensing in Arabidopsis. *Nature* **514**:367–371. doi:10.1038/nature13593

Zhang H, Zhu J, Gong Z, Zhu J-K. 2022. Abiotic stress responses in plants. *Nat Rev Genet* **23**:104–119.

doi:10.1038/s41576-021-00413-0

Zhang R, Wang Y, Hussain S, Yang S, Li R, Liu S, Chen Y, Wei H, Dai Q, Hou H. 2022. Study on the

Effect of Salt Stress on Yield and Grain Quality Among Different Rice Varieties. *Front Plant Sci*

13:918460. doi:10.3389/fpls.2022.918460

1271

1272
1273