Comparative population genomics unveils congruent secondary suture zone in Southwest Pacific Hydrothermal Vents

AUTHORS :

Adrien Tran Lu Y[1,3]*, Stéphanie Ruault[2], Claire Daguin-Thiébaut[2], Anne-Sophie Le Port[2], Marion Ballenghien[2], Jade Castel[2], Pierre-Alexandre Gagnaire[1], Nicolas Bierne[1], Sophie Arnaud-Haond[3], Camille Poitrimol[2], Eric Thiébaut[2], François H Lallier[2], Thomas Broquet[2], Didier Jollivet[2], François Bonhomme[1] & Stéphane Hourdez[4]

Affiliations :

1 : ISEM, Univ Montpellier, CNRS, IRD, Montpellier, France

2 : Sorbonne Université, CNRS, Station Biologique de Roscoff, UMR 7144 Adaptation and diversity in the marine environment, Place Georges Teissier, 29 680 Roscoff, France

3 : MARBEC UMR 248, Université de Montpellier, Ifremer, IRD, CNRS; Avenue Jean Monnet CS 30171, 34203, Sète, France

4 : UMR 8222 LECOB CNRS-Sorbonne Université, Observatoire Océanologique de Banyuls, Avenue du Fontaulé, 66650, Banyuls-sur-mer, France

*Corresponding author : dr.adrien.tranluy@gmail.com

Abstract

How the interplay of biotic and abiotic factors shapes current genetic diversity at the community level remains an open question, particularly in the deep sea. Comparative phylogeography of multiple species can reveal the influence of past climatic events, geographic barriers, and species life history traits on spatial patterns of genetic structure across lineages.

Here, we conducted a comparative population genomic study on seven hydrothermal vent species co-distributed in the Back-arc Basins of the Southwest Pacific region. Using ddRAD-seq, we compared distribution range-wide genomic diversity patterns across species and discovered a shared zone of phylogeographic break. Although species exhibit congruent patterns of spatial structure, they also show variation in the degree of divergence among lineages across the suture zone. Additionally, two species have a sympatric contact zone in the Woodlark Basin.

Demogenetic inference revealed shared histories of lineage divergence and secondary contact. Low levels of asymmetric gene flow probably occurred in most species between the Woodlark and North Fiji basins, but the exact location of contact zones varied from species to species. Results show that gene flow is heterogeneous across the genome, indicating possible partial reproductive isolation between lineages and early speciation.

Our comparative study sheds light on the factors that shape genetic variation at the community level, and our findings enrich our understanding of deep-sea biogeography patterns. Emphasizing the pivotal role of historical and contemporary factors, our research underscores the necessity for a holistic approach, and in particular filling in the gaps regarding life history traits of deep-sea species (generation time, development type, duration of the larval phase).

Introduction

Hydrothermal vents are one of the most emblematic chemosynthesis-based ecosystems that host a highly specialized fauna. This vent fauna depends on local hydrothermal activity and is likely to share historical patterns of colonization linked to the tectonic history of the ridge system (Plouviez et al., 2009, Matabos et al., 2011, Matabos & Jollivet., 2019). In contrast to other deep-sea ecosystems, vents represent a linear but highly fragmented and relatively unstable ecosystem based on chemosynthetic primary producers, which cannot thrive elsewhere. Hydrothermal activity is linked to specific geological features associated with the volcanic and tectonic activities of ocean ridges or submarine volcanoes (Hourdez and Jollivet, 2020). Plate tectonics has previously been cited as a driver of most of the biogeographic distribution of the vent fauna (Tunnicliffe, 1991), that may lead to allopatric speciation with possible secondary contacts (Hurtado et al. 2004, Johnson et al. 2006, 2013, Faure et al. 2009, Plouviez et al., 2009, Matabos et al. 2011). On the East Pacific Rise, these previous studies pointed toward the emergence of a transition zone between biogeographic provinces initially separated.

Unlike other hydrothermal ecosystems, the fauna of the vents of the South-West Pacific is distributed across several geological Back-Arc-Basins (BABs) separated by abyssal plains, ridges and volcanic arcs, forming a fragmented and discontinuous complex. These BABs formations are estimated to be between 12 to 1 million years (My) old (Schellart et al., 2006). A Previous study highlighted contrasting phylogeographic patterns for few species, including some closely related species (Poitrimol et al., 2022). This suggested alternative dispersal strategies and evolutionary history to cope with fragmentation despite a common geological history of the vent habitat in this region. This geological dynamics provided a completely different and discontinuous situation of hydrothermal environments, in sharp contrast to the linear setting of mid-oceanic ridges such as the Mid-Atlantic Ridge (MAR) or the East Pacific Rise (EPR). These unique features raise questions about how abiotic and biotic factors shape the distribution of genetic diversity and the connectivity between populations of different lineages.

The vent fauna and communities inhabiting these Southwest Pacific BABs appear as a single biogeographic unit (Bachraty et al., 2009; Moalic et al., 2012; Tunnicliffe et al., 2024). In contrast to other hydrothermal communities, which are mainly composed of tubeworms,

mussels and shrimps, the BAB fauna is mainly composed of large symbiotic Provannidae

gastropods, such as *Ifremeria nautilei* or *Alviniconcha spp.*, and deep-sea *Bathymodiolus*

mussels. These large engineer species create specific habitats for a wide assemblage of

invertebrate species, including annelids from different families (e.g. Polynoidae, Alvinellidae,

Siboglinidae), limpets (*Lepetodrilus* spp., *Shinkailepas* spp.), barnacles, holothurians, and

crustaceans (copepods, amphipods, shrimps, and crabs) (Desbruyères et al., 2006).

While these vents form thriving oases of life, they also produce metallic sulfide deposits, which

attract the interest of deep-sea mining companies. The upcoming management of the

Southwest Pacific vent fauna will rely on understanding population delimitation and

parameters, which are of crucial importance in conservation biology (Gena, 2013; Niner et al.,

2018; Van Dover, 2011; Van Dover et al., 2017; Washburn et al., 2019). Anthropogenic

exploitation of vent resources has already begun on a Japanese site in the Northwest Pacific

and has been first planned in the Manus BAB (Solwara prospects) but recently abandoned,

although the potential consequences of these activities are not yet known (Carver et al., 2020).

Connectivity and renewal of vent populations is mostly driven by larval dispersal due to the

sedentary nature and the strict relationship with the vent fluid. While some vagile fauna, such

as fish, crabs, or shrimp, may contribute to connectivity through adult migration in response

to local environmental changes, but only to very limited spatial scale (vent fields) (Lutz et al.,

1994; Shank et al., 1998). Direct connectivity assessment is not technically feasible for minute

larvae numbering in millions and their life-traits characteristics (Levin, 1990, Vrijenhoek,

2010). As consequence, demographic connectivity needs to be assessed by indirect methods

such as population genetics, larval dispersal modeling or recent method of elemental

fingerprints tracking (Mouchi et al. 2024).

Dispersal modeling in the region suggested possible but limited larval exchange between

distant back-arc basins (Mitarai et al., 2016). However, in the context of the unstable and

fragmented habitat of deep-sea hydrothermal vents, metapopulation theory predicts long-

distance dispersal to mitigate risks of inbreeding and local extinction (Hamilton & May, 1977;

McPeek & Holt, 1992). Initial genetic analyses of several vent species along mid-oceanic ridges

however displayed conflicting evidence regarding dispersal capabilities, with some suggesting

almost panmictic populations at the ridge scale while others hint at patterns of isolation by

distance and stepwise (re)colonization (Teixeira et al., 2012; Audzijonyte & Vrijenhoek, 2011).

Present-day geographical distribution of genetic diversity reflects the complex interplay of connectivity patterns across space and time. Studying and disentangling the origin of these variations is the main objective of phylogeography (Avise, 2000, 2009; Avise et al., 1987). Expanding this approach, to a multispecies comparative framework within a given biome or ecosystems can highlight the effect of several factors shaping the genetic diversity (Hickerson et al., 2010; Papadopoulou & Knowles, 2016). Using new methods and approaches for large genomic dataset, we can start to disentangle past and present connectivity patterns and offer a unique opportunity to describe species distribution patterns at the community level and to improve scientific guidelines for conservation (De Jode et al., 2023, Gagnaire, 2020).

Our study aimed to elucidate genetic diversity and connectivity patterns across a biogeographic hydrothermal unit in the South West Pacific. We conducted a comparative population genomics analysis on seven vent species spanning various taxonomic groups inhabiting the same environment and geographic range, covering most basins of the South West Pacific region. Phylogeographic patterns exhibited by these species, each characterized by distinct life-history traits, were compared at a regional scale using a genome-wide analysis. This analysis revealed a clear phylogeographic break encompassing all seven species around the Solomon-Vanuatu archipelago islands, alongside a contact zone on the Woodlark ridge observed in two species. Based on inferred demogenetic histories, we propose a scenario of vicariance for all species. Furthermore, estimating genetic diversity and gene flow at the community scale underscores the importance of comprehending population connectivity across different geographic scales. This understanding is vital for informing potential management strategies, particularly in the context of future deep-sea mining activities, and for gaining more insights into biogeography of deep-sea species.

Material and methods

Sampling

Seven hydrothermal-vent species from four main vent habitats have been sampled over five Western Pacific regions (the Manus, Woodlark, North Fiji, Lau back-arc basins) and the Futuna volcanic arc (see SI Table 1 and Figure 1). These species include the emblematic symbiotic snails *Ifremeria nautilei* and *Alviniconcha kojimai*, the vent mussel *Bathymodiolus manusensis*, the limpets *Shinkailepas tollmanni*, *Lepetodrilus schrolli*, and *L. aff. schrolli* (recently described as *L. fijiensis* by Chen & Sigwart 2023 in North Fiji, Lau and Futuna BAB), which live on the shells of *I. nautilei, Bathymodiolus manusensis* and *B. septemdierum,* the cirriped *Eochionelasmus ohtai*, mostly found on the edge of diffuse venting areas, and, finally, the large scaleworm *Branchinotogluma segonzaci*, found on the walls of vent chimneys. Animal collections were made during the Chubacarc cruise in 2019 (chief scientists S. Hourdez & D. Jollivet) on board the RV *L'Atalante* with the ROV Victor 6000 (Hourdez & Jollivet 2019). All species, with the exception of *B. segonzaci,* were sampled from diffuse venting areas with the tele-manipulated arm of the ROV and brought back to the surface in thermally insulated boxes. The mobile chimney scaleworms *B. segonzaci* were collected using the slurp gun of the ROV and kept in 5 L bottles until the ROV recovery. On board, large animals (provannid gastropods, mussels, cirripedes and scaleworms) were dissected to separate tissues and individually preserved in 80% ethanol, and/or directly used for DNA extractions. A hierarchical sampling scheme was used for all species by sampling two replicate sites within each vent field (locality) and one to three vent fields per basin, for a total number of 21 sampling localities for each vent community (SI Table 1). A total number of 24 individuals were preserved for DNA extraction (or extracted directly) for each locality replicate and species.

This sampling scheme was however not possible for *B. manusensis*. It was only found at Manus, Futuna and one site of Lau (at the Mangatolo locality) in sympatry with *Bathymodiolus septemdierum* (previously known as *B. brevior*). It was not found at the site La Scala in the Woodlark Basin (no mussels), nor at the Southern localities of the Lau Basin and along the North Fiji Ridge, where *B. septemdierum* was only found. DNA extractions were directly performed on board for *I. nautilei* (foot tissue), *A. kojimai* (foot tissue), *B. manusensis* (mantle tissue), *S. tollmanni* and *L. schrolli & L. aff schrolli (now L. fijiensis)* (whole individual without the shell) and *E. ohtai* (whole individuals without the skeleton). For *B. segonzaci,* DNA

extractions were done later in the laboratory on ethanol-preserved tissues. DNA extractions
were either performed with a modified CTAB 2%/PVPP 2% protocol (see Jolly et al. 2003) or
the NucleoSpin®Tissue 96 kit (Macherey-Nagel, Karlsruhe, Germany).

Figure 1: Sampling areas in the South-west Pacific Ocean. Colors represent the different areas   (BABs and volcanic arc). Each
square box contains species found and sampled for each sampling area. 1: I.nautilei, 2: A. kojimai, 3: B. manusensis 4: B.
segonzaci, 5: E. ohtai, 6: S. tollmanni and 7: L. schrolli (& fijiensis). Small points represent hydrothermal vents. Red, active and
confirmed. Yellow, Active and inferred. Green, Inactive. Vents activity data taken from InterRidge Vents Database V3.4.

Preparation of the ddRAD Libraries.
The preparation of ddRAD genomic libraries was standardized by following the protocol
described in Daguin-Thiébaut et al. (2021) and used for *I. nautilei* in Tran Lu Y et al. (2022).
These seven libraries comprise 294 individuals for *A. kojimai* (from Castel et al. (2022)), 469
individuals for *S. tollmanni*, 282 individuals for *E. ohtai*, 195 individuals for *B. manusensis*, 262
individuals for *B. segonzaci*, and 522 individuals for *L. schrolli & L. fijiensis*. All libraries were
produced from gDNA digested with the enzymes *Pst1* and *Mse1,* except for *B. manusensis* that
was digested with *Pst1* and *Msp1*.

These libraries also included 8 to 47 replicates (individuals replicated twice or three times as
controls) used for quality control and parameter calibration. Single-end (only for *B.*
manusensis) or paired-end 150 sequencing (all other taxa) was performed on HiSeq 4000 (*I.*
nautilei) or Novaseq 6000 Illumina (all other taxa), by the Genoscope, France (*I. nautilei*), or
Novogene Europe (Cambridge, UK; all other taxa). The Fastqc Software (V.0.1.19) was used to
check the sequence quality of the raw reads prior to the *de novo* assembly for each species.

An independent assembly was performed from each species library with the "*de novo*" Stacks2
module (Rochette et al., 2019) after demultiplexing individuals with the Process_radtags
module. Parameter calibration approach followed the recommendations of Mastretta-Yanes
et al. (2015) and Paris et al. (2017), (see SI Calibration). For all species, the parameter
calibration approach, data filtering, and the Stacks modules used were consistent with the
methods described in Tran Lu Y et al. (2022). However, for *S. tollmanni* and *L. schrolli* & *L.*
fijiensis, the data filtering was slightly modified to reduce the loss of SNPs and individuals while
minimizing the proportion of missing data due to a greater allele divergence (SI Table 2).
Additionally, for each species, the minor allele frequency filter was replaced by a masking of
singletons in the demogenetic inference datasets (i.e. used with ∂a∂i). Finally, the "*de novo*"
assembly of *B. manusensis* data was realized from single-end reads (R1, 150 bp).

Population structure and admixture across the southwest Pacific

For each species, a series of independent population genomics analyses were performed to
explore the population structure, phylogeographic and admixture patterns. Principal
Component Analysis (PCA) was performed on individual genotypes with SNPRelate (V.1.21.7)
to explore the spatial distribution of genetic diversity (Zheng et al., 2012). Further analyses to
explore population differentiation and admixture proportions were performed with the
software Admixture (V.1.3.0) (Alexander & Lange, 2011). Population trees including migration
edges and F_3 statistics were computed with the software Treemix (V.1.13) (Pickrell & Pritchard,
2012) with ten replicates and migration events from 0 to 5.

The degree of genetic differentiation was estimated with pairwise F_{st} values between groups
of individuals (genetic units or metapopulations, basins or localities) with Arlequin (V.3.5.2.2)
(Excoffier & Lischer, 2010) and their statistical significance was assessed with 10,000
permutations of genotypes between populations. All plots were produced with R (V.4.0.3) (R

Core Team, 2023) and ggplot2 (V.3.3.6) (Wickham, 2016). For *S. tollmanni,* Woodlark individuals were split into two groups according to their genetic assignment (See results).

In addition, the net divergence between two populations (D_a) was calculated from the measure of absolute divergence (D_{xy}) corrected by the average nucleotide diversity (π) estimated using Stacks2 (Rochette et al., 2019) by following the formula of Nei & Li (1979). The genetic diversity indices (*He*, *Ho*, π) were also calculated for each species with the Stacks (V.2.52) population module on the final dataset. Indices were estimated for each genetic unit found by PCA and the Admixture analysis, which number depends on the species considered and on geography (five geographic regions/BABs listed in SI Table 1).

Evolutionary history of vent species and metapopulation connectivity

Relative gene flow orientation.

Gene flow orientation was assessed with the module Divmigrate (Sundqvist et al., 2016) contained in the R package diveRsity (V.1.9.90) (Keenan et al., 2013). This method uses allele frequencies and several estimators derived from the Fst statistics to calculate a migration matrix between populations, which is then normalized by the highest gene flow value found in the pairwise comparisons, setting gene flow value between 0 and 1 (where 1 represents 100% gene flow and 0, no gene flow). Statistical significance of the difference between gene flow directions in a given pair of populations was estimated with 1000 bootstraps.

Demogenetic history of species metapopulations

To understand the demographic history of populations and identify the best model of population divergence, we fitted joint allele frequency spectra to specific population models with the *∂a∂i* software (V2.1.0) (Gutenkunst et al., 2009) for each taxon independently, considering the case of an ancestral population splitting into two daughter populations with or without migration. Our choice was dictated by the fact that populations of all vent species analyzed (with the exception of *L. schrolli & L. fijiensis*, see results) are only subdivided into two genetic units (Eastern vs. Western populations). Although, *L. schrolli & L. fijiensis* displays further minor subdivisions between basins, for the sake of the comparison with other species, we considered Eastern (NF/L) vs Western populations (M) (see discussion). One of the main advantages of the *∂a∂i* approach is that it takes into consideration the effects of linked selection and heterogeneous migration across the genome. Ignoring these effects in

demographic inference may lead to erroneous reconstructions of the evolutionary history of populations (Ewing & Jensen, 2016; Ravinet et al., 2017). To discuss patterns of divergence and past and present genetic connectivity of all species, we re-used the models implemented in Tran Lu Y et al. (2022) with a similar population design for all taxa. These models considered a total of 28 possible scenarios extended from the 4 major divergence models (Strict Isolation (SI), Isolation with Migration (IM), Ancient Migration (AM) and Secondary Contact (SC)) that were initially developed and used in Rougeux et al. (2017) (also see Tran Lu Y et al., 2022 for further details). Briefly, different demographic and evolutionary processes have been considered within these models, such as the increase or the contraction in size of the two derived populations (G), the effect of barrier loci due to either hybrid counterselection or local adaptation by simulating heterogeneous migration along the genome (2m) and the effect of linked selection (2N).

The ancestral allele of each locus could not be identified because no external groups were available (or too divergent) for any of the taxa. Hence, we used folded joint allele frequency spectra (folded JAFS). All models were fitted 10 times independently for each species to check for model convergence. The Akaike Information Criterion (AIC) was used to compare models for each simulation. Given the gaps in the knowledge of the biology of these taxa, we estimated model parameters and divergence times (i.e. Ts since divergence, Tsc since secondary contact, and the absolute divergence time, which can be Ts or Ts+Tsc) using an average mutation rate of 10^{-8}. Parameter uncertainties were calculated using the Fisher Information Matrix (FIM) on the best fit model for each species. The uncertainties are calculated at the 95% confidence level.

Results

Calibration parameters and filtering steps.

The 150 bp paired-end sequencing yielded an average of 3.7, 3.3, 3.0 and 2.8 million of paired reads per individual for *S. tollmanni*, E. *ohtai, B. segonzaci* and *L. schrolli & L fijiensis*, respectively. For *B. manusensis,* single read data were obtained with 2.9 million reads per individual. The raw reads used from *A. kojimai* and *I. nautilei* were obtained following the protocol described in Castel et al. (2022) and Tran Lu Y et al. (2022).

For each species, we identified the most appropriate set of assembly parameters with Stacks (V.2.52) by applying the same pipeline of the *de novo* assembly of bi-allelic loci used for *I. nautilei* in Tran Lu Y et al (2022) following the procedure recommended by Mastretta-Yanes et al. (2015) and Paris et al. (2017). After running several sets of parameter combinations, we decided to use the assembly parameters that ranged between 4 and 6 for m, 4 and 11 for M and 5 and 11 for n, depending on the species under scrutiny (see SI Calibration & SI Table 3). In addition, we applied the same methodology, analyses, modules and filtering parameters that we used for *I. nautilei* in Tran Lu Y et al. (2022) to better compare genetic patterns between species (see SI Calibration). The *de novo* assemblies and subsequent filtering steps resulted in a variable number of SNPs (2 904 to 47 547) obtained for 159 to 414 individuals retained for each of the 7 species (see SI Table 2 & 3).

Population structure and admixture.

Population structure analyses revealed similar patterns for the spatial distribution of genetic diversity for the seven species. The PCA showed a common genetic structure, separating the Manus individuals from those of the Eastern zones (i.e. North Fiji, Futuna and Lau, hereafter NF/F/L) on the first component (PC1), with 1.94% to 26.03% of the total variance explained (Figure 2). However, the site La Scala on the Woodlark Ridge (newly discovered, Boulart et al., 2021) showed contrasting results. First, *I. nautilei*, *A. kojimai*, and *E. ohtai* are clearly subdivided into two genetic groups with individuals of Manus-Woodlark (M/W) on one hand, and individuals from NF/F/L on the other hand (Figure 2 A, B, D). However, the genetic relationships of individuals from Woodlark differ slightly for the other species. *S. tollmanni* was also subdivided into two genetic groups but individuals from Woodlark clustered with either the Manus or with the NF/F/L genetic groups, with one admixed individual positioned in between (Figure 2 E). For *L. schrolli & L fijiensis,* all individuals from Woodlark are positioned as intermediates between Manus (*L. schrolli)* and NF/F/L (*L fijiensis)*. These intermediate individuals are genetically closer to the North Fiji individuals, which slightly differed from the Futuna/Lau (F/L) individuals on PC1 (Figure 2 G). A comparable situation also holds for the species *B. segonzaci* for which the Woodlark individuals are well separated from either individuals of Manus or those of the NF/F/L group but are a little bit closer to the latter (Figure 2 F). For *B. manusensis,* which is absent from La Scala, a similar situation is observed (Figure 2 C).

The second component (PC2) accounting for 0.62-0.76% of the total genetic variance, also showed slight differences at the regional scale for some species (Figure 2 B, F, G). Individuals from North Fiji exhibit a slight genetic differentiation from the group F/L on PC2 for both *A. kojimai* and *L. fijiensis*, and the same pattern is observed between Woodlark and Manus individuals for *B. segonzaci* and *L. schrolli.* According to the first two principal components of the PCA, the only species exhibiting a more basin-specific signature was the species complex *L. schrolli/L. fijiensis* (Figure 2 G and SI Figure 1).

This shared distribution of the genetic variation was also captured by admixture analyses for all species. The optimal number of clusters identified was always k = 2 (SI Figure 2). These two genetic units are composed, as previously found, of M/W individuals on one side, and NF/F/L on the other side. Except for some individuals from Woodlark and North Fiji BABs for which genome admixture ranged between 10 to 50% depending on the species under scrutiny, the proportion of shared ancestry between these two groups is very low for most individuals (Figure 2). Woodlark individuals for *L. schrolli/L. fijiensis, S. tollmanni* and *B. segonzaci* display shared ancestry from both genetic clusters. In *S. tollmanni*, the Woodlark population is composed of a mixture of parental types (both genetic backgrounds) and a putative F1 hybrid. In contrast*,* all Woodlark individuals displayed almost equal (~50%) shared ancestry from both *L. schrolli* (Manus group) and *L. fijiensis* (Eastern group) when k = 2. Admixture analyses also discriminate additional clusters for Woodlark and then North Fiji in the *L. schrolli/L.fijiensis* species complex with increasing values of k (K = 3 to 5, SI Figure 3 & 4).

The F_3 statistics showed significant negative values, indicating admixture in Woodlark, for two species only. The first species is *S. tollmanni*, with individuals from the Western (Woodlark2) population (Manus type) with source populations coming from both Manus (M) and the Eastern group (NF/F/L) and Woodlark1 (Lau type). The Woodlark *Lepetodrilus* individuals displayed a similar situation with source populations coming also from Manus (M) and the Eastern group (NF/F/L) (SI Figure 5).

Examining the overall genetic differentiation (F_{st}) and the net divergences (D_a) between the Western and Eastern groups provided three main patterns (Table 1). The first group of species (i.e. the limpets *Lepetodrilus* and *Shinkailepas*) is characterized by high values of differentiation (0.271-0.360) and divergence (0.013-0.019). The species of the second group (*I. nautilei, E. ohtai* and *B. manusensis*) also have high values of genetic differentiation (0.203-

0.387) but low to moderate divergences (0.002-0.007). Finally, the last group comprising *B.*

segonzaci and *A. kojimai* has low values of genetic differentiation (0.018-0.038) and moderate

divergences (0.007).

Pairwise Fst also captured differentiation between localities, when the five regions were

considered as separate populations, with F_{st} ranging from 0 to 0.364, depending on the species

analyzed and the pair of populations compared (SI Table 4). Most cases of significant genetic

differentiation between BABs were observed in pairwise comparisons between the Western

and Eastern BABs (Manus or Woodlark against Lau, North Fiji and Futuna). Some

differentiation was also observed at the basin scale within a group with some specific

separation of the populations from North Fiji and Futuna/Lau for the species *A. kojimai* and *L.*

fijiensis. No population differentiation between Lau and Futuna has been however observed

except for *L. fijiensis*, where differentiation is extremely low but significant.

354
355 *Figure 2: All PCA (PC1 & PC2) and Admixture plots for the best number of genetic clusters (K = 2) for each species (A, B, C, D,*
356 *E, F , G). Colors in PCA plots represent regions (Manus, Woodlark, North Fiji and Lau Back-Arc-Basins, and the Futuna Volcanic*
357 *Arc). Open circles represent the multivariate normal distribution of each group of points (basins) at 95% in PCA plots. Colors*
358 *in Admixture plots represent each inferred genetic cluster. M: Manus, W: Woodlark, NF: North Fiji, F: Futuna, L: Lau.*

Table 1: Fixation index (Fst) and net nucleotide divergence (Da) measured between the two (Eastern and Western) metapopulations for each species.

Species	F_{st} (M/W vs NF/F/L)	Da (M/W vs NF/F/L)
Lepetodrilus schrolli & L. fijensis	0.360	0.019
Shinkailepas tollmanni	0.271	0.013
Ifremeria nautilei	0.387	0.008
Eochionelasmus ohtai	0.203	0.007
Branchinotogluma segonzaci	0.038	0.007
Alviniconcha kojimai	0.018	0.007
Bathymodiolus manusensis	0.206	0.002

Treemix analyses revealed a consistent two-group pattern of population differentiation across species, with varying optimal numbers of migration events (ranging from 0 to 2) between the two regions (SI Figure 6 and 7). These findings highlight the occurrence of a primary migration event between the Eastern and Western groups for all species, excluding *E. ohtai* and *B. manusensis*. Furthermore, a secondary migration event was exclusively observed for *I. nautilei*, *A. kojimai*, and *S. tollmanni*, occurring between two different basins of the same genetic group (i.e., metapopulation). Notably, these interacting basins were not consistently the same across different species. *B. segonzaci* exhibited a distinctive pattern, featuring two migration events between the Eastern and Western groups: the first one from Futuna to Woodlark and the second from Lau to Manus.

Genetic diversity of species

Regardless of the statistics used (Ho, He and π: SI figure 8 & 9), *S. tollmani* displayed a gene diversity twice higher than that of the other species. The level of genetic diversity of species slightly differed between the Eastern and Western groups, but not always in the same direction. *I. nautilei*, *B. manusensis* and *L. schrolli/L.fijiensis* exhibit a slightly higher gene diversity in M/W compared with NF/F/L whereas it was the opposite for the other species (*A. kojimai*, *S. tollmani*, *E. ohtai* and *B. segonzaci*). These statistics displayed exactly the same pattern of distribution when calculated per basin (see SI Figure 9).

Evolutionary history of populations and connectivity at the multispecies scale

Relative migration rates

The Divmigrate analysis between the Eastern and Western groups revealed a robust and common pattern of bidirectional but asymmetrical gene flow in all species. The main direction of gene flow predicted by the analysis is westward from NF/F/L (1) to M/W (2), while gene flow in the opposite direction was about half to a two-third (SI Figure 10). The species complex *L. schrolli/L. fijiensis*, however, displayed a more complex bidirectional pattern with a much stronger gene flow between Futuna and Lau than between Futuna/Lau and North Fiji and virtually no gene flow between Woodlark, Manus and the others BABs.

Demo-genetic inferences

Modes of divergence

The folded JAFS (Figure 3) show the distribution of allelic variants between the Eastern and western metapopulations for each species. Of the 28 models tested for each species, the SC (Secondary Contact) model was almost always the best-fitting model selected by the Weighted AIC. The exceptions are *I. nautilei* (Tran Lu Y et al., 2022) and *B. manusensis*, for which the SC model is slightly, but not significantly, better than the IM (Isolation with Migration) model. The Strict Isolation (SI) model was the worst model for all species. This indicated that all species are still able to maintain low levels of gene flow at the regional scale despite the large geographical distances and the abyssal plain separating the different regions. Increasing the complexity of the SC or IM population model by the addition of genomic heterogeneity (2N and/or 2m) or demographic change (G) parameters improved the model fit (see Figure 4 and SI Figure 11). Capturing linked selection (2N) improved the model fit more than the G or 2m parameters for most species except *E. ohtai*. Alternatively, adding heterogeneous gene flow (2m) to the SC model improved the model fit for *E. ohtai* (SC2mG), although the two metapopulations appeared to be well separated. This situation of semi-permeable barrier also holds for *S. tollmanni* and *L. schrolli/L. fijensis,* for which genetic admixture is locally suspected. For these two latter species but also for *A. kojimai*, *B. segonzaci*, and *I. nautilei,* the SC2N2mG was the best model after evaluating all possible parameter combinations. For *B. manusensis*, it was however not possible to discriminate

between the models IM2N, SC2N, and SC2N2m, and both SC2mG and SC2NG performed similarly as well as the SC2N2mG model for *B. segonzaci* (Figure 4 and SI Figure 11 & 12).

Figure 3: Folded Joint Allele Frequency Spectrum (JAFS) plots for all species. Each four plots represent, Observed Joint allele frequency spectrum (JAFS) between lineages East and West. NF: North Fiji, F: Futuna, L: Lau, M: Manus, W: Woodlark

Timing of divergence and gene flow

Depending on the best model, we estimated divergent times Ts (time since the population split), Tsc (time since the secondary contact after the primary divergence), and Ttotal (which could be Ts or Ts+Tsc), using a fixed mutation rate of 10^{-8} (Table 2). Ts ranged by a factor of ten thousand generations to about one hundred thousand generations according to the species analyzed (40 000 for *A. kojimai* to 116 000 for *L. schrolli/L. fijensis*). Similarly, the time elapsed since secondary contact (Tsc) changes also by a factor of 10 from 6,386 generations for *S. tollmanni* to 69,371 generations for *I. nautilei* and *L. schrolli/L. fijensis*. However, the total time since the ancestral population split (Ts or Ts+Tsc) varied less, with the lowest values being 40,892 and 42,117 generations for *A. kojimai* and *B. segonzaci* respectively, and the highest values being 94,071, 116,711 and 101,718 generations for *E. ohtai*, *L. schrolli/L. fijensis* and *B. manusensis*, respectively. For *B. manusensis*, the time of divergence however varied up to two-fold (43,644 to 101,718

generations) depending on the best models used. *I. nautilei* and *S. tollmanni* produced intermediate values of spitting times.

All species display heterogeneous gene flow (2m). Migration rate parameters estimated from *∂a∂i* show that present-day gene flow is stronger from NF/F/L to M/W (East to West) than the opposite, with the exception of *L. schrolli/L. fijiensis* for both the two classes of gene flow ("neutral" migration m and "reduced" (due to barrier loci) migration me, in relative genomic proportions P and 1-P; Table 3). For *B. manusensis*, migration rates were similar in both directions. *I. nautilei* and *E. ohtai* had about half of their genome characterized by reduced gene flow (P ~=1-P). Only *A. kojimai* displayed high neutral gene flow (with m>me and P>>1-P) whereas *S. tollmanni, B. segonzaci, B. manusensis and L. schrolli/L. fijiensis)* exhibited a larger proportion of barrier loci (1-P >>P) which strongly reduced gene flow (me <<m) between the genome of the two genetic groups (Table 3).

Table 2: Estimation of divergence times (Times are estimated with a fixed mutation rate per generation and site, 10^{-8}) and estimation of mutation rate per site and generation for all species. Ts: time since the ancestral population subdivided into two populations; Tsc: time since secondary contact. Ttotal: Ts and Ts + Tsc. For I. nautilei using the AM2N2mG model, Ts corresponds to the time of the strict split after ancient migration ended (Tsc).

Species	Best model	Ts	Tsc	Ttotal
I. nautilei	IM2N2mG		66 951	66 951
I. nautilei	SC2N2mG	16 279	54 017	70 295
I. nautilei	AM2N2MG	41	69 372	69 413
A. kojimai	SC2N2mG	29 157	11 735	40 892
S. tollmanni	SC2N2m	63 335	5 419	68 754
S. tollmanni	SC2N2mG	42 683	22 326	65 009
E. ohtai	SC2mG	37 278	56 793	94 071
B. segonzaci	SC2N2mG	18 194	15 375	33 569
B. manusensis	IM2N	101 718		101 718
B. manusensis	SC2N2m	40 752	2 892	43 644
B. manusensis	SC2N	56 394	37 940	94 335
L. schrolli/L. fijiensis)	SC2N2mG	48 381	68 331	116 712

451   *Table 3: Gene flow parameters estimated from ∂a∂i (m and me are the "neutral" and "reduced" migration rate parameters.*

452   *P the proportion of the genome characterized by migration rate m, and 1-P the proportion of the genome affected by reduced*

453   *migration me). 1 : East population (NF/F/L); 2 : West population (M/W).*

Species	Best model	m1<-2	m2<-1	P	me1<-2	me2<-1	1-P
I.nautilei	IM2N2mG	0,444	0,825	0,483	0,038	0,283	0,517
I.nautilei	SC2N2mG	0,422	0,810	0,439	0,038	0,270	0,561
I.nautilei	AM2N2MG	0,461	0,825	0,471	0,039	0,300	0,529
A. kojimai	SC2N2mG	4,741	4,389	0,851	0,384	0,841	0,149
S. tollmanni	SC2N2m	0,321	3,314	0,014	4,334	13,866	0,986
S. tollmanni	SC2N2mG	0,268	1,048	0,205	1,048	2,800	0,795
E. ohtai	SC2mG	0,364	1,386	0,508	0,013	0,124	0,492
B. segonzaci	SC2N2mG	0,579	6,489	0,000	2,195	5,412	1,000
B. manusensis	IM2N	0,436	0,566				
B. manusensis	SC2N2m	3,988	4,891	0,323	0,227	0,646	0,677
B. manusensis	SC2N	0,520	0,537				
L. schrolli/L. fijensis)	SC2N2mG	0,878	0,308	0,180	0,009	0,007	0,820

Figure 4 : Weighted AIC value on the best runs for all models and parameters per species. Highest value represents the best
model fit (the lowest AIC value). Ifremeria nautilei data are derived from Tran Lu Y et al. (2022).

Discussion

Population genomics allowed us to gain a deeper understanding of factors that have shaped
genetic variation within and between populations in a context of a discontinuous ridge system
in a well-delimited biogeographic region. Our study was aiming to "put the geography (and
more) into comparative population genomics" (from Edwards et al., 2022) by applying such
an approach to seven very different taxa strictly associated with the hydrothermal habitat.
Using the same sampling scheme across all species, we show that these species share a
common biogeographic break and a similar demographic history despite their different life-
history traits. This pattern of differentiation was however slightly more complex for the limpet

L. schrolli/L. fijiensis, S. tollmanni and the scaleworm *B. segonzaci*, for which a further genetic

subdivision between the Manus Basin and the Woodlark Ridge was also observed.

The multispecies transition zone between the Eastern and Western metapopulations is

located on the Woodlark Ridge or between it and the western part of the North Fiji BAB,

isolating the Manus Basin to the West and the North Fiji and Lau Basins, as well as the Futuna

volcanic arc to the East. Based on demogenetic inferences, we found that after a primary

allopatric divergence, these metapopulations were secondarily connected, resulting in weak,

asymmetric, and predominantly westward ongoing gene flow. Levels of divergence and

differentiation, however, varied between taxa, probably because of different generation times

and life-history traits. Heterogeneous gene flow was also found in all species in various

proportions, suggesting that primary divergence could have led to the generation of genomic

incompatibilities, as commonly found in hybrid zones (Bierne et al., 2011; Matute et al., 2010).

A suture zone around the Woodlark back-arc Basin.

All seven species share a clear genetic break into two main metapopulations across their

geographical range, with the Manus/Woodlark (M/W) BABs on one hand and the North Fiji,

and Lau BABs and Futuna Volcanic Arc (NF/F/L) on the other hand. Despite this common

phylogeographic pattern, the strength of differentiation between these Eastern and Western

genetic pools estimated by F_{st} varied from very high (0.387) to very low (0.020), depending on

the species. A transition zone, a region where previously isolated lineages come into contact

and exchange genetic material, appears to be located somewhere between North Fiji and

Woodlark BAB or even at Woodlark itself, where both lineages are found in sympatry for some

species. This is clearly observed for the two limpet species complexes, *S. tollmanni* and *L.*

schrolli/L. fijiensis, for which the Western and Eastern lineages co-occur in Woodlark and

hybrid individuals, are detected. This signature is typical of the existence of a tension zone,

where selection can operate against hybrid genotypes, probably due to the existence of

underdominant genetic incompatibilities (Bierne et al., 2011) or other reproductive barriers.

Variable levels of admixture detected between Western and Eastern lineages in all species

present at Woodlark, and, to a lesser extent at North Fiji (Tran Lu Y et al. 2023), reinforced this

view. The species variability in terms of population structuring can stem from diverse origins:

differences in life history traits (generation time, dispersal capability, fecundity, ...) or in the

genomic architecture of species (intensity of linked selection and number of barrier loci

(genetic incompatibilities) that whole genome information could help better visualize. These

latter factors are noticeably linked with the depth of evolutionary history of divergence and

the reproductive mode. Our study suggests two possible hypotheses. First, geophysical

rearrangements at a given time may have limited (and are still currently limiting) effective

dispersal for a number of species with different life history traits. Leading this common pattern

of isolation, regardless of the timing of their origin in the SouthwesternPacific (isolation with

migration, IM, one of the models supported by *∂a∂i* for *Ifremeria nautilei* and *Bathymodiolus*

manusensis). Alternatively, these species may have experienced one or more vicariance

events, resulting in a period of primary divergence in allopatry followed by secondary contacts

(SC) with a gene flow restart. The latter scenario is the most likely, supported by the *∂a∂i*

analyses for most species (Figures 3 & 4).

While evolutionary processes unfold independently in each species, our demogenetic

reconstruction indeed indicates a scenario of secondary contact (SC) for five species. For two

species, the model cannot decisively distinguish between secondary contact and isolation-

with-migration (IM), possibly due to a prolonged period of secondary contact. These results

nevertheless strongly suggest an allopatric initial divergence for all species. Modeling also

underscored the detection of a genomic heterogeneity of differentiation in all species, with

both heterogeneous gene flow across the genome (2m) and linked selection (2N). This finding

suggests that the initial divergence was extensive enough to generate genomic

incompatibilities between populations in a genome characterized by substantial background

selection and low recombination. This pattern can be a feature of hydrothermal-vent species,

which may undergo strong purifying selection in this challenging environment (Chevaldonné

et al. 2002, Fontanillas et al. 2017, Thomas-Bulle et al. 2022).

The position of the contact zone between Western and Eastern lineages, today located at

Woodlark or closeby, may have moved since the secondary contact took place, probably along

the subduction arcs of Vanuatu or Solomon. Nevertheless, the Manus Basin appears more

diversified in terms of species (Poitrimol et al., 2022) and could have served as a source of

biodiversity for part of the Western Pacific hydrothermal fauna, suggesting a possible "out of

Manus" hypothesis. In line with this idea, recent work on taxa network analysis redescribes

the Western Pacific region for hydrothermal fauna not as a single biogeographic province, as

suggested by Moalic et al. (2012), but as two distinct provinces, the North West Pacific and

the South West Pacific, with the Manus Basin as a possible hub connecting both (Tunnicliffe et al., 2024).

Gene flow between BABs

The connectivity and dispersal of hydrothermal vent species may be influenced by their life history traits, particularly the larval phase. Species with planktotrophic larvae, which can have an extended planktonic phase, should exhibit higher dispersal abilities and greater connectivity between populations. In contrast, species with lecithotrophic larvae, which rely on yolk reserves and usually have a shorter planktonic phase in coastal waters, are expected to show more limited dispersal and reduced connectivity (but see: Young 1994 for the deep-sea fauna). Our findings, with a consistent asymmetrical gene flow (lower eastward gene flow), unequivocally reject complete isolation between East and West metapopulation/lineages for all species, despite their contrasting dispersal life-traits. It partially also aligns with prior studies indicating very low gene flow between Manus and Lau BABs (Breusing et al., 2021; Plouviez et al., 2019; Thaler et al., 2011, 2014). However some other studies have reported a lack of genetic differentiation for *S. tollmanni* with some mito-nuclear incongruities (Yahagi etl., 2019, Poitrimol et al., 2022). Additionally, our demogenetic modeling indicates semi-permeability (2m) of gene flow for all species, probably from the accumulation of genomic incompatibilities and other barrier loci during primary divergence preceding secondary contact (Barton & Bengtsson, 1986).

The orientation of gene flow also partially aligns with previous simulations of larval dispersal at intermediate depth (Mitarai et al., 2016). However, the bi-directional nature of the gene flow suggests potential dispersal near the surface, where the main counter-current flows eastward, in addition to the deep-water westward dispersal. The current genetic data and the lack of information on the biology of the species are insufficient to assess the role of life-history traits, particularly the larval developmental mode of the species, in relation to the degrees of differentiation, divergence, or even the orientation of dispersal at this stage. Further analyses are required with a better understanding of the biology of these species. Depending on the life-history traits and potential presence of intermediate populations in the Vanuatu and Solomon archipelagos, the frequency and duration of secondary contacts may have varied between the Eastern and Western lineages. Nevertheless, present-day connectivity remains highly limited due to the existence of both genetic and physical barriers to dispersal. In this scenario, species with high dispersal abilities and/or long generation times

may have experienced less divergence (e.g., *A. kojimai, B. segonzaci)* compared to those with low dispersal abilities and short generation times (e.g., *L. schrolli/L. fijiensis*). This situation is well illustrated in vent copepods in the Lau Basin, which possess both of the latter attributes (Diaz-Recio Lorenzo et al., 2024).

Species specific variation

While the West/East divergence is clear for all species, few species display some basin peculiarities.

B. segonzaci shows slight differentiation between Woodlark and Manus Basin populations, not solely due to allelic introgression from other Eastern regions. The North Fiji Basin population of *A. kojimai* differs slightly from the Lau/Futuna regions, despite the overall lack of genetic differentiation (Fst= 0.018) at the regional scale. *B. manusensis* has been newly identified in the Eastern regions (north of Lau Basin and Futuna volcanic arc). This unusual presence in the East is notable, as it locally co-occurred with *B. septemdierum*: the only mussel species typically found in that area. These Eastern scattered populations may represent a leading edge of range expansion but lack the expected reduced diversity (Dupoué et al., 2020). The slight basin differentiation for *B. segonzaci* and *A. kojimai* raises questions about limiting factors on connectivity, potentially influenced by larval dispersal depth, demographic turn-over, or putative depth selection in North Fiji Basin and Woodlark populations.

While most of our species exhibit a relatively low level of genetic divergence, the limpets *L. schrolli/L. fijiensis* and *S. tollmanni* show a high degree of divergence and differentiation, as well as a highly reduced gene flow between the Western and Eastern groups. *S. tollmanni* displays a clear admixture signal in Woodlark, where both lineages are in sympatry and one first generation hybrid was identified. Previous studies have shown a lack of genetic differentiation at the regional scale with the mitochondrial *Cox1* (Poitrimol et al., 2022; Yahagi et al., 2020). Based on genomic data, the two lineages West/East are strongly isolated and display very reduced and heterogeneous gene flow, where alleles can be exchanged at very few loci. This incongruity is consistent with our results. Indicating a strong reduction in gene flow, but where few loci can still be exchanged and captured, including mitochondrial genomes.

The species *L. schrolli* and *L. fijiensis* display a nearly identical divergence but with a clearly
admixed population between the Manus and NF/F/L lineages at Woodlark. This result is in line
with mtDNA *Cox1* results for Woodlark individuals, where half of them reflect Manus *Cox1*
haplotype lineage and the other half, the NF/F/L haplotype one (Poitrimol et al., 2022). This
is also consistent with previous results using 42 nuclear markers, which indicate very low and
asymmetric gene flow between Manus and Lau BABs (Plouviez et al 2019). It indicates
segregation for mtDNA markers in a sympatric population, consistent with the possible partial
accumulation of genomic incompatibilities in which the mitochondrial genome is this time
included. Our findings therefore reinforce the recent taxonomic revision of *L. schrolli* (M) and
L. aff schrolli, now *L. fijiensis* (NF/F/L), as separate species (Chen & Sigwart 2023). This also
suggests that the taxonomy of *S. tollmanni* may need some revision.

As previously mentioned, dispersal-related life history traits potentially influence connectivity.
However, when we look at what we know about the life history traits of our seven species,
there is no consistency with the type of larval development. *B. segonzaci* and *A. kojimai*, both
colonizing the hottest parts of the West Pacific vent environment, display the lowest genetic
differentiation and divergence between Eastern and Western populations, despite having
different larval development modes. *B. segonzaci* is a free-living, mobile annelid with small
local population sizes. Its large oocytes (ca. 150 µm) suggest lecithotrophic larval development
that can remain for long times in cold oligotrophic waters. *A. kojimai*, on the other hand, has
larger patchily distributed populations and produces much smaller oocytes, implying
planktotrophic larval development in surface waters (Warèn & Bouchet, 1993, Sommer et al.
2017, Kim et al., 2022). In these two species, the low level of regional differentiation therefore
could be linked to the specific distribution of 'hot' vent emissions when compared to colder
diffuse venting, provided that they are both able to widely disperse.

B. manusensis and *E. ohtai* also display a low level of divergence, but exhibit an intermediate
level of differentiation between metapopulations. *E. ohtai* is a common sessile hydrothermal
vent cirriped forming dense populations in diffuse areas. Its oocyte diameter (ca. 500 µm)
again suggests lecithotrophic development (~500 µm; SH unpublished data, Yamaguchi &
Newman 1997, Tyler & Young 1999). In contrast, deep-sea bivalves such *as B. manusensis*
typically have small oocytes and larvae are expected to be planktotrophic, as larvae of
Gigantidas childressi (formerly known as "*Bathymodiolus*" *childressi*) have been detected in
surface water and known to live up to 18 months (Arellano and Young, 2009).

Although the focus was on the Southwest Pacific BABs, the Kermadec Basin, hundreds of

nautical miles south of the Lau Basin, has a very distinctive vent fauna. But, *B. segonzaci* and

L. fijiensis are among the very few species shared with the most northerly location sampled

here (SH unpublished data). It therefore appears that these species may have spread for a

longer time and/or have greater habitat flexibility than other species. The addition of some

individuals from Kermadec to the dataset showed no differentiation from the NF/F/L

population for *B. segonzaci*, but did show some differentiation from NF/F/L for *L. fijiensis* (see

details in SI Figures 13, 14, 15, data not shown).

Timing divergences and the hypothesis of vicariance

The comparative study highlights common phylogeographic patterns across the Southwest

Pacific Ocean for seven vent species. This pattern has probably been generated by a shared

initial divergence event, due to either geological or climatic factors, generating two

metapopulations, with the separation lying somewhere between Woodlark and North Fiji

BABs, if we consider the populations have remained in place during the isolation process. This

barrier is semi permeable to gene flow, allowing occasionally metapopulations to exchange

genetic material through secondary contacts but not at the same rate and not over the same

regions of their genomes. This reconnection may also be modulated by species-specific life-

history traits, including type of larval development (planktotrophic vs. lecithotrophic), depth

of larval dispersion, longevity, age of first reproduction and mean generation time or habitat

fragmentation.

The estimation of net nucleotide divergence (*Da*), shows that species have undergone

different periods of divergence. In particular, *L. schrolli/L. fijiensis)* and *S. tollmanni* both

display particularly high net divergence values compared to the other species. This indirect

approach suggests that these two species did undergo a longer period of isolation in allopatry

or a much higher number of generations since a possible vicariant event. These results

naturally pose the question of cryptic species and speciation processes. This level falls into the

grey zone of speciation of Roux et al. (2017), where reproductive isolation between

populations varies widely. Our demogenetic inferences shed some light not only on the

intensity and genomic heterogeneity of gene flow (as previously discussed) but also on the

timing of divergence.

The divergence time since the ancestral lineage split, in generations, indeed varies by a factor of three between species even if they all share the same population history. In this approach, time estimation strongly depends on the per generation nucleotide mutation rate and the mean generation time considered for each species. However, these biological parameters not yet known for the species studied here. This is a common limitation for many non-model organisms (and especially in the deep-sea), and has led us to make assumptions for both parameters that we have set to the same value for all species, although we acknowledge this is highly unlikely. Thus, a factor of three in the divergence time is not so great and can be largely due to the distinct species life history traits. Some studies on growth rates show that larger species have slower growth rates (Schöne & Giere, 2005), while small ones have higher rates (Poitrimol et al., 2024), suggesting discrepancy in the generation time and first age at maturity between species. In addition, bigger species often have smaller population sizes than smaller ones and we cannot exclude the possibility that they would have otherwise undergone relatively simultaneous demographic changes.

Resulting estimates of primary divergence may have begun between ~ 40,892 to 101,718 generations, while the secondary contact ranges from 6,386 to 69,372 generations, depending on the species. Considering an average mutation rate of 10^{-8} and a quite short time of one generation per year, this primary divergence and secondary contact would take place in the Holocene between climatic oscillations. The initial split may be more specifically around the Last Glacial Maxima for the region (11,500-20,300 years ago (Tongo Glaciation), 62,000 years ago (Komia Glaciation) and 130,600-158,000 years ago (Mengane Glaciation) (Barrows et al., 2011)).

At the other end of the range. If we consider a ten times lower mutation rates (10^{-9}) that has been largely invoked in the molecular dating of the vent fauna as the result of strong purifying selection (Chevaldonné et al. 2002, Johnson et al. 2006, Matabos and Jollivet 2019). The magnitude of divergence would be in the order of 400,000 to 1,000,000 years and 63,860 - 693,720 for the secondary contact. This primary divergence timeframe falls partly at the start of magmatic accretion in some actual active BAB ridges, such as Lau or Manus (Schellart et al., 2006). The geological structures that currently separate the two genetic entities are the Woodlark Basin, North-Fiji and the other geological structures between them (Vanuatu Trough (formerly New Hebrides), Vanuatu and Solomon Islands volcanic arc). These formations have a rather older geotectonic history with accretion times of several million

years (Woodlark: ~ 6 Ma, North Fiji: ~3Ma, Vanuatu Trough & volcanic arc: 12Ma, Solomon

volcanic arc: Eocene, ~ 40 Ma) (Schellart et al., 2006). While this primary divergence period

better fits these geodynamic features, the time frame for secondary contact still remains in

the Holocene period and may be the consequences of climatic oscillations and possible

changes in the overall Pacific water-mass circulation due to the ocean elevation.

Timeframe estimates remain very difficult to interpret due to many unknowns on the biology

of the species, but all species were affected, suggesting a major climatic or geological

reorganization of connectivity that probably initiated the primary divergence and the

secondary contact. However, we cannot reject the hypothesis that some secondary contacts

are much older than others since life-traits history varies between species.

Limits of the method and other hydrothermal species

The present study primarily focused on the most common and emblematic species in

hydrothermal communities. Earlier studies on two of our target species and other

hydrothermal vent inhabitants have yielded findings consistent with our results (Lee et al.,

2019; Plouviez et al. 2019; Poitrimol et al., 2022; Thaler et al., 2014). Like all our species, they

exhibit a 'common' pattern of genetic differentiation across the Western and Eastern parts of

this region, implying a shared vicariant event. However, it is important to note that these

results may primarily reflect the evolutionary history of the most highly abundant vent

species. As mentioned earlier, these ecosystems are home to numerous other species. Many

of them probably have lower population densities, are less dependent on vent fluids and have

diversified much more rapidly towards endemism. Consequently, these less abundant and

often smaller species may demonstrate more contrasted phylogeographic patterns. For

example, based on *Cox1* data, some limpets, barnacles, and copepods have been shown to

have a wide distribution, while others (sometimes closely related) have a basin-specific

occurrence (Boulart et al., 2022; Poitrimol et al., 2022, Diaz-Recio Lorenzo et al. 2024).

Beyond the idea that we did not examine rare and occasional species, our comparative

phylogeographic study also has some limitations. First, our sampling design, although quite

extensive, focused on a fraction of the existing hydrothermal vents, lacking fine-grained

representation within and across BABs. This limitation is common in deep-sea research, where

potential habitats and populations are known or suspected but remain unsampled. In our

case, the study area comprised a few unsampled - documented vents (e.g. Nifonea in

Vanuatu), and other 'ghost' undiscovered vent sites possibly located on seamounts along

volcanic arcs. Additional data from these sites should not affect our main conclusions, but

they would provide useful information to refine our patterns of population connectivity and

the timing of contact zones. Second, we generated ddRAD datasets across a wide range of

species with significant divergence between geographic lineages. These varying contexts of

divergence may have influenced our results, as divergent lineages may share fewer ddRAD loci

due to allele dropout and preferential bias towards less divergent loci to avoid missing data,

although the number of remaining loci was still very high.

Implications for conservation and future directions

As previously shown, cases that correspond to geographically separated cryptic species need

to be managed separately (e.g. *L. schrolli/L. fijiensis* or *S. tollmanni*). Other species depict much

lower divergence but with some variation in population differentiation. Although sporadic and

possibly rare, there is now clear evidence of present-day genetic connectivity between the

Western and Eastern metapopulations with an apparent high genetic homogeneity within

each of them. If larval exchange does not seem to be limited within each geographic group,

the inter-metapopulation migration rate is probably too low to allow any regional rescue

effect. Discussing effective migration and thus dispersal is however still quite challenging

(Lowe & Allendorf, 2010; Waples & Gaggiotti, 2006). Recent studies based on simulations of

particle dispersion at different depths (500 and 1000 meters) for *I. nautilei* and *Alviniconcha*

spp. suggested low but possible migration between BABs depending on the Pelagic Larval

Duration (PLD) and depth (Breusing et al., 2021; Mitarai et al., 2016). However, based on our

finding any rescue effect is likely to be basin-specific and highly sensitive to increasing local

extinctions magnified by deep-sea mining.

Most of our knowledge on the stability of vent ecosystems through time is derived from times

series established on the East Pacific Rise, a fast-spreading mid-oceanic ridge with a one-

dimensional stepping-stone axis of colonization (Dupreez & Fisher 2018, Audzijonyte &

Vrijenhoek 2011) and, some punctual physical barriers to dispersal (Plouviez et al. 2009, 2010,

2013). There, the fast extinction and recolonization rates of active sites are likely to select for

species, which can disperse far, and grow and reproduce fast. In back-arc basins, the ridge

spreading rate is rather low but varies between basins (Dick 2019). Extinction and

recolonization events are likely less common, which led to concerns about the ability of the

populations to recover if the metal sulfide deposits formed by the hydrothermal vent activity

are mined (Dupreez & Fisher 2018). Within each of the two metapopulations, high genetic

homogeneity of local populations can arise from either a substantial population size mitigating

genetic drift or the presence of a sufficient number of migrants exchanged within BABs.

Consequently, dispersal seems to be effective between nearby sites at the scale of either the

Western or Eastern regions, but much more limited between them. Because BAB zones are

spatially limited with a restricted number of active vent sites, mining the already known sites

should compromise any local 'rescue' effect. The fact that introgressed alleles between

metapopulations appear capable of reaching the Woodlark and North Fiji BABs only, suggests

that the inter-basin dispersal will not compensate for population bottlenecks within each

metapopulation. As shown by Bailleul et al. (2018), the apparent genetic homogeneity of

demes within a metapopulation is often maintained by a limited genetic drift, even in the face

of low inter-deme migration rates.

Conclusion

We have identified a strong phylogeographic break for several hydrothermal species of the

Southwest Pacific back-arc basins, consistent with a common suture zone between the

Woodlark and Lau basins. Although sharing a common pattern of population structuring,

variability in the degree of population differentiation is observed among these species,

potentially related to life history traits and species-specific demographic histories.

Our initial hypothesis that hydrothermal species of the Southwestern Pacific back-arc basins

may have evolved toward long-distance dispersal strategies to cross over non-hydrothermal

abyssal zones associated with a discontinuous system of oceanic ridges may only be valid for

a few species. In fact, the western Pacific vent communities are composed of species with

highly contrasted life-history traits and dispersal strategies, which are likely to promote the

good replenishment of populations locally in spite of vent instability, fragmentation and

geological discontinuities. Although some uncertainty remains about the timing of divergence

and secondary contacts, connectivity patterns between the two geographic groups are similar

among species, with asymmetric but bidirectional gene flow favoring the Western direction

(with the exception of *L. schrolli/L. fijiensis* for which a Manus origin of colonization is

suspected). This study clearly identified the existence of genetic barriers at some intermediate

locations that are likely to slow down gene flow for some species. It also confirmed that most

of the vent assemblages sustainability rather depends on the strength of the network of local

populations that makes up each regional metapopulation, rather than long distance dispersal. Ensuring the resilience of these communities requires sustainable management of their populations at the level of each biogeographic unit or back-arc basin, bearing in mind that the majority of current genetic exchange between the Eastern and Western basins are more specifically redirected towards the Manus Basin.

Acknowledgements

The research was funded by the ANR CERBERUS project (ANR-17-CE02-0003). We would like to thank the captains and crews of the French research vessel *L'Atalante* and the ROV *Victor* team for the CHUBACARC cruise (Hourdez and Jollivet, 2019, https://doi.org/10.17600/18001111). Sampling would not have been possible without their dedication. Shiptime and scientist travels were supported by the Flotte Océanographique Française (FOF) and the Centre National de la Recherche Scientifique (CNRS). This work benefited from access to the Genomer and ABIMS platforms part of the Biogenouest network, at Station Biologique de Roscoff, an EMBRC-France and EMBRC-ERIC site. We also warmly thank Cindy L. Van Dover for sharing with us some of her Manus 2009 samples in order to perform preliminary ddRAD libraries over the different species to choose restriction enzymes and check the number of usable loci.

Author contributions

Stephane Hourdez and Didier Jollivet designed the CHUBACARC and CERBERUS projects, François Bonhomme supervised the genetic work. Adrien Tran Lu Y, Stéphanie Ruault, Claire Daguin-Thiébaut, Anne-Sophie le Port, Marion Ballenghein, Sophie Arnaud-Haond, Jade Castel, Camille Poitrimol, Eric Thiébaut, François Lallier, Thomas Broquet, François Bonhomme, Didier Jollivet and Stéphane Hourdez performed laboratory work. Adrien Tran Lu Y performed bioinformatic statistical analyses with the contribution of François Bonhomme, Didier Jollivet, Pierre-Alexandre Gagnaire, Nicolas Bierne and Thomas Broquet. Adrien Tran Lu Y, François Bonhomme, Thomas Broquet, Didier Jollivet and Stephane Hourdez wrote the manuscript with feedback and inputs from all authors. All authors approved the manuscript

Conflict of interest

The authors have no conflicts of interest.

Data availability statement

Raw sequence reads (Individual fastq files) are available at the European Nucleotide Archive (bioproject PRJEB47533; *I. nautilei*) and the NCBI sequence read archive (PRJNA768636 for A. kojimai; PRJNA779874 for *L. schrolli*; PRJNA772682 for *S. tollmanni*; PRJNA1044574 for *B. manusensis*; PRJNA1030156 for *E. ohtai*; PRJNA1044042 for *B. segonzaci*). Metadata relative to the samples are also available with Biosamples accessions and linked to the sequence reads accessions. Scripts used in this study (R, $\partial a \partial$i) are available on a public Github repository: (https://github.com/Atranluy/Scripts-Ifremeria). VCFs and associated metadata will be available on public repository upon peer-review and publication.

Benefit-sharing statement

In order to obtain the requested authorizations to work in national waters and in agreement with the Nagoya protocol, we contacted the authorities of the different countries (Papua-New Guinea, Fiji, and Tonga) and territories (Wallis and Futuna) for benefit sharing where sampling was performed. The data generated will be accessible on public databases (see above). The results obtained will also be communicated to these authorities which may have to make decisions regarding conservation of deep-sea hydrothermal vent communities in their EEZs. Observers for the different countries who took part in the on-board activities will be informed of our findings.

Bibliography

Alexander, D. H., & Lange, K. (2011). Enhancements to the ADMIXTURE algorithm for individual ancestry estimation. *BMC Bioinformatics*, *12*(1), 246. https://doi.org/10.1186/1471-2105-12-246

Arellano, S. M., & Young, C. M. (2009). Spawning, Development, and the Duration of Larval Life in a Deep-Sea Cold-Seep Mussel. *The Biological Bulletin*, *216*(2), 149-162. https://doi.org/10.1086/BBLv216n2p149

Audzijonyte, A., & Vrijenhoek, R. C. (2010). When Gaps Really Are Gaps : Statistical Phylogeography of Hydrothermal Vent Invertebrates. *Evolution*, *64*(8), 2369-2384. https://doi.org/10.1111/j.1558-5646.2010.00987.x

Avise, J. C. (2000). *Phylogeography : The History and Formation of Species*. Harvard University
Press. https://doi.org/10.2307/j.ctv1nzfgj7

Avise, J. C. (2009). Phylogeography : Retrospect and prospect. *Journal of Biogeography*, *36*(1),
3-15. https://doi.org/10.1111/j.1365-2699.2008.02032.x

Avise, J. C., Arnold, J., Ball, R. M., Bermingham, E., Lamb, T., Neigel, J. E., Reeb, C. A., &
Saunders, N. C. (1987). INTRASPECIFIC PHYLOGEOGRAPHY : The Mitochondrial DNA Bridge
Between Population Genetics and Systematics. *Annual Review of Ecology and Systematics*,
18(1), 489-522. https://doi.org/10.1146/annurev.es.18.110187.002421

Bachraty, C., Legendre, P., & Desbruyères, D. (2009). Biogeographic relationships among deep-
sea hydrothermal vent faunas at global scale. *Deep Sea Research Part I: Oceanographic*
Research Papers, *56*(8), 1371-1378. https://doi.org/10.1016/j.dsr.2009.01.009

Bailleul, D., Mackenzie, A., Sacchi, O., Poisson, F., Bierne, N., & Arnaud-Haond, S. (2018). Large-
scale genetic panmixia in the blue shark (*Prionace glauca*) : A single worldwide population, or
a genetic lag-time effect of the "grey zone" of differentiation? *Evolutionary Applications*,
11(5), 614-630. https://doi.org/10.1111/eva.12591

Barrows, T. T., Hope, G. S., Prentice, M. L., Fifield, L. K., & Tims, S. G. (2011). Late Pleistocene
glaciation of the Mt Giluwe volcano, Papua New Guinea. *Quaternary Science Reviews*,
30(19-20), 2676-2689. https://doi.org/10.1016/j.quascirev.2011.05.022

Barton, N., & Bengtsson, B. O. (1986). The barrier to genetic exchange between hybridising
populations. *Heredity*, *57*(3), Article 3. https://doi.org/10.1038/hdy.1986.135

Beaulieu, S. E., Baker, E. T., German, C. R., & Maffei, A. (2013). An authoritative global database
for active submarine hydrothermal vent fields. *Geochemistry, Geophysics, Geosystems*,
14(11), 4892-4905. https://doi.org/10.1002/2013GC004998

Bierne, N., Welch, J., Loire, E., Bonhomme, F., & David, P. (2011). The coupling hypothesis :
Why genome scans may fail to map local adaptation genes. *Molecular Ecology*, *20*(10),
2044-2072. https://doi.org/10.1111/j.1365-294X.2011.05080.x

Breusing, C., Johnson, S. B., Mitarai, S., Beinart, R. A., & Tunnicliffe, V. (2021). Differential
patterns of connectivity in Western Pacific hydrothermal vent metapopulations : A

comparison of biophysical and genetic models. *Evolutionary Applications*.
https://doi.org/10.1111/eva.13326

Carver, R., Childs, J., Steinberg, P., Mabon, L., Matsuda, H., Squire, R., McLellan, B., & Esteban,
M. (2020). A critical social perspective on deep sea mining : Lessons from the emergent
industry in Japan. *Ocean & Coastal Management*, *193*, 105242.
https://doi.org/10.1016/j.ocecoaman.2020.105242

Castel, J., Hourdez, S., Pradillon, F., Daguin-Thiébaut, C., Ballenghien, M., Ruault, S., Corre, E.,
Tran Lu Y, A., Mary, J., Gagnaire, P.-A., Bonhomme, F., Breusing, C., Broquet, T., & Jollivet, D.
(2022). Inter-Specific Genetic Exchange Despite Strong Divergence in Deep-Sea Hydrothermal
Vent Gastropods of the Genus Alviniconcha. *Genes*, *13*(6), Article 6.
https://doi.org/10.3390/genes13060985

Chen, C., & Sigwart, J. D. (2023). The lost vent gastropod species of Lothar A. Beck. *Zootaxa*,
5270(3), Article 3. https://doi.org/10.11646/zootaxa.5270.3.2

Chevaldonné, P., Jollivet, D., Desbruyères, D., Lutz, R., & Vrijenhoek, R. (2002). Sister-species
of eastern Pacific hydrothermal vent worms (Ampharetidae, Alvinellidae, Vestimentifera)
provide new mitochondrial COI clock calibration. *CBM - Cahiers de Biologie Marine*, *43*(3-4),
367-370. https://archimer.ifremer.fr/doc/00000/895/

Daguin-Thiébaut, C., Ruault, S., Roby, C., Broquet, T., Viard, F., & Brelsford, A. (2021,
septembre 21). *Construction of individual ddRAD libraries*. Protocols.Io.
https://www.protocols.io/view/construction-of-individual-ddrad-libraries-bv4tn8wn

De Jode, A., Le Moan, A., Johannesson, K., Faria, R., Stankowski, S., Westram, A. M., Butlin, R.
K., Rafajlović, M., & Fraïsse, C. (2023). Ten years of demographic modeling of divergence and
speciation in the sea. *Evolutionary Applications*, *16*(2), 542-559.
https://doi.org/10.1111/eva.13428

Desbruyères, D., Hashimoto, J., & Fabri, M.-C. (2006). Composition and Biogeography of
Hydrothermal Vent Communities in Western Pacific Back-Arc Basins. In *Back-Arc Spreading*
Systems : Geological, Biological, Chemical, and Physical Interactions (p. 215-234). American
Geophysical Union (AGU). https://doi.org/10.1029/166GM11

Diaz-Recio Lorenzo, C., Tran Lu Y, A., Brunner, O., Arbizu, P. M., Jollivet, D., Laurent, S., &
Gollner, S. (2024). Highly structured populations of copepods at risk to deep-sea mining :
Integration of genomic data with demogenetic and biophysical modelling. *Molecular Ecology*,
33(9), e17340. https://doi.org/10.1111/mec.17340

Dick, G. J. (2019). The microbiomes of deep-sea hydrothermal vents : Distributed globally,
shaped locally. Nature Reviews Microbiology, 17(5), 271-283.
https://doi.org/10.1038/s41579-019-0160-2

Du Preez, C., & Fisher, C. R. (2018). Long-Term Stability of Back-Arc Basin Hydrothermal Vents.
Frontiers in Marine Science, *5*. https://doi.org/10.3389/fmars.2018.00054

Dupoué, A., Trochet, A., Richard, M., Sorlin, M., Guillon, M., Teulieres-Quillet, J., Vallé, C.,
Rault, C., Berroneau, M., Berroneau, M., Lourdais, O., Blaimont, P., Bertrand, R., Pottier, G.,
Calvez, O., Guillaume, O., Le Chevalier, H., Souchet, J., Le Galliard, J.-F., … Aubret, F. (2021).
Genetic and demographic trends from rear to leading edge are explained by climate and forest
cover in a cold-adapted ectotherm. *Diversity and Distributions*, *27*(2), 267-281.
https://doi.org/10.1111/ddi.13202

Edwards, S. V., Robin, V. V., Ferrand, N., & Moritz, C. (2022). The Evolution of Comparative
Phylogeography : Putting the Geography (and More) into Comparative Population Genomics.
Genome Biology and Evolution, *14*(1), evab176. https://doi.org/10.1093/gbe/evab176

Ewing, G. B., & Jensen, J. D. (2016). The consequences of not accounting for background
selection in demographic inference. *Molecular Ecology*, *25*(1), 135-141.
https://doi.org/10.1111/mec.13390

Excoffier, L., & Lischer, H. E. L. (2010). Arlequin suite ver 3.5 : A new series of programs to
perform population genetics analyses under Linux and Windows. *Molecular Ecology*
Resources, *10*(3), 564-567. https://doi.org/10.1111/j.1755-0998.2010.02847.x

Faure, B., Jollivet, D., Tanguy, A., Bonhomme, F., & Bierne, N. (2009). Speciation in the Deep
Sea : Multi-Locus Analysis of Divergence and Gene Flow between Two Hybridizing Species of
Hydrothermal Vent Mussels. *PLOS ONE*, *4*(8), e6485.
https://doi.org/10.1371/journal.pone.0006485

Gagnaire, P.-A. (2020). Comparative genomics approach to evolutionary process connectivity.

Evolutionary Applications, *13*(6), 1320-1334. https://doi.org/10.1111/eva.12978

Gena, K. (2013). Deep Sea Mining of Submarine Hydrothermal Deposits and its Possible

Environmental Impact in Manus Basin, Papua New Guinea. *Procedia Earth and Planetary*

Science, *6*, 226-233. https://doi.org/10.1016/j.proeps.2013.01.031

Gutenkunst, R. N., Hernandez, R. D., Williamson, S. H., & Bustamante, C. D. (2009). Inferring

the Joint Demographic History of Multiple Populations from Multidimensional SNP Frequency

Data. *PLOS Genetics*, *5*(10), e1000695. https://doi.org/10.1371/journal.pgen.1000695

Hadley Wickham. (2016). *Ggplot2* (Springer-Verlag New York). https://ggplot2.tidyverse.org

Hamilton, W. D., & May, R. M. (1977). Dispersal in stable habitats. *Nature*, *269*(5629), 578-581.

https://doi.org/10.1038/269578a0

Hickerson, M. J., Carstens, B. C., Cavender-Bares, J., Crandall, K. A., Graham, C. H., Johnson, J.

B., Rissler, L., Victoriano, P. F., & Yoder, A. D. (2010). Phylogeography's past, present, and

future : 10 years after Avise, 2000. *Molecular Phylogenetics and Evolution*, *54*(1), 291-301.

https://doi.org/10.1016/j.ympev.2009.09.016

Hourdez, S., & Jollivet, D. (2019). *CHUBACARC Cruise Report*.

https://doi.org/10.17600/18001111

Hourdez, S., & Jollivet, D. (2020). Chapter 2. In *Metazoan adaptation to deep-sea*

hydrothermal vents (G. di Prisco, A. Huiskes and C. Ellis-Evanz (eds.), p. 42-67). Ecological

reviews, Cambridge Univ. Press.

https://books.google.com/books?hl=en&lr=&id=YtX7DwAAQBAJ&oi=fnd&pg=PA42&dq=info

:isvZ-xX6CFwJ:scholar.google.com&ots=Iv3McwTtw5&sig=3I841J3xRdhy_EzuiOlrPwbZgSA

Hurtado, L. A., Lutz, R. A., & Vrijenhoek, R. C. (2004). Distinct patterns of genetic differentiation

among annelids of eastern Pacific hydrothermal vents. *Molecular Ecology*, *13*(9), 2603-2615.

https://doi.org/10.1111/j.1365-294X.2004.02287.x

Johnson, S. B., Won, Y.-J., Harvey, J. B., & Vrijenhoek, R. C. (2013). A hybrid zone between

Bathymodiolus mussel lineages from eastern Pacific hydrothermal vents. *BMC Evolutionary*

Biology, *13*(1), 21. https://doi.org/10.1186/1471-2148-13-21

Johnson, S. B., Young, C. R., Jones, W. J., Warén, A., & Vrijenhoek, R. C. (2006). Migration,
Isolation, and Speciation of Hydrothermal Vent Limpets (*Gastropoda; Lepetodrilidae*) Across
the Blanco Transform Fault. *The Biological Bulletin*, *210*(2), 140-157.
https://doi.org/10.2307/4134603

Keenan, K., McGinnity, P., Cross, T. F., Crozier, W. W., & Prodöhl, P. A. (2013). diveRsity : An R
package for the estimation and exploration of population genetics parameters and their
associated errors. *Methods in Ecology and Evolution*, *4*(8), 782-788.
https://doi.org/10.1111/2041-210X.12067

Kim, M., Kang, J.-H., and Kim, D. (2022). Holoplanktonic and meroplanktonic larvae in the
surface waters of the Onnuri vent field in the Central Indian Ridge. Journal of Marine Science
and Engineering, 10(2), 158. https://doi.org/10.3390/jmse10020158

Lee, W.-K., Kim, S.-J., Hou, B. K., Dover, C. L. V., & Ju, S.-J. (2019). Population genetic
differentiation of the hydrothermal vent crab *Austinograea alayseae* (*Crustacea :
Bythograeidae*) in the Southwest Pacific Ocean. *PLOS ONE*, *14*(4), e0215829.
https://doi.org/10.1371/journal.pone.0215829

Levin, L. A. (1990). A review of methods for labeling and tracking marine invertebrate larvae.
Ophelia, *32*(1-2), 115-144. https://doi.org/10.1080/00785236.1990.10422028

Lowe, W. H., & Allendorf, F. W. (2010). What can genetics tell us about population
connectivity? *Molecular Ecology*, *19*(15), 3038-3051. https://doi.org/10.1111/j.1365-
294X.2010.04688.x

Lutz, R. A., Shank, T. M., Fornari, D. J., Haymon, R. M., Lilley, M. D., Von Damm, K. L., &
Desbruyeres, D. (1994). Rapid growth at deep-sea vents. *Nature*, *371*(6499), 663-664.
https://doi.org/10.1038/371663a0

Mastretta-Yanes, A., Arrigo, N., Alvarez, N., Jorgensen, T. H., Piñero, D., & Emerson, B. C.
(2015). Restriction site-associated DNA sequencing, genotyping error estimation and de novo
assembly optimization for population genetic inference. *Molecular Ecology Resources*, *15*(1),
28-41. https://doi.org/10.1111/1755-0998.12291

Matabos, M., & Jollivet, D. (2019). Revisiting the *Lepetodrilus elevatus* species complex
(*Vetigastropoda : Lepetodrilidae*), using samples from the Galápagos and Guaymas
hydrothermal vent systems. *Journal of Molluscan Studies*, *85*(1), 154-165.
https://doi.org/10.1093/mollus/eyy061

Matabos, M., Plouviez, S., Hourdez, S., Desbruyères, D., Legendre, P., Warén, A., Jollivet, D., &
Thiébaut, E. (2011). Faunal changes and geographic crypticism indicate the occurrence of a
biogeographic transition zone along the southern East Pacific Rise. *Journal of Biogeography*,
38(3), 575-594. https://doi.org/10.1111/j.1365-2699.2010.02418.x

Matute, D. R., Butler, I. A., Turissini, D. A., & Coyne, J. A. (2010). A Test of the Snowball Theory
for the Rate of Evolution of Hybrid Incompatibilities. *Science*, *329*(5998), 1518-1521.
https://doi.org/10.1126/science.1193440

McPeek, M. A., & Holt, R. D. (1992). The Evolution of Dispersal in Spatially and Temporally
Varying Environments. *The American Naturalist*, *140*(6), 1010-1027.
https://doi.org/10.1086/285453

Mitarai, S., Watanabe, H., Nakajima, Y., Shchepetkin, A. F., & McWilliams, J. C. (2016).
Quantifying dispersal from hydrothermal vent fields in the western Pacific Ocean. *Proceedings*
of the National Academy of Sciences, *113*(11), 2976-2981.
https://doi.org/10.1073/pnas.1518395113

Moalic, Y., Desbruyères, D., Duarte, C. M., Rozenfeld, A. F., Bachraty, C., & Arnaud-Haond, S.
(2012). Biogeography Revisited with Network Theory : Retracing the History of Hydrothermal
Vent Communities. *Systematic Biology*, *61*(1), 127-127.
https://doi.org/10.1093/sysbio/syr088

Mouchi, V., Pecheyran, C., Claverie, F., Cathalot, C., Matabos, M., Germain, Y., Rouxel, O.,
Jollivet, D., Broquet, T., & Comtet, T. (2024). A step towards measuring connectivity in the
deep sea : Elemental fingerprints of mollusk larval shells discriminate hydrothermal-vent sites.
Biogeosciences, *21*(1), 145-160. https://doi.org/10.5194/bg-21-145-2024

Nei, M., & Li, W. H. (1979). Mathematical model for studying genetic variation in terms of
restriction endonucleases. *Proceedings of the National Academy of Sciences*, *76*(10),
5269-5273. https://doi.org/10.1073/pnas.76.10.5269

Niner, H. J., Ardron, J. A., Escobar, E. G., Gianni, M., Jaeckel, A., Jones, D. O. B., Levin, L. A., Smith, C. R., Thiele, T., Turner, P. J., Van Dover, C. L., Watling, L., & Gjerde, K. M. (2018). Deep-Sea Mining With No Net Loss of Biodiversity—An Impossible Aim. *Frontiers in Marine Science*, *5*. https://www.frontiersin.org/article/10.3389/fmars.2018.00053

Papadopoulou, A., & Knowles, L. L. (2016). Toward a paradigm shift in comparative phylogeography driven by trait-based hypotheses. *Proceedings of the National Academy of Sciences*, *113*(29), 8018-8024. https://doi.org/10.1073/pnas.1601069113

Paris, J. R., Stevens, J. R., & Catchen, J. M. (2017). Lost in parameter space : A road map for stacks. *Methods in Ecology and Evolution*, *8*(10), 1360-1373. https://doi.org/10.1111/2041-210X.12775

Pickrell, J. K., & Pritchard, J. K. (2012). Inference of Population Splits and Mixtures from Genome-Wide Allele Frequency Data. *PLOS Genetics*, *8*(11), e1002967. https://doi.org/10.1371/journal.pgen.1002967

Plouviez, S., Faure, B., Guen, D. L., Lallier, F. H., Bierne, N., & Jollivet, D. (2013). A new barrier to dispersal trapped old genetic clines that escaped the easter microplate tension zone of the Pacific vent mussels. *PLOS ONE*, *8*(12), e81555. https://doi.org/10.1371/journal.pone.0081555

Plouviez, S., LaBella, A. L., Weisrock, D. W., Meijenfeldt, F. A. B. von, Ball, B., Neigel, J. E., & Dover, C. L. V. (2019). Amplicon sequencing of 42 nuclear loci supports directional gene flow between South Pacific populations of a hydrothermal vent limpet. *Ecology and Evolution*, *9*(11), 6568-6580. https://doi.org/10.1002/ece3.5235

Plouviez, S., Le Guen, D., Lecompte, O., Lallier, F. H., & Jollivet, D. (2010). Determining gene flow and the influence of selection across the equatorial barrier of the East Pacific Rise in the tube-dwelling polychaete *Alvinella pompejana*. *BMC Evolutionary Biology*, *10*(1), 220. https://doi.org/10.1186/1471-2148-10-220

Plouviez, S., Shank, T. M., Faure, B., Daguin-Thiebaut, C., Viard, F., Lallier, F. H., & Jollivet, D. (2009). Comparative phylogeography among hydrothermal vent species along the East Pacific Rise reveals vicariant processes and population expansion in the South. *Molecular Ecology*, *18*(18), 3903-3917. https://doi.org/10.1111/j.1365-294X.2009.04325.x

Poitrimol, C., Thiébaut, É., Daguin-Thiébaut, C., Port, A.-S. L., Ballenghien, M., Y, A. T. L., Jollivet, D., Hourdez, S., & Matabos, M. (2022). Contrasted phylogeographic patterns of hydrothermal vent gastropods along South West Pacific : Woodlark Basin, a possible contact zone and/or stepping-stone. *PLOS ONE*, *17*(10), e0275638. https://doi.org/10.1371/journal.pone.0275638

Poitrimol, C., Matabos, M., Veuillot, A., Ramière, A., Comtet, T., Boulart, C., Cathalot, C., & Thiébaut, É. (2024). Reproductive biology and population structure of three hydrothermal gastropods (*Lepetodrilus schrolli, L. fijiensis* and *Shinkailepas tollmanni*) from the South West Pacific back-arc basins. *Marine Biology*, *171*(1), 31. https://doi.org/10.1007/s00227-023-04348-4

R Core Team. (2023). https://www.R-project.org/

Ravinet, M., Faria, R., Butlin, R. K., Galindo, J., Bierne, N., Rafajlović, M., Noor, M. a. F., Mehlig, B., & Westram, A. M. (2017). Interpreting the genomic landscape of speciation : A road map for finding barriers to gene flow. *Journal of Evolutionary Biology*, *30*(8), 1450-1477. https://doi.org/10.1111/jeb.13047

Rochette, N. C., Rivera-Colón, A. G., & Catchen, J. M. (2019). Stacks 2 : Analytical methods for paired-end sequencing improve RADseq-based population genomics. *Molecular Ecology*, *28*(21), 4737-4754. https://doi.org/10.1111/mec.15253

Rougeux, C., Bernatchez, L., & Gagnaire, P.-A. (2017). Modeling the Multiple Facets of Speciation-with-Gene-Flow toward Inferring the Divergence History of Lake Whitefish Species Pairs (*Coregonus clupeaformis*). *Genome Biology and Evolution*, *9*(8), 2057-2074. https://doi.org/10.1093/gbe/evx150

Schellart, W. P., Lister, G. S., & Toy, V. G. (2006). A Late Cretaceous and Cenozoic reconstruction of the Southwest Pacific region : Tectonics controlled by subduction and slab rollback processes. *Earth-Science Reviews*, *76*(3), 191-233. https://doi.org/10.1016/j.earscirev.2006.01.002

Schöne, B. R., & Giere, O. (2005). Growth increments and stable isotope variation in shells of the deep-sea hydrothermal vent bivalve mollusk *Bathymodiolus brevior* from the North Fiji

Basin, Pacific Ocean. *Deep Sea Research Part I: Oceanographic Research Papers*, *52*(10),

1896-1910. https://doi.org/10.1016/j.dsr.2005.06.003

Shank, T. M., Fornari, D. J., Von Damm, K. L., Lilley, M. D., Haymon, R. M., & Lutz, R. A. (1998).

Temporal and spatial patterns of biological community development at nascent deep-sea

hydrothermal vents (9°50′N, East Pacific Rise). *Deep Sea Research Part II: Topical Studies in*

Oceanography, *45*(1), 465-515. https://doi.org/10.1016/S0967-0645(97)00089-1

Sommer, S. A., Van Woudenberg, L., Lenz, P. H., Cepeda, G., & Goetze, E. (2017). Vertical

gradients in species richness and community composition across the twilight zone in the North

Pacific   Subtropical   Gyre.   *Molecular   Ecology*,   *26*(21),   6136-6156.

https://doi.org/10.1111/mec.14286

Sundqvist, L., Keenan, K., Zackrisson, M., Prodöhl, P., & Kleinhans, D. (2016). Directional

genetic differentiation and relative migration. *Ecology and Evolution*, *6*(11), 3461-3475.

https://doi.org/10.1002/ece3.2096

Teixeira, S., Serrão, E. A., & Arnaud-Haond, S. (2012). Panmixia in a Fragmented and Unstable

Environment : The Hydrothermal Shrimp *Rimicaris exoculata* Disperses Extensively along the

Mid-Atlantic Ridge. *PLOS ONE*, *7*(6), e38521. https://doi.org/10.1371/journal.pone.0038521

Thaler, A. D., Plouviez, S., Saleu, W., Alei, F., Jacobson, A., Boyle, E. A., Schultz, T. F., Carlsson,

J., & Dover, C. L. V. (2014). Comparative Population Structure of Two Deep-Sea Hydrothermal-

Vent-Associated Decapods (*Chorocaris sp. 2* and *Munidopsis lauensis*) from Southwestern

Pacific   Back-Arc   Basins.   *PLOS   ONE*,   *9*(7),   e101345.

https://doi.org/10.1371/journal.pone.0101345

Thaler, A. D., Zelnio, K., Saleu, W., Schultz, T. F., Carlsson, J., Cunningham, C., Vrijenhoek, R. C.,

& Van Dover, C. L. (2011). The spatial scale of genetic subdivision in populations of *Ifremeria*

nautilei, a hydrothermal-vent gastropod from the southwest Pacific. *BMC Evolutionary*

Biology, *11*(1), 372. https://doi.org/10.1186/1471-2148-11-372

Tran Lu Y, A., Ruault, S., Daguin-Thiébaut, C., Castel, J., Bierne, N., Broquet, T., Wincker, P.,

Perdereau, A., Arnaud-Haond, S., Gagnaire, P.-A., Jollivet, D., Hourdez, S., & Bonhomme, F.

(2022). Subtle limits to connectivity revealed by outlier loci within two divergent

metapopulations of the deep-sea hydrothermal gastropod *Ifremeria nautilei*. *Molecular Ecology*, *31*(10), 2796-2813. https://doi.org/10.1111/mec.16430

Tunnicliffe, V. (1992). The Nature and Origin of the Modern Hydrothermal Vent Fauna. *PALAIOS*, *7*(4), 338-350. https://doi.org/10.2307/3514820

Tunnicliffe, V., Chen, C., Giguère, T., Rowden, A. A., Watanabe, H. K., & Brunner, O. (2024). Hydrothermal vent fauna of the western Pacific Ocean : Distribution patterns and biogeographic networks. *Diversity and Distributions*, *30*(2), e13794. https://doi.org/10.1111/ddi.13794

Van Dover, C. L. (2011). Mining seafloor massive sulphides and biodiversity : What is at risk? *ICES Journal of Marine Science*, *68*(2), 341-348. https://doi.org/10.1093/icesjms/fsq086

Van Dover, C. L., Ardron, J. A., Escobar, E., Gianni, M., Gjerde, K. M., Jaeckel, A., Jones, D. O. B., Levin, L. A., Niner, H. J., Pendleton, L., Smith, C. R., Thiele, T., Turner, P. J., Watling, L., & Weaver, P. P. E. (2017). Biodiversity loss from deep-sea mining. *Nature Geoscience*, *10*(7), Article 7. https://doi.org/10.1038/ngeo2983

Vrijenhoek, R. C. (2010). Genetic diversity and connectivity of deep-sea hydrothermal vent metapopulations. *Molecular Ecology*, *19*(20), 4391-4411. https://doi.org/10.1111/j.1365-294X.2010.04789.x

Waples, R. S., & Gaggiotti, O. (2006). INVITED REVIEW : What is a population? An empirical evaluation of some genetic methods for identifying the number of gene pools and their degree of connectivity. *Molecular Ecology*, *15*(6), 1419-1439. https://doi.org/10.1111/j.1365-294X.2006.02890.x

Warèn, A., & Bouchet, P. (1993). New records, species, genera, and a new family of gastropods from hydrothermal vents and hydrocarbon seeps*. *Zoologica Scripta*, *22*(1), 1-90. https://doi.org/10.1111/j.1463-6409.1993.tb00342.x

Washburn, T. W., Turner, P. J., Durden, J. M., Jones, D. O. B., Weaver, P., & Van Dover, C. L. (2019). Ecological risk assessment for deep-sea mining. *Ocean & Coastal Management*, *176*, 24-39. https://doi.org/10.1016/j.ocecoaman.2019.04.014

Yahagi, T., Fukumori, H., Warén, A., & Kano, Y. (2019). Population connectivity of

hydrothermal-vent limpets along the northern Mid-Atlantic Ridge (*Gastropoda :*

Neritimorpha: Phenacolepadidae). *Journal of the Marine Biological Association of the United*

Kingdom, *99*(1), 179-185. https://doi.org/10.1017/S0025315417001898

Yahagi, T., Thaler, A. D., Dover, C. L. V., & Kano, Y. (2020). Population connectivity of the

hydrothermal-vent limpet *Shinkailepas tollmanni* in the Southwest Pacific (*Gastropoda :*

Neritimorpha: Phenacolepadidae). *PLoS One*, *15*(9), e0239784.

https://doi.org/10.1371/journal.pone.0239784

Young, C. M. (1994). A tale of two dogmas: the early history of deep-sea reproductive biology.

Reproduction, larval biology, and recruitment in the deep-sea benthos. Columbia University

Press, New York, 1-25.

Zheng, X., Levine, D., Shen, J., Gogarten, S. M., Laurie, C., & Weir, B. S. (2012). A high-

performance computing toolset for relatedness and principal component analysis of SNP data.

Bioinformatics, *28*(24), 3326-3328. https://doi.org/10.1093/bioinformatics/bts606