Some like it hot: efficiency of the type III secretion system and

activation of plant defenses have multiple thermosensitive

behaviors in the *Pseudomonas syringae* complex

E. Caullireau[1,2], D. Danzi[1], V.M. Tempo[1], M. Pandolfo[1], C.E. Morris[2], E. Vandelle[1*]

[1] Department of Biotechnology, University of Verona, Verona, Italy.

[2] UR0407 Plant Pathology, INRAE, Montfavet, France.

*Corresponding author:

Email: elodiegenevieve.vandelle@univr.it (EV)

Abstract

The *Pseudomonas syringae* species complex is a very important plant pathogenic bacterium,

causing damage and economic losses on many crops. Research efforts for the understanding of

the determinants of the host-range of a given strain are considerable. Recently, we showed that

the inability of a *P. syringae* pv. *actinidiae* strain to trigger ETI in *A. thaliana* is due to an inefficient

T3SS and not to the absence of a recognized effector. In this context, we compared several *P.*

syringae strains belonging to different phylogroups and carrying the same plasmid-born

avirulence gene for their ability to induce an HR in *A. thaliana* Col-0, as a marker of T3SS efficiency.

Pto DC3000 *AvrB* and *Pma* M6 *AvrB* consistently triggered a strong HR while other strains induced

it at different intensities significantly depending on temperature. Both behaviors of low and warm

temperature-dependency for T3SS efficiency were observed among thermosensitive strains,

irrespective of their *in-vitro* growth optimum. Surprisingly, differences were also observed among

quasi-clonal strains. These results reveal a strongly strain-specific regulatory role of temperature

in effector injection and reinforce the notion that the presence/absence of effectors is not

sufficient to predict the outcome of plant-bacteria interactions. Moreover, this work highlights the

necessity to study bacterial virulence in a broader set of strains insofar as *Pto* DC3000 is a reliable

model strain but not representative of the *P. syringae* complex.

Author summary

Pseudomonas syringae is a bacterium ubiquitous in the environment and responsible for

important crop losses worldwide. Moreover, *P. syringae* serves as a model organism for studying

plant-pathogen interactions. Bacteria rely on various virulence mechanisms to successfully infect

their host, one of the most important of which is the Type III Secretion System (T3SS). Although it

is acknowledged that environmental conditions play a significant role in the plant-pathogen

interactions, our understanding of their influence on both partners is still limited. Regarding

temperature, contrasting findings have been reported about its influence on T3SS efficiency. While

most efforts have focused on a few strains of *P. syringae*, the phylogenetic diversity of this

bacterium is enormous. Consequently, we decided to characterize the role of temperature on T3SS

efficiency in *P. syringae* in a broader way. Our results showed that bacterial behavior in response

to temperature is highly variable among strains, even among very closely related ones. Our study

provides new insights into the role of temperature in regulating bacterial virulence and further

evidence of the need to investigate the diversity of pathogenic organisms beyond model strains.

Introduction

Pseudomonas syringae is a gram-negative bacterium that poses a threat to global agricultural

production with new outbreaks that emerge regularly [1,2]. *P. syringae* has been divided into 13

phylogenetic groups (phylogroups, PG), further divided into 23 clades [3]. During the infection

process, *P. syringae* relies on an arsenal of secreted virulence factors, such as secretion systems,

quorum sensing (QS), production of phytotoxins (coronatine, syringomycin…), hormones,

exopolysaccharides (EPS), siderophores etc. [4,5]. *P. syringae* pathogenesis is predominantly

mediated by the type III secretion system (T3SS), a highly conserved needle-like specialized

protein secretion machinery, found in numerous gram-negative bacterial pathogens affecting

plants and animals. T3SS forms a continuous channel, guiding the translocation of diverse

proteinaceous effectors (T3SEs) from the bacterium directly into the host cell or apoplast [6–9].

T3SEs facilitate the invasion of plant host cells, suppress plant immunity, or interfere with essential

plant cellular processes [10–15]. *P. syringae* strains in the phylogroups with the canonical T3SS

commonly harbor a repertoire of approximately 15-30 T3SEs, playing distinct roles within the host

plant [16–21]. Of note, there is no direct relationship between the size or content of the T3SE

repertoire and host range [22,23]. When considered individually, numerous T3SEs are dispensable

for *P. syringae* virulence. However, collectively, they redundantly contribute to disarming plant

immunity [19,24].

Throughout their co-evolution with pathogens, plants have evolved an extensive array of defenses

to recognize them and block pathogenesis. Common bacterial features such as flagellin,

lipopolysaccharide (LPS), peptidoglycan and elongation factor Tu (EF-Tu) contain conserved

molecular motifs (Pathogen or Microbe-Associated Molecular Patterns; P/MAMPs) that are

sensed by the plant as non-self, through their binding to Pattern Recognition Receptors (PRRs)

situated on the plant cell surface [25,26]. Non-self detection by PRRs is the first step of the PTI or

MTI, for PAMP-/MAMP-Triggered Immunity, implying then various cellular responses to inhibit

pathogen growth and virulence mechanisms, such as T3SS activation for bacterial pathogens

[27,28]. On the other hand, T3SEs serve the primary function of subverting PTI, targeting key

signaling components to interfere with their function. In response, plants have evolved

cytoplasmic receptors called 'resistance' (R) proteins to detect some of these effectors, either

directly or indirectly, leading to the activation of the Effector-Triggered Immunity (ETI) [29]. R

genes typically encode for NLR proteins (Nucleotide-binding oligomerization domain-Like

Receptor), consisting of a nucleotide-binding site (NBS) and a leucine-rich repeat (LRR) domain

[30]. PTI and ETI are unlikely to be separate processes as suggested in the 'zigzag' model [31], but

both non-self and virulence effector recognition by membrane and cytoplasmic receptors are

simultaneously involved in a global surveillance system to respond effectively to pathogen attack

[32,33]. ETI is a potentiation of PTI, often leading to rapid and localized programmed cell death,

known as the hypersensitive response (HR). HR generally occurs within 2 to 6 hours post initial

host-pathogen contact, inhibiting the propagation of (hemi)-biotrophic pathogens, such as *P.*

syringae [34,35].

According to the 'disease triangle' paradigm [36], plant diseases arise from the interplay between a susceptible plant and a virulent pathogen, under conducive environmental conditions, resulting in a compatible reaction. Overall plant-pathogen interaction mechanisms are highly influenced by environmental conditions, which can either favor, have neutral effects, or hinder both partners [21,37]. Among the environmental factors, temperature has a regulatory role of virulence in plant-associated bacteria (reviewed in references [37,38]).

Within the *P. syringae* species complex, temperature variations impact various traits crucial for bacterial fitness, including phytotoxins, EPS biosynthesis, motility or quorum-sensing [38–52]. T3SS has also been investigated concerning its possible thermoregulation. In *P. syringae* pv. *syringae* and pv. *tomato* DC3000, the *in-vitro* secretion of HopPsyA and AvrPto effectors was impaired at temperatures above 22°C and 20°C, indicating some kind of low-temperature dependency for their T3SS efficiency [53]. In contrast, recent findings in strain DC3000 revealed that the translocation of T3SEs in *A. thaliana* cells is enhanced at 30°C compared to 23°C, indicating the possible T3SS efficiency even at elevated temperatures [54]. However, in tomato, the expression of T3SS-related genes *hrpL* and *hrpA* in this same strain was not affected by a temperature rise from 26°C to 31°C, although the expression of the avirulence gene *AvrPtoB* was significantly reduced [48]. On the other hand, for *P. syringae* pv. *actinidiae* strains, it was shown that there was no difference in *hrpA1* promoter activity *in vitro* across a temperature range of 18-28°C, nonetheless, substantial differences were evident at a functional level, demonstrating a low-temperature dependency for T3SS efficiency for these strains [55]. However, the complete implications of warm temperatures on *P. syringae* virulence in plants remain poorly understood [39,47]. Studies indicate complex interactions with plant immune signaling pathways,

emphasizing the need for further research to understand their implications in terms of disease

[54,56].

Thermosensing principles in phytopathogens are less understood than those of pathogens of

warm-blooded organisms. They were assumed to be different, given the ecological differences

between these groups of pathogens [38,57]. Nevertheless, some common mechanisms have been

identified in both plant- and mammal-associated bacteria. Various molecular strategies are

involved in sensing temperature, such as two-component systems (TCSs) that typically involve a

membrane-bound sensor histidine kinase that detects environmental stimuli and a cytoplasmic

response regulator influencing gene expression upon phosphorylation of an aspartic acid [57].

Several TCS were shown to be involved in the thermoregulation of virulence mechanisms, such as

the VirA/G system in *Agrobacterium tumefaciens* [58], the GacS/GacA system in *P. syringae* pv.

phaseolicola [59,60] and the CorRSP system in *P. syringae* pv. *glycinea* [61].

Evidence of the influence of temperature in regulating virulence factors in *P. syringae* species

complex has been documented, however, there have not been studies dedicated to comparing

strains under identical conditions. Consequently, we wanted to further understand the role of the

temperature on the function/activation dynamics of the T3SS, investigating a broader range of

diversity within the *P. syringae* species complex under comparable conditions. For this purpose,

we transformed 13 strains of *P. syringae* to carry the effector AvrB and infiltrated them into

Arabidopsis thaliana Col-0 leaf disks. In this model plant, AvrB is indirectly recognized by the

resistance protein RPM1, through RIN4 modification, which triggers HR [62–64]. Infiltrated leaf

disks were incubated at different temperatures and the quantification of electrolyte leakage

served as the indicator of HR onset. Conductivity increase was used as a proxy for T3SS efficiency,

and this allowed us to effectively discriminate strains based on a robust phenotype directly linked

to the activation of the T3SS. Based on the different behaviors we observed in response to

temperature, we investigated the genomes of the strains in an attempt to decipher the origin of

the thermosensing mechanisms.

134 Results

135 The intensity of HR induction is not strictly reproducible for individual

136 strains.

To evaluate T3SS functionality at different temperatures, *AvrB*-expressing *P. syringae* strains were

infiltrated into *A. thaliana* Col-0 leaf disks. As a T3SS-dependent mechanism, HR was quantified

by measuring the increase in ion leakage due to dying cells. Conductivity levels measured among

the diverse replicated experiments were highly variable. The conductivity value measured at 24

hours post-infection (hpi) for a given modality (*i.e.*, one strain infiltrated and incubated at one

temperature) doubled or tripled among replicated experiments (*e.g*. conductivity values ranged

from 80 µS.cm^{-1} to 226 µS.cm^{-1} for strain DC3000 *AvrB* when incubated at 18°C) (Fig 1). This

variability is illustrated in Fig 2 for the seven replicated experiments for strain USA007 *AvrB* at 18°C

(ratio of areas under the curves calculated for the replicated experiments that induced the most

and the least = 2.8). Moreover, conductivity showed considerable heteroscedasticity, *i.e.* there

was a positive correlation between the mean area under the curves and their variance with

Spearman's coefficient = 0.91; p < 0.05 (S1 Fig). Four examples of the relationship between

variability and mean conductivity are shown in Fig 2. This large variability raised the question of

how the HR induction could be quantified to compare the strains, as addressed in the next section.

Fig 1. Variability in the increase of conductivity measured at 24 hours post-inoculation (hpi) over

the replicated experiments performed for each *AvrB*-expressing strain, at 18°C (A), 24°C (B) and

28°C (C). Bars represent the mean increase in conductivity for each of the independent biological

replicates (3 ≤ n ≤ 27). Values are sorted from left to right in a descending order.

Fig 2. Electrolyte leakage curves obtained post-infiltration for strain M6 *AvrB* at 18°C (A), USA007

***AvrB* at 18°C (B), LAB0041 *AvrB* at 24°C (C) and CC1498 *AvrB* at 28°C (D).** Each curve represents

the mean increase in conductivity measured over time for one independent biological replicate

performed with three technical replicates each. Error bars represent standard error.

Intensity of HR induction varies among strains.

To compare HR induction intensity among strains, we accounted for the variability described

above by fitting Linear Mixed effects Models to our data. This statistical approach allowed us to

classify strains as inducers or non-inducers at each of the temperatures that we tested. This

revealed that almost all the *P. syringae* strains induced HR except for J35 *AvrB* at all the

temperatures, MAFF302273 *AvrB* at 24°C and 28°C, B728a *AvrB* at 24°C, and 1448A and CRA-FRU

8.43 at 28°C (Table 1). The inducer strains led to conductivity levels that were significantly greater

than those measured in the mock treatment (10mM $MgCl_2$ or *hrp*-inducing medium HIM) (p <

0.05) according to Linear Mixed-effects Models. More specifically, our results demonstrated that

all the inducer strains varied in the intensity of the HR induction they caused. The model estimated

the mean difference in points of conductivity measured between the tested modality and the

mock treatment. The values presented in Table 1 ("Estimate") are proxies of the intensity of HR

induction for each strain. DC3000 *AvrB* and M6 *AvrB* displayed the highest values ("Estimate" > 65

regardless of the temperature tested), whereas some others such as the three quasi-clonal strains

CC0073 *AvrB*, CC0094 *AvrB* and CC1498 *AvrB* averaged less than 20. These differences are also

reflected in the mean areas under conductivity progress curves (AUCPC) at each temperature as

shown in Fig 3. This proxy of the HR induction intensity was correlated neither with the *AvrB*

expression levels (S2 Fig) nor with the bacterial growth capacity at a given temperature (S3 Fig).

Table 1. Statistics associated with linear mixed-effects models fitted to ion leakage experiment

results.

PHYLOGROUP	STRAIN	INDUCER AT A GIVEN TEMPERATURE						TEMPERATURE	
		18°C		24°C		28°C			
		Estimate	*p-value*	*Estimate*	*p-value*	*Estimate*	*p-value*	*p-value*	*Estimate*
1a	DC3000	69.97	**2.00E-16**	73.12	**1.25E-15**	65.27	**8.46E-15**	8.02E-01	-0.19
	M6	95.14	**2.00E-16**	89.06	**5.35E-16**	73.05	**1.59E-14**	1.95E-01	-2.07
	T1	28.98	**2.00E-16**	29.07	**2.00E-16**	13.45	**2.81E-08**	**1.42E-02**	**-1.26**
	LAB0041	31.36	**4.24E-11**	29.56	**2.00E-16**	16.83	**2.83E-09**	5.17E-02	-1.22
1b	J35	3.35	5.43E-02	0.18	9.19E-01	-1.27	4.46E-01	NA	
	CRA-FRU 8.43	14.09	**4.69E-11**	4.97	**4.33E-04**	0.97	4.45E-01	**3.80E-03**	**-1.28**
	USA007	52.24	**2.00E-16**	46.69	**8.58E-16**	34.15	**2.26E-12**	**3.35E-02**	**-1.70**
2d	CC0073	15.22	**5.97E-07**	15.28	**7.62E-07**	19.77	**3.42E-09**	3.55E-01	0.63
	CC0094	6.86	**6.44E-08**	9.72	**2.61E-06**	14.49	**8.91E-08**	**9.76E-04**	**0.97**
	CC1498	7.89	**4.96E-06**	12.69	**2.24E-05**	19.96	**1.52E-08**	**2.58E-02**	**1.36**
	B728a	3.12	**2.08E-02**	2.22	1.61E-01	4.75	**8.07E-03**	**4.27E-02**	**0.37**
	MAFF302273	3.17	**1.73E-02**	2.88	9.68E-02	1.78	3.77E-01	NA	
3a	1448A	5.14	**1.07E-02**	6.73	**9.73E-03**	2.46	2.41E-01	6.41E-01	0.53

The effect of strains was calculated for each mutant bacterium at each temperature by comparing

with the results obtained for the mock treatment (10mM $MgCl_2$). Significant values of the estimate

parameter ($p < 0.05$, in bold) are associated with HR inducing conditions. The temperature effect

was evaluated for each *AvrB*-expressing strain, inducing HR at least at two temperatures (non-

inducer strains are highlighted in grey), comparing the electrolyte leakage curves obtained at

these different temperatures. Strains insensitive to the temperature ($p > 0.05$) are highlighted in

green, strains leading to minor conductivity levels as the temperature increases (negative

estimate, $p < 0.05$) are highlighted in blue, and strains displaying a warm-dependency (positive

estimate, $p < 0.05$) are highlighted in red. The number of replicated experiments for each modality

(strain x temperature condition) varied from 3 up to 27.

Fig 3. Mean area under conductivity progress curve (AUCPC) calculations for all the replicated

experiments ($3 \leq n \leq 27$) of each *AvrB*-expressing strain at 18°C (A), 24°C (B) and 28°C (C). Error

bars represent standard error.

Intensity of HR induction is temperature-sensitive in half of the inducer

strains.

With Linear Mixed-effects Models, we categorized the HR inducer strains according to the

temperature-sensitivity of HR induction. Six out of eleven strains were insensitive to a change in

temperature (DC3000 *AvrB*, M6 *AvrB*, LAB0041 *AvrB*, CC0073 *AvrB*, 1448A *AvrB*) ($p > 0.05$) (Table

1). The other HR inducer strains were considered temperature-sensitive ($p < 0.05$) and could be

categorized into two groups according to whether the "Estimate" values were negative or positive.

In one group, strains showed higher conductivity with lower temperatures (T1 *AvrB*, CRA-FRU 8.43

AvrB and USA007 *AvrB*). In the case of CRA-FRU 8.43 *AvrB* strain, the decrease in conductivity

observed with increasing temperature led the strain to be categorized as a non-inducer at 28°C

(Table 1). In the other group, strains showed higher conductivity with higher temperatures

(CC0094 *AvrB*, CC1498 *AvrB* and B728a *AvrB*). Temperature-sensitivity of the *AvrB*-expressing

strains was further confirmed by the analysis of HrpZ secretion *in vitro* as an assessment of T3SS

functionality. Interestingly, the strain MAFF302273 *AvrB* displayed a functional T3SS, while it

constantly did not lead to conductivity increase once infiltrated into leaf disks, suggesting T3SS

inactivation in *A. thaliana* (S4 Fig).

Temperature responsiveness patterns vary among strains but remain

consistent within the same phylogroup.

A given phylogroup could contain strains that induced HR in temperature-sensitive and

temperature-insensitive ways (Table 1). However, temperature-sensitive strains within a same

phylogroup showed the same trend relative to increasing temperature. Indeed, the group of

strains whose T3SS was more efficient at the cooler temperatures (*i.e.*, low-temperature-

dependent; T1 *AvrB*, CRA-FRU 8.43 *AvrB* and USA007 *AvrB*) and the group whose T3SS was more

efficient at warmer temperatures (*i.e.*, warm-temperature-dependent; CC0094 *AvrB*, CC1498 *AvrB*

and B728a *AvrB*) belong to the phylogroups 1 and 2d, respectively (Table 1). Additionally, HrpZ

secretion by MAFF302273 *AvrB* was more important as the temperature rose, revealing a pattern

of warm-temperature dependency for T3SS efficiency, similar to the other thermosensitive strains

belonging to the same phylogroup 2d (S4 Fig). Nevertheless, dramatic differences among strains

even very closely related phylogenetically were still observed, as illustrated by the three quasi-

clonal *AvrB*-expressing strains from phylogroup 2d, CC0073, CC0094 and CC1498 (Table 1) (Fig 4).

CC0073 *AvrB* was temperature-independent in its ability to induce HR while CC0094 *AvrB* and

CC1498 *AvrB* both showed clearly greater capacity at warmer temperatures. Moreover, in terms

of intensity, CC1498 *AvrB* led to higher values of conductivity measured at 24 hpi than CC0073

AvrB and CC0094 *AvrB*. Finally, in terms of kinetics, CC0073 *AvrB* was responsible for a faster HR

induction at the beginning of the interaction – no matter the temperature concerned – that

stalled, whereas CC0094 *AvrB* and CC1498 *AvrB* were much slower in inducing HR (Fig 4).

Fig 4. Electrolyte leakage curves over time obtained post-infiltration for CC0073 *AvrB* (A),

CC0094 *AvrB* (B) and CC1498 *AvrB* (C). Infiltrated leaf disks were incubated at 18°C (blue), 24°C

(pink) and 28°C (red). Conductivity following infiltration of the mock treatment (10mM $MgCl_2$) and

incubation at 24°C is represented by the black dotted line. Data represent one single

representative biological replicate with all strains and temperature conditions evaluated

simultaneously with three technical replicates each. Error bars represent standard error.

Strains from the *P. syringae* complex display diverse speeds of HR

induction.

To account for dynamic aspects in the behaviors of *AvrB*-expressing *P*. *syringae* strains, the ratio

of conductivity values measured at 6 hpi and 24 hpi was calculated for each replicated experiment

for all modalities that were found to be HR-inducing. This parameter was used as a proxy for HR

induction velocity. An analysis of variance of these ratios revealed significant differences among

AvrB-expressing strains (ANOVA1, p < 0.05). DC3000 *AvrB* and M6 *AvrB* proved to be the fastest

HR inducers with more than 70% of their total induction already observable at 6 hpi (Fig 5). Yet

again, the diversity of behaviors within phylogroups was revealed, even among the three *AvrB*-

expressing quasi-clonal strains (Fig 5, post-hoc Dunn's test showed that CC0073 *AvrB* and CC0094

AvrB and CC1498 *AvrB* groups differed significantly at p < 0.05). Furthermore, 6 hpi/24 hpi-ratios

were compared across the different temperatures overall revealing that higher temperatures

correlated with faster electrolyte leakage, consistent with the results observed for bacterial *in-*

vitro growth (S5 Fig).

Fig 5. **Estimation of velocity of HR induction.** The proxy for velocity was calculated as the ratio of

conductivity increase measured at 6 hpi and 24 hpi for each *AvrB*-expressing strain inducing HR at

least at one temperature condition. Strains are sorted by phylogroup from left to right, PG1a

(purple), PG1b (green), PG2d (yellow), PG3a (red). Boxes were calculated from 5 up to 73

replicated experiments per strain (single dots). Boxes associated with different letters represent

significantly different means (p < 0.05, according to a Kruskal-Wallis's test followed by post-hoc

Dunn's test).

Whole genome sequences suggest a few candidate genes for

temperature-sensitivity.

As we wanted to decipher the origin of sensitivity to the temperature for HR induction, we applied

a comparative genomic approach focusing on the three quasi-clonal strains CC0073, CC0094 and

CC1498. These strains displayed at least 99.84% homology (ANI calculation). Among the whole set

of strains, 286 putative genes were specific to the three CC-strains, and 76 were common only to

the two warm temperature-dependent ones (CC0094 and CC1498) (Fig 6). By comparing the

putative genes present in CC0094 and CC1498 but missing in CC0073 (regardless of their

presence/absence in the other strains) the list was extended to 173 genes, with 94% of the

proteins encoded by these genes unknown (neither the name nor the function) and only 10

putative genes with a known or putative function (S1 Table).

Fig 6. Comparative genomics performed for the 13 strains characterized in ion leakage

experiments. Vertical bars correspond to the number of genes shared by groups of strains or

exclusively present in single strains (group/strain-specific genes). Horizontal color bars represent

the type of behavior of each strain in response to temperature, as identified in Table 1. Common

putative genes specific to the 3 quasi-clonal strains CC0073, CC0094 and CC1498 are in yellow,

common putative genes specific to CC0094 and CC1498 are in red.

Discussion

New perspectives on virulence thermoregulation processes.

The diverse behaviors of T3SS efficiency observed in relation to incubation temperature indicate

that temperature plays a crucial role in regulating bacterial virulence. This observation is another

piece of evidence out of the many reported over the past 30 years regarding the role of

temperature in virulence regulation in plant-associated bacteria (reviewed in references [37,38]).

Our work has presented new examples of relatively warm-temperature dependency for virulence

efficiency, consistent with previous findings about gene expression beyond and within the *P.*

syringae species complex [45,46,54,65–67]. Altogether, these pieces of evidence do not align with

the long-standing notion according to which "almost all virulence genes of plant pathogenic

bacteria [...] exhibit increased transcription at temperatures well-below the respective growth

optima" [38]. Such misconception has led some authors to conclude that "the increase of bacterial

growth in plants at elevated temperature cannot be explained by variations in intrinsic bacterial

growth rate or secretion of effector proteins at different temperatures. A slow HR is also unlikely

to be caused by a slow accumulation of bacteria or effector proteins leading to a slower detection

by plants" [68]. Such statements are strongly contradicted by our results, and no longer included

in the conclusions of other important and more recent articles that present an implicitly updated

perspective with which our work aligns [37,54,60].

Effector repertoire role in the lack of HR induction.

Although *AvrB* expression was assessed for all the strains, two of them (J35 *AvrB* and MAFF302273

AvrB) were categorized as non-inducers for at least two of the tested temperatures. While HrpZ

secretion data *in vitro* confirmed that J35 *AvrB*'s T3SS was unactive, the contrary was

demonstrated for MAFF302273 *AvrB*. Consequently, the lack of HR induction could result from the

T3SS inactivation in *A. thaliana*, potentially because of a quick PTI onset [27]. On the other hand,

it could also be a matter of speed of ETI onset, whose rapidity prevents cell death occurrence [69].

Otherwise, strain MAFF302273 *AvrB* may contain in its repertoire some suppressive effector(s)

blocking the ability of AvrB to trigger the HR [70], or some effector(s) targeting a molecular actor

involved in both PTI and ETI, or simply blocking the PTI – required in the first place for ETI

activation. Consistently we checked for effector repertoire composition for strain MAFF302273

(PsyTEC [71]). To the best of our knowledge, no evidence of direct or indirect antagonism between

AvrB and other effectors naturally present was found. However, it is not possible to exclude the

occurrence of 'meta-effector' interactions [20,72]. Overall, the in-depth-study of the effector

repertoires of all the strains used in our work could help to explain the differences observed in

terms of HR intensity, with one or more effectors that may inhibit HR but also enhance it

conversely – as observed for strain CC1498 *AvrB* versus strain CC0094 *AvrB*, for example.

Temperature and plant immunity.

Using HR induction as a proxy for T3SS efficiency, we implicitly accounted for plant defense

mechanisms that play a role in the response [69]. In *A. thaliana*, the role of temperature in plant

immunity is more complex than what reported by previous studies. In 2009, Wang and colleagues

concluded that a mild increase in temperature impaired both PTI and ETI, suggesting a general

deficiency in the plant defense system at elevated temperatures (28-30°C) [68]. Subsequently, a

straightforward dichotomy was established: less efficient ETI (effective from 10 to 23°C, with a

peak at 18°C) versus more efficient PTI (effective from 23 to 32°C, with a peak at 28°C) as the

temperature increases [56]. This model was later qualified by the demonstration that, since ETI

relies on distinct processes (*i.e.*, virulence suppression and programmed cell death/HR), only HR

was inhibited by elevated temperatures as the virulence suppression was maintained [73].

However, despite their demonstration that ion leakage from *A. thaliana* leaf disks infiltrated with

the avirulent DC3000 strain is completely inhibited at 28°C-30°C, our results for this *AvrB*-strain

and CC0094 *AvrB* and CC1498 *AvrB* contradict this finding.

Temperature variations may affect the kinetics of effector translocation and/or binding to R

proteins and hence impact the speed and efficiency of signaling cascades involved in the defense

response. Comparison of the ratio of conductivity values measured at 6 hpi and 24 hpi among

temperatures for a given HR-inducer strain revealed that the elevation of the temperature led to

an acceleration of electrolyte leakage as temperature increased. As this observation was

maintained across the strains that induced HR at different temperatures, it is tempting to associate

this general effect of accelerated response with the effect of the higher temperature on the plant,

and not on the T3SS activation or efficiency. Nevertheless, across all strains, we demonstrated that

some were faster than others to induce HR, also indicating some pathosystem-specific behaviors.

Temperature and effector secretion properties.

Interestingly, a few examples of temperature-dependency for T3SE secretion have been shown to

be 'protein/effector specific' – mentioning some specific secretion properties of AvrB [53].

Likewise, the two avirulent DC3000 strains with which HR suppression at elevated temperatures

was demonstrated carried either the HopZ1a or AvrRpt2 effector [73]. HopZ1a, AvrRpt2 and AvrB

are recognized by the NLRs ZAR1, RPS2 and RPM1, respectively, which are all members of the

subgroup of CC-NLRs. The unclear variability of temperature-sensitivity across different NLR

subgroups (CC-NLRs versus TIR-NLRs) [73,74] suggests that sensitivity may not be conserved in all

NLRs or CC-NLRs, given their distinct structural and functional properties. In this light, it is

reasonable to attribute the differences observed between our work and the one of Menna and

colleagues [73] to the different 'ligand-receptor' couples used. Echoing the differences in

'protein/effector specific' secretion efficiency, there are likely to be differences in temperature-

sensitivity for effector recognition by R proteins. Overall, we assume that the impact of

temperature on T3SS-secretion/ETI/HR-triggering efficiencies can be highly pathosystem-specific.

Illustrating this high specificity, DC3000 *AvrB* and M6 *AvrB* displayed a unique behavior among all

the tested strains. Being such strong and fast HR inducers probably reflects the fact that they are

highly aggressive on *A. thaliana* Col-0 (in their WT version), so the exacerbated response may also

be a sign of the global host compatibility between these two strains and the model plant [75,76].

365   Thermosensing origin and distribution among strains.

We attempted to identify the mechanism(s) that may be responsible for the thermosensitivity at

the origin of the different behaviors we observed. Two-component systems such as thermosensor

CorRSP responsible for the low-temperature-dependency of coronatin production in *P. syringae*

pv. *glycinea* could be involved in the regulation of T3SS-related gene expression in response to

temperature [61]. Alternatively, if evidence hints that gene expression is not directly modified by

temperature [55], thermoregulation might act later in the transcription/translation chain, during

T3SS assembly, or on translocation efficiency. In addition, the study of potential differences in

membrane fluidity between strains is promising. More specifically, membrane fluidity is

influenced by functional lipid microdomains ('FMM' or lipid rafts) that are enriched in proteins

involved in signal transduction and protein secretion and have the capacity to organize in space

and time components of transport machineries [77]. Likely, T3SS proteins co-localize on lipid rafts

to allow their proper assembly, and temperature would impair or favor the process, by changing

membrane fluidity.

We observed that the behavior of high temperature-preference/dependency is exclusive to strains

from phylogroup 2d, whereas the opposite case is specific to strains belonging to phylogroups 1a

and 1b. This suggests that temperature responsiveness patterns might vary among strains but

remain consistent within the same phylogroup. While the temperature sensitivity feature is widely

distributed among the strains, the preference/dependency on low or high temperatures in case

of thermosensitivity is conserved within a same phylogroup. This implies that the genes or targets

responsible for both processes may be distinct, making their identification more challenging

through comparative genomics. Beyond the identification of the mechanisms involved in

thermosensitivity, preference/dependency on low or high temperatures could be linked to the

evolutionary history of pathogens, which may have been shaped by temperature-related selection

pressures. Investigating this question would help shed light on the origin of the phenomenon.

Virulence thermoregulation in relation to disease development.

Since the same thermoregulation components may control various virulence factors, one

intriguing question is whether the effect of temperature on virulence factors is consistent (i) for a

given strain, and (ii) with temperature conditions in which disease occurrence is generally

reported. Crossing observations between temperature optimum for virulence and

epidemiological data regarding our selected strain revealed a high level of complexity within the

P. syringae species complex and does not allow us to draw any clearer conclusions. Concerning *P.*

syringae pv. *phaseolicola*, the low-temperature-dependency was established for phaseolotoxin

production and T6SS regulation, in line with the development of halo blight of bean favored at

temperatures below 25°C [51,78]. However, T3SS-related gene expression inhibition has been

reported at 18°C [50], and our results indicate that the strain 1448A is thermo-insensitive in its

ability to induce HR. Regarding *P. syringae* pv. *actinidiae*, our results confirmed a low-temperature

dependency for T3SS efficiency [55], which is entirely in line with the environmental conditions –

in late winter or early spring – under which *P. syringae* pv. *actinidiae* is reported to cause disease

or damage in the kiwifruit orchards [79,80]. On the other hand, concerning the strains CC0073

and CC0094 which were isolated from cantaloupe during blight epidemics in southern France in

the 90s, the authors have reported that the climatic conditions that favored disease outbreak and

development were heavy rainfall associated with a mean minimum temperature below 12-13°C

(under which the physiology of cantaloupe comes to a standstill) in 95% of cases, whereas our

results demonstrated a major T3SS efficiency at higher temperatures [81–83]. Finally, strain

DC3000 displays a consistent behavior, exhibiting both unaltered T3SS efficiency and coronatine

production in response to temperature, at least within the 18°C to 28°C range [44]. Its consistency

in response to various environmental parameters associated with high reproducibility of data

related to it have certainly contributed to making it the model strain it has become. However, this

strain was generated in the laboratory as a rifampicin-resistant derivative of the strain DC52,

which originated from Guernsey in 1960 [21,84]. Therefore, no conclusive connection could be

established between its ecology and its virulence behavior in response to temperature, as this

model strain has evolved for many years in laboratories, far from a field context.

Another important consideration is that our results highlighted that growth optima – at least *in*

vitro – are not correlated with T3SS efficiency. Therefore, providing a definitive answer to the

question of whether the temperature conditions that favor virulence (specifically T3SS) are

identical to those reported as favorable for disease development is challenging due to both

supporting and opposite examples: several traits characterizing phytopathogenic bacteria are

thermo-sensitive, and each pathogen is characterized by growth and virulence mechanisms that

are activated at precise temperatures, generating a complex scenario in which temperature can

alter the outcome of the interaction between plant and bacterium [60]. Consequently, we state

that efforts to simplify or generalize the behavior of different strains would not necessarily help

our understanding.

Interstrain variability.

One important conclusion of our work is the significant variability in behavior among strains, even

when closely related at the genomic level (as illustrated with the 3 'CC' strain-quasi-clones). Over

the years of research on these issues, examples of such inherent variability in terms of virulence

have been largely overlooked, with only a few cases reported in literature, both beyond and within

the *P. syringae* species complex [44,85–87]. This might be attributed to the commendable aim of

simplifying our understanding of the biological models and plant-pathogen interactions under

study. However, we emphasize the urgent need to study the diversity among strains, since

evidence of their inherent variability has been documented. Although investigating model strains

has facilitated the understanding and characterization of intricate molecular mechanisms in plant-

pathogen interactions, it is now imperative to consider the full spectrum of *P. syringae* species

complex diversity [3,88], both within and outside of the agronomic context.

Material & Methods

Bacterial strains and culture conditions

Thirteen *Pseudomonas syringae* strains were selected for this study (Table 2). These strains

represented four phylogroups/clades (1a, 1b, 2d, and 3a). The geographical origin and isolation

source of the strains varied, encompassing annual crops (*e.g.* M6, T1) [89,90], woody plants (*e.g.*

J35, CRA-FRU 8.43, MAFF302273) [91–93], and environmental reservoirs like rivers or snow (*e.g.*

LAB0041, USA007) [3,94]. Genomic relatedness within a phylogroup was also considered, aiming

for a balance of very divergent and closely related strains, such as the three 'CC' strains (CC0073,

CC0094, and CC1498) that are quasi-clones [95–97]. Model strains DC3000, B728a and 1448A

were also included in the set as there is a plethora of information about their biology [84,98,99].

Table 2. Set of selected *P. syringae* strains.

PHYLOGROUP	STRAIN	FULL NAME	ALIAS IN COLLECTIONS	ISOLATION SOURCE	COUNTRY	YEAR	REFERENCE
1a	DC3000	*P. syringae* pv. *tomato*	CFBP 7438, ICMP 18429	*Solanum lycopersicum*	United-Kingdom	1960	[84]
	M6	*P. syringae* pv. *maculicola*	CFBP 4314, NCPPB 1766, LMG 5560	*Brassica oleracea* var. Botrytis	United-Kingdom	1965	[90]
	T1	*P. syringae* pv. *tomato*	-	*Solanum lycopersicum*	Canada	1986	[89]
	LAB0041	*P. syringae*	-	Epilithic biofilm	France	2009	[3]
1b	J35	*P. syringae* pv. *actinidiae*	CFBP 4909, ICMP 9617, NCPPB 3739	*Actinidia deliciosa* cv. Hayward	Japan	1984	[93]
	CRA-FRU 8.43	*P. syringae* pv. *actinidiae*	-	*Actinidia chinensis* cv. Hort16	Italy	2008	[91]
	USA007	*P. syringae*	CFBP 8574	Creek water	United States of America	2007	[94]
2d	CC0073	*P. syringae* pv. *aptata*	CFBP 5430	*Cucumis melo*	France	1997	[95]
	CC0094	*P. syringae* pv. *aptata*	CFBP 8529	*Cucumis melo*	France	1997	[95]
	CC1498	*P. syringae*	-	Snowfall	France	2006	[96]
	B728a	*P. syringae* pv. *syringae*	CFBP 8502, ICMP 18427, NCPPB 4487, LMG 26717	*Phaseolus vulgaris*	United States of America	1987	[99]
	MAFF302273	*P. syringae* pv. *aceris*	CFBP 2339, ICMP 2802, NCPPB 958, LMG 2106	*Acer* sp.	United States of America	1939	[92]
3a	1448A	*P. syringae* pv. *phaseolicola*	CFBP 7087, NCPPB 4478	*Phaseolus vulgaris*	Ethiopia	1985	[98]

456   Wild-type (WT) *P. syringae* strains were grown at 28°C on King's B agar (KB) [100] – supplemented

457   with rifampicin at 50µg/mL for DC3000 and B728a strains. Kanamycin at 50µg/mL was added for

458   the culture of *AvrB* mutants. Single colonies of each strain were picked and grown overnight in

459   liquid KB medium – supplemented with rifampicin at 50μg/mL for DC3000 and B728a strains and

460   with kanamycin at 50μg/mL for *AvrB* mutants, with agitation at 190rpm at 28°C.

AvrB-transformation

Four mL of fresh WT bacterial suspension were centrifuged (5min at 4500g at 4°C) and pellets

were washed three times in 1mL and then resuspended in 30-50μL of sterile 10% glycerol solution

kept on ice. Once the pure plasmid carrying *AvrB* gene (pVB01 Km^R [62]) was added to the

suspensions, they underwent then an electroporation process. Once recovered, the suspensions

were streaked on petri dishes containing KB agar – supplemented with kanamycin at 50μg/mL for

all strains plus rifampicin at 50μg/mL for DC3000 and B728a strains. Plates were incubated for 48

up to 72 hours a 28°C. Mutant colonies were picked and *AvrB* presence was checked by PCR using

Promega GoTaq™ DNA polymerase (Fisher Scientific, USA) and specific primers (FOR:

ATGGGCTGCGTCTCGTCAAAAAGCA, REV: TTAAAAGCAATCAGAATCTAGCAAG, annealing

temperature = 65°C) followed by electrophoresis of the amplicons on a 1-1.5% (w/v) agarose gel.

AvrB expression control

Relative expression levels of *AvrB* among the mutant strains were evaluated by real-time PCR. Four

mL of fresh suspension of *AvrB*-transformed *P. syringae* strains grown overnight in selective

medium were centrifuged (5min at 4500g at room temperature) and total RNA was extracted from

bacterial pellets using TRIzol substitute (EuroGold TriFast™ Kit, EuroClone, Italy) according to

manufacturer instructions. Extraction yield and quality of the transcripts were checked by

spectrophotometric reading (NanoDrop^{TM}, Thermo Fisher Scientific, USA). Samples with

absorbance ratios A_{260nm}/A_{230nm} or A_{260nm}/A_{280nm} outside the range of [2 ± 0.5] were cleaned

following a lithium chloride RNA precipitation treatment (solution made of 7.5M LiCl, 50mM EDTA,

pH 8.0). 1-2µg RNA per sample underwent DNase treatment (TURBO^{TM} DNase, Life Technologies,

USA) then the single-stranded complementary DNA synthesis was performed from 0.5-1µg of

template RNA using SuperScript^{TM} III Reverse Transcriptase enzyme (Invitrogen, USA) according to

manufacturer instructions. Samples get 10-fold diluted and procedure was performed using Luna®

Universal qPCR Master Mix (New England Biolabs, USA) and *rpoD* housekeeping gene as reference,

and amplification was run on a QuantStudio^{TM} 3 real-time PCR system (Applied Biosystems, USA)

(primers *AvrB* FOR: ATACTCTGCATTCGCCTCCG, REV: TCAACGACTCTGGCAAGGAC; primers *rpoD*

FOR: AACTTGCGTCTGGTGATCTC, REV: ATCAAGCCGATGTTGCCTTC). *AvrB* expression was

determined using the $2^{-\Delta\Delta Ct}$ method and normalized against *rpoD*, and fold-change for each strain

was established relative to the strain DC3000 *AvrB.*

Ion-leakage experiments

Hypersensitive response (HR) induction in *Arabidopsis thaliana* Col-0 was assessed by quantifying

programmed cell death through ion-leakage measurement, as previously published [101,102]

with slight modifications. *A. thaliana* Col-0 plants were cultivated in a mixture of peat soil and

perlite (4/1, v/v) in a growth chamber set at 24°C/21.5°C thermoperiod and 8 h light–16 h dark

photoperiod with 70% of relative humidity for 6 to 8 weeks. Three healthy plants with six 5-mm

leaf disks detached from a same plant with a cork borer were used for each experimental

condition. To recognize the plant they were detached from, leaf disks were marked with a colored

felt-tip and were then transferred into a 50mL tube containing 10mM MgCl$_2$ solution while

bacterial suspension preparation. Ten mL of fresh suspension of *AvrB*-expressing *P. syringae*

strains grown overnight in selective medium were centrifuged (5min at 4500g at room

temperature) and pellets were washed three times in 10mL of sterile 10mM MgCl$_2$ solution.

Optical density at 600nm was measured by spectrophotometer (Evolution™, ThermoFisher

Scientific, USA) and bacterial cells were finally resuspended in sterile 10mM MgCl$_2$ solution to

reach 10^8 CFU/mL (OD$_{600nm}$ = 0.1; final volume = 30mL). Leaf disks were infiltrated with the

bacterial suspensions using a 50mL syringe or a vacuum chamber. Sterile 10mM MgCl$_2$ solution or

hrp-inducing medium (HIM; using 27mM glycerol as carbon source; pH 5.5) [103] were used as

negative controls. Infiltrated leaf disks were rinsed twice for 15m in 40mL of milliQ water under

gentle agitation (90 rpm) at room temperature to eliminate suspension/mock solution trace and

ions leaked due to mechanical damage. Infiltrated leaf disks were then distributed into 12-well

plate containing 2mL of milliQ water per well in order to have: (i) six disks per well, (ii) two disks

from each of the three different plants per well, (iii) three identical wells (one plate column) per

each experimental modality. Initial conductivity values (μS.cm^{-1}) in each well was measured prior

plates incubation (120 rpm; constant light of 80 μmol.m^{-2}.s^{-1}) at different temperatures (18°C, 24°C

and 28°C) using a compact conductometer (LAQUAtwin EC-11, HORIBA Scientific, Japan).

Conductivity increase was measured every 1.5-2 hours during the first 8 hours of the experiment,

and then twice between 18- and 24-hours post-inoculation (hpi), obtaining from 6 to 9

conductivity measures on average per well. Each experimental modality (*i.e.* one specific *AvrB*-

expressing strain infiltrated and then incubated at a certain temperature) was reproduced at least

three times (at least 3 `biological replicates' or `replicated experiments').

Statistical analysis

For each technical replicate, the conductivity value from the first measure was subtracted to

values measured at each successive time points for normalization [55]. For each strain-

temperature modality within each replicated experiment, conductivity was calculated as the

average of the values of the three technical replicates for each time point, and ion-leakage curves

over time were established accordingly. Area under each single conductivity progress curve was

calculated (AUCPC), as well as variance and mean AUCPC per modality. Moreover, conductivity

values corresponding to precisely 6 hpi and 24 hpi were calculated via linear interpolation, and

the ratio 6 hpi/24 hpi was used as a dynamic parameter, indicating how fast a given strain induced

HR at a given temperature. Comparison of strains with each other based on the 6 hpi/24 hpi

conductivity ratio was assessed by analysis of variance (Kruskal-Wallis's test) followed by a post-

hoc Dunn's test. The same procedure was conducted to compare the three different temperatures

regardless of the strain. Data were represented graphically by boxplots.

Each single conductivity progress curve constituted from 6 to 9 time points and was considered

as one statistical individual (*i.e.*, one iteration) to fit Linear Mixed-effects Models (LMM). This

allowed us to account for the important variability among the diverse experimental replicates of

a same modality and, most importantly, to consider the entire time course of electrolyte leakage,

rather than focusing solely on isolated time points. The mathematical representation of LMM is:

$Y = X\beta + Z\upsilon + \varepsilon$; where Y represents the dependent variable (the measured feature, *i.e.* conductivity

increase), $X\beta$ stands for the fixed effects (*i.e.* the potential explanatory variables we want to assess

the effect on Y), $Z\upsilon$ stands for the random effects (all the unknown parameters that may be

responsible for the huge variability among the different replicated experiments and which we are

not directly interested to), and ε represents the residual. Using LMM in RStudio, we evaluated (i)

the strain effect (*i.e.* if the ion-leakage following *AvrB*-expressing strain infiltration was significantly

different from the ion-leakage due to mock treatment infiltration) to categorize each strain as 'HR-

inducer' or 'non-inducer' at each of the tested temperatures; and (ii) the temperature effect (*i.e.*

if the ion-leakage following *AvrB*-expressing strain infiltration was significantly different as the

incubation temperature changed) on the strains considered as HR-inducers at least at two of the

tested temperatures. In this way we estimated the values of model parameters and the associated

p-values.

In-vitro HrpZ secretion

Secretion of HrpZ *in vitro* was assessed for each strain used in ion leakage experiments. Twenty

mL of fresh suspension of *AvrB*-expressing *P. syringae* strains grown overnight in selective medium

were centrifuged (5min at 4500g at room temperature) and pellets were washed three times in

10mL of sterile minimal medium (HIM). Optical density at 600nm was measured with a

spectrophotometer (Evolution™, ThermoFisher Scientific, USA) and bacterial cells were

resuspended in sterile HIM to reach 7 x 10^8 CFU/mL (OD_{600nm} = 0.7; final volume = 150mL).

Suspensions were divided into three portions of 50mL, each was separately incubated at 18°C,

24°C and 28°C with agitation at 190 rpm. After 5 and 24 hours of incubation, 25mL of each

bacterial suspension was sampled and absorbance at 600nm was measured again (Evolution™,

ThermoFisher Scientific, USA). Cell-free culture supernatant was obtained by centrifugation and

filtration, and proteins of supernatant were then precipitated with trichloroacetic acid (TCA) as

previously described [104,105] with TCA final concentration at 5%.

Samples were mixed with loading buffer (50mM DTT) and run (10 min at 100V; 80 min at 130V)

through two 1.5mm 12% sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE),

either loading 25µL of each sample, or adjusting volume to load the supernatant extracted from

ca. 10^7 cells (OD$_{600nm}$ = 0.6, final volume = 17.5µL). 180kDa-PageRuler^{TM} (ThermoFisher Scientific,

USA) was used as protein ladder. One gel was used to assess total soluble protein content in each

sample and thus confirm the correct loading ratio: after rinsing with milli-Q water, the gel was

stained for 20 minutes in Coomassie R-250 staining solution (25% isoprophyl alcohol, 10% acetid

acid, 0.05% Coomassie R-250) and destained with 10% acetic acid until complete removal of the

background. The second gel was transferred onto nitrocellulose blotting membrane (GE

HealthCare, USA) for Western Blot analysis. The membrane was immersed for one hour at 37°C in

a blocking solution made of phosphate-buffered saline solution (PBS) (137mM NaCl, 2.7mM KCl,

10mM Na$_2$HPO$_4$, 1.8mM KH$_2$PO$_4$), supplemented with 0.05% Tween® 20 and 5% non-fat dry milk.

Then, the membrane was probed with the specific primary anti-HrpZ antibody [106], diluted 1:500

in the same blocking solution, and incubated at room temperature for 1h and a half. Following

primary antibody incubation, membrane was washed 3 times for 15min in PBS supplemented with

0.05% Tween® 20 under gentle agitation in the dark. Then, the membrane was probed for 1h at

room temperature in the dark using the conjugated secondary antibody to horseradish peroxidise

(Amersham Biosciences, USA), diluted 1:10000 in the same blocking solution. Finally, after three

additional washings in PBS (supplemented with 0.05% Tween® 20 for the first two), proteins were

visualised (Chemidoc^{TM}, ThermoFisher Scientific, USA) by enhanced chemiluminescent (ECL)

detection (GE HealthCare, USA). Blot pictures were analyzed using ImageLab software (BioRad,

USA). The intensity of each band was normalized to the intensity of the 70kDa band from the

ladder. When necessary, relative quantity was then normalized to the optical density measured

by the end of the incubation period to adjust the results to the number of cells initially present in

each sample.

In-vitro growth measurement

In-vitro bacterial growth under different temperatures was assessed for each strain. Four mL of

fresh suspension of *AvrB*-expressing *P. syringae* strains grown overnight in selective medium were

centrifuged (5min at 4500g at room temperature) and pellets were washed three times in 1mL of

fresh liquid KB. Optical density at 600nm was measured by spectrophotometer (Evolution™,

ThermoFisher Scientific, USA) and bacterial cells were then resuspended to reach 2 x 10^7 CFU/mL

(OD_{600nm} = 0.02; final volume = 2mL). Suspensions were aliquoted in a microtiter plate (200μL per

well), with 6 technical replicates per strain, and plates were incubated at 18°C, 24°C and 28°C.

Liquid KB (supplemented or not with the antibiotics) was used as negative control. Absorbance at

600nm was measured every 1.5-2 hours during the first 8 hours of the experiment, and then two

or three times between 18 and 24 hpi, obtaining from 6 to 9 absorbance measures per well. To

eliminate unspecific absorbance signals due to the color of KB medium and the antibiotics, the

value from the first measure was subtracted from values measured at each successive time points.

Absorbance was then calculated as the average of the values of the six technical replicates for

each time point, and bacterial growth curves over time were established accordingly. Each strain-

temperature modality was reproduced at least twice, except for T1.

Genome comparison

Genomes of the 13 *AvrB*-expressing strains used in ion-leakage experiments were retrieved mostly from public sequence databases (Pseudomonas Genome Database and NCBI); draft genomes of strains LAB0041, CC0073 and CC1498 were provided by Dr. Cindy Morris. To analyze the genomes, a custom bioinformatic pipeline was created, with the help of the Nextflow framework [107]. The pipeline was composed of several steps: i) the FastANI tool was used for rapid many-to-many ANI calculation [108]; ii) BUSCO was used to assess both the genome quality and protein annotations quality [109]; iii) the Prokka pipeline was used for the protein annotation [110]; iv) the EggNog-mapper tool was used for the functional annotation [111]; and v) the Panaroo pipeline was implemented for the creation of a pangenome of the 13 strains [112]. Then, the Panaroo output table of gene presence/absence was exploited via an R script to produce interactive tables to determine the number of total genes and specific genes per strain. These tables allowed for the selection of genes present in a given strain, and queries could be performed to obtain information for genes shared by strains.

Acknowledgments

The authors thank Pr. Boris A. Vinatzer, Dr. Perrine Portier, Charlotte Chandeysson and Dr. Marco Scortichini for kindly providing the *P. syringae* strains used in this study. We also thank Pr. Sheng-Yang He for providing the antibody anti-HrpZ. Special thanks to Pr. Roberto Chignola and Dr. Davide Martinetti for their help in analyzing ion-leakage data through statistical approaches. Finally, we

632   also thank the members of the laboratory of plant pathology of the University of Verona for their

633   technical support in conductivity measurements.

References

1.   Lamichhane JR, Messéan A, Morris CE. Insights into epidemiology and control of diseases

of annual plants caused by the *Pseudomonas syringae* species complex. J Gen Plant Pathol.

2015;81: 331–350. doi:10.1007/s10327-015-0605-z

2.   Lamichhane JR, Varvaro L, Parisi L, Audergon J-M, Morris CE. Disease and frost damage of

woody plants caused by *Pseudomonas syringae*. Advances in Agronomy. 2014. pp. 235–

295. doi:10.1016/B978-0-12-800132-5.00004-3

3.   Berge O, Monteil CL, Bartoli C, Chandeysson C, Guilbaud C, Sands DC, et al. A user's guide

to a data base of the diversity of *Pseudomonas syringae* and its application to classifying

strains   in   this   phylogenetic   complex.   PLoS   One.   2014;9:   e105547.

doi:10.1371/journal.pone.0105547

4.   Sharma A, Gupta AK, Devi B. Current trends in management of bacterial pathogens

infecting plants. Antonie Van Leeuwenhoek. 2023;116: 303–326. doi:10.1007/s10482-023-

01809-0

5.   Melotto M, Kunkel BN. Virulence strategies of plant pathogenic bacteria. In: Rosenberg E,

DeLong EF, Lory S, Stackebrandt E, Thompson F, editors. The Prokaryotes. Berlin,

Heidelberg: Springer Berlin Heidelberg; 2013. pp. 61–82. doi:10.1007/978-3-642-30141-

4_62

6.   Roine E, Wei W, Yuan J, Nurmiaho-Lassila EL, Kalkkinen N, Romantschuk M, et al. Hrp pilus:

an *hrp*-dependent bacterial surface appendage produced by *Pseudomonas syringae* pv.

tomato DC3000. Proc Natl Acad Sci U S A. 1997;94: 3459–3464.

doi:10.1073/pnas.94.7.3459

7.   Büttner D, He SY. Type III protein secretion in plant pathogenic bacteria. Plant Physiol.

2009;150: 1656–1664. doi:10.1104/PP.109.139089

8.   Jin Q, He S-Y. Role of the Hrp pilus in type III protein secretion in *Pseudomonas syringae*.

Science. 2001;294: 2556–2558. doi:10.1126/science.1066397

9.   Li CM. The Hrp pilus of *Pseudomonas syringae* elongates from its tip and acts as a conduit

for translocation of the effector protein HrpZ. EMBO J. 2002;21: 1909–1915.

doi:10.1093/emboj/21.8.1909

10.   Cunnac S, Lindeberg M, Collmer A. *Pseudomonas syringae* type III secretion system

effectors: repertoires in search of functions. Curr Opin Microbiol. 2009;12: 53–60.

doi:10.1016/j.mib.2008.12.003

11.   Galán JE, Collmer A. Type III secretion machines: bacterial devices for protein delivery into

host cells. Science. 1999;284: 1322–1328. doi:10.1126/science.284.5418.1322

12.   Guo M, Tian F, Wamboldt Y, Alfano JR. The majority of the type III effector inventory of

Pseudomonas syringae pv. *tomato* DC3000 can suppress plant immunity. Mol Plant-

Microbe Interact. 2009;22: 1069–1080. doi:10.1094/MPMI-22-9-1069

13.   Mudgett MB, Staskawicz BJ. Protein signaling via type III secretion pathways in

phytopathogenic bacteria. Curr Opin Microbiol. 1998;1: 109–115. doi:10.1016/S1369-

5274(98)80150-1

14.   Hueck CJ. Type III protein secretion systems in bacterial pathogens of animals and plants.

Microbiol Mol Biol Rev. 1998;62: 379–433. doi:10.1128/mmbr.62.2.379-433.1998

15.   Lee CA. Type III secretion systems: machines to deliver bacterial proteins into eukaryotic

cells? Trends Microbiol. 1997;5: 148–156. doi:10.1016/S0966-842X(97)01029-9

16.   Guttman DS, Vinatzer BA, Sarkar SF, Ranall M V., Kettler G, Greenberg JT. A functional

screen for the type III (Hrp) secretome of the plant pathogen *Pseudomonas syringae*.

Science. 2002;295: 1722–1726. doi:10.1126/science.295.5560.1722

17.   Chang JH, Urbach JM, Law TF, Arnold LW, Hu A, Gombar S, et al. A high-throughput, near-

saturating screen for type III effector genes from *Pseudomonas syringae*. Proc Natl Acad

Sci U S A. 2005;102: 2549–2554. doi:10.1073/pnas.0409660102

18.   Xin X-F, Nomura K, Aung K, Velásquez AC, Yao J, Boutrot F, et al. Bacteria establish an

aqueous living space in plants crucial for virulence. Nature. 2016;539: 524–529.

doi:10.1038/nature20166

19.   Lindeberg M, Cunnac S, Collmer A. *Pseudomonas syringae* type III effector repertoires: last

words in endless arguments. Trends Microbiol. 2012;20: 199–208.

doi:10.1016/j.tim.2012.01.003

20.   Martel A, Laflamme B, Breit-McNally C, Wang P, Lonjon F, Desveaux D, et al. Metaeffector

interactions modulate the type III effector-triggered immunity load of *Pseudomonas*

syringae. PLoS Pathog. 2022;18: e1010541. doi:10.1371/JOURNAL.PPAT.1010541

21.   Xin X-F, Kvitko B, He SY. *Pseudomonas syringae*: what it takes to be a pathogen. Nat Rev

Microbiol. 2018;16: 316–328. doi:10.1038/nrmicro.2018.17

22.   Morris CE, Lamichhane JR, Nikolić I, Stanković S, Moury B. The overlapping continuum of

host range among strains in the *Pseudomonas syringae* complex. Phytopathol Res. 2019;1:

4. doi:10.1186/s42483-018-0010-6

23.   Baltrus DA, McCann HC, Guttman DS. Evolution, genomics and epidemiology of

Pseudomonas syringae. Mol Plant Pathol. 2017;18: 152–168. doi:10.1111/MPP.12506

24.   Ruiz-Bedoya T, Wang PW, Desveaux D, Guttman DS. Cooperative virulence via the collective

action of secreted pathogen effectors. Nat Microbiol. 2023;8: 640–650.

doi:10.1038/s41564-023-01328-8

25.   Cui H, Xiang T, Zhou JM. Plant immunity: a lesson from pathogenic bacterial effector

proteins. Cell Microbiol. 2009;11: 1453–1461. doi:10.1111/J.1462-5822.2009.01359.X

26.   Boller T, Felix G. A renaissance of elicitors: perception of Microbe-Associated Molecular

Patterns and danger signals by Pattern-Recognition Receptors. Annu Rev Plant Biol.

2009;60: 379–406. doi:10.1146/annurev.arplant.57.032905.105346

27.   Crabill E, Joe A, Block A, van Rooyen JM, Alfano JR. Plant immunity directly or indirectly

restricts the injection of type III effectors by the *Pseudomonas syringae* type III secretion

system. Plant Physiol. 2010;154: 233–244. doi:10.1104/pp.110.159723

28.   Anderson JC, Wan Y, Kim Y-M, Pasa-Tolic L, Metz TO, Peck SC. Decreased abundance of

type III secretion system-inducing signals in *Arabidopsis mkp1* enhances resistance against

Pseudomonas syringae. Proc Natl Acad Sci. 2014;111: 6846–6851.

doi:10.1073/pnas.1403248111

29.   Thomma BPHJ, Nürnberger T, Joosten MHAJ. Of PAMPs and effectors: the blurred PTI-ETI

dichotomy. Plant Cell. 2011;23: 4–15. doi:10.1105/tpc.110.082602

30.   Monteiro F, Nishimura MT. Structural, functional, and genomic diversity of plant NLR

proteins: an evolved resource for rational engineering of plant immunity. Annu Rev

Phytopathol. 2018;56: 243–267. doi:10.1146/annurev-phyto-080417-045817

31.   Jones JDG, Dangl JL. The plant immune system. Nature. 2006;444: 323–329.

doi:10.1038/nature05286

32.   Ngou BPM, Ahn H-K, Ding P, Jones JDG. Mutual potentiation of plant immunity by cell-

surface and intracellular receptors. Nature. 2021;592: 110–115. doi:10.1038/s41586-021-

03315-7

33.   Yuan M, Jiang Z, Bi G, Nomura K, Liu M, Wang Y, et al. Pattern-recognition receptors are

required for NLR-mediated plant immunity. Nature. 2021;592: 105–109.

doi:10.1038/s41586-021-03316-6

34.   Balint-Kurti P. The plant hypersensitive response: concepts, control and consequences. Mol

Plant Pathol. 2019;20: 1163–1178. doi:10.1111/mpp.12821

35.   Coll NS, Epple P, Dangl JL. Programmed cell death in the plant immune system. Cell Death

Differ. 2011;18: 1247–1256. doi:10.1038/cdd.2011.37

36.   Stevens RB. JG Horsfall and AE Dimond, eds. Plant Pathology, an Advanced Treatise, Vol 3.

Academic Press, NY; 1960. pp. 357–429.

37.   Velásquez AC, Castroverde CDM, He SY. Plant–pathogen warfare under changing climate

conditions. Current Biology. Cell Press; 2018. pp. R619–R634.

doi:10.1016/j.cub.2018.03.054

38.   Smirnova A, Li H, Weingart H, Aufhammer S, Burse A, Finis K, et al. Thermoregulated

expression of virulence factors in plant-associated bacteria. Arch Microbiol. 2001;176:

393–399. doi:10.1007/s002030100344

39.   Hockett KL, Burch AY, Lindow SE. Thermo-regulation of genes mediating motility and plant

interactions in *Pseudomonas syringae*. PLoS One. 2013;8: e59850. doi:10.1371/journal.pone.0059850

40. Aguilera S, Alvarez-Morales A, Murillo J, Hernández-Flores JL, Bravo J, De la Torre-Zavala S. Temperature-mediated biosynthesis of the phytotoxin phaseolotoxin by *Pseudomonas syringae* pv. *phaseolicola* depends on the autoregulated expression of the *phtABC* genes. PLoS One. 2017;12: e0178441. doi:10.1371/journal.pone.0178441

41. Palmer DA, Bender CL. Effects of environmental and nutritional factors on production of the polyketide phytotoxin coronatine by *Pseudomonas syringae* pv. *glycinea*. Appl Environ Microbiol. 1993;59: 1619–1626. doi:10.1128/aem.59.5.1619-1626.1993

42. Ullrich MS, Schergaut M, Boch J, Ullrich B. Temperature-responsive genetic loci in the plant pathogen *Pseudomonas syringae* pv. *glycinea*. Microbiology. 2000;146: 2457–2468. doi:10.1099/00221287-146-10-2457

43. Braun Y, Smirnova A V., Weingart H, Schenk A, Ullrich MS. Coronatine gene expression *in vitro* and *in planta*, and protein accumulation during temperature downshift in *Pseudomonas syringae*. Sensors. 2009;9: 4272–4285. doi:10.3390/s90604272

44. Weingart H, Stubner S, Schenk A, Ullrich MS. Impact of temperature on *in planta* expression of genes involved in synthesis of the *Pseudomonas syringae* phytotoxin coronatine. Mol Plant-Microbe Interact. 2004;17: 1095–1102. doi:10.1094/MPMI.2004.17.10.1095

45. Krishna PS, Woodcock SD, Pfeilmeier S, Bornemann S, Zipfel C, Malone JG. *Pseudomonas syringae* addresses distinct environmental challenges during plant infection through the coordinated deployment of polysaccharides. J Exp Bot. 2022;73: 2206–2221. doi:10.1093/jxb/erab550

764   46.   Peñaloza-Vázquez A, Kidambi SP, Chakrabarty AM, Bender CL. Characterization of the

765   alginate biosynthetic gene cluster in *Pseudomonas syringae* pv. *syringae*. J Bacteriol.

766   1997;179: 4464–4472. doi:10.1128/jb.179.14.4464-4472.1997

767   47.   Li H, Schenk A, Srivastava A, Zhurina D, Ullrich MS. Thermo-responsive expression and

768   differential secretion of the extracellular enzyme levansucrase in the plant pathogenic

769   bacterium *Pseudomonas syringae* pv. *glycinea*. FEMS Microbiol Lett. 2006;265: 178–185.

770   doi:10.1111/J.1574-6968.2006.00486.X

771   48.   Scalschi L, Fernández-Crespo E, Pitarch-Marin M, Llorens E, González-Hernández AI,

772   Camañes G, et al. Response of tomato-*Pseudomonas* pathosystem to mild heat stress.

773   Horticulturae. 2022;8: 174. doi:10.3390/horticulturae8020174

774   49.   Bender CL, Alarcón-Chaidez F, Gross DC. *Pseudomonas syringae* phytotoxins: mode of

775   action, regulation, and biosynthesis by peptide and polyketide synthetases. Microbiol Mol

776   Biol Rev. 1999;63: 266–292. doi:10.1128/MMBR.63.2.266-292.1999

777   50.   Arvizu-Gómez JL, Hernández-Morales A, Aguilar JRP, Álvarez-Morales A. Transcriptional

778   profile of *P. syringae* pv. *phaseolicola* NPS3121 at low temperature: physiology of

779   phytopathogenic bacteria. BMC Microbiol. 2013;13: 81. doi:10.1186/1471-2180-13-81

780   51.   Nüske J, Fritsche W. Phaseolotoxin production by *Pseudomonas syringae* pv. *phaseolicola*:

781   the   influence   of   temperature.   J   Basic   Microbiol.   1989;29:   441–447.

782   doi:10.1002/JOBM.3620290713

783   52.   Aguilera S, López-López K, Nieto Y, Garcidueñas-Piña R, Hernández-Guzmán G, Hernández-

784   Flores JL, et al. Functional characterization of the gene cluster from *Pseudomonas syringae*

785   pv. *phaseolicola* NPS3121 involved in synthesis of phaseolotoxin. J Bacteriol. 2007;189:

2834. doi:10.1128/JB.01845-06

53.   Van Dijk K, Fouts DE, Rehm AH, Hill AR, Collmer A, Alfano JR. The Avr (effector) proteins

HrmA (HopPsyA) and AvrPto are secreted in culture from *Pseudomonas syringae* pathovars

via the Hrp (type III) protein secretion system in a temperature- and pH-sensitive manner.

J Bacteriol. 1999;181: 4790–4797. doi:10.1128/jb.181.16.4790-4797.1999

54.   Huot B, Castroverde CDM, Velásquez AC, Hubbard E, Pulman JA, Yao J, et al. Dual impact

of elevated temperature on plant defence and bacterial virulence in *Arabidopsis*. Nat

Commun. 2017;8: 1808. doi:10.1038/s41467-017-01674-2

55.   Puttilli MR, Danzi D, Correia C, Brandi J, Cecconi D, Manfredi M, et al. Plant signals

anticipate the induction of the type III secretion system in *Pseudomonas syringae* pv.

actinidiae, facilitating efficient temperature-dependent effector translocation. Microbiol

Spectr. 2022;10. doi:10.1128/spectrum.02073-22

56.   Cheng C, Gao X, Feng B, Sheen J, Shan L, He P. Plant immune response to pathogens differs

with changing temperatures. Nat Commun. 2013;4: 2530. doi:10.1038/ncomms3530

57.   Klinkert B, Narberhaus F. Microbial thermosensors. Cellular and Molecular Life Sciences.

Springer; 2009. pp. 2661–2676. doi:10.1007/s00018-009-0041-3

58.   Jin S, Song YN, Deng WY, Gordon MP, Nester EW. The regulatory VirA protein of

Agrobacterium tumefaciens does not function at elevated temperatures. J Bacteriol.

1993;175: 6830–6835. doi:10.1128/JB.175.21.6830-6835.1993

59.   Arvizu-Gómez JL, Hernández-Morales A, Jofre-Garfias AE, Álvarez-Morales A. Temperature,

the GacS/GacA system, and host metabolites regulate the type VI secretion system of

Pseudomonas syringae pv. *phaseolicola*. J Gen Plant Pathol. 2013;79: 194–197.

doi:10.1007/s10327-013-0443-9

60.   Tribelli PM, López NI. Insights into the temperature responses of *Pseudomonas* species in

beneficial and pathogenic host interactions. Applied Microbiology and Biotechnology.

2022. pp. 7699–7709. doi:10.1007/s00253-022-12243-z

61.   Ullrich M, Penaloza-Vazquez A, Bailey AM, Bender CL. A modified two-component

regulatory system is involved in temperature- dependent biosynthesis of the *Pseudomonas*

syringae phytotoxin coronatine. J Bacteriol. 1995;177: 6160–6169.

doi:10.1128/jb.177.21.6160-6169.1995

62.   Innes RW, Bisgrove SR, Smith NM, Bent AF, Staskawicz BJ, Liu Y. Identification of a disease

resistance locus in *Arabidopsis* that is functionally homologous to the *RPG1* locus of

soybean. Plant J. 1993;4: 813–820. doi:10.1046/j.1365-313X.1993.04050813.x

63.   Mudgett MB. New insights to the function of phytopathogenic bacterial type III effectors

in plants. Annual Review of Plant Biology. Annual Reviews; 2005. pp. 509–531.

doi:10.1146/annurev.arplant.56.032604.144218

64.   Mackey D, Holt BF, Wiig A, Dangl JL. RIN4 interacts with *Pseudomonas syringae* type III

effector molecules and is required for RPM1-mediated resistance in *Arabidopsis*. Cell.

2002;108: 743–754. doi:10.1016/S0092-8674(02)00661-X

65.   Wei M, Yokoyama T, Minamisawa K, Mitsui H, Itakura M, Kaneko T, et al. Soybean seed

extracts preferentially express genomic loci of *Bradyrhizobium japonicum* in the initial

interaction with soybean, *Glycine max* (L.) Merr. DNA Res. 2008;15: 201–214.

doi:10.1093/dnares/dsn012

66.   Wei M, KeisukeTakeshima, Yokoyama T, Minamisawa K, Mitsui H, Itakura M, et al.

Temperature-dependent expression of type III secretion system genes and its regulation in

Bradyrhizobium japonicum. Mol Plant-Microbe Interact. 2010;23: 628–637.

doi:10.1094/MPMI-23-5-0628

67. Czajkowski R, Kaczyńska N, Jafra S, Narajczyk M, Lojkowska E. Temperature-responsive

genetic loci in pectinolytic plant pathogenic *Dickeya solani*. Plant Pathol. 2017;66: 584–594.

doi:10.1111/PPA.12618

68. Wang Y, Bao Z, Zhu Y, Hua J. Analysis of temperature modulation of plant defense against

biotrophic microbes. Mol Plant-Microbe Interact. 2009;22: 498–506. doi:10.1094/MPMI-

22-5-0498

69. Künstler A, Bacsó R, Gullner G, Hafez YM, Király L. Staying alive – is cell death dispensable

for plant disease resistance during the hypersensitive response? Physiol Mol Plant Pathol.

2016;93: 75–84. doi:10.1016/j.pmpp.2016.01.003

70. Jamir Y, Guo M, Oh H, Petnicki-Ocwieja T, Chen S, Tang X, et al. Identification of

Pseudomonas syringae type III effectors that can suppress programmed cell death in plants

and yeast. Plant J. 2004;37: 554–565. doi:10.1046/j.1365-313X.2003.01982.x

71. Laflamme B, Dillon MM, Martel A, Almeida RND, Desveaux D, Guttman DS. The pan-

genome effector-triggered immunity landscape of a host-pathogen interaction. Science.

2020;367: 763–768. doi:10.1126/science.aax4079

72. Kubori T, Shinzawa N, Kanuka H, Nagai H. *Legionella* metaeffector exploits host proteasome

to temporally regulate cognate effector. PLoS Pathog. 2010;6: e1001216.

doi:10.1371/journal.ppat.1001216

73. Menna A, Nguyen D, Guttman DS, Desveaux D. Elevated temperature differentially

852   influences effector-triggered immunity outputs in *Arabidopsis*. Front Plant Sci. 2015;6: 1–

853   7. doi:10.3389/fpls.2015.00995

854 74.   Hua J. Temperature and plant immunity. Temperature and Plant Development. Wiley;

855   2014. pp. 163–180. doi:10.1002/9781118308240.ch7

856 75.   Preston GM. *Pseudomonas syringae* pv. *tomato* : the right pathogen, of the right plant, at

857   the right time. Mol Plant Pathol. 2000;1: 263–275. doi:10.1046/j.1364-3703.2000.00036.x

858 76.   Xin XF, He SY. *Pseudomonas syringae* pv. *tomato* DC3000: a model pathogen for probing

859   disease susceptibility and hormone signaling in plants. Annu Rev Phytopathol. 2013;51:

860   473–498. doi:10.1146/annurev-phyto-082712-102321

861 77.   Bramkamp M, Lopez D. Exploring the existence of lipid rafts in bacteria. Microbiol Mol Biol

862   Rev. 2015;79: 81–100. doi:10.1128/mmbr.00036-14

863 78.   Goss RW. The relation of temperature to common and halo blight of beans.

864   Phytopathology. 1940;30: 258–264.

865 79.   Kim GH, Jung JS, Koh YJ. Occurrence and epidemics of bacterial canker of kiwifruit in Korea.

866   Plant Pathol J. 2017;33: 351–361. doi:10.5423/PPJ.RW.01.2017.0021

867 80.   Donati I, Cellini A, Sangiorgio D, Vanneste JL, Scortichini M, Balestra GM, et al.

868   *Pseudomonas syringae* pv. *actinidiae*: ecology, infection dynamics and disease

869   epidemiology. Microb Ecol. 2020;80: 81–102. doi:10.1007/s00248-019-01459-8

870 81.   Morris CE, Pitrat M. La bactériose du melon : connaissances acquises et travaux en cours.

871   PHM Rev Hortic. 1998; 44–47.

872 82.   Riffaud C. La bactériose du melon : écologie et stratégies de lutte contre *Pseudomonas*

873   *syringae* pv.*aptata*. Université Claude Bernard Lyon I. 2002.

83.   Mention P, Lavigne D, Leix-Henry F, Bouchu S, Morris CE, Prior P, et al. Bactériose du melon:

acquis et perspectives. PHM Rev Hortic. 2004; 22–26.

84.   Cuppels DA. Generation and characterization of Tn5 insertion mutations in *Pseudomonas*

syringae pv. *tomato*. Appl Environ Microbiol. 1986;51: 323–327.

doi:10.1128/aem.51.2.323-327.1986

85.   Baron C, Domke N, Beinhofer M, Hapfelmeier S. Elevated temperature differentially affects

virulence, VirB protein accumulation, and T-pilus formation in different *Agrobacterium*

tumefaciens and *Agrobacterium vitis* strains. J Bacteriol. 2001;183: 6852–6861.

doi:10.1128/JB.183.23.6852-6861.2001

86.   Hasegawa H, Chatterjee A, Cui Y, Chatterjee AK. Elevated temperature enhances virulence

of *Erwinia carotovora* subsp. *carotovora* strain EC153 to plants and stimulates production

of the quorum sensing signal, N -acyl homoserine lactone, and extracellular proteins. Appl

Environ Microbiol. 2005;71: 4655–4663. doi:10.1128/AEM.71.8.4655-4663.2005

87.   Rohde BH, Pohlack B, Ullrich MS. Occurrence of thermoregulation of genes involved in

coronatine biosynthesis among various *Pseudomonas syringae* strains. J Basic Microbiol.

1998;38: 41–50. doi:10.1002/(SICI)1521-4028(199803)38:1<41::AID-JOBM41>3.0.CO;2-6

88.   Gomila M, Busquets A, Mulet M, García-Valdés E, Lalucat J. Clarification of taxonomic

status within the *Pseudomonas syringae* species group based on a phylogenomic analysis.

Front Microbiol. 2017;8: 320397. doi:10.3389/fmicb.2017.02422

89.   Whalen MC, Innes RW, Bent AF, Staskawicz BJ. Identification of *Pseudomonas syringae*

pathogens of *Arabidopsis* and a bacterial locus determining avirulence on both *Arabidopsis*

and soybean. Plant Cell. 1991;3: 49. doi:10.1105/TPC.3.1.49

90.   Debener T, Lehnackers H, Arnold M, Dangl JL. Identification and molecular mapping of a

single *Arabidopsis thaliana* locus determining resistance to a phytopathogenic

Pseudomonas syringae isolate. Plant J. 1991;1: 289–302. doi:10.1046/J.1365-

313X.1991.T01-7-00999.X

91.   Ferrante P, Scortichini M. Molecular and phenotypic features of *Pseudomonas syringae* pv.

actinidiae isolated during recent epidemics of bacterial canker on yellow kiwifruit (*Actinidia*

chinensis) in central Italy. Plant Pathol. 2010;59: 954–962. doi:10.1111/j.1365-

3059.2010.02304.x

92.   Sawada H, Suzuki F, Matsuda I, Saitou N. Phylogenetic analysis of *Pseudomonas syringae*

pathovars suggests the horizontal gene transfer of *argK* and the evolutionary stability of

hrp gene cluster. J Mol Evol. 1999;49: 627–644. doi:10.1007/PL00006584

93.   Takikawa Y, Serizawa S, Ichikawa T, Tsuyumu S, Goto M. *Pseudomonas syringae* pv.

actinidiae pv. nov.: the causal bacterium of canker of kiwifruit in Japan. Japanese J

Phytopathol. 1989;55: 437–444. doi:10.3186/jjphytopath.55.437

94.   Morris CE, Sands DC, Vanneste JL, Montarry J, Oakley B, Guilbaud C, et al. Inferring the

evolutionary history of the plant pathogen *Pseudomonas syringae* from its biogeography

in headwaters of rivers in North America, Europe, and New Zealand. MBio. 2010;1.

doi:10.1128/mBio.00107-10

95.   Morris CE, Glaux C, Latour X, Gardan L, Samson R, Pitrat M. The relationship of host range,

physiology, and genotype to virulence on cantaloupe in *Pseudomonas syringae* from

cantaloupe blight epidemics in France. Phytopathology. 2000;90: 636–646.

doi:10.1094/PHYTO.2000.90.6.636

96.   Morris CE, Sands DC, Vinatzer BA, Glaux C, Guilbaud C, Buffière A, et al. The life history of

the plant pathogen *Pseudomonas syringae* is linked to the water cycle. ISME J. 2008;2: 321–

334. doi:10.1038/ismej.2007.113

97.   Monteil CL, Yahara K, Studholme DJ, Mageiros L, Méric G, Swingle B, et al. Population-

genomic insights into emergence, crop adaptation and dissemination of *Pseudomonas*

syringae pathogens. Microb Genomics. 2016;2: e000089. doi:10.1099/mgen.0.000089

98.   Taylor JD, Teverson DM, Allen DJ, Pastor-Corrales MA. Identification and origin of races of

Pseudomonas syringae pv. *phaseolicola* from Africa and other bean growing areas. Plant

Pathol. 1996;45: 469–478. doi:10.1046/J.1365-3059.1996.D01-147.X

99.   Loper JE, Lindow SE. Lack of evidence for *in situ* fluorescent pigment production by

Pseudomonas syringae pv. *syringae* on bean leaf surfaces. Phytopathology. 1987;77: 1449.

doi:10.1094/phyto-77-1449

100. King EO, Ward MK, Raney DE. Two simple media for the demonstration of pyocyanin and

fluorescin. J Lab Clin Med. 1954;44: 301–307.

101. Imanifard Z, Vandelle E, Bellin D. Measurement of hypersensitive cell death triggered by

avirulent bacterial pathogens in *Arabidopsis*. Methods Mol Biol. 2018;1743: 39–50.

doi:10.1007/978-1-4939-7668-3_4

102. Johansson ON, Nilsson AK, Gustavsson MB, Backhaus T, Andersson MX, Ellerström M. A

quick and robust method for quantification of the hypersensitive response in plants. PeerJ.

2015;3: e1469. doi:10.7717/peerj.1469

103. Huynh T V., Dahlbeck D, Staskawicz BJ. Bacterial blight of soybean: regulation of a pathogen

gene determining host cultivar specificity. Science. 1989;245: 1374–1377.

doi:10.1126/science.2781284

104.   Flaugnatti N, Journet L. Identification of effectors: precipitation of supernatant material.

Bacterial Protein Secretion Systems. 2017. pp. 459–464. doi:10.1007/978-1-4939-7033-

9_31

105.   Kobayashi T, Kishigami S, Sone M, Inokuchi H, Mogi T, Ito K. Respiratory chain is required

to maintain oxidized states of the DsbA-DsbB disulfide bond formation system in

aerobically growing *Escherichia coli* cells. Proc Natl Acad Sci U S A. 1997;94: 11857–11862.

doi:10.1073/pnas.94.22.11857

106.   Alfano JR, Bauer DW, Milos TM, Collmer A. Analysis of the role of the *Pseudomonas*

syringae pv. *syringae* HrpZ harpin in elicitation of the hypersensitive response in tobacco

using functionally non-polar *hrpZ* deletion mutations, truncated HrpZ fragments, and *hrmA*

mutatio. Mol Microbiol. 1996;19: 715–728. doi:10.1046/j.1365-2958.1996.415946.x

107.   Di Tommaso P, Chatzou M, Floden EW, Barja PP, Palumbo E, Notredame C. Nextflow

enables reproducible computational workflows. Nature Biotechnology. Nature Publishing

Group; 2017. pp. 316–319. doi:10.1038/nbt.3820

108.   Jain C, Rodriguez-R LM, Phillippy AM, Konstantinidis KT, Aluru S. High throughput ANI

analysis of 90K prokaryotic genomes reveals clear species boundaries. Nat Commun.

2018;9: 5114. doi:10.1038/s41467-018-07641-9

109.   Manni M, Berkeley MR, Seppey M, Simão FA, Zdobnov EM. BUSCO update: novel and

streamlined workflows along with broader and deeper phylogenetic coverage for scoring

of eukaryotic, prokaryotic, and viral genomes. Mol Biol Evol. 2021;38: 4647–4654.

doi:10.1093/molbev/msab199

110. Seemann T. Prokka: rapid prokaryotic genome annotation. Bioinformatics. 2014;30: 2068–

2069. doi:10.1093/bioinformatics/btu153

111. Cantalapiedra CP, Herṇandez-Plaza A, Letunic I, Bork P, Huerta-Cepas J. eggNOG-mapper

v2: functional annotation, orthology assignments, and domain prediction at the

metagenomic scale. Mol Biol Evol. 2021;38: 5825–5829. doi:10.1093/molbev/msab293

112. Tonkin-Hill G, MacAlasdair N, Ruis C, Weimann A, Horesh G, Lees JA, et al. Producing

polished prokaryotic pangenomes with the Panaroo pipeline. Genome Biol. 2020;21: 1–21.

doi:10.1186/s13059-020-02090-4

Supporting information captions

S1 Fig. Variance of the mean area under conductivity progress curve (AUCPC) values obtained

for each 39 modalities *AvrB*-expressing strain x temperature. Number of replicated experiments

for each strain varied from 3 up to 27. Linear regression trend curve was fitted to the data, with

y=323.89x − 150932 and R^2=0.8588, showing a positive correlation between mean AUCPC and

their variance (Spearman's coefficient = 0.91; $p < 0.05$).

S2 Fig. *AvrB* expression among the mutant strains. RNA was extracted from bacterial suspensions

grown overnight at 28°C in liquid KB medium supplemented with kanamycin (50µg/mL) and

rifampicin (50µg/mL, for DC3000 *AvrB* and B728a *AvrB* only). Real-time PCR was performed using

rpoD as the housekeeping gene. Expression levels were established using the $2^{-\Delta\Delta Ct}$ method using

DC3000 *AvrB* as a reference (fold change = 1). Error bars represent standard error. Data represent

the mean of three independent biological replicates, each repeated at least twice.

S3 Fig. *In-vitro* growth curves for DC3000 *AvrB* (A), USA007 *AvrB* (B) and CC0094 *AvrB* (C).

Bacterial growth was measured for 24 hours at 18°C (blue), 24°C (pink) and 28°C (red). Overnight

grown bacteria were resuspended in liquid KB medium supplemented with kanamycin (50μg/mL)

and rifampicin (50μg/mL, for DC3000 *AvrB* only) at an initial load of 10^7 CFU/mL (optical density

measured at 600nm = 0.01). Data represent one single representative biological replicate with all

strains and temperature conditions evaluated simultaneously with six technical replicates each.

Error bars represent standard error.

S4 Fig. HrpZ secretion profiles across a range of temperature. *AvrB*-expressing strains were

incubated in *hrp*-inducing medium (HIM) at 18°C, 24°C or 28°C for 5 hours. Cell-free culture

supernatant was obtained by centrifugation and filtration. Proteins of bacterial culture

supernatant were then precipitated with trichloroacetic acid (TCA) and analyzed by Western blot

using an anti-HrpZ antibody. The expected molecular weight is *ca*. 40kDa. A mix of samples proven

to contain HrpZ in previous replicates was used as the positive control (C+) at two different

concentrations (1X and 2X). Samples were run in different gels that were transferred onto

membranes whose pictures were cropped and merged into a single figure.

1002

1003   **S5 Fig. Ratios of conductivity increase measured at 6 hpi and 24 hpi for all *AvrB*-expressing**

1004   **strains inducing HR at least at one temperature condition, at each of the temperatures.** Boxes

1005   were calculated from 70 to 97 replicated experiments per temperature (single dots). Boxes

1006   associated with different letters represent significantly different means ($p < 0.05$, according to a

1007   Kruskal-Wallis's test followed by post-hoc Dunn's test).

1008

1009   **S1 Table. List of the 10 genes with available annotation present in strains CC0094 and CC1498**

1010   **and missing in strain CC0073.** Four out of these ten genes are also present in other strains.