Multi-omic analyses unveil contrasting composition and spatial distribution of specialized metabolites in seeds of *Camelina sativa* and other Brassicaceae

Léa Barreda[1,±], Céline Brosse[1,±], Stéphanie Boutet[1], Nicolas Klewko[1], Delphine De Vos[1], Tracy Francois[1], Boris Collet[1], Damaris Grain[1], Céline Boulard[1], Jean Chrisologue Totozafy[1], Benoît Bernay[2], François Perreau[1], Loïc Lepiniec[1], Loïc Rajjou[1], Massimiliano Corso[1*]

[1] Université Paris-Saclay, INRAE, AgroParisTech, Institut Jean-Pierre Bourgin for Plant Sciences (IJPB), 78000, Versailles, France

[2] Plateforme Proteogen, SFR ICORE 4206, Université de Caen Basse-Normandie, Esplanade de la paix, 14032 Caen cedex, France

[±] Equal contribution

[*] Correspondence to: Massimiliano Corso massimiliano.corso@inrae.fr

E-mails:

LB lea.barreda@inrae.fr

CB celine.brosse@inrae.fr

SB stephanie.boutet@inrae.fr

NK nicolas.klewko57@gmail.com

DDV delphine.de-vos@inrae.fr

TF tracy.francois@inrae.fr

BC boris.collet@inrae.fr

DG damaris.grain@inrae.fr

CB celine.boulard@inrae.fr

JCT jean-chrisologue.totozafy@inrae.fr

BB benoit.bernay@unicaen.fr

FP francois.perreau@inrae.fr

LL loic.lepiniec@inrae.fr

LR loic.rajjou@inrae.fr

MC massimiliano.corso@inrae.fr

Running head: Specialized metabolites in camelina seed tissues

Keywords: *Camelina sativa*, glucosinolates, seed tissues, Brassica, Arabidopsis, seed development, seed germination, untargeted metabolomics, transcriptomics, proteomics

Summary

Seeds of Brassicaceae produce a large diversity of beneficial and antinutritional specialized metabolites (SMs) that influence their quality and provide resistance to stresses. While the distribution of these compounds has been described in leaves and roots tissues, limited information is available about their spatio-temporal accumulation in seeds.

Camelina sativa (camelina) is an oilseed Brassicaceae cultivated for human and animal nutrition, and for industrial uses. While we previously explored SM diversity and plasticity, no information is available about SM distribution and expression of related proteins and genes in camelina seeds.

In this study, we used a multi-omic approach, integrating untargeted metabolomics, data-independent acquisition proteomics, and transcriptomics to investigate the synthesis, modifications and degradations of SMs accumulated in the different seed tissues (i.e. seed coat, endosperm, and embryo) at 6 developmental and 2 germination stages. Our results showed distinct patterns of SMs and their related pathways, highlighting significant contrasts in seed composition and spatial distribution for the defence-related and antinutritional glucosinolate (GSL) compounds among camelina, *Arabidopsis thaliana,* and *Brassica napus,* three closely-related Brassicaceae species. Notably, the variation in GSL spatial distributions was primarily driven by differences in their structure and transport mechanisms. Long chain C8-C11 methylsulfinylalkyl GSLs were predominantly accumulated in the seed coat and endosperm, while mid- and short-chain C3-C7 methylsulfinylalkyl GSLs were accumulated in the embryo.

Characterizing the spatial dynamics of seed SMs provides valuable insights that can guide the development of crops with optimized distribution of beneficial and toxic metabolites, improving seed nutritional profiles for feed and food.

Introduction

Plants produce a myriad of specialized metabolites (SMs) with tremendous interest for agriculture, nutrition and health (Arimura and Maffei, 2016). The large diversity of chemical structures and, hence, of functions observed among specialized metabolic classes arises from different functional groups (decorations or modifications) commonly added to SM core structures (Barreda et al., 2024). SM structures can indeed be modified by the addition or removal of functional groups, hydroxylation, methylation, glycosylation and acylation being the most frequent modification reactions.

Various SMs are widely accumulated in seeds of Brassicaceae model and crop species, which thus constitute an interesting model to study SMs diversity and functions (Corso et al., 2020; Corso et al., 2021; Baud et al., 2023). Major metabolic classes that are accumulated in seeds of Brassicaceae species, including the model species *Arabidopsis thaliana* and the crops *Camelina sativa* and *Brassica napus* (rapeseed), are the nitrogen-containing compounds glucosinolates, and many phenylpropanoids, including flavonols, proanthocyanidins (PAs, or condensed tannins), cinnamic acids and monolignols (Halkier and Gershenzon, 2006; Quéro et al., 2016; Barreda et al., 2024).

While some SM classes can be directly synthetized in seeds, others are synthetized in different organs or tissues of the mother plant and then transported to seeds. In *A. thaliana*, the flavonoids proanthocyanidins (PAs) and flavonols are synthetized in seeds (Lepiniec et al., 2006; Corso et al., 2020). In particular, the aglycone and decorated quercetin and kaempferol flavonols are accumulated at high concentration in the seed coat, endosperm and embryo. The localization of flavonols in *A. thaliana* seeds depends on their modifications, particularly on the glycosylation pattern, the monoglycosylated flavonols being accumulated in the seed coat and/or endosperm, and di and tri –glycosylated flavonoids accumulated in the embryo of Arabidopsis seeds (Routaboul et al., 2006). PAs are specifically synthetized in the seed coat, where they are accumulated at high concentrations at early Arabidopsis seed developmental stages, starting from 1-2 days after pollination (Lepiniec et al., 2006). Differently from flavonoids, glucosinolates (GSLs) are synthetized in maternal tissues outside of the seeds and then transported to seeds in *A. thaliana* (Nour-Eldin and Halkier, 2009; Sanden et al., 2023; Xu et al., 2023). Recent results nicely showed that, in *A. thaliana* seeds, GSL glucoiberin and glucoraphanin are synthetized in the funiculus and then transported to the seed embryo, where they accumulate and are subject to modifications (Sanden et al., 2023; Xu et al., 2023). In particular, the side chain of those two methionine-derived aliphatic GSL is hydroxylated by the 2-oxoglutarate-dependent dioxygenase AOP3 to form hydroxylated/hydroxylalkyl GSL (OH-GSL) (Kliebenstein et al., 2001). Previous reports suggested that GSL sinapoylation and benzoylation occurring on the -OH from the side chain structure are catalysed by SCPL17, a serine carboxypeptidase-like, while the synthesis of benzoyl compounds is catalysed by the benzoate-CoA ligase BZO1 (Reichelt et al., 2002; Kliebenstein et al., 2007; Lee et al., 2012; Sanden et al., 2023). Many abundant cinnamic acids in Brassicaceae seeds derive from sinapic acid core structure and decorated metabolites, mainly sinapate esters (COO-R) (Baumert et al., 2005; Corso et al., 2020). In *A. thaliana*, the synthesis of sinapate esters starts with the glycosylation of sinapic

acid to form sinapoyl-glucose (sinapoyl-G), that constitutes the source of sinapate for the production of

other sinapate-esters. These include the conversion of sinapoyl-G to sinapoyl-malate by the

sinapoylglucose: malate sinapoyltransferase (SMT), the synthesis of 1,2-di-sinapoyl-G from two

molecules of sinapoyl-G catalyzed by the sinapoyl-glucose:sinapoyl-glucose sinapoyltransferase or the

synthesis of sinapoylcholine (sinapine) by 1-O-sinapoyl-beta-glucose:choline sinapoyltransferase

(SCT) (Fraser et al., 2007; Stehle et al., 2008; Barreda et al., 2024).

While many of the above cited SMs are essential for Brassicaceae seeds to cope with abiotic and biotic

stresses (Corso et al., 2020), many efforts have been done by breeders to drastically reduce their content

in seeds since some of them are considered antinutritional compounds for human and animal

consumption (Di Gioia et al., 2019). Among them sinapine (an alkaloidal amine) is one of the most

abundant metabolites synthetized in Brassicaceae model and crop species and can constitute up to 2%

of seed weight in these species (Menard et al., 2022). Sinapine reduces protein quality and digestibility

and, because of its bitter taste, makes the meal obtained from Brassicaceae seeds less suitable and

palatable for animal consumption (Bell, 1993; Huang et al., 2008; Menard et al., 2022). Similarly to

other SMs accumulated in seeds, GSLs are part of the arsenal of compounds that plant species use to

defend themselves against insects, vertebrate herbivores and pathogens (Dalling et al., 2020). GSLs,

such as glucoraphanin, glucobrassicin and sinigrin, are considered antinutritional because of their bitter

taste or their potential toxicity due to the accumulation of degradation products such as isothiocyanates,

thiocyanates, nitriles and epithionitriles. Nevertheless, many reports highlighted a wide range of

beneficial effects on human health, including but not exclusively due to their anticancerogenic,

antibacterial, antidiabetic, and antioxidant activities (Di Gioia et al., 2019). In addition, despite GSLs

were considerably reduced in Brassicaceae seed by breeding programs, limited information is available

about the activity and/or toxicity of a large set of aglycone and decorated (e.g. glycosylated, sinapoylated

and benzoylated) GSLs, together with their related degradation products.

Seeds from Brassicaceae crop species are used for oil and protein extraction, and for valorisation of co-

products, including many SMs, but less information is available about SM diversity in these seeds and,

hence, modifications, tissue-distribution and accumulation during seed development, unlike the

extensive studies conducted in *A. thaliana*. Among these species *Camelina sativa* (camelina) is an

oilseed crop cultivated for human and animal nutrition, and for industrial uses, increasingly used as

model for research on Brassicaceae species. The characteristics of camelina make it a valuable crop for

diversifying and intensifying production in agroecological cropping systems (Zubr, 1997; Neupane et

al., 2022). Camelina seeds oil is rich in omega-3, polyunsaturated fatty acids (PUFA) and vitamin E

(tocopherol), and is used in food, cosmetics and as a biofuel (Faure and Tepfer, 2016; Zanetti et al.,

2017). Besides oil, camelina seeds accumulate high levels of SMs, including many phenylpropanoids

and GSLs with a wide range of biological functions (Quéro et al., 2016; Alberghini et al., 2022; Boutet

et al., 2022). *Camelina sativa* is hexaploid (2n=6x=40), with three sets of chromosomes and therefore

three genomes (Kagale et al., 2014). *C. sativa* genome (n = 20, N^6N^7H) originated from a hybridization

between the parental genomes *C. neglecta*–like (n = 13, N^6N^7) and *C. hispida* (n = 7, H) (Mandáková et
al., 2020). In particular, every chromosome or chromosomal region of *A. thaliana*, was represented in
three independent chromosomes in the genome of *C. sativa*. We recently analyzed the seed SM
landscape in camelina seeds, highlighting the accumulation of a large set of SM classes, including GSLs,
flavonoids, cinnamic acids, monolignols, terpenoids and amino acids and derivatives (Boutet et al.,
2022). This work documented major effects of the environment on the stimulation of seed SMs in *C.*
sativa seeds and showed that SMs, including GSLs and phenylpropanoids, showed a higher
environmentally-induced plasticity with respect to most of the primary metabolites, including some
sugars, fatty acids, proteins and lipids (Alberghini et al., 2022; Boutet et al., 2022). While we explored
in detail SM diversity and plasticity, few information is available about SM distribution and expression
of SM-related proteins and/or genes in developing and germinating seed tissues, including coat,
endosperm and embryo, of camelina and other oilseed crop species (Fang et al., 2012). Knowing the
spatiotemporal synthesis, transport and degradation of SMs in a crop species such as camelina would be
indeed of main importance to develop new tools to finely modulate the ratio between beneficial and
antinutritional compounds in seeds.

In this study we used untargeted metabolomics, proteomics and transcriptomics to analyze the synthesis,
transport, modifications and degradations of SMs accumulated in the different seed tissues of *C. sativa*
(i.e. seed coat, endosperm, and embryo), during seed development and germination. This multi-omic
approach allowed to identify specific signatures of SMs, proteins and transcripts associated with the
development of the camelina seed coat/endosperm and embryo. In addition, we highlighted contrasting
seed composition and spatial distributions for the defence-related and antinutritional GSL compounds
among closely related Brassicaceae species: Camelina, Arabidopsis and Brassica. While camelina plants
produced long chain (C8-C11) GSL, Arabidopsis and Brassica accumulate mid- to short-chain (C3-C7)
GSL (Czerniawski et al., 2021; Missinou et al., 2022). Our data further revealed specific GSL
distributions within Brassicaceae seed tissues, with long chain methylsulfinylalkyl GSLs primarily
accumulating in the seed coat and endosperm, while mid- to short-chain methylsulfinylalkyl GSLs were
mainly localized in the embryo.

Results

Camelina sativa **seed coat and embryo development**

Camelina sativa seeds reach maturity in about 50 days from the onset of flowering. Plants were

cultivated in a greenhouse (see materials and methods for more detailed information). Seeds were

harvested at 13, 21, 28, 35, 42 Days After Flowering (DAF), at dry seeds, and at two germination stages

(6 hours and 24 hours from seed imbibition) (Figure 1). Seed coat and endosperm (SCE), and seed

embryo (SE) were separated and used for multi-omic analyses. As for the ratio between the two seed

tissues, at 13 DAF 77% of the seed weight is constituted by the SCE, at 21 DAF the two tissues represent

together 50% of the seed weight, while from 28 DAF the SE represents around 70% of the seed weight

(Figure 1).

Specialized metabolite diversity in camelina seed coats and embryos

Untargeted SM profiling revealed a large diversity in camelina seeds (Figure 2; Table S1 and S2A).

Untargeted metabolomic analyses (positive and negative ElectroSpray Ionization, ESI+ and ESI-)

allowed the detection of 4061 peaks of putative metabolites (i.e. metabolic features) (Table S2B).

Molecular network analyses enabled the clustering of SM compounds according to their MS/MS spectra

(Figure 2A, Table S2A) and was used to improve the annotation of the unknown metabolic features

(Olivon et al., 2018; Boutet et al., 2022). Annotated SMs in camelina seed coats and embryos belong to

several classes, among which amino acids (AA) and derivatives (208 SMs), cinnamic acids (176 SMs)

and flavonols (64 SMs) were the most numerous, followed by nucleosides and derivatives, terpenoids,

flavan-3-ols and proanthocyanidins (PAs), fatty amides, carbohydrates, carboxylic acids, glucosinolates

(GSLs), flavonoids, sulfoxides, lignans, glutathione and derivatives, isothiocyanates (ITCs) and amines

(Figure 2B and Table S2A).

The average intensity of SMs in seed coats and embryos allowed the identification of the top-10

metabolites with highest relative intensity (Table S1). Sinapine (cinnamic acids/alkaloids) was the SM

showing the highest intensity for both seed coat/endosperm (SCE) and embryos (SE). In addition, two

ITCs (9-Methylsulfinylnonyl ITC and 10-Methylsulfinyldecyl ITC) deriving from the degradation of

two major GSL compounds, rutin (flavonol) and adenosine (nucleosides and derivatives) were highly

accumulated metabolites common to seed coats and embryos. Among the SMs with high intensity and

specific to the SCE, there were one ITC (11-Methylsulfinylundecyl ITC), the GSL glucoarabin (9-

methylsulfinylnonyl GSL) and a GSL derivative (p2553; proposed formula: $C_{17}H_{32}NO_7S_2^+$), together

with the epicatechin (flavan-3-ol) and a cinnamic acid with unknown formula (p8). Among the seed-

embryo–specific SMs with the highest intensity there were a triglycosylated flavonol, putatively

annotated as quercetin-(xyloside-rhamnoside-glucoside), a sulfoxide (p29, proposed formula

$C_{13}H_{25}NOS$) and three AA and derivatives (gamma-Glu-Leu, p13 with proposed formula $C_{11}H_{20}N_2O_5S$

and (2S)-2-amino-5-[[(1S)-1-carboxy-2-methylsulfanylethyl]amino]-5-oxopentanoic acid).

Camelina seed specialized metabolome is shaped by the tissue and the developmental stage

The untargeted metabolomic data obtained from seed coats and embryos harvested at six developmental

and two germination stages were used to perform a principal component analysis (PCA). 24,3 % and

19,6 % of the variability observed among the samples can be explained respectively by the axis PC1,

which is associated to the type of tissue, and the axis PC2, which is associated to the developmental

stage (Figure 2C). The PCA performed allowed to clearly separate the samples according to the tissue,

confirming that seed coat/endosperm and embryo present contrasting specialized metabolome

signatures. Hence, Differentially Accumulated Metabolites (DAMs) according to the "developmental

stage" (13 days after flowering (DAF) to dry seeds, and germinating seeds) factor were identified

(ANOVA, FDR<0.05) separately for the two seed tissues considered (SCE and SE). The statistical

analysis highlighted that the accumulation of 62% (2523) and 73% (2973) of SMs was impacted by the

seed developmental stage in the SCE and SE, respectively (Figure 2D).

A core-set of specialized metabolites is regulated during seed embryo and coat-endosperm
development and maturation

In order to identify stage-specific DAMs, a hierarchical clustering analysis was conducted for SCE and

SE tissues using DAMs intensities. The analysis allowed the separation of the DAMs into five clusters

for seed coat-endosperm (I to V; Figure 3A) and four clusters for seed embryo (VI to IX; Figure 3B).

The clusters allowed the identification of DAMs accumulated from early (13 DAF) to late (42 DAF and

dry seeds) seed developmental stages, and of germination-specific DAMs. Afterwards, using the DAMs

belonging to each cluster, an enrichment analysis was performed to identify the metabolic categories

that are characteristics of each tissue and developmental stage. The enrichment analysis was performed

using the metabolic categories identified with the molecular network analysis, both the 'annotated' and

'not-annotated' (or 'unknown') ones (Table S2A, column "Metabolite class"). In particular, not-

annotated metabolites that clustered together in the molecular network analysis (Figure 2A) and

therefore have similar chemical structures, were regarded as a distinct metabolic category. Besides the

SMs assigned to known metabolic categories, several not-annotated SMs showing similar MS/MS

profiles were enriched in different clusters displaying specificities to some seed tissue and/or

developmental stage (Figure 3A and B).

Regarding the SCE specialized metabolome (Figure 3A), carboxylic acids and flavan-3-ols and PAs

were found to be enriched and specific to cluster I, characterized by DAMs with higher accumulations

at 21 DAF and 28 DAF. In addition, glucosinolates, and nucleosides and derivatives SMs were found to

be enriched in cluster II, which included DAMs with high intensity at 13 DAF. Cinnamic acids, AA and

derivatives, flavonols and lignans categories were enriched in cluster III, which was the biggest for SCE

tissues and showed SMs with increasing accumulation from 28 to 42 DAF. Finally, AA and derivatives,

sulfoxides, amines and carbohydrates were enriched in clusters IV and V that are specific to seeds during

germination stages.

Concerning the seed embryo (SE) SMs (Figure 3B), no enriched metabolic categories were found for

SMs from cluster VI, which is characterized by DAMs more accumulated during germination 24h after

imbibition. Cluster VII, which includes DAMs with the highest accumulation mainly at the early seed

developmental stage (13 DAF), is enriched for fatty amides and terpenoids. Cluster VIII is the largest

one and contains SMs with increasing intensity from 35 DAF to dry seeds, and showed an enrichment

for several metabolic classes, including cinnamic acids, AA and derivatives, flavonols, carboxylic acids

and flavan-3-ols and PAs. From work on Arabidopsis we know that flavan-3-ols and derivatives are not

produced in the embryo (Lepiniec et al., 2006) and that these metabolites could originate from degraded

cells of the endothelium and, hence, will not be further discussed in this article. Finally, AA and

derivatives and isothiocyanates (GSLs degradation products) were found to be enriched in cluster IX,

which is characterized by DAMs with higher accumulation predominantly at 21 and 28 DAF.

According to the results highlighted by the enrichment analysis and the broad interest for seed biology

and quality, we will focus on phenylpropanoid (flavonoids, cinnamic acids and monolignols) and GSL

metabolic categories, and on genes or enzyme categories involved in their modifications (i.e.

glycosylation, methylation, hydroxylation and acylation) (Barreda et al., 2024).

Proteomic analyses allowed the identification of SM-related proteins with specific expression pattern during seed development

To get a better understanding of the regulation and accumulation dynamics of SM-related pathways in

camelina SCE and SE, data-independent acquisition (DIA) proteomic combined with ion mobility

spectrometry analyses were performed on the same samples used for untargeted metabolomics. These

analyses allowed to measure the abundance of 11462 proteins (Table S3). The PCA performed with

proteomic data separated the samples according to seed developmental and germination stages (Figure

4A). As for metabolomic data, Differentially Accumulated Proteins (DAPs) impacted by the

"developmental stage" factor were identified (ANOVA, FDR<0.05) separately for SCE and SE (Figure

S1). The statistical analysis highlighted that the accumulation of 14% (995) and 33% (2418) DAPs was

specifically affected by the seed developmental stage in SCE and SE, respectively, while 53% (3912)

of detected proteins were differentially expressed and common to SCE and SE (Figure 4B and S1). Next,

the proteins involved in GSLs, flavonoids, cinnamic acids and monolignols biosynthesis/pathways, and

the enzymes putatively involved in SM modifications that were differentially expressed in SCE (333

DAPs) and SE (374 DAPs) during development, were identified. Among these DAPs, 273 were in

common between SCE and SE (Table S3 and Figure 5). A hierarchical clustering analysis was conducted

for SCE and SE seed tissues using GSLs, flavonoids, cinnamic acids and monolignols –related DAPs

(Figure 5A and 5B). DAPs were separated into six clusters for SCE (Figure 5A) and seven clusters for

seed embryo (Figure 5B), according to their accumulation patterns. In SCE, most DAPs belonging to

selected SM categories were accumulated at earlier seed developmental stages, either at 13 and/or 21

DAF (clusters IV and VI). Similar results were observed for SE, except for proteins involved in SM

glycosylation (belonging to glycosyltransferase GT1 group) that were accumulated at several seed

developmental and maturation stages, particularly during seed germination (cluster VII).

Multi-omic data revealed contrasting glucosinolate landscapes and spatial distribution in seeds of

camelina and other Brassicaceae species

Glucosinolates and their related degradation products were further characterized in Camelina SE and

SCE (Figures 6, 7). The results revealed that GSLs and their degradation products are differentially

accumulated in the SCE and/or the SE of camelina seeds (Figure 7). Most GSLs that were differentially

accumulated during seed development, including the seed-specific aliphatic GSLs 8-methyl-sulfinyl-

octyl-GSL,   glucoarabin   (9-Methylsulfinylnonyl-GSL;   GSL9),   glucocamelinin   (10-

Methylsulfinyldecyl-GSL;   GSL10)   and   gluconesliapaniculatin   (11-Methylsulfinylundecyl-GSL;

GSL11), were strongly accumulated in the SCE and showed the highest accumulation level at early seed

developmental stage (13 DAF) and/or at later stage (35 DAF) (Figure 7A). The indolic GSLs

glucobrassicin and neoglucobrassicin showed their highest accumulation at 13 DAF. In contrast, most

of the GLS degradation products (ITCs) were differentially accumulated only in camelina embryos,

where they reached their peak levels at dry mature seed stage in the SE (Figure 7A, 7B). The contrasting

GSL distributions between the SCE and SE is further highlighted by comparing the relative

accumulation levels of the main annotated and differentially accumulated GSLs and their ITC

degradation products in these tissues (Figure 7B). Glucoarabin (GSL9) was the GSL with the highest

relative abundance, peaking at 35 DAF in SCE. Consistently, the main GSL degradation product, 9-

Methylsulfinylnonyl ITC, reached its maximum relative intensity at 13 DAF in the SCE and in the dry

seed embryo. Gluconesliapaniculatin (GSL11) was the second most abundant GSL, with its peaked at

35 DAF, followed by glucocamelinin (GSL10), which was characterized by two peaks at 13 and 35

DAF (Figure 7B). The corresponding degradation products of GSL10 and GSL11,10-

Methylsulfinyldecyl and 11-Methylsulfinylundecyl ITCs, were differentially and highly accumulated in

the SE of dry mature seeds (Figure 7B). Finally, 8-Methylsulfinyloctylglucosinolate was also specific

to the SCE, but showed lower intensity compared to GSL9, GSL10 and GSL11.

In most cases, the accumulation of proteins involved in GSL side-chain elongation and core-structure

formation was higher at earlier seed developmental stages (13 and 21 DAF) in both SCE and SE (Figure

6). Among the proteins related to GSL modifications, we identified two Flavin-Monooxygenases,

namely FMOGS-OX5 and FMOGS-OX7, putatively implicated in GSL hydroxylation. These proteins

showed the highest accumulation at early (FMOGS-OX5, 13-21 DAF) or late (FMOGS-OX7, 35 DAF

to dry seeds, specific to SE) seed developmental stages. FMOGS-OX5 was also found in the SE during

germination. Proteins playing a role in GSL degradation were accumulated during the whole seed

development and germination in both SCE and SE, with some specificities and exceptions. For example,

PYK10 myrosinase showed the highest accumulation during germination, while TGG1 myrosinase was

accumulated throughout seed development in both SCE and SE, and also during germination specifically
in SCE tissues (Figure 6).

These results mainly highlighted that in *C. sativa* GSLs compounds were unexpectedly accumulated in
both the seed coat and endosperm, while their corresponding degradation products, ITCs, were present
at high level in the embryo. Next, GSL and ITC accumulations were characterized in the seed
coats/endosperms and embryos of two others Brassicaceae species, *A. thaliana* and *B. napus*. These
results showed that camelina seeds are characterized by different GSL accumulation patterns and
expression of related genes/proteins compared to *A. thaliana* and *B. napus* (Figure 8a; Table S4). **First**,
we showed that *C. sativa* exclusively accumulated long carbon(C)-chain GSLs, notably C9-C11
methylsulfinylalkyl GSLs, in the seed coat and the endosperm, while Arabidopsis and Brassica seeds
did not synthetize these compounds (Matsuda et al., 2010; Missinou et al., 2022). **Second**, we showed
that the long chain C8 methylsulfinylalkyl GSLs are accumulated in the seed coat and endosperm of *C.*
sativa and in *A. thaliana*, but are not synthetized and accumulated in *B. napus* seeds. **Third**, mid- and
short-chain C3-C7 methylsulfinylalkyl GSLs are accumulated in the embryo of *B. napus* and in *A.*
thaliana, but are not synthetized and accumulated in *C. sativa* seeds. **Fourth**, ITCs (GSL degradation
products) are accumulated in the embryo in *C. sativa* and in *B. napus*, and in the seed coat and endosperm
in *A. thaliana*. **Fifth**, benzoylated and sinapoylated GSLs were found exclusively in *A. thaliana* seeds.
In accordance with this finding, no Camelina orthologue of the SCPL17 acyltransferase enzyme,
involved in GSL side-chain benzoylation and sinapoylation in *A. thaliana*, has been found in our
proteomic data (Figure 6, Table S3 and Table S5). These multi-omic data revealed a complex and fine-
tuned regulation of GSL synthesis, modification and transport that has been summarized in Figure 8b.

Multi-omic data highlighted seed coat-endosperm and embryo -specific accumulation patterns of
proteins and metabolites related to glucosinolates, flavonoids and cinnamic acids
To get a better overview about other major SM categories, the accumulations of metabolites and proteins
belonging to flavonoids (Figures 9) and cinnamic acids and monolignols (Figures S2 and S3) categories
were further analysed in camelina seed tissues. Given that *C. sativa* has a hexaploid genome, three
homeolog genes could potentially encode proteins involved in a specific enzymatic activity. The
proteomic results showed in this study allowed the identification of the proteins copies specifically
accumulated in SE and SCE and involved in SM biosynthesis, modifications and degradation (Table S3;
Figure 5). For GSLs, cinnamic acids and monolignols, one gene copy was expressed for most proteins
involved in the same enzymatic activity in both SCE and SE. In contrast, flavonoid-related proteins were
mostly accumulated with one gene copy for SCE and two copies for SE (Figure S4).
As observed for GSLs, the accumulation of different flavonoid classes showed marked differences
depending on the seed tissue and the developmental stage (Figure 9B). In particular, in SCE tissue
flavan-3-ols and derivatives were accumulated from 13 to 28 DAF in their aglycone form (epicatechin,

catechin and procyanidin B1 and B2), while glycosylated flavan-3-ols (epicatechin and catechin glycosides) showed the highest accumulation level at 35 and 42 DAF. Most flavonol metabolites, including quercetin and kaempferol mono-, di- and tri-glycosides, were accumulated at later seed developmental stages in SCE (35-42 DAF) and SE (dry mature seeds). In contrast, isorhamnetin (i.e. methylated quercetin) -glycoside was predominantly accumulated at earlier developmental stages in both tissues (Figure 9B). Interestingly, when the isorhamnetin presented a second -glycosyl group (isorhamnetin-(Rha-Glc), it was mainly accumulated at later seed developmental stages in both tissues (Figure 9B). Enzymes involved in the biosynthesis of the core flavonoid pathway and in flavonol specific pathway were expressed at 13-21 DAF in both SCE and SE, with a shift between the two tissues for some of them (Figure 7). This is the case of flavonol synthase 1 (FLS1) which showed the highest expression at 21 DAF in the SCE and at 13 DAF in the SE. On the contrary, flavonol synthase 3 (FLS3), showed an earlier expression in SCE compared to the SE. Anthocyanidin reductase (ANR), involved in flavan-3-ols synthesis showed its highest expression in the SCE at early developmental stages. As concerns proteins involved in flavonoid glycosylation, it is worth mentioning that the UDP-dependent glycosyltransferase (UGT), F7GlcT, was specific to the embryo and showed the highest expression during the germination stage.

While the metabolite landscape and composition of GSLs and flavonoids differed markedly between the SCE and SE, fewer differences between in cinnamic acid composition were observed between these two tissues (Figure S3). Sinapine was the most abundant metabolite belonging to this category in both tissues (Table S1), and showed the highest intensity at 35-42 DAF for SCE and in dry seeds for SE. Similarly, other cinnamic acid derivatives, e.g. sinapoyl, feruloyl and coumaroyl glycosides, were accumulated at 35-42 DAF in the SCE and at dry seeds in the embryo. In addition, guaiacyl-ferulic acid-glucose was more abundant during germination in SCE and SE. Some differentially accumulated proteins involved in cinnamic acids synthesis or decoration showed specific expression in seed embryo (Figure S2). This is the case of the sinapic acid:UDP-glucose glucosyltransferase (SGT1), involved in sinapic acid glycosylation, of the Sinapoylglucose:malate sinapoyltransferase (SMT), which produced sinapoyl-malate, and of 1-O-sinapoyl-beta-glucose:choline sinapoyltransferase, catalysing the synthesis of sinapine.

Transcriptomic analyses in seed embryo.

Transcriptomics were performed on a subset of samples compared to those used for metabolomic and proteomic analyses. In particular, seed embryo at 13, 21, 28, 35, 42 DAF and at 6 hours after germination stage were subjected to RNA-Seq analyses (Figure 10, Figure S5 and Table S5). As observed for proteomic and untargeted metabolomic analyses, large differences in gene expression (FDR < 0.05) were observed between seed developmental stages, and between seed development at 42 DAF and at 6 hours of germination. The PCA confirms these results and the quality of the data obtained since it allowed a clear separation of embryos from different developmental stages and showed that the three

biological replicates were clustered together (Figure 10). Next, we extracted Differentially Expressed Genes (DEGs) involved in GSL, flavonoid, and cinnamic acid synthesis, transport, decorations (or modifications), degradation and regulation (Figure 10 and Figure S6). Many DEGs related to GSLs were expressed at 13 DAF and showed generally an earlier expression peak compared to what has been observed for corresponding proteomic data. As for flavonoids, all DEGs involved in their biosynthesis and most DEGs involved in their decoration showed the highest expression level at 13 DAF. *MYB11* and *MYB111* transcription factors (TFs) involved in the regulation of flavonol metabolites, were expressed at 13DAF, while *TTG1* TF showed its expression peak at germination stage. DEGs related to cinnamic acids were characterized by larger differences concerning their expression pattern. In particular, genes coding for Serine-carboxypeptidase-like enzymes involved in sinapic acid ester biosynthesis showed the highest expression level at 13 DAF (*SCPL12-FTP1*), 21 DAF (*SCPL9-SNG2-SCT*), 28 DAF (*SCPL7*), 42 DAF (*SGT1*) and during germination for *SST* (*SCPL9*).

Discussion

The accumulations of SMs has a significant impact on the quality of plant-derived products. While the spatial distribution of these compounds in plant leaves and roots tissues has been described for some model and crop species (Brown et al., 2003; Czerniawski et al., 2021), very limited information and mainly restricted to primary metabolites is available about their spatial accumulation in seed tissues (Zhang et al., 2021). Given their major impact on seed quality, elucidating SM distribution in seed coat, endosperm and embryo during development and germination is of main interest in Brassicaceae species, including the oilseed crop *C. sativa* (Kagale et al., 2014; Faure and Tepfer, 2016; Zanetti et al., 2021). In the present study we combined untargeted metabolomic, proteomic and transcriptomic analyses to unveil seed SM accumulation (synthesis, transport and/or modifications/decorations) in developing seed coat and endosperm, and seed embryo. Our results emphasized a large diversity of SM in camelina seeds, as shown in a previous study (Boutet et al., 2022).

A progressive but significant evolution of the specialized metabolome can be observed during embryo development. The sample replicates of the earliest stage of seed development studied (13 DAF) and the samples replicates of the latest seed development stage (dry seeds) are the most distant; and intermediary developmental stage samples are distributed between these two samples, following a chronological order. Interestingly, the embryo specialized metabolome of 6h and 24h germinating seeds appear to be similar to those of seed embryo in the early and middle seed developmental stages, respectively (Figure 2C). Hence, these data suggest that there is a considerable specialized metabolome change/shift occurring in the embryo from dry seed to germinating seed and that the metabolome is afterwards following an evolution equivalent to the one that takes place during seed development. Similar results have been observed for the SCE, for which seeds harvested at 13 DAF and germinating seeds at 24h after imbibition showed a similar specialized metabolome.

RNA-Seq transcriptomic analysis revealed that the expression of a majority of genes involved in the biosynthetic pathways of GSLs, flavonoids, cinnamic acids, and monolignols peaks at the early seed developmental stage (13 DAF) in the embryo (Figure 10 and Figure S6). Consistently, a large part of the enzymes involved in the synthesis of GSL, flavonoid, cinnamic acid and monolignol were accumulated in the embryo between 13 DAF to 21 DAF (Figure 5B), reflecting a delay between gene expression and the accumulation of the corresponding proteins. Interestingly, SM-related enzymes from SCE were also mostly accumulated at early stage of seed development (Figure 5B; Table S3). However, SMs displayed contrasting accumulation patterns between both seed tissues (Figure 3). Indeed, whereas both SE and SCE had some groups of SMs highly accumulated at 13 DAF, a large part of SCE SMs were mainly accumulated at 35 and 42 DAF stages, whereas the majority of embryo SMs were highly accumulated at dry seed stage (Figure 3). Both of those SCE and SE SM clusters were enriched in AA and derivatives, cinnamic acids and flavonols, indicating that those metabolic categories are synthesised and/or stored in SCE during seed development and transported to SE between the maturation (42 DAF) and dry seed stage (Figure 3). Hence, those results emphasize the importance of SCE tissues, at the

interface between the mother plant and the embryo, in the support of embryonic growth (Yan et al., 2014). PCA analyses on untargeted metabolomic data highlighted that, while both factors are impacting SMs diversity, metabolite signature seems more similar within the same tissue during seed development than in different tissues at a specific developmental stage (Figure 2C). For example, proanthocyanidins (PAs) are specifically accumulated in *Arabidopsis thaliana* seed coat from the early seed developmental stages (1-2 DAF) and reach the maximum peak at 6-7 DAF for Col-0 ecotype and 12 DAF for Cvi-0 ecotype (Routaboul et al., 2012; Corso et al., 2020). In addition, PA accumulation is paralleled by the expression of the *BANYULS* gene, which is involved in their synthesis (Debeaujon et al., 2003). This trend is confirmed in camelina seeds, where the anthocyanidin reductase (ANR) protein, encoded by a *BANYULS* homolog (Xie et al., 2003), showed the highest expression peak at 13 DAF in SCE samples (Figure 9A) and, consistently, the flavan-3-ols derivatives (catechin, epicatechin and procyanidin) are highly accumulated at 13, 21 and 28 DAF (Figure 9B). Differently from the aglycone forms, catechin and epicatechin glucosides (as most glycosylated and acylated flavonoids) are accumulated at later stages of seed coat development in camelina, as shown for *Vitis vinifera* seeds (Zerbib et al., 2018) (Figure 9B), consistent with an increase stability of these glycosylated metabolites, as previously shown by Raab et al. (2010).

Besides flavonoids, the defensive and antinutritional GSLs and related ITC compounds, deriving from GSL degradation, showed contrasting accumulation pattern when comparing SCE to embryo tissues, in camelina seeds (Figure 6). While some GSLs have a negative impact on seed oil and protein quality and digestibility, and are in some cases toxic for animals and humans, others are powerful antioxidants with nutritional and health benefits (Traka, 2016; Di Gioia et al., 2019). Hence, they have been a target of domestication for some Brassicaceae (e.g. Brassica and Camelina species) for which many efforts are still being done to modulate the content or nature of GSLs in seeds (Nour-Eldin et al., 2017; Hölzl et al., 2023; Mann et al., 2023). Nevertheless, GSLs are antioxidants accumulated in response to abiotic stresses (Rao et al., 2021; Mácová et al., 2022) and are essential for plant resistance to pathogens, so much so that their systematic elimination could prove counterproductive.

Our data highlighted that the different GSL spatial distribution, accumulation and, hence, functions in seeds, observed among *C. sativa*, Arabidopsis and Brassica species were mainly due to GSL side-chain length, GSL transport and potentially GSL chemical modifications (decorations). Long chain C8-C11 methylsulfinylalkyl GSLs, synthetized in camelina (C8-C11) and Arabidopsis (only C8), were indeed accumulated in the seed coat and endosperm, while mid- and short-chain C3-C7 methylsulfinylalkyl GSLs were accumulated in the embryo of Brassica and Arabidopsis species studied in this work. These differences could be due to the affinity of GSL transporters for metabolites depending on the side-chain length.

The contrasting pattern of GSL and ITC accumulation among these three Brassicaceae species could be partially due to the absence of benzoylated and sinapoylated GSLs in camelina and brassica

seeds, whereas these compounds accumulate at high level in Arabidopsis embryos (Kliebenstein et al., 2007; Lee et al., 2012; Bussell et al., 2014; Sanden et al., 2023). In particular, sinapoyl- and benzoyl-GSLs could be more stable and less toxic than non-decorated GSLs and might impact their degradation to ITC and nitriles and the transport to other seed tissues and/or cellular/vacuolar compartments, thus facilitating their accumulation in the embryo. Nevertheless, no clear information is available regarding the biological activity and role of acylated-GSLs in seeds, which deserves further investigation. Regarding the distribution of ITCs, it could be suggested that long-chain ITCs (C>8) can be accumulated in the embryo rather than the SC because they may present lower toxicity levels than shorter ITCs. Indeed, some reports show that the ITC toxicity against pathogens varies depending on ITC chemical structure, including the carbon chain length (Matthäus and Zubr, 2000; Wilson et al., 2013; Abdel-Massih et al., 2023).

Interestingly, fewer SMs showed peaks of accumulation at germination stages (6h Germ and 24h Germ) compared to the seed developmental stages for both tissues (Figure 3). This suggests that many SMs are degraded rather than synthetized during germination. Similarly, there were fewer proteins strongly accumulated at germination stages compared to seed developmental stages for both tissues (Figure 5). However, it is worth mentioning that glycosyltransferases were found to be highly expressed in SE at both 6h and 24h germination stages (Figure 5). This could be linked to starch and sugar remobilization occurring during the germination and establishment of the seedling. Likewise, carbohydrates metabolic category was found to be specifically enriched upon seed germination (Figure 3). Moreover, 9-Methylsulfinylnonyl, 10-Methylsufnyldecyl and 11-Methylsulfinylundecyl ITCs were found to decrease progressively from dry seed stage to 24h Germination stage (Figure 7). This decrease of ITCs was concomitant with an increase in expression of several enzymes involved in GSL degradation, and thus potentially ITC production (Figure 10), suggesting that ITCs may have a role during camelina seed germination by providing for instance an important source of AA, sulfur atoms or other molecules.

To conclude, GSLs, flavonoids, cinnamic acids, and monolignols are largely synthetized and accumulated in camelina seeds and, therefore, determine their nutritional quality and play a key role in seed interaction with biotic and abiotic environments. Characterizing the spatial dynamics of seed SMs will contribute to the development of crops with an optimized distribution of beneficial and toxic metabolites for seeds quality and animal nutrition or to the development of new strategies (e.g. fractionation or germination) to reduce the level of undesired compounds after harvest.

     Experimental procedures

Plant culture and seed harvest

Camelina sativa (L.) "Céline" genotype was used for the plant culture and analyses. Seeds of the three

biological replicates (10 plants/replicate) were sown with a gap of two weeks between each replicate

and were cultivated in different part of the greenhouse. Plants were cultivated in 16 cm diameter pots

(from TEKU) in a greenhouse: 16h light (23-25 °C) / 8h dark (15°C) (Osram Powerstar E40 HQI-BT

400W lights). Relative humidity room was set on 45%. A nutrient solution containing 140 milligrams

of nitrogen per liter (140 mg N L^{-1}) was used to irrigate the plants.

Flowers were tagged just before opening to count opened flowers and harvest seeds at defined stages

(Sakai et al., 2018). Seed embryos and coats/endosperms were separated and harvested at 13, 21, 28, 35,

42 DAF, at dry mature seed stage, and at 6 hours and 24 hours, after imbibition, during germination.

For seed pictures acquisition, siliques from each time-point were carefully dissected to harvest the seeds.

Chloral hydrate solution (chloral hydrate:H_2O_2:glycerol (8:2:1, w/v/v) was used to bleach seed coats and

fix the samples that were placed between slide and slip (Boisson et al., 2001). Seeds were then stored

for 24h to 72h at 4°C before imaging. Images were acquired on Axioplan 2 imaging epifluorescence

microscope after several weeks of seed coat clearing (Zeiss, Göttingen, Germany).

Arabidopsis thaliana "Columbia-0" (Col-0) genotype was used in this study. Seeds were sown directly

in fertilized soil (3 biological replicates, three plants/replicate). Plants were cultivated in 7.5 x 7.5 cm

pots in a growth chamber with long day photoperiod: 16h light (21°C) - 8h dark (19°C). Dry seeds were

harvested and kept at -20°C until dissection.

For *Brassica napus* species, "Aviso" and "Major" genotype, showing contrasting glucosinolate

accumulation, were used. Seeds were sown directly in fertilized soil. After 4 weeks the plants were

transferred in a vernalisation room for 10 weeks (7,5 x 7,5 cm pots, at 4°C, with a 16/8h day/night

photoperiod). Next, plants were transferred in 19L pots and cultivated during 5 months in a growth

chamber with long day photoperiod: 14h of light (21°C) and 10h dark (13°C). Plants were watered 2

times a day with a nutritional solution (140mg of nitrogen / liter). Dry seeds were harvested and kept at

-20°C until the dissection.

Specialized metabolites extraction and analysis

Metabolite extraction and injection. Untargeted metabolomic data analyses and date were described

in (Boutet et al., 2022). Briefly, metabolites were extracted from 20 mg of seed coats and embryos using

a MeOH: Methyl-tert-butyl: H_2O (1:3:1) buffer, while polar and semipolar metabolites were separated

from the oil and protein fractions using a MeOH: H2O (1:3) buffer. Polar and semipolar metabolite

fraction was used for metabolomic analyses. Untargeted metabolomic data were acquired using a

UHPLC system (Ultimate 3000 Thermo) coupled to quadrupole time of flight mass spectrometer (Q-

Tof Impact II Bruker Daltonics, Bremen, Germany). For chromatographic separation a Nucleoshell RP

18 plus reversed-phase column was used. Samples were injected in positive and negative ionisation

mode (ESI+ and ESI-).

Data processing and ion annotation. ESI+ and ESI- were processed using MZmine 2.52 software

(http://mzmine.github.io/).

Metabolite annotation was performed in four steps: i) LC-MS/MS data were compared with a home-

made library (IJPB chemistry and metabolomic platform) containing more than 400 standards or

experimental common features; ii) LC-MS/MS data were also searched against the available MS2

spectral libraries (Massbank NA, GNPS Public Spectral Library, NIST14 Tandem, NIH Natural Product

and MS-Dial); iii) a molecular network analysis was used to assign not-annotated metabolites to a

chemical family (see Boutet *et al.*, 2022 and Olivon *et al.*, 2018 for further information); iv) Sirius

software (https://bio.informatik.uni-jena.de/software/sirius/) was used to assign a putative annotation to

metabolic features that were not annotated during the previous steps. Raw data were normalised on the

internal standard (Apigenin) and weight of seeds used for the extraction.

Statistical analyses of metabolomic data. Statistical analyses were performed using Metaboanalyst 5.0

software (Pang et al., 2021; Pang et al., 2022). In particular, an ANOVA has been conducted to identify

differentially accumulated metabolites among seed developmental stages, separately for SCE and SE

(adjusted p value < 0.05).

Transcriptomic analyses (RNA-Seq)

RNA extraction. RNAs were extracted with the PicoPure™ RNA Isolation Kit (ThermoFisher)

following the corresponding protocol with slight modifications. Briefly, 2 to 5 mg of seeds were grinded

with some polyvinylpolypyrrolidone (PVPP; phenolic compound absorbing agent) in liquid nitrogen.

100 µL of extraction buffer was added to each sample. Following, samples were vortexed, incubated at

42°C for 30 min (500 rpm) using a ThermoMixer™ C (Eppendorf), and centrifuged (2 min, 3000 g).

Supernatants were collected, without picking-up the pelleted material, and placed in the pre-conditioned

RNA purification columns to be centrifuged (1 min, 16 000 g). The cell extracts obtained were mixed

with 100 µL of 70 % ethanol by pipetting up and down. The mixtures were placed into the pre-

conditioned purification columns which were centrifuged 2 min at 100 g, to allow RNA binding to the

columns, immediately followed by a centrifugation to remove flow-through (30 sec, 16 000 g). The

columns were washed by adding 100 µL of wash buffer 1 and centrifugation (1 min, 8 000 g). DNase

treatment was made by adding 40 µL of DNase solution mix (10µl DNase I Stock solution + 30µl Buffer

RDD; RNase-Free DNase Set, Qiagen) to each column, followed by 15 min of incubation (ambient

temperature) and washing of the column (addition of 40 µL of wash buffer 1 and centrifugation 30 sec

at 8 000 g). Columns were washed twice with 100 µL of wash buffer 2 (2 min at 8 000 g) and a last

centrifugation was made after flow-through waste discard to remove all traces of wash buffer (1 min,

16 000 g). 12 µL of elution buffer were added to each column, which were incubated 1 min at ambient

temperature before RNA elution centrifugations (1 min at 1 000 g followed by 1 min at 16 000 g).

Obtained RNAs were stored at -80 °C until sequencing.

RNA sequencing. Sequencing of messenger RNAs (mRNAs) was performed by BGI Genomics (Hong

Kong, https://www.bgi.com). Prior to cDNA library construction (Eukaryotic Strand-specific

Transcriptome), quality and concentrations of mRNAs were checked again with the bioanalyzer from

BGI platform. The sequencing (paired-end reads of 150bp, 70M clean reads per sample) was performed

with DNA Nanoballs Technology (DNBSEQ^{TM}). After sequencing, raw reads were filtered by the BGI

platform with the filtering software SOAPnuke (version 1.5.2, https://github.com/BGI-

flexlab/SOAPnuke) (Cock et al., 2009; Chen et al., 2018) (Parameters used: l 15 -q 0.2 -n 0.05) to reach

a Phred+33 fastq quality score.

Annotation and statistical analysis of transcriptomic data. High quality raw reads (Céline strain)

were aligned against the *Camelina sativa* genome (strain: DH55, genome assembly:

GCA_000633955.1, annotation: https://www.camelinadb.ca/downloads/: gff and fasta to be made

available if no more available in this site) using GSNAP (gmap-2021-12-17) with a set of options

allowing to find novel splicing and to be tolerant to SNP between DH55 and Céline (VCF files, to be

made available). htseq-count (HTSeq-0.12.4-python2) was used with the following options (-m

intersection-nonempty --secondary-alignments=ignore -a 20 -s no) to count the number of reads by gene.

Orthologous genes of *Arabidopsis thaliana* were searched in *Camelina sativa* using Plant.ensembl

website (using Biomart) and CamRegBase (synthelogs data).

Statistical analyses were performed to identify Differentially Expressed Genes using the Edge R package

(Robinson et al., 2010). Statistical analyses were performed to identify genes that were differentially

expressed among developmental and germination stages in the SCE and SE separately (FDR < 0.05).

Proteomic analyses (DIA)

Sample preparation and analysis. Five µg of each protein extract were prepared using a modified Gel-

aided Sample Preparation protocol (https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4409837/).

Samples were digested with trypsin/Lys-C overnight at 37°C. For nano-LC fragmentation, protein or

peptide samples were first desalted and concentrated onto a µC18 Omix (Agilent) before analysis.

The chromatography step was performed on a NanoElute (Bruker Daltonics) ultra-high-pressure nano

flow chromatography system. Approximatively 200ng of each peptide sample were concentrated onto a

C18 pepmap 100 (5mm x 300µm i.d.) precolumn (Thermo Scientific) and separated at 50°C onto a

reversed phase Reprosil column (25cm x 75µm i.d.) packed with 1.6µm C18 coated porous silica beads

(Ionopticks). Mobile phases consisted of 0.1% formic acid, 99.9% water (v/v) (A) and 0.1% formic acid

in 99.9% ACN (v/v) (B). The nanoflow rate was set at 250 nl/min, and the gradient profile was as

follows: from 2 to 30% B within 70 min, followed by an increase to 37% B within 5 min and further to

85% within 5 min and reequilibration.

MS experiments were carried out using a TIMS-TOF pro mass spectrometer (Bruker Daltonics) with a

modified nano electrospray ion source (CaptiveSpray, Bruker Daltonics). A 1400 spray voltage with a

capillary temperature of 180°C was typically employed for ionizing. MS spectra were acquired in the

positive mode in the mass range from 100 to 1700 m/z and 0.60 to 1.60 1/k0 window. In the experiments

described here, the mass spectrometer was operated in PASEF DIA mode with exclusion of single

charged peptides. The DIA acquisition scheme consisted of 16 variable windows ranging from 400 to

1200 m/z.

Protein identification. Database searching and LFQ quantification (using XIC) was performed using

DIA-NN (version 1.6.1) (Demichev et al., 2020). The UniProt *Camelina sativa* database was used for

library-free search / library generation. For RT prediction and extraction mass accuracy, we used the

default parameter 0.0, which means DIA-NN performed automatic mass and RT correction. Top six

fragments (ranked by their library intensities) were used for peptide identification and quantification.

The FDR was set to 1% at the peptide precursor level. The variable modifications allowed were as

follows: Nterm-acetylation and Oxidation (M). In addition, C-Propionoamide was set as fix

modification. "Trypsin/P" was selected. Data were filtering according to a FDR of 1%. Cross-run

normalisation was performed using RT-dependent.

Identification of differentially expressed proteins. To quantify the relative levels of protein

abundance between different groups, datas from DIA-NN were then analysed using DEP package from

R. Briefly, proteins that are identified in 2 out of 3 replicates of at least one condition were filtered,

missing data were imputed using random draws from a manually defined left-shifted Gaussian

distribution and differential enrichment analysis was based on linear models and empherical Bayes

statistic. A 1.2-fold increase in relative abundance and a 0.01 pvalue were used to determine enriched

proteins. ANOVA were performed from PERSEUS to determine those enriched proteins. A 1.2-fold

increase in relative abundance and a 0.05 FDR were used.

Glucosinolate distribution in Brassicaceae seeds

Seed embryos and coats/endosperms were separated from dry seeds of *Arabidopsis thaliana* (L.)

"Columbia-0" (Col-0) ecotype and *Brassica napus* (L.) "Major" and "Aviso" genotypes.

Specialized metabolite extraction and metabolomic analyses were performed as presented previously

(see "Specialized metabolites extraction and analysis"). 200 seeds and 25 mg were extracted for

Arabidopsis and Brassica samples respectively.

Arabidopsis LC-MS/MS data were first normalized according to the seed tissue proportion (Seed coat

and endosperm: 10 % and Seed embryo: 90 %), following data were normalized by the total sum of

accumulation per sample in each mode (ESI + and ESI -), multiplied by 1000.

Clustering analyses

The R package 'pheatmap' was used to perform hierarchical clustering analysis (hclust) and create

heatmaps using omics data.

Acknowledgements

This work was supported by AgroParisTech (grant "Appel à projets scientifiques 2020" to M.C. and L.

R.), the "Département de Biologie et Amélioration des Plantes" (BAP) of INRAE (grant "Appel à projets

scientifiques BAP 2020" to M.C.), by the project 'Seed MetSpe' of the Labex Saclay Plant Sciences-

SPS (ANR-10-LABX-0040-SPS to L.L. and M.C.) and by a project financed by Plant2Pro Carnot

institute ('BrassiMet' to M.C.).

The IJPB benefits from the support of Saclay Plant Sciences-SPS (ANR-17-EUR-0007). This work has

benefited from the support of IJPB's Plant Observatory platforms PO-Chem and PO-Plants. We are

especially thankful to Patrick Grillot (IJPB, INRAE-Versailles) for his help during the plant culture in

the greenhouse.

The authors would like to thank as well Dr Mark Tepfer (retired, former IJPB, INRAE) for the

constructive discussion on *Camelina sativa* species.

Conflicts of interest

The authors declare that they have no competing interests.

Data availability

The metabolomic data and metadata on *Camelina sativa* have been deposited at the MassiVE data

repository portal with the identifier MSV000094688 (doi:10.25345/C5HH6CH59).

The transcriptomic data have been deposited at the National Center for Biotechnology Information

(NCBI) Transcriptome Shotgun Assembly Sequence Database (TSA) with BioProject identification

PRJNA835304.

The proteomic data have been deposited at the iProX (integrated Proteome resources) data repository

portal with the identifier IPX0008673001

The metabolomic data and metadata on *Arabidopsis thaliana* have been deposited at the MassiVE data

repository portal with the identifier MSV000095844 (doi:10.25345/C51J97K84).

The metabolomic data and metadata on *Brassica napus* have been deposited at the MassiVE data

repository portal with the identifier MSV000095843 (doi:10.25345/C5599ZD2H).

Author contributions

MC and LR directed and designed the research. MC, DDV, TF, DG and CB grew the plants and

harvested seed coat and embryos at defined developmental stages. SB and JCT generated the LC-

MS/MS data and performed the initial quality analyses on the metabolomic data. SB, FP, LB, NK, CB

and JCT performed the metabolites annotation with MetGem, SIRIUS and using MS/MS spectra. MC,

LB, CB, SB, FP, NK and LL analysed and interpret the untargeted metabolomic data. BC prepared the

samples for proteomic analyses. BB performed the proteomic analyses. MC analysed transcriptomic

data. MC, LB and CB performed the statistical analyses on multiomic data. MC, LB and CB generated

all Fig.s and supplemental material. MC and LB wrote the paper, which was edited by CB, LR and LL.

All the authors commented on and approved the manuscript.

 Figure legend

Figure 1. *Camelina sativa* seeds used in this study.

Seed coat and embryo samples used in this study were harvested at 13, 21, 28, 35, 42 days after

flowering. Dry mature and germinating seeds (6h and 24h) were also used.

Figure 2. Untargeted metabolomics in developing and germinating SE and SCE

A) The molecular networks are shown for metabolite analyses carried out in positive (ESI+) and

negative (ESI-) ionization modes. Different colours correspond to different metabolic classes.

Metabolites are grouped based on their chemical structures. Cosine similarity scores of 0.85 and 0.65

were used for ESI+ and ESI-, respectively. **B)** Histogram showing the annotated/known metabolic

categories. The number of metabolites that belong to each category is indicated in the figure. Categories

with less than two metabolite features are not shown. **C)** Principal component analyses using the

untargeted metabolomic data of SCE and SE harvested at several developmental stages and during

germination. DAF = Days After Flowering. **D)** Venn diagram indicating the number of Differentially

Accumulated Metabolite features (DAMf) according to the tissue and developmental stage factors.

Figure 3. Specialized metabolites signatures and stage-specific enrichment of metabolic categories

during SE and SCE development and germination

Hierarchical clustering and enrichment analyses of Differentially Accumulated Metabolites (DAMs)

specific to SCE **(A)** and SE **(B).** Annotated and unknown enriched (hypergeometric test, p < 0.05)

metabolic categories of DAMf clusters specific to seed coats and embryos are showed.

Figure 4. Proteomic analyses in developing SCE and SE

A) Principal component analyses using the proteomic data of SCE and SE harvested at several

developmental stages and during germination. DAF = Days After Flowering. **B)** Histogram indicating

the number of Differentially accumulated proteins (DAPs) according to the developmental stage in seed

coat and embryo.

Figure 5. Accumulation pattern of specialized metabolites -related proteins

A) Hierarchical clustering of SM-related DAPs in SCE and SE. **B)** Expression behaviour of proteins

involved in the biosynthesis, transport and degradation of the major specialized metabolite pathways,

and of the main SM-decoration/modification categories. GT1 = glycosyltransferases, group 1.

Figure 6. Expression of proteins involved in glucosinolate pathways

Expression of proteins involved in GSL biosynthesis, side-chain elongation and modifications, and

degradation in camelina SCE and SE. Only differentially accumulated proteins (DAPs, ANOVA, FDR

< 0.05) are showed in the figure.

Figure 7. Accumulation profiles of glucosinolates and isothiocyanates metabolites in SCE and SE

A) Heatmap showing the glucosinolates and isothiocyanates degradation products that are differentially

accumulated during seed development in SCE and SE tissues. **B)** Histograms showing the major

glucosinolates and isothiocyanates specialized metabolites accumulated in SCE and/or SE.

Figure 8. Distribution of glucosinolates and related degradation products in different Brassicaceae

species

A) Heatmap showing the glucosinolates and related degradation products present in *Camelina sativa*,

Brassica napus and *Arabidopsis thaliana* seed coat/endosperm and embryo tissues.

B) Model summarizing GSL metabolites distribution and expression of related enzymes in seed tissues

of *C. sativa*, *B. napus* and *A. thaliana*. Abbreviations: I3M, 3-indolylmethyl GSL (Glucobrassicin);

1OHI3M GSL, 1-hydroxyindol-3-ylmethyl GSL; 4OHI3M GSL, 4-hydroxyindol-3-ylmethyl GSL;

1MOI3M GSL, 1-methoxyindole-3-ylmethyl GSL (Neoglucobrassicin); 4MOI3M GSL, 4-

methoxyindole-3-ylmethyl GSL; 3MTP GSL, 3-methylthiopropyl GSL (Glucoiberverin); 4MTB GSL,

4-methylthiobutyl GSL (Glucoerucin); 5MTP GSL, 5-methylthiopentyl GSL (Glucoberteroin); 6MTH

GSL, 6-methylthiohexyl GSL (Glucolesquerellin); 7MTH GSL, 7-methylthioheptyl GSL; 8MTO GSL,

8-methylthiooctyl GSL; 9MTN GSL, 9-methylthiononyl GSL; 10MTD GSL, 10-methylthioecyl GSL;

11MTU GSL, 11-methylthioundecyl GSL; 6'Sin-4MTB, 6'sinapoyl-4-methylthiobutyl GSL; 3MSOP

GSL, 3-methylsulfinylpropyl GSL (Glucoiberin); 4MSOB GSL, 4- methylsulfinylbutyl GSL

(Glucoraphanin); 5MSOP GSL, 5-methylsulfinylpentyl GSL (Glucoalyssin); 6MSOH GSL, 6-

methylsulfinylhexyl GSL (Glucohesperin); 7MSOH GSL, 7-methylsulfinylheptyl GSL (Glucoibarin);

8MSOO GSL, 8-methylsulfinyloctyl GSL (Glucohirsutin); 9MSON GSL, 9-methylsulfinylnonyl GSL;

10MSOD GSL, 10-methylsulfinyldecyl GSL; 11MSOU GSL, 11-methylsulfinylundecyl GSL; 6'Sin-

4MSOB, 6'sinapoyl-4-methylsulfinyl GSL; 3OHP GSL, 3-hydroxypropyl GSL; 4OHB GSL, 4-

hydroxybutyl GSL; 3BUT GSL, 3-butenyl GSL (Gluconapin); 2OH3BUT GSL, (2R)-2-hydroxy 3-

butenyl GSL (Progoitrin); 3BZO GSL, 3-benzoyloxypropyl GSL (Glucomalcomiin); 3SIN GSL, 3-

sinapoyloxypropyl GSL; 4BZO GSL, 4-benzoyloxybutyl GSL; 4SIN GSL, 4-sinapoyloxybutyl GSL;

6'Bnz-4BZO GSL, 6'benzoyl-4-benzoyloxybutyl GSL; 3MSOP ITC, 3-methylsulfinylpropyl ITC

(Iberin); 4MSOB ITC, 4- methylsulfinylbutyl ITC (Sulforaphane); 5MSOP ITC, 5-methylsulfinylpentyl

ITC; 6MSOH ITC; 6-methylsulfinylhexyl ITC; 7MSOH ITC, 7-methylsulfinylheptyl ITC; 8MSOO

ITC; 8-methylsulfinyloctyl ITC; 9MSON ITC; 9-methylsulfinylnonyl ITC; 10MSOD ITC, 10-

methylsulfinyldecyl ITC; 11MSOU ITC, 11-methylsulfinylundecyl ITC; 9MSON NIT, 9-

methylsulfinylnonylnitrile.

Figure 9. Accumulation profiles of flavonoids proteins and metabolites in SCE and SE

A) Expression of proteins involved in flavonoid biosynthesis, degradation, transport and glycosylation in camelina SCE and SE. Only differentially accumulated proteins (DAPs, ANOVA, FDR < 0.05) are showed in the figure. B) Heatmap showing flavonoid metabolites that are differentially accumulated during seed development in SCE and SE tissues.

Figure 10. RNA-Seq transcriptomic analyses and expression of flavonoid and GSL-related genes
A) Principal component analyses using the RNA-Seq transcriptomic data of SCE and SE harvested at several developmental stages and during germination. DAF = Days After Flowering. B) Heatmap showing the expression behaviour of GSL and flavonoid –related genes during seed development in SCE and SE tissues.

SUPPORTING INFORMATION
Table S1. Top-10 most accumulated specialized metabolites in *C. sativa* seed coats and endosperms, and in the embryos. Average intensity of each metabolic feature is expressed as relative intensity according to the internal standard Apigenin.

Table S2. *Camelina sativa* **seed specialized metabolome (LC-MS/MS)**
(a) Annotation and information on Camelina sativa seed coat/endosperm and embryo specialized metabolites. Identification (ID), ionization mode (mode), m/z, retention time (RT), Molecular Network clusters, Annotation (based on homemade library, standards and public databases), metabolite class (based on molecular network clusters and annotation), and source of the annotation are indicated for each metabolite. n = metabolite or cluster from negative ionization mode, p = metabolite or cluster from positive ionization mode.
(b) Relative accumulation of Camelina sativa seed coat and embryo specialized metabolites (LC-MS/MS). Data are showed for two tissues, eight developmental stages, and three biological replicates considered in this study. The QCs (quality control) were obtained by mixing all the samples considered in the analyses. SCE = seed coat and endosperm, SE = seed embryo. DAF = days after flowering.

Table S3. *Camelina sativa* **seed proteome (DIA)**
Relative accumulation of *Camelina sativa* seed coat/endosperm and embryo proteins (DIA, TIMS-TOF). Data are showed for two tissues, eight developmental stages, and three biological replicates considered in this study. Corresponding *A. thaliana* (At ID) and protein description are showed when available. SCE = seed coat and endosperm, SE = seed embryo. DAF = days after flowering

Table S4. Annotation, information and accumulation on *Brassica napus* **(a) and** *Arabidopsis thaliana* **(b) seed coat/endosperm and embryo glucosinolates and related degradation products**

(LC-MS/MS).

Identification (ID), ionization mode (mode), m/z, retention time (RT), Annotation (based on homemade

library, standards and public databases), metabolite class (based on molecular network clusters and

annotation), and source of the annotation are indicated for each metabolite

n = metabolite or cluster from negative ionization mode, p = metabolite or cluster from positive

ionization mode.

Table S5. *Camelina sativa* **seed transcriptome (RNA-Seq)**

Expression of Camelina sativa seed coat/endosperm and embryo genes (RNA-Seq). Data are showed

for embryo tissue, six developmental stages, and three biological replicates considered in this study.

Corresponding *A. thaliana* (At ID) and gene description are showed when available. SE = seed embryo.

DAF = days after flowering.

Figure S1. Accumulation profiles of Differentially accumulated proteins (DAPs)

Heatmaps showing DAPs that are differentially accumulated during seed development in SCE and SE

tissues.

Figure S2. Expression of proteins involved in cinnamic acids and monolignols pathways

Expression of proteins involved in cinnamic acids and monolignols biosynthesis and decorations in

camelina SCE and SE. Only differentially accumulated proteins (DAPs, ANOVA, FDR < 0.05) are

showed in the figure.

Figure S3. Accumulation profiles of cinnamic acids and monolignols metabolites in SCE and SE

Heatmap showing cinnamic acids and monolignols metabolites that are differentially accumulated

during seed development in SCE and SE tissues.

Figure S4. Expression of SM-related homeologs proteins

The number of copies expressed for proteins differentially expressed (FDR < 0.05) in SE and SCE and

involved in GSLs, flavonoids, and cinnamic acids and monolignols pathways.

Figure S5. Differentially Expressed Genes (DEGs) in the embryo (SE) during seed development

Figure S6. Expression of cinnamic acids and monolignols -related genes

Heatmap showing the expression behaviour of cinnamic acids and monolignols –related genes during

seed development in SCE and SE tissues.

REFERENCES

Abdel-Massih RM, Debs E, Othman L, Attieh J, Cabrerizo FM (2023) Glucosinolates, a natural chemical arsenal: More to tell than the myrosinase story. Front Microbiol **14**: 1–15

Alberghini B, Zanetti F, Corso M, Boutet S, Lepiniec L, Vecchi A, Monti A (2022) Camelina [Camelina sativa (L .) Crantz] seeds as a multi-purpose feedstock for bio-based applications. Ind Crop Prod **182**: 114944

Arimura G, Maffei M (2016) Plant Specialized Metabolism: Genomics, Biochemistry, and Biological Functions, Taylor and. Taylor and Francis, New York, NY

Barreda L, Brosse C, Boutet S, Perreau F, Rajjou L, Lepiniec L, Corso M (2024) Specialized metabolite modifications in Brassicaceae seeds and plants: diversity, functions and related enzymes. Nat Prod Rep **41**: 834–859

Baud S, Corso M, Debeaujon I, Dubreucq B, Job D, Marion-Poll A, Miquel M, North H, Rajjou L, Lepiniec L (2023) Recent progress in molecular genetics and omics-driven research in seed biology. Comptes Rendus - Biol **345**: 61–110

Baumert A, Milkowski C, Schmidt J, Nimtz M, Wray V, Strack D (2005) Formation of a complex pattern of sinapate esters in Brassica napus seeds, catalyzed by enzymes of a serine carboxypeptidase-like acyltransferase family? Phytochemistry **66**: 1334–1345

Bell JM (1993) Factors affecting the nutritional value of canola meal: A review. Can J Anim Sci **73**: 689–697

Boisson M, Gomord V, Audran C, Berger N, Dubreucq B, Granier F, Lerouge P, Faye L, Caboche M, Lepiniec L (2001) Arabidopsis glucosidase I mutants reveal a critical role of N-glycan trimming in seed development. EMBO J **20**: 1010–1019

Boutet S, Barreda L, Perreau F, Totozafy J-C, Mauve C, Gakière B, Delannoy E, Martin-Magniette M-L, Monti A, Lepiniec L, et al (2022) Untargeted metabolomic analyses reveal the diversity and plasticity of the specialized metabolome in seeds of different Camelina sativa genotypes. Plant J **110**: 147–165

Brown PD, Tokuhisa JG, Reichelt M, Gershenzon J (2003) Variation of glucosinolate accumulation among different organs and developmental stages of Arabidopsis thaliana. Phytochemistry **62**: 471–481

Bussell JD, Reichelt M, Wiszniewski AAG, Gershenzon J, Smith SM (2014) Peroxisomal ATP-binding cassette transporter COMATOSE and the multifunctional protein ABNORMAL INFLORESCENCE MERISTEM are required for the production of benzoylated metabolites in arabidopsis seeds. Plant Physiol **164**: 48–54

Chen Y, Chen Y, Shi C, Huang Z, Zhang Y, Li S, Li Y, Ye J, Yu C, Li Z, et al (2018) SOAPnuke: A MapReduce acceleration-supported software for integrated quality control and preprocessing of high-throughput sequencing data. Gigascience **7**: 1–6

Cock PJA, Fields CJ, Goto N, Heuer ML, Rice PM (2009) The Sanger FASTQ file format for sequences with quality scores, and the Solexa/Illumina FASTQ variants. Nucleic Acids Res **38**: 1767–1771

Corso M, Perreau F, Mouille G, Lepiniec L (2020) Specialized Phenolic compounds in seeds: structures, functions, and regulations. Plant Sci **296**: 110471

Corso M, Perreau F, Rajjou L, Ben Malek R, Lepiniec L, Mouille G (2021) Specialized metabolites in seeds. Adv. Bot. Res. pp 35–70

Czerniawski P, Piasecka A, Bednarek P (2021) Evolutionary changes in the glucosinolate biosynthetic capacity in species representing Capsella, Camelina and Neslia genera. Phytochemistry **181**: 112571

Dalling JW, Davis AS, Arnold AE, Sarmiento C, Zalamea PC (2020) Extending Plant Defense Theory to Seeds. Annu Rev Ecol Evol Syst **51**: 123–141

Debeaujon I, Nesi N, Perez P, Devic M, Grandjean O, Caboche M, Lepiniec L (2003) Proanthocyanidin-Accumulating Cells in Arabidopsis Testa: Regulation of Differentiation and Role in Seed Development. Plant Cell **15**: 2514–2531

Demichev V, Messner CB, Vernardis SI, Lilley KS, Ralser M (2020) DIA-NN: neural networks and interference correction enable deep proteome coverage in high throughput. Nat Methods **17**: 41–44

Fang J, Reichelt M, Hidalgo W, Agnolet S, Schneider B (2012) Tissue-Specific Distribution of Secondary Metabolites in Rapeseed (Brassica napus L.). PLoS One. doi: 10.1371/journal.pone.0048006

Faure J-D, Tepfer M (2016) Camelina, a Swiss knife for plant lipid biotechnology. OCL **23**: D503

Fraser CM, Thompson MG, Shirley AM, Ralph J, Schoenherr JA, Sinlapadech T, Hall MC, Chapple C (2007) Related arabidopsis serine carboxypeptidase-like sinapoylglucose acyltransferases display distinct but overlapping substrate specificities. Plant Physiol **144**: 1986–1999

Di Gioia F, Pinela J, Bailón ADH, Fereira ICFR, Petropoulos SA (2019) The dilemma of "good" and "bad" glucosinolates and the potential to regulate their content. Glucosinolates Prop Recover Appl. doi: 10.1016/B978-0-12-816493-8.00001-9

Halkier BA, Gershenzon J (2006) Biology and biochemistry of glucosinolates. Annu Rev Plant Biol **57**: 303–333

Hölzl G, Rezaeva BR, Kumlehn J, Dörmann P (2023) Ablation of glucosinolate accumulation in the oil crop Camelina sativa by targeted mutagenesis of genes encoding the transporters GTR1 and GTR2 and regulators of biosynthesis MYB28 and MYB29. Plant Biotechnol J **21**: 189–201

Huang J, Rozwadowski K, Bhinu VS, Schäfer U, Hannoufa A (2008) Manipulation of sinapine, choline and betaine accumulation in Arabidopsis seed: Towards improving the nutritional value of the meal and enhancing the seedling performance under environmental stresses in oilseed crops. Plant Physiol Biochem **46**: 647–654

Kagale S, Koh C, Nixon J, Bollina V, Clarke WE, Tuteja R, Spillane C, Robinson SJ, Links MG, Clarke C, et al (2014) The emerging biofuel crop Camelina sativa retains a highly undifferentiated hexaploid genome structure. Nat Commun **5**: 3706

Kliebenstein DJ, D'Auria JC, Behere AS, Kim JH, Gunderson KL, Breen JN, Lee G, Gershenzon J, Last RL, Jander G (2007) Characterization of seed-specific benzoyloxyglucosinolate mutations in Arabidopsis thaliana. Plant J **51**: 1062–1076

Kliebenstein DJ, Lambrix VM, Reichelt M, Gershenzon J, Mitchell-Olds T (2001) Gene duplication in the diversification of secondary metabolism: Tandem 2-oxoglutarate-dependent dioxygenases control glucosinolate biosynthesis in arabidopsis. Plant Cell **13**: 681–693

Lee S, Kaminaga Y, Cooper B, Pichersky E, Dudareva N, Chapple C (2012) Benzoylation and sinapoylation of glucosinolate R-groups in Arabidopsis. Plant J **72**: 411–422

Lepiniec L, Debeaujon I, Routaboul J-M, Baudry A, Pourcel L, Nesi N, Caboche M (2006) Genetics and Biochemistry of Seed Flavonoids. Annu Rev Plant Biol **57**: 405–430

Mácová K, Prabhullachandran U, Štefková M, Spyroglou I, Pěnčík A, Endlová L, Novák O, Robert HS (2022) Long-Term High-Temperature Stress Impacts on Embryo and Seed Development in Brassica napus. Front Plant Sci **13**: 844292

Mandáková T, Pouch M, Brock JR, Al-Shehbaz IA, Lysak MA (2020) Origin and Evolution of Diploid and Allopolyploid Camelina Genomes Were Accompanied by Chromosome Shattering. Plant Cell **31**: 2596–2612

Mann A, Kumari J, Kumar R, Kumar P, Pradhan AK, Pental D, Bisht NC (2023) Targeted editing of multiple homologues of GTR1 and GTR2 genes provides the ideal low-seed, high-leaf glucosinolate oilseed mustard with uncompromised defence and yield. Plant Biotechnol J 1–14

Matsuda F, Hirai MY, Sasaki E, Akiyama K, Yonekura-Sakakibara K, Provart NJ, Sakurai T, Shimada Y, Saito K (2010) AtMetExpress Development: A Phytochemical Atlas of Arabidopsis Development. Plant Physiol **152**: 566–578

Matthäus B, Zubr J (2000) Variability of specific components in Camelina sativa oilseed cakes. Ind Crops Prod **12**: 9–18

Menard GN, Langdon M, Bhunia RK, Shankhapal AR, Noleto-Dias C, Lomax C, Ward JL, Kurup S, Eastmond PJ (2022) Diverting phenylpropanoid pathway flux from sinapine to produce industrially useful 4-vinyl derivatives of hydroxycinnamic acids in Brassicaceous oilseeds. Metab Eng **70**: 196–205

Missinou AA, Ferreira de Carvalho J, Marnet N, Delhaye T, Hamzaoui O, Abdel Sayed D, Guitton Y, Lebreton L, Langrume C, Laperche A, et al (2022) Identification and Quantification of Glucosinolates and Phenolics in a Large Panel of Brassica napus Highlight Valuable Genetic Resources for Chemical Ecology and Breeding . J Agric Food Chem. doi: 10.1021/acs.jafc.1c08118

Neupane D, Lohaus RH, Solomon JKQ (2022) Realizing the Potential of Camelina sativa as a Bioenergy Crop for a Changing Global Climate. Plants **11**: 772

Nour-Eldin HH, Halkier BA (2009) Piecing together the transport pathway of aliphatic glucosinolates. Phytochem Rev **8**: 53–67

Nour-Eldin HH, Madsen SR, Engelen S, Jørgensen ME, Olsen CE, Andersen JS, Seynnaeve D, Verhoye T, Fulawka R, Denolf P, et al (2017) Reduction of antinutritional glucosinolates in *Brassica* oilseeds by mutation of genes encoding transporters. Nat Biotechnol **35**: 377–382

Olivon F, Elie N, Grelier G, Roussi F, Litaudon M, Touboul D (2018) MetGem Software for the Generation of Molecular Networks Based on the t-SNE Algorithm. Anal Chem **90**: 13900–13908

Pang Z, Chong J, Zhou G, Anderson De Lima Morais D, Chang L, Barrette M, Gauthier C, Jacques P-´ E, Li S, Xia J (2021) MetaboAnalyst 5.0: narrowing the gap between raw spectra and functional insights. Nucleic Acids Res. doi:

10.1093/nar/gkab382

Pang Z, Zhou G, Ewald J, Chang L, Hacariz O, Basu N, Xia J (2022) Using MetaboAnalyst 5.0 for LC–HRMS spectra
processing, multi-omics integration and covariate adjustment of global metabolomics data. Nat Protoc **17**: 1735–1761

Quéro A, Molinié R, Mathiron D, Thiombiano B, Fontaine JX, Brancourt D, Van Wuytswinkel O, Petit E, Demailly
H, Mongelard G, et al (2016) Metabolite profiling of developing Camelina sativa seeds. Metabolomics **12**: 1–14

Raab T, Barron D, Arce Vera F, Crespy V, Oliveira M, Williamson G (2010) Catechin Glucosides: Occurrence,
synthesis, and stability. J Agric Food Chem **58**: 2138–2149

Rao SQ, Chen XQ, Wang KH, Zhu ZJ, Yang J, Zhu B (2021) Effect of short-term high temperature on the accumulation
of glucosinolates in Brassica rapa. Plant Physiol Biochem **161**: 222–233

Reichelt M, Brown PD, Schneider B, Oldham NJ, Stauber E, Tokuhisa J, Kliebenstein DJ, Mitchell-Olds T,
Gershenzon J (2002) Benzoic acid glucosinolate esters and other glucosinolates from Arabidopsis thaliana.
Phytochemistry **59**: 663–671

Robinson MD, Mccarthy DJ, Smyth GK (2010) edgeR: a Bioconductor package for differential expression analysis of
digital gene expression data. Bioinforma Appl NOTE **26**: 139–140

Routaboul J-M, Dubos C, Beck G, Marquis C, Bidzinski P, Loudet O, Lepiniec L (2012) Metabolite profiling and
quantitative genetics of natural variation for flavonoids in Arabidopsis. J Exp Bot **63**: 3749–3764

Routaboul JM, Kerhoas L, Debeaujon I, Pourcel L, Caboche M, Einhorn J, Lepiniec L (2006) Flavonoid diversity and
biosynthesis in seed of *Arabidopsis thaliana*. Planta **224**: 96–107

Sakai K, Taconnat L, Borrega N, Yansouni J, Brunaud V, Paysant-Le Roux C, Delannoy E, Martin Magniette M-L,
Lepiniec L, Denis Faure J, et al (2018) Combining laser-assisted microdissection (LAM) and RNA-seq allows to
perform a comprehensive transcriptomic analysis of epidermal cells of Arabidopsis embryo. Plant Methods **14**: 10

Sanden NCH, Kanstrup C, Crocoll C, Schulz A, Nour-Eldin HH, Halkier BA, Xu D (2023) An UMAMIT-GTR
transporter cascade controls glucosinolate seed loading in Arabidopsis. Nat Plants doi: 10.1038/s41477-023-01598-4

Stehle F, Brandt W, Schmidt J, Milkowski C, Strack D (2008) Activities of Arabidopsis sinapoylglucose:malate
sinapoyltransferase shed light on functional diversification of serine carboxypeptidase-like acyltransferases.
Phytochemistry **69**: 1826–1831

Traka MH (2016) Health Benefits of Glucosinolates. Adv Bot Res **80**: 247–279

Wilson AE, Bergaentzlé M, Bindler F, Marchioni E, Lintz A, Ennahar S (2013) In vitro efficacies of various
isothiocyanates from cruciferous vegetables as antimicrobial agents against foodborne pathogens and spoilage
bacteria. Food Control **30**: 318–324

Xie D-Y, Sharma SB, Paiva NL, Ferreira D, Dixon RA (2003) Role of anthocyanidin reductase, encoded by BANYULS in
plant flavonoid biosynthesis. Science (80-) **299**: 396–399

Xu D, Christian N, Sanden H, Hansen LL, Belew ZM, Madsen SR, Meyer L, Jørgensen ME, Hunziker P, Veres D, et
al (2023) Export of defensive glucosinolates is key for their accumulation in seeds. Nature **617**: 132–138

Yan D, Duermeyer L, Leoveanu C, Nambara E (2014) The functions of the endosperm during seed germination. Plant
Cell Physiol **55**: 1521–1533

Zanetti F, Alberghini B, Marjanovi A, Grahovac N, Rajkovi D, Kiprovski B, Monti A (2021) Camelina, an ancient
oilseed crop actively contributing to the rural renaissance in Europe. A review. Agron Sustain Dev **41**: 2

Zanetti F, Eynck C, Christou M, Krzyżaniak M, Righini D, Alexopoulou E, Stolarski MJ, Van Loo EN, Puttick D,
Monti A (2017) Agronomic performance and seed quality attributes of Camelina (Camelina sativa L. crantz) in multi-
environment trials across Europe and Canada. Ind Crops Prod **107**: 602–608

Zerbib M, Cazals G, Enjalbal C, Saucier C (2018) Identification and Quantification of Flavanol Glycosides in Vitis
vinifera Grape Seeds and Skins during Ripening. Molecules. doi: 10.3390/molecules23112745

Zhang S, Ghatak A, Bazargani MM, Bajaj P, Varshney RK, Chaturvedi P, Jiang D, Weckwerth W (2021) Spatial
distribution of proteins and metabolites in developing wheat grain and their differential regulatory response during the
grain filling process. Plant J **107**: 669–687

Zubr J (1997) Oil-seed crop: Camelina sativa. Ind Crops Prod **6**: 113–119

Camelina sativa seed development

Days after flowering

| 13 | 21 | 28 | 35 | 42 | Dry Mature | Germination 6h | 24h |

1 mm

1 mm

% Weight of SCE and SE during seed development

SCE = Seed Coat and Endosperm
SE = seed embryo

Untargeted metabolomics

A Molecular network

ESI + ESI -

- Aminoacids and derivatives
- Carbohydrates
- Carotenoids
- Chalcones
- Cinnamic acids
- Flavonoids
- Glucosinolates
- Lignans
- Nucleosides and derivatives
- Phenylpropanoids
- Terpenoids
- Unknown / not annotated

B Annotated Metabolic Categories

Nr features

Category	Value
Amines	6
Isothiocyanates	7
GSH	7
Lignans	9
Sulfoxides	11
Flavonoids	17
Glucosinolates	18
Carboxylic acids	23
Carbohydrates	25
Fatty amides	29
Flavan-3-ols and PAs	32
Terpenoids	39
Ns and derivatives	47
Flavonols	64
Cinnamic acids	176
AA and derivatives	208

AA = aminoacids
Ns = nucleosides
PAs = proanthocyanidins

C PCA – untargeted metabolomic data

PC2 = 19,6%
PC1 = 24,3%

Tissue
- Seed coat and endosperm
- Seed embryo

Developmental stage
- 13 DAF
- 21 DAF
- 28 DAF
- 35 DAF
- 42 DAF
- Dry seed
- Germ 6h
- Germ 24h

D Differentially Accumulated Metabolites

Nr Features

	Seed coat and endosperm	Seed Embryo
DAMs	2523 (62%)	2973 (73%)
Not-Significant	1538	1088

Not-Significant   DAMs

Clustering of differentially accumulated metabolic features

A — Seed coat and endosperm (SCE)

Enriched Categories

Cluster	Metabolic Class	Nr features	% SM class
I	Carboxylic acids	13	57
	Flavan-3-ols and PAs	13	89
	p_72	3	100
	p_28	3	60
II	Glucosinolates	10	56
	Ns and derivatives	14	30
	p_74	3	100
	p_76	3	100
	p_82	3	100
III	Cinnamic acids	114	65
	AA and derivatives	90	43
	Flavonols	31	48
	Lignans	9	100
	p_22	7	100
	p_16	7	78
	n_1a	7	64
	p_45	4	100
	p_35	4	80
IV	AA and derivatives	17	8
V	Sulfoxides	5	45
	Amines	4	67
	Carbohydrates	4	16

Relative accumulation

Lower ——— Higher
0   50   100

B — Seed Embryo (SE)

Enriched Categories

Cluster	Metabolic Class	Nr features	% SM class
VI	No enrichment		
VII	Fatty amides	18	59
	Terpenoids	17	46
	n_29	3	100
VIII	Cinnamic acids	111	63
	AA and derivatives	105	50
	p_22	7	100
	Flavonols	36	56
	Carboxylic acids	16	70
	Flavan-3-ols and PAs	19	59
	GSH	6	86
	Sulfoxides	8	73
	p_45	4	100
	Amines	5	83
	p_74	3	100
IX	AA and derivatives	17	8
	Isothiocyanates	4	31
	p_72	3	100

DAF = days after flowering
AA = aminoacids
Ns = nucleosides
PAs = proanthocyanidins

Proteomics

A

PCA – proteomic data

Tissue
- ◉ Seed coat
- ◯ Seed embryo

Developmental Stage
- ● 13 DAF
- ● 21 DAF
- ● 28 DAF
- ● 35 DAF
- ● 42 DAF
- ● Dry seed
- ● Germ 6h
- ● Germ 24h

B

Differentially Accumulated Proteins (DAPs)
FDR < 0.05

Nr DEPs

4907 (43%)

6330 (55%)

Seed Coat and endosperm

Seed Embryo

DAPs SM SCE — 59 (14%)

273 (63%)

DAPs SM SE — 101 (23%)

Specialized Metabolites (SM) –related Proteins

A

Accumulation behaviour of SM–related DAPs

Seed coat and endosperm (SCE)
(333 DAPs)

B   Accumulation behavior of SM categories

Seed embryo (SE)
(377 DAPs)

Relative Expression

Lower 0 — 50 — Higher 100

DAPs = Differentially Accumulated Proteins

% related to the highest value

0 — 50 — 100

(a) GSL and GSL-degradation-products distribution in Brassicaceae seeds

Camelina sativa Céline — 1mm

Brassica napus Aviso and Major — 1mm

Arabidopsis thaliana Col-0 — 0.5mm

(b) Comparison of seed GSL metabolites and genes-enzymes among Brassicaceae species

Transcriptomic analyses (RNA-Seq) - Seed Embryo